The inflammatory accumulation of lipids and ROS in human Nrf1α-deficient hepatoma cells is
ameliorated by 2-bromopalmitate

Ze Zheng[1], Meng Wang[1], Shaofan Hu[1], Rongzhen Deng[1], Jing Feng[1], and Yiguo Zhang[1]

[1]The Laboratory of Cell Biochemistry and Topogenetic Regulation, College of Bioengineering and Faculty of Medical Sciences,
Chongqing University, No. 174 Shazheng Street, Shapingba District, Chongqing 400044, China
* Correspondence: yiguozhang@cqu.edu.cn, or eaglezhang64@gmail.com

Abstract: Since Nrf1 and Nrf2 are essential for regulating the lipid metabolism pathways, their dysregulation was also
shown to be critically involved in the non-controllable inflammatory pathology into cancer development. However, it
is unknown that the interaction between Nrf1 and Nrf2 in the regulation of lipid metabolism, especially in hepatoma
cells. Here, we have further explored the molecular mechanisms underlying their distinct regulation of lipid
metabolism, by comparative analysis of changes in those lipid metabolism-related genes in $Nrf1\alpha^{-/-}$ and/or $Nrf2^{-/-}$
cell lines relative to wild-type controls. The results revealed that loss of Nrf1 leads to disordered lipid metabolism; its
lipid synthesis pathway was up-regulated by JNK-Nrf2-AP1 signaling, while its lipid decomposition pathway was
down-regulated by the nuclear receptor PPAR-PGC1 signaling, resulting in severe accumulation of lipids as deposited
in lipid droplets. By contrast, knockout of Nrf2 gave rise to decreases in lipid synthesis and uptake capacity. These
demonstrate that Nrf1 and Nrf2 contribute to significant differences in the cellular lipid metabolism regulatory
profiles and relevant pathological responses. Further experiments unraveled that lipid deposition in $Nrf1\alpha^{-/-}$ cells was
resulted from CD36 upregulation by activating the PI3K-AKT-mTOR pathway, leading to induction of the inflammatory
response. Following treatment of $Nrf1\alpha^{-/-}$ cells with 2-bromopalmitate (2BP), it enabled the yield of lipid droplets to
be strikingly alleviated, as companied by substantial abolishment of CD36 and critical inflammatory cytokines. Such
$Nrf1\alpha^{-/-}$-led inflammatory accumulation of lipids and ROS was significantly ameliorated by 2BP. Overall, this study
provides a potential strategy for cancer prevention and treatment by precision targeting of Nrf1, Nrf2, or both.
Key words: Nrf1; Nrf2; lipid metabolism; lipid droplets; JNK; AP1; mTOR; CD36; IL-6; ROS; 2-bromopalmitate

1. Introduction

Lipids as biological macromolecules are essential for energy storage and structural membranes, and also crucial for a
variety of biological functions in most life processes, including cell growth, proliferation, differentiation, and
movement. Therein, many types of lipids act as important biological signal molecules and thus exert diverse
functionality in energy metabolism and cell homeostasis during the healthy life process. Contrarily, abnormal lipid
metabolism can lead to a variety of physiological diseases. For instance, obesity caused by over-nutrition,
nonalcoholic steatohepatitis (NASH), and type 2 diabetes are all related to the imbalance of lipid metabolism status
(1). In addition to specific metabolic syndromes, the development of hepatocellular carcinoma (HCC) is also
inseparable from lipid disorders (2). This is because most cancer cells were shown to have a higher rate of *de novo*
lipid synthesis, leading to an increase in newly-synthesized fatty acids (e.g., glycerol phospholipids and cholesterol)
that serve the raw materials so as to meet the needs for malignant growth and proliferation of cancer cells. Besides,
the decomposing metabolism of lipids by β-oxidation in the mitochondria can also provide enough energy, when
required for cancer cells (3).
In almost all living organisms, the lipids exist as a major stored form of triglycerides and cholesterol in lipid
droplets (LDs), a kind of hydrophobic organelle with monolayer membrane structure that can mobilize fatty acids
during specific life processes (4). Since the liver is the main metabolic organ of lipids in vertebrates, these lipids play a
vital role in the liver's physiological homeostasis and pathological processes of many relevant diseases. Of note,
liver-specific knockout of *Nrf1* [also called *Nfe2l1*, encoding a highly conserved membrane-bound transcription factor
of the cap'n'collar (CNC) basic-region leucine zipper (bZIP) family(5)] in the mouse leads to spontaneous development

of NASH and malignant transformation into hepatoma (6). Such hepatocyte-specific loss of Nrf1 caused up-regulation of the cytochrome p450 family *CYP4A* and *CYP1A1* genes, as fatty acid (FA) and cystine/cysteine contents increased, followed by severe steatosis, apoptosis, necrosis, and inflammation and fibrosis, eventually worsening and developing into liver cancer (7,8). During this pathogenesis, significant activation of two lipid metabolism-related genes *Lipin1* and *PGC-1β* was also shown in the murine liver-specific *Nrf1⁻/⁻* cells, implying that Nrf1 negatively regulates these two genes by its direct binding to the putative antioxidant response elements (AREs) located in their promoter regions (9).

Nrf2 (also encoded by *Nfe2l1*, as a water-soluble transcription factor of the CNC-bZIP family) has been proved to regulate lipid metabolism by mediating the expression of a number of lipid metabolism-related genes, such as those encoding ACLY, ACC1 and ELOVL6. It was found by Huang *et al* (10) that *Nrf2⁻/⁻* mice were manifested with a loss of their liver weight at six months of age, as accompanied by a marked decrease of the triacylglycerol contents in their livers. Such changes were shown to be attributable to *Nrf2⁻/⁻*-leading up-regulation of lipogenic genes including *PPARγ*, *FASN*, *SCD1* and *SREBP1*, albeit they were proved to serve direct target genes of Nrf2 (10). The adipose-specific ablation of Nrf2 caused a transient delay of high-fat diet-induced obesity by altering glucose, lipid and energy metabolism in the mice (11). Altogether, these demonstrate that Nrf2 acts as an important regulatory player in lipid metabolism. However, how Nrf1 and Nrf2 differentially govern (and inter-regulate) the lipid metabolism process, in particular liver cancer cells, is poorly understood to date.

In this study, we focus to determine the lipid synthesis and catabolism, as well as the mechanism related to lipid deposition, by conducting various experiments to examine yield of lipid droplets in *Nrf1α⁻/⁻* cells and relevant changes in its lipid metabolism gene expression as compared with those obtained from in *Nrf2⁻/⁻* cells. The results unveiled that Nrf1 and Nrf2 have performed their respective regulatory functions in the lipid metabolism process of hepatoma cells through distinct molecular mechanisms. Further experiments also revealed that both factors are differentially involved in putative cytoprotection against the inflammatory responses to abnormal lipid metabolism caused by *Nrf1α⁻/⁻* and/or *Nrf2⁻/⁻*. Interestingly, such lipid deposition and related inflammation responses are significantly reduced by a palmitic acid analogue, 2-bromopalmitate (2BP). Collectively, this study provides a novel potential strategy for the prevention and treatment of NASH and hepatoma by precision targeting of Nrf1, Nrf2, or both.

2. Materials and Methods

2.1 Cell lines and culture

The human hepatocellular carcinoma HepG2 cells (i.e., *Nrf1/2⁺/⁺*) were obtained originally from the American Type Culture Collection (ATCC, Manassas, VA, USA). Additional two derived *Nrf1α⁻/⁻* and *Nrf2⁻/⁻* cell lines were established in our laboratory by Qiu *et al* (12). The fidelity was confirmed by its authentication profiling and STR (short tandem repeat) typing map (by Shanghai Biowing Applied Biotechnology Co., Ltd). Before experimentation, they were allowed for growth in Dulbecco's modified Eagle's medium (DMEM), containing 25 mmol/L high glucose, 10% (v/v) fetal bovine serum (FBS), 100 units/mL penicillin-streptomycin, by being cultured in a 37℃ incubator with 5% CO2.

2.2 Real-time quantitative PCR analysis

After all experimental cell lines reached 80% of their confluence, the total RNAs were firstly extracted with no treatment. In the parallel experiments, these cell lines were subjected to different treatments with specific inhibitors or siRNA (in order to interfere with cognate target gene expression). Subsequently, total RNAs were extracted by using an RNA extraction kit (TIANGEN, Beijing, China) and then reversely transcribed into single-stranded cDNA by a Reverse-Transcription Kit (Promega, USA). The mRNA expression level of each target gene in these cell lines was

87 determined by real-time quantitative PCR (RT-qPCR) with each of the indicated pairs of their forward and reverse
88 primers (in Table 1).

89 **Table 1. All oligonucleotides and antibodies used in this study.**
90 (i)   The primer pairs used for RT-qPCR analysis.

Gene Name	Forward Primers (5' to 3')	Reverse Primers (5' to 3')
β-actin	GCGGCTACAGCTTCACCAC	GCCATCTCTTGCTCGAAGTCCA
Nrf1	CCCCTGCTCTTCAAAACAACC	ATCTTGCCACTGCTGTTCC
Nrf2	AGTCCAGAAGCCAAACTGACA	ATGCTGCTGAAGGAATCCTCA
FASN	ATTCGACCTTTCTCAGAACCAC	AAGAACGCATCCAGTAGGAC
SCD1	ATGCTAATTCAATGCCGCAAC	CTTCCCCAGCAGAGACCAC
ACCα	CTCAGAGGCGAGAAACTCCA	CATCAATCGCTCTACCCAGT
SREBP1	GCCATCGACTACATTCGCTTT	GCACGTCTGTGTTCCCTCC
HO1	CAGTGCCACCAAGTTCAAGCA	CTTGGCCTCTTCTATCACCCT
JUN	ATGGAAACGACCTTCTATGACGA	CGTTGCTGGACTGGATTATCA
NQO1	ATTCCTTAGCCTGGATTTCCT	GCACGAATACAGTCGATTCCC
GCLM	ACGAGAATCGCTTGAACCT	AGTTTTCCCAGGCAAATCACA
GCLC	CCCGATATGACTCAATAGACAGC	ATGATCAATGCCTTCCTGCAAC
CD36	TACAGATGCAGCCTCATTTCCAC	ACAGCATAGATTGACCTGCAAA
IL-6	CAAATGTAGCATGGGCACCT	AACATAAGTTCTGTGCCCAGT
IL-1β	CCTGCCCACAGACCTTCCAG	AAGCCTCGTTATCCCATGTGT
NFκB1	CGTCAGAAGCTCATGCCCAA	GCCACTACCAAACATGCCTCC
NFκB2	CCTGAAGCCAGTCATCTCCC	GCACCTTGTCACAAAGCAGA
TNFα	CCCTCAACCTCTTCTGGCTCA	GCCACACATTCCTGAATCCCA

91 (ii)   Oligonucleotides sequences of siRNA.

Name	Forward Primers (5' to 3')	Reverse Primers (5' to 3')
siNrf2	GUAAGAAGCCAGAUGUUAAdTdT	UUAACAUCUGGCUUCUUACdTdT
siJUN	GCAUGGACCUAACAUUCGAdTdT	UCGAAUGUUAGGUCCAUGCdTdT

92 (iii)   The key antibodies used in this work.

Antibodies	Identifier	Source
Nrf1	N/A	Zhang's(13)
Nrf2	ab62352	Abcam
ACCα	D155300	Sangon Biotech
FASN	D262701	Sangon Biotech
SCD1	D162163	Sangon Biotech
SREBP1	14088-1-AP	Proteintech
ATGL	ab109251	Abcam
PPARα	D161086	Sangon Biotech
PPARγ	ab272718	Abcam
PPARδ	ab178866	Abcam
PGC-1α	D262041	Sangon Biotech
PGC-1β	D162042	Sangon Biotech
JNK	ab208035	Abcam
p-JNK	ab124956	Abcam
JUN	10024-2-AP	Proteintech
FOS	ab134122	Abcam
HO1	ab68477	Abcam
AKT1	ab32505	Abcam
p-AKT	ab38449	Abcam
p-mTOR^{2448}	ab109268	Abcam
p-mTOR^{2481}	ab137133	Abcam
PI3K	ab245781	Abcam
CD36	ab133625	Abcam
COX2	ab62331	Abcam
IL-6	ab233706	Abcam
NFκB	ab32360	Abcam
p-NFκB	ab140751	Abcam
β-actin	TA-09	ZSGB-BIO

2.3 Small interfering RNAs and kinase-specific inhibitors

To determine the effect of a given gene on the downstream signaling, the JNK function inhibitor SP600125 (Selleck,
Shanghai, China), the PI3K inhibitor NVP-BKM120 (Cayman Chemical, USA), and the mTOR inhibitor RAPA (rapamycin,
Sigma, USA) were employed respectively. Two pairs of small-interfering RNAs (i.e., siRNAs, with their oligonucleotide
sequences listed in Table 1) were designed for targeting Nrf2 and JUN that interfere with their mRNA expression
levels, before being synthesized by Tsingke (Beijing, China).

2.4 Western blotting analysis of key proteins

All experimental cells were subjected to extraction of total proteins in a lysis buffer, containing a protease inhibitor
cOmplete Tablets EASYpack or another phosphatase inhibitor PhosSTOP EASYpack (each tablet per 15 mL buffer,
Roche, Basel, Switzerland). The proteins in cell lysates were denatured by boiling at 100℃ for 10 min, disrupted by
ultrasound, and then diluted in 3× loading buffer (187.5 mmol/L Tris-HCl, pH 6.8, 6% SDS, 30% Glycerol, 150 mmol/L
DTT, and 0.3% Bromphenol Blue) at 100℃ for 5 min. Thereafter, equal amounts of proteins were separated by
SDS-PAGE gels containing 8-12% polyacrylamide and visualized by Western blotting with distinct primary antibodies
(listed in Table 1).

2.5 Determination of intracellular triglyceride contents

All the relevant experimental cells were subjected to determination of the triglyceride (TG) contents, as
conducted according to the manufacture's instruction of the TG-enzymatic assay kit (APPLYGEN, Beijing, China).

2.6 Oil Red O-staining of lipid droplets in cells

After all experimental cells reached 70% of their confluence, the cells were subjected to different treatments with
either different concentrations of oleic acid (OA, Solarbio, Beijing, China) or 2-bromopalmitate (2BP, Sigma, USA).
Then, the cells were fixed in a tissue fixative buffer containing 4% paraformaldehyde (Boster Biological Technology,
Wuhan, China) for 30 min, prior to being stained for additional 30 min in another solution containing 3 g/L of oil red
O (Sangon Biotech, Shanghai, China). Subsequently, the stained cells were rinsed three times by using 60% of
isopropyl alcohol (Kelong, Chengdu, China), before the red lipid droplets were visualized by microscopy and
photographed as shown.

2.7 Analysis of cell ROS by flow cytometry

After all experimental cells reached 70% of their confluence, the cells were transferred to be cultured in a medium
containing 2BP or without this chemical for 24 h and then prepared according to the manufacture's instruction of the
ROS assay kit (Beyotime, Shanghai, China). Thereafter, the intracellular ROS levels were determined by flow
cytometry. The FlowJo 7.6.1 software, and the resulting data were analyzed as described previously (14).

2.8 Statistical analysis

Significant differences were statistically determined using the Student's *t*-test and Multiple Analysis of Variations
(MANOVA), except for somewhere indicated. The data are herein shown as a fold change (mean ± S.D.), each of
which represents at least three independent experiments that were each performed in triplicate.

3. Results
3.1. Knockout of Nrf1α leads to lipid deposition while knockout of Nrf2 does the opposite in hepatoma cells.

A striking feature in the pathology of NASH is lipid deposition, which is also a hallmark of liver cancer
development from the murine *Nrf1*$^{-/-}$-disordered lipid metabolism (7,8). Herein, wild-type HepG2 cells (*Nrf1/2*$^{+/+}$)
and two derived cell lines (*Nrf1α*$^{-/-}$and *Nrf2*$^{-/-}$) were employed in the following experiments to investigate distinct
functionality of Nrf1 and Nrf2 in human intracellular lipid metabolism. These cell lines were further validated by
Western blotting and real-time qPCR analyses of Nrf1 and Nrf2 (Fig. 1, A & B); the results were consistent with our

previous data (12,14). When compared with wild-type $Nrf1/2^{+/+}$ controls, $Nrf1\alpha^{-/-}$ cells exhibited no expression of

Nrf1α-derived proteins, but other shorter isoforms of Nrf1ß and Nrf1γ were almost unaltered (Fig.1 A, *upper panel*).

By contrast, Nrf2 protein abundance (Fig.1 A, *middle panel*) and its basal mRNA levels (Fig. 2B) were significantly

increased in $Nrf1\alpha^{-/-}$ cells. Conversely, Nrf1α-derived proteins were slightly enhanced in $Nrf2^{-/-}$ cells (lacking its two

major proteins), but this was accompanied by a modest decrease in basal mRNA expression of *Nrf1* (Fig. 1, A & B).

These indicate putative inter-regulation of between Nrf1α and Nrf2 could be executed at distinct layers.

After 24-h treatment of distinct cell lines with oleic acid (OA) at different concentrations (0, 50, 100 μmol/L), the

yields of lipid droplets were evaluated by oil red O staining. The resulting images were photographed and shown in

(Fig. 1D). When compared with wild-type $Nrf1/2^{+/+}$ controls, much more and larger lipid droplets were yielded in

$Nrf1\alpha^{-/-}$ cells, whereas the least and smallest lipid droplets were in $Nrf2^{-/-}$ cells. Also, yields of lipid droplets after OA

treatment of the above three cell lines were increasing to different extents, with incrementing concentrations of OA

from 50 to 100 μmol/L (Fig. 1D). Such an OA-increased trend of lipid droplets in $Nrf1\alpha^{-/-}$ cells was significantly higher

than that of $Nrf1/2^{+/+}$ controls, while the lowest trend of increased lipid droplets was manifested in $Nrf2^{-/-}$ cells.

These unraveled that loss of Nrf1α led to an aberrant accumulation of lipids, whereas similar lipid deposition

appeared to be alleviated by Nrf2 inactivation. This notion was further evidenced by an intracellular triglyceride (TG)

enzymatic detection assay of the above experimental cell lines that had been treated with 100 μmol/L OA (Fig. 1C).

Their distinct TG contents (in particular lipid droplets) were also visualized by oil red O staining, as shown

photographically in (Fig. 1E). These distinct results demonstrated that loss of Nrf1α enables the lipid deposition to be

augmented in $Nrf1\alpha^{-/-}$ hepatoma cells, but a similar lipid-accumulating phenomenon was not observed in $Nrf2^{-/-}$ cells

and even its lipid synthesis and/or uptake may be also suppressed by Nrf2-specific knockout of its transactivation

domain rather than its DNA-binding domain.

3.2 Loss of Nrf1α or Nrf2 causes different effects on the expression of genes responsible for lipid metabolism.

It had been shown that those genes governing the *de novo* lipid synthesis were highly expressed in cancer cells

to considerably greater extents than their equivalents in normal cells (15). Such key rate-limiting enzymes (e.g., FASN,

ACCα, SCD1) of *de novo* lipid synthesis and regulated transcriptional factor SREBP1 (sterol regulatory element-binding

protein 1) were examined herein. As anticipated, the results revealed that loss of $Nrf1\alpha^{-/-}$ caused substantial

increases in the protein expression of such key genes encoding FASN (fatty acid synthase), ACCα (acetyl-CoA

carboxylase alpha), SCD1 (stearoyl-Coenzyme A desaturase 1) and SREBP1 (Fig. 2A), when compared with wild-type

$Nrf1/2^{+/+}$ cells. Similar augmented expression of their mRNA levels in $Nrf1\alpha^{-/-}$ cells were also determined by real-time

qPCR (Fig. 2B). By sharp contrast, $Nrf2^{-/-}$ cells gave rise to down-expressed levels of these genes to greater or fewer

degrees (Fig. 2, A & B).

Further examinations unraveled a considerably lower expression level of adipose triglyceride lipase (ATGL, also

called PNPLA2, which is the catalytic rate-limiting enzyme in the first step of TG hydrolysis during oxidation catabolic

mobilization (16)) in $Nrf1\alpha^{-/-}$ cells (Fig. 2, C & D). Intriguingly, another modestly decreased expression of ATGL was

also exhibited in $Nrf2^{-/-}$ cells as compared with wild-type cells. Hence, to clarify this issue, further experimental

analysis of those nuclear receptor-types (i.e. PPARα, γ, δ) transcription factors governing oxidative degradation of

fatty acids (17) and their interacting transcriptional co-activators (i.e. PGC-1α, β) (18) was conducted herein. The

results unraveled that all the examined proliferator activated receptors (PPARα, γ, δ) and PPARγ co-activators (PGC-1α,

-1β) were significantly down-regulated in $Nrf1\alpha^{-/-}$ cells (retaining a hyper-active Nrf2), and thus were roughly

unaffected in $Nrf2^{-/-}$ cells but with an exception of down-expressed PPARα only (Fig. 2, C & D). Taken altogether,

these collective data demonstrate that aberrant deposition of lipids in $Nrf1\alpha^{-/-}$ cells result from up-regulation of lipid

anabolism, along with substantial down-expression of lipid catabolism-controlled genes, but their opposite effects

were elicited in $Nrf2^{-/-}$ cells.

Such differential expression profiles of lipid synthesis- and catabolism-related genes in between *Nrf1α*- and
Nrf2-deficient cell lines were further explored. The results showed a markedly increased abundance of
phosphorylated JNK and its cognate target JUN in *Nrf1α*$^{-/-}$ cells (with a hyper-active Nrf2) (Fig. 2E, *e2 to e4*), although
the total protein level of JUK was relatively down-expressed (Fig. 2E, *e1*), when compared with wild-type controls. By
sharp contrast, all those examined protein levels were strikingly diminished upon inactivation of *Nrf2*$^{-/-}$ (Fig. 2E). Such
similar expression trends of Nrf2, p-JUNK and JUN suggest a comparable pattern to those of the aforementioned
lipogenic genes, thus providing a new clue for the further study of their relevant gene regulation.

3.3 The JNK-Nrf2-AP1 signaling is required for up-regulation of lipogenic genes in Nrf1α-deficient cells.
Here, treatment of *Nrf1α*$^{-/-}$ cells with a specific inhibitor SP600125 (to block the activity of phosphorylated JNK,
as shown in Fig. 3A, *a1*) caused substantial down-expression of all seven other examined proteins, including Nrf2,
JUN (a direct substrate of JNK), HO-1, FASN, ACCα, SCD1 and SREBP1 (Fig. 3A, *a2 to a8*). Of note, the lipid
synthesis-related ACCα and SCD1 displayed a profound inhibitory trend similar to those of JUN and Nrf2, albeit
SREBP1 was only partially inhibited by SP600125. This implies that the JUK signaling to Nrf2 and/or AP1 is involved in
the aberrant lipogenesis of *Nrf1α*$^{-/-}$ cells. As such, the treated cells still retained a remnant fraction of JUN and Nrf2,
but their direct downstream HO-1 expression was almost completely abolished, implicating an integrative effect of
both factors on HO-1.
Further experiments revealed that silencing of Nrf2 by its siNrf2 (to interfere with its expression in *Nrf1α*$^{-/-}$ cells)
also led to obvious down-regulation of AP1 (i.e. a heterodimer of JUN and FOS), HO-1 (a direct target of Nrf2 and
AP1), and those lipogenic enzymes FASN, ACCα and SCD1 (Fig. 3B, *b1 to b6*). Next, the real-time qPCR analysis
unraveled that knockdown of Nrf2 by its siNrf2 caused significantly reduced expression of *JUN, FASN, ACCα* and *SCD1*
(Fig. 3C). This indicates a predominant requirement for Nrf2 in regulating AP1 and the key enzymes in lipid synthesis,
besides its cognate target genes *HO-1, NQO1, GCLC* and *GCLM* (Fig. 3C). However, SREBP1 appeared to be roughly
unaffected by silencing of Nrf2 (Fig. 3, B & C).
Additionally marked decreases in the expression of FASN, ACCα and SCD1 at their protein and mRNA levels
resulted from siJUN-led knockdown of AP1 activity in *Nrf1α*$^{-/-}$ cells (Fig. 3, D & E). Nonetheless, SREBP1 was almost
unaltered by silencing of JUN. Further examination of *Nrf1α*$^{-/-}$ cells by transfecting expression constructs for JUN
alone or plus FOS revealed that such forced expression of AP1 led to striking up-regulation of SCD1, but FASN was
unaffected (Fig. 3F, *c.f. f5 with f3*). Intriguingly, only a marginal increase of ACCα was caused by overexpression of JUN
alone, but conversely, a substantial decrease of this enzyme was dictated by co-expression of JUN and FOS (Fig. 3F, *f4*).
Such dual opposite regulation of ACCα by JUN alone or plus FOS may be determined by a putative functional
heterodimer of JUN with another partner, competitively together with the AP1 factor, depending on distinct
experimental settings.

3.4 Nrf1α$^{-/-}$ cell-autonomous induction of the PI3K-AKT-mTOR signaling leads to abnormal hyper-expression of
CD36.
An important nutritional metabolism mechanism sets into account by signaling of the PI3K (phosphatidylinositol
3-kinase) to its downstream AKT (also called protein kinase B) and mTOR (mechanistic target of rapamycin kinase)
(19). Upon activation of mTOR by stimulation of nutritional signals, it enables for up-regulation of CD36 (acting as a
regulator of fatty acid uptake and its transport (20). As anticipated, a significant increase in basal constitutive
expression of CD36 was determined in *Nrf1α*$^{-/-}$ cells, when compared with wild-type controls (Fig. 4A, *a4*). Such
aberrant hyper-expression of CD36 was accompanied by dominant cell-autonomous activation of PI3K^{P110Δ} and AKT,
as well as of phosphorylated AKT and mTOR at distinct sites (Fig. 4A, *a1 to a5*). In addition, *Nrf1α*$^{-/-}$ cells gave rise to
the basally-augmented expression of a general inflammatory marker COX2 (cyclooxygenase 2) (Fig. (Fig. 4A, *a7*); this
enzyme was also called prostaglandin-G/H synthase (PTGS2), involved in the arachidonic acid metabolism.

In order to determine whether $Nrf1\alpha^{-/-}$ cell-autonomous activation of the PI3K-AKT-mTOR signaling is required

for up-regulation of CD36 by its deficiency, a PI3K-specific inhibitor NVP-BKM120 (Fig. 4B) and another mTOR-specific

inhibitor rapamycin (RAPA, Fig. 4C) were allowed for treatment of $Nrf1\alpha^{-/-}$ and $Nrf1/2^{+/+}$ cell lines respectively. As

expected, the results revealed that two phosphorylated protein levels of mTOR were substantially diminished by

NVP-BKM120 and RAPA in their treated cell lines (Figs. 4B, *b1 & b2* and 4C, *c1 & c2*). In the meantime, basal

expression abundances of COX2 and CD36 were also almost abrogated by NVP-BKM120 and RAPA (Figs. 4B, *b7 & b8*

and 4C, *c8 & c9*). However, no or less effects of both inhibitors on all other examined proteins (e.g., JNK, Nrf1, Nrf2

and HO-1) were detected (Figs. 4B, *b3 to b6* and 4C, *c3 to c7*). Altogether, these demonstrate that loss of Nrf1α

causes a non-controllable activation status of CD36, as well as COX2, primarily by the PI3K-AKT-mTOR signaling

pathway.

Such up-regulation of CD36 can also enable it to assist the Toll-like receptor 4 (TLR4) with synergistic activation

of the inflammatory factor NFκB and cytokines IL-6, IL-1β and TNFα (21). Next, we thus determine whether their

changed expression levels in $Nrf1\alpha^{-/-}$ cells have relevance to the hyper-expression of CD36. The results unveiled that

dramatic activation of CD36 by loss of Nrf1α was accompanied by varying enhancements of NFκB, IL-6, IL-1ß, and TNF

α at their protein and/or mRNA expression levels in $Nrf1\alpha^{-/-}$ cells, when compared with those equivalents of

$Nrf1/2^{+/+}$ cells (Fig. 4, D & E). Overall, it is postulated that $Nrf1\alpha^{-/-}$-disordered lipid metabolism may be attributable to

aberrant activation of CD36, with a concomitant inflammatory response.

3.5 The inflammatory accumulation of lipids and ROS in Nrf1α$^{-/-}$ cells is alleviated by 2-bromopalmitate.

As shown schematically in Fig. 5A, 2BP is a palmitic acid analogue with one bromine (Br) atom at the β-C

position, allowing for dramatic improvement of its stability more than that of the prototypic palmitate so as to make

it difficult to proceed to the next step. Thereby, this physicochemical property of 2BP can enable this palmitic acid

analogue to block those biological processes in which palmitic acids should have participated; for example, 2BP was

employed as a natural ligand of PPARδ receptor in fatty acid oxidation (22). Herein, it was found that the triglyceride

(TG) accumulation in $Nrf1\alpha^{-/-}$ cells was significantly inhibited by 2BP (Fig. 5B). Similar treatments with 2BP caused

almost abolishment of basal CD36 expression in both $Nrf1\alpha^{-/-}$ and $Nrf1/2^{+/+}$ cell lines (Fig. 5C), and also enabled for

substantial suppression of lipid droplets in $Nrf1\alpha^{-/-}$ cells to considerably lower levels (Fig. 5D). These findings

demonstrate that 2-bromopalmitate has the capability to alleviate severe lipid deposition mediated by CD36 in

$Nrf1\alpha^{-/-}$ cells.

Further examination revealed that transcriptional expression levels of the inflammatory cytokines IL-6 and TNFα

in $Nrf1\alpha^{-/-}$ cells were significantly suppressed or even completely abolished by 2BP, respectively, when compared with

wild-type controls (Fig. 5E). Also, the basal IL-6 abundance and active phosphorylated NFκB were apparently

repressed by 2BP (Fig. 5C). This implies that the informatory response to the loss of $Nrf1\alpha^{-/-}$ may be ameliorated by

2-bromopalmitate. Flow cytometry analysis unraveled a marked accumulation of the intracellular reactive oxygen

species (ROS, that could be involved in the informatory response) in $Nrf1\alpha^{-/-}$ cells (Fig. 5F, *left panel*, with an

enhanced peak shifted to the right), but such severe ROS accumulation was also obviously mitigated by 2BP (Fig. 5F,

right panel, with a reduced peak shifted to the left). However, no similar ROS changes were observed in $Nrf1/2^{+/+}$

cells (*middle panel*). Collectively, these indicate that 2-bromopalmitate can enable effective alleviation of the

inflammatory accumulation of lipids by enhanced biosynthesis and uptake of fatty acids (through up-regulation of

CD36) in $Nrf1\alpha^{-/-}$ cells. This occurs concomitantly with a reduction of ROS by 2-bromopalmitate to mitigate the

inflammatory response caused by loss of $Nrf1\alpha$.

4. Discussion

In the present study, we have provided the evidence unraveling different or even opposite contributions of Nrf1

and Nrf2 have been made in human lipid metabolism, in addition to our previously-reported glucose metabolism

(14,23), protein homeostasis (proteostasis) (12,24), redox balance and signaling control (25,26). It is of crucial

significance to note that the experimental cell model of $Nrf1\alpha^{-/-}$ with pathophysiological phenotypes appears to

largely resemble those manifested by the liver-specific Nrf1 knockout mice. Such loss of $Nrf1^{-/-}$ results in the

spontaneous development of NASH and eventual liver cancer, which is attributed to the rapid accumulation of a large

amount of lipids, as a key etiology, in the pathological process (6). Here, our experimental evidence demonstrated

that various piles of lipids (i.e., triglyceride contents) were markedly deposited in distinct sizes of the intracellular

lipid droplets widely distributed within human $Nrf1\alpha^{-/-}$ hepatoma cells, particularly following treatment with oleic

and palmitic acids. By contrast, $Nrf2^{-/-}$ cells seemed to be considered destitute of lipid droplets, even after they were

treated with oleic acids.

Interestingly, we have also discovered that aberrant lipid deposition in $Nrf1\alpha^{-/-}$ cells was substantially prevented

by a palmitic acid analogue, i.e., 2-bromopalmitate, through significant down-regulation of CD36 (acting as an

essential translocator of fatty acids). The concomitant inflammatory response and oxidative stress were ameliorated

significantly by 2-bromopalmitate. As a preventive consequence, key inflammatory cytokines and relevant

transcription factor NFκB were markedly down-regulated, following apparent reduction of ROS and lipids in

2-bromopalmitate-treated $Nrf1\alpha^{-/-}$ cells (as illustrated in Fig. 5). However, it should also be noted that, in addition to

the secondary yield of ROS stimulated by the accumulation of lipids, the primary amount of ROS is produced by

$Nrf1\alpha^{-/-}$ cell-autonomous reprogramming of redox metabolism (14,23,26). Thereby, such $Nrf1\alpha^{-/-}$ cell-autonomous

yield of ROS is inferable to be almost unaffected by 2-bromopalmitate. In fact, our evidence has been presented

herein, showing that even though the putative concomitant ROS resulting from severe lipid deposition may be

diminished by this palmitic acid analogue, the entire oxidative stress is not completely mitigated and recovered to the

levels measured from wild-type cells.

We have also provided evidence revealing that aberrant accumulation of lipids in $Nrf1\alpha^{-/-}$ cells result from

incremented uptake of fatty acids through constitutive activation of the PI3K-AKT-mTOR signaling to up-regulation of

CD36 (as shown in Fig. 5). The CD36-augmented uptake of lipids is fully consistent with those results obtained from

the tissue-specific $Nrf1^{-/-}$ mice (7,27). These two independent groups also unraveled that Nrf1 is essential for

controlling translocation of fatty acids and cystine/cysteine contents by governing a complex hierarchical regulatory

network in the redox and lipid metabolism. Therein, some cytochrome family members, besides CD36, were also

up-regulated by loss of $Nrf1^{-/-}$, thus leading to a certain yield of ROS from the oxidative metabolism of fatty acids, in

the murine deficient liver cells (7,8). In addition, it is also quite interesting to notice our previous work unraveling

that Nrf1 and Nrf2 are contributable to dual opposing regulation of the PTEN-AKT signaling to be selectively

repressed or activated, respectively, in the pathogenesis of NASH and liver cancer (12). Here, we further validated

that the extended PI3K-AKT-mTOR signaling to the inflammatory marker COX, as well as those key inflammatory

responsive factors along with CD36, was also constitutively activated by loss of $Nrf1\alpha$ in human hepatoma cells.

In this study, we have further provided a revealing insight into the molecular mechanisms of $Nrf1\alpha^{-/-}$-leading

lipid metabolism disorders, along with a cascade of the enzymatic reactions leading to lipid deposition during the

development of NASH and hepatoma. The resulting evidence demonstrates that those key enzymes including FASN,

SCD1 and ACCα and relevant transcriptional factor SREBP1 involved in the de novo biosynthesis of lipids were

significantly up-regulated in $Nrf1\alpha^{-/-}$ cells with hyper-active Nrf2, but they were hence unaffected or even

down-regulated by inactivation of Nrf2 in $Nrf2^{-/-}$ cells. Further experimental evidence unraveled that such

up-regulation of the lipid biosynthesis pathway by $Nrf1\alpha^{-/-}$-deficiency may occur through the JNK-Nrf2-AP1 signaling.

Of note, a few of different experimental results may be determined by distinct functional heterodimers of Nrf2 with

each of sMaf or other bZIP proteins, thus competitively with JUN and FOS against the heterodimeric AP1 factor for

selective physical binding to the consensus ARE sites within their target gene promoter regions (28).

Furthermore, we also found that ATGL and the nuclear receptor PPAR-PGC1 signaling towards the lipid oxidative

decomposition pathway were markedly down-regulated in $Nrf1\alpha^{-/-}$ cells, but they were unaffected or even
conversely up-regulated in $Nrf2^{-/-}$ cells. Such activation of the PPAR-PGC1 signaling was also supported by the
evidence obtained from the tissue-specific $Nrf1^{-/-}$ mice (9,29). Therein, Hirotsu *et al* (9) had shown that liver-specific
Nrf1 deficiency cannot only lead to obviously decreased expression of Lipin1 and PGC-1β, but conversely, the
presence of Nrf1 enabled this CNC-bZIP factor to directly mediate transcriptional activation of Lipin1 and PGC-1β by
binding to the ARE sequences in their promoter regions. Collectively, these indicate that Nrf1 is essential for
regulating critical genes involved in lipid catabolism. Upon loss of $Nrf1\alpha^{-/-}$, constitutive activation of the PPAR-PGC1
signaling is implicated in the severe aberrant accumulation of lipids. The ensuing metabolic pathological process may
be prevented by 2-bromopalmitate, because this chemical had also been shown to serve as a natural ligand of PPARδ
receptor involved in the fatty acid oxidation (22), albeit detailed molecular mechanisms remain elusive.

In summary, this work demonstrates that both Nrf1 and Nrf2 make different or even opposite contributions to
the intracellular lipid metabolism, as shown in the model of Fig. 6. Upon knockout of $Nrf1\alpha$, the de novo biosynthesis
of lipids is incremented by significant induction of the JNK-Nrf2-AP1 signaling pathway in $Nrf1\alpha^{-/-}$cells, while the lipid
oxidative decomposition pathway was down-regulated by the PPAR-PGC1 signaling. $Nrf1\alpha^{-/-}$cell-autonomous
activation of PI3K-AKT-mTOR signaling towards up-regulation of CD36 leads to augmented uptake of fatty acids.
These together result in severe accumulation of lipids as deposited in lipid droplets. By contrast, $Nrf2^{-/-}$ cells gave rise
to obvious decreases in both lipid synthesis and uptake capacity, along with enhanced oxidative metabolism of fatty
acids. Importantly, such $Nrf1\alpha^{-/-}$-leading lipid metabolism disorders and relevant inflammatory responses are
substantially ameliorated by 2-bromopalmitate. Overall, this study provides a potential strategy for prevention of the
inflammatory transformation into cancer and relevant treatment by precision targeting of Nrf1, Nrf2, or both in
translational medicine.

Author contributions: Z.Z. performed the experiments with help of M.W., S.H., and J.F., collected all the relevant data,
made draft of this manuscript with most figures and supplemental information with assistance of. R.D. Lastly, Y.Z. designed
and supervised this study, analyzed all the data, helped to prepare all figures with cartoons, wrote and revised the paper.

Acknowledgments: We are greatly thankful to Drs. Lu Qiu (Zhengzhou University, Henan, China) and Yonggang Ren (North
Sichuan Medical College, Sichuan, China) for their involvement in establishing the indicated cell lines used in this study. We
also thank to all those other present and past members of Prof. Zhang's laboratory (at Chongqing University, China) for
giving critical discussion and invaluable help with this work. Notably, this study was also funded by the National Natural
Science Foundation of China (NSFC, with two project grants 81872336 and 82073079) awarded to Prof. Yiguo Zhang.

Conflicts of Interest: The authors declare no conflict of interest. Besides, it should be noted that the preprinted version of
this paper had been initially posted at *BioRxiv,* doi: https://doi.org/10.1101xxx

References

1.   Cheng, C., Zhuo, S. M., Zhang, B., Zhao, X., Liu, Y., Liao, C. L., Quan, J., Li, Z. Z., Bode, A. M., Cao, Y., and Luo, X. J. (2019)
Treatment implications of natural compounds targeting lipid metabolism in nonalcoholic fatty liver disease, obesity and
cancer. *Int J Biol Sci* **15**, 1654-1663

2.   Alannan, M., Fayyad-Kazan, H., Trezeguet, V., and Merched, A. (2020) Targeting Lipid Metabolism in Liver Cancer.
Biochemistry-Us **59**, 3951-3964

3.   Santos, C. R., and Schulze, A. (2012) Lipid metabolism in cancer. *Febs J* **279**, 2610-2623

4.   Walther, T. C., and Farese, R. V., Jr. (2012) Lipid droplets and cellular lipid metabolism. *Annu Rev Biochem* **81**, 687-714

5.   Zhang, Y., and Xiang, Y. (2016) Molecular and cellular basis for the unique functioning of Nrf1, an indispensable transcription
factor for maintaining cell homoeostasis and organ integrity. *Biochem J* **473**, 961-1000

6.   Parola, M., and Novo, E. (2005) Nrf1 gene expression in the liver: A single gene linking oxidative stress to NAFLD, NASH and
hepatic tumours. *J Hepatol* **43**, 1096-1097

7.   Tsujita, T., Peirce, V., Baird, L., Matsuyama, Y., Takaku, M., Walsh, S. V., Griffin, J. L., Uruno, A., Yamamoto, M., and Hayes, J. D.

(2014) Transcription Factor Nrf1 Negatively Regulates the Cystine/Glutamate Transporter and Lipid-Metabolizing Enzymes.

Mol Cell Biol **34**, 3800-3816

8. Xu, Z. R., Chen, L. Y., Leung, L., Yen, T. S. B., Lee, C., and Chan, J. Y. (2005) Liver-specific inactivation of the Nrf1 gene in adult

mouse leads to nonalcoholic steatohepatitis and hepatic neoplasia. *P Natl Acad Sci USA* **102**, 4120-4125

9. Hirotsu, Y., Hataya, N., Katsuoka, F., and Yamamoto, M. (2012) NF-E2-Related Factor 1 (Nrf1) Serves as a Novel Regulator of

Hepatic Lipid Metabolism through Regulation of the Lipin1 and PGC-1 beta Genes. *Mol Cell Biol* **32**, 2760-2770

10. Huang, J. S., Tabbi-Anneni, I., Gunda, V., and Wang, L. (2010) Transcription factor Nrf2 regulates SHP and lipogenic gene

expression in hepatic lipid metabolism. *Am J Physiol-Gastr L* **299**, G1211-G1221

11. Zhang, L., Dasuri, K., Fernandez-Kim, S. O., Bruce-Keller, A. J., and Keller, J. N. (2016) Adipose-specific ablation of Nrf2

transiently delayed high-fat diet-induced obesity by altering glucose, lipid and energy metabolism of male mice. *Am J Transl*

Res **8**, 5309-+

12. Qiu, L., Wang, M., Hu, S. F., Ru, X. F., Ren, Y. G., Zhang, Z. W., Yu, S. W., and Zhang, Y. G. (2018) Oncogenic Activation of Nrf2,

Though as a Master Antioxidant Transcription Factor, Liberated by Specific Knockout of the Full-Length Nrf1 alpha that Acts as

a Dominant Tumor Repressor. *Cancers* **10**

13. Zhang, Y. G., Lucocq, J. M., and Hayes, J. D. (2009) The Nrf1 CNC/bZIP protein is a nuclear envelope-bound transcription factor

that is activated by t-butyl hydroquinone but not by endoplasmic reticulum stressors. *Biochemical Journal* **418**, 293-310

14. Zhu, Y. P., Zheng, Z., Hu, S., Ru, X., Fan, Z., Qiu, L., and Zhang, Y. (2019) Unification of Opposites between Two Antioxidant

Transcription Factors Nrf1 and Nrf2 in Mediating Distinct Cellular Responses to the Endoplasmic Reticulum Stressor

Tunicamycin. *Antioxidants* **9**

15. Snaebjornsson, M. T., Janaki-Raman, S., and Schulze, A. (2020) Greasing the Wheels of the Cancer Machine: The Role of Lipid

Metabolism in Cancer. *Cell Metab* **31**, 62-76

16. Kronenberg, F., Schoenborn, V., Heid, I. M., Vollmert, C., Lingenhel, A., Adams, T. D., Hopkins, P. N., Illig, T., Zimmermann, R.,

Zechner, R., and Hunt, S. C. (2005) The ATGL gene is associated with free fatty acids, trigylcerides and type 2 diabetes. *Genet*

Epidemiol **29**, 259-259

17. Memon, R. A., Tecott, L. H., Nonogaki, K., Beigneux, A., Moser, A. H., Grunfeld, C., and Feingold, K. R. (2000) Up-regulation of

peroxisome proliferator-activated receptors (PPAR-alpha) and PPAR-gamma messenger ribonucleic acid expression in the liver

in murine obesity: Troglitazone induces expression of PPAR-gamma-responsive adipose tissue-specific genes in the liver of

obese diabetic mice. *Endocrinology* **141**, 4021-4031

18. Miura, S., Kai, Y., Kamei, Y., and Ezaki, O. (2008) Isoform-specific increases in murine skeletal muscle peroxisome

proliferator-activated receptor-gamma coactivator-1 alpha (PGC-1 alpha) mRNA in response to beta 2-adrenergic receptor

activation and exercise. *Endocrinology* **149**, 4527-4533

19. Gao, N., Zhang, Z., Jiang, B. H., and Shi, X. L. (2003) Role of PI3K/AKT/mTOR signaling in the cell cycle progression of human

prostate cancer. *Biochem Bioph Res Co* **310**, 1124-1132

20. Wang, C., Hu, L., Zhao, L., Yang, P., Moorhead, J. F., Varghese, Z., Chen, Y., and Ruan, X. Z. (2014) Inflammatory stress increases

hepatic CD36 translational efficiency via activation of the mTOR signalling pathway. *Plos One* **9**, e103071

21. Cao, D. Y., Luo, J., Chen, D. K., Xu, H. F., Shi, H. P., Jing, X. Q., and Zang, W. J. (2016) CD36 regulates lipopolysaccharide-induced

signaling pathways and mediates the internalization of Escherichia coli in cooperation with TLR4 in goat mammary gland

epithelial cells. *Sci Rep-Uk* **6**

22. Berger, J., and Moller, D. E. (2002) The mechanisms of action of PPARs. *Annu Rev Med* **53**, 409-435

23. Zhu, Y. P., Zheng, Z., Xiang, Y., and Zhang, Y. (2020) Glucose Starvation-Induced Rapid Death of Nrf1alpha-Deficient, but Not

Nrf2-Deficient, Hepatoma Cells Results from Its Fatal Defects in the Redox Metabolism Reprogramming. *Oxid Med Cell Longev*

2020, 4959821

24. Chen, J., Wang, M., Xiang, Y., Ru, X., Ren, Y., Liu, X., Qiu, L., and Zhang, Y. (2020) Nrf1 Is Endowed with a Dominant

Tumor-Repressing Effect onto the Wnt/beta-Catenin-Dependent and Wnt/beta-Catenin-Independent Signaling Networks in

the Human Liver Cancer. *Oxid Med Cell Longev* **2020**, 5138539

25.   Zhang, S., Deng, Y., Xiang, Y., Hu, S., Qiu, L., and Zhang, Y. (2020) Synergism and Antagonism of Two Distinct, but Confused,

Nrf1 Factors in Integral Regulation of the Nuclear-to-Mitochondrial Respiratory and Antioxidant Transcription Networks. *Oxid*

Med Cell Longev **2020**, 5097109

26.   Wufur, R., Fan, Z., Liu, K., and Zhang, Y. (2021) Differential yet integral contributions of Nrf1 and Nrf2 in the human

antioxidant cytoprotective response against tert-butylhydroquinone as a pro-oxidative stressor. *bioRxiv* **2021.06.05.447190**,

doi: https://doi.org/10.1101/2021.1106.1105.447190

27.   Widenmaier, S. B., Snyder, N. A., Nguyen, T. B., Arduini, A., Lee, G. Y., Arruda, A. P., Saksi, J., Bartelt, A., and Hotamisligil, G. S.

(2017) NRF1 Is an ER Membrane Sensor that Is Central to Cholesterol Homeostasis. *Cell* **171**, 1094-1109 e1015

28.   Newman, J. R., and Keating, A. E. (2003) Comprehensive identification of human bZIP interactions with coiled-coil arrays.

Science **300**, 2097-2101

29.   Hou, Y., Liu, Z., Zuo, Z., Gao, T., Fu, J., Wang, H., Xu, Y., Liu, D., Yamamoto, M., Zhu, B., Zhang, Y., Andersen, M. E., Zhang, Q.,

and Pi, J. (2018) Adipocyte-specific deficiency of Nfe2l1 disrupts plasticity of white adipose tissues and metabolic

homeostasis in mice. *Biochem Biophys Res Commun*

Figure 1

Figure 1. Significant discrepancies between Nrf1 and Nrf2 in lipid metabolism are observed in their genotypic knockout
cell lines. (A) Distinct protein abundances of Nrf1 and Nrf2 in three different genotypic cell lines *Nrf1/2$^{+/+}$*, *Nrf1α$^{-/-}$* and
Nrf2$^{-/-}$ were determined by Western blotting with their specific antibodies, respectively. (B) Distinct mRNA expression
levels of *Nrf1* and *Nrf2* were examined by real-time qPCR analysis of the aforementioned cell lines. The results are
calculated as fold changes (mean ± S.D) and shown graphically with significant increases (\$, $P < 0.05$) and significant
decreases (*, $P < 0.05$; **, $P < 0.01$). (C) The triglyceride (TG) contents were measured after those experimental cell lines
were pretreated with 100 μmol/L oleic acid (OA). The results were calculated as a concentration of nmol/mg protein, with
significant increases (\$, $P < 0.01$; \$\$, $P < 0.001$) or significant decreases (*, $P < 0.01$). (D) These cell lines were subjected to
pretreatment with distinct concentrations of oleic acid (OA), followed by the oil red O staining. The resulting images (scale
bar = 25 μm) were then photographed under a microscope. (E) Appropriate sizes of those individual cells that had been
treated as described above, were screened by microscopy and photographed. Scale bar = 10 μm.

Figure 2

Figure 2. Different expression levels of lipid metabolism-related genes mediated by Nrf1 and Nrf2 in distinct genotypic cell lines. (A) Different protein abundances of key genes involved in *de novo* biosynthesis of lipids in *Nrf1/2$^{+/+}$*, *Nrf1α$^{-/-}$* and *Nrf2$^{-/-}$* cell lines were determined by Western blotting. (B) Different mRNA expression levels of the indicated genes were also determined by real-time qPCR analysis of the aforementioned cell lines. The results are calculated as fold changes (mean ± S.D) with significant increases ($, $P < 0.05$; $$, $P < 0.01$) and significant decreases (**,$P < 0.01$).(C,D) Distinct expression levels of key enzymes and transcription factors involved in lipid catabolism were evaluated by Western blotting (C). The intensity of the indicated immunoblots was calculated at fold changes (mean ± S.D) and shown graphically with significant decreases (*, $P < 0.05$; **, $P < 0.01$). (E) Basal protein expression levels of JNK, p-JNK, Nrf2 and JUN were also examined in *Nrf1/2$^{+/+}$*, *Nrf1α$^{-/-}$* and *Nrf2$^{-/-}$* cell lines.

Figure 3

**Figure 3. The JNK-Nrf2-AP1 signaling pathway is required for increased expression of lipid synthesis genes in *Nrf1α*^{-/-}
cells.** (A) *Nrf1α*^{-/-} cells were or were not treated with the JNK-specific inhibitor SP600125 (20 μmol/L) for 24 h and then
subjected to Western blotting of the JNK-Nrf2-AP1 signaling towards lipid biosynthesis pathway. (B, C) *Nrf1α*^{-/-} cells were
transfected with siNC (i.e., small-interfering RNA negative control) or siNrf2 (to specifically interfere with Nrf2), before
being determined by (B)Western blotting of the protein abundances as described above or (C) real-time qPCR analysis of
their mRNA expression levels. The results were calculated as fold changes (mean ± S.D) with significant decreases (*, $P <$
0.01). (D, E) *Nrf1α*^{-/-} cells were transfected with siJUN (to specifically interfere with JUN) or siNC, and subjected to Western
blotting analysis (D) and real-time qPCR analysis (E), as described above. (F) Wild-type *Nrf1/2*^{+/+} cells were allowed for
transient transfection with JUN (2 μg of DNA) alone or plus FOS (i.e., JUN+FOS, every 1 μg of DNA) and then subjected to
visualization by Western blotting analysis of those indicated proteins.

Figure 4

Figure 4. *Nrf1α*[−/−] cell-autonomous up-regulation of CD36 and COX2 through the PI3K-AKT-mTOR signaling pathway. (A) Western blotting analysis of those indicated protein abundances in both *Nrf1/2*[+/+] and *Nrf1α*[−/−] cell lines was carried out as described in the legend for Figure 3. (B, C) These two cell lines were treated with (B) 0.5 μmol/L of the PI3K-specific inhibitor NVP-BKM120 or (C) 200 nmol/L of the mTOR inhibitor rapamycin (RAPA) for 24 h and then determined by Western blotting with those indicated antibodies. (D) The mRNA expression levels of key inflammatory factors and cytokines in *Nrf1/2*[+/+] and *Nrf1α*[−/−] cell lines were measured by real-time qPCR analysis. The results were calculated as fold changes (mean ± S.D) with significant increases ($, $P < 0.01$; $$, $P < 0.001$). (E) Western blotting of some inflammatory factors and cytokines revealed changed levels of their protein expression in *Nrf1/2*[+/+] and *Nrf1α*[−/−] cell lines.

Figure 5

Figure 5. Accumulation of lipids and ROS along with the inflammatory response in *Nrf1α^-/-* cells are substantially alleviated by 2-bromopalmitate. (A) The schematic shows two molecular structures of palmitic acid (PA) and 2-bromopalmitate (2BP). (B) The triglyceride contents were measured following treatment of both *Nrf1/2^+/+* and *Nrf1α^-/-* cell lines with 50 µmol/L 2BP. Thereafter, such 2BP-treated cell lines were subjected, respectively, to (C) Western blotting analysis of CD36, IL-6 and p-NFκB; (D) visualization by the Oil red O staining as shown in the photographed images (scale bar = 10 µm); (E) real-time qPCR analysis of IL-6, NFκB1, NFκB2 and TNFα at their mRNA expression levels, the resulting data of which were calculated as fold changes (mean ± S.D) with a significant increase ($, *P* < 0.01, relative to the wild-type controls), and another significant decrease (*, *P* < 0.01, when compared to those measured from the untreated cells); and (F) flow cytometry determination of ROS as described in the section of "Materials and Methods".

Figure 6

Figure 6. A model proposed for the molecular mechanisms underlying the inflammatory accumulation of lipids and ROS
in *Nrf1α*$^{-/-}$ cells. This cartoon provides a better explanation of molecular mechanisms by which the inflammatory
accumulation of lipids and ROS occurs in *Nrf1α*$^{-/-}$ cells. Of note, *Nrf1α*$^{-/-}$ cell-autonomous induction of the PI3K-AKT-mTOR
signaling to up-regulation of CD36 results in augmented uptake of fatty acids. The *de novo* biosynthesis of lipids is
up-regulated by the JNK-Nrf2-AP1 signaling to transactivation of their target genes encoding key metabolic rate-limiting
enzymes and the relevant transcription factors, while the lipid oxidative decomposition pathway is down-regulated by the
PPAR-PGC1 signaling. These together result in the inflammatory accumulation of lipids and ROS in *Nrf1α*$^{-/-}$ cells, but this
pathological process is alleviated substantially by 2-bromopalmitate (2BP) to down-regulate the CD36 expression. In
addition, the yield of ROS in *Nrf1α*$^{-/-}$ cells, was discussed in the relevant main text.