Characterizing Genetic Mechanisms Involved in Measuring Day-length in *Drosophila*

melanogaster

Harjit Khaira[1] and Kwangwon Lee[1,2*]

[1] Department of Biology, Rutgers, The State University of New Jersey, Camden, New Jersey,

United States of America

[2] Center for Computational & Integrative Biology, Rutgers, The State University of New Jersey,

Camden, New Jersey, United States of America

Short Title: *PKCdelta* on Photoperiodism

[*] Corresponding author

E-mail: kwangwon.lee@rutgers.edu

Abstract

Many organisms are known to regulate seasonal behaviors and physiological processes in response

to day length changes through photoperiodism. Extreme changes in photoperiods have detrimental

effects on human health, which can impair development and serve as the origin of adult diseases.

Since the seminal work by Bünning in 1936, there are studies supporting the view that organisms

can measure the day length through an endogenous 24-hour cellular circadian clock. However, the

mechanisms involved in measuring seasonal or day-length changes are not understood. In the

current study, we performed a genome-wide association study (GWAS) on photoperiodism using

the *Drosophila* Genetic Reference Panel. The GWAS identified 32 candidate genes responsible

for photoperiodic regulations. The knockout mutants of the top four candidate genes (*Protein*

Kinase C delta (Pkcdelta), Glucuronyltransferase-P (*GlcAT-P*), *Brain-specific homeobox* (*Bsh*),

and *Diuretic hormone 31 Receptor (Dh31-R1))* were analyzed for their photoperiod and circadian

period phenotypes. *PKCdelta* and *GlcAT-P* mutants show a significantly different photoperiod

response compared to that of the wild type strain, and also had an altered circadian period

phenotype. Further molecular characterization revealed that the mutant two independent mutant

alleles of *PKCdelta* with a defective catalytic domain had distinct photoperiod responses. Taken

these data together, we concluded that there is overlap between the circadian clock and

photoperiodic regulations in *Drosophila*, and *PKCdelta* is a component that is involved in both

circadian and photoperiodic regulations. By identifying novel molecular components of

photoperiod, the current study provides new insights into the genetic mechanisms of determining

the seasonal changes.

Keywords Photoperiodism; Circadian Clock; GWAS; *Drosophila*

Introduction

Most organisms experience predictable environmental changes; daily, tidal, lunar, and seasonal.

These changes mean different availability of resources and stressors at predictable time intervals.

It is crucial for an organism to prepare for the incoming season for its survival in nature. Thus, it

is not surprising that organisms have evolved time-measuring cellular devices. Among the known

biological rhythms, seasonal rhythm plays a major role in many biological processes such as

growth, development, reproduction, migration, and dormancy. A robust response to these rhythms

can increase the survival success of an organism (Bradshaw and Holzapfel, 2010). However, little

is known about the molecular and genetic mechanism of how an organism interprets and adapts to

seasonal environmental changes.

Photoperiodism is the physiological response of an organism to changes in day length.

Photoperiod is one of the most important factors that control seasonal breeding. Few species show

extreme behaviors like hibernation while many species synchronize breeding and birth of their

offspring to the most advantageous time of the year for better survival. Based on the gestation

period, photoperiodic species can be categorized as long day (LD) or short day (SD) breeders.

Animals like sheep and goats with long gestation periods begin to breed as day length decreases,

while animals like mice and hamsters breed as day length increases (Ortavant et al., 1964). In

humans there are some psychological and psychiatric consequences in response to the

photoperiodic changes. There also is seasonality in birth rates among humans and onset of certain

diseases like measles in the population (Martinez-Bakker et al., 2014). Patients with bipolar

disorders have more depressive episodes during the late autumn and early winter, while there is an

increase of manic episodes during spring and summer (Maruani et al., 2018). In flowering plants,

some are flowering at specific season of the year, and thus grouped as SD or LD flowering plants.

The molecular and genetic pathways of photoperiodism have been studied in plants (Johansson

and Staiger, 2015). It is also known that the circadian clock controls the activity of CONSTANS

(CO), which is a key component of the photoperiod pathway. CO activates the transcription

of *FLOWERING LOCUS T* (*FT*) which the ultimately leads to flowering in plants (Song et al.,

2010).

Photoperiodic responses are well documented in insects (Bradshaw and Holzapfel, 2010).

Although *Drosophila melanogaster* is the successful model species for many biological processes,

the species was regarded as less optimal for studying photoperiodism due do a less pronounced

photoperiod response (Tauber and Kyriacou, 2001). Most of the studies of photoperiodism in

insects utilized the diapause as a phenotype. Diapause occurs when female insect ovaries shrink

due to photoperiod changes as an indicator of photoperiodic phenotype. On the contrary to the

concern in using *D. meanogaster* for studying photoperiodism, it also has been shown that *D.*

melanogaster has a robust photoperiod response under semi-natural or SD conditions (Nagy et al.,

2018; Saunders et al., 1989).

One compelling hypothesis regarding the seasonal rhythm is that an organism can

'interpret' the season by measuring the day length. A crucial assumption is that an organism knows

the season by measuring photoperiods is due to the presence of an endogenous cellular daily clock,

called the circadian clock. Circadian clocks are endogenous timekeeping networks that drive 24hr

circadian rhythms. Bünning reported that the rate of leaf movement speed changes during different

seasons and proposed the circadian clock controlled this via photoperiodism (Bünning, 1936).

Since then, a number of studies supported Bunning's hypothesis in certain organisms while

rejecting his hypothesis in others. Studies on aphids and spider mites conclude that their

photoperiod mechanisms do not seem to depend on endogenous 24-h rhythms (Lees, 1960).

Mutation of the classical clock gene *period* (*per)* still allowed *D. melanogaster* to distinguish

between SD and LD . It was concluded that the clock gene *per* cannot be involved in night-length

measurement mechanism (Saunders, 1990). However, another study in *Drosophila* suggests that

the circadian clock is necessary for a photoperiodic response, as arrhythmic clock mutants *per^{01}*,

tim^{01}, and *Clk^{JRK}* have altered photoperiodic responses from wild-type *Drosophila* (Pegoraro et al.,

2014). For the last few decades, the genetic and molecular mechanisms of circadian clocks have

been characterized in many organisms (Partch et al.2014, Nohales and Kay 2016). However, it is

still not well understood on how the circadian clock is involved in measuring photoperiodism or

day length changes.

There are at least two different hypotheses on how circadian clocks are involved in

measuring seasonal changes: the external coincidence model and the internal coincidence model.

In the external coincidence model, the photoperiodic response can be generated by the circadian

clock that is sensitive to light at certain phases (Sawa et al., 2008). If the light signal is detected at

this sensitive phase, then the photoperiodic response will be displayed such as flowering in plants.

Light can entrain the clock and set the phase of photoperiodic response like flowering, but light

can also directly induce flowering. In the internal coincidence model, two (or more) rhythms are

produced by the clock. Under the internal coincidence model, responses, such as flowering, are

only generated in conditions that bring both rhythms in phase with each other (Sawa et al., 2008).

Various studies show one or the other hypotheses at work in a range of organisms. In *Arabidopsis*,

external coincidence models can explain long day-specific dusk *FT* (a florigen gene) expression

(Song et al., 2018). Studies using the parasitic wasp *Nasonia vitripennis* support the internal

coincidence model. Using the Nanda-Hamner protocol it was found that *N.vitripennis* exhibit two

peaks in diapause, a dawn and dusk peak, which is suggestive of two circadian oscillators

(Saunders, 2012). In some bacterial species the hourglass model is suggested to be responsible for

timekeeping (Johnson et al., 2017). When bacteria exposed to more regular environmental

conditions such as consistent light-dark cycles or constant photoperiod, hourglass model is at play.

However, when the environmental conditions vary such as inconsistent light-dark cycles or

photoperiods leads to a sustained oscillator (Johnson et al., 2017).

In our current study, we aim to better understand the genetic mechanisms of

photoperiodism and the roles of circadian clocks in photoperiodism using the eukaryotic model

organism *D. melanogaster*. *Drosophila* Genetic Reference Panel (DGRP) is a community resource

of 205 inbred lines created from 40 wild type parents for studying population genomics and

quantitative traits (Mackay et al., 2012). Because the DGRP is inbred to homozygosity, it is

possible to identify SNP variants responsible for the trait of interest. DGRP is a fully sequenced

population with annotated high-quality SNPs, indels, and chromosomal inversions (Huang et al.,

2014; Mackay et al., 2012). DGRP has been used to study various complex traits such as starvation

resistance, startle response, chill coma recovery (CCR) time, and circadian behavior (Mackay et

al., 2012; Harbison et al., 2019).

Although diapause has been the phenotype of choice in characterizing the photoperiodic

response, the diapause assay is challenging to perform in a high throughput manner. Therefore, we

optimized the CCR assay as a phenotype for an organism's ability to respond to the photoperiodic

changes to perform the initial DGRP Genome-Wide Association Study (GWAS). We propose that

our approach would identify genes involved in both photoperiodism and the circadian clock or

only in photoperiodism. Our current study identified 32 candidate photoperiod genes from which

we assessed the photoperiod response of the top 4 genes using CCR. We further characterized the

role of allelic mutants of *PKCdelta* in photoperiodism using CCR, diapause, and molecular

techniques.

Materials and Methods

Fly stocks

We purchased the DGRP strains from the Bloomington Drosophila Stock Center (Indiana

University, Bloomington, IN). In the laboratory, the fly stock was maintained at room temperature

(25°C) and in 12L:12D photoperiod conditions. The following are the *PKCdelta* mutants

BDSC#18258 Pbac[RB]PKCdelta[e04408], BDSC#59433 Mi[MIC]PKCdelta[MI13377]. The

BDSC#62582 is *GlcAT-P* knockout mutant Pbac[GlcAT-P]0027-G4, BDSC#27467 Bsh mutant

EPbsh[G3075]. BDSC#22718 is the *Dh31-R1* knockout mutant ET1Dh31-R1[MB00175].

Photoperiod conditions and Chill coma recovery

Chill Coma Recovery (CCR) is a traditional method used to assess the photoperiod phenotype in

Drosophila species. CCR is expressed as the time (in seconds) it takes an organism to recover from

a paralysis like state after being kept under extremely low temperature. *D. melanogaster* enter the

chill coma at 7.0 ± 0.9 °C (Hosler et al., 2000). It has been clearly demonstrated that flies raised

under short-day (SD) photoperiod conditions compared to long-day (LD) photoperiod conditions

have an advanced (or shorter) recovery time (Lanciani et al., 1992). DGRP fly lines were reared

under LD and SD in a chamber, 25 ± 0.5 °C and 70% humidity. The LD consisted of 16 hours of

light and 8 hours of dark (16h:8h), while the SD consisted of 8 hours of light and 16 hours of dark

(8h:16h). The flies were reared under these conditions for two weeks. Chill coma was then

performed by placing the flies in 4 °C for 60 minutes. Next, the flies were placed into 5 mm glass

tubes and into the *Drosophila* activity monitors (DAM2, Trikinetics) to measure the time it takes

for the flies to recover. For an accurate measure of the recovery time, the flies were placed in the

middle of the glass tube near the IR sensor to capture the exact time the fly recovered from the

coma.

Diapause Assay

The diapause assay was performed by surgically removing the ovaries after rearing female flies

under different photoperiods. Previous research suggested that female *Drosophila* placed under

SD enter reproductive diapause which results in underdeveloped ovaries. The SD chamber

condition mimics the winter-like days when most organisms cease to reproduce to conserve

energy, therefore, diapause is induced in female flies (Saunders et al., 1989). Female *Drosophila*

placed under LD condition do not enter diapause. The *PKCdelta* mutants, *PKCdelta*:Pbac and

PKCdelta:MIC were reared under SD and LD photoperiodic conditions along with DGRP208

(WT) strain as a control. Newly hatched female flies were placed under the photoperiod conditions

for 12 days, 25˚ C. Overies were dissected in 1x PBS, the image was taken with a light microscope

(LEICA microsystems, LEICA EC3, Model:MEB126) under 4x magnification.

qPCR

The expression level of the candidate gene *PKCdelta* was characterized under SD and LD. The

RNA was prepared using the Zymo Research Direct-Zol RNA extraction kit (Cat#R2050). The

cDNA was prepared using the New England Biolabs (NEB) protoscript II first strand cDNA

synthesis kit (Cat#E65605). For qPCR the Applied Biosytems PowerUp SYBER Green master

mix was used (Cat#A25742) in the Applied Biosystems QuantStudio 6 Rlex Real-Time PCR

system according to the manufacture's manual. The following primer set was used; ACTIN served

as a control, (Fw; GCAAGTACTCTGTCTGGATC) (Rv; CCAGCAGAATCAAGACCATCC).

HK-5 primer set which targeted the RD transcript of *PKCdelta* (Fw;

CGATCGATTGGGGTTTGCTGG) (Rv:GTGATGTGAGGATTTGTGTAGG), expected

product size 283 bp. HK6 primer set targeted the RE transcript of *PKCdelta* (Fw;

TCGAGCTGAAGACAACACCT) (Rv: CTTTCTTAGCTTCGACTTGGTT) expected to product

size 160bp (Figure S4, Table S2).

Statistical Analysis

Cohen's D

CCR has been used successfully by other groups as the phenotype reflecting photoperiodic

response (Pegoraro et al., 2014). Our primary interest is measuring the 'differential' response of

each genotype to two different photoperiodic conditions. Cohen's D (CD) is used to measure the

effect size of the photoperiodic response in each genotype (Lakens, 2013):

199
$$CD = \frac{(CCR_{SD} - CCR_{LD})}{Average(STDV_{CCR_{SD}}, STDEV_{CCR_{LD}})}$$

For each of the DGRP strains, the average CCR time was calculated from 20 replicates per

strain in SD (CCR$_{SD}$) and LD (CCR$_{LD}$). The CD value of each genotype was calculated by

subtracting CCR$_{LD}$ from CCR$_{SD}$, and divided by the average of the two standard deviations of

CCR$_{LD}$ and CCR$_{SD}$. The CD values can be divided into three groups: large effect size, CD > 0.8;

medium effect size, 0.8 > CD > 0.5; and small effect size, CD < 0.2. The CD values for the mutants

were obtained by subtracting equinox photoperiod CCR value from that of SD or that of LD

respectively. This was done to ensure that the data was normalized to the EQ photoperiod.

$$CD_{SD} = \frac{(CCR_{SD} - CCR_{EQ})}{Average(STDV_{CCR_{SD\&EQ}})} \quad , CD_{LD} = \frac{(CCR_{LD} - CCR_{EQ})}{Average(STDEV_{CCR_{LD\&EQ}})}$$

CD value provides directionality by positive or negative values. Thus, CD value shows if a

photoperiodic condition provided a positive effect on the strain or a negative effect in the CCR

assay.

Genotype Effect Size

To measure the difference between the effect size of CD in different genotypes, we calculated the

Genotype Effect Size (GES) by subtracting the CD value of the wild type from that of the mutant.

$$CD_{GenotypeX} - CD_{Wild\ type} = Genetic\ Effect\ Size\ (GES)$$

In calculating the GES values, DGRP208 (WT) was used as the baseline. GES value allowed us

to determine which candidate gene had the most differential response to the photoperiod in

comparison to that of the wild type.

GWAS Analysis

After calculating CD for all the strains under different photoperiodic conditions, we performed the

GWAS using the web-based tool provided by DGRP2 (Mackay et al., 2012; Huang et al., 2014).

The 1,881,979 select SNPs were analyzed for their association with the photoperiod response. All

the variants from the GWAS were graphed in a manhattan plot with their chromosome on the X-

axis and -log10(p) of the p values on the Y-axis. There are two significant thresholds -log10(5e-5)

and -log10(5e-8) (Figure 2). The SNPs above the -log10(5e-8) were considered significant SNPs.

Heritability

The HiLMM R package was used to calculate narrow-sense heritability of photoperiod response

in the DGRP population. The HiLMM package provides an estimation of heritability with

confidence intervals in linear mixed models.

Results

Variation in CCR in the DGRP

Figure 1. Variation in chill coma recovery in the DGRP population. A. The overall distribution of the CD values within the DGRP population, Shapiro-Wilks normality test $p = 3.763e-09$, $p < 0.05$, and $W = 0.9194$. B. The Cohen's D (CD) of the DGRP population, CD = (CCR_{SD} - CCR_{LD})/average ($STDV\_CCR_{SD}$, $STDV\_CCR_{LD}$). CD > 0.8 is large effect size, CD of 0.5 is medium effect size, and CD < 0.2 is small effect size. The majority of the DGRP strains fall under medium effect size total of 93 strains, while 63 strains fall under large effect size and 49 under small effect size. Thick horizontal lines represent mean and thin lines represent standard deviation. C. The CCR of the two representative strains from the DGRP population, DGRP28140 and DGRP28152 under Short Day (SD, circle) and Long Day (LD, square) photoperiods.

236
To test our hypothesis that genetic variations exist in photoperiodic response in *D. melanogaster*,

we analyzed the CCR phenotype in DGRP, and found significant variations in CCR among DGRP

strains (Figure 1, S1). The distribution of Cohen's D values in DGRP, Shapiro-Wilks p = 3.763e-

09, and W = 0.9194 (Figure 1A). To determine the differential response of each DGRP genotype

to the two different photoperiods, CD values were calculated for each genotype (Materials and

Methods). The majority of the DGRP strains fall under medium effect size (n = 93), large effect

size (n = 63), and small effect size (n= 49) (Figure 1B). There is more variation in CD values

among the large effect size DGRP strains than the strains that fall under medium and small effect

size. The overall average CCR in LD is 941 sec while CCR in SD is 1095 sec in the whole DGRP

population. Two representative strains are shown along with their recovery time under SD and LD

(Figure 1C). DGRP28140 has a delayed recovery time under SD in comparison to that of LD,

while DGRP28152 has a faster recovery time in SD compared to LD. To determine how much of

the variation is due to genetic factors in the DGRP population, we calculated the heritability.

Heritability (h^2) for the long day condition was 0.51 and h^2 for the short-day condition was 0.45.

It was previously reported that there is a sex difference in the CCR of *Drosophila* (Pegoraro et al.,

2014). However, there was no strong correlation between sex and CCR in our experimental

condition. The data from both sexes were grouped together for our GWAS analysis (Figure S2).

GWAS Analysis for Photoperiodism

Figure 2. Manhattan plot for SNPs in the GWAS analysis. Each point represents a single SNP, and the height of the SNP represent the strength of association with photoperiod (CCR) response expressed as –log10(p). The horizontal lines represent the genome-wide Bonferroni significance threshold (blue line p = -log10(5e-5), red line p = -log10(5e-8). The variant rs1037063 is associated with GlcAT-P, rs554277 is associated with Dh31-R1, rs1753776 is associated with PKCdelta, rs277132 and rs1244560 are not associated with any particular gene since they are in the intergenic regions.

We performed GWAS analysis on CCR using it as a readout for photoperiodic response (Materials

and methods) and identified genetic variants that were significant, p = 1E-07 (Figure 2). We also

identified the 32 most promising variants strongly associated with the photoperiod response (Table

S1). The enrichment analysis of these candidate genes shows a list of biological phenotypes

including jumping, flight phenotypes, and circadian rhythms (Figure S3).

We characterized the top four candidate genes that are strongly associated with photoperiod

CCR responses, p ≥1E-07; *Glucuronyltransferase-P* (*GlcAT-P*); p = 6.63E-08, *Brain-specific*

homeobox (*Bsh*); p = 8.46E-07, *Protein Kinase C delta (Pkcdelta)*; p = 8.22E-07, and *Diuretic*

hormone 31 Receptor (Dh31-R1); p = 3.49E-07. *GlcAT-P* belongs to the glycosyltransferase

family which are involved in catalyzing the conjugation of glucuronic acid from UDP-glucuronate

to an acceptor. This particular gene is involved in the biosynthesis of L2/HNK-1 carbohydrate

epitopes on both glycolipids and glycoproteins (Kim et al., 2003). *Bsh* encodes a putative

transcription factor, and it is needed for the specification of neural identities in the adult brain. The

rat ortholog of *Bsh* (*Bsx)* is suggested to be under circadian regulations and is expressed in the

mature pineal gland of rat (Carstensen et al., 2019). *Dh31-R1* is *Diuretic hormone 31 Receptor*

which encodes a protein that exhibits diuretic hormone receptor activity, and it is involved in G

protein-coupled receptor signaling pathways. *Dh31-R1* is also known to be involved in time

measurement and activity rhythms through the circadian clock system (Dubowy and Cavanaugh,

2014). *PKCdelta* belongs to a family PKC consisting of serine/threonine kinases which are

activated by extracellular signaling. PKCs are divided into subgroups of Classical PKCs, Novel

PKCs, and Atypical PKCs. *PKCdelta* belongs to the class of Novel PKCs which happen to be

known for their two C1 domain and activation via 1,2-diacylglycerol (DAG) and phosphatidyl-

serine (PS). While the cPKC family requires DAG and PS and aPKC only requires PS as a cofactor

for activation (Shieh et al., 2002; Suh et al., 2003). We identified two independent SNP variants

in *PKCdelta*, which strongly suggest that the alleles of *PKCdelta* is associated with the variation

of CCR in the DGRP population.

Photoperiodic responses of the knockouts of the candidate genes

Figure 3. The photoperiod response and clock period phenotype of the top four candidate knockout strains A. Cohen's D values of the knockout mutants of the four candidate genes under LD and SD photoperiods compared to the wild type DGRP 208 strain (WT). The CD is normalized to equinox (EQ) photoperiod (Materials and Methods). The WT, *Dh31-R1* and *Bsh* have negative CD in LD (open) and SD (filled), while *PKCdelta* mutant *PKCdelta:*Pbac and *GlcAT-P* have positive CD under both conditions. B. The Genetic Effect Size of the candidate gene mutants, *PKCdelta:*Pbac and *GlcAT-P* mutants show a different photoperiod response compared to the WT under LD (open) and SD (filled). C. The circadian period of the candidate gene mutants compare to the WT with the period of 24 hr. *GlcAT-P* has a significantly longer period compared to WT (t-test: $p < 0.05$, $p = 1.16E-13$) and *PKCdelta:*MIC mutant of the *PKCdelta* has longer period compared to the WT (t-test: $p < 0.05$, $p = 1.28E-3$). Error bars represents standard deviation. N > 20.

To further characterize the top 4 candidate genes, first, we analyzed CCR response of the knockout

strains; *PKCdelta* mutant PKCδ^{e04408} (BDSC# 18258), *GlcAT-P* mutant $GLCAT-P^{0027-G4}$

(BDSC#62582), *Bsh* mutant Bsh^{G3075} (BDSC#27467) and *Dh31-R1* mutant $Dh31-R^{MB00175}$

(BDSC#22718). As a control, one DGRP parent strain DGRP208 (WT) was used to compare

between the Cohen's D values. For this analysis equinox (EQ) condition, a 12:12 hr light:dark

cycle was used along with SD and LD photoperiods. The CD was obtained by subtracting (or

normalizing) the EQ photoperiod CCR from SD & LD CCR. The WT strain has a negative CD

value of -0.69 (Figure 3A). The *Dh31-R1* mutant has a similar photoperiod response to the WT

with negative CD values in both SD and LD photoperiod. *Bsh* mutant also has negative CD value

though the effect is not as exaggerated. Both *GlcAT-P* and *PKCdelta;Pbac* mutants has positive

CD values (Figure 3A). To determine how different these mutants' photoperiod responses were

from the WT control, the CD value of the WT was subtracted from the CD values of the mutant,

the final value is termed Genetic Effect Size (GES) (Figure 3B, Materials and Methods). *Bsh*,

GlcAT-P and *PKCdelta*:Pbac have different photoperiod responses compared to the WT with

PKCdelta being the most different. The GES of *PKCdelta*:Pbac is not significantly different

between LD and SD. However, *GlcAT-P, Bsh,* and *Dh31-R1* have a difference in GES between

LD and SD (Figure 3B).

Next, we tested the hypothesis that there are shared genetic regulators between the

circadian clock and photoperiodism. We examined the period of the candidate photoperiod knock

out genes using WT as a control (Figure 3C). Compared to the WT which had a period of 24 hr,

GlcAT-P and *PKCdelta*:MIC mutants have a significantly longer period (t-test: $p < 0.001$, $p<0.01$).

We concluded that *GlcAT-P* and *PKCdelta*:MIC genes play a role in both circadian clock

regulation and photoperiodism. We decided to investigate *PKCdelta* further for three reasons. The

two *PKCdelta* mutant alleles showed a significantly different photoperiodic CCR response from

that of the WT (Figure 3B), one of the allelic mutants of *PKCdelta* had an altered circadian period

(Figure 3C), and GWAS analysis showed that there are two independent SNP variants associated

with the photoperiodic CCR response for this gene (Figure 2 and File S3).

PKCdelta Candidate Gene

PKCdelta is a kinase known to be involved in cellular signaling pathways like cell growth,

apoptosis, tumor promotion and carcinogenesis (Suh et al., 2003). We examined two parental

DGRP strains one with a large positive CD value (DGRP379) and another with a large negative

CD value (DGRP391). *PKCdelta* transcribes three transcripts, RC, RD, and RE (Figure S4). The

transcripts RC and RD are similar to each other except with a slight difference in the UTR region;

the RD isoform has a shorter UTR region than the RC isoform. The RC and RD transcript have

regulatory domain which contain C2, C1 and pseudosubstrate domains toward the N-terminal.

Toward the C-terminal there is a catalytic kinase domain which is shared with all three isoforms

(Figure S5). There are two different transcription start sites, one that starts the beginning of

isoforms RD and RC, and the other starts at the RE isoform. The RE isoform is a nested gene with

its transcription start site residing in one of the introns of RC and RD transcripts (Kumar, 2009).

We found that there are two non-synonymous SNPs in the exon 6 which resides in the RE

transcript in the DGRP379 strain but not in DGRP391. The first SNP position is at 12445745:

Guanine to Adenosine, which changes Alanine to Threonine. The second SNP position is at

12446453: Guanine to Cytosine, which changes Alanine to Proline. The first SNP position is in a

ployQ region, which is conserved in the *Drosophila* species. The second SNP causes a repeat of

the sequence surrounding the region (Figure S6A, S6B). We found that RE isoform is a nested

gene which is only conserved in the *Drosophila* genus (Figure S7).

Diapause and CCR in PKCdelta mutant alleles

Figure 4. The photoperiod and period response of *PKCdelta*:MIC and *PKCdelta*:Pbac mutants compared to WT. A. The diapause response of candidate gene *PKCdelta* mutants compared to WT in LD (open) vs SD (filled). There is a significant difference in *PKCdelta*:Pbac mutant diapause response, compared to WT in both photoperiods (Z-test: p= 0.0014 (LD), 0.010 (SD)). *PKCdelta*:MIC diapause response compared to WT in each photoperiod is not significantly different (Z-test: p = 0.27 (LD), 0.30 (SD)). However, there is a significant difference in *PKCdelta*:Pbac LD and SD diapause response (Z-test p = 0.002) and in WT (Z-test: p = 0.007),

but not in *PKCdelta*:MIC (Z-test: p = 0.29). B. Cohen's D of *PKCdelta*;Pbac and *PKCdelta*;MIC compared to WT, the CD values were obtained by subtracting EQ CCR response from LD (open) and SD (filled) photoperiod response. WT has a negative CD, while *PKCdelta*:MIC and *PKCdelta*:Pbac are positive. C. GES of the mutants compared to the WT, both mutant alleles show significantly different photoperiod response from that of the WT, LD (open), SD (filled). D. Circadian period of the *PKCdelta* mutants compared to that of WT separated by homozygous females (open) and hemizygous males (filled). Error bars represent SD, 10 < n < 20, homozygous vs hemizygous (t-test, p > 0.05).

To further explore the role of *PKCdelta* in photoperiodism, we performed the diapause assay using

the *PKCdelta*:Pbac and *PKCdelta*:MIC allelic mutants under different photoperiodic conditions.

The *PKCdelta*:Pbac mutant has an insertion of a *piggy bac* element in an intron close to the 3' end

of the gene. The *PKCdelta:*MIC mutant has an insertion in the 5' end of the gene and contains a

minos-based construct which was created by the Gene Disruption Project (GDP) (Venken et al.,

2011) (Table S4). We found that both mutant alleles have a mutation in the catalytic domain of the

enzyme *PKCdelta* (Figure S9, Table S3 and S4)

Using WT as a control, the diapause assay was carried out under SD and LD conditions

(Figure 4A). As expected, the WT has increased diapause incidences under SD conditions

compared to the LD condition (Z-test, p < 0.05). There was a significant difference in

PKCdelta:Pbac mutant's diapause response compared to that of WT in both photoperiods (Z-test,

p = 0.0014 (LD), p = 0.01 (SD)). The *PKCdelta*:MIC mutant's diapause response compared to WT

in each photoperiod is not significantly different (Z-test, p = 0.27 (LD), p = 0.30 (SD)). However,

there is a significant difference in *PKCdelta*:Pbac LD and SD diapause response (Z-test, p =

0.002), and in WT (Z-test, p = 0.007), but not in *PKCdelta*:MIC (Z-test, p = 0.29) (Figure 4A).

The diapause response in *PKCdelta*:MIC mutant is not statistically significant from that of the

WT. We interpreted this as the mutant allele *PKCdelta*:MIC cannot distinguish SD from LD

because the diapause response between photoperiods is not significantly different (Z-test: p =

0.29).

To determine the necessity of *PKCdelta* for the photoperiodic CCR response, we

preformed the CCR assay using the allelic mutants of *PKCdelta* (Figure 4B). The CD values of

PKCdelta:Pbac and *PKCdelta*:MIC are positive compared to WT. However, *PKCdelta*:MIC have

lower CD values from that of *PKCdelta*:Pbac. The GES of the mutants compared to the WT show

that *PKCdelta*:Pbac have significantly different photoperiod response (Figure 4C).

Since *PKCdelta* is located on the X-chromosome, both of the *PKCdelta* mutants contain

hemizygous males and homozygous females. The CCR was performed to determine whether there

will be a difference in photoperiod response between hemizygous and homozygous flies under SD

and LD photoperiod. Both *PKCdelta*:MIC and *PKCdelta*:Pbac mutants show a difference in

photoperiod response between hemizygous and homozygous flies (Figure S8). To determine if

there was a difference in period phenotype between hemizygous and homozygous flies in the

PKCdelta:Pbac and MIC mutants, we performed the period assay. We found no significant

difference between the period phenotype between hemizygous and homozygous flies (Figure 4D).

WT flies have an expected period of 24 hr with no difference between hemizygous and

homozygous flies (t-test, p > 0.05). *PKCdelta*:MIC hemizygous flies have a period of 30 hr and

homozygous flies have a period of 29 hr but there was no statistically significant difference

between hemizygous and homozygous flies (t-test, p > 0.05). There was no significant difference

between hemizygous and homozygous flies of *PKCdelta*:Pbac in the circadian period (t-test, p >

0.05). Based on these data together, we concluded that MIC mutant allele affected the ability to

distinguish between LD from SD, and also altered the circadian period.

Next, we predicted that *PKCdelta*:Pbac has knockdown of all three transcripts of

PKCdelta, while *PKCdelta*:MIC has knockdown of RC & RD based on the reported locations of

the insertion. The RE isoform is believed to be mainly responsible for photoperiod response

because the *PKCdelta*:MIC mutant shows a decrease in the photoperiod response in the CCR and

diapause assays. We also predicted that the transcripts RC and RD could play a role in interpreting

photoperiod response, though they might not be major players in interpreting photoperiods. These

predictions were then confirmed by performing a qPCR experiment using the *PKCdelta*:MIC and

PKCdelta:Pbac mutants under SD and LD condition (Fig.5).

Expression of PKCdelta

Figure 5. Expression level of *PKCdelta* allelic mutants under LD (open) and SD (filled) compared to DGRP 208 WT. In the WT flies there is no difference in expression of RD and RE transcripts under LD and SD (n=3-5, t-test, p > 0.05). In *PKCdelta*:MIC the expression of RD

is significantly reduced compared to WT (n=3-5, t-test, p<0.001). However, there is a reduction of RE transcript under LD compared to SD in *PKCdelta*:MIC (n=3-5, t-test, p < 0.001). In the *PKCdelta*:Pbac background, the expression of RD and RE is overall reduced compared to WT (n=3-5, t-test, p<0.05). There was no statistical difference in the expression levels under LD and SD photoperiods (n=3-5, t-test, p > 0.05).

We measured the expression levels of *PKCdelta* under different photoperiod conditions (Materials

and Methods). WT flies under LD and SD photoperiods did not show statistically significant

change in *PKCdelta* RD and RE transcript levels (t-test, p > 0.05). *PKCdelta*:MIC which was

predicted to have reduced expression of RD and normal expression of RE, show overall decrease

in expression of RD in both LD and SD (t-test, p > 0.05). However, there is an alteration in the

expression levels of the RE between the photoperiod conditions in *PKCdelta*:MIC. The

*PKCdelta:*MIC mutant under LD show a significant decrease in its expression level from that of

SD (t-test, p < 0.001). *PKCdelta*:Pac mutant show an overall decrease in RD and RE expression

from that of WT (t-test p < 0.05). Though there is no significant difference in the expression levels

of RD and RE between the photoperiods in *PKCdelta*:Pac mutant, (t-test p > 0.05). This suggests

that there is an alteration in the expression of the RE transcript in the *PKCdelta:*MIC mutant due

to long day photoperiod condition. Based on this data and the diapause data, we concluded that

RD transcript could be involved in distinguishing between SD and LD. Since *PKCdelta*:MIC

which has reduced expression of RD transcript show no difference in the diapause response

between photoperiods (Figure 4A & 5). Alternatively, the *PKCdelta*:Pbac mutant which has

reduced expression of both transcripts shows a greater proportion of diapause response compared

to WT in both LD and SD suggesting that expression *PKCdelta* and diapause response is

negatively correlated in this mutant.

Next, we reasoned that the GWAS analysis of the DGRP CCR response itself as a trait

under LD and SD conditions will lead us identifying the downstream genes involved in CCR. We

performed the GWAS analysis on the CCR trait as a phenotype rather than the photoperiodic

response (CD values) as a trait discussed earlier (Files S4 - S7). There were six genes that were

highly associated with the CCR phenotypes both in SD and LD conditions: *Sif, Comm2, Bru-3,*

Fife, *CG32111* and *Spab*. *Sif* is involved in actin cytoskeleton organization; visual perception and

it localizes to the synapse. *Comm2* is essential for nerve cord development and axon guidance.

Bru-3 is known to mediate mRNA translation repression. While *Fife* is responsible for encoding

synaptic scaffolding protein, which promotes neurotransmitter release. *CG32111* is a long non-

coding RNA though much information is not known about this gene. *Spab* is a protein coding gene

from the neuropeptide gene group, however not much is known about this gene (Flybase.org). As

expected, most of the genes responsible for the CCR response play a role in the nervous system.

However, *Sif* and *CG32111* are known to be a target of the circadian clock transcription complex

CLK/CYC, suggesting that CCR photoperiod response could be regulated by the clock (Abruzzi

et al., 2011). *Sif* is also known to interact with *Fas2* which than interacts with *PKCdelta* (Rees et

al., 2011).

Discussion

In this study, we were able to establish that there is a variation in photoperiod response measured

by the CCR phenotype within the DGRP population. By calculating the effect size (Cohen's D

value) for each DGRP genotype we determined how each genotype's photoperiod response was

different from one another. The GWAS analysis on CD values allowed us to narrow down the 32

photoperiod candidate genes. We chose 4 top candidate photoperiod genes for genetic analysis.

The allelic mutant of *PKCdelta* showed a significantly different photoperiod response from WT.

To test the necessity of the gene for the photoperiodic response, we analyzed two different mutant

alleles: *PKCdelta*:Pbac and *PKCdelta*:MIC. By performing a qPCR assay under LD and SD it was

confirmed that *PKCdelta*:Pbac has decreased expression of RD and RE transcripts of the gene.

The *PKCdelta*:MIC mutant which show reduced expression of the RD transcript, show a

photoperiod effect under LD in the RE transcript levels. The CCR assay showed that

PKCdelta:Pbac mutants have a greater effect size compared to *PKCdelta*:MIC. Finally, to test the

hypothesis whether there is an overlap in the genetic mechanisms between the circadian clock and

photoperiodism, the circadian periods of our candidate genes were analyzed. *GlcAT-P* knockout

and *PKCdelta*;MIC mutants had a significantly longer period phenotype from the WT. Our data

supports the view that *PKCdelta* is involved in both circadian clock and photoperiodism.

Diapause and CCR are two different output pathways that are under the photoperiod

regulation in insects. Our data showed that the two different mutant alleles of *PKCdelta* manifested

the altered photoperiod response in different ways. The flies with *PKCdelta*:MIC allele could not

distinguish LD from SD, whereas the flies with *PKCdelta*:Pbac allele had an elevated instances of

diapause under both photoperiods (Figure 4A). The CCR data suggests that the *PKCdelta*:Pbac

mutation plays a role in increasing the sensitivity to photoperiod in the flies overall. Both the

mutants have positive CD values indicting their recovery time was faster than that of WT. While

PKCdelta:Pbac compared to *PKCdelta*:MIC has a greater effect size (Figure 4B).

Taken together, we have a working model for the photoperiodism in *D. melanogaster*. The

photoperiod candidate gene *PKCdelta* is hypothesized to be regulated by the clock. There are

certain genetic components that are under clock control which interact with *PKCdelta*. The

circadian transcription complex CLK/CYC is known to directly regulates the *Mef2* transcription

which then has been identified to target *Fas2*. *Mef2* is known for its role in myogenesis and

development (Sivachenko et al., 2013). However, *Mef2* is shown to be present in clock neurons

and fluctuation in *Mef2* are rhythmic (Sivachenko et al., 2013). *Mef2* is a major component of the

CLK/CYC, which is important for circadian behavior of neuronal morphology. A Chip-Chip

analysis determined that *Fascilin2* (*Fas2*) is a direct target of *Mef2* (Sivachenko et al., 2013). *Fas2*

is known to be involved in neural cell adhesion, and there is a rhythmic oscillation of *Fas2* in

pigment-dispersing factor (PDF) cells types, suggesting that *Fas2* is under circadian clock control

(Sivachenko et al., 2013). *PKCdelta* is proposed to interact with *Fas2* (Rees et al., 2011). PDF is

known to effect *Drosophila* circadian clock where certain mutations in PDF can shorten the period

of certain clock neurons while lengthening the period others neurons (Wülbeck et al., 2008). The

diapause assay showed that *PKCdelta*:MIC mutants lose the ability to distinguish between day-

lengths. *PKCdelta*:MIC mutant's loss of photoperiod sensitivity is similar to the well-studied

PDF^{01} mutant (Ojima et al., 2018).Our data reject the model that circadian clock and

photoperiodism are separate entities with no shared regulations. The current study sheds light onto

potential genes that could be further studied to characterize the mechanisms of photoperiodism,

and its relationship to the circadian clock.

Data Availability

All DGRP lines are available at the Bloomington Drosophila Stock Center, and the genotype

information of all lines are available at http://dgrp2.gnets.ncsu.edu. The raw CCR and CD data for

all DGRP lines are available under File S1. The Files S2, S4, & S6 contain the top GWAS

associations along with qq-plots, and input data used to obtain the GWAS results from CD, LD,

and SD data sets. The Files S3, S5, & S7 contain all associations found in the GWAS study using

the CD, LD and SD data sets. All Supplemental Materials are available at figShare portal,

gsajournals.figshare.com.

Acknowledgements

The authors are grateful for Drs. Nir Yakoby, Nathan Fried, and Charot Rodeget for critical reading

and insightful suggestions. We are grateful for the Lee lab members, especially Sienna Casciato

and Somya Patro, for helpful discussions. The authors would also like to thank Cody Stevens for

helping us navigate through some of the challenging aspects of the project. The work was funded

by the NIH MARC U-STAR, T34GM127154.

Supplemental Material

Figure S1. The distribution of core 38 set DGRP parent strains.

Figure S2. Sex difference in CCR assay among the select DGRP strains.

Figure S3. The Gene enrichment results of the 32 hits using flymodEnrichr.

Figure S4. Diagram of the two *PKCdelta* mutant alleles, Pbac and MIC.

Figure S5. Overview of the *PKCdelta* gene structure and the position of the oligos for PCR

products.

Figure S6. Sequence variations in *PKCdelta* RE transcript.

Figure S7. Phylogenetic analysis of *PKCdelta* RE transcript.

Figure S8. Cohen's D and Genome Effect Size of *PKCdelta* mutants.

Figure S9. The regulatory and catalytic domains of *PKCdelta*.

Table S1. The top 32 hits of the GWAS analysis sorted by the significant p values.

Table S2. Oligos used in the current study.

Table S3. Predicted changes *PKCdelta*:Pbac mutant.

Table S4. Predicted changes in *PKCdelta*:MIC mutant.

File S1. Raw CCR and Cohen's D data for all the DGRP lines. It's divided into three tabs;

Cohen's values, CCR in LD and CCR in SD.

File S2. TOP GWAS hits along with images of QQ plots.

File S3. SNP associations found in the GWAS study.

File S4. TOP GWAS hits using the CCR data in LD.

File S5. All GWAS hits using the CCR data in LD.

File S6. TOP GWAS hits using the CCR data in SD.

File S7. All GWAS hits using the CCR data in SD.

References

Abruzzi, K.C., Rodriguez, J., Menet, J.S., Desrochers, J., Zadina, A., Luo, W., Tkachev, S., and

510   Rosbash, M. (2011). Drosophila CLOCK target gene characterization: implications for

511   circadian tissue-specific gene expression. Genes & Development *25*, 2374–2386.

512   Bradshaw, W.E., and Holzapfel, C.M. (2010). What Season Is It Anyway? Circadian Tracking

513   vs. Photoperiodic Anticipation in Insects. J. Biol. Rhythms *25*, 155–165.

514   Carstensen, M.B., Hertz, H., Bering, T., Møller, M., Rohde, K., Klein, D.C., Coon, S.L., and

515   Rath, M.F. (2019). Circadian regulation and molecular role of the *Bsx* homeobox gene in

516   the adult pineal gland. J. Pineal Res.

Chiu, J.C., Low, K.H., Pike, D.H., Yildirim, E., and Edery, I. (2010). Assaying Locomotor

Activity to Study Circadian Rhythms and Sleep Parameters in Drosophila. JoVE J. Vis.

Exp. e2157.

Dubowy, C.M., and Cavanaugh, D.J. (2014). Sleep: A Neuropeptidergic Wake-Up Call for Flies.

Curr. Biol. *24*, R1092–R1094.

Harbison, S.T., Kumar, S., Huang, W., McCoy, L.J., Smith, K.R., and Mackay, T.F.C. (2019).

Genome-Wide Association Study of Circadian Behavior in Drosophila melanogaster.

Behav. Genet. *49*, 60–82.

Hosler, J.S., Burns, J.E., and Esch, H.E. (2000). Flight muscle resting potential and species-

specific differences in chill-coma. J. Insect Physiol. *46*, 621–627.

Huang, W., Massouras, A., Inoue, Y., Peiffer, J., Ramia, M., Tarone, A.M., Turlapati, L.,

Zichner, T., Zhu, D., Lyman, R.F., et al. (2014). Natural variation in genome architecture

among 205 Drosophila melanogaster Genetic Reference Panel lines. Genome Res. *24*,

1193–1208.

Johnson, C.H., Zhao, C., Xu, Y., and Mori, T. (2017). Timing the day: what makes bacterial

clocks tick? Nat Rev Microbiol *15*, 232–242.

Johansson, M., and Staiger, D. (2015). Time to flower: interplay between photoperiod and the

circadian clock. Journal of Experimental Botany *66*, 719–730.

Kim, B.-T., Tsuchida, K., Lincecum, J., Kitagawa, H., Bernfield, M., and Sugahara, K. (2003).

Identification and Characterization of Three Drosophila melanogaster

Glucuronyltransferases Responsible for the Synthesis of the Conserved

Glycosaminoglycan-Protein Linkage Region of Proteoglycans. J. Biol. Chem. *278*, 9116–

9124.

Kumar, A. (2009). An Overview of Nested Genes in Eukaryotic Genomes. Eukaryot. Cell *8*,

1321–1329.

Lanciani, C.A., Lipp, K.E., and Giesel, J.T. (1992). The effect of photoperiod on cold tolerance

in Drosophila melanogaster. Journal of Thermal Biology *17*, 147–148.

Larkin A, Marygold SJ, Antonazzo G, Attrill H, dos Santos G, Garapati PV, Goodman JL,

Gramates LS, Millburn G, Strelets VB, Tabone CJ, Thurmond J and the FlyBase

Consortium (2021) FlyBase: updates to the Drosophila melanogaster knowledge

base. Nucleic Acids Res. 49(D1) D899–D907

Lees, A.D. (1960). Some Aspects of Animal Photoperiodism. Cold Spring Harb. Symp. Quant.

Biol. *25*, 261–268.

Mackay, T.F.C., Richards, S., Stone, E.A., Barbadilla, A., Ayroles, J.F., Zhu, D., Casillas, S.,

Han, Y., Magwire, M.M., Cridland, J.M., et al. (2012). The Drosophila melanogaster

Genetic Reference Panel. Nature *482*, 173–178.

Martinez-Bakker, M., Bakker, K.M., King, A.A., and Rohani, P. (2014). Human birth

seasonality: latitudinal gradient and interplay with childhood disease dynamics. Proc. R.

Soc. B Biol. Sci. *281*, 20132438.

Maruani, J., Anderson, G., Etain, B., Lejoyeux, M., Bellivier, F., and Geoffroy, P.A. (2018). The

neurobiology of adaptation to seasons: Relevance and correlations in bipolar disorders.

Chronobiol. Int. *35*, 1335–1353.

Mukai, A., and Goto, S.G. (2016). The clock gene period is essential for the photoperiodic

response in the jewel wasp Nasonia vitripennis (Hymenoptera: Pteromalidae). Applied

Entomology and Zoology *51*, 185–194.

Nagy, D., Andreatta, G., Bastianello, S., Martín Anduaga, A., Mazzotta, G., Kyriacou, C.P., and

Costa, R. (2018). A Semi-natural Approach for Studying Seasonal Diapause in

Drosophila melanogaster Reveals Robust Photoperiodicity. J. Biol. Rhythms *33*, 117–

125.

Ojima, N., Hara, Y., Ito, H., and Yamamoto, D. (2018). Genetic dissection of stress-induced

reproductive arrest in Drosophila melanogaster females. PLOS Genetics *14*, e1007434.

Ortavant, R., Mauleon, P., and Thibault, C. (1964). Photoperiodic Control of Gonadal and

Hypophyseal Activity in Domestic Mammals. Ann. N. Y. Acad. Sci. *117*, 157–193.

Pegoraro, M., Gesto, J.S., Kyriacou, C.P., and Tauber, E. (2014). Role for circadian clock genes

in seasonal timing: testing the Bünning hypothesis. PLoS Genet. *10*, e1004603.

Saunders, D.S. (1990). The Circadian Basis of Ovarian Diapause Regulation in Drosophila

melanogaster: Is the period Gene Causally Involved in Photoperiodic Time

Measurement? J. Biol. Rhythms *5*, 315–331.

Saunders, D.S. (2012). Insect photoperiodism: seeing the light. Physiol. Entomol. *37*, 207–218.

Saunders, D.S., Henrich, V.C., and Gilbert, L.I. (1989). Induction of diapause in Drosophila

melanogaster: photoperiodic regulation and the impact of arrhythmic clock mutations on

time measurement. Proc. Natl. Acad. Sci. *86*, 3748–3752.

Sawa, M., Kay, S.A., and Imaizumi, T. (2008). Photoperiodic flowering occurs under internal

and external coincidence. Plant Signal. Behav. *3*, 269–271.

Shieh, B.-H., Parker, L., and Popescu, D. (2002). Protein Kinase C (PKC) Isoforms in

Drosophila. J. Biochem. (Tokyo) *132*, 523–527.

Song, Y.H., Ito, S., and Imaizumi, T. (2010). Similarities in the circadian clock and

photoperiodism in plants. Curr Opin Plant Biol *13*, 594–603.

Song, Y.H., Kubota, A., Kwon, M.S., Covington, M.F., Lee, N., Taagen, E.R., Laboy Cintrón, D.,

586   Hwang, D.Y., Akiyama, R., Hodge, S.K., et al. (2018). Molecular basis of flowering under

587   natural long-day conditions in Arabidopsis. Nat. Plants *4*, 824–835.

588   Sivachenko, A., Li, Y., Abruzzi, K.C., and Rosbash, M. (2013). The transcription factor Mef2, links the

589   Drosophila core clock to Fas2, neuronal morphology and circadian behavior. Neuron *79*.

590   Suh, K.S., Tatunchak, T.T., Crutchley, J.M., Edwards, L.E., Marin, K.G., and Yuspa, S.H.

591   (2003). Genomic structure and promoter analysis of PKC-δ. Genomics *82*, 57–67.

592   Tauber, E., and Kyriacou, B.P. (2001). Insect Photoperiodism and Circadian Clocks: Models and

593   Mechanisms. J. Biol. Rhythms *16*, 381–390.

594   Vaziri, P., Ryan, D., Johnston, C.A., and Cripps, R.M. (2020). A Novel Mechanism for

595   Activation of Myosin Regulatory Light Chain by Protein Kinase C-Delta in *Drosophila*.

596   Genetics *216*, 177–190.

597   Venken, K.J.T., Schulze, K.L., Haelterman, N.A., Pan, H., He, Y., Evans-Holm, M., Carlson,

598   J.W., Levis, R.W., Spradling, A.C., Hoskins, R.A., et al. (2011). MiMIC: a highly

599   versatile transposon insertion resource for engineering Drosophila melanogaster genes.

600   Nat. Methods *8*, 737–743.

601   Wülbeck, C., Grieshaber, E., and Helfrich-Förster, C. (2008). Pigment-Dispersing Factor (PDF)

602   Has Different Effects on Drosophila's Circadian Clocks in the Accessory Medulla and in

603   the Dorsal Brain. J. Biol. Rhythms *23*, 409–424.

605