1 **Pan-cancer analysis of patient tumor single-cell transcriptomes identifies promising**
2 **selective and safe CAR targets in head and neck cancer**
3

4 Sanna Madan[1,2], Sanju Sinha[2], J. Silvio Gutkind[3,4], Ezra E. W. Cohen[3,5], Alejandro A.
5 Schäffer[2,**], Eytan Ruppin[2,**]
6
7 [1] Dept. of Computer Science, University of Maryland, College Park MD 20742 USA
8 [2] Cancer Data Science Laboratory, National Cancer Institute, Bethesda, MD 20892 USA
9 [3] Moores Cancer Center, University of California San Diego, La Jolla, CA 92093 USA
10 [4] Department of Pharmacology, University of California San Diego, La Jolla, CA 92093 USA
11 [5] Department of Medicine, University of California San Diego, La Jolla, CA 92037 USA
12
13
14 **To whom correspondence should be addressed at alejandro.schaffer@nih.gov or
15 eytan.ruppin@nih.gov
16

Keywords
CAR-T, single-cell RNA-sequencing, immunotherapy, head and neck cancer,

glioblastoma, immunoreceptors, targets, selective, safe, solid tumors

Abstract

BACKGROUND: Chimeric antigen receptor (CAR) T cell therapies have yielded

transformative clinical success for patients with blood tumors, but their full potential remains to

be unleashed against solid tumors. One challenge is finding selective targets: cell surface

proteins that are expressed widely by cancer cells and minimally by healthy cells in the tumor

microenvironment and other normal tissues.

METHODS: Analyzing pan-cancer patient tumor single cell transcriptomics data, we first

define and quantify selectivity and safety scores of existing CAR targets for indications in which

they are in clinical trials or approved. Selectivity scores are computed by the ability of a given

surfaceome gene to classify tumor from nontumor cells in the tumor microenvironment. Safety

scores are computed by mining healthy tissue transcriptomics and proteomics atlas data. Second,

we identify new candidate cell surface CAR targets that have better selectivity and safety scores

than the leading targets among those currently being tested, in an indication-specific manner.

RESULTS: Remarkably, in almost all cancer types, we cannot find such better targets, testifying

to the overall near optimality of the current target space. However, in HPV-negative head and

neck squamous cell carcinoma (HNSC), for which there is currently a dearth of existing CAR

targets, we find five new targets that have both superior selectivity and safety scores. Among the

HNSC new targets, we find a few that additionally are strongly essentiality in HNSC cell lines.

CONCLUSIONS: The newly identified cell surface targets lay a basis for future investigations

of better HNSC CAR treatments.

Background

Chimeric antigen receptor T cell (CAR-T) therapy has revolutionized cancer treatment. After a

patient's T cells are extracted, they are modified to recognize a tumor-specific protein via the

addition of a synthetic CAR on the cell membrane, and then infused back into the patient. CAR-

T treatment has resulted in durable remissions in patients, particularly in those with B-cell

malignancies. Recently, CARs directed against the B-cell marker CD19 have been declared

curative, with observed remission rates as high as 90% [1]. The physiological phenomenon that

makes B cell tumors more amenable to CAR-T therapy than other tumors is that one can kill all

the B cells – both cancerous and not cancerous – and the patient is likely to survive, especially if

treated with intravenous immunoglobulin (IVIG) to replace the missing antibodies.

As of September 2021, the Trialtrove repository of clinical trials [2] had recorded more

than 1,500 clinical trials involving cells engineered to target various membrane surface proteins

or other antigens in specific indications that have either taken place, are currently in progress, or

are proposed/planned. Most of these trials involve modified CAR T cells and some involve NK

cells; because our analysis applies to both T and NK cells, we use "CAR" without a suffix in

most places. The CAR targets that are currently FDA-approved or are being tested include a

variety of proteins, *e.g.*, CD19 and CD22 in B-cell malignancies, and HER2, IL-13 receptor α2,

EGFRvIII, carbonic anhydrase IX, and MUC1 in various solid tumors [3].

While CAR-T treatments have been remarkably successful against B-cell tumors, several

challenges remain for realizing their potential in the realm of solid tumors. Two key challenges

are: (1) heterogeneity of cell surface proteins expressed by a tumor's constituent malignant cells,

and (2) expression of the CAR target protein on the cell surfaces of normal tissues elsewhere in

the body, posing a toxicity risk. These challenges may be addressed by studying single-cell RNA

sequencing (scRNA-seq), which has generated many patient tumor transcriptomes in recent

years. We hypothesized that we could harness these single-cell measurements to profile the

expression of genes encoding targets of CAR therapies and more broadly, cell surface proteins,

at a cellular resolution – thus enabling us to identify suitable surface proteins that had not

previously been considered seriously as targets for CAR therapy. Indeed, recent studies have

harnessed scRNA-seq to investigate toxicities of CAR-T therapies [4,5] and highlighted the

importance of optimizing surface target selection by utilizing scRNA-seq [6–8]. Furthermore, a

notable recent analysis leveraged public patient tumor and reference atlas scRNA-seq data to

first construct an integrated "meta-atlas" of tumor and normal cells and then identify tumor-

specific combinations of target antigens via logic gates [9]. Hopefully, identifying optimal cell

surface targets in a systematic, data-driven manner may translate in the future to improved

patient outcomes, leading to more durable remissions and reduced toxicities.

Zeroing in on the challenges listed above, we asked whether mining scRNA-seq datasets

of solid tumors can enable us to: (1) First, chart the landscape of existing cell-surface targets

against which CARs are currently aimed at in clinical trials, and estimate their selectivity and

safety with quantitative scores. (2) Second, identify alternative target proteins that are

differentially highly expressed by tumor cells and lowly expressed by non-tumor cells within the

tumor microenvironment (TME) and additionally in normal tissues across the body. Our goal is

to identify new targets for CAR therapies that have better safety and selectivity scores than the

targets currently tested, in an indication-specific manner.

We use the term "indication" to refer to an ordered pair of (cancer type, CAR target) that

has been proposed in at least one clinical trial; we use the opposite term "non-indication" to

denote a (cancer type, CAR target) for which we could not find any proposed clinical trial

(**Methods**). Our analysis proceeds in the following steps: (1) First, we mine clinical trial

databases and literature, we identify the current CAR-T targets in clinics mapped to the solid

tumors in which they are indicated / targeted against. For each (cancer type, CAR target) pair, we

compute a **tumor selectivity score**, *i.e.*, the selectivity of the expression of a given candidate

target gene in tumor cells versus all other cells in the TME. This is quantified by the ability of a

classifier to separate tumor cells from non-tumor cells based on the single cell expression data of

this gene in the TME of the respective cancer indication. (2) Second, we compute a **safety score**

for each candidate target, which is computed by quantifying the level of protein and scRNA-seq

expression across normal human tissues. We additionally investigate the *in vitro* essentiality of

genes encoding CAR targets that are either approved or being tested in clinics as well as the

candidate new targets yielded by our analysis in CRISPR knockout screens. The overview of our

approach showing all three resource inputs is shown in **Figure 1**.

The first part of our analysis evaluates the indications that are currently studied in clinical

trials (or already approved) for their tumor selectivity and safety metrics. We then compare

untested, candidate CAR target surface protein versus these current state-of-the-art benchmarks.

Our goal is to identify new target genes that match or exceed the leading extant targets'

selectivity and safety scores in each respective cancer type, thus forming promising new target

tumor antigens for further possible consideration when thinking about engineering new CARs. In

difference from previous related studies [7,9], the unique aim of our study is to find novel targets

for CARs that are strictly better than the best CAR target antigens in clinics based on the

selectivity and safety metrics. This analysis, quite surprisingly, identifies only a few but highly

promising candidates for clinical translation.

Figure 1: The left panel shows some solid cancer locations and CAR-T targets being tested for them. The right panel shows that our study uses single-cell RNA data from patients, healthy cell atlases, and CRISPR knockout screens as inputs to help us select CAR-T targets on the right.

Methods

Clinical trial curation

To obtain a list of CAR clinical trials and targets in solid tumors, we first obtained a list of single clinical CAR targets from MacKay *et al.*, who carried out their ClinicalTrials.gov query on March 3, 2019 [10]. To obtain a complete up-to-date list of clinical trials since then, an additional search was carried out on ClinicalTrials.gov on August 20, 2021 for CAR-T clinical trials posted on March 3, 2019 and after. The following query was used: "*CAR-T OR CAR OR chimeric antigen receptor OR chimeric immunoreceptors OR artificial T-cell receptors | Recruiting, Not yet recruiting, Active, not recruiting, Completed, Enrolling by invitation,*

Suspended, Terminated Studies | Interventional Studies | (cancer OR carcinoma OR solid tumor

OR solid tumors OR melanoma OR sarcoma) NOT (hematological OR lymphoid OR liquid

cancers OR lymphoma OR leukemia OR CD19 OR B cell malignancy OR myeloma OR

Hematologic OR blood cancer) | First posted from 03/03/2019 to 08/20/2021". CAR trials and

targets were further identified through manual curation, and clinical trial statuses were cross-

checked on TrialTrove (https://citeline.informa.com/trials/results) to ensure consistency [2].

Access to TrialTrove requires a license.

CAR protein targets were then selected for further analyses if they had a corresponding single

gene that could be analyzed for gene expression. The following molecules were left out of the

analysis: Claudin 18.2, CD44v6, EGFRvIII, k-IgG, LeY, CEA, and GD2 because i) they are not

proteins or ii) they are protein isoforms whose presence and abundance cannot easily be

estimated from single cell gene expression data. For each target, its indications were identified as

the specific tumor types which were mentioned in its ClinicalTrials.gov records, excluding 'solid

tumors' and 'metastases'. The following mapping of clinical trial cancer types to our datasets

was used: brain cancer or glioma → glioblastoma; pancreatic cancer → pancreatic

adenocarcinoma; stomach cancer or gastric cancer → stomach adenocarcinoma; liver or

hepatocellular carcinoma → liver hepatocellular carcinoma; head and neck cancer → head and

neck squamous cell carcinoma; melanoma → skin cutaneous melanoma; colon cancer or

colorectal cancer → colorectal cancer; ovarian cancer → ovarian serous cystadenocarcinoma;

lung cancer or non-small cell lung cancer (but not small cell lung cancer) → non-small cell lung

cancer; breast cancer → breast invasive carcinoma. Liquid tumor CAR targets were taken from

MacKay *et al*. [10]

Single-cell transcriptomics data collection

We downloaded single cell transcriptomics datasets from the Tumor Immune Single-cell Hub

(TISCH) 1.0 database [11]. This resource contains 76 uniformly processed scRNA-seq datasets

spanning 28 cancer types and containing data from over 2 million cells. We selected datasets

from human tissues which contained *both* malignant and non-malignant cells. This yielded a total

of 26 unique solid tumor datasets: two of breast cancer (BRCA), one of colorectal cancer (CRC),

eleven of glioblastoma (GBM), one of head and neck squamous cell carcinoma (HNSC), one of

liver hepatocellular carcinoma (LIHC), four of non-small cell lung cancer (NSCLC), one of

ovarian cancer (OV), two of pancreatic adenocarcinoma (PAAD), two of skin cutaneous

melanoma (SKCM), and one of stomach adenocarcinoma (STAD). Additionally, an acute

myeloid leukemia (AML) dataset was analyzed. These data were all uniformly processed by

TISCH as $\log_2(\text{TPM}/10+1)$ values (TPM stands for transcripts per million).

Single-cell data analysis

In-house programs written in Python 3 were used to carry out analyses on the scRNA-seq

datasets. A cell was labeled 'tumor' if it was labeled 'malignant' by TISCH, and 'non-tumor'

otherwise. We then partitioned the cells in each dataset by patient (if such annotation was

available); the partition by patient is so that each patient counts equally in the downstream

analysis regardless of how many cells were sampled from that patient. As quality control, only

patient samples which contained at least 25 tumor cells and 25 non-tumor cells were selected for

analysis. After this filtering step, only the cancer types where we had samples from at least three

patients remaining were subsequently considered, excluding breast cancer entirely from our

analysis, for which we only had two patients.

A list of 3,559 cell surface protein encoding genes was obtained from Hu *et al* [7]. Genes whose

symbols differed in that reference in TISCH and in DepMap (see later subsection) were resolved

by using the Human Gene Nomenclature Committee table including gene aliases (downloaded

from https://www.genenames.org) We generated patient-specific-AUCs of tumor-vs-non-tumor

cell discrimination for each gene whose respective protein was listed in a CAR clinical trial and

for all cell surface protein encoding genes at large. For each existing (CAR target, indication) we

compute a mean tumor selectivity score over all patient-specific selectivity scores. We took the

top-ranking target in each cancer type by the mean tumor selectivity scores as the threshold to

which we compared all other cell surface protein encoding genes. In doing these comparisons,

we used the Wilcoxon signed rank one-sided test comparing matched patient-level AUCs, with

Benjamini-Hochberg FDR p-value < 0.1 denoting statistical significance, repeating this

procedure for each cancer type separately.

Reference atlas analysis: computing safety scores

For computing the safety score of a cell surface protein in the Human Protein Atlas (HPA), an

approach similar to that outlined in MacKay *et al.* [10] was taken. Individual tissue types were

grouped into tissue groups, with the following mapping:

Tissue Group	Tissue types
Adipose and soft	adipose tissue, soft tissue
Endocrine	adrenal gland, parathyroid gland, thymus, thyroid gland
Nervous system	caudate, cerebellum, cerebral cortex, choroid

	plexus, hippocampus, hypothalamus, pituitary gland, substantia nigra, dorsal raphe
Circulatory	heart muscle
Immune tissue	appendix, lymph node, spleen, tonsil
Immune cells	bone marrow
Female tissue	breast, cervix, endometrium, fallopian tube, lactating breast, ovary, placenta, vagina
Lungs	bronchus, lung, nasopharynx
Colon	colon
Gastrointestinal	duodenum, oral mucosa, rectum
Male tissue	epididymis, prostate, seminal vesicle, testis
Esophagus	esophagus
Eye	eye, retina
Gallbladder	gallbladder
Skin	hair, skin, sole of foot
Urinary	kidney, urinary bladder
Liver	liver
Pancreas	pancreas
Salivary gland	salivary gland
Muscle	skeletal muscle, smooth muscle
Small intestine	small intestine
Stomach	stomach

HPA protein measurements for a given tissue type are either 'Not detected', 'Low', 'Medium',

or 'High'. For a tissue group, if the measurements of a protein over all tissue types only ranged

from 'Not detected' to 'Low', then the tissue group was assigned 'ND-L'. If the measurements

over all tissue types only ranged from 'Medium' to 'High', then the group was assigned 'M-H'.

If values from either 'Not detected' or 'Low' AND 'Medium' or 'High' were present in the

measurements over all tissue types, then the group was assigned 'Variable.' Thus, in determining

the overall HPA safety score of a protein, the scoring weight is in the following order: ND-L >

Variable > M-H. For each tissue group, a score of 2 was added for each 'ND-L', 1 was added for

each 'Variable', and 0 was added for each 'M-H.' The mean of the resulting sum was taken to

obtain a representative measure over all tissue groups that were measured for the protein. As

such, the scoring values ranged from 0 to 2.

For the Tabula Sapiens safety score computation, the tissues measured were taken as is, with no

further grouping, as the following: Bladder, Blood, Bone Marrow, Eye, Fat, Heart, Kidney,

Large intestine, Liver, Lung, Lymph node, Mammary Gland, Muscle, Pancreas, Prostate,

Salivary Gland, Skin, Small intestine, Spleen, Thymus, Tongue, Trachea, Uterus, and

Vasculature. To compute the safety score of a surfaceome gene, first, within each tissue, the

percentage of cells expressing that gene was computed. Next, for a tissue, if the percentage was

< 1%, 10 was added to the final score; if it fell between 1-10%, 9 was added to the final score; if

it fell between 11-20%, 8 was added to the final score; if it fell between 21-30%, 7 was added to

the final score; if it fell between 31-40%, 6 was added to the final score; if it fell between 41-

50%, 5 was added to the final score; if it fell between 51-60%, 4 was added to the final score; if

it fell between 61%-70%, 3 was added to the final score; if it fell between 71-80%, 2 was added

to the final score; if it fell between 81-90%, 1 was added to the final score; and finally, if it fell

between 91-100%, 0 was added to the final score. The mean of this sum was taken over the total

number of tissue groups for which the gene had been measured. As such, the scoring values

ranged from 0 to 10.

DepMap essentiality scores

We used the DepMap Public 22Q2, specifically 'CRISPR_gene_dependency.csv' was

downloaded from https://depmap.org/portal/. This file contains the probability of dependence

scores for each (cell line,gene) pair. The DepMap defines a probability of ≥ 0.5 as essential, and

as such, our analysis defined a cell line as dependent on a gene if its p(dependent) ≥ 0.5.

Cell lines were partitioned by cancer type for each clinical or novel CAR target gene. For CRC,

we analyzed cell lines labeled 'Colon/Colorectal Cancer' as the primary disease; for GBM, we

analyzed cell lines labeled 'Brain Cancer' as primary disease and 'Glioblastoma' as the subtype;

for NSCLC, we analyzed cell lines labeled 'Lung Cancer' as primary disease and 'Non-Small

Cell Lung Cancer (NSCLC)' as the subtype; for PAAD we analyzed cell lines labeled

'Pancreatic cancer' as the primary disease and 'Adenocarcinoma' as the subtype; for STAD we

analyzed cell lines labeled 'Gastric Cancer' as the primary disease and 'Adenocarcinoma' as the

subtype; for OV we analyzed cell lines labeled 'Ovarian Cancer' as the primary disease with no

constraint on subtype; for SKCM we analyzed cell lines labeled 'Skin Cancer' with 'Melanoma'

as the subtype; for HNSC we analyzed cell lines labeled 'Head and Neck Cancer' as the primary

disease and 'Squamous Cell Carcinoma' as the subtype; for B-cell Non-Hodgkin's Lymphoma

we analyzed cell lines labeled 'Lymphoma' as the primary disease and 'B-cell, Non-Hodgkin's'

as the subtype; for B-cell Hodgkin's Lymphoma we analyzed cell lines as primary 'Lymphoma'

and subtype 'B-cell, Hodgkin's'; for T-cell, Non-Hodgkin's Lymphoma, we analyzed cells

labeled 'Lymphoma' as primary disease and 'T-cell, Non-Hodgkin's' as subtype; for T-cell

Lymphoma we analyzed cell lines with 'Lymphoma' primary and 'T-cell' subtype; for B-cell

Lymphoma, we analyzed cell lines with 'Lymphoma' primary and 'B-cell' subtype; for Acute

Myelogenous Leukemia (AML) we analyzed cell lines with 'Leukemia' primary and 'Acute

Myelogenous Leukemia' in the subtype; for Chronic Myelogenous Leukemia we analyzed cell

lines with 'Leukemia' as the primary and 'Chronic Myelogenous Leukemia' as the subtype; for

B-cell Chronic Lymphoblastic Leukemia (B-CLL), we analyzed cell lines with 'Leukemia' as

the primary and 'Chronic Lymphoblastic Leukemia' as the subtype; for B-cell Acute Myeloid

Leukemia (AML) we analyzed cell lines with 'Leukemia' as the primary and 'Acute

Lymphoblastic Leukemia (ALL), B-cell' as the subtype; for T-cell AML, we analyzed cell lines

with 'Leukemia' as the primary and 'Acute Lymphoblastic Leukemia (ALL), T-cell' as the

subtype; for unspecified B-cell Leukemia we analyzed cell lines with 'Leukemia' as the primary

and 'B-cell, unspecified' as the subtype; for Myeloma we analyzed cell lines with 'Myeloma' as

the primary.

For each (cancer type, gene) indication, the mean p(essentiality) of the gene was simply taken

over all p(essential) values in the cell lines of the cancer type indication as mapped above.

The DepMap (https://depmap.org/portal/documentation/) defines a 'strongly selective' gene "as

having a dependency distribution that is at least 100 times more likely to have been sampled

from a skewed t-distribution than a normal distribution." *i.e.*, its skewed-LRT value exceeds 100.

A 'common essential' (*i.e.*, 'pan-essential') gene "is defined as a gene which, in a large, pan-

cancer screen, ranks in the top X most depleting genes in at least 90% of cell lines. X is chosen

empirically using the minimum of the distribution of gene ranks in the 90th percentile least

depleting lines."

Results

Overview of the analysis

We analyzed nine cancer types that have sufficient data in the scRNA-seq Tumor Immune

Single-cell Hub (TISCH) 1.0 collection: colorectal cancer, glioma, (HPV-negative) head and

neck cancer, liver cancer, non-small cell lung cancer, ovarian cancer, pancreatic

adenocarcinoma, melanoma, and stomach cancer [11] (**Methods**). To quantify the *tumor*

selectivity of a given gene encoding either an extant or new candidate CAR target cell surface

protein, we used the "area under the curve" (AUC) metric, which quantifies the extent by which

its expression levels can be used to discriminate between tumor and non-tumor cells in the TME

(**Methods**). We use 'AUC' and '*tumor selectivity score*' interchangeably. The tumor selectivity

score is computed for each patient individually, such that for a given cell surface protein target in

a cancer type, we obtain a distribution of tumor selectivity scores across all patients, whose mean

and standard deviation can be quantified.

In parallel, to quantify the *overall safety* of a target gene encoding a cell surface protein, a

two-part *safety score* was computed, considering separately the Human Protein Atlas (HPA)

proteomics data [12] and the scRNA-seq Tabula sapiens gene expression data [13]. Each of these

two *sub-safety scores* is computed at the tissue level and averaged over the set of pertaining

samples; for each resource, the safety measures are computed for each tissue separately and then

the arithmetic mean of safety scores across all tissues is taken as the final safety score

(**Methods**). The HPA safety scores range from 0-2, and the Tabula Sapiens safety scores range

from 0-10, following each resource's scoring.

The quantified tumor selectivity and safety landscape of the human surfaceome

To obtain a birds-eye view of the tumor selectivity and safety metrics outlined in the previous

section, we quantified their values across all surfaceome genes (curated from Hu *et al*. [7]). The

cancer-type-specific distributions of tumor selectivity scores across all surfaceome genes are

shown in **Figure 2A**. The distributions are highly similar across cancer types: all are symmetric

and unimodal with AUC values tightly concentrated around 0.50.

We examined how many highly tumor-specific (AUC > 0.75) surfaceome genes were

upregulated in each cancer type (**Figure 2B**). Interestingly, the number of such surfaceome genes

varies considerably by cancer type, with the lowest values in GBM (n=2), OV (n=6), and LIHC

(n=7), and highest values in SKCM (n=42), CRC (n=50), and HNSC (n=84). This suggests that

the latter subset of solid tumor types may contain a larger space of surface targets that may be

considered for CAR therapeutic development, compared to the former subset of solid tumor

types. Finally, the distribution of safety scores derived from the HPA and Tabula sapiens atlases

are shown in **Figure 2C** and **Figure 2D**, respectively. Both distributions are strongly skewed to

the left; most surfaceome genes and proteins have considerably high safety scores (HPA safety

mean=1.38, median=1.55, standard deviation=0.53; Tabula sapiens safety mean=8.56,

median=9, standard deviation=1.54). Additionally, the quantified cancer-type-specific selectivity

scores and safety scores derived from each resource (with tissue-specific resolution) for all

surfaceome genes are provided in **Additional File 2: Tables S1-S3**.

Figure 2: The landscape of tumor selectivity and safety scores of human surfaceome genes.
(A) 1D scatterplots depicting the tumor selectivity scores (AUCs) of all surfaceome genes within each cancer type (red curves). Within each cancer-type-plot, the top N (where N is < 10) genes are labeled and represented by green points instead of red. Genes are sorted in ascending order on the x-axis from left to right. **(B)** The number of tumor-specific surfaceome genes (AUC > 0.75) in each cancer type, in ascending order from top to bottom. Histograms depicting the distributions of HPA-derived safety scores and Tabula sapiens-derived safety scores are shown in **(C)** and **(D)**, respectively.

The selectivity and safety landscape of targets of approved and currently studied

CARs

Mining TrialTrove, ClinicalTrials.gov, and existing literature [10], we assembled a table

summarizing the genes encoding proteins that are currently being targeted by CARs in clinical

trials in each cancer type, including those already approved **(Table 1 & Methods)**. We term

those *clinical* CAR targets. For almost all the solid tumors we have considered, we find at least

10-20 unique genes encoding cell surface proteins that are currently targeted by CARs in various

stages of active or planned clinical trials. The most frequently studied genes among those are

PSCA, *MUC1*, *CD274*, *EGFR*, and *ERBB2*. Notably, however, head and neck cancer (HNSC)

has a dearth of unique CAR targets, with only two targets currently in clinical trials, *EGFR* and

ERBB2, suggesting that there may be a critical unmet need to identify additional cell surface

targets for HNSC. Following this assessment, we charted the distribution of tumor selectivity

scores for each cancer type's pertaining clinical CAR targets (**Figure 3A, 3B**).

Cancer type	Genes encoding clinical CAR targets
CRC	MUC1, CD274, PROM1, ROR1, PSCA, MET, EPCAM, EGFR, FOLH1, ERBB2, MSLN, CD276, TM4SF1, CEACAM5, GUCY2C
GBM	MUC1, CD274, PROM1, ROR1, MPP2, EPHA2, PSCA, IL13RA2, EGFR, ERBB2, MSLN, CD276, BSG
LIHC	MUC1, CD274, PROM1, ROR1, AFP, TNFRSF10B, PTGS2, PSCA, MET, EPCAM, EGFR, GPC3, ERBB2, MSLN, CD276, BSG
NSCLC	MUC1, CD274, ROR1, CD80, CD86, MAGEA1, MAGEA4, PSCA, EGFR, FOLH1, GPC3, ERBB2, MSLN, CD276, TM4SF1
PAAD	MUC1, CD274, PROM1, CD70, ROR2, CLDN18, PSCA, EPCAM, EGFR, ERBB2, MSLN, TM4SF1, GUCY2C
STAD	MUC1, ROR2, PSCA, EPCAM, EGFR, ERBB2, MSLN, CD276, TM4SF1, GUCY2C
OV	PROM1, CD70, FOLR1, MUC16, MET, EGFR, ERBB2, MSLN, ALPP
SKCM	CD70, PMEL, KDR, MET, IL13RA2
HNSC	EGFR, ERBB2

343 **Table 1: Current solid tumor CAR targets in clinics.** Genes encoding cell surface proteins
344 that are currently targeted by CARs in clinical trials, for the nine solid tumor types we studied.
345

Cancer type	Gene with highest tumor selectivity score (0-1)	Gene with highest HPA Safety score (0-2)	Gene with highest TS safety score (0-10)
STAD	TM4SF1 (0.80)	PSCA (2.0)	GUCY2C (9.83)
HNSC	EGFR (0.73)	ERBB2 (1.55)	ERBB2 (7.92)
OV	MSLN (0.75)	MUC16 (1.84)	ALPP (10.0)
SKCM	PMEL (0.84)	PMEL (1.95)	PMEL (9.79)
CRC	EPCAM (0.90)	PSCA (2.0)	GUCY2C (9.83)
GBM	EGFR (0.74)	PSCA (2.0)	IL13RA2 (9.79)
PAAD	EPCAM (0.86)	PSCA (2.0)	CLDN18 (9.83)
LIHC	GPC3 (0.65)	AFP (2.0)	AFP (9.79)
NSCLC	MUC1 (0.80)	PSCA (2.0)	MAGEA1 (10.0)

346
347 **Table 2: The top selectivity and safety scores of current clinical CAR targets within each**
348 **cancer type.**
349

Next, for each cancer type studied, we identified: 1) the leading clinical CAR target based

on its tumor selectivity score, as derived from TISCH scRNA-seq data, 2) the leading clinical

CAR target based on its safety score, as derived from the Human Protein Atlas (HPA), and 3) the

leading clinical CAR target based on its safety score, as derived from the Tabula sapiens (TS)

scRNA-seq reference atlas (**Table 2**). The scores of the leading targets by each criterion are

further used as threshold parameters by which any gene encoding cell surface proteins can be

compared to quantify its potential to serve as a CAR targets.

We find that the tumor selectivity scores of genes encoding clinical CAR targets is higher

than other surfaceome genes in a statistically significant manner in six cancer types (Wilcoxon

rank-sum p-value < 0.05) (**Figure 3A**). Notably, this testifies to the utility of the single-cell

tumor selectivity metric we have derived, as this measure is demonstrably higher among genes

encoding surface proteins that have already been vetted to be targeted by CARs. We similarly

chart the distributions of safety score values, also encouragingly revealing that the safety scores

we derived from the HPA are higher among clinical CAR targets than all surfaceome proteins

(**Figure 3C**, Wilcoxon rank-sum p=0.11). However, we note that the median Tabula sapiens-

derived safety scores of all surfaceome genes and genes encoding clinical CAR targets are

roughly equivalent (**Figure 3D**, Wilcoxon rank-sum p=0.95). We continue to use this sub-safety

score in what follows to adopt a cautious approach and provide as comprehensive a picture as

possible, but we note that it is not discriminatory as the tumor selectivity and HPA safety scores

are. We additionally plotted the distributions of safety scores in all surfaceome genes and cancer-

type-stratified clinical CAR targets (**Additional File 1: Figure S1**), as well as comparisons

between genes encoding clinical CAR targets and the most tumor-selective surfaceome genes in

each cancer type (**Additional File 1: Figure S2**).

Figure 3: Surveying the selectivity and safety scores of known CAR targets.

(A) Violin plots with nested boxplots depicting the tumor selectivity scores of genes encoding
clinical CAR-T targets (shown in red) versus all surfaceome genes (shown in green) for each

cancer type. Wilcoxon rank-sum test with p-values shown in each subplot, $p < 0.05$ significant. **(B)** 1D scatterplots depicting tumor selectivity scores (AUCs) of genes encoding clinical CAR targets within each cancer type. Genes whose selectivity scores are in the top 10% are represented by green points and are labeled, for each cancer type-plot. Genes are sorted in ascending order on the x-axis by AUC from left to right. **(C-D)** Distribution of safety scores of all clinical CAR target encoding genes versus all surfaceome genes. Tabula sapiens derived safety scores **(C)** and HPA derived safety scores **(D)**. Wilcoxon rank-sum comparison test p-values shown at the bottom of each plot.

Identifying novel CAR targets with higher estimated selectivity and safety scores

In the previous subsection, for each cancer type, we identified the top-ranking *existing* CAR

targets based on their tumor selectivity and safety scores. We now pivot to identify new

candidate targets that have better selectivity and safety scores than those of the current therapies.

We proceeded in three steps: (1) identifying highly selective CAR targets (with yet moderate

toxicity levels); (2) identifying high safety scoring CAR targets (with yet moderate selectivity

levels), and finally, (3) identifying the targets that have both very high selectivity and safety

scores (**Figure 4A**). These three steps were conducted for every indication, comparing the scores

of the new targets to the best scoring CAR targets in that cancer type.

        We first aimed to identify novel *highly selective* CAR targets yet having potentially

acceptable low toxicity. We searched for surface proteins that are encoded by genes whose

selectivity scores across patients in a cancer type are significantly higher than the respective top

currently established CAR target (employing a one-sided Wilcoxon signed rank test comparing

patient-level selectivity scores, with FDR p-value < 0.1 denoting significance). To exclude

targets with potentially extreme toxicity levels and only consider those that are most likely

reasonable/safe, we applied lower bound thresholds to each of the sub-safety scores: 1.5 (out of

2) for HPA and 7.5 (out of 10) for Tabula sapiens. Notably, the only cancer types where we find

new higher scoring selective targets are HPV-negative HNSC (seven targets) and glioblastoma

(GBM; one target) (**Figure 4B & Additional File 2: Table S4**).

Second, studying a complementary perspective, we next searched for *highly safe targets*,

that is those that have higher safety scores that the best approved/studied CAR target in a cancer

type. We concomitantly required that these targets are at least moderately selective, with mean

tumor selectivity scores ≥ 0.7. We required that new candidate targets safety scores surpass *both*

atlases' safety scores of the pertaining cancer-type-specific top target(s). Notably, here, we found

highly targets only for HPV-negative HNSC. The eighteen newly suggested targets are shown in

Figure 4C and **Additional File 2: Table S5** along with the established target *ERBB2*.

Figure 4: Identification of novel CAR targets with improved selectivity and safety compared to leading clinical CAR targets. **(A) 2D scatterplots of all surfaceome genes with HPA safety score on the x-axis and tumor selectivity score on the y-axis.** Red points represent clinical CAR target encoding genes for the respective cancer type. Horizontal thresholds denote the highest tumor selectivity score of clinical CAR targets pertaining to each cancer type, while vertical thresholds denote the highest respective HPA safety score of clinical CAR targets pertaining to the cancer type. Clinical CAR targets are shown in red, surfaceome genes which surpass both the horizontal and vertical thresholds (thus appearing in the upper right quadrant) are shown in dark purple. These points emerged only for HNSC. The remaining surfaceome genes are shown in transparent blue. Tabula sapiens safety scores were excluded for simplicity of visualization.
(B) Top selective (and yet safe) new targets. Stems in red denote the leading clinical target by selectivity for each of the two cancer types, HNSC (left) and GBM (right), for which we found new selective targets. Stems in green denote the seven total surfaceome genes which emerged from our analysis as having greater tumor selectivity than the leading target in each cancer type.
(C) Top safe (and yet selective) new targets. Stems colored red denote the leading clinical target by safety score, derived from HPA (left) and Tabula sapiens (right). Stems colored green denote the surfaceome genes which emerged from our analysis as having greater safety scores than the leading clinical CAR targets derived from each atlas. Both plots depict the same set of 18 newly suggested surfaceome genes for HNSC as well as the established target gene *ERBB2*.
(D) Top selective and safe new targets. New surfaceome targets appearing as both highly selective and highly safe from the previous two iterations of the analysis (shown in **B & C**). Rows sorted by descending tumor selectivity scores.

Finally, thirdly, we searched for target genes that have selectivity *and safety scores* that are both higher than those observed for the current targets studied *w*ithin each cancer type. We find five such targets, all for HPV-negative HNSC (**Figure 4D**). Indeed, charting the tumor selectivity scores and HPA safety scores of all surfaceome genes in two dimensions (Tabula sapiens safety scores were excluded to enable clearer visualization, but are provided in **Additional File 2: Table S3**), along with the cancer-type-specific thresholds for both metrics, revealed that HNSC is the only cancer type for which there exist any surface targets that surpass both thresholds and are thus more selective and safer than the leading existing CAR targets (**Figure 4A, 4D**).

Further ranking CAR targets by their essentiality in tumor cells

To evaluate the potential of a surface protein to serve as a CAR target, it is of further interest to study its essentiality in cancer cells. A strong dependency on a target in its pertaining cancer type testifies to a biological dependency on that protein by the tumor cells and subsequently, points to a possible decreased potential for immune evasion via downregulation of the gene or loss of this antigen on the cell surface. To our knowledge, the dependency in cell lines of genes encoding current clinical CAR targets has not yet been studied; we thus analyzed the CRISPR DepMap [37] essentiality scores for these genes (**Figure 5A&B**, **Additional File 2: Table S6-7**) along with the untested candidate targets yielded by our pipeline (**Figure 5C**, **Additional File 2: Table S8**).

To assess the essentiality of a given (cancer type, gene) pair, we analyzed the DepMap cell lines belonging to that indication (**Methods**) and considered a cell line dependent on that gene if p(dependence) ≥ 0.50, as defined by DepMap. For each target gene, we computed the mean p(dependence) over all cancer cell lines of that indication, and the fraction of cell lines in which the gene was deemed essential. We also note that DepMap subcategorizes 'essential' genes as either as 'strongly selective' or 'common-essential." The former denotes strong essentiality in a subset of cell lines and the latter denotes broad, pan-cancer essentiality in almost every cell line.

Intriguingly, most CAR targets in clinical trials exhibit quite low dependency scores across cell lines of their respective indications. One notable exception is the FDA-approved CAR-T target-encoded gene *CD19*, which is essential in 75% (9/12) of non-Hodgkin's B-cell lymphoma cell lines, 25% (3/12) of B-cell acute lymphoblastic leukemia (B-ALL) cell lines, and 100% (3/3)

of unspecified B-cell leukemia cell lines (**Figure 5B, Additional File 2: Table S6**). *CD19* is also

deemed 'strongly selective' by DepMap. Interestingly, the other blood tumor clinical CAR-T

target genes, *CD22* (in Phase 1 trials) and *TNFRSF17* (encoding the BCMA protein; FDA-

approved against multiple myeloma), are not essential in *any* cell line of their respective

indications analyzed here.

In solid tumors the emerging overall picture is similar. Almost all clinical CAR targets are not

essential in any of the cell lines in the pertaining indications, with the striking exceptions of

EGFR and *ERBB2,* which are highly essential across all their indications. For each cancer type

they are indicated in, *EGFR* and *ERBB2* consistently rank the highest both in terms of fraction of

cell lines they are essential in and the mean and median p(dependency) across cell lines (**Figure**

5A). *EGFR* is essential in 76% of HNSC cell lines and *ERBB2* is essential in 52% of HNSC cell

lines (**Additional File 2: Table S7**). Out of the 35 solid tumor clinical CAR target genes that we

collected and analyzed, *EGFR*, *ERBB2*, and *MET* are the only ones that are deemed essential by

the CRISPR DepMap, all 'strongly selective.'

Of the candidate CAR targets yielded by our pipeline, the vast majority are not essential in

any cell lines of their respective cancer indication. However, two targets stand out: *IGSF3* and

SLC2A1, which are essential in 80% and 45% of HNSC cell lines, respectively (**Figure 5C,**

Additional File 2: Table S8). *IGSF3* is classified as 'common-essential' (*i.e.*, 'pan-essential')

and *SLC2A1* is classified as 'strongly selective' by DepMap.

Figure 5: the dependency of genes encoding known CAR targets and our candidate new CAR targets in DepMap cell lines. (A) Probability of essentiality/dependency by cell lines on genes encoding existing solid tumor clinical CAR targets, for each of the nine cancer types studied. The vertical dashed line is at p(0.50), denoting the threshold above which a cell line is considered to be "dependent" on a gene. **(B)** Probability of essentiality/dependency by cell lines on genes encoding existing clinical liquid tumor CAR targets, for lymphomas, leukemias, and myeloma. Vertical dashed line is at p(0.50), denoting the threshold where a cell line is considered to be "dependent" on a gene. **(C)** Probability of essentiality/dependency by cell lines on genes encoding untested CAR targets we proposed for HNSC and GBM. Vertical dashed line is at p(0.50), denoting the threshold where a cell line is considered to be "dependent" on a gene.

Discussion

Overall, there are twenty unique genes encoding surface proteins that were identified by our pipeline across all three filtering configurations (top selective, top safe, and requiring both concomitantly). The twenty genes, the surface proteins they encode, and the families/classes of the respective proteins are shown in **Table 3**. Here we present a narrative summary of seven yielded by our analysis: the five superior targets that are both highly selective and safe, and two that are already existing targets of CAR therapies or antibody-drug conjugates in different indications. Analogous narratives for the remaining candidate surface targets can be found in **Additional File 1: Supp Text**.

We begin with *TACSTD2*, having a very high selectivity score in HNSC (mean AUC=0.94, one-sided Wilcoxon signed rank test FDR p-value=0.065). It encodes the tumor associated calcium signal transducer 2, also known as "Trop-2" and "epithelial glycoprotein-1 antigen", a calcium signal transducing cell surface receptor and known tumor-associated antigen. Among all cancers in The Cancer Genome Atlas (TCGA) [14], HNSC harbors the third highest median *TACSTD2* expression levels (265.2 FPKM). In the HPA Pathology Atlas [15], antibody staining

revealed medium to high membranous and/or cytoplasmic positivity in two of four HNSC

patients. Biallelic loss-of-function mutations in *TACSTD2* are known to cause gelatinous drop-

like corneal dystrophy [16], which suggests that expression of this gene is functionally important

only the eye. Notably, Trop-2 is the target of sacituzumab govitecan [17], an antibody-drug

conjugate that is FDA-approved to treat metastatic triple-negative breast, HR-positive breast, and

urothelial cancers. The existence of this approved drug may provide a basis for expedited anti-

Trop2 CAR development for patients with HNSC and other solid tumors.

Next, *DSG3* both improved tumor selectivity (AUC=0.93, FDR p=0.065) *and* safety (Tabula

sapiens score=9.13; HPA score=1.79) over the leading extant targets in HNSC. This gene

encodes the transmembrane glycoprotein desmoglein-3, which belongs to both the desmoglein

family and the cadherin cell adhesion molecule group. *DSG3* median gene expression levels

were the highest in HNSC among all TCGA cancer types (median=155 FPKM), and its encoded

protein was detected in two of four HNSC patients via membranous and/or cytoplasmic antibody

staining in the HPA Pathology Atlas. Additionally, *DSG3* was previously found to be

overexpressed at both the RNA and protein levels in head and neck cancer [18]. Inactivating

mutations in the *DSG3* gene can lead to the formation of acantholytic blisters in the oral and

laryngeal mucosa [19]. Importantly, the autoimmune disease pemphigus vulgaris is caused by

autoantibodies targeting desmoglein-3 [20], resulting in blistering and erosions of the skin and

mucosae (accordingly, desmoglein-3 is also known as "pemphigus vulgaris antigen"). Given this

critical caveat, we caution that developing CARs against all DSG3-expressing cells is likely

risky, and that a better strategy is likely to target only DSG3-overexpressing cells.

CDH3 improved upon both the leading clinical tumor selectivity and safety scores in HNSC

(AUC=0.92, FDR p-value=0.065; TB safety=9.21; HPA safety=1.58). It encodes the cadherin-3

protein, which is a member of the cadherin superfamily, cell adhesion proteins that are dependent

on calcium and enable cells to stick together by binding to each other. *CDH3* exhibited the

highest median RNA expression in HNSC among all cancer types in the TCGA (median=90.9

FPKM). Moreover, HPA Pathology antibody staining revealed all four out of four HNSC

patients exhibited medium/high CDH3 protein expression. Previous studies have reported *CDH3*

overexpression and association with poor prognosis in HNSC [21–23]. Biallelic mutations in

CDH3 can cause recessive congenital hypotrichosis with juvenile macular dystrophy [24] a rare

disorder characterized by sparse hair and progressive vision loss, as well as ectodermal

dysplasia, ectrodactyly, and macular dystrophy syndrome, similar to the former but additionally

causing congenital limb malformation [25].

DSC3, another cadherin gene, improves upon both tumor selectivity and safety in HNSC

(AUC=0.91, FDR p=0.065; TS safety=8.96; HPA safety=1.68) and encodes the desmocollin-3

protein. As a desmosomal protein, it is mostly present in epithelial cells and is essential for the

formation and adherence of desmosome cell-cell junctions. In the HPA Pathology Atlas, *DSC3* is

"cancer enhanced" in head and neck cancer and cervical cancer, with highest median expression

in HNSC (94.5 FPKM) out of all cancer types. Moderate/strong cytoplasmic positivity was seen

in HNSC, while DSC3 was generally lowly expressed in most other cancers [15]. While studies

on this protein's role in HNSC have been limited to the best of our knowledge, reduced

expression of DSC3 in oral squamous carcinomas has been previously correlated with

histological grade [26]. Biallelic inactivating mutations in *DSC3* cause skin blistering and

hypotrichosis [27].

Next, *PKP3*, encoding the protein plakophilin-3, improves upon both the leading clinical

tumor selectivity and safety scores for HNSC (AUC=0.89, FDR p=0.065; TB safety=8.67; HPA

safety=1.55). It belongs to the plakophilin family of desmosomal proteins and links the

desmosomal plaque to cytoskeletal intermediate filaments. While detected in many TCGA

cancer types, it displayed highest median expression in HNSC (82.8 FPKM). Moderate to high

cytoplasmic immunoreactivity of this protein was yielded by the majority of HNSC cancers

stained in the HPA Pathology Atlas. Intriguingly, while the upregulation of *PKP3* has been

observed during development of many cancer types, its potential tumor suppressive role was also

reported in HNSC, with its expression prognostically favorable, denoting a paradoxical role of it

and other desmosomal proteins in HNSC [28]. There are no known monogenic disorders caused

by mutations in *PKP3*.

The next candidate target, *COL17A1*, encodes the collagen type XVII alpha 1 chain. This

gene has high tumor selectivity and safety scores for HNSC (AUC=0.88, FDR p=0.084; TS

safety=9.13; HPA safety=1.89). The encoded protein is part of hemidesmosomes, which are

protein complexes mediating the adhesion of the dermis and epidermis. The HPA Pathology

Atlas reports *COL17A1* as specifically "cancer enriched" in HNSC (164.9 FPKM), while its

expression was detected in many cancer types. Moderate to strong cytoplasmic and membranous

staining of the encoded protein was observed in HNSC samples. Collagen XVII has been found

to frequently present in HNSCs, especially those with an oral cavity or larynx origin(29).

Biallelic inactivating mutations in *COL17A1* cause intermediate junctional epidermolysis bullosa

4 [30], a genetic disorder characterized by skin blisters and erosions, nail dystrophy, and

alopecia. *COL17A1* mutations can also cause epithelial current erosion dystrophy, a genetic

disorder characterized by corneal erosion [31].

MAGEA4 has higher safety scores in HNSC (TS safety=10; HPA safety=1.95). It encodes the

melanoma-associated antigen 4, which regulates cell proliferation by stopping cell cycle arrest at

G1 phase and decreases apoptosis mediated by p53. Importantly, the MAGE-A4 protein is

already an extant CAR-T target against non-small cell lung cancer (NSCLC) [10] but has not

been studied in HNSC patients. Thus, repurposing CAR therapies targeting this protein to

patients with MAGEA4+ HNSC may be low cost and high yield. HPA Pathology Atlas defines

MAGEA4 to be "cancer enhanced" in lung cancer in the TCGA, with comparable expression in

head and neck cancer. Medium to high protein expression was observed in HNSC samples by

antibody staining. Its widespread expression has been previously observed in a variety of head

and neck cancers [32–36]. There are no known monogenic disorders caused by mutations in

MAGEA4.

Row	Cancer Type	Surface protein	Encoding gene	Protein family/group(s)
1	HNSC	Tumor Associated Calcium Signal Transducer 2; Trop-2; epithelial glycoprotein-1 antigen (EGP-1)	*TACSTD2*	Epithelial cell adhesion molecule family; transmembrane glycoprotein

2	**HNSC**	Desmoglein-3	*DSG3*	Desmoglein family; cadherin superfamily; transmembrane glycoprotein
3	**HNSC**	Cadherin-3	*CDH3*	Classical cadherin; cadherin superfamily
4	**HNSC**	Glucose transporter 1; solute carrier family 2	*SLC2A1*	Solute carrier 2A family
5	**HNSC**	Desmocollin 3	*DSC3*	Desmosomal cadherin; cadherin superfamily
6	**HNSC**	Plakophilin 3	*PKP3*	Plakophilin family; arm-repeat (armadillo) family; desmosomal protein
7	**HNSC**	Collagen type XVII alpha 1 chain	*COL17A1*	Collagen family; hemidesmosomal protein
8	**HNSC, GBM**	Protein Tyrosine Phosphatase Receptor Type Z1	*PTPRZ1*	Receptor-type tyrosine-protein phosphatase family
9	**HNSC**	Melanoma-associated antigen 4	*MAGEA4*	MAGE family of tumor-associated antigens
10	**HNSC**	FAT Atypical Cadherin 2	*FAT2*	Protocadherin; FAT family
11	**HNSC**	Chloride Channel Accessory 2	*CLCA2*	Calcium-activated chloride channel (CLCA) family
12	**HNSC**	Cholinergic Receptor Muscarinic 3; M3 Muscarinic receptor	*CHRM3*	Muscarinic acetylcholine receptors family
13	**HNSC**	Solute carrier family 2 member 9	*SLC2A9*	Solute carrier 2A family

14	**HNSC**	Cadherin-13	*CDH13*	Non-classical cadherin; cadherin superfamily
15	**HNSC**	Hydroxycarboxylic Acid Receptor 2	*HCAR2*	Hydroxycarboxylic acid receptor family; G-protein coupled receptor 1 family
16	**HNSC**	Immunoglobulin Superfamily Member 3	*IGSF3*	Immunoglobulin superfamily
17	**HNSC**	Podoplanin	*PDPN*	Podoplanin family; transmembrane receptor glycoprotein
18	**HNSC**	LY6/PLAUR Domain Containing 3	*LYPD3*	LY6/PLAUR domain-containing family; GPI-anchored metastasis associated protein
19	**HNSC**	Cluster of differentiation 109	*CD109*	Alpha-2-macroglobulin family
20	**HNSC**	Solute Carrier Family 7 Member 8	*SLC7A8*	Solute carrier 7A family

617
618 **Table 3.** Families/classes of the twenty surface proteins encoded by genes resulting from our
619 analysis as candidate CAR targets.
620

We additionally explored the *in vitro* dependency by tumors on genes encoding known CAR

targets (to our knowledge, the first systematic interrogation) and our newly uncovered candidate

CAR targets. The strong essentiality of *CD19* in a significant fraction of B-cell lymphomas and

leukemias is unsurprising given that CD19 is a B-cell marker and has important function in B

cell receptor signaling. However, other blood tumor clinical CAR-T target genes, *CD22* and

TNFRSF17, are not essential in any cell line of their respective indications. In solid tumors, the

high essentiality of *EGFR* and *ERBB2* may suggest that these genes are highly important to the

viability of cancer cells in their respective solid tumor indications. The essentiality of *IGSF3* and

SLC2A1 in a significant fraction of HNSC cell lines may indicate their importance to the

viability of HNSC tumor cells. Consequently, the ability of these cancer cells to downregulate

IGSF3 and *SLC2A1* to evade CAR cell killing might be limited.

In closing, we note that our findings are aligned with those reported in a recent study that

has characterized the dependence of the genes constituting the surfaceome in cancer cell line

growth and found that the percentage of surfaceome genes defined as essential in DepMap

"[was] substantially less than" the percentage of non-surfaceome genes defined as essential; only

4.1% of the former were essential versus 14.0% of the latter [7]. We note, however, that DepMap

quantified *in vitro* gene essentiality, whose relation to *in vivo* essentiality observed in patients'

tumors remains to be further elucidated.

The key finding of our study is a set of genes encoding cell surface proteins that may

constitute promising new targets for engineering new CARs, mostly to treat HNSC and to a

lesser extent GBM tumors. The targets we found have higher tumor selectivity and safety scores

versus those of leading extant targets in clinics. Identified primarily by the analysis of tumors

expression at the single cell level, most of our proposed targets have additional supportive

evidence from human reference atlases, TCGA expression, pathology staining and previous

gene-specific publications. Our analysis of dependency scores in the DepMap illuminates the

intriguing phenomenon that the vast majority clinical CAR targets are not biologically essential

to cancers, except for: *CD19*, *EGFR*, *ERBB2*, and *MET*. Of the targets yielded by our analysis,

IGSF3 and *SLC2A1* exhibited strong essentiality in HNSC cell lines.

Our study has a few limitations. First and foremost, we have used gene expression as a

proxy for protein expression, but it is ultimately proteins that are targeted by CARs. When

advances in high-throughput single-cell proteomics technology come to the forefront in the

future, it will be pertinent to apply the approach presented in this study here to evaluate the

discriminatory potential of membrane protein receptors as viable CAR targets. In the meantime,

with the accumulating plethora of scRNA-seq datasets of tumors, gene expression is the best

representation of cell genomics available to us. Second, the use of the AUC metric to evaluate

target discrimination potential has limitations. In practice, optimal discrimination relies on

knowing antigen detection thresholds of CAR-T cells. Currently, our control in setting antigen

detection density of CARs is poor, but emerging efforts may allow us to tune antigen detection

cutoffs[5]. This would allow us to evaluate CAR target viability more practically. Finally, in future

studies, it would be of great interest to evaluate dual- and multi-targeted CARs, based on the

approach outlined herewith. The computational framework enabling such combinatorial

investigations has already been laid out in MadHitter [6], but for simplicity and practical

applicability, we aimed to focus here on finding optimal single CAR surfaceome targets.

In summary, employing a systematic, data-driven analysis, we first chart the landscape of

existing CAR targets in the clinic, identifying the leading targets in each cancer type based on

tumor selectivity and safety metrics that we have computed. Next, from single cell

transcriptomics data, we have performed a genome wide search across many different cancer

types to identify new and candidate CAR targets with both better selective and safety scores than

those of extant clinical targets. Remarkably, in almost all cancer types, we could not find such

better targets, testifying to the near optimality of the current target space, at least in accordance

with our measures. However, one striking exception is HPV-negative HNSC, for which there is a

dearth of existing targets in clinical trials. Still, bearing that in mind, specifically, our

investigation has discovered 20 new and untested CAR targets for treating HNSC and GBM

tumors more precisely and safely than extant targets, which are each followed up with in a

thorough literature survey and an analysis of their *in vitro* indication specific essentiality. Five of

these 20 have selectivity and safety scores that are both higher than those computed for the

existing CAR targets, whose further investigation may lead to better CAR treatments in HNSC.

Additional files

Additional File 1 (.txt)

Supp Text: Literature review on candidate targets (the fourteen not mentioned in the main text).

Fig S1: the distributions of safety scores in surfaceome genes versus indication-specific CAR

targets. (A) shows HPA safety scores, (B) shows Tabula Sapiens safety scores.

Fig S2: the distributions of safety scores in indication-specific CAR targets versus top 10 tumor-

selective surfaceome genes in each indication. Wilcoxon rank-sum test comparison p-values

shown at bottom of each plot. (A) shows HPA safety scores, (B) shows Tabula Sapiens safety

scores.

Additional File 2 (.xlsx)

Table S1: (gene x cancer type) cells list all the patient-specific tumor selectivity scores (aka

tumor-vs-nontumor classification AUCs) of each surfaceome gene in the respective cancer type.

'N/A' denotes genes that weren't profiled for single cell datasets pertaining to the respective

cancer type.

Table S2: (gene x tissue group) values denote levels of protein abundance measured in tissue

group, such that: ND-L < Variable < M-H. Safety scores shown in rightmost column.

Table S3: All (gene x organ) cells contain fraction of cells expressing gene in respective organ.

Overall safety score provided in rightmost column.

Table S4: Top selective (and yet safe) new targets. The first two rows (marked by *) denote the

leading clinical targets by selectivity for the two indications, HNSC and GBM, for which we

found new targets. The table is sorted by descending tumor selectivity scores.

Table S5: Top safe (and yet selective) new targets. The first row (marked by *) denotes the

leading clinical target by safety for HNSC. The second row (marked by **) denotes a protein

which is already a CAR target for the NSCLC indication (MacKay et al.). The table is sorted

primarily by descending Tabula Sapiens safety scores, with ties broken in order of descending

HPA safety scores.

Table S6: Dependency in cell lines of genes encoding CAR-T targets in blood tumors.

Table S7: Supplementary Table 7. Dependency in cell lines of genes encoding CAR-T targets in

solid tumors.

Table S8: Supplementary Table 8. Dependency in cell lines of each candidate target gene

yielded by our pipeline. Rows highlighted in green are genes that are classified as 'strongly

selective' or 'pan-essential' by DepMap and are considerably more essential than all other genes

in the Table.

Ethics approval and consent to participate

Not applicable.

Consent for publication

723   Not applicable.

Availability of data and materials

726   All data analyzed in this study are publicly available via their respective cited sources. TISCH

727   single-cell RNA-seq data were downloaded from http://tisch1.comp-genomics.org/. HPA data

728   were downloaded from https://www.proteinatlas.org/about/download. Tabula sapiens data were

729   downloaded from https://figshare.com/projects/Tabula_Sapiens/100973. CRISPR DepMap data

730   were downloaded from https://figshare.com/articles/dataset/DepMap_22Q2_Public/19700056/2.

Competing interests

Funding

738   This research is supported in part by the Intramural Research Program of the National Institutes

739   of Health, NCI, CCR. S.M. is supported by the NCI-UMD Partnership for Integrative Cancer

740   Research Program.

Author's contributions

AAS and ER supervised the study. SM, AAS, and ER conceived and designed the study. SM and SS analyzed data and designed the figures. SM, JSG, EEWC, AAS, and ER drafted and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The NIH HPC Biowulf cluster (http://hpc.nih.gov) was used for computational analyses.

References

1. Melenhorst JJ, Chen GM, Wang M, Porter DL, Chen C, Collins MA, et al. Decade-long leukaemia remissions with persistence of CD4+ CAR T cells. Nature. 2022 Feb;602(7897):503–9.

2. Stergiopoulos S, Getz KA, Blazynski C. Evaluating the Completeness of ClinicalTrials.gov. Ther Innov Regul Sci. 2019 May 1;53(3):307–17.

3. Milone MC, Xu J, Chen SJ, Collins MA, Zhou J, Powell DJ, et al. Engineering-enhanced CAR T cells for improved cancer therapy. Nat Cancer. 2021 Aug;2(8):780–93.

4. Jing Y, Liu Y, Li Q, Ye Y, Diao L, Huang Y, et al. Expression of chimeric antigen receptor therapy targets detected by single-cell sequencing of normal cells may contribute to off-tumor toxicity. Cancer Cell. 2021 Dec 13;39(12):1558–9.

5. Parker KR, Migliorini D, Perkey E, Yost KE, Bhaduri A, Bagga P, et al. Single-Cell Analyses Identify Brain Mural Cells Expressing CD19 as Potential Off-Tumor Targets for CAR-T Immunotherapies. Cell. 2020 Oct 1;183(1):126-142.e17.

6. Ahmadi S, Sukprasert P, Vegesna R, Sinha S, Schischlik F, Artzi N, et al. The landscape of receptor-mediated precision cancer combination therapy via a single-cell perspective. Nat Commun. 2022 Mar 25;13(1):1613.

7. Hu Z, Yuan J, Long M, Jiang J, Zhang Y, Zhang T, et al. The Cancer Surfaceome Atlas integrates genomic, functional and drug response data to identify actionable targets. Nat Cancer. 2021 Dec 13;1–17.

8. Lareau CA, Parker KR, Satpathy AT. Charting the tumor antigen maps drawn by single-cell genomics. Cancer Cell. 2021 Dec 13;39(12):1553–7.

9. Kwon J, Kang J, Jo A, Seo K, An D, Baykan MY, et al. Single-cell mapping of combinatorial
target antigens for CAR switches using logic gates. Nat Biotechnol. 2023 Feb 16;1–13.

10. MacKay M, Afshinnekoo E, Rub J, Hassan C, Khunte M, Baskaran N, et al. The therapeutic
landscape for cells engineered with chimeric antigen receptors. Nat Biotechnol. 2020
Feb;38(2):233–44.

11. Sun D, Wang J, Han Y, Dong X, Ge J, Zheng R, et al. TISCH: a comprehensive web
resource enabling interactive single-cell transcriptome visualization of tumor
microenvironment. Nucleic Acids Res. 2021 Jan 8;49(D1):D1420–30.

12. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, et al. Tissue-
based map of the human proteome. Science. 2015 Jan 23;347(6220):1260419.

13. THE TABULA SAPIENS CONSORTIUM. The Tabula Sapiens: A multiple-organ, single-
cell transcriptomic atlas of humans. Science. 2022 May 13;376(6594):eabl4896.

14. Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, et al. The
Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet. 2013 Oct;45(10):1113–20.

15. Uhlen M, Zhang C, Lee S, Sjöstedt E, Fagerberg L, Bidkhori G, et al. A pathology atlas of
the human cancer transcriptome. Science. 2017 Aug 18;357(6352):eaan2507.

16. Tsujikawa M, Kurahashi H, Tanaka T, Nishida K, Shimomura Y, Tano Y, et al.
Identification of the gene responsible for gelatinous drop-like corneal dystrophy. Nat Genet.
1999 Apr;21(4):420–3.

17. Bardia A, Hurvitz SA, Tolaney SM, Loirat D, Punie K, Oliveira M, et al. Sacituzumab
Govitecan in Metastatic Triple-Negative Breast Cancer. N Engl J Med. 2021 Apr
22;384(16):1529–41.

18. Chen YJ, Chang JT, Lee L, Wang HM, Liao CT, Chiu CC, et al. DSG3 is overexpressed in
head neck cancer and is a potential molecular target for inhibition of oncogenesis. Oncogene.
2007 Jan 18;26(3):467–76.

19. Kim JH, Kim SE, Park HS, Lee SH, Lee SE, Kim SC. A Homozygous Nonsense Mutation in
the DSG3 Gene Causes Acantholytic Blisters in the Oral and Laryngeal Mucosa. J Invest
Dermatol. 2019 May 1;139(5):1187–90.

20. Amagai M, Klaus-Kovtun V, Stanley JR. Autoantibodies against a novel epithelial cadherin
in pemphigus vulgaris, a disease of cell adhesion. Cell. 1991 Nov 29;67(5):869–77.

21. Dasgupta S, Tripathi PK, Qin H, Bhattacharya-Chatterjee M, Valentino J, Chatterjee SK.
Identification of molecular targets for immunotherapy of patients with head and neck
squamous cell carcinoma. Oral Oncol. 2006 Mar 1;42(3):306–16.

22. Keck MK, Zuo Z, Khattri A, Stricker TP, Brown CD, Imanguli M, et al. Integrative Analysis
of Head and Neck Cancer Identifies Two Biologically Distinct HPV and Three Non-HPV
Subtypes. Clin Cancer Res. 2015 Feb 16;21(4):870–81.

23. Wu T, Xiao Z, Li Y, Jiao Z, Liang X, Zhang YongQ, et al. CDH3 is associated with poor
prognosis by promoting the malignancy and chemoresistance in oral squamous cell
carcinoma. Asian J Surg. 2022 Dec 1;45(12):2651–8.

24. Sprecher E, Bergman R, Richard G, Lurie R, Shalev S, Petronius D, et al. Hypotrichosis with
juvenile macular dystrophy is caused by a mutation in CDH3, encoding P-cadherin. Nat
Genet. 2001 Oct;29(2):134–6.

25. Kjaer KW, Hansen L, Schwabe GC, Marques-de-Faria AP, Eiberg H, Mundlos S, et al.
Distinct CDH3 mutations cause ectodermal dysplasia, ectrodactyly, macular dystrophy
(EEM syndrome). J Med Genet. 2005 Apr 1;42(4):292–8.

26. Wang L, Liu T, Wang Y, Cao L, Nishioka M, Aguirre RL, et al. Altered expression of
desmocollin 3, desmoglein 3, and beta-catenin in oral squamous cell carcinoma: correlation
with lymph node metastasis and cell proliferation. Virchows Arch Int J Pathol. 2007
Nov;451(5):959–66.

27. Onoufriadis A, Ahmed N, Bessar H, Guy A, Liu L, Marantzidis A, et al. Homozygous
Nonsense Mutation in DSC3 Resulting in Skin Fragility and Hypotrichosis. J Invest
Dermatol. 2020 Jun 1;140(6):1285–8.

28. Liu YQ, Zou HY, Xie JJ, Fang WK. Paradoxical Roles of Desmosomal Components in Head
and Neck Cancer. Biomolecules. 2021 Jun 20;11(6):914.

29. Jones VA, Patel PM, Gibson FT, Cordova A, Amber KT. The Role of Collagen XVII in
Cancer: Squamous Cell Carcinoma and Beyond. Front Oncol. 2020 Mar 19;10:352.

30. Gatalica B, Pulkkinen L, Li K, Kuokkanen K, Ryynänen M, McGrath JA, et al. Cloning of
the human type XVII collagen gene (COL17A1), and detection of novel mutations in
generalized atrophic benign epidermolysis bullosa. Am J Hum Genet. 1997 Feb;60(2):352–
65.

31. Jonsson F, Byström B, Davidson AE, Backman LJ, Kellgren TG, Tuft SJ, et al. Mutations in
Collagen, Type XVII, Alpha 1 (COL17A1) Cause Epithelial Recurrent Erosion Dystrophy
(ERED). Hum Mutat. 2015;36(4):463–73.

32. Cuffel C, Rivals JP, Zaugg Y, Salvi S, Seelentag W, Speiser DE, et al. Pattern and clinical
significance of cancer-testis gene expression in head and neck squamous cell carcinoma. Int
J Cancer. 2011;128(11):2625–34.

33. Ishihara M, Kageyama S, Miyahara Y, Ishikawa T, Ueda S, Soga N, et al. MAGE-A4, NY-
ESO-1 and SAGE mRNA expression rates and co-expression relationships in solid tumours.
BMC Cancer. 2020 Jun 29;20(1):606.

34. Melo DH, Mamede RCM, Neder L, Saggioro FP, Figueiredo DLA, Araújo da Silva Jr W, et
al. Expression of mage-A4 and mage-C1 tumor-associated antigen in benign and malignant
thyroid diseases. Head Neck. 2011;33(10):1426–32.

35. Montoro JR de MC, Mamede RCM, Neder Serafini L, Saggioro FP, Figueiredo DLA, Silva
Jr WA da, et al. Expression of cancer-testis antigens MAGE-A4 and MAGE-C1 in oral
squamous cell carcinoma. Head Neck. 2012;34(8):1123–8.

36. Prasad ML, Jungbluth AA, Patel SG, Iversen K, Hoshaw-Woodard S, Busam KJ. Expression
and significance of cancer testis antigens in primary mucosal melanoma of the head and
neck. Head Neck. 2004;26(12):1053–7.

37. DepMap 22Q2 Public [Internet]. figshare; 2022 [cited 2023 Mar 27]. Available from:
https://figshare.com/articles/dataset/DepMap_22Q2_Public/19700056/2

Solid Tumor CAR-T Targets in Clinics

EGFR, IL13RA2, PSCA

EGFR, ERBB2

MUC1, PSCA, MAGEA1

PMEL

GPC3, AFP

TM4SF1, PSCA, GUCY2C

EPCAM, PSCA, GUCY2C

Surfaceome genes

Patient tumor scRNA-seq → **Selectivity**: Tumor-upregulated genes

Healthy tissue scRNA-seq and proteomics → **Safety**: Genes/proteins lowly expressed in normal tissues

CRISPR knockout screens → **Dependency**: Tumor-essential genes

Cancer-type-specific CAR targets