Bile Acid Regulates the Colonization and Dissemination of *Candida albicans* from the

Gastrointestinal Tract by Controlling Host Defense System and Microbiota

Shankar Thangamani [a, b, c]#, Ross Monasky[c], Jung Keun Lee[c], Vijay Antharam [d], Harm

HogenEsch [a, b], Tony Hazbun[e], Yan Jin[f], Haiwei Gu[f, g], Grace L. Guo[h, i]

[a]Department of Comparative Pathobiology, College of Veterinary Medicine, Purdue University,

West Lafayette, Indiana, USA

[b]Purdue Institute for Immunology, Inflammation and Infectious Diseases (PI4D), West

Lafayette, Indiana, USA

[c]College of Veterinary Medicine, Midwestern University, Glendale, AZ, USA

[d]Department of Chemistry, School of Science and Human Development, Methodist University,

Fayetteville, NC, USA

[e]Department of Medicinal Chemistry and Molecular Pharmacology, College of Pharmacy, Purdue

University, West Lafayette, IN, USA

[f]Arizona Metabolomics Laboratory, College of Health Solutions, Arizona State University,

Phoenix, AZ, USA

[g]Center for Translation Science, Department of Environmental Health Sciences, Robert Stempel

College of Public Health and Social Work, Florida International University, Port St. Lucie, FL,

USA

[h]Department of Pharmacology and Toxicology, Earnest Mario School of Pharmacy, Rutgers

University, Piscataway, NJ, USA

[i]Department of Veterans Affairs New Jersey Health Care System, East Orange, NJ, USA

Running Title: Bile Acid Regulation of *Candida albicans*

#Address correspondence to Shankar Thangamani, **sthangam@purdue.edu**

Abstract

Candida albicans (**CA**), a commensal and opportunistic eukaryotic organism, frequently inhabits

the gastrointestinal (GI) tract and causes life-threatening infections. Antibiotic-induced gut

dysbiosis is a major risk factor for increased CA colonization and dissemination from the GI tract.

We identified a significant increase of taurocholic acid **(TCA),** a major bile acid in antibiotic-

treated mice susceptible to CA infection. *In vivo* findings indicate that administration of TCA

through drinking water is sufficient to induce colonization and dissemination of CA in wild type

and immunosuppressed mice. Treatment with TCA significantly reduced mRNA expression of

immune genes *ang4* and *Cxcr3* in the colon. In addition, TCA significantly decreased the relative

abundance of three culturable species of commensal bacteria, *Turicibacter sanguinis*,

Lactobacillus johnsonii, and *Clostridium celatum,* in both cecal contents and mucosal scrapings

from colon. Taken together, our results indicate that TCA promotes fungal colonization and

dissemination of CA from the GI tract by controlling host defense system and intestinal microbiota

that play a critical role in regulating CA in the intestine.

Importance

Broad-spectrum antibiotics, FDA-approved bile acid drugs, and probiotics used to control

metabolic and infectious diseases profoundly alter the level of TCA in the gut. Furthermore, TCA

level is highly altered in a subset of cancer, colitis and surgery patients who are highly susceptible

to CA infection. Inadvertently, these therapies and disease conditions could be either promoting

CA colonization and dissemination. Our findings indicate that TCA alone can induce fungal

colonization and dissemination from the intestine. Results from this study will have a significant

impact in understanding how bile acids interact with the microbiota and host in regulating

49   invasive fungal infections that originate from the intestine and to develop potential new

50   antifungal therapeutics.

Introduction

Candida albicans (**CA**), a commensal and opportunistic eukaryotic organism, frequently inhabits

the gastrointestinal (GI) tract and can cause life-threatening infections [1-3]. CA is present in

small numbers in the healthy GI tract of humans; thus, CA is harmless in immunocompetent

humans [4, 5]. While dysregulation in host defense system contribute to invasive CA infections,

use of broad-spectrum antibiotics is a major predisposing risk factor for increased fungal

colonization and subsequent dissemination from the intestine [6-13]. CA is normally absent in the

GI tract of adult mice; however, antibiotic treatment leads to CA colonization and dissemination

in mice and closely resembles CA infection in human patients [4, 5, 13-21]. Because colonization

of the GI tract is necessary for dissemination of CA and CA colonization resistance is nullified by

antibiotic treatment, identifying the factors that play a critical role in CA colonization may pave

the way to develop novel approaches to limit CA dissemination [5, 19].

Antibiotics alter the intestinal microbiota and lead to changes in the composition of the gut

microbial metabolites that play a critical role in controlling several enteric bacterial pathogens [22-

24]. However, there is limited knowledge about the potential role of microbial metabolites in the

regulation of CA colonization and pathogenesis. To address this gap in knowledge, we previously

performed targeted metabolomics and identified several groups of metabolites that are

differentially regulated in the gut contents of cefoperazone-treated CA-susceptible mice and

control CA-resistant mice [25]**.** A bile acid, taurocholic acid (TCA), was identified as one of the

major class of metabolites that was significantly increased in the antibiotic-treated mice [25]**.**

Following synthesis in the liver, TCA enters the intestine and undergoes two major chemical

modifications (*deconjugation and dihydroxylation*) carried out by gut microbes [26-28]. Antibiotic

treatment depletes TCA-metabolizing commensal bacteria leading to increased levels of TCA in

the gut contents [25]. Furthermore, the gut concentration of TCA is considerably altered in

different diseases. Specifically, TCA is highly upregulated in immunocompromised cancer

patients [29-31], drug-induced liver injury [32-34], high fat diets [35], and liver cirrhosis [36-41],

conditions that are associated with severe morbidity and mortality caused by CA. Numerous FDA-

approved drugs and probiotics currently used to treat various diseases also profoundly alter bile

acid levels including TCA in the gut [42-51]. Therefore, understanding the role of TCA in the

regulation of CA will gain insights into the bile-mediated regulation of gut fungi and will form a

platform to develop novel therapies to control and treat invasive CA infections that arise as a result

of antibiotic-induced dysbiosis and to control CA-associated infections in individuals with liver

cirrhosis. Therefore, in this study, we investigated the role of TCA on CA colonization and

dissemination from the intestine. Our findings indicate that TCA promotes fungal colonization and

dissemination by altering intestinal defense system and microbiota.

Materials and Methods

Strains and Reagents. *Candida albicans* SC5314 was provided by Dr. Andrew Koh at the

University of Texas Southwestern Medical Center [52]. This study used Yeast Peptone Dextrose

(YPD) (242810, BD Difco) agar (BP1423, Fisher) as well as Bacteroides Bile Esculin Agar (BBE)

(AS114, Anaerobe Systems) for various assays. Additionally, antibiotics kanamycin (J61272, Alfa

Aesar), ampicillin (69-52-3, IBI Scientific), and streptomycin (S6501, Sigma-Aldrich) were used

in media for fungal enumeration and cefoperazone (J65185, Alfa Aesar) was used in antibiotic

pre-treatment. Taurocholic acid (TCA) (16215, Cayman Chemicals) and cyclophosphamide

(PHR1404, Millipore Sigma) were purchased for these studies. Additional kits and reagents used

included: hydrogen sulfide gas assay (ZAN-5084, Cell Biolabs), fluorescein isothiocyanate-

dextran (FITC-dextran) (FD4, Millipore Sigma), taurine (A12403, Alfa Aesar), and sodium

formate (A17813, Alfa Aesar).

Mouse studies. Male and female C57BL/6J mice (000664) and BALB/cJ mice (000651) of ages

6-12 weeks were purchased from the Jackson Laboratory. All animal studies conducted were

approved by the Institutional Animal Care and Use Committee (IACUC) at Midwestern University

under MWU IACUC Protocol #4014.

Fungal colonization and dissemination in the immunosuppressed mouse model. Studies

specifically examining survival and dissemination of mice infected with *C. albicans* SC5314 under

immunosuppressive conditions were performed with BALB/c mice. Male and female mice (age 8

to 12 weeks old) were treated with or without cefoperazone (0.5mg/mL) via drinking water, and

drinking water was replaced with fresh antibiotic water every two days. After 7 days, all mice were

infected with *Candida albicans* SC5314 via oral gavage with 1-2 x 10^8 CFU/mouse. Three days

post infection, all mice were treated with 150 mg of cyclophosphamide (cyclo) per kilogram of

mouse body weight by intraperitoneal injection. A second and third dose of cyclophosphamide

were given again at 5- and 7-days post infection, respectively. Mice were monitored for survival,

and euthanized if mice were moribund. After euthanasia, cecal contents, liver, and kidneys were

collected and processed for fungal burden to assess the fungal dissemination as described before

[25]. Briefly, gut contents and tissue homogenates were serially diluted and plated onto YPD agar

plates containing broad spectrum antibiotics and incubated at 30°C for 24 hours to determine the

colony forming unites (CFU).

Effect of TCA on fungal colonization and dissemination in mice. Male and female wild type

C57B/6 mice were infected with *C. albicans* SC5314 via oral gavage with 1-2 x 10^8 CFU/mouse.

One group of mice was treated with 1% TCA in the drinking water and the water was changed

every day. Control groups received sterile water. Fecal contents were collected at the indicated

time points to determine the fungal load in feces. Mice were monitered for survival and cecal

contents, liver, and kidneys were processed to determine the fungal load and dissemination as

described before [25]. To determine the effect of TCA on fungal colonization and dissemination

in the immunosuppressed mouse model, male and female BALB/cJ mice (age 10 to 12 weeks old)

were infected as described above and all mice were treated with 150 mg of cyclophosphamide

(cyclo) per kilogram of mouse body weight through intraperitoneal injection. Second and third

dose of cyclophosphamide were given again at 5- and 7-days post-infection respectively.

FITC-Dextran permeability assay. Gut permeability was measured in infected mice using a

FITC-dextran assay. Wild type C57BL/6 mice were infected with *Candida albicans* SC5314 via

oral gavage with 1-2 x 10^8 CFU/mouse and treated with or without 1% TCA through drinking

water. After 10 days post-infection, mice were administered orally 150 uL PBS containing 15mg

FITC-dextran. The mice were anesthetized four hours later and blood was collected via retro-

orbital route. Blood samples were then centrifuged to collect serum. Serum samples were

processed via a 2-fold serial dilution in a 96-well plate and fluorescence was measure via plate

reader (excitation: 485nm; emission: 520nm). Standard curves were made using serially diluted

FITC-Dextran in PBS to determine the FITC-Dextran levels in serum samples.

RNA sequencing and analysis. Wild type C57BL/6 mice were infected with *C. albicans* SC5314

via oral gavage with 1-2 x 10^8 CFU/mouse and treated with or without 1% TCA through drinking

water. At 10 days post-infection, colon was collected and flash frozen in liquid nitrogen. Total

RNA was collected using Zymo Research RNA kit as per the manufacturer's instructions. Total

RNA-Seq libraries were constructed from 500 ng of total RNA and rRNA was removed. Libraries

were prepared using the Zymo-Seq RiboFree Total RNA Library Prep Kit (Cat # R3000) according

to the manufacturer's instructions. RNA-Seq libraries were sequenced on an Illumina NovaSeq to

a sequencing depth of at least 30 million read pairs (150 bp paired-end sequencing) per sample.

Sequence Data Alignments and Differential Expression Analysis: NovaSeq paired-end 150-bp

reads from Total RNA-Seq data files were first adaptor trimmed, and then analyzed using the

STAR program (version 2.6.1d) for alignment of short reads to the genome of interest. Transcript

counts were inferred from alignment files. All transcripts with either 0 or 1 counts were removed.

Gene expression was measured using EdgeR. The DESeq2 R package was used for differential

expression analysis using the gene feature.

Microbiome sequencing and analysis. Wild type C57BL/6 mice were infected with *C. albicans*

SC5314 via oral gavage with 1-2 x 10^8 CFU/mouse and treated with or without 1% TCA through

drinking water. After 10 days post-infection, cecal contents and mucosal scrapings from colon was

collected using glass slides to determine the microbiome composition. Microbiome sequencing

was carried out at Zymo Research corporation (Irvine, CA). The ZymoBIOMICS® Microbial

Community Standard (Zymo Research, Irvine, CA) was used as a positive control for each DNA

extraction, if performed. The final library was sequenced on Illumina® MiSeqTM with a v3

reagent kit (600 cycles) and the relative abundance of bacteria were analyzed as described

elsewhere [25] .

Immunofluorescence staining. Colon from untreated and TCA treated mice infected with CA

were collected after 10-12 days of infection. Colon tissue was fixed in 10% neutral buffered

formalin solution and sectioned at 5 µm, and stained with DAPI and ZO-1 Antibody (21773-1-

AP, Thermofischer). Stained tissues were imaged (40X) using a Keyence BZ-X700 microscope.

Metabolomics. Bile acid metabolites (TCA and DCA) levels in the cecal contents were

determined as described previously [25]**.**

Statistical analysis. Statistical analyses were performed using GraphPad Prism 6.0 (Graph Pad

Software, La Jolla, CA). *P* values were calculated using the Mann-Whitney U test or unpaired

Student's t- test or one way-ANOVA followed by Bonferroni comparison or Kaplan-Meier (log

rank) survival test as indicated. *P* values of (*$\leq$ 0.05) (**P$\leq$ 0.01) (***P$\leq$ 0.001) (****P$\leq$ 0.0001)

were considered as significant.

Results

TCA is the major bile acid metabolite up-regulated in the cefoperazone-treated mice

susceptible to CA infection. To identify the metabolites that regulate CA colonization and

dissemination, we treated BALB/c mice with or without a broad-spectrum antibiotic

(cefoperazone), orally infected them with CA, and then injected them with cyclophosphamide

intraperitoneally. As expected, all mice in the antibiotic-treated group died within 6 days of

infection (**Fig. 1A**). Furthermore, the fungal load was significantly higher in the cecal contents and

liver of antibiotic-treated mice, thus confirming increased CA colonization and dissemination from

the GI tract of antibiotic-treated mice compared to control groups (**Fig. 1B**). Previous findings

from our lab indicates that more than 200 metabolites were differentially regulated in the cecal

contents of control and antibiotic-treated mice [25]. Bile acids are one of the major metabolites

differentially altered in the CA-susceptible mice compared to control groups [25]. Among bile acid

metabolites, our results indicate that six bile acids were significantly upregulated in the antibiotic-

treated CA susceptible mice (**Fig. 1C-D**). Interestingly, only the abundance of TCA was

significantly higher than all other bile acid metabolites that were increased in the antibiotic-treated

CA susceptible mice compared to untreated control groups (**Fig. 1D**). These results indicate that

TCA is the highly upregulated bile acid in the antibiotic-treated CA- susceptible mice compared

to control groups.

TCA alone induces fungal colonization and dissemination from the GI tract in the absence

of antibiotics and immunosuppressive agents. To dissect the specific role of TCA in regulating

CA colonization and dissemination *in vivo*, C57BL/6J mice infected with CA were treated orally

with or without 1% TCA in the drinking water (**Fig. 2A)**. The fungal load in stool samples and

mortality were monitored. Mice treated with TCA alone died 15 days post-infection (**Fig. 2B)**.

Fungal load in the feces of the TCA group was significantly higher starting day 4 post-infection

compared to the untreated control groups (**Fig. 2C**). Furthermore, the TCA-treated group had

significantly higher fungal loads in the cecal content, liver, and kidney, whereas no CA was

detected in the liver and kidney from the untreated mice (**Fig. 2D)**. These results indicate that TCA

alone can induce fungal colonization and dissemination from the GI tract in the absence of

antibiotics and (or) immunosuppression.

TCA induces fungal colonization and dissemination from the GI tract of immunosuppressed

mice in the absence of antibiotic treatment. Since antibiotic induced gut dysbiosis is necessary

to induce CA colonization and dissemination in immunosuppressed mice (**Fig. 1A**) [5], we tested

whether administration of 1% TCA in the drinking water in immunosuppressed mice can promote

fungal colonization and dissemination **(Fig. 3A)**. Interestingly, BALB/c mice that received TCA

started to die 6 days post-cyclophosphamide injection and all the mice were dead by day 9 after

the first dose of cyclophosphamide. On the other hand, the mice that were infected with CA and

received three doses of cyclophosphamide survived **(Fig. 3B).** The fungal load was significantly

increased in cecal contents and liver of the TCA treated group compared to immunosuppressed

mice that received only sterile drinking water **(Fig. 3C)**. These results suggest that an increased

concentration of TCA in the intestine promotes fungal colonization, dissemination and mortality

even in the absence of antibiotic treatment in immunosuppressed mice.

TCA enhanced intestinal permeability and reduced expression of a tight junction protein.

The intestinal barrier function is critical to prevent enteric pathogens from disseminating from the

GI tract to systemic organs [5, 53]. Since TCA treatment induced CA dissemination from the GI

tract, we evaluated if TCA dysregulated the intestinal barrier function with a FITC-Dextran

permeability assay. The TCA-treated group had significantly increased levels of FITC-Dextran in

the blood compared to the untreated control groups (**Fig. 4A**). Furthermore, expression of the ZO-

1 tight junction protein was decreased in the colon of TCA-treated mice (**Fig. 4B**). Altogether,

these results indicate that TCA dysregulates the intestinal barrier function which may contribute

to CA dissemination from the intestinal tract.

TCA downregulates *ang4* and *Cx3cr1* expression in the colon tissue. CA can cause severe

invasive disease and mortality once critical components of the host defense system are

compromised [5, 53-56]. To identify if TCA dysregulates the intestinal host defense system to

induce CA colonization and dissemination, colonic tissue from untreated and TCA-treated mice

collected after 10 days of CA infection and treatment were RNA sequenced to examine the

expression of host defense genes (**Fig. 5A**). Among the host defense genes examined, the relative

expression of angiogenin 4 (*ang4*), an antimicrobial peptide, and CX3CR1 (*Cx3cr1*), a chemokine

receptor, was significantly downregulated in TCA-treated mice compared to untreated mice

infected with CA (**Fig. 5A and 5B**). Bile acids receptors play an important role in bile acid

signaling [49, 57-61]. Further, our findings indicate that among the receptors examined, the

relative expression of *Slc10a2* and *Nr1h4* that encode apical sodium-dependent bile acid

transporter (ASBT) and farnesoid X receptor (FXR) receptors, respectively, was increased in the

TCA-treated mice compared to untreated groups infected with CA (**Fig. 5C**).

TCA alters microbial composition in both luminal and mucosal parts of the GI tract. Next,

we examined if TCA changes the composition of intestinal microbiota, as commensal bacteria are

also essential to provide colonization resistance to CA [5, 25]. We analyzed the intestinal

microbiota composition from the cecal contents and mucosal scrapings from colon in the untreated

control and TCA treated mice infected with CA. Our results revealed that several culturable and

unculturable bacterial members were significantly altered in cecal contents and colon mucosal

scrapings in both groups (**Supplementary Figures 1-4**, **Fig. 6A and 6B**). Out of the top 20

statistically significant culturable and unculturable bacterial members found in cecal contents and

colon mucosal scrapings, we focused on culturable bacterial members whose 16s reads could be

mapped at species-level resolution. The relative abundance of three culturable bacterial members:

Turicibacter sanguinis (cecum: 5.2% versus 0.90%; colonic mucosa: 5.2% versus 0.65%),

Lactobacillus johnsonii (cecum: 26.4% versus 8.0%; colonic mucosa: 24.6% versus 6.8%), and

Clostridium celatum (cecum: 5.3% versus 0.76%; colonic mucosa: 4.3% versus 0.44%) were

significantly decreased in both cecal content and colon scrapings from TCA-treated mice

compared to untreated mice (**Fig. 6C and 6D**). In mucosal scrapings, *L. johnsonii* constituted close

to 25% of 16s reads but was three times more depleted in mice fed with TCA-infused drinking

water. Comparatively, the diminishment as a result of TCA was 6-7-fold for reads mapping to *T.*

sanguinis and *C. celatum*. Taken together, the relative abundance of three culturable bacterial

members (*T. sanguinis*, *L. johnsonii*, and *C. celatum*) were significantly decreased in both cecal

content and colon scrapings from TCA-treated mice compared to untreated mice infected with CA.

Discussion

Our findings indicate that TCA is one of the most strongly upregulated major bile acids in the

antibiotic-treated mice susceptible to CA infection. After synthesis in the liver, TCA enters the

small intestine and undergoes deconjugation and dihydroxylation, which are carried out by bile

salt hydrolases and bile acid inducible enzymes present in gut microbes. These two chemical

modifications convert TCA to DCA in the cecum [47, 57, 62-67]. Treatment with antibiotics

decreases the microbial metabolism of TCA and induces an increase of this bile acid. Treatment

of wild type and immunosuppressed mice with TCA alone promoted fungal colonization and

dissemination from the GI tract in the absence of antibiotic treatment. Our studies revealed three

primary mechanisms that underlie the TCA-induced enhanced CA colonization and dissemination

from the GI tract including: (i) deficiencies in host defense mechanisms, (ii) disruption of the

commensal microbiota which permits intestinal overgrowth of CA and (iii) intestinal barrier

dysfunction [5, 53].

TCA significantly decreased the expression of *ang4* and *Cx3cr1* in the colon. Interestingly, both

angiogenin-4 and CX3CR1 are involved in antifungal defense [55, 68]. Mice have three

orthologous *ang* antimicrobial peptide genes (*ang* 1, 3, and 4). However, only *ang4* is highly

expressed in the colon and small intestine and is produced by goblet and Paneth cells [68-70].

Humans only have one orthologous protein (ANG). Both *ang4* and ANG exhibit antifungal activity

in vitro [68]. Similarly, CX3CR1 mainly expressed by macrophages in the intestine plays a critical

role in antifungal defense to CA [55, 71]. Furthermore, recent evidence indicates that CX3CR1

expressing macrophages are important to maintain intestinal barrier function to prevent bacterial

translocation from the gut [72]. Mice dosed with 1% TCA orally had significantly decreased

abundance of *L. johnsonii, T. sanguinis*, and *C. celatum* in the cecal contents and colon mucosal

scrapings. Interestingly, previous findings from our lab indicates that all these three bacterial

genera were also decreased in antibiotic-treated mice susceptible to CA [25]. Commensal bacteria

such as *L. johnsonii* plays a critical role in maintaining the intestinal barrier function [73-76] and

can directly inhibit the growth of fungi [77]. Since *ang4* and *Cx3cr1* expression is also regulated

by the microbiota [68, 78], it is possible that TCA indirectly regulates CA by controlling *ang4* and

Cx3cr1 expression through one of the three commensal bacteria modulated by TCA.

In the intestine, TCA is mainly absorbed through apical sodium-dependent bile acid transporter

(ASBT), encoded by the *Slc10a2* gene in ileal and colonic epithelial cells [49, 57, 58]. After

absorption, TCA binds to FXR, a nuclear receptor encoded by *Nr1h4*, expressed in enterocytes

and macrophages resulting in the transcription of target genes [49, 57, 59-61]. The expression of

Slc10a2 and *Nr1h4* was also significantly increased in the colon of antibiotic treated mice [49, 79].

Therefore, these findings along with others [55, 68, 71, 72] suggest that TCA may induces CA

colonization and dissemination by controlling intestinal defense system and barrier function

through ASBT and FXR. Future studies to understand how TCA impacts these factors in regulating

CA will form a strong platform to reduce fungal colonization and dissemination from the GI tract.

Under normal physiological conditions, intestine and cecum holds a bile salt concentration

gradient ranging from 2% and 0.05% [80, 81], therefore we used an average concentration of 1%

TCA for mouse studies. Therefore, future studies to understand the effect of TCA at different

concentrations and dose will gain insights into the role of TCA on CA colonization and

dissemination. For example, a lower concentration (0.1 or 0.5%) may affect only microbiota or

host and potentially can enhance only CA colonization without dissemination. Further, in order to

rule the possibility that the metabolic transformation of orally administered TCA to DCA may

contribute to host and microbiota alterations observed in our study, we also examined the DCA

levels in TCA treated mice. Surprisingly, mice that received TCA had considerably reduced levels

of DCA compared to untreated groups (Supplementary Figure 5). This may be because TCA may

directly inhibit certain species of bile-metabolizing bacteria (*Lactobacillus*). Previous findings

from our lab indicates that TCA promotes, whereas DCA inhibits, CA growth *in vitro* [25, 82, 83].

However, surprisingly our *in vivo* findings indicate that orally administration with TCA alone in

mice induces fungal colonization and dissemination suggesting that regulating host and microbiota

is important in order to cause intestinal barrier dysfunction leading to CA dissemination and

mortality [5, 53, 74, 84]. In addition to host and microbiota effect, TCA and (or) DCA may have

a direct effect on CA. Future studies warrants further investigation on the direct effect of TCA and

DCA on CA, and the *in vivo* role of DCA on fungal colonization and dissemination.

Given the significant alteration in bile acid levels during various therapies and diseases [29-51,

85-106], understanding how bile acids regulate CA will provide a new approach for to manipulate

and control the fungal-bile communication system in favor of the host. Furthermore, CA in the

gut plays an active role in inflammatory bowel disease [95, 116, 118], the microbiota-gut-brain

axis [110, 119], and treating various pathogens including *Clostridium difficile* [120, 121]. Our

findings will have a significant impact in understanding how microbial metabolites interact with

the microbiota and host in regulating gut fungi.

Given the high morbidity and mortality associated with invasive fungal infections, the limited

arsenal of antifungals, and the continuing emergence of treatment-resistant strains, new approaches

to treating and preventing invasive fungal infections in patients are desperately needed. Though

TCA and DCA are well-known metabolites, their specific role in the colonization and pathogenesis

of CA is unknown. Addressing this critical gap in our knowledge could pave the way to use these

naturally occurring bile acid metabolites to reduce fungal colonization and ultimately decrease

dissemination in the host. In depth understanding of the role of bile acid metabolites on fungal

colonization and dissemination will form a strong platform to modulate bile acids or bile-acid

metabolizing commensal bacteria or to modulate the host defense system through bile acids to

prevent and treat invasive fungal infections originating from the GI tract.

References

1.   Angarone M: **Fungal infections in cancer patients**. *Cancer Treat Res* 2014, **161**:129-155.
2.   Low CY, Rotstein C: **Emerging fungal infections in immunocompromised patients**.
354   *F1000 Med Rep* 2011, **3**:14.
355   3.   Perfect JR, Hachem R, Wingard JR: **Update on epidemiology of and preventive**
356   **strategies for invasive fungal infections in cancer patients**. *Clinical infectious diseases*
357   *: an official publication of the Infectious Diseases Society of America* 2014, **59 Suppl**
358   **5**:S352-355.
359   4.   Neville BA, d'Enfert C, Bougnoux ME: **Candida albicans commensalism in the**
360   **gastrointestinal tract**. *FEMS Yeast Res* 2015, **15**(7).
361   5.   Fan D, Coughlin LA, Neubauer MM, Kim J, Kim MS, Zhan X, Simms-Waldrip TR, Xie
362   Y, Hooper LV, Koh AY: **Activation of HIF-1alpha and LL-37 by commensal bacteria**
363   **inhibits Candida albicans colonization**. *Nat Med* 2015, **21**(7):808-814.
364   6.   Huppert M, Cazin J, Jr.: **Pathogenesis of Candida albicans infection following antibiotic**
365   **therapy. II. Further studies of the effect of antibiotics on the in vitro growth of**
366   **Candida albicans**. *J Bacteriol* 1955, **70**(4):435-439.
367   7.   Krause R, Krejs GJ, Wenisch C, Reisinger EC: **Elevated fecal Candida counts in patients**
368   **with antibiotic-associated diarrhea: role of soluble fecal substances**. *Clin Diagn Lab*
369   *Immunol* 2003, **10**(1):167-168.
370   8.   Krause R, Schwab E, Bachhiesl D, Daxbock F, Wenisch C, Krejs GJ, Reisinger EC: **Role**
371   **of Candida in antibiotic-associated diarrhea**. *J Infect Dis* 2001, **184**(8):1065-1069.
372   9.   Schulte DM, Sethi A, Gangnon R, Duster M, Maki DG, Safdar N: **Risk factors for**
373   **Candida colonization and Co-colonization with multi-drug resistant organisms at**
374   **admission**. *Antimicrob Resist Infect Control* 2015, **4**:46.
375   10.   Guiot HF, Fibbe WE, van 't Wout JW: **Risk factors for fungal infection in patients with**
376   **malignant hematologic disorders: implications for empirical therapy and**
377   **prophylaxis**. *Clin Infect Dis* 1994, **18**(4):525-532.
378   11.   Guastalegname M, Russo A, Falcone M, Giuliano S, Venditti M: **Candidemia subsequent**
379   **to severe infection due to Clostridium difficile: is there a link?** *Clin Infect Dis* 2013,
380   **57**(5):772-774.
381   12.   Nerandzic MM, Mullane K, Miller MA, Babakhani F, Donskey CJ: **Reduced acquisition**
382   **and overgrowth of vancomycin-resistant enterococci and Candida species in patients**
383   **treated with fidaxomicin versus vancomycin for Clostridium difficile infection**. *Clin*
384   *Infect Dis* 2012, **55 Suppl 2**:S121-126.
385   13.   Samonis G, Gikas A, Anaissie EJ, Vrenzos G, Maraki S, Tselentis Y, Bodey GP:
386   **Prospective evaluation of effects of broad-spectrum antibiotics on gastrointestinal**
387   **yeast colonization of humans**. *Antimicrob Agents Chemother* 1993, **37**(1):51-53.
388   14.   Wiesner SM, Jechorek RP, Garni RM, Bendel CM, Wells CL: **Gastrointestinal**
389   **colonization by Candida albicans mutant strains in antibiotic-treated mice**. *Clin*
390   *Diagn Lab Immunol* 2001, **8**(1):192-195.
391   15.   Ekenna O, Sherertz RJ: **Factors affecting colonization and dissemination of Candida**
392   **albicans from the gastrointestinal tract of mice**. *Infect Immun* 1987, **55**(7):1558-1563.

16. Koh AY: **Murine models of Candida gastrointestinal colonization and dissemination**. *Eukaryot Cell* 2013, **12**(11):1416-1422.

17. Prieto D, Pla J: **Distinct stages during colonization of the mouse gastrointestinal tract by Candida albicans**. *Front Microbiol* 2015, **6**:792.

18. Vautier S, Drummond RA, Chen K, Murray GI, Kadosh D, Brown AJ, Gow NA, MacCallum DM, Kolls JK, Brown GD: **Candida albicans colonization and dissemination from the murine gastrointestinal tract: the influence of morphology and Th17 immunity**. *Cell Microbiol* 2015, **17**(4):445-450.

19. Conti HR, Huppler AR, Whibley N, Gaffen SL: **Animal models for candidiasis**. *Curr Protoc Immunol* 2014, **105**:19 16 11-17.

20. Mason KL, Erb Downward JR, Mason KD, Falkowski NR, Eaton KA, Kao JY, Young VB, Huffnagle GB: **Candida albicans and bacterial microbiota interactions in the cecum during recolonization following broad-spectrum antibiotic therapy**. *Infect Immun* 2012, **80**(10):3371-3380.

21. Erb Downward JR, Falkowski NR, Mason KL, Muraglia R, Huffnagle GB: **Modulation of post-antibiotic bacterial community reassembly and host response by Candida albicans**. *Sci Rep* 2013, **3**:2191.

22. Suez J, Elinav E: **The path towards microbiome-based metabolite treatment**. *Nat Microbiol* 2017, **2**:17075.

23. Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H, Kinnebrew M, Viale A *et al*: **Precision microbiome reconstitution restores bile acid mediated resistance to Clostridium difficile**. *Nature* 2015, **517**(7533):205-208.

24. Kohli N, Crisp Z, Riordan R, Li M, Alaniz RC, Jayaraman A: **The microbiota metabolite indole inhibits Salmonella virulence: Involvement of the PhoPQ two-component system**. *PLoS One* 2018, **13**(1):e0190613.

25. Gutierrez D, Weinstock A, Antharam VC, Gu H, Jasbi P, Shi X, Dirks B, Krajmalnik-Brown R, Maldonado J, Guinan J *et al*: **Antibiotic-induced gut metabolome and microbiome alterations increase the susceptibility to Candida albicans colonization in the gastrointestinal tract**. *FEMS Microbiol Ecol* 2020, **96**(1).

26. Chiang JYL: **Bile Acid Metabolism and Signaling**. *Comprehensive Physiology* 2013, **3**(3):1191-1212.

27. Staels B, Fonseca VA: **Bile Acids and Metabolic Regulation: Mechanisms and clinical responses to bile acid sequestration**. *Diabetes Care* 2009, **32**(Suppl 2):S237-S245.

28. Sayin SI, Wahlstrom A, Felin J, Jantti S, Marschall HU, Bamberg K, Angelin B, Hyotylainen T, Oresic M, Backhed F: **Gut microbiota regulates bile acid metabolism by reducing the levels of tauro-beta-muricholic acid, a naturally occurring FXR antagonist**. *Cell Metab* 2013, **17**(2):225-235.

29. Song WS, Park HM, Ha JM, Shin SG, Park HG, Kim J, Zhang T, Ahn DH, Kim SM, Yang YH *et al*: **Discovery of glycocholic acid and taurochenodeoxycholic acid as phenotypic biomarkers in cholangiocarcinoma**. *Sci Rep* 2018, **8**(1):11088.

30. Manieri E, Folgueira C, Rodriguez ME, Leiva-Vega L, Esteban-Lafuente L, Chen C, Cubero FJ, Barrett T, Cavanagh-Kyros J, Seruggia D *et al*: **JNK-mediated disruption of bile acid homeostasis promotes intrahepatic cholangiocarcinoma**. *Proc Natl Acad Sci U S A* 2020, **117**(28):16492-16499.

31. Kuhn T, Stepien M, Lopez-Nogueroles M, Damms-Machado A, Sookthai D, Johnson T, Roca M, Husing A, Maldonado SG, Cross AJ *et al*: **Prediagnostic Plasma Bile Acid**

Levels and Colon Cancer Risk: A Prospective Study. *J Natl Cancer Inst* 2020,
112(5):516-524.

32. Tian Q, Yang R, Wang Y, Liu J, Wee A, Saxena R, Wang L, Li M, Liu L, Shan S *et al*: **A**
High Serum Level of Taurocholic Acid Is Correlated With the Severity and
Resolution of Drug-induced Liver Injury. *Clin Gastroenterol Hepatol* 2020.

33. Luo L, Aubrecht J, Li D, Warner RL, Johnson KJ, Kenny J, Colangelo JL: **Assessment of**
serum bile acid profiles as biomarkers of liver injury and liver disease in humans.
PLoS One 2018, **13**(3):e0193824.

34. Ridlon JM, Wolf PG, Gaskins HR: **Taurocholic acid metabolism by gut microbes and**
colon cancer. *Gut Microbes* 2016, **7**(3):201-215.

35. Devkota S, Wang Y, Musch MW, Leone V, Fehlner-Peach H, Nadimpalli A, Antonopoulos
DA, Jabri B, Chang EB: **Dietary-fat-induced taurocholic acid promotes pathobiont**
expansion and colitis in Il10-/- mice. *Nature* 2012, **487**(7405):104-108.

36. Liu Z, Zhang Z, Huang M, Sun X, Liu B, Guo Q, Chang Q, Duan Z: **Taurocholic acid is**
an active promoting factor, not just a biomarker of progression of liver cirrhosis:
evidence from a human metabolomic study and in vitro experiments. *BMC*
Gastroenterol 2018, **18**(1):112.

37. Bajaj JS, Liu EJ, Kheradman R, Fagan A, Heuman DM, White M, Gavis EA, Hylemon P,
Sikaroodi M, Gillevet PM: **Fungal dysbiosis in cirrhosis**. *Gut* 2018, **67**(6):1146-1154.

38. Maraolo AE, Scotto R, Zappulo E, Pinchera B, Schiano Moriello N, Nappa S, Buonomo
AR, Gentile I: **Novel strategies for the management of bacterial and fungal infections**
in patients with liver cirrhosis: focus on new antimicrobials. *Expert Rev Anti Infect*
Ther 2020, **18**(3):191-202.

39. Li B, Yang C, Qian Z, Huang Y, Wang X, Zhong G, Chen J: **Spontaneous Fungal Ascites**
Infection in Patients with Cirrhosis: An Analysis of 10 Cases. *Infect Dis Ther* 2021.

40. Fiore M, Leone S: **Spontaneous fungal peritonitis: Epidemiology, current evidence and**
future prospective. *World J Gastroenterol* 2016, **22**(34):7742-7747.

41. Ferrarese A, Cattelan A, Cillo U, Gringeri E, Russo FP, Germani G, Gambato M, Burra P,
Senzolo M: **Invasive fungal infection before and after liver transplantation**. *World J*
Gastroenterol 2020, **26**(47):7485-7496.

42. Ethanic M, Stanimirov B, Pavlovic N, Golocorbin-Kon S, Al-Salami H, Stankov K, Mikov
M: **Pharmacological Applications of Bile Acids and Their Derivatives in the**
Treatment of Metabolic Syndrome. *Front Pharmacol* 2018, **9**:1382.

43. Li T, Chiang JYL: **Bile acid-based therapies for non-alcoholic steatohepatitis and**
alcoholic liver disease. *Hepatobiliary Surg Nutr* 2020, **9**(2):152-169.

44. Chiang JYL, Ferrell JM: **Bile Acid Biology, Pathophysiology, and Therapeutics**. *Clin*
Liver Dis (Hoboken) 2020, **15**(3):91-94.

45. Zhang Y, Limaye PB, Renaud HJ, Klaassen CD: **Effect of various antibiotics on**
modulation of intestinal microbiota and bile acid profile in mice. *Toxicol Appl*
Pharmacol 2014, **277**(2):138-145.

46. Culpepper T, Rowe CC, Rusch CT, Burns AM, Federico AP, Girard SA, Tompkins TA,
Nieves C, Jr., Dennis-Wall JC, Christman MC *et al*: **Three probiotic strains exert**
different effects on plasma bile acid profiles in healthy obese adults: randomised,
double-blind placebo-controlled crossover study. *Benef Microbes* 2019, **10**(5):497-509.

47. Begley M, Hill C, Gahan CG: **Bile salt hydrolase activity in probiotics**. *Appl Environ*
Microbiol 2006, **72**(3):1729-1738.

485 48. Vrieze A, Out C, Fuentes S, Jonker L, Reuling I, Kootte RS, van Nood E, Holleman F, Knaapen M, Romijn JA *et al*: **Impact of oral vancomycin on gut microbiota, bile acid metabolism, and insulin sensitivity**. *J Hepatol* 2014, **60**(4):824-831.

488 49. Zarrinpar A, Chaix A, Xu ZZ, Chang MW, Marotz CA, Saghatelian A, Knight R, Panda S: **Antibiotic-induced microbiome depletion alters metabolic homeostasis by affecting gut signaling and colonic metabolism**. *Nat Commun* 2018, **9**(1):2872.

491 50. Mooranian A, Zamani N, Mikov M, Golocorbin-Kon S, Stojanovic G, Arfuso F, Al-Salami H: **Stability and biological testing of taurine-conjugated bile acid antioxidant microcapsules for diabetes treatment**. *Ther Deliv* 2019, **10**(2):99-106.

494 51. Khatun Z, Nurunnabi M, Reeck GR, Cho KJ, Lee YK: **Oral delivery of taurocholic acid linked heparin-docetaxel conjugates for cancer therapy**. *J Control Release* 2013, **170**(1):74-82.

497 52. Fan D, Coughlin LA, Neubauer MM, Kim J, Kim MS, Zhan X, Simms-Waldrip TR, Xie Y, Hooper LV, Koh AY: **Activation of HIF-1α and LL-37 by commensal bacteria inhibits Candida albicans colonization**. *Nature medicine* 2015, **21**(7):808.

500 53. Koh AY, Kohler JR, Coggshall KT, Van Rooijen N, Pier GB: **Mucosal damage and neutropenia are required for Candida albicans dissemination**. *PLoS Pathog* 2008, **4**(2):e35.

503 54. Netea MG, Joosten LA, van der Meer JW, Kullberg BJ, van de Veerdonk FL: **Immune defence against Candida fungal infections**. *Nat Rev Immunol* 2015, **15**(10):630-642.

505 55. Leonardi I, Li X, Semon A, Li D, Doron I, Putzel G, Bar A, Prieto D, Rescigno M, McGovern DPB *et al*: **CX3CR1(+) mononuclear phagocytes control immunity to intestinal fungi**. *Science* 2018, **359**(6372):232-236.

508 56. Drummond RA, Gaffen SL, Hise AG, Brown GD: **Innate Defense against Fungal Pathogens**. *Cold Spring Harb Perspect Med* 2014, **5**(6).

510 57. Wahlstrom A, Sayin SI, Marschall HU, Backhed F: **Intestinal Crosstalk between Bile Acids and Microbiota and Its Impact on Host Metabolism**. *Cell Metab* 2016, **24**(1):41-50.

513 58. Dekaney CM, von Allmen DC, Garrison AP, Rigby RJ, Lund PK, Henning SJ, Helmrath MA: **Bacterial-dependent up-regulation of intestinal bile acid binding protein and transport is FXR-mediated following ileo-cecal resection**. *Surgery* 2008, **144**(2):174-181.

517 59. Fiorucci S, Biagioli M, Zampella A, Distrutti E: **Bile Acids Activated Receptors Regulate Innate Immunity**. *Front Immunol* 2018, **9**:1853.

519 60. Vavassori P, Mencarelli A, Renga B, Distrutti E, Fiorucci S: **The bile acid receptor FXR is a modulator of intestinal innate immunity**. *J Immunol* 2009, **183**(10):6251-6261.

521 61. Li L, Zhang Q, Peng J, Jiang C, Zhang Y, Shen L, Dong J, Wang Y, Jiang Y: **Activation of farnesoid X receptor downregulates monocyte chemoattractant protein-1 in murine macrophage**. *Biochem Biophys Res Commun* 2015, **467**(4):841-846.

524 62. Ridlon JM, Kang DJ, Hylemon PB, Bajaj JS: **Bile acids and the gut microbiome**. *Curr Opin Gastroenterol* 2014, **30**(3):332-338.

526 63. Ramirez-Perez O, Cruz-Ramon V, Chinchilla-Lopez P, Mendez-Sanchez N: **The Role of the Gut Microbiota in Bile Acid Metabolism**. *Ann Hepatol* 2017, **16**(Suppl. 1: s3-105.):s15-s20.

529 64. Ridlon JM, Harris SC, Bhowmik S, Kang DJ, Hylemon PB: **Consequences of bile salt biotransformations by intestinal bacteria**. *Gut Microbes* 2016, **7**(1):22-39.

65. Ridlon JM, Hylemon PB: **Identification and characterization of two bile acid coenzyme A transferases from Clostridium scindens, a bile acid 7alpha-dehydroxylating intestinal bacterium**. *J Lipid Res* 2012, **53**(1):66-76.

66. Ridlon JM, Kang DJ, Hylemon PB: **Bile salt biotransformations by human intestinal bacteria**. *J Lipid Res* 2006, **47**(2):241-259.

67. O'Flaherty S, Briner Crawley A, Theriot CM, Barrangou R: **The Lactobacillus Bile Salt Hydrolase Repertoire Reveals Niche-Specific Adaptation**. *mSphere* 2018, **3**(3).

68. Hooper LV, Stappenbeck TS, Hong CV, Gordon JI: **Angiogenins: a new class of microbicidal proteins involved in innate immunity**. *Nat Immunol* 2003, **4**(3):269-273.

69. Forman RA, deSchoolmeester ML, Hurst RJ, Wright SH, Pemberton AD, Else KJ: **The goblet cell is the cellular source of the anti-microbial angiogenin 4 in the large intestine post Trichuris muris infection**. *PLoS One* 2012, **7**(9):e42248.

70. Walker CR, Hautefort I, Dalton JE, Overweg K, Egan CE, Bongaerts RJ, Newton DJ, Cruickshank SM, Andrew EM, Carding SR: **Intestinal intraepithelial lymphocyte-enterocyte crosstalk regulates production of bactericidal angiogenin 4 by Paneth cells upon microbial challenge**. *PLoS One* 2013, **8**(12):e84553.

71. Doron I, Leonardi I, Li XV, Fiers WD, Semon A, Bialt-DeCelie M, Migaud M, Gao IH, Lin WY, Kusakabe T *et al*: **Human gut mycobiota tune immunity via CARD9-dependent induction of anti-fungal IgG antibodies**. *Cell* 2021, **184**(4):1017-1031 e1014.

72. Medina-Contreras O, Geem D, Laur O, Williams IR, Lira SA, Nusrat A, Parkos CA, Denning TL: **CX3CR1 regulates intestinal macrophage homeostasis, bacterial translocation, and colitogenic Th17 responses in mice**. *J Clin Invest* 2011, **121**(12):4787-4795.

73. Liu HY, Roos S, Jonsson H, Ahl D, Dicksved J, Lindberg JE, Lundh T: **Effects of Lactobacillus johnsonii and Lactobacillus reuteri on gut barrier function and heat shock proteins in intestinal porcine epithelial cells**. *Physiol Rep* 2015, **3**(4).

74. Antonini M, Lo Conte M, Sorini C, Falcone M: **How the Interplay Between the Commensal Microbiota, Gut Barrier Integrity, and Mucosal Immunity Regulates Brain Autoimmunity**. *Front Immunol* 2019, **10**:1937.

75. Guo W, Zhou X, Li X, Zhu Q, Peng J, Zhu B, Zheng X, Lu Y, Yang D, Wang B *et al*: **Depletion of Gut Microbiota Impairs Gut Barrier Function and Antiviral Immune Defense in the Liver**. *Front Immunol* 2021, **12**:636803.

76. Hiippala K, Jouhten H, Ronkainen A, Hartikainen A, Kainulainen V, Jalanka J, Satokari R: **The Potential of Gut Commensals in Reinforcing Intestinal Barrier Function and Alleviating Inflammation**. *Nutrients* 2018, **10**(8).

77. Charlet R, Bortolus C, Sendid B, Jawhara S: **Bacteroides thetaiotaomicron and Lactobacillus johnsonii modulate intestinal inflammation and eliminate fungi via enzymatic hydrolysis of the fungal cell wall**. *Sci Rep* 2020, **10**(1):11510.

78. Honda M, Surewaard BGJ, Watanabe M, Hedrick CC, Lee WY, Brown K, McCoy KD, Kubes P: **Perivascular localization of macrophages in the intestinal mucosa is regulated by Nr4a1 and the microbiome**. *Nat Commun* 2020, **11**(1):1329.

79. Out C, Patankar JV, Doktorova M, Boesjes M, Bos T, de Boer S, Havinga R, Wolters H, Boverhof R, van Dijk TH *et al*: **Gut microbiota inhibit Asbt-dependent intestinal bile acid reabsorption via Gata4**. *J Hepatol* 2015, **63**(3):697-704.

80. Islam KB, Fukiya S, Hagio M, Fujii N, Ishizuka S, Ooka T, Ogura Y, Hayashi T, Yokota
A: **Bile acid is a host factor that regulates the composition of the cecal microbiota in**
rats. *Gastroenterology* 2011, **141**(5):1773-1781.
81. Ruiz L, Margolles A, Sanchez B: **Bile resistance mechanisms in Lactobacillus and**
Bifidobacterium. *Front Microbiol* 2013, **4**:396.
82. Guinan J, Villa P, Thangamani S: **Secondary bile acids inhibit Candida albicans growth**
and morphogenesis. *Pathog Dis* 2018, **76**(3).
83. Guinan J, Thangamani S: **Antibiotic-induced alterations in taurocholic acid levels**
promote gastrointestinal colonization of Candida albicans. *FEMS Microbiol Lett* 2018,
365(18).
84. Chelakkot C, Ghim J, Ryu SH: **Mechanisms regulating intestinal barrier integrity and**
its pathological implications. *Exp Mol Med* 2018, **50**(8):1-9.
85. Meadows V, Kennedy L, Kundu D, Alpini G, Francis H: **Bile Acid Receptor**
Therapeutics Effects on Chronic Liver Diseases. *Front Med (Lausanne)* 2020, **7**:15.
86. Dayan SH, Humphrey S, Jones DH, Lizzul PF, Gross TM, Stauffer K, Beddingfield FC,
3rd: **Overview of ATX-101 (Deoxycholic Acid Injection): A Nonsurgical Approach for**
Reduction of Submental Fat. *Dermatol Surg* 2016, **42 Suppl 1**:S263-S270.
87. Beer K, Weinkle SH, Cox SE, Rubin MG, Shamban A, Somogyif C: **ATX-101**
(Deoxycholic Acid Injection) for Reduction of Submental Fat: Results From a 12-
Month Open-Label Study. *J Drugs Dermatol* 2019, **18**(9):870-877.
88. Daruich A, Picard E, Boatright JH, Behar-Cohen F: **Review: The bile acids urso- and**
tauroursodeoxycholic acid as neuroprotective therapies in retinal disease. *Mol Vis*
2019, **25**:610-624.
89. Gordon SC, Wu KH, Lindor K, Bowlus CL, Rodriguez CV, Anderson H, Boscarino JA,
Trudeau S, Rupp LB, Haller IV *et al*: **Ursodeoxycholic Acid Treatment Preferentially**
Improves Overall Survival Among African Americans With Primary Biliary
Cholangitis. *Am J Gastroenterol* 2020, **115**(2):262-270.
90. Xiao Y, Zhou K, Lu Y, Yan W, Cai W, Wang Y: **Administration of antibiotics**
contributes to cholestasis in pediatric patients with intestinal failure via the alteration
of FXR signaling. *Exp Mol Med* 2018, **50**(12):1-14.
91. Singla P, Dalal P, Kaur M, Arya G, Nimesh S, Singh R, Salunke DB: **Bile Acid Oligomers**
and Their Combination with Antibiotics To Combat Bacterial Infections. *J Med Chem*
2018, **61**(22):10265-10275.
92. Winston JA, Theriot CM: **Diversification of host bile acids by members of the gut**
microbiota. *Gut Microbes* 2020, **11**(2):158-171.
93. Sinha SR, Haileselassie Y, Nguyen LP, Tropini C, Wang M, Becker LS, Sim D, Jarr K,
Spear ET, Singh G *et al*: **Dysbiosis-Induced Secondary Bile Acid Deficiency Promotes**
Intestinal Inflammation. *Cell Host Microbe* 2020, **27**(4):659-670 e655.
94. Trojanowska D, Zwolinska-Wcislo M, Tokarczyk M, Kosowski K, Mach T, Budak A: **The**
role of Candida in inflammatory bowel disease. Estimation of transmission of C.
albicans fungi in gastrointestinal tract based on genetic affinity between strains. *Med*
Sci Monit 2010, **16**(10):CR451-457.
95. Gerard R, Sendid B, Colombel JF, Poulain D, Jouault T: **An immunological link between**
Candida albicans colonization and Crohn's disease. *Crit Rev Microbiol* 2015,
41(2):135-139.

96. Iliev ID, Funari VA, Taylor KD, Nguyen Q, Reyes CN, Strom SP, Brown J, Becker CA, Fleshner PR, Dubinsky M *et al*: **Interactions between commensal fungi and the C-type lectin receptor Dectin-1 influence colitis**. *Science* 2012, **336**(6086):1314-1317.

97. Leonardi I, Paramsothy S, Doron I, Semon A, Kaakoush NO, Clemente JC, Faith JJ, Borody TJ, Mitchell HM, Colombel JF *et al*: **Fungal Trans-kingdom Dynamics Linked to Responsiveness to Fecal Microbiota Transplantation (FMT) Therapy in Ulcerative Colitis**. *Cell Host Microbe* 2020, **27**(5):823-829 e823.

98. Wajszczuk CP, Dummer JS, Ho M, Van Thiel DH, Starzl TE, Iwatsuki S, Shaw B, Jr.: **Fungal infections in liver transplant recipients**. *Transplantation* 1985, **40**(4):347-353.

99. Brumble L, Keaveny AP: **Editorial: The Risky Business of Fungal Infections in Patients with Cirrhosis**. *Am J Gastroenterol* 2018, **113**(4):564-566.

100. Habib S, Yarlagadda S, Carreon TA, Schader LM, Hsu CH: **Fungal Infection in Acutely Decompensated Cirrhosis Patients: Value of Model for End-Stage Liver Disease Score**. *Gastroenterology Res* 2020, **13**(5):199-207.

101. Hassan EA, Abd El-Rehim AS, Hassany SM, Ahmed AO, Elsherbiny NM, Mohammed MH: **Fungal infection in patients with end-stage liver disease: low frequency or low index of suspicion**. *Int J Infect Dis* 2014, **23**:69-74.

102. Pramar YV, Mandal TK, Bostanian LA, Nguyen AT, Miller V, Morris TC, Graves RA: **Stability of Compounded Ursodiol Suspensions in PCCA Base, SuspendIt**. *Int J Pharm Compd* 2019, **23**(1):70-76.

103. Simental-Mendia M, Sanchez-Garcia A, Simental-Mendia LE: **Effect of ursodeoxycholic acid on liver markers: A systematic review and meta-analysis of randomized placebo-controlled clinical trials**. *Br J Clin Pharmacol* 2020.

104. Phaw NA, Dyson JK, Jones D: **Emerging drugs for the treatment of primary biliary cholangitis**. *Expert Opin Emerg Drugs* 2020:1-12.

105. Vlasova AN, Kandasamy S, Chattha KS, Rajashekara G, Saif LJ: **Comparison of probiotic lactobacilli and bifidobacteria effects, immune responses and rotavirus vaccines and infection in different host species**. *Vet Immunol Immunopathol* 2016, **172**:72-84.

106. Ciorba MA: **A gastroenterologist's guide to probiotics**. *Clin Gastroenterol Hepatol* 2012, **10**(9):960-968.

107. Nash AK, Auchtung TA, Wong MC, Smith DP, Gesell JR, Ross MC, Stewart CJ, Metcalf GA, Muzny DM, Gibbs RA *et al*: **The gut mycobiome of the Human Microbiome Project healthy cohort**. *Microbiome* 2017, **5**(1):153.

108. Lai GC, Tan TG, Pavelka N: **The mammalian mycobiome: A complex system in a dynamic relationship with the host**. *Wiley Interdiscip Rev Syst Biol Med* 2019, **11**(1):e1438.

109. Chacon MR, Lozano-Bartolome J, Portero-Otin M, Rodriguez MM, Xifra G, Puig J, Blasco G, Ricart W, Chaves FJ, Fernandez-Real JM: **The gut mycobiome composition is linked to carotid atherosclerosis**. *Benef Microbes* 2018, **9**(2):185-198.

110. Enaud R, Vandenborght LE, Coron N, Bazin T, Prevel R, Schaeverbeke T, Berger P, Fayon M, Lamireau T, Delhaes L: **The Mycobiome: A Neglected Component in the Microbiota-Gut-Brain Axis**. *Microorganisms* 2018, **6**(1).

111. Forbes JD, Bernstein CN, Tremlett H, Van Domselaar G, Knox NC: **A Fungal World: Could the Gut Mycobiome Be Involved in Neurological Disease?** *Front Microbiol* 2018, **9**:3249.

112. Hernandez-Santos N, Klein BS: **Through the Scope Darkly: The Gut Mycobiome Comes into Focus**. *Cell Host Microbe* 2017, **22**(6):728-729.

113. Huseyin CE, O'Toole PW, Cotter PD, Scanlan PD: **Forgotten fungi-the gut mycobiome in human health and disease**. *FEMS Microbiol Rev* 2017, **41**(4):479-511.

114. Sam QH, Chang MW, Chai LY: **The Fungal Mycobiome and Its Interaction with Gut Bacteria in the Host**. *Int J Mol Sci* 2017, **18**(2).

115. Szabo G: **Gut-Liver Axis Beyond the Microbiome: How the Fungal Mycobiome Contributes to Alcoholic Liver Disease**. *Hepatology* 2018, **68**(6):2426-2428.

116. Chu Y, Jiang MZ, Xu B, Wang WJ, Chen D, Li XW, Zhang YJ, Liang J: **Specific changes of enteric mycobiota and virome in inflammatory bowel disease**. *J Dig Dis* 2018, **19**(1):2-7.

117. Ward TL, Dominguez-Bello MG, Heisel T, Al-Ghalith G, Knights D, Gale CA: **Development of the Human Mycobiome over the First Month of Life and across Body Sites**. *mSystems* 2018, **3**(3).

118. Stamatiades GA, Ioannou P, Petrikkos G, Tsioutis C: **Fungal infections in patients with inflammatory bowel disease: A systematic review**. *Mycoses* 2018, **61**(6):366-376.

119. Kantarcioglu AS, Kiraz N, Aydin A: **Microbiota-Gut-Brain Axis: Yeast Species Isolated from Stool Samples of Children with Suspected or Diagnosed Autism Spectrum Disorders and In Vitro Susceptibility Against Nystatin and Fluconazole**. *Mycopathologia* 2016, **181**(1-2):1-7.

120. Zuo T, Wong SH, Cheung CP, Lam K, Lui R, Cheung K, Zhang F, Tang W, Ching JYL, Wu JCY *et al*: **Gut fungal dysbiosis correlates with reduced efficacy of fecal microbiota transplantation in Clostridium difficile infection**. *Nat Commun* 2018, **9**(1):3663.

121. Tso GHW, Reales-Calderon JA, Tan ASM, Sem X, Le GTT, Tan TG, Lai GC, Srinivasan KG, Yurieva M, Liao W *et al*: **Experimental evolution of a fungal pathogen into a gut symbiont**. *Science* 2018, **362**(6414):589-595.

Figure Legends

Figure 1. Taurocholic acid (TCA) is the major bile acid metabolite up-regulated in the cefoperazone-treated mice susceptible to CA infection. (A) BALB/c mice were fed sterile water in the presence or absence of cefoperazone (0.5 mg/mL) for 7 days and then infected with ~2 × 10^8 CFU CA SC5314 via oral gavage. Antibiotic treatment was continued until the end of the experiment. Three days post-infection, all mice were injected with three doses of

cyclophosphamide intraperitoneally (150mg/kg body weight) and monitored for survival. A log

rank test was performed using 95% confidence intervals; statistical significance was calculated to

compare the antibiotic-treated and untreated control groups. (B) Fungal load from cecum, liver

and kidney were collected immediately after death in the antibiotic-treated groups and mice

euthanized after 6 days post-cyclophosphamide treatment from control groups. Statistical

significance was evaluated using the Mann-Whitney U test. (C) Relative Log_2 fold-change of bile

acid metabolites in antibiotic-treated C57BL/6 mice relative to control groups. (D) Relative

abundance of bile acid metabolites that were highly upregulated and significant in the antibiotic-

treated group relative to control groups. At least five mice per group was used and the data

represents mean ± SEM. one way-ANOVA followed by a multiple comparison using Bonferroni

correction. *P* values of (*≤ 0.05) (**P≤ 0.01) (***P≤ 0.001) (****P≤ 0.0001) were considered as

significant.

Figure 2. TCA alone induces fungal colonization and dissemination in the absence of

antibiotics and immunosuppression. (A) Experimental outline. C57BL/6 mice infected with ~2

$\times 10^8$ CFU CA SC5314 via oral gavage. Untreated mice received sterile drinking water (untreated

group); treatment group received sterile water containing 1% TCA (TCA group). **(B)** Mice were

monitored for survival. A log rank test was performed using 95% confidence intervals; statistical

significance was calculated to compare the antibiotic-treated and untreated control groups. **(C)**

Fungal load from feces collected from untreated and TCA groups 1-, 4-, 7-, and 12-days post-

infection. **(D)** Fungal load from cecum, liver, and kidney from dead mice collected immediately

in the TCA-treated groups and mice euthanized 15 days post-infection for untreated groups. Ten

mice per group was used and the data represents mean ± SEM. Statistical significance was

evaluated using the Mann-Whitney U test. *p* values ≤0.05 (*) or ≤0.01 (**) were considered

statistically significant.

Figure 3. TCA induces fungal dissemination from the GI tract of immunosuppressed mice

in the absence of antibiotic treatment. **(A)** Experimental outline. BALB/c mice infected with

~2 × 10^8 CFU CA SC5314 via oral gavage received sterile water with or without containing 1%

TCA. Three days post-infection, all mice were injected with three doses of cyclophosphamide

intraperitoneally (150mg/kg body weight). **(B)** Mice were monitored for survival and log rank test

was performed using 95% confidence intervals; statistical significance was calculated to compare

the antibiotic-treated and untreated control groups. **(C)** Fungal load from cecum, liver, and kidney

collected immediately after death in the TCA+ cyclo treated groups and mice euthanized 9 days

post-infection for cyclo groups. 4-5 mice per group was used and the data represents mean ± SEM.

Statistical significance was evaluated using the Mann-Whitney U test. *p* values ≤0.05 (*) or ≤0.01

(**) were considered statistically significant.

Figure 4. TCA increases the intestinal permeability leading to fungal dissemination from the

GI tract. **(A)** Gut permeability was measured in infected mice using a FITC-dextran assay. At 10

days post-infection, male and female BL57/6J mice in both control and TCA-treated groups

infected with CA were given an oral gavage of 150 μL PBS containing 15 mg FITC-dextran. Four

hours after administering FITC-dextran, mice were anesthetized, and blood was collected via retro-

orbital bleed. Blood samples were processed via a two-fold serial dilution in a 96-well plate and

fluorescence was measure via a plate reader (excitation: 485 nm; emission: 520 nm). Data

represent mean ± SEM. Data represent mean ± SEM. Statistical significance was evaluated using

the Mann-Whitney U test. *p* values ≤0.05 (*) or ≤0.01 (**) were considered statistically significant.

(**B**) ZO-1 tight junction protein expression in untreated and TCA-treated mice. Colon tissue from

untreated and TCA-treated mice from CA-infected mice (10 days post-infection) were stained with

ZO-1 and DAPI antibodies. Representative images are shown here.

Figure 5. TCA inhibits the expression of *ang4* and CX3CR1 in the colon *in vivo*. Colon tissue

from untreated and TCA-treated mice collected after 10 days of CA infection and treatment were

RNA sequenced to examine the expression level of host defense genes. (**A**) Average relative

expression of host defense genes in untreated and TCA treated mice were shown. (**B**) Relative

expression of *ang4* and *Cx3cr1* in untreated and TCA treated mice were shown. (**C**) Average

relative expression of intestinal metabolite receptors in untreated and TCA treated mice were

shown. Four to five mice per group was used. Data represent mean ± SEM. Statistical significance

was evaluated using the Student's t test.

Figure 6. TCA alters the relative abundance of intestinal microbiota. (A and **B)** C57BL/6 mice

were infected with ~2 × 10^8 CFU CA SC5314 via oral gavage. Control mice received sterile

drinking water (untreated group); the treatment group received sterile water containing 1% TCA

(TCA group). Ten days post-infection, cecal contents and colon scrapings were collected to

determine the relative abundance of intestinal microbiota. Bacterial members that are significantly

altered in the TCA group compared to the untreated control groups both in cecal content and colon

mucosal scrapings were shown here. Five mice per group was used. Data represent mean ± SEM.

Statistical significance was evaluated using the Student's t test.

Figure 1

Figure 2

(A)

(B)

(C)

Figure 3

Figure 4

Figure 5

Figure 6