Why can we detect lianas from space?

Marco D. Visser[1,2,*], Matteo Detto[1], Félicien Meunier[3,4], Jin Wu[5], Jane R. Foster[6,7], David C.

Marvin[8], Peter M. van Bodegom[2], Boris Bongalov[9], Matheus Henrique Nunes[9,10], David

Coomes[9,11], Hans Verbeeck[4], J. Antonio Guzmán Q[12], Arturo Sanchez-Azofeifa[12], Chris J.

Chandler[13], Geertje M.F van der Heijden[13], Doreen S. Boyd[13], Giles M. Foody[13], Mark

E.J.Cutler[14], Eben N. Broadbent[15], Shawn P. Serbin[16], Stefan Schnitzer[17], M. Elizabeth

Rodríguez-Ronderos[18], Frank Sterck[19], José A. Medina-Vega[20], and Steve Pacala[1]

1. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ.

2. Institute of Environmental Sciences, Leiden University, Einsteinweg 2, 2333 CC Leiden, The

Netherlands.

3. Computational and Applied Vegetation Ecology, Department of Environment, Ghent University,

Ghent, Belgium.

4. Department of Earth and Environment, Boston University, Boston, USA

5. University of Hong Kong, Hong Kong.

6. Rubenstein School of Environment and Natural Resources, University of Vermont, Burlington,

VT, USA

7. Southern Research Station, US Forest Service, Knoxville, TN, USA.

8. Salo Sciences, Inc., San Francisco, CA USA.

9. Department of Plant Sciences, Forest Ecology and Conservation group, University of

Cambridge, Cambridge, UK.

10. Department of Geographical Sciences, University of Maryland, College Park, MD 20742, USA.

11. Department of Plant Sciences, The Conservation Research Institute, University of Cambridge,

Cambridge CB2 3QZ, UK.

2412. Centre for Earth Observation Sciences (CEOS), Earth and Atmospheric Sciences Department,
25 University of Alberta, Edmonton, Alberta, Canada T6M 2R7.

2613. School of Geography, University of Nottingham, University Park, Nottingham, United Kingdom

2714. School of Social Sciences, University of Dundee, Dundee, United Kingdom DD1 4HN.

2815. School of Forest Resources and Conservation, University of Florida, Gainesville, Florida, USA.

2916. Terrestrial Ecosystem Science & Technology Group, Environmental Sciences Department,
30 Brookhaven National Laboratory, Upton, NY 11973-5000, USA.

3117. Department of Biological Sciences, Marquette University, Milwaukee, WI 53201, USA.

3218. Department of Biological Sciences, National University of Singapore, Singapore.

3319. Forest Ecology and Management Group, Wageningen University, Wageningen 6708 PB the
34 Netherlands.

3520. Forest Global Earth Observatory, Smithsonian Tropical Research Institute, PO Box 37012,
Washington, DC 20013, USA.

* corresponding author. m.d.visser@cml.leidenuniv.nl.

Open data statement: Of the 17 datasets used, 10 are published and publicly accessible, with

links provided in this submission (Appendix S1: Section S1). Upon acceptance, remaining seven

datasets will be provided via Smithsonian's Dspace. The open-source model code is available as

R-package ccrtm (https://cran.r-project.org/web/packages/ccrtm/index.html) and on github

(https://github.com/MarcoDVisser/ccrtm). Code will be archived in Zenodo should the

manuscript be accepted for publication

Keywords: leaf optics, leaf traits, leaf angles, radiative transfer models, biodiversity, albedo

Abstract

Lianas, woody vines acting as structural parasites of trees, have profound effects on the composition and structure of tropical forests, impacting tree growth, mortality, and forest succession. Remote sensing offers a powerful tool for quantifying the scale of liana infestation, provided the availability of robust detection methods. We analyze the consistency and global specificity of spectral signals from liana-infested tree crowns and forest stands, examining the underlying mechanisms. We compiled a database, including leaf reflectance spectra from 5424 leaves, fine-scale airborne reflectance data from 999 liana-infested canopies, and coarse-scale satellite reflectance data covering hectares of liana-infested forest stands. To unravel the mechanisms of the liana spectral signal, we applied mechanistic radiative transfer models across scales, corroborated by field data on liana leaf chemistry and canopy structure. We find a consistent liana spectral signature at canopy and stand scales across sites. This signature mainly arises at the canopy level due to direct effects of leaf angles, resulting in a larger apparent leaf area, and indirect effects from increased light scattering in the NIR and SWIR regions, linked to lianas' less costly leaf construction compared to trees. The existence of a consistent global spectral signal for lianas suggests that large-scale quantification of liana infestation is feasible. However, because the traits identified are not exclusive to lianas, accurate large-scale detection requires rigorously validated remote sensing methods. Our models highlight challenges in automated detection, such as potential misidentification due to leaf phenology, tree life-history, topography, and climate, especially where the scale of liana infestation is less than a single remote sensing pixel. The observed cross-site patterns also prompt ecological questions about lianas' adaptive similarities across environments, indicating possible convergent evolution due to shared constraints on leaf biochemical and structural traits.

Introduction

Lianas are woody vines that act as structural parasites of trees and have a strong influence on forest dynamics (Visser et al. 2018b), decreasing tree growth and survival (Ingwell et al. 2010) and arresting forest succession (Tymen et al. 2016). Lianas are increasing in abundance in many forests (Schnitzer and Bongers 2011, Wright et al. 2015), raising concerns about the potential negative impact on carbon sequestration (Durán et al. 2013, van der Heijden et al. 2015). However, the spatial extent of this increase is uncertain, and the causes remain mostly unknown (Schnitzer 2015, Schnitzer et al. 2020). Remote sensing could provide a powerful tool to quantify this trend on a broad range of spatiotemporal scales and elucidate the climatic and bio-geographical factors contributing to liana proliferation (van der Heijden et al. 2022). Yet, as of today we do not know whether a globally distinct liana spectral-signal exists that could make such research possible.

Ecologists have utilized various remote sensing platforms and methods to detect lianas, including multispectral and hyperspectral sensors on drones, aircraft, and satellites (Foster et al. 2008, Tymen et al. 2016, Marvin et al. 2016, Li et al. 2018, Waite et al. 2019, Chandler et al. 2021, Kaçamak et al. 2022). These studies suggest that lianas have distinct detectable spectral signals at the canopy and stand scales. However, differentiating lianas and trees based on the spectral properties of leaves might depend on forest type, with detectable differences observed in dry forests but not in wet forests (Sánchez-Azofeifa et al. 2009a, Asner and Martin 2012, Guzmán Q. et al. 2018) suggesting that climatic factors influence the chemical and structural properties of liana leaves compared to their hosts (Asner and Martin 2012, Werden et al. 2018, Medina-Vega et al. 2021b).

The influence of climate indicates exceptions and contextual dependencies rather than a

distinct global liana spectral signal. This raises doubts about the applicability of automated

classifiers for quantifying liana abundances on large spatiotemporal domains. Data-driven

automated classifiers can capture complex relationships but may struggle to generalize to

scenarios outside the training data, particularly when crucial relationships are missing (Féret et

al. 2019, Meyer and Pebesma 2022, Willard et al. 2022). The major challenge in robustly

applying automated classification techniques is collecting training data representative of all

conditions, which may not be feasible across large spatial domains with different climates or

floristic compositions. Collecting representative training data in tropical forests is especially

daunting due to confounders and high biodiversity. For instance, many leaf and canopy traits are

known to vary systematically among plant groups with life history and leaf phenology, but

differences can be small compared to the large variability across interspecific, intraspecific,

phenotypic, and ontogenetic levels of natural vegetation (Castro-Esau et al. 2004, Kitajima et al.

2005, Zhang et al. 2006, Sánchez-Azofeifa et al. 2009a, Wu et al. 2018, Werden et al. 2018,

Detto and Xu 2020).

At a more fundamental level, pure data-driven automated classifiers do not explain the

underlying reasons responsible for a plant spectral signal. Alternatively, a process-based

modeling approach can provide mechanistic insights into how the signal changes across regions

and scales (Wu et al. 2018). Physics-based models, although unbiased towards out-of-sample,

tend to have lower precision (Féret et al. 2019, Gumiere et al. 2020). A rapidly growing

technique combines both approaches, with so-called physics-informed machine-learning models

showing improved prediction accuracy and precision in situations with low data coverage

(Willard et al., 2022). However, a thorough mechanistic understanding of the liana signal

spanning multiple scales is a prerequisite for applying such hybrid classifiers.

We explore the global liana spectral signal across leaf, canopy, and stand scales in the

visible, near-infrared, and shortwave-infrared spectrum (400 - 2500 nm). Our goal is to

understand the mechanisms behind the spectral signal of liana leaves and canopies, compared to

host trees, by using radiative transfer models to link species traits to reflectance. Our objectives

include: 1) determining if a consistent liana spectral signal exists at various scales using global

datasets and different remote sensing tools; 2) identifying traits that create a liana signal via

radiative transfer modelling; 3) validating results with independent field data; 4) quantifying trait

importance; and 5) evaluating the effectiveness of spectral indices and classifiers in detecting

lianas at different scales using simulated spectra.

Methods

Overview. To test all objectives, we compiled liana reflectance data in the 400 - 2500 nm range

from multiple sites, sensors, and scales (Figure 1, Table S1). We first compared liana reflectance

at the leaf level with other plant growth forms and examined case studies of successful

discrimination between tropical lianas and host trees on canopy and stand scales (objective 1).

Next, we inversely fitted mechanistic radiative transfer models to quantify differences in leaf

biophysical, leaf chemical, and canopy architectural properties between lianas and host trees and

validate them with ground measurements (objectives 2 & 3). We assessed the relative importance

of each trait in generating the liana spectral distribution through iterative model experiments

(objective 4). Finally, using forward simulations of the fitted models, we evaluated the potential

of different platforms to discriminate lianas and trees under various scenarios (objective 5).

This paper integrates diverse data from 17 independent sources spanning multiple sites

and scales. Due to space and legibility concerns, details on data, field, and laboratory protocols

are provided in the supplemental information (SI) and the original published sources.

Study sites, species and data.

Spectral datasets. The spectral data (Figure 1, Table S1) span three distinct spatial scales.

Leaf scale ($<<1m^2$ scale) data encompassed 11 published reflectance spectra datasets

covering 5424 leaves of 720 species. Our focus is primarily on tropical species, with 648 tropical

tree species and 66 liana species. To place liana leaf reflectance globally, we also included data

on 72 crops, shrubs, and herbaceous species in the SI, as all groups could potentially lead to

confounding in large classification studies (e.g., Tropek et al., 2014). In all datasets, reflectance

was measured in the 300 – 2500 nm range (<2 nm step) with laboratory or field

spectroradiometers (Table S1 and Appendix S1). Unique species counts were obtained by cross-

referencing names with the GBIF database and tallying unique GBIF keys. Geographic

distributions, were also sourced from GBIF, and cleaned following Zizka et al. (2019).

Canopy scale (1-4 m^2 scale) data were obtained from two published airborne

hyperspectral campaigns in Panama and Malaysia, covering reflectance from 999 tree crowns

with varying liana infestation levels. Further details on the crown scale data, sites and

instruments can be found in Appendix S1, Chandler et al. (2021) and Marvin et al. (2016).

Stand scale (900 m^2 scale) data used satellite imagery from previous studies to contrast

liana-infested and liana-sparse forest patches in Bolivia and French Guiana (Foster et al., 2008;

Tymen et al., 2016). In both cases, we obtained the original images used, which included a 30-m

resolution Landsat Thematic Mapper (L1TP) image (French Guiana) and hyperspectral Hyperion

imagery from NASA's EO-1 satellite (Bolivia). Reflectance data span over 775 hectares of forest

(40.5 in French Guiana, 734.76 in Bolivia) and was delineated in low liana density forest and

adjacent heavily infested stands (Foster et al., 2008; Tymen et al., 2016). In French Guiana, the

liana forest was delineated using lidar, aerial photography, and field campaigns, while in Bolivia,

it was contrasted with random forest patches using high-resolution videography and field

surveys. Both images were georeferenced and surface reflectance corrected. The Hyperion image

was corrected for the smile effect. Detailed information on the methods and sites can be found in

Tymen et al. (2016), Foster et al. (2008), and Appendix S1.

Leaf and canopy traits

To achieve objectives 2 and 3, we combined published and newly collected field data on leaf and

canopy traits, including data on lianas and host tree species from Panama and Malaysia.

Leaf biochemical and biophysical trait data. We obtained measurements of leaf

Chlorophyll (C_{ab}, μg/cm^2), Carotenoids (C_{ar}, μg/cm^2), water content (C_w, g/cm^2), leaf thickness

(μm) and leaf mass area (C_m, g/cm^2). Trait measurements followed typical protocols where

pigment contents were determined from leaf discs punched from fresh leaves, and water content

was measured by recording fresh weight before drying. A subset of the data included matched

leaf spectral reflectance and physically measured traits for the same leaves (Table 1). The paired

leaf data were used to benchmark the inversion of leaf radiative transfer models (RTMs; see

appendices). Average trait values for lianas and trees were also available for dataset 3 in

Sánchez-Azofeifa et al. (2009a). Finer details on sampling, extraction, and quantification can be

found in Appendix S1.

Plant area index (PAI). The contribution of liana leaves to PAI in infested canopies was

estimated using a plant canopy analyzer (LAI2000, LI-COR) (dataset 16 in Appendix S1). The

estimation capitalized on a liana removal experiment in central Panama, where the contribution

of lianas to PAI was estimated from the difference between control and removal plots (Appendix

1). Details regarding the LI-COR measurements are given in Appendix S1 and Rodríguez-

Ronderos et al. (2016), while van der Heijden et al. (2015) describe the removal experiment.

Note that we use PAI to refer to empirical measurements, and LAI to refer to the PROSAIL

model parameter as per convention.

Leaf angles. Leaf angle distribution was measured for canopies of 40 tree species and 10

liana species in Panama (dataset 17). Samples were taken in the canopies of trees and lianas

using leveled-digital photography (Ryu et al. 2010, Pisek et al. 2011). Images were taken at

different locations, including canopy cranes, telecommunication towers, eddy-covariance flux

towers, and canopy gaps. We measured sun and shaded leaves for trees, but only sun leaves for

lianas, as the vast majority of liana leaves are in the canopy (Avalos et al. 1999, Medina-Vega et

al. 2021a). Leaf angles were defined as the angular orientation of the leaf surface normal relative

to the zenith (Ross 1981) and measured from digital pictures using imageJ (Rueden et al. 2017).

Whenever the leaves were compound, the angle from the petiole to the furthest leaf tip relative to

the zenith was taken as the leaf angle. A total of 1540 individual leaves (646 tree and 894 liana

leaves) were measured.

Radiative transfer models (RTMs)

To address objectives 1-5, we utilized RTMs from the PROSPECT and SAIL families, which

have been extensively validated and applied in reflectance modeling studies (e.g. Zhang et al.

2005, Shiklomanov et al. 2016, Wu et al. 2018). All models were programmed in R, optimized

for speed following Visser et al. (2015), and refactored in C++ using Rcpp (Eddelbuettel and

François 2011). Model code is available as the open-source R-package ccrtm (coupled chain

radiative transfer models; version 0.1.6; Visser, 2020). We provide a short overview as these

models are exhaustively described elsewhere (see e.g. Jacquemoud and Ustin, 2019).

Leaf model. Within the PROSPECT family of models, we evaluated PROSPECT 5, 5b,

and D (Féret et al. 2017). for their ability to capture independently measured trait data. We fitted

each model to reflectance data for individual leaves and selected the model (5, 5b, or D) that

closely reproduced the validation data. Traits, including water mass per unit area, leaf mass area,

chlorophyll content, and carotenoids, were validated using paired reflectance and trait data from

datasets 1, 2, and 11. Leaf thickness (µm) indirectly validated the leaf structure parameter (N) in

datasets 1, 2, 3, and 11. Some parameters, such as brown pigments (C_{br}) and anthocyanin content

(C_{an}), lacked validation data and were treated as calibration parameters: they were only kept if

their inclusion improved both the quality of fit and the validation of other parameters (see also

Féret et al., 2017). Details on the fit procedure are given below, and all parameters and their units

are given in Table 1. Our benchmarks among leaf models showed PROSPECTD performed the

best with regards to reducing errors in both the fit and validation datasets (Appendix S4 & 6) and

therefore further analyses focus solely on PROSPECTD inversions.

Canopy model. We adapted 4SAIL2, which can represent vertically separated vegetation

types while maintaining analytical tractability (Verhoef and Bach 2003, 2007, Zhang et al.

2005). Lianas typically form a canopy on top of their host crown (Stevens 1987, Avalos et al.

1999), with empirical studies finding the vast majority of liana leaves in the upper-most leaf

layers (Medina-Vega et al. 2021b). This vertical mixing of vegetation types has nonlinear

impacts on reflectance, rendering linear mixing inappropriate (Asner 1998). The 4SAIL2 model

simulates the reflectance of two-component canopies with distinct vertically separated layers,

accounting for differences in leaf optics, optical thickness (LAI), and leaf inclination angles

(Braswell et al. 1996). We provide a short description specific to the liana-host tree model

implemented here in Figure 2.

Coupled chain models: the PROSPECT and SAIL families of models can be coupled

together in a chain, typically known as PROSAIL, where the outputs from the former are used as

inputs for the latter. Here, we coupled PROSPECT with 4SAIL2, to simulate canopy and stand-

scale reflectance in liana-infested forests. The full PROSAIL2 model, in its most complete form,

is given by:

$$\hat{\tau}_i, \hat{\rho}_i = PROSPECT(\lambda, N_i, C_{ab,i}, C_{ar,i}, C_{an,i}, C_{br,i}, C_{w,i}, C_{m,i}) \quad i = 1,2 \tag{1a}$$

$$\hat{\rho}_d, \hat{\rho}_s = 4SAIL2(\lambda, \hat{\tau}_1, \hat{\rho}_1, \hat{\tau}_2, \hat{\rho}_2, \text{LAI}, \text{L} \times f_2, \Omega_1, \Omega_2, D, \kappa, C_v, \zeta, \theta_s, \theta_o, \psi) \tag{1b}$$

$$\hat{\rho} = S(\lambda, \theta_s) \hat{\rho}_d + (1 - S(\lambda, \theta_s))\hat{\rho}_s \tag{1c}$$

here $\hat{\tau}_i$ and $\hat{\rho}_i$ are the predicted leaf transmittance and reflectance ($i = 1$ for lianas, 2 for trees) at

wavelength λ, while $\hat{\rho}_s$ and $\hat{\rho}_d$ represent bidirectional (direct) and hemispherical directional

reflectance (diffuse). The parameter f_2 and L divide the total leaf area (LAI) between the top and

bottom layers, with L being the infestation intensity index (ranging between 0 and 1). The optical

thickness of the top liana layer is assumed to be proportional to the liana infestation intensity

(L $\times$ $(1 - f_2)$). This results in a maximum optical thickness of $(1 - f_2)$ for the top layer when

$L = 1$, and the canopy reverts to a single layer with only tree leaves when the infestation is zero

($L = 0$). The fraction of LAI assigned to lianas is then LAI $\times$ L $(1 - f_2)$. The final predicted

reflectance of the target pixel $\hat{\rho}$ depends on the fraction of diffuse light, $S(\lambda, \theta_s)$, at wavelength λ

and solar zenith angle (θ_s) (following Danner et al., 2019; François et al., 2002). Parameters

detail and units are given in Table 1 and Figure 2.

Inverse modelling of the liana signal

Spectral inverse modeling aims to estimate physical parameter values consistent with observed

reflectance data (objective 2). We employed a Bayesian framework for spectral inverse modeling

(Shiklomanov et al. 2016), where coupled chain RTM-models are particularly useful, as models

fitted at one scale can inform models at another scale. We first fit the PROSPECTD model to all

5424 individual leaf-reflectance spectroscopy data, decomposing variation among groupings,

using a hierarchical modelling approach:

$$T_{ijk} \sim \beta_0 + \beta_i + \beta_{jk} + \beta_k + \varepsilon \qquad (2)$$

where T_{ijk} represents one of the RTM model parameters in equation 1a, fit for growth form i

(liana or tree) from species j nested in study k. We included study to account for variation due to

factors associated with specific methodologies, such as the usage of different spectroradiometers

(Meireles et al. 2020). All β parameters were assumed to be normally distributed with

uninformative hyperpriors. We utilized leaf-level means and variances to inform priors for leaf

parameters at canopy and stand scales, assuming them to be normally distributed. The

informative priors for both lianas and trees largely overlapped, with their means falling within

one standard deviation of each other in every instance. All other parameters had uniform flat

priors (see Appendix S2 for details). For models fit at all scales, we sampled the joint posterior

distribution of each RTM with Monte-Carlo Markov Chain (MCMC) methods using the

differential evolution adaptive metropolis algorithm, implemented in the 'BayesianTools' R

package (Hartig et al. 2019). In the inversion procedure we excluded the multispectral data from

French Guiana (dataset 13) because parameter uncertainty is sensitive to spectral resolution

(Shiklomanov et al. 2016). Therefore, we focus on hyperspectral data (datasets 12, 13, and 15),

and included each sensor's spectral response functions in the fitting procedure. Finer details on

model inversion are given in Appendix S2. We conducted several model robustness checks,

including simulations to determine parameter identifiability and the effect of measurement error

(Appendix S3).

Spectral differences and the detection of the liana signal

Kullback–Leibler divergence (KLD) is a measure from information theory that quantifies the

difference between probability distributions, commonly used for optimization in machine

learning (Kosheleva and Kreinovich 2017). As a measure of feature discernibility for classifiers,

KLD helps explore differences in spectral distribution between plant groups, the impact of traits

on spectral distributions, and the theoretical ability to distinguish lianas from trees (objectives 1,

4, 5). At the leaf level, we used KLD to compare different plant groups, with tropical liana leaves

as the reference (0). KLD is interpreted as the information lost when encoding the liana spectral

distribution (0) with another plant group's distribution (k). KLD was defined as:

$$d_{k0}(u_k | u_0) = \int\limits_{400}^{2500} \int\limits_{0}^{1} u_k(\rho, \lambda) \log \frac{u_k(\rho, \lambda)}{u_0(\rho, \lambda)} d\lambda d\rho \tag{3}$$

281   where $u_k(\rho,\lambda)$ is the probability of observing a reflectance value ρ at wavelength λ, and $u_0(\rho,\lambda)$ is

282   a reference distribution. Assuming normality, the spectral distribution can be estimated with the

283   wavelength specific mean and standard deviation, $\mu(\lambda)$ and $\sigma(\lambda)$ respectively. In this case KLD

284   (eqn. 3) reduces to:

$$d_{k0}(u_k | u_0) = \int\limits_{400}^{2500} log \left(\frac{\sigma(\lambda)_0}{\sigma(\lambda)_k} + \frac{\sigma(\lambda)_0^2 + (\mu(\lambda)_k - \mu(\lambda)_0)^2}{2\sigma(\lambda)_0^2} - \frac{1}{2} \right) d\lambda \tag{4}$$

285   The smaller the KLD value, the more comparable the groups are, decreasing the likelihood of

286   robust classification (Kosheleva and Kreinovich 2017). The expected KLD between two

287   identically distributed signals with equal means and variances is, by definition, zero.

288   *Model sensitivity experiment.* KLD quantifies the loss or gain of information due to

289 individual traits by comparing modelled spectral distributions that include or exclude one or

290 more parameters. This provides a metric of individual traits' importance (objective 4). KLD was

291 calculated with predicted reflectance $\hat{\rho}(\lambda)$ as the mean and variance estimated from the residual

292 error from the inverse fits (σ, Appendix S2). Starting with a model parameterized with mean tree

293 leaf and canopy traits as the focal distribution $u_k(\hat{\rho},\lambda)$, we iteratively changed single parameters

294 to liana-specific values and calculated the change in KLD using the liana spectral distribution as

295 the reference distribution $u_0(\hat{\rho},\lambda)$. This determines the parameters with the greatest information

296 loss or gain towards the liana spectral distribution. To understand influential parameters at

297 different wavelengths, we also evaluated the KLD at each wavelength separately and assessed

298 combinations of two parameters to identify interactions that most strongly recreated the liana

299 pattern.

300   *The theoretical ability of classifiers to detect the liana signal.* In theory, sensors and

301 spectral indices that sample at bands maximizing KLD between trees and lianas optimize

302 discernibility between these groups. Hence, sensor sampling regions of maximum KLD are

303 expected to provide optimal information for modeling and classification (Wang et al. 2015).

304 Therefore, we compare how well the spectral response functions of several common

305 multispectral remote sensing platforms maximize KLD. Given the strong spatial clustering of

306 lianas at different scales (Ledo and Schnitzer 2014) and their varying infestation patterns from a)

307 incidental crown infestation (Marvin et al. 2016) to b) arrested succession (Schnitzer et al. 2000)

308 and c) liana forests (Tymen et al. 2016), we investigate how the spatial aggregation scale

309 interacts with sensor spatial resolution in determining the expected KLD. We simulated

310 1000×1000 pixel hyperspectral image scenes with various scales and autocorrelation degrees in

liana infestation for scenarios (a-c). We started with the finest scale (1 m^2) as pure signals and

then recalculated the expected KLD between feature (liana) and non-feature (liana-free) areas,

gradually reducing spatial resolution from 1×1 m to 250×250 m via resampling, repeating this 50

times per scenario. Each scenario maintained a roughly equal number of feature pixels. The

sensors modeled included Landsat TM, Worldview 2's WV110 camera, and Hyperion.

Note that we focus solely on the aspects of sensor choice readily controllable by the user.

We explicitly ignore atmospheric effects that decrease the signal-to-noise ratio of most sensors

(Landgrebe and Makaret 1986). Consequently, the results should be considered a "best-case

scenario".

320   **Results**

Objective 1: Does a globally consistent liana signal exist at the leaf, canopy and stand scales?

The distribution of liana leaf-scale reflectance, measured from 718 leaves from 66 species,

exhibited significant overlap with trees, shrubs, crops, and grasses (Figure S6). This finding was

further supported by inversely fit PROSPECT leaf models (Figure 3 A-C). Kullback-Leibler

divergence (KLD) analyses on spectral distributions and predicted posterior distributions

confirmed the indistinguishable nature of tree and liana leaf reflectance across all wavelengths

(Figure 3). In contrast to leaf reflectance, however, predicted posterior distributions of leaf

transmission and absorption demonstrated substantial differences, particularly in the NIR and

SWIR regions (Figure 3 E-J). Liana leaves had higher transmission and lower absorption

throughout the evaluated spectrum, leading to pronounced distributional divergence (Figure 3 G

& J).

Liana-infested canopies showed consistently higher reflectance than liana-free canopies

at both canopy and stand scales across various sites, platforms, and wavelengths (Figure 1B &

C). The reflectance difference ($\mathbf{R_{lianas}}$ - $\mathbf{R_{trees}}$) had an average 89% correlation between sites

(range 74% to 99%) when matched to Landsat 5TM bands (used in French Guiana). For

hyperspectral images at overlapping wavelengths, the mean correlation for reflectance difference

was 96% (range 95%-98%). These results reveal a distinct and detectable spectral signal for liana

reflectance at canopy and stand scales, stable across instrumental noise and sensor differences.

Objective 2: Inverse modelling of the liana spectral signal at the leaf, canopy and stand scales.

The leaf signal was accurately reproduced by PROSPECTD exhibiting a low overall error (R^2 >

0.94 for 5424 individual leaves; Appendix S4). Inverse fits of the PROSAIL2 model to canopy

and stand scale data closely approximated the spectral reflectance of heavily infested (>50%

cover) and lightly infested canopies (<50%) in Bolivia, Malaysia, and Panama (R^2 > 0.98; Figure

4A-C). The model also successfully replicated the difference in reflectance between heavily and

lightly infested canopies ($\mathbf{R_{lianas}}$ - $\mathbf{R_{trees}}$), explaining between 46% and 92% of the variation (inset

plots in Figure 4A-C). Model posteriors can be found in Appendix S5.

Objective 3: Model validation.

The PROSPECTD model, fitted to leaf reflectance data, accurately and unbiasedly replicated

independently measured leaf traits such as C_{ar}, C_{ab}, C_m, and C_w, with observed and predicted

traits showing agreement ranging from R^2 = 0.4 to 0.73 (Appendix S6). The leaf structural

parameter strongly correlated with leaf thickness (R^2 = 0.54), suggesting it captured meaningful

physical variation.

Figure 5 depicts the predicted differences between lianas and trees based on inverse fits

of PROSAIL2 (black dots and whiskers) for parameters showing non-overlapping credible

intervals (CI) at least once across sites. The pattern of differences between lianas and trees in

inversely estimated parameters was generally consistent across sites, although not all parameters

exhibited significant differences at all sites based on 95% CI (Figure 5). Regarding leaf

biophysical and chemical parameters, all models indicated that lianas had more cheaply

constructed leaves on average, characterized by thinner leaves, lower photosynthetic pigment

content (Chlorophyll and Carotenoids) per unit leaf area, and lower dry mass and water mass per

unit leaf area. Note that water mass C_w (g/cm²) should not be equated with leaf water content

(%): lianas had higher water content compared to trees (77% vs 70% on average). Inverse fits

with PROSPECTD confirmed the independent predictions of PROSAIL2 whenever leaf-level

spectral data were available (blue dots and whiskers in Figure 5), except for C_m in Malaysia,

which was lower according to PROSAIL2 compared to C_m estimated from leaf reflectance

(PROSPECTD) and direct measurements. However, all estimates of C_m in Malaysia aligned with

the general trend of lower leaf mass per unit area for lianas compared to trees (Figure 5). At the

canopy and stand scales, the models predicted that trees contribute 66% (f_2) to the total LAI

across sites. Indicating that liana leaves contributed an average of 34% ($1-f_2$) when a tree crown

is fully infested, with site averages ranging from 16% to 55% (Figure 5). Liana leaves were also

predicted to have leaf angles that were 23.5% flatter compared to trees on average, with

parameter Ω_1 being 4% to 59% lower than Ω_2 across sites (Figure 5).

The inverse fits with the PROSAIL2 model were supported by independently collected

verification data where available. Lab measurements of leaf thickness, Chlorophyll and

Carotenoids content, dry mass, and water content per unit area for liana leaves were all found to

be lower on average for liana leaves (red dots and whiskers in Figure 5). Canopy and stand scale

validation data also showed consistency with the model predictions. The estimation of the

fraction of PAI attributed to trees vs lianas (f_2 vs $1-f_2$) using LI-COR data in plots where lianas

were removed showed that trees contribute 65.3% ($f_2 = 0.653 \pm 0.087$ SE) to the PAI when their

canopies are infested. Indicating that lianas contribute on average 34.7% - a value close to the

inverse model estimate. Additionally, the inverse model results predicted flatter leaf angles for

all sites (Figure5), and measurements conducted in Panama confirmed that lianas have, on

average, 26.2% flatter leaf angles compared to trees (mean angle of $27.9° \pm 0.76$ SE vs $37.8° \pm$

0.64 SE).

Objective 4: what causes the liana signal?

The influence of parameters on the liana spectral distribution varied depending on the solar

spectrum region (Figure 6A). In the visible spectrum, leaf angles (Ω) and photosynthetic pigment

content (C_{ar}, C_{ab}) played significant roles, with flatter leaf angles and lower pigment content

leading to increased reflectance. In the near-infrared, leaf angles (Ω), leaf mass area (C_m), and

leaf structure (N) consistently showed importance, although the relative significance of leaf

angles (Ω) versus leaf mass area (C_m) differed among sites (Figure S8A, C, E). Leaf water

content (C_w) became increasingly influential with wavelength and dominated the signal in the

short-wave infrared (Figure 6A), particularly at the tropical dry site in Bolivia (Figure S8A). The

most influential traits overall were leaf mass area (C_m), leaf angles (Ω), and leaf water content

(C_w) (Figure6B). Leaf angles (Ω) were consistently important across sites, while the importance

of C_m and C_w varied among sites (Figure S8B, D, F). Substituting the leaf structural parameter

(N) of trees with that of lianas resulted in predictions further deviating from the liana spectral

distribution (negative contribution), especially when interacting with other parameters (light

colors in Figure 6B).

Objective 5: theoretical limits to the detection of liana.

The KLD between liana-infested and liana-free spectral distributions was highest in the NIR and

SWIR bands, which are covered by several multispectral spaceborne sensors (Figure 7A). The

performance of a sensor in discriminating lianas from trees improved with an increased number

of bands within the optimal range (Figure 7B). The spatial resolution of the sensor interacts with

liana spatial aggregation (Figure 7C-E). In a crown infestation scenario (liana pixel aggregation

~ 350 m²), the divergence between feature and background decreases rapidly (Figure 7C). For

larger spatial aggregation resembling arrested forest succession (2000 m²), the divergence

dropped at a similar pace to random feature pixels (Figure 7D). For highly aggregated lianas

across extensive forest areas (~30,000 m²), the divergence decreased slower than the random

expectation.

Discussion

We demonstrate a unique spectral distribution in liana-infested canopies and forest-stands

(Figures 1, 4 & 7), showing higher reflectance than liana-free areas (Figure 1 & 4), particularly

in the NIR and SWIR spectrum sections (Figure 7). This distinct spectral distribution, observed

across four global sites (Figure 1), has implications for liana evolution and forest energy budgets

- and provides a basis for expanded research on lianas at pantropical scales. However, the traits

underpinning this distribution might not be exclusive to lianas, posing a clear risk of confusion

with other phenomena.

The mechanisms behind the liana spectral signal

Our analysis reveals a consistent liana spectral distribution across sites, observable at larger

scales (≥ 1 m^2; Figure 1 & 4), but absent in leaf-scale reflectance ($<< 1$m^2; Figure 3). Model

predictions indicate leaf traits alone do not account for the consistent reflectance trend between

liana and tree leaves across sites (Figure 2A & Figure S7). Previous research confirms varying

reflectance distributions and optical traits among liana species and sites (Sánchez-Azofeifa et al.,

2009a; Asner and Martin, 2012; Medina-Vega et al., 2021a). Therefore, we conclude that leaf

reflection alone is unlikely to be critical. In contrast, differences in leaf traits between lianas and

trees do affect transmission and absorption profiles (Figure 2) and, by interacting with canopy

traits, likely drive the distinct distribution observed across multiple sites and platforms (Figure 1

& 4). Our findings are supported by radiative transfer model inversions at the canopy and stand

scales (Figure 5 & 6).

Differences in leaf traits between lianas and trees yield consistent trends in leaf

absorption and transmission (Figure 2), and we predict lower leaf absorption and increased

transmission for liana leaves at the canopy scale across evaluated sites (Figure S7). Plausible

mechanisms for the distinct liana spectral distribution involve reduced light extinction rates in

liana-infested canopies. In addition, our model experiment highlights the impact of flatter leaf

angles in liana leaves compared to host trees (Figure 5, 6). We propose a two-component

mechanism for the "liana signal": 1) higher projected leaf area, which, at equal leaf reflectance,

leads to increased reflected energy at the canopy level, and 2) multiple scattering from greater

transmission and lower leaf absorption, which indirectly increases reflectance. The latter is an

effect similar to what is hypothesized to cause canopy greening (Wu et al. 2018).

Building on our mechanistic models, we can derive theoretical expectations for the

generation of liana canopy-level spectral distribution. The first component arises as lianas retain,

on average ~ 24% flatter leaf angles (27.9^o vs 37.8^o), which should result in roughly an 11%

increase in projected leaf area as seen from a polar angle and a proportional increase in the

reflected energy. The second "multiple scattering" component arises from liana leaves having

lower average absorption (Figure 3 & S7). This results in slower decay of light energy

(Appendix S9) and increased scattering, including in the observer's direction, leading to higher

reflectance when integrated across the canopy. Multiple scattering also explains the regions

where the liana spectral distribution differs most from trees. Since the projected leaf area does

not change with wavelength, its importance is relatively uniform across the spectrum (Figure6A,

parameter Ω). The importance of multiple scattering, however, peaks when leaf transmission is

maximized, in the SWIR and NIR parts of the spectrum. This corresponds to the point of

maximal discernibility of the liana distribution (Figure 7), and where leaf traits appear the most

influential in our sensitivity analysis (Figure 6).

In conclusion, a distinct liana spectral distribution emerges from interactions between

multiple leaves within canopies. This multi-leaf dependency explains why the signal is observed

at the canopy scale (Figure1 & 4; Chandler et al 2021, Waite et al 2020, Foster et al 2008,

Tymen et al 2016) and the lack of a consistent signal when only leaf reflectance is considered

(Figure3, Sánchez-Azofeifa et al., 2009a).

Ecological consequences of the liana spectral signal

By increasing forest canopy albedo (Figure 2), lianas have the potential to impact forest energy

budgets and understory light conditions. As tropical forests are typically light-limited and liana

abundance is increasing in the Neotropics (Schnitzer and Bongers 2011, Wright et al. 2015),

liana proliferation could have broader impacts beyond current knowledge.

Meunier et al. (2021) integrated liana optical properties into the ED2 dynamic vegetation

model suggesting several consequences. With approximately 25% liana leaf area, the model

predicted an increase in reflected energy of 11.7-17.1% across the PAR and IR spectrum. This

resulted in a reduction of 3.6 Wm^{-2} in absorbed energy, leading to darker understories (up to 50%

darker) and slightly cooler soils (-0.5°C in the topsoil). These findings highlight the broader

impacts of the liana spectral signal on forest functioning, including temperature-dependent

processes such as metabolic respiration and photosynthesis. Our results support Meunier et al.'s

findings and indicate a global characteristic of higher albedo in liana-infested areas. We believe

this shows a direct and urgent need for empirical validation of these model predictions to better

understand the potential long-term changes in forested ecosystems with increasing liana

prevalence.

The evolution of a global liana spectral distribution.

Lianas consistently display cost-effective leaves with flatter angles than their tree hosts, as

shown across diverse tropical regions in Asia and South America (Figure 5). This pattern,

uniformly revealed through our inverse modeling at leaf, canopy, and stand scales, is notable

given lianas' independent evolution in various angiosperm families (Putz and Mooney 1991,

Gentry 1992). This prompts a fundamental question: why do lianas possess a similar spectral

signal across vastly different sites? We propose that this phenomenon may be an example of

convergent evolution.

Lianas rely on structural parasitism to access the heights and light conditions required for

seed production and dispersal. This, however, increases their host's mortality (Ingwell et al.

2010, Visser et al. 2018a), which consequently shortens their residency in the canopy. This

strongly mirrors pathogen virulence evolution (Anderson and May 1982), and evolutionary

theory suggests that lianas should, therefore, evolve to be less harmful to hosts, extending their

canopy residency and reproductive time. Such adaptation would involve slower growth and

reduced resource usurpation from hosts, entailing minimized host-shading (e.g. Ichihashi and

Tateno 2011). Slow growth implies greater investment in leaves (Osnas et al. 2013) and steeper

leaf angles for better light penetration, akin to tree strategies that optimize their leaf angles to

allow optimal light levels to reach lower leaf layers (Falster and Westoby 2003). The fact that

these expectations are not realized, is probably related to the fact that lianas seldom exploit their

hosts in isolation: tree canopies are typically infested by two or more liana species (Visser et al.

2018b).

In such multi-parasite systems, theory predicts a "tragedy of the commons", where any

parasite limiting host exploitation becomes susceptible to invasion by a more exploitative variant

(Frank 1996). This leads to heavy competition for resources and time at the expense of the host:

reproductive success requires maximized light capture and rapid growth to avoid being

overshadowed by other lianas. We suggest that this "tragedy of the commons" is a common

evolutionary force among forest lianas, driving rapid growth, turnover and metabolism and

resulting in lower leaf longevity and construction costs among sun-exposed lianas compared to

trees (Slot et al. 2013, Wyka et al. 2013). In contrast, the most common hosts for lianas are

shade-tolerant trees (Visser et al., 2018a), which typically have costlier, long-living leaves and

high Leaf Area Index (LAI) values (up to 5 in an individual canopy; Kitajima 1994) - creating a

distinct spectral contrast to lianas.

Robust detection of lianas.

Our simulations highlight two primary insights regarding liana detection. First, many sensor

platforms often miss optimal bands distinguishing lianas from trees. Satellite platforms might

possess just one optimal band (Figure 7A). Second, as crown infestation levels increase, the

liana spectral distribution decays non-linearly (Figure 7C-E). High spatial resolution is essential

for detecting lianas within tree crowns, where signal weakening due to spectral mixing occurs

faster than expected (Figure 7C). This means that moderate-scale platforms (around 20-30m

resolution) are less suitable for detecting crown infestations but should be able to identify larger

liana-infested regions where the signal decays slower (Figure7E). Our simulation results align

with previous studies. These two simulation findings echo patterns from earlier studies.

Moderate resolution platforms like Hyperion and Landsat effectively identified large liana-

dominated areas (Foster et al. 2008, Tymen et al. 2016), whereas high-resolution hyperspectral

images were best for detecting heavy tree crown infestations (Marvin et al. 2016, Chandler et al.

2021). Hence, our findings can inform future liana remote sensing research and sensor selection.

However, caution is advised as this analysis, being a "best-case scenario", overlooks atmospheric

factors potentially weakening satellite signals (Landgrebe and Makaret 1986).

Confounding variables. Our analysis of the liana spectral distribution prompts careful use of

classifiers due to potential signal confusion. Notable confounders are:

       1) Leaf phenology. Seasonal shifts in phenology can confound classification. Like lianas,

528   younger tree leaves exhibit lower C_m, C_w, and pigment concentrations, causing them to

529   resemble liana signals by being more transmissive (Wu et al. 2018).

       2) Tree life history. Light-demanding species possess flatter leaf angles and thinner leaves,

531   risking liana misclassification. Concurrently, liana infestation can obscure tree signals,

532   causing mapping inaccuracies (Zhang et al. 2006).

       3) Topography and sun-sensor geometry. Another possible confounder is areas sloped

534   towards the sun. These will appear to have relatively higher reflectance across the

535   spectrum.

       4) Climate. Some key leaf-trait differences between lianas and trees identified here, lower

537   C_m, and C_w, remain but become less pronounced in wetter climates (Sánchez-Azofeifa et

538   al. 2009a, Asner and Martin 2012, Medina-Vega et al. 2021b). As these traits are mostly

539   responsible for the maximal divergent signal from trees in the NIR and SWIR portions

(Figure7), the spectral difference between liana-infested and free canopies may also

become less pronounced.

These confounding factors in detecting lianas can be addressed uniquely. Seasonal leaf

phenology changes contrast with lianas' longer-term growth, aiding detection, especially in dry

forests where lianas retain leaves longer (Schnitzer 2005). Light-demanding tree species, less

prone to liana infestation and with fewer leaf layers (Kitajima 1994, Visser et al. 2018a), can be

distinguished using lidar structural data alongside spectral information (sensu Rao et al., 2020).

This approach is particularly relevant given the substantial structural changes caused by liana

infestation, such as canopy wood area index alteration (Sánchez-Azofeifa et al. 2009b).

Furthermore, topography and sun-sensor geometry issues might be mitigated with multiangle

sensors (i.g. CHRIS Proba-1) or matching-resolution digital elevation models. Thus, automated

classifiers should be applied with caution over large areas, ensuring robustness against these and

other confounding factors.

Caveats and limitations

The main result from model inversion at three sites was backed up by independent measurements

in the field and previous studies (Sánchez-Azofeifa and Castro-Esau 2007, Mello et al. 2020).

Separate measurements on leaf reflectance and in situ measurements of leaf traits and angles

confirmed that lianas tend to have relatively flatter leaf angles and more cheaply constructed

leaves with less dry matter and pigments. However, the model's predictions were not always

perfectly in line with ground truth data. On an absolute scale, ground measurements of mean leaf

angles were lower than those predicted by PROSAIL2 (Appendix S6). One explanation for this

is that we do not consider the effect of leaf clustering on stems and branches in our field

campaign, while the PROSAIL2 result is more likely to reflect the integrated angles of leaves

clustered on stems and branches. Direct measurements (i.e. not derived from remotely sensed

data) on leaf angles, stems and branches collected from multiple sites would confirm the

robustness of this result.

Conclusions

Our work reveals a unique spectral signature for lianas at the canopy scale, detectable by aerial

and spaceborne sensors. Models attribute the signature to lianas' larger projected leaf area, lower

leaf light absorption and higher transmission, leading to increased canopy-level light scattering.

We observe the signal, characterized by higher forest albedo, in four distinct tropical sites,

suggesting a consistent global pattern. If further confirmed, this could significantly impact our

understanding of liana proliferation's effects on forest energy budgets and understory dynamics.

The identification of a general spectral signal for liana infestation across sites underscores the

importance of developing accurate classifiers for quantifying liana abundance. Our successful

use of radiative transfer models highlights the potential for creating robust, potentially physics-

informed classifiers. In conclusion, fertile ground exists for future research into the liana spectral

signal, its robust detection from remote sensing platforms, and the ecological consequences of

liana proliferation.

Acknowledgements.

We thank Princeton University's Carbon Mitigation Initiative, the European Research Council

Grant 637643 (TREECLIMBERS), FWO (grants 1214720N/1214723N), and various Natural

Environment Research Council grants [NE/P004806/1; NE/I528477/1; NE/L002604/1], along

with the University of Nottingham Anne McLaren Research Fellowship. We acknowledge the

References

Anderson, R. M., and R. M. May. 1982. Coevolution of hosts and parasites. Parasitology

85:411–426.

Asner, G. P. 1998. Biophysical and biochemical sources of variability in canopy reflectance.

Remote Sensing of Environment 64:234–253.

Asner, G. P., and R. E. Martin. 2012. Contrasting leaf chemical traits in tropical lianas and trees:

Implications for future forest composition. Ecology Letters 15:1001–1007.

Avalos, G., S. Mulkey, and K. Kitajima. 1999. Leaf optical properties of trees and lianas in the

outer canopy of a tropical dry forest. Biotropica 31:517–520.

Braswell, B. H., D. S. Schimel, J. L. Privette, B. Moore, W. J. Emery, E. W. Sulzman, and A. T.

Hudak. 1996. Extracting ecological and biophysical information from AVHRR optical data:

An integrated algorithm based on inverse modeling. Journal of Geophysical Research

Atmospheres 101:23335–23348.

Castro-Esau, K. L., G. A. Sánchez-Azofeifa, and T. Caelli. 2004. Discrimination of lianas and

trees with leaf-level hyperspectral data. Remote Sensing of Environment 90:353–372.

Chandler, C. J., G. M. F. Van Der Heijden, S. Doreen, C. E. Waite, M. E. J. Cutler, H. Costa, R.

Nilus, and G. M. Foody. 2021. Remote sensing liana infestation in an aseasonal tropical

forest: addressing mismatch in spatial units of analyses. Journal of remote sensing and

conservation.

Danner, M., K. Berger, M. Wocher, W. Mauser, and T. Hank. 2019. Fitted PROSAIL

Parameterization of Leaf Inclinations, Water Content and Brown Pigment Content for

Winter Wheat and Maize Canopies. Remote Sensing 11:1150.

Detto, M., and X. Xu. 2020. Optimal leaf life strategies determine $V_{c,max}$ dynamic during

ontogeny. New Phytologist 228:361–375.

Durán, S. M., E. Gianoli, S. M. Duran, and E. Gianoli. 2013. Carbon stocks in tropical forests

decrease with liana density. Biology letters 9:20130301.

Eddelbuettel, D., and R. François. 2011. Rcpp: Seamless R and C++ integration. Journal of

Statistical Software 40:1–18.

Falster, D. S., and M. Westoby. 2003. Plant height and evolutionary games. Trends in Ecology

and Evolution 18:337–343.

Féret, J. B., A. A. Gitelson, S. D. Noble, and S. Jacquemoud. 2017. PROSPECT-D: Towards

modeling leaf optical properties through a complete lifecycle. Remote Sensing of

Environment 193:204–215.

Féret, J. B., G. le Maire, S. Jay, D. Berveiller, R. Bendoula, G. Hmimina, A. Cheraiet, J. C.

Oliveira, F. J. Ponzoni, T. Solanki, F. de Boissieu, J. Chave, Y. Nouvellon, A. Porcar-

Castell, C. Proisy, K. Soudani, J. P. Gastellu-Etchegorry, and M. J. Lefèvre-Fonollosa.

2019. Estimating leaf mass per area and equivalent water thickness based on leaf optical

properties: Potential and limitations of physical modeling and machine learning. Remote

Sensing of Environment 231:110959.

Foster, J. R., P. A. Townsend, and C. E. Zganjar. 2008. Spatial and temporal patterns of gap

dominance by low-canopy lianas detected using EO-1 Hyperion and Landsat Thematic

Mapper. Remote Sensing of Environment 112:2104–2117.

François, C., C. Ottlé, A. Olioso, L. Prévot, N. Bruguier, and Y. Ducros. 2002. Conversion of

400-1100 nm vegetation albedo measurements into total shortwave broadband albedo using

a canopy radiative transfer model. Agronomie 22:611–618.

Frank, S. a. 1996. Models of parasite virulence. The Quarterly review of biology 71:37–78.

Gentry, A. H. 1992. The distribution and evolution of climbing plants. Pages 3–49 *in* F. E. Putz

and H. A. Mooney, editors. The biology of vines. Cambridge University Press, Cambridge,

UK.

Gumiere, S. J., M. Camporese, A. Botto, J. A. Lafond, C. Paniconi, J. Gallichand, and A. N.

Rousseau. 2020. Machine Learning vs. Physics-Based Modeling for Real-Time Irrigation

Management. Frontiers in Water 0:8.

Guzmán Q., J. A., B. Rivard, and G. A. Sánchez-Azofeifa. 2018. Discrimination of liana and tree

leaves from a Neotropical Dry Forest using visible-near infrared and longwave infrared

reflectance spectra. Remote Sensing of Environment 219:135–144.

Hartig, F., F. Minunno, S. Paul, and D. Cameron. 2019. BayesianTools: General-Purpose

MCMC and SMC Samplers and Tools for Bayesian Statistics. R package version 0.1.7.

van der Heijden, G. M. F., A. D. C. Proctor, K. Calders, C. J. Chandler, R. Field, G. M. Foody,

S. M. Krishna Moorthy, S. A. Schnitzer, C. E. Waite, and D. S. Boyd. 2022. Making

(remote) sense of lianas. Journal of Ecology 110:498–513.

van der Heijden, G. M., J. S. Powers, and S. A. Schnitzer. 2015. Lianas reduce carbon

accumulation and storage in tropical forests. Proceedings of the National Academy of

Sciences 112:13267–13271.

Ichihashi, R., and M. Tateno. 2011. Strategies to balance between light acquisition and the risk

of falls of four temperate liana species: To overtop host canopies or not? Journal of Ecology

99:1071–1080.

Ingwell, L. L., S. J. Wright, K. K. Becklund, S. P. Hubbell, and S. A. Schnitzer. 2010. The

impact of lianas on 10 years of tree growth and mortality on Barro Colorado Island,

Panama. Journal of Ecology 98:879–887.

Jacquemoud, S., and S. L. Ustin. 2019. Leaf Optical Properties. Cambridge University Press,

Cambridge.

Kaçamak, B., N. Barbier, M. Aubry-Kientz, E. Forni, S. Gourlet-Fleury, D. Guibal, J. J.

Loumeto, S. Pollet, V. Rossi, N. Rowe, Y. van Hoef, and M. Réjou-Méchain. 2022. Linking

Drone and Ground-Based Liana Measurements in a Congolese Forest. Frontiers in Forests

and Global Change 5:31.

Kitajima, K. 1994. Relative importance of photosynthetic traits and allocation patterns as

correlates of seedling shade tolerance of 13 tropical trees. Oecologia 98:419–428.

Kitajima, K., S. Mulkey, and S. Wright. 2005. Variation in crown light utilization characteristics

among tropical canopy trees. Annals of Botany 95:535–547.

Kosheleva, O., and V. Kreinovich. 2017. Why Deep Learning Methods Use KL Divergence

Instead of Least Squares: A Possible Pedagogical Explanation Recommended Citation.

Page Part of the Computer Sciences Commons Comments.

Landgrebe, D. A., and E. Makaret. 1986. Noise in Remote-Sensing Systems: The Effect on

Classification Error. IEEE Transactions on Geoscience and Remote Sensing GE-24:294–

300.

Ledo, A., and S. A. Schnitzer. 2014. Disturbance and clonal reproduction determine liana

distribution and maintain liana diversity in a tropical forest. Ecology 95:2169–2178.

Li, W., C. Campos-Vargas, P. Marzahn, and A. Sanchez-Azofeifa. 2018. On the estimation of

tree mortality and liana infestation using a deep self-encoding network. International

Journal of Applied Earth Observation and Geoinformation 73:1–13.

Marvin, D. C., G. P. Asner, and S. A. Schnitzer. 2016. Liana canopy cover mapped throughout a

tropical forest with high-fidelity imaging spectroscopy. Remote Sensing of Environment

176:98–106.

Medina-Vega, J. A., F. Bongers, L. Poorter, S. A. Schnitzer, and F. J. Sterck. 2021a. Lianas have

more acquisitive traits than trees in a dry but not in a wet forest. Journal of Ecology

109:2367–2384.

Medina-Vega, J. A., F. Bongers, S. A. Schnitzer, and F. J. Sterck. 2021b. Lianas explore the

forest canopy more effectively than trees under drier conditions. Functional Ecology

35:318–329.

Meireles, J. E., J. Cavender-Bares, P. A. Townsend, S. Ustin, J. A. Gamon, A. K. Schweiger, M.

E. Schaepman, G. P. Asner, R. E. Martin, A. Singh, F. Schrodt, A. Chlus, and B. C.

O'Meara. 2020. Leaf reflectance spectra capture the evolutionary history of seed plants.

New Phytologist 228:485–493.

Mello, F. N. A., S. Estrada-Villegas, D. M. Defilippis, and S. A. Schnitzer. 2020. Can Functional

Traits Explain Plant Coexistence? A Case Study with Tropical Lianas and Trees. Diversity

2020, Vol. 12, Page 397 12:397.

Meunier, F., M. D. Visser, A. Shiklomanov, M. C. Dietze, J. Q. Antonio Guzmán, A. Sanchez-

Azofeifa, H. P. T De Deurwaerder, S. M. Krishna Moorthy, S. A. Schnitzer, D. C. Marvin,

M. Longo, L. Chang, E. N. Broadbent, A. M. Almeyda Zambrano, H. Muller-Landau, M.

Detto, and H. Verbeeck. 2021. Liana optical traits increase tropical forest albedo and reduce

ecosystem productivity. bioRxiv.

Meyer, H., and E. Pebesma. 2022. Machine learning-based global maps of ecological variables

and the challenge of assessing them. Nature Communications 2022 13:1 13:1–4.

Osnas, J. L. D. J., J. W. J. Lichstein, P. P. B. Reich, and S. W. S. Pacala. 2013. Global leaf trait

relationships: mass, area, and the leaf economics spectrum. Science 340:741–4.

Pisek, J., Y. Ryu, and K. Alikas. 2011. Estimating leaf inclination and G-function from leveled

digital camera photography in broadleaf canopies. Trees - Structure and Function 25:919–

924.

Putz, F. E., and H. A. Mooney. 1991. The Biology of Vines. Cambridge University Press,

Cambridge, United Kingdom.

Rao, K., A. P. Williams, J. F. Flefil, and A. G. Konings. 2020. SAR-enhanced mapping of live

fuel moisture content. Remote Sensing of Environment 245:111797.

Rodríguez-Ronderos, M. E., G. Bohrer, A. Sanchez-Azofeifa, J. S. Powers, and S. A. Schnitzer.

2016. Contribution of lianas to plant area index and canopy structure in a Panamanian

forest. Ecology 97:3271–3277.

Ross, J. 1981. The radiation regime and architecture of plant stands. Page The radiation regime

and architecture of plant stands. Springer Netherlands, The Hague.

Rueden, C. T., J. Schindelin, M. C. Hiner, B. E. DeZonia, A. E. Walter, E. T. Arena, and K. W.

Eliceiri. 2017. ImageJ2: ImageJ for the next generation of scientific image data. BMC

Bioinformatics 18:529.

Ryu, Y., O. Sonnentag, T. Nilson, R. Vargas, H. Kobayashi, R. Wenk, and D. D. Baldocchi.

2010. How to quantify tree leaf area index in an open savanna ecosystem: A multi-

instrument and multi-model approach. Agricultural and Forest Meteorology 150:63–76.

Sánchez-Azofeifa, G. A., K. Castro, S. J. Wright, J. Gamon, M. Kalacska, B. Rivard, S. A.

Schnitzer, and J. L. Feng. 2009a. Differences in leaf traits, leaf internal structure, and

spectral reflectance between two communities of lianas and trees: Implications for remote

sensing in tropical environments. Remote Sensing of Environment 113:2076–2088.

Sánchez-Azofeifa, G. A., and K. Castro-Esau. 2007. Canopy observations on the hyperspectral

properties of a community of tropical dry forest lianas and their host trees Canopy

observations on the hyperspectral properties of a community of tropical dry forest lianas.

International Journal of Remote Sensing 27:2101–2109.

Sánchez-Azofeifa, G. A., M. Kalácska, M. M. do Espírito-Santo, G. W. Fernandes, and S.

Schnitzer. 2009b. Tropical dry forest succession and the contribution of lianas to wood area

index (WAI). Forest Ecology and Management 258:941–948.

Schnitzer, S. A. 2005. A Mechanistic Explanation for Global Patterns of Liana Abundance and

Distribution. the american naturalist 166.

Schnitzer, S. A. 2015. Increasing liana abundance in neotropical forests: causes and

consequences. Pages 451–464 *in* S. A. Schnitzer, F. Bongers, R. J. Burnham, and F. E. Putz,

editors. Ecology of Lianas. John Wiley & Sons, Chichester, UK.

Schnitzer, S. A., and F. Bongers. 2011. Increasing liana abundance and biomass in tropical

forests: Emerging patterns and putative mechanisms. Ecology Letters 14:397–406.

Schnitzer, S. A., S. Estrada☐Villegas, and S. J. Wright. 2020. The response of lianas to 20 yr of

nutrient addition in a Panamanian forest. Ecology 101.

Shiklomanov, A. N., M. C. Dietze, T. Viskari, P. A. Townsend, and S. P. Serbin. 2016.

Quantifying the influences of spectral resolution on uncertainty in leaf trait estimates

through a Bayesian approach to RTM inversion. Remote Sensing of Environment 183:226–

238.

Slot, M., S. J. Wright, and K. Kitajima. 2013. Foliar respiration and its temperature sensitivity in

trees and lianas: in situ measurements in the upper canopy of a tropical forest. Tree

Physiology 33:505–515.

Stevens, G. C. 1987. Lianas as structural parasites: the Bursera simaruba example.

Tropek, R., O. Sedláček, J. Beck, P. Keil, Z. Musilová, I. Šímová, and D. Storch. 2014.

Comment on "High-resolution global maps of 21st-century forest cover change." Science

344.

Tymen, B., M. Réjoul-Méchain, J. W. Dalling, S. Fauset, T. R. Feldpausch, N. Norden, O. L.

Phillips, B. L. Turner, J. J. Viers, J. J. Chave, M. Réjou-Méchain, J. W. Dalling, S. Fauset,

T. R. Feldpausch, N. Norden, O. L. Phillips, B. L. Turner, J. J. Viers, and J. J. Chave. 2016.

Evidence for arrested succession in a liana-infested Amazonian forest. Journal of Ecology

104:149–159.

Verhoef, W., and H. Bach. 2003. Remote sensing data assimilation using coupled radiative

transfer models. Physics and Chemistry of the Earth 28:3–13.

Verhoef, W., and H. Bach. 2007. Coupled soil-leaf-canopy and atmosphere radiative transfer

modeling to simulate hyperspectral multi-angular surface reflectance and TOA radiance

data. Remote Sensing of Environment 109:166–182.

Visser, M. D. 2020. ccrtm: Coupled Chain Radiative Transfer Models. https://cran.r-

project.org/web/packages/ccrtm/index.html.

Visser, M. D., H. De Kroon, H. C. Muller-landau, S. A. Schnitzer, and E. Jongejans. 2018a. A

host – parasite model explains variation in liana infestation among co- occurring tree

species:2435–2445.

Visser, M. D., S. M. McMahon, C. Merow, P. M. Dixon, S. Record, and E. Jongejans. 2015.

Speeding Up Ecological and Evolutionary Computations in R; Essentials of High

Performance Computing for Biologists. PLoS computational biology 11:e1004140.

Visser, M. D., S. J. Wright, E. Muller-Landau, H. C. Jongejans, L. S. Comita, H. de Kroon, and

S. Schnitzer. 2018b. Tree species vary widely in their tolerance for liana infestation: a case

study of differential host response to generalist parasites. Journal of Ecology 106:781–794.

Waite, C. E., G. M. F. Heijden, R. Field, and D. S. Boyd. 2019. A view from above: Unmanned

aerial vehicles (<scp>UAV</scp> s) provide a new tool for assessing liana infestation in

tropical forest canopies. Journal of Applied Ecology 56:902–912.

Wang, Z., A. K. Skidmore, T. Wang, R. Darvishzadeh, and J. Hearne. 2015. Applicability of the

PROSPECT model for estimating protein and cellulose + lignin in fresh leaves. Remote

Sensing of Environment 168:205–218.

Werden, L. K., B. G. Waring, C. M. Smith-Martin, and J. S. Powers. 2018. Tropical dry forest

trees and lianas differ in leaf economic spectrum traits but have overlapping water-use

strategies. Tree Physiology 38:517–530.

Willard, J., X. Jia, S. Xu, M. Steinbach, and V. Kumar. 2022. Integrating Scientific Knowledge

with Machine Learning for Engineering and Environmental Systems. ACM Computing

Surveys 55.

Wright, S. J., I.-F. I. Sun, M. Pickering, C. D. C. Fletcher, and Y.-Y. Y. Chen. 2015. Long-term

changes in liana loads and tree dynamics in a Malaysian forest. Ecology 96:27482757.

Wu, J., H. Kobayashi, S. C. Stark, R. Meng, K. Guan, N. N. Tran, S. Gao, W. Yang, N.

Restrepo-Coupe, T. Miura, R. C. Oliviera, A. Rogers, D. G. Dye, B. W. Nelson, S. P.

Serbin, A. R. Huete, and S. R. Saleska. 2018. Biological processes dominate seasonality of

remotely sensed canopy greenness in an Amazon evergreen forest. New Phytologist

217:1507–1520.

Wyka, T. P., J. Oleksyn, P. Karolewski, and S. A. Schnitzer. 2013. Phenotypic correlates of the

lianescent growth form: a review. Annals of Botany 112:1667–1681.

Zhang, J., B. Rivard, A. Sánchez-Azofeifa, and K. Castro-Esau. 2006. Intra- and inter-class

spectral variability of tropical tree species at La Selva, Costa Rica: Implications for species

identification using HYDICE imagery. Remote Sensing of Environment 105:129–141.

Zhang, Q., X. Xiao, B. Braswell, E. Linder, F. Baret, and B. Moore. 2005. Estimating light

absorption by chlorophyll, leaf and canopy in a deciduous broadleaf forest using MODIS

data and a radiative transfer model. Remote Sensing of Environment 99:357–371.

Zizka, A., D. Silvestro, T. Andermann, J. Azevedo, C. Duarte Ritter, D. Edler, H. Farooq, A.

Herdean, M. Ariza, R. Scharn, S. Svantesson, N. Wengström, V. Zizka, and A. Antonelli.

2019. CoordinateCleaner: Standardized cleaning of occurrence records from biological

collection databases. Methods in Ecology and Evolution 10:744–751.

 Tables and figures

Figure 1. Datasets used in the study to compare the tropical liana and tree spectral signals. Panel

A shows the geographic distribution of plants species in the leaf spectroscopy dataset, including

the focal tropical liana (orange) and tree species (green). Note that no sampling was conducted in

Africa, but three species are common to both the Americas and Africa: *Parinari excelsa,*

Symphonia globulifera and the liana *Byttneria catalpifolia*. The locations and remote sensing

platform types of canopy scale data are given in blue and purple with reference numbers

corresponding to Table S1. Panels B and C display liana spectral signals from airborne and

spaceborne sensors at focal sites. Top rows show surface reflectance signals of lightly infested

forests (R_{tree}) and heavily-infested forests (R_{liana}) at each location. Bottom rows show average

differences between R_{lianas} and R_{trees}.

Figure 2. Illustration of the coupled radiative transfer model, merging PROSPECT and

FOURSAIL2 to study light behavior in vegetation canopies. PROSPECT addresses leaf radiative

transfer, using semi-transparent stacked plates to represent leaf absorption (α), transmittance (τ),

and reflectance (ρ). It employs the generalized plate model to determine wavelength-dependent

transmissivity ($k(\lambda)$), based on empirical absorption coefficients and leaf biochemical

concentrations (see table 1). FOURSAIL2 simulates bi-directional reflectance at the canopy

level, using differential equations for direct and hemispherical radiance. It splits the canopy into

two vertical and infinite horizontal layers with randomly distributed leaves. The model integrates

leaf properties (α, ρ, τ) from PROSPECT and allows for different particles, either layered or

mixed, using the canopy dislocation parameter (D). In our study, D was set to 1, placing liana

leaves predominantly above host leaves. Canopy layers are defined by sensor-sun geometry (θ_s,

θ_o, ψ), affecting radiance flux length and the angular projection of leaf area. This projection

relies on mean leaf angle (Ω_1, Ω_2) and leaf area index (LAI), split between top (liana; 1- f_2) and

bottom (tree; f_2) layers. The model includes a soil layer with variable reflectance and a moisture

parameter (ω), and considers the hotspot effect, where equal observer and solar zenith angles

increase reflectance. The hotspot's intensity is calculated as canopy coarseness (leaf width to

canopy height ratio, $\kappa = b/H$), and the model also accounts for crown cover (Cv) and shape (ζ).

However, we only study dense tropical forests, so Cv was set to 1, simplifying the model by

negating ζ's impact on reflectance. Illustrations by M. D. Visser.

Figure 3. Spectal properties of liana and tree leaves in the solar spectrum (400 to 2500 nm).

Panels A to J show the predicted posterior distributions of reflectance ($\hat{\rho}$), transmission ($\hat{\tau}$) and

absorption ($\hat{\alpha}$) of liana and tree leaves (A, E, H), and the difference between the posterior

distributions (B, F, I) – here solid lines depict means and the shaded envelop shows 95% credible

intervals. Kullback-Leiber Divergence (KLD) quantifies how the overall spectral probability

distribution of trees diverges from liana for reflectance (C), transmission (G), and absorption (J)

– withe zero indicating an identical distribution. The violin plots in panel K, show differences

between lianas and trees in predicted posterior distributions of leaf traits. From right to leaf;

chlorophyll content (C_{ab}), carotenoids (C_{ar}), water mass per unit area (C_w), leaf mass area (C_m),

anthocyanin content (C_{an}), brown pigments (C_{br}) and leaf structure (N). Fractions within each

violin plot incate the fraction of posterior samples that overlap with zero (i.e. no difference).

Figure 4. Model predictions for high and low levels of lianas coverage compared to their

observed reflectance from airborne and satellite hyperspectral signals. Dotted lines and colored

envelopes give mean and inter quartile range (IQR) respectively. Model fits performance was

uniformly high, as measured by the coefficient of determination ($R^2 > 0.98$). Inset plots show

model predictions of the differences in the signal between high and low liana coverage, which all

compared well with the observed difference (R^2 = 0.46, 0.73, 0.92), respectively for Bolivia,

Malaysia and Panama).

Figure 5. Leaf and canopy traits of liana-covered and liana-free forest canopies. Inverse

estimates for lianas and trees from full canopy models are compared at each site (black) with

independent leaf model data from spectrometers (blue) and, where available, independent field

and lab measurements (red). Traits are shown as percentages of average tree traits for easy

comparison, irrespective of units. Absolute values and ground truth comparisons are in Appendix

S6. The fraction of leaf area index due to lianas is noted as 1-f2. Mean values and 95% credible

intervals are represented with dots and whiskers. Parameters not significantly different between

groups across any sites are omitted. Full details on parameters and their posterior distributions

are in Appendix S5.

Figure 6. The relative importance of different traits in generating the liana signal across sites.

Here, spectral traits of the top layer were initially set as tree traits and gradually replaced with

liana traits either one by one (additive) or in pairs (interactive effects). The Kullback-Leiber

divergence score was calculated at each iteration. Panel A displays the average relative absolute

impact of changing a single trait in the top layer at each wavelength across all sites. Panel B

shows the total impact of additive (darker colors) and interactive effects (lighter colors). Positive

impact indicates a change towards the liana signal, while negative impact indicates a divergence.

Figure S7 presents the same data for individual regions (Bolivia, Malaysia, and Panama).

Figure 7. The theoretical capacity of classifiers to distinguish lianas using the expected

Kullback-Leiber divergence (KLD). Panel A predicts KLD across wavelengths for 100% liana

infestation, with model-based variance estimation, and overlays spectral sampling regions of

various remote sensors to assess their effectiveness in liana detection. Panel B presents the

cumulative KLD for each sensor, with mean divergences of 0.63 (Hyperion), 0.54 (WorldView

2), and 0.45 (Landsat 5). Panels C-E explore the impact of sensor spatial resolution on liana

discernibility at different scales: crown-scale infestation (C), forest gap arrestation (D), and liana

forests (E). They include simulated scenes and resampling results, showing the effect on KLD.

Liana cluster sizes in these simulations are 350, 2000, and 30,000 m², corresponding to

infestation scales from prior studies (Marvin et al. 2016; Chandler et al. 2021; Schnitzer et al.,

2000; Foster et al. 2006; Tymen et al. 2016). The colormap indicates liana infestation levels,

while line colors in the graphs show different features. Grey vertical lines mark different

resolutions, and a black dashed line shows the expected KLD decline for random feature

distribution. Each scenario has a roughly equal number of feature pixels.

Scale	Trait	parameter	units	Description	Validation dataset
Leaf	Leaf stuctural parameter	N	unitless	Average number of air-cell wall interfaces (where light passes between the refractive index for water to air) within the mesophyll. Expected to correlate strongly with leaf thickness.	1,2,3,11
Leaf	Chlorophyll A & B content per unit leaf area	C_{ab}	$\mu g/cm^2$	Primary photosynthetic pigment.	1,2,3
Leaf	Carotenoid content per unit leaf area	C_{ar}	$\mu g/cm^2$	Secondary photosynthetic pigment.	1,2,3
Leaf	Water mass per unit leaf area.	C_w	g/cm^2	Known as the "equivalent water thickness", the hypothetical thickness of water distributed over the area of a remotely sensed pixel, because the parameter's units reduce to cm when "divided" by the density of pure water (1 g/cm^3).	1,2,3,11
Leaf	Leaf dry matter per area	C_m	g/cm^2	Oven dried weight over fresh area.	1,2,3,11
Leaf	Anthocyanin content	C_{an}	$\mu g/cm^2$	Protective flavonoids shielding photostems from excess light and herbivory.	estimated (calibration parameter)
Leaf	brown pigments content	C_{br}	arbitrary	Colored residu of degraded pigments during senescence or after stress.	estimated (calibration parameter)
Canopy	Leaf Area Index	LAI	unitless	Projected area of all leaves over a unit of land (m^2 / m^2)	13
Canopy	Leaf inclination angle	Ω	degrees	The average angular orientation of the leaf normal relative to the zenith	17
Canopy	Fraction secondary particles	f_2	unitless	Proportion of secondary (tree) leaf particles in canopy as a fraction of the total LAI, with 1-f_2 giving the fraction of primary leaves (lianas).	16
Canopy	Canopy dissociation factor	D	unitless	Mixing of primary (liana) and secondary (tree) particles, ranging between 0 and 1, with 0 a perfectly homogeneous mixture and 1 all primary leaves in top layer.	fixed (set to 1)
Canopy	hotspot	κ	unitless	Particle coarseness, defined as the ratio of the mean (projected) particle size over the width of the canopy layer (b/H; Fig. 2).	estimated
Canopy	Soil moisture	ω	unitless	Soil layer reflectance is influenced by soil moisture (0 to 1).	estimated
Canopy	Level of infestation	L	unitless	liana infestation index (0 to 1) known from data.	fixed
Stand	Verticle crown cover fraction	Cv	unitless	Fraction ground area covered by foliage at nadir view (C_{area}/A; Fig. 2)	fixed (set to 1)
Stand	Crown shape	ζ	unitless	Tree height over crown diameter (C_d/A; Fig. 2)	fixed
Sensor	Solar zenith angle	θ_s	degrees	Compliment of sun elevation in degrees from horizon (90^o - sun elevation)	fixed
Sensor	Observer zenith angle	θ_o	degrees	Sensor zenith angle (datasets often nadir corrected)	fixed
Sensor	Sun-sensor relative azimuth angle	ψ	degrees	Relative azimuth angle, or difference between sun and oberver azimuth angles	fixed
All	Error	σ	unitless	residual error on predicted reflectance $\hat{\rho}$	estimated

Table 1: Model parameters and associated canopy and leaf traits

Figure 1: caption in maintext

Figure 2: caption in maintext

Figure 3: caption in maintext

Figure 4: caption in maintext

Figure 5: caption in maintext

Figure 6: caption in maintext

Figure 7: caption in maintext