A hymenopteran odorant alerts flies to bury eggs

Shaun M. Davis[1], Gregory T. Chism[2], Megan M. Maurer[3], Julio E. Trejo[1], Ricardo J. Garcia[1], and Todd A. Schlenke[1]

[1]Department of Entomology, University of Arizona, Tucson, AZ [2]Entomology and Insect Science Graduate Interdisciplinary Program, University of Arizona, Tucson, AZ; [3]Bioscience Mass Spectrometry Core Facility, Arizona State University, Tempe, AZ

Correspondence: shaundavis@email.arizona.edu, schlenke@email.arizona.edu

Abstract: Ants are ubiquitous and consume insects at all life stages, presumably creating a strong selective pressure for ant avoidance behaviors across insects. The insect egg stage can be especially defenseless against predation given that eggs are usually immobile and unguarded, suggesting insect mothers may have evolved oviposition strategies to minimize the ant predation risk to their offspring. Given the lack of parental care in most insects, these oviposition strategies would likely be innate rather than learned, since insect mothers are not usually present to assess predation of their eggs. Here, we use the vinegar fly *Drosophila melanogaster* as a model system for examining parental defensive responses to ant presence. Flies usually lay eggs partially inserted into the food substrate, although some are laid on top of the food and a few are inserted deeply into the food. We found that exposure to ants significantly alters fly oviposition depth: the proportion of eggs on the food surface decreased while the proportion of buried eggs increased. Buried eggs survive ant foraging bouts better than surface eggs, showing that this oviposition depth behavior is adaptive. This induced behavior is conserved across the genus Drosophila and is dependent on the fly olfactory system: anosmic mutant flies fail to bury their eggs in the presence of ants, and ant odor extracts are sufficient to induce egg burying. By fractionating ant body washes and using GC-MS to identify fraction constituents, we identified the saturated, long-chain alcohol 1-octadecanol as the odorant flies use to sense ant presence. To further delineate the ant lineages to which flies respond, we exposed flies to the odors from numerous species of ants and other insects. Surprisingly, flies buried their eggs in response to the odors of nearly all hymenopterans tested, including hymenopteran groups that flies rarely interact with in nature like bees and paper wasps. Our data suggest that 1-octadecanol is a conserved and ancient hymenopteran odorant, and that drosophilids evolved a mechanism for sensing this odorant early in their evolution as a means of protecting their offspring from ant predation. This study sheds light on the ecology and mechanisms underlying a common biotic interaction in nature, that between insect parents and the ants that would consume their offspring.

Introduction:

One of the greatest threats that animals face in nature is predation, being killed and eaten by another organism (Bowman & Hacker, 2020). A common type of defense against predation are defensive behaviors, e.g. where animals hide or escape from nearby predators. Organisms have also evolved to assess and behaviorally respond to the mere risk of predation, not just direct predator exposure (Hermann & Landis, 2017). Because predation results in death, anti-predation behaviors, especially in non-social organisms, are usually innate rather than learned, encoded in an organism's germ line and brain (Baker et al., 2001; Ren & Tao, 2020). The neurogenetic basis of how naïve prey organisms perceive and respond to predators to which they have never been exposed remains poorly understood. Therefore, it would be useful to develop a model system for studying the mechanistic basis of innate anti-predation behaviors.

The vinegar fly *Drosophila melanogaster* has been a genetic 'model organism' for more than a century, and has already proven useful for understanding interactions with a different kind of biotic threat: parasitism (Lemaitre & Hoffmann, 2007). Furthermore, recent technological innovations have made it possible to genetically manipulate the activity of relatively small groups of neurons in the fly nervous system, and thus define the neurological circuitry underlying defensive behaviors (Venken et al., 2011). These tools have been used to uncover the neurological basis of fly responses to simulated predation scenarios, such as a visual 'looming' stimulus (Ache et al., 2019; Morimoto et al., 2020; Muijres et al., 2014). Nevertheless, experiments with natural predators may yield a more nuanced and complete picture of fly defense behaviors. Because flies and their potential predators are relatively small, these natural predator-prey interactions can be studied in controlled lab settings. However, one drawback of developing flies as a model for anti-predation behaviors is that natural history data describing the relative importance of different predator types is limited for natural populations of *Drosophila* (Markow, 2015; Reaume & Sokolowski, 2006; Soto-Yéber et al., 2018).

A large diversity of generalist predators likely consume *D. melanogaster* in nature. We have observed adults being caught in the air or picked off surfaces by predatory flies, dragonflies, spiders, lizards, and hummingbirds. Flies have evolved constitutively erratic flight patterns, as well as induced behaviors like evasive flight maneuvers, avoidance, freezing, jumping, and posturing to escape capture in these contexts (Combes et al., 2012; de la Flor et al., 2017; Parigi et al., 2019). Less is known about predation of *D. melanogaster* eggs, larvae, and pupae in nature, although we have observed rove beetles, predatory beetle larvae, predatory fly larvae, and ants consuming these juvenile stages. Given that fly eggs and pupae are immobile and that larvae are relatively slow crawlers, options for behavioral defenses against predators in these life stages appear limited. However, some constitutive behaviors have presumably evolved to limit predation of juvenile flies, such as adult females preferring to oviposit in food crevices and chemically masking their eggs with pheromones, larvae preferring darker (more hidden) parts of the food and often burrowing as they are eating, and pupating larvae dispersing away from the food and gluing themselves to a substrate (Borne et al., 2021; Narasimha et al., 2019; Rockwell & Grossfield, 1978; Sawin-McCormack et al., 1995; Soto-Yéber et al., 2018; Vijendravarma et al., 2013)

Female oviposition choices are particularly interesting because they can represent trans-
generational anti-predation behaviors (a type of parental care) (Refsnider & Janzen, 2010). The
decision about where, when, and how to lay an egg is complicated and relies on multiple kinds
of sensory information (Cury et al., 2019; Richmond & Gerking, 1979; Rockwell & Grossfield,
1978; Wang et al., 2020; Zhang et al., 2020). This information includes the mating and
reproductive status of the female, aspects of the abiotic environment like the weather, the time
of day and season, aspects of the food source like water content, stiffness, color, odor, and
taste, and aspects of the biotic environment like presence of conspecifics and potentially the
presence of biotic threats. Might flies alter their oviposition choices (i.e. trigger an innate
induced behavior) in the presence of predatory threats to their offspring?

Ants are ubiquitous and occupy diverse niches in nearly all natural ecosystems
(Hölldobler & Wilson, 1990). Many are facultative or obligate predators of other organisms,
including insects of all life stages (Fernandes et al., 2012), which provide necessary proteins and
fats for colony growth (Hölldobler & Wilson, 1990). A number of studies have shown that
presence of ants, and specifically visual or olfactory ant signals, cause diverse insect species to
avoid oviposition in the ant-infested area (Freitas & Oliveira, 1996; Sendoya et al., 2009; Taylor
et al., 1998; Van Mele et al., 2009). Although published data on interactions between *D.
melanogaster* and ants are scarce (Soto-Yéber et al., 2018), ants have been shown to be
important predators of the eggs and larvae of other *Drosophila* species (Escalante & Benado,
1990; Lewis & Worthen, 1992; Worthen et al., 1993). Anecdotally, we have noticed that rotting
fruit traps meant to attract *D. melanogaster* attract far fewer flies when ants are present, and
we have seen ants carrying off what appear to be *D. melanogaster* eggs and larvae.

Here we tested whether naïve flies alter their oviposition behavior in the presence of
predatory ants. We discovered that flies push their eggs deeply into the food substrate when
exposed to ants, which protects eggs from ant predation. This innate, induced oviposition
depth behavior is a conserved trait across the genus Drosophila and is exhibited in response to
diverse ant species, as well as other hymenopterans. Unlike previous work with other fly
predators (de la Flor et al., 2017), olfaction, but not vision, is required for the oviposition depth
behavior. We identified the saturated long chain alcohol 1-octadecanol as a conserved
hymenopteran odorant that alerts flies to ant presence. This system can serve as a model for
how innate threat recognition and downstream defensive behaviors are encoded in the
germline and hardwired into the brain.

Materials and Methods:

Species used:

Flies: All flies were raised on cornmeal molasses food (10L water, 75g agar, 275g yeast, 520g
cornmeal, 110g sugar, 1046g molasses, 45mL propionic acid, and 100mL 20% (w/v) Tegosept)
and maintained at 24°C in ~60% humidity with a 16:8 light:dark cycle. We used a *D.
melanogaster* Oregon R (OreR) strain as our wild-type strain. The following *D. melanogaster*
mutants were obtained from the Bloomington Drosophila Stock Center (with stock number
given): vision mutants GMR-hid (5771) and ninaB[1] (24776), olfaction mutant orco[2] (23130), the

driver strain Or49a-GAL4 (9985), and the responder strains UAS-Kir2.1 (6595) and UAS-hid
(65403). The mutant chromosomes from these strains were crossed into the OreR background
to reduce genetic variability. White-eyed w[1118] flies were provided by Daniela Zarnescu and
used as hosts for growing parasitoid wasps. Of the other *Drosophila* species used, wild *D.*
simulans were caught in Tucson, AZ in 2018 and maintained in the lab as a single strain. *D.*
yakuba and *D. virilis* were acquired from the Drosophila Species Stock Center (stock numbers
14021-0261.01 and 15010-1051.87, respectively).

Ants: A laboratory culture of *Pheidole hyatti* was originally started in June 2018 from a single
queen collected post-nuptial flight and kept in a small 5ml cotton-stopped tube. Once the
queen's first brood emerged the housing tube was placed into 17.5cm x 12.5cm x 6cm plastic
container with inside walls coated in 'insect-a-slip' (BioQuip product #2871A) to prevent
escape. The colony was given cotton ball stopped, water-filled 5 ml plastic tubes, and were fed
ad libitum weekly with both a 2 ml microcentrifuge tubes of honey water (1/4 teaspoon per 50
mL water), and 1/8 of a fresh-frozen cockroach (approximately 0.075g) (*Nauphoeta cinerea*).
During acclimation, the colony was kept in a laboratory at 20°C with a 12:12 h light cycle and
20-25% relative humidity. Once the colony outgrew the initial nest chamber, larger 3.5 cm
diameter water-filled glass tubes, stopped with cotton, and a housing container measuring
31.5cm x 23.5cm x 10.5cm was provided. Other species of wild ants were trapped around
Tucson, AZ using small amounts of protein (tuna) and sugar (honey) arranged on pieces of
cardboard placed near active ant trails. After roughly 1 hour, ant-covered cardboards were
transferred to gallon-sized zip lock bags and brought back to the lab, where most were frozen
at -20°C for a later experiment (Fig. 5). Because the laboratory *P. hyatti* colony collapsed during
the course of these experiments, we turned to readily-available *Forelius mccooki* wild ants for
many live exposure experiments. Live *F. mccooki* workers were collected and placed in a
31.5cm x 23.5cm x 10.5cm bin (walls coated with insect-a-slip) and provided with 2 cm and 3.5
cm diameter cotton-stopped glass tubes of water. Live ants were never reused across
experiments.

Ant species were identified morphologically to genus using a dichotomous key (Fisher &
Cover, 2007), and identified to species by regional species level keys (see Supplementary Table
1 for all keys and sources). No species used in this study are protected or endangered. To
confirm morphological identifications, DNA was extracted from the ant samples for Sanger
sequencing. First, 10-50 ants were homogenized in 20-30 μL extraction buffer (0.1M Tris-HCL,
pH 9.0, 0.1M EDTA, 1% SDS) with a Kontes pellet pestle in a 1.5 mL Eppendorf tube. Additional
extraction buffer was added to bring the volume up to 500 μL, then the samples were
incubated for 30 min in a 70°C water bath. Cellular material was precipitated and pelleted by
adding 70 μL of 8M potassium acetate to each sample, incubating on ice for 30 min, and
centrifuging at 13000 rpm for 15 min at 4°C. The supernatant was transferred to a fresh
Eppendorf tube, then the DNA was precipitated and pelleted by adding 300 μL isopropanol and
centrifuging at 13000 rpm for 5 min at room temperature. After a second clean-up spin, the
resulting DNA pellet was washed with 1 mL 70% ethanol and centrifuged at 13000 rpm for 5
min at room temperature to re-pellet. The pellet was allowed to air dry for 1-2 minutes, then
resuspended in 75 μL nuclease-free water and stored at -20°C. The mitochondrial *cytochrome*
oxidase I (*COI*) gene was sequenced to identify the different ant species. Following the GoTaq

Green Master Mix (Promega, WI, USA) protocol, we generated 25μL reactions containing 12.5
μL GoTaq, 7.5 μL nuclease-free water, 2 μL of both LCO1490 and HC02198 primers (Folmer et
al., 1994), and 1 μL extracted DNA. Polymerase chain reaction (PCR) conditions were as follows:
94°C for 5 min, 37 cycles of 94°C for 1 min, 51°C for 1 min, 72°C for 1 min, followed by 72°C for
2 min and a hold at 4°C. The PCR products were visualized on a 1% agarose gel with SYBR safe
DNA gel stain (Thermo Fisher Scientific) to confirm single DNA bands were present. PCR
products were purified using the QIAquick PCR Purification Kit (Qiagen) and then sequenced at
the University of Arizona Genetics Core using the PCR primers. All sequences can be found in
Supplementary File 1.

Wasps: The *Drosophila* parasitoid wasp *Leptopilina heterotoma* (strain Lh14) and *L. boulardi*
(strain Lb17) (Schlenke et al., 2007) were used in various wasp exposure assays, and a number
of other live parasitoid species maintained in the Schlenke lab were used in a later experiment
(Fig. 5). To culture most of these wasps, adult *D. melanogaster w*^{1118} flies were allowed to lay
eggs in vials containing *Drosophila* media for 3-4 days, after which the flies were replaced with
5-10 female and 2-3 male wasps. The wasps *L. clavipes*, *Ganaspis brasiliensis*, and *Asobara*
tabida were reared the same way, except that they were grown on the host species *D. virilis*
and in bottles rather than vials. The adult wasps were aged for at least 2 days before being used
in exposure experiments. Live wasps were never reused across experiments.

Other species: A variety of other arthropod species were used in an experiment to determine
the range of organisms to which *D. melanogaster* responds (Fig. 5). Many of these species were
collected around Tucson, AZ, and identified by morphology to the most specific taxonomic
group possible. Several other species were provided by research labs at the University of
Arizona. See Supplementary Table 2 for more information about all the species used. The NCBI
taxonomy ID was used to create a phylogeny of the arthropods used using phyloT and iTOL
software (Letunic & Bork, 2021).

Exposure experiments: Flies were grown in 6oz square bottom culture bottles (Genessee
Scientific). On day 0, all adults were cleared from the bottles, the flies were allowed to eclose
for three days, then collected and kept in standard molasses food vials for roughly 24 hours
until day 4. New vials were prepared with 0.5g ± 1mg Instant Drosophila Medium (Carolina
Biological Supply Company) and hydrated with 1.6 mL water with 1% red food coloring
(McCormick) (to enhance contrast between the food and eggs). The vials were supplemented
with 3-4 drops of hydrated yeast to promote egg development in female flies. The flies were
sorted into groups of ten females (of the appropriate genotype) plus two OreR males per vial
and kept overnight on this food until day 5 to allow them to recover from CO_2 exposure before
the experiment. The experimental vials were also prepared using the same red instant food but
without the added yeast. For live insect exposure trials, on the day prior to the start of the
experiment eight live insects, either *F. mccooki* or *P. hyatti* ants, or Lh14 or Lb17 wasps, were
added to the experimental vials to allow the odorous compounds to accumulate (day 4).
Control, unexposed vials contained no insects. The following day (day 5), the 2-5 day old flies
were flipped to the experimental vials, with or without live insects or insect odors. For the
whole-body wash odor exposure trials (see below), 50 μL of the solvent (control) or odor
extract was added to each experimental vial. Once 1-octadecanol was identified as the odorant

of interest, we similarly used 50 μL of 1-octadecanol standard (MilliporeSigma, 258768
ReagentPlus® 99%, Burlington, MA, USA) in hexane at indicated concentrations in each
experimental vial. The vials were kept in a fume hood to allow the solvent to evaporate for
fifteen minutes before the flies were added to start the experiment.

For all exposure experiments, we ran five replicates per treatment. The flies were kept
in the experimental vials for 24 hours at 24°C in ~60% humidity with a 16:8 light:dark cycle,
then all insects were removed and the vials were moved to -20°C to stop egg development. The
following parameters were used to categorize each egg position: 'surface' eggs were those for
which their entire circumference, from anterior to posterior ends, was entirely visible; 'buried'
eggs were those where the position at which the dorsal appendages emerges from the eggshell
was located below the surface of the substrate; 'partial' eggs were those that were positioned
in the food substrate, but not to the extent of the buried eggs. Unlike *D. melanogaster* eggs, *D.
virilis* eggs have two pairs of dorsal appendages, so the more posterior pair was used to
determine the 'buried' position. To calculate the egg position index (EPI), we used the following
equation: EPI = $(S - B)/T$ where S is the number of eggs on the surface, B is the number of eggs
that are buried, and T is the total number of eggs across all three categories.

Odor extracts: All insects were frozen at -20°C for at least 24 hours prior to odor extraction. A
freeze-thaw cycle was shown to improve detection of ant volatile compounds (Chen, 2017). We
used 1.5 mL Eppendorf tubes, washed two times with 500 μL solvent, for the odor extraction.
Initial experiments with ants or parasitoid wasps used 70 insects washed with 350 μL solvent.
Tubes were vortexed for 1-2 minutes, centrifuged briefly, then 50 μL aliquots (representing the
odor equivalent of ten insects) were pipetted out of the tube and added to each of the five
replicate experimental vials. For control vials, the pure solvent underwent the same procedure
but without insects. For the ant and wasp fractionation experiments, we increased the odorant
concentration five-fold by reducing the solvent-to-insect ratio to 1 μL solvent per insect. For
odor extractions from especially small or large arthropod species, we used body mass instead of
number of individuals to determine the solvent-to-insect ratio. A minimum of 30 mg was
washed with 350 μL solvent. See Supplementary Table 1 for more information about how body
washes were generated.

Ant predation experiment: OreR flies laid eggs on standard molasses food in 5.5cm diameter
petri dishes for up to two hours. Eggs were gently removed and manually positioned on 5.5cm
diameter petri dish filled with 0.5% agarose gel with red food coloring. Surface eggs were set on
top of the agarose gel, partially buried eggs were inserted lengthwise roughly 50% into the gel,
and the buried eggs were inserted until the gel covered the point at which the dorsal
appendages connect to the eggshell (twenty eggs per category for a total of sixty eggs). The *P.
hyatti* laboratory colony was used under the assumption that it displays normal foraging
activity. A secondary 17.5cm x 12.5cm x 6cm plastic container, with walls coated in 'insect-a-
slip,' was connected to the main colony chamber by a 2cm (length) x 1.5cm (diameter) plastic
tube plugged with cotton. The cotton separator was then removed, and ants were allowed to
freely explore the smaller chamber for thirty minutes before the dish with fly eggs was added.
The ants were allowed to forage for thirty minutes before removal of the egg dish, at which
point the remaining eggs were counted. The connection between the colony chamber and the

foraging chamber was then blocked, the remaining ants were transferred back to the colony
chamber, and the bottom of the foraging chamber was washed with ethanol to remove any ant
pheromone trails. After one hour, the cotton separator was removed and the next replicate
trial began. Two replicates were run per day over the course of ten days, for a total of twenty
replicates. After counting the remaining fly eggs, we scored egg hatching rate 48 hours after the
ant exposure to confirm that the remaining eggs were viable.

Testing tradeoffs to deep oviposition: To test whether buried fly eggs show lower survival or
slower development to the adult stage, we had OreR flies lay eggs on standard molasses food in
5.5cm diameter petri dishes for two hours. Groups of thirty eggs were each transferred to a vial
with molasses food and manually positioned such that all eggs were either on the surface or
buried deeply in the food. Starting eleven days post egg lay, the number of flies that eclosed
per vial was counted daily. To test whether buried fly eggs are more susceptible to a biotic risk
(being further buried by foraging fly larvae), twenty white-eyed (w^{1118}) third instar larvae were
added to the vials of fly eggs, and eclosion was assayed. To test for cold temperature risk, vials
with fly eggs were placed overnight (~15 hours) at 4°C, then transferred to 24°C to allow the
flies to continue development, before assaying eclosion. To test for risk of drowning, vials with
fly eggs were sprayed with water using a 4oz fine mist sprayer bottle such that the bottom of
the vial had a thin layer of water, before assaying eclosion.

To test whether there is a tradeoff to adult female flies when ovipositing deeply into the
food, we measured the duration of the oviposition phase and the clean-and-rest phase post-
oviposition by taking videos of groups of ovipositing flies. Batches of five OreR females plus one
male were collected from bottles three days after being cleared and kept on molasses food
overnight in rectangular 3 cm (length) x 3 cm (width) x 6.3 cm (height) chambers. The following
day, five female wasps (strain Lh14) were added and kept co-housed with the flies overnight.
The next day, all insects were transferred to a new rectangular chamber with molasses food.
Two strips of parafilm were placed in parallel on top of the food, leaving a 2-3 mm wide area
available for oviposition at a depth of field amenable to video capture. Multiple 1-hour videos
were recorded per day using a Basler 2.3-megapixel acA1920-155µm camera (Graftek Imaging).
Frames were captured using Pylon Viewer (version 6.2.4.9387 64-bit) at 15 Hz, and converted
to .avi video format using MATLAB code (Chowdhury et al., 2021). Egg position was scored,
then the oviposition and clean-and-rest phase durations were determined by examination of
the video files.

GC-MS: N-hexane (Optima grade), acetonitrile (Optima grade), and dichloromethane (Optima
grade) were purchased from Fisher Scientific (Pittsburg, PA, USA). 1-octadecanol (99%), 1,3-di-
tert-butylbenzene (97%), and C7 - C30 Saturated Alkanes (certified reference material) were
purchased from Sigma-Aldrich (St. Louis, MO, USA).

Ant and wasp body washes were fractionated using a Thermo Vanquish LC system
(Waltham, MA USA) with an Agilent Technologies AdvanceBio RP-mAb C4, 4.6 × 50 mm, 3.5 µm
column (Santa Clara, CA, USA). Mobile phase A was acetonitrile and mobile phase B was 50/50
dichloromethane/hexane. The flow rate was 0.830 mL/min. 50 µL of sample was injected onto
the column and fractions were collected using an Ultimate 3000 Automated Fraction Collector
(Thermo Scientific, Waltham, MA, USA). The gradient was started with 100% phase A and was

held for 1 minute. The solvent composition was changed linearly to 100% phase B at 9 minutes
and was held for 2 minutes. The solvent was immediately returned to 100% phase A and
allowed to re-equilibrate for 5 minutes. The fractions were collected every 1 minute until a total
of 11 fractions had been collected. Fractions were dried down under a gentle stream of
nitrogen and reconstituted in hexane.

GC-MS analysis was performed using an Agilent Technologies 7890B GC with 7000D
Triple Quadrupole MS (Santa Clara, CA, USA) with MassHunter Acquisition software (v.
B.07.06.2704, Agilent Technologies, Santa Clara, CA, USA). Helium was used as the carrier gas at
a flow rate of 2 mL/min. Inlet temperature was 250 °C. 1 µL of sample was injected onto a DB-5
30m × 0.25 mm × 0.25 µm column (Agilent Technologies, Santa Clara, CA, USA). Initial oven
temperature was set to 40 °C and held for 2 minutes, then increased at 5 °C/min to 300 °C and
held for 2 minutes. The transfer line was set to 230 °C. The electron impact ionization source
was set at 70 eV and the source temperature was set to 230 °C. The quench flow was set to 2.5
mL/min. The mass range was scanned from 35-500 *m/z* at a scan time of 200 ms.

Data analysis was performed using MassHunter Qualitative Analysis Workflows software
(v. B.08.00, Agilent Technologies, Santa Clara, CA, USA). Peaks were identified using the Agile
integrator and compounds were identified using the NIST 17 database with a reverse match
score of at least 50.00 (National Institute of Standards and Technology 2017). Compounds
identified as siloxanes or other silicone containing molecules were ignored as these compounds
are from the sample vial septa or the inlet septum. Compounds that were identified in both the
ant and wasp fractions constituting a significant portion of the peak area were confirmed by
matching retention times to known standards.

Statistical analysis: All graphs and statistical tests were performed using GraphPad Prism 9 (v
9.2.0) software. All data were tested for normality using the Shapiro-Wilk test. Oviposition
rates, egg position index, and eclosion rates were tested using the unpaired, two-tailed t-test
with Welch's correction or two-way ANOVA followed by Dunnett's multiple comparison test
against the control condition. The Brown-Forsythe and Welch's ANOVA with Dunnett's multiple
comparison test was used on egg position data with multiple, independent samples. Fly egg
survival after ant predation, oviposition timing, and clean-and-rest phase timing were
compared using the nonparametric Kruskal-Wallis test with Dunn's multiple comparison test.
Throughout the paper, statistical notation are as follows: * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$,
****$p < 0.0001$, ns: not significant.

Results:

Flies bury their eggs when exposed to ants

To test whether adult female flies change their oviposition behavior in the presence of
predatory ants, we co-housed flies with either wild-caught *Forelius mccooki* or laboratory raised
Pheidole hyatti ants for 24 hours. Unlike with exposure to parasitoid wasps (Supp. Fig. 1A and C)
(Kacsoh et al., 2015; Lefèvre et al., 2012; Lynch et al., 2016), female flies did not reduce
oviposition in the presence of these ants (Fig. 1A and B). The non-significant trend towards
reduced egg numbers from the *P. hyatti* exposure is likely due to the ants actively capturing the

adult flies during the experiment, as exemplified by flies that had sections missing from their wings or legs at the end of the experimental period (Fig. 1C). Ants of the genus *Pheidole* have been described as highly aggressive in terms of prey capture (Traniello, 2010).

Although there was no reduced oviposition, we noticed while counting the fly eggs that the eggs of ant-exposed flies tended to be less visible than those of control unexposed flies. Flies oviposit their eggs either fully on the surface of the food substrate, partially inserted into the food, or completely submerged beneath the surface (Fig. 1D-E). Further examination revealed that the majority of the eggs, regardless of ant exposure condition, were laid partially submerged into the food substrate (Fig. 1F), but that the second most common category switched depending upon ant exposure. Unexposed flies lay more of their eggs on the surface while those exposed to ants push more of their eggs deeply into the food substrate (Fig 1F). To better describe this alteration in oviposition depth, we calculated the egg position index, which is the proportional difference between the surface vs buried eggs (see methods). A positive value indicates a stronger preference to lay eggs on the surface of the food while a negative value indicates a stronger preference for burying eggs under the food. Exposure to both *F. mccooki* and *P. hyatti* ants significantly altered the flies' oviposition preference towards the buried egg category (Fig. 1G-H). Interestingly, this behavioral modification also occurs when flies were exposed to parasitoid wasps (Supp. Fig. 1B and D).

Buried eggs survive ant predation

We hypothesized that eggs laid deeper into the food substrate would be protected against ant consumption. We provided the laboratory culture of *P. hyatti* ants with fly eggs manually positioned at three different depths, similar to the three natural depth categories (Fig. 1D-E, 2A). After a 30 minute ant foraging bout, the remaining fly eggs were counted. Eggs positioned on the surface of the media were the most susceptible to predation, while eggs positioned beneath the surface were strongly protected (Fig. 2B). Indeed, ants readily removed the eggs on the surface but struggled (attempted and often failed) to remove the deeply positioned eggs (Supp. videos 1 and 2). Of the eggs remaining after the ant foraging bout, hatching rates were high and similar across the depth categories (Fig. 2C), showing that the ants were not consuming or otherwise harming the buried eggs on-site. These results suggest that the fly oviposition depth behavior is an adaptation to protect fly offspring from ant predation.

Given that laying eggs deeply into the substrate protects them from predation, why don't fly mothers perform this behavior constitutively? We tested whether buried eggs suffer some fitness cost compared with eggs laid on the substrate surface. First, we tested whether there was any reduction in offspring developmental time or survival for flies that were buried in the egg stage, but did not find any difference in eclosion time or eclosion success between flies manually positioned on the food surface versus buried at the egg stage (Supp. Fig. 2A). We also tested whether the presence of older fly larvae churning the food, reduced temperature, or simulated rain (water misting) harmed the buried eggs, but once again found no difference in survival to eclosion (Supp. Fig. 2B-D). These results suggest that any cost of deep oviposition is not incurred by the offspring, but instead may be borne by fly mothers. The *D. melanogaster* oviposition program has been described as a progression of stereotyped phases starting with a searching phase, then the oviposition phase, and finally the clean-and-rest phase (C.-H. Yang et

al., 2008). We hypothesized that female flies might require more time to lay buried eggs or to
rest after laying buried eggs. However, video analysis of the fly oviposition program during
parasitoid wasp exposure showed no difference in the average length of either of these phases
when the flies laid a surface egg versus laid a buried egg (Supp. Fig. 2E-F). Curiously, female flies
spent significantly longer periods laying partially buried eggs as compared to surface eggs or
fully buried eggs (Supp. Fig. 2E), suggesting that partially buried eggs need more care to
position. In summary, we were unable to detect any negative consequences of the induced egg
burying behavior. Either there is no fitness cost associated with this behavior, or there is a cost
that we have not yet identified.

Egg burying is a conserved trait across *Drosophila*

To test whether ant-induced egg burying is a conserved defensive behavior in flies, we
assayed the behavior in three other *Drosophila* species of varying evolutionary distances from
D. melanogaster: D. simulans (~3MY since the most recent common ancestor), *D. yakuba*
(~5MY), and *D. virilis* (~30MY) (Powell, 1997). Like *D. melanogaster*, the other *Drosophila*
species maintained their egg production level when exposed to *F. mccooki* ants (Fig. 3A), and
also like *D. melanogaster*, all three *Drosophila* species consistently laid their eggs deeper in the
substrate in the presence of ants (Fig. 3B). This result holds even though *D. virilis*, and to some
extent *D. yakuba*, tend to lay more deeply buried eggs than *D. melanogaster* in the unexposed
state (Fig. 3B). These data suggest that although normal egg depth behavior varies across
species, the switch to laying more buried eggs to prevent egg predation is a conserved trait.

Olfaction mediates ant detection

Fly mothers sense and respond to the presence of ants by burying their eggs, but how
do they know that ants are present? Flies have been shown to detect parasitoid wasps through
the combined effects of the visual and olfactory systems (Ebrahim et al., 2015; Kacsoh et al.,
2015, 2013; Lynch et al., 2016), and given that flies also bury their eggs in the presence of
wasps (Supp. Fig. 1) we tested both of these sensory modalities. *GMR-hid* flies express a pro-
apoptotic factor in the eye, making their eyes significantly reduced in size (Grether et al., 1995).
These sight-deficient flies maintained their ability to respond to the presence of *F. mccooki* ants
by burying their eggs (Fig. 4A). *GMR-hid* as well as *ninaB[1]* mutant flies (which are blind due to
loss of photoreceptors) also maintained their ability to sense and respond to *L. heterotoma*
parasitoid wasps (Supp. Fig. 3A-B). These data suggest that vision is dispensable for the altered
egg depth behavior. We next tested the necessity of the olfactory system: *orco[2]* mutant flies
lack the odorant receptor co-receptor and therefore lack most olfactory ability (Larsson et al.,
2004). Heterozygous *orco[2]* mutant flies showed a strong induced egg burying response in the
presence of *F. mccooki* ants, but the anosmic homozygous *orco2* mutant flies showed a much
weaker (albeit still significant) response (Fig. 4B). Exposing the anosmic flies to either *P. hyatti*
ants or *L. heterotoma* parasitoid wasps resulted in a complete failure to alter egg depth (Supp.
Fig. 3C-D). To further test the necessity of olfaction for the egg burying behavior, we manually
ablated the two main olfactory sensing organs from experimental flies either individually or in
tandem. While ablating the maxillary palps had no effect on fly ability to bury eggs in the
presence of *F. mccooki* ants, ablating the fly antennae abolished their ability to respond to the
ants (Supp. Fig. 3E). Altogether, these data demonstrate that flies require olfactory input

through their antennae to sense ants in their environment and alter oviposition depth behavior
accordingly.

It was previously shown that the fly odorant receptors Or49a and Or85f, which are
expressed in the same olfactory receptor neurons in adult flies, detect specific odorants from
parasitoid wasps in the genus *Leptopilina*, in turn driving avoidance behaviors during fly
oviposition (Ebrahim et al., 2015). One of these compounds, iridomyrmecin, was first isolated
from the Argentine Ant, *Linepithema humile* (formerly *Iridomyrmex humilis*) (Pavan, 1948)
suggesting flies may also detect ants via this mechanism. Using an *Or49a-Gal4* driver, we
inhibited activity of Or49a/Or85a-expressing olfactory receptor neurons by driving expression
of *Kir2.1* (which causes neuronal membrane hyperpolarization) in these neurons, or we
completely ablated these neurons by expressing in them the proapoptotic gene *hid*. Neither
modification blocked the fly egg burying behavior in the presence of *L. heterotoma* parasitoid
wasps (Supp. Fig. 3F-G). These data suggest that flies are using a different type of olfactory
sensory neuron and olfactory receptor to detect a novel odorant associated with ants.

If an ant olfactory stimulus is sufficient to induce the fly oviposition depth switch, we
hypothesized that ant body-wash extracts could replace live ants as the egg burying stimulus.
We used the solvent hexane to extract odorants from the bodies of *P. hyatti* ants and added the
odors directly onto the food substrate. While flies that were exposed to live *P. hyatti* ants as a
positive control showed a strong induced egg burying behavior, flies exposed to ant odors alone
showed a weaker, though still significant, response (Fig. 4C). There are at least three reasons
why odor extracts do not fully recapitulate the direct ant exposure condition. First, it is possible
that the body wash extracts only possess a subset of multiple distinct odorants required for the
full behavioral response. To test this, we used different solvents that varied in their polarity to
target different molecules. Interestingly, extracts collected in non-polar hexane (Fig 4C, Supp.
Fig. 4A) and weakly polar dichloromethane (Supp. Fig. 4B) were both sufficient to induce the
partial egg burying behavior, while use of more strongly polar solvents failed to induce any fly
response (Supp. Fig. 4C-E). However, combining body wash odorants extracted using hexane
and dichloromethane did not induce the full egg burying behavior (data not shown), showing
that no additive effect is achieved by combining the odorant subsets extracted by each solvent.
Second, it is possible that exposure to insect body wash extracts fails to fully recapitulate the
live insect exposure results because the body wash odorants dissipate over the course of the
experiment without being replaced. To test this idea, we exposed flies to both dead *L.*
heterotoma parasitoid wasps, and to vials in which the wasps had been housed, both
treatments of which should have the full wasp odorant profile but no new odorants being
produced. In both cases, flies showed a partial egg burying response similar to their response to
insect body wash extracts, supporting the odorant dissipation hypothesis (data not shown).
Third, it is possible that flies are using some other sensory modality besides olfaction or vision
to detect ant/wasp presence, such as gustation, a hypothesis that remains untested.
Regardless, we were able to extract an odorant from insect bodies that flies detected, causing
them to induce at least a partial egg burying response.

Flies sense the odorant 1-octadecanol

To identify the odorant (or odorants) in insect body washes that induces flies to bury
their eggs, we used regular-phase column chromatography to generate fractions from the
whole-body washes. These fractions are expected to pull out different subsets of the molecules
from the whole-body washes, allowing us to narrow down and identify the causal odorant(s)
using gas-chromatography mass-spectrometry (GC-MS). Eleven fractions were generated from
F. mccooki ant whole-body washes and assayed for phenotypic effects on flies. As expected,
none of the fractions altered fly oviposition rates (Fig. 5A), but fraction 9 did induce the flies to
lay eggs significantly deeper in the food than control hexane treatments (Fig. 5B). Fraction 9
from a *L. heterotoma* parasitoid wasp body wash showed a similar result (data not shown). GC-
MS of the *F. mccooki* whole-body wash compared to the fraction 9 sample revealed that the
fraction contained a small subset of the molecules found in the whole-body wash, and in
particular contained one shared high intensity peak (Fig. 5C-D). This peak was identified as the
saturated, linear eighteen-carbon alcohol, 1-octadecanol (Fig. 5C-D, peak 2). While the *F.
mccooki* whole-body wash contained iridomyrmecin (Fig. 5C, peak 1), an odorant found in
parasitoid wasps and previously shown to affect fly oviposition (Ebrahim et al., 2015), fraction 9
did not possess this odorant, consistent with the idea that the fly egg burying behavior is
triggered by a novel odorant. To confirm the role of 1-octadecanol in the egg burying behavior,
we purchased pure 1-octadecanol and exposed flies to various concentrations. At
concentrations as low as 10^{-8} M (similar to 1-octadecanol concentration in the whole-body
wash sample as measured by a standard), flies responded to 1-octadecanol as strongly as they
responded to whole ant body washes (Fig. 5E). Thus, we concluded that 1-octadecanol is the
ant odorant (shared with *L. heterotoma* parasitoid wasps) that alerts flies to the presence of
dangerous predators, inducing flies to bury their eggs in order to protect their offspring.

Flies respond to most hymenopterans

We have shown that flies lay eggs more deeply into the food substrate when exposed to
ants (Fig. 1), and that this behavior protects the eggs from ant predation (Fig. 2). Flies also
induce this behavior when exposed to a parasitoid wasp, although it is unclear what benefit
there is to burying eggs in this context given that *L. heterotoma* does not attack fly offspring
until they reach the larval stage, when they are crawling on top of the food. To determine the
breadth of ants and potentially other insects that induce flies to bury their eggs, we exposed
flies to body washes or live samples of numerous arthropods. Flies responded to the vast
majority of ant species tested representing the three ant subfamilies Dolichoderinae,
Formicinae, and Myrmicinae (Fig. 6). Exceptions included the Myrmicinae species *Monomorium
cyaneum*, as well as some trials from several other ant species (Fig. 6). Parasitoid wasps
represent another large branch of hymenopterans that impose strong selective pressures on fly
offspring survival. Flies that were exposed directly to parasitoid wasps that infect fly larvae
(genus *Leptopilina*, *Ganaspis*, and *Asobara*) or fly pupae (*Pachycrepoideus* and *Trichopria*)
significantly altered their oviposition depth (Fig. 6). However, flies also responded to the body
wash extract of the whitefly parasitoid, *Encarsia inaron*, as well as extracts from every other
hymenopteran tested (bees and paper wasps), even though *D. melanogaster* has no known
ecological interactions with these insects (Fig. 6).

The odor extracts from non-hymenopteran insects generally did not induce flies to alter
their oviposition behavior, but there were two exceptions. First, odor extract from the leaf-

footed bug, *Leptoglossus zonatus*, did induce the fly egg burying behavior in two separate trials,
despite body washes from other hemipterans showing no effect on the flies (Fig. 6). Second,
odor extract from the nymph stage of the Madagascar hissing cockroach, *Gromphadorhina*
portentosa, also affected fly oviposition behavior. In sum, these data suggest that a common
odorant, 1-octadecanol, evolved early in the hymenopterans and has been maintained over
time, with sporadic gains and losses in the broader insect group. Further, our data suggest that
flies evolved to sense this odorant as a means of protecting their offspring from predatory ants.

Discussion:

Recognition of environmental threats, and execution of an appropriate response, is
critical to organismal survival. Here, we have shown that flies detect ant presence through
olfactory sensing of the hymenopteran odorant 1-octadecanol, and switch their oviposition
behavior to more deeply position their eggs in the food substrate. This protects fly eggs from
ant predation.

Egg laying behavior follows a similar stereotyped pattern across different fly species. It is
first characterized by a search-like, informational gathering processes about the local nutrient
availability and substrate stiffness through labellum and leg contact (Bräcker et al., 2019; C.-H.
Yang et al., 2008; Zhang et al., 2020). Once a suitable site is found, flies switch to a more refined
search behavior, including ovipositor contact with the substrate (Bräcker et al., 2019), before an
egg is finally laid. These discrete micro-behaviors are conserved across *Drosophila*, with minor
variations depending on the ecology of each species. For example, the agricultural pest *D.*
suzukii spends significantly more time than *D. melanogaster* contacting the food substrate with
its ovipositor and expelling its eggs (Bräcker et al., 2019), presumably because it tends to lay
eggs more deeply into less ripened fruits (Karageorgi et al., 2017). Furthermore, *Drosophila*
species that oviposit on mushrooms tend to deeply bury their eggs (Rouquette & Davis, 2003),
perhaps because mushroom flesh is soft, or because ants commonly visit mushrooms (Lewis &
Worthen, 1992). While fly oviposition behaviors are well-studied (Bräcker et al., 2019; Cury et
al., 2019; Karageorgi et al., 2017; Van Mele et al., 2009; Wang et al., 2020; C.-H. Yang et al.,
2008; Zhang et al., 2020), this is the first report describing an induced change in oviposition
depth.

Our measure of oviposition depth, the egg position index, is a blunt tool. While egg
depth is a continuous trait, we bin the eggs into one of three depth categories, and most of the
eggs are classified as 'partially buried'. A lot of the behavior may be missed when an egg that is
only 20% inserted into the substrate is scored the same as an egg that is 80% inserted,
especially if these eggs have different levels of protection against foraging ants. Ideally a more
sensitive measure of egg depth could be devised. Nevertheless, the fact that we see significant
differences in the egg position index across treatments indicates that egg burying is a robust
and important fly behavior.

Several examples exist of organisms altering oviposition behavior in response to biotic
threats. For example, water striders (*Aquarius paludum insularis*) oviposit their eggs deeper in
the water column after exposure to parasitoid wasps, which limits egg parasitism (Hirayama &

Kasuya, 2009). Newts (*Taricha granulosa*) lay their eggs attached to plants higher in the water
column to avoid egg predator caddisfly larvae (*Limnephilus flavastellus*) (Gall et al., 2012). There
are even examples in fruit flies: *Drosophila* species avoid ovipositing at sites infested by toxic
microbes, which they detect via the odorant geosmin (Stensmyr et al., 2012). They also alter
oviposition in the presence of parasitoid wasps (Carton et al., 1986) using both visual and
olfactory cues: female flies lay fewer eggs during forced exposures, they preferentially choose
non-infested sites when given a choice by sensing the wasp odorant iridomyrmecin, and they
preferentially lay their eggs in more toxic (alcoholic) environments when exposed (Kacsoh et al.,
2013; Lefèvre et al., 2012; Lynch et al., 2016; Ebrahim et al., 2015). All of these are examples of
threat-induced behavioral changes that organisms make about 'when' or 'where' to oviposit,
whereas the new behavior described here is about 'how' flies choose to oviposit. Furthermore,
we know that the fly responses to geosmin and iridomyrmecin, and likely 1-octadecanol as well,
are hardwired into the fly, using specific olfactory receptors and neuronal circuits in the brain. It
will be interesting to determine how these different environmental cues are signaled through
the brain to regulate different types of egg laying behaviors, and whether there is any overlap
in each circuit.

Ants are a ubiquitous presence across diverse habitats, and many insects have evolved
ant avoidance behaviors triggered by distinct sensory modalities. For example, honeybees avoid
flowers in the field that have live ants or their odors nearby (Li et al., 2014; Sidhu & Rankin,
2016), and tephritid fruit flies avoid fruits perfumed with ant pheromones (Van Mele et al.,
2009). Furthermore, butterflies (*Eunica bechina*) were reported to avoid oviposition sites due to
visual detection of ant presence, although the fact that ants were pinned to plant leaves
suggests that ant odors may also have contributed to the butterfly avoidance (Sendoya et al.,
2009). Surprisingly, termites (*Coptotermes acinaciformis*) were shown to detect and avoid ants
based on the vibrational pattern of ants walking through their woody substrates (Oberst et al.,
2017). This may reflect a keen sense of vibration detection in termites, which have been shown
to communicate alarm signals via similar vibrational cues (Inta et al., 2009). These examples
show that detection of ant threats is common in the insect world, and that insects have evolved
diverse sensory mechanisms for identifying ant presence. Given that *D. melanogaster* invests
more resources into its olfactory system than other sensory systems (Keesey et al., 2019), it
may not be surprising that they detect ant predators through olfactory signals. In the future, it
will be interesting to identify the odorant receptor(s) responsible for fly sensing of ant odors
and the neural circuitry that responds to odor sensing by altering oviposition choices (Wang et
al., 2020).

Although olfaction is necessary for fly detection of ants, we could not prove that ant
odors are sufficient to induce the full fly egg burying phenotype (Fig. 4). We believe there are
two likely reasons why providing flies with ant body wash odorant extracts does not fully
recapitulate the direct insect-exposure results. First, the ant odor extract experiments likely
started with a high concentration of ant odors (like 1-octadecanol), but over the course of the
24 hour experiment these volatile odorant concentrations may have declined to a point where
fly behavior was no longer altered. In contrast, the live-ant exposures may have maintained a
high concentration of odorants throughout the experiments due to constant release from the
insects. It would be interesting to conduct an experiment in which ant odors are constantly

passed over the fly habitat to test whether exposure to a constant concentration of ant odors
induces a more robust egg burying phenotype. Second, it is possible that fly oviposition
behaviors like egg burying require multimodal sensory integration such that ant odors suppress
the preference for 'surface' eggs, but a second stimulus enhances the magnitude of preference
for 'buried' eggs. This second stimulus might include gustatory or auditory cues from the ants.
This kind of gating mechanism has been observed for fly larval rolling behavior during parasitoid
attack, where vibrational cues enhance the fly rolling response to mechanical poking by the the
wasp ovipositor (Ohyama et al., 2015). Further testing of fly sensory systems will help identify
additional inputs into the oviposition depth behavioral response.

There are sporadic reports in the literature about insects producing or behaviorally
responding to 1-octadecanol. For example, 1-octadecanol is a component in the pheromone
blend secreted from the male hairpencil glands of various male lepidopterans to attract females
(Bober & Rafaeli, 2010; Teal & Tumlinson, 1989). However, 1-octadecanol has been reported to
have repulsive effects on other insects: aphids (*Rhopalosiphum padi*) avoid leaves coated with
Trichoderma fungi, which produce 1-octadecanol (Ganassi et al., 2016), and the beet
armyworm, *Spodoptera exigua,* avoids ovipositing on roses infected with the powdery mildew
Podosphaera pannosa (Fa-zhong Yang et al., 2013), due to increased plant production of
volatiles like 1-octadecanol (Fazhong Yang et al., 2019). In our system, flies do not seem
deterred from ovipositing on food substrates with ant odors or 1-octadecanol, they just lay
their eggs deeper into the substrate. Still, allowing flies a choice between oviposition sites with
and without 1-octadecanol would offer interesting insight into potential tradeoffs – do flies
prefer to spend time searching for non-contaminated oviposition sites or potentially spend
more energy laying eggs on a contaminated oviposition site?

We showed that the ability of flies to sense 1-octadecanol from ants, and induce the egg
burying behavior, is beneficial to the flies due to the increased survival of their eggs during ant
predation events (Fig. 2). However, flies also induce the egg burying behavior in response to the
odors of other hymenopterans like wasps and bees, even though wasps and bees are not
known to harm fly eggs. This suggests that 1-octadecanol is a conserved odorant dating back to
the common ancestor of all hymenopterans. Likewise, the ant odor-induced egg burying
behavior is conserved across the genus Drosophila (Fig. 3). Given that the expansion of ant,
wasp, and bee lineages seems to coincide with the expansion of dipterans in the Jurassic period
over 150 million years ago (Misof et al., 2014; Peters et al., 2017), it is possible that fly
recognition of hymenopteran threats via 1-octadecanol sensing is ancient. In the future, it will
be interesting to determine if other dipterans mount behavioral defenses in response to 1-
octadecanol, and if the neurogenetic basis of this sensing is conserved across dipterans.

Acknowledgments:

We would like to thank Goggy Davidowitz, Anna Dornhaus, Martha Hunter, Michael Riehle,
Patricia Stock, Bruce Tabashnik, and Kathleen Walker from the University of Arizona, Vanessa
Corby-Harris from the USDA, and Elizabeth Tibbetts from the University of Michigan for
providing insects for odor extraction. This research was funded by the University of Arizona
BIO5 Institute Postdoctoral Fellowship to SMD and NSF grant IOS-1257469 to TAS. The authors

acknowledge resources and support from the Knowledge Enterprise Biosciences Mass
Spectrometry Core Facility at Arizona State University.

References:

Ache, J. M., Polsky, J., Alghailani, S., Parekh, R., Breads, P., Peek, M. Y., Bock, D. D., Von Reyn, C.
R., & Card, G. M. (2019). Neural Basis for Looming Size and Velocity Encoding in the
Drosophila Giant Fiber Escape Pathway. *Current Biology*, *29*, 1073–1081.
https://doi.org/10.1016/j.cub.2019.01.079
Baker, B. S., Taylor, B. J., & Hall, J. C. (2001). Are Complex Behaviors Specified by Dedicated
Regulatory Genes? Reasoning from Drosophila. *Cell*, *105*(1), 13–24.
https://doi.org/10.1016/S0092-8674(01)00293-8
Bober, R., & Rafaeli, A. (2010). Gene-silencing reveals the functional significance of pheromone
biosynthesis activating neuropeptide receptor (PBAN-R) in a male moth. *Proceedings of the
National Academy of Sciences*, *107*(39), 16858–16862.
https://doi.org/10.1073/PNAS.1008812107
Borne, F., Prigent, S. R., Molet, M., & Courtier-Orgogozo, V. (2021). Drosophila glue protects
from predation. *Proc. R. Soc. B*, *288:* https://doi.org/10.1098/rspb.2021.0088
Bowman, W. D., & Hacker, S. D. (2020). Predation. In *Ecology* (5th ed.). Oxford University Press.
Bräcker, L. B., Schmid, C. A., Bolini, V. A., Holz, C. A., Prud'homme, B., Sirota, A., & Gompel, N.
(2019). Quantitative and discrete evolutionary changes in the egg-laying behavior of single
Drosophila females. *Frontiers in Behavioral Neuroscience*, *13*.
https://doi.org/10.3389/fnbeh.2019.00118
Carton, Y., Bouletreau, M., Alphen, J. J. M. van, & Lenteren, J. C. van. (1986). The Drosophila
Parasitic Wasps. In M. Ashburner, E. Novitski, H. L. Carson, & J. N. Thompson (Eds.), *The
Genetics and Biology of Drosophila* (pp. 347–394). Academic Press.
Chen, J. (2017). Freeze–Thaw Sample Preparation Method Improves Detection of Volatile
Compounds in Insects Using Headspace Solid-Phase Microextraction. *Analytical Chemistry*,
89(16), 8366–8371. https://doi.org/10.1021/ACS.ANALCHEM.7B01622
Chowdhury, B., Wang, M., Gnerer, J. P., & Dierick, H. A. (2021). The Divider Assay is a high-
throughput pipeline for aggression analysis in Drosophila. *Communications Biology*, *4*(1).
https://doi.org/10.1038/S42003-020-01617-6
Combes, S. A., Rundle, D. E., Iwasaki, J. M., & Crall, J. D. (2012). Linking biomechanics and
ecology through predator–prey interactions: flight performance of dragonflies and their
prey. *Journal of Experimental Biology*, *215*(6), 903–913.
https://doi.org/10.1242/JEB.059394
Cury, K. M., Prud'homme, B., & Gompel, N. (2019). A short guide to insect oviposition: when,
where and how to lay an egg. *Journal of Neurogenetics*, *33*(2), 75–89.
https://doi.org/10.1080/01677063.2019.1586898
de la Flor, M., Chen, L., Manson-Bishop, C., Chu, T. C., Zamora, K., Robbins, D., Gunaratne, G., &
Roman, G. (2017). Drosophila increase exploration after visually detecting predators. *PLoS
ONE*, *12*(7), 1–17. https://doi.org/10.1371/journal.pone.0180749
Ebrahim, S. A., Dweck, H. K., Stokl, J., Hofferberth, J. E., Trona, F., Weniger, K., Rybak, J., Seki, Y.,
Stensmyr, M. C., Sachse, S., Hansson, B. S., & Knaden, M. (2015). Drosophila Avoids
Parasitoids by Sensing Their Semiochemicals via a Dedicated Olfactory Circuit. *PLoS Biol*,

13(12), e1002318. https://doi.org/10.1371/journal.pbio.1002318

Escalante, A., & Benado, M. (1990). *Predation on the Cactophilic Fly, Drosophila starmeri, in the*

Columnar Cactus, Pilosocereus lanuginosus. *22*(1), 48–50.

Fernandes, W. D., Sant'Ana, M. V., Raizer, J., & Lange, D. (2012). Predation of fruit fly larvae

Anastrepha (Diptera: Tephritidae) by ants in grove. *Psyche*, *2012*, 1–7.

https://doi.org/10.1155/2012/108389

Fisher, B. L., & Cover, S. P. (2007). *Ants of North America : a Guide to the Genera.* University of

California Press.

Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification

of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates.

Molecular Marine Biology and Biotechnology, *3*(5), 294–299.

Freitas, A. V. L., & Oliveira, P. S. (1996). Ants as Selective Agents on Herbivore Biology: Effects

on the Behaviour of a Non-Myrmecophilous Butterfly. *Journal of Animal Ecology*, *65*(2),

205–210. https://doi.org/10.2307/5723

Gall, B. G., Brodie III, E. D., & Brodie Jr, E. D. (2012). Fine-scale selection by ovipositing females

increases egg survival. *Ecology and Evolution*, *2*(11), 2763–2774.

https://doi.org/10.1002/ece3.389

Ganassi, S., Grazioso, P., De Cristofaro, A., Fiorentini, F., Sabatini, M. A., Evidente, A., &

Altomare, C. (2016). Long Chain Alcohols Produced by Trichoderma citrinoviride Have

Phagodeterrent Activity against the Bird Cherry-Oat Aphid Rhopalosiphum padi. *Frontiers*

in Microbiology, *7*, 297. https://doi.org/10.3389/FMICB.2016.00297

Grether, M. E., Abrams, J. M., Agapite, J., White, K., & Steller, H. (1995). The head involution

defective gene of Drosophila melanogaster functions in programmed cell death. *Genes and*

Development, *9*(14), 1694–1708. https://doi.org/10.1101/gad.9.14.1694

Hermann, S. L., & Landis, D. A. (2017). Scaling up our understanding of non-consumptive effects

in insect systems. *Current Opinion in Insect Science*, *20*, 54–60.

https://doi.org/10.1016/j.cois.2017.03.010

Hirayama, H., & Kasuya, E. (2009). Oviposition depth in response to egg parasitism in the water

strider: high-risk experience promotes deeper oviposition. *Animal Behaviour*, *78*(4), 935–

941. https://doi.org/10.1016/j.anbehav.2009.07.019

Hölldobler, B., & Wilson, E. O. (1990). *The ants*. Belknap Press of Harvard University Press.

Inta, R., Evans, T. A., & Lai, J. C. S. (2009). Effect of vibratory soldier alarm signals on the

foraging behavior of subterranean termites (Isoptera: Rhinotermitidae). *Journal of*

Economic Entomology, *102*(1), 121–126. https://doi.org/10.1603/029.102.0117

Kacsoh, B. Z., Bozler, J., Ramaswami, M., & Bosco, G. (2015). Social communication of predator-

induced changes in Drosophila behavior and germline physiology. *ELife*, *4*, 1–36.

https://doi.org/10.7554/eLife.07423

Kacsoh, B. Z., Lynch, Z. R., Mortimer, N. T., & Schlenke, T. A. (2013). Fruit flies medicate

offspring after seeing parasites. *Science*, *339*(6122), 947–950.

https://doi.org/10.1126/science.1229625

Karageorgi, M., Bräcker, L. B., Lebreton, S., Minervino, C., Cavey, M., Siju, K. P., Grunwald

Kadow, I. C., Gompel, N., & Prud'homme, B. (2017). Evolution of Multiple Sensory Systems

Drives Novel Egg-Laying Behavior in the Fruit Pest Drosophila suzukii. *Current Biology*,

27(6), 847–853. https://doi.org/10.1016/J.CUB.2017.01.055

Keesey, I. W., Grabe, V., Gruber, L., Koerte, S., Obiero, G. F., Bolton, G., Khallaf, M. A., Kunert, G., Lavista-Llanos, S., Valenzano, D. R., Rybak, J., Barrett, B. A., Knaden, M., & Hansson, B. S. (2019). Inverse resource allocation between vision and olfaction across the genus Drosophila. *Nature Communications 2019 10:1*, *10*(1), 1–16. https://doi.org/10.1038/s41467-019-09087-z

Larsson, M. C., Domingos, A. I., Jones, W. D., Chiappe, M. E., Amrein, H., & Vosshall, L. B. (2004). Or83b encodes a broadly expressed odorant receptor essential for Drosophila olfaction. *Neuron*, *43*(5), 703–714. https://doi.org/10.1016/j.neuron.2004.08.019

Lefèvre, T., de Roode, J. C., Kacsoh, B. Z., & Schlenke, T. A. (2012). Defence strategies against a parasitoid wasp in Drosophila: Fight or flight? *Biology Letters*, *8*(2), 230–233. https://doi.org/10.1098/rsbl.2011.0725

Lemaitre, B., & Hoffmann, J. (2007). The Host Defense of Drosophila melanogaster. *Annual Review of Immunology*, *25*, 697–743. https://doi.org/10.1146/ANNUREV.IMMUNOL.25.022106.141615

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Research*, *49*(W1), W293–W296. https://doi.org/10.1093/NAR/GKAB301

Lewis, G. P., & Worthen, W. B. (1992). Effects of Ant Predation and Mushroom Desiccation on the Survival of Mycophagous Drosophila tripunctata Larvae. *Oikos*, *64*(3), 553–559. Retrieved from https://www.jstor.org/stable/3545175

Li, J., Wang, Z., Tan, K., Qu, Y., & Nieh, J. C. (2014). Giant Asian honeybees use olfactory eavesdropping to detect and avoid ant predators. *Animal Behaviour*, *97*, 69–76. https://doi.org/10.1016/j.anbehav.2014.08.015

Lynch, Z. R., Schlenke, T. A., & de Roode, J. C. (2016). Evolution of behavioural and cellular defences against parasitoid wasps in the Drosophila melanogaster subgroup. *Journal of Evolutionary Biology*, *29*(5), 1016–1029. https://doi.org/10.1111/jeb.12842

Markow, T. A. (2015). The secret lives of Drosophila flies. *ELife*, *4*(JUNE). https://doi.org/10.7554/ELIFE.06793

Misof, B., Liu, S., Meusemann, K., Peters, R. S., Donath, A., Mayer, C., Frandsen, P. B., Ware, J., Flouri, T., Beutel, R. G., Niehuis, O., Petersen, M., Izquierdo-Carrasco, F., Wappler, T., Rust, J., Aberer, A. J., Aspöck, U., … Friedrich, F. (2014). Phylogenomics resolves the timing and pattern of insect evolution. *Source: Science*, *346*(6210), 763–767. https://doi.org/10.2307/24917799

Morimoto, M. M., Nern, A., Zhao, A., Rogers, E. M., Wong, A. M., Isaacson, M. D., Bock, D. D., Rubin, G. M., & Reiser, M. B. (2020). Spatial readout of visual looming in the central brain of drosophila. *ELife*, *9*, 1–102. https://doi.org/10.7554/ELIFE.57685

Muijres, F. T., Elzinga, M. J., Melis, J. M., & Dickinson, M. H. (2014). Flies evade looming targets by executing rapid visually directed banked turns. *Science*, *344*(6180), 172–177. https://doi.org/10.1126/SCIENCE.1248955

Narasimha, S., Nagornov, K. O., Menin, L., Mucciolo, A., Rohwedder, A., Humbel, B. M., Stevens, M., Thum, A. S., Tsybin, Y. O., & Vijendravarma, R. K. (2019). Drosophila melanogaster cloak their eggs with pheromones, which prevents cannibalism. *PLOS Biology*, *17*(1), e2006012. https://doi.org/10.1371/JOURNAL.PBIO.2006012

Oberst, S., Bann, G., Lai, J. C. S., & Evans, T. A. (2017). Cryptic termites avoid predatory ants by

756   eavesdropping on vibrational cues from their footsteps. *Ecology Letters*, *20*(2), 212–221.
757    https://doi.org/10.1111/ELE.12727
758  Ohyama, T., Schneider-Mizell, C. M., Fetter, R. D., Aleman, J. V., Franconville, R., Rivera-Alba,
759    M., Mensh, B. D., Branson, K. M., Simpson, J. H., Truman, J. W., Cardona, A., & Zlatic, M.
760    (2015). A multilevel multimodal circuit enhances action selection in Drosophila. *Nature*,
761    *520*(7549), 633–639. https://doi.org/10.1038/nature14297
762  Parigi, A., Porter, C., Cermak, M., Pitchers, W. R., & Dworkin, I. (2019). The behavioral repertoire
763    of Drosophila melanogaster in the presence of two predator species that differ in hunting
764    mode. *PLoS ONE*, *14*(5), 1–21. https://doi.org/10.1371/journal.pone.0216860
765  Pavan, M. (1948). Iridomyrmecin, an antibiotic substance extracted from the Argentine ant
766    (Iridomyrmex pruinosus humilis Mayz.) VIII Internat. *Cong. Entomol., Stock-Holm*, 863–865.
767  Peters, R. S., Krogmann, L., Mayer, C., Donath, A., Gunkel, S., Meusemann, K., Kozlov, A.,
768    Podsiadlowski, L., Petersen, M., Lanfear, R., Diez, P. A., Heraty, J., Kjer, K. M., Klopfstein, S.,
769    Meier, R., Polidori, C., Schmitt, T., … Niehuis, O. (2017). Evolutionary History of the
770    Hymenoptera. *Current Biology*, *27*(7), 1013–1018.
771    https://doi.org/10.1016/J.CUB.2017.01.027
772  Powell, J. R. (1997). *Progress and prospects in evolutionary biology : the Drosophila model*.
773    Oxford University Press.
774  Reaume, C. J., & Sokolowski, M. B. (2006). The nature of Drosophila melanogaster. *Current*
775    *Biology*, *16*(16), R623–R628. https://doi.org/10.1016/J.CUB.2006.07.042
776  Refsnider, J. M., & Janzen, F. J. (2010). Putting Eggs in One Basket: Ecological and Evolutionary
777    Hypotheses for Variation in Oviposition-Site Choice. *Annual Review of Ecology, Evolution,*
778    *and Systematics*, *41*, 39–57. https://doi.org/10.1146/ANNUREV-ECOLSYS-102209-144712
779  Ren, C., & Tao, Q. (2020). Neural Circuits Underlying Innate Fear. *Advances in Experimental*
780    *Medicine and Biology*, *1284*, 1–7. https://doi.org/10.1007/978-981-15-7086-5_1
781  Richmond, R. C., & Gerking, J. L. (1979). Oviposition site preference in Drosophila. *Behavior*
782    *Genetics*, *9*(3), 233–241. https://doi.org/10.1007/BF01066978
783  Rockwell, R. F., & Grossfield, J. (1978). Drosophila: Behavioral Cues for Oviposition. *The*
784    *American Midland Naturalist*, *99*(2), 361–368.
785  Rouquette, J., & Davis, A. J. (2003). Drosophila species (Diptera: Drosophilidae) oviposition
786    patterns on fungi: The effect of allospecifics, substrate toughness, ovipositor structure and
787    degree of specialisation. In *Eur. J. Entomol* (Vol. 100).
788    https://doi.org/10.14411/eje.2003.056
789  Sawin-McCormack, E. P., Sokolowski, M. B., & Campos, A. R. (1995). Characterization and
790    genetic analysis of Drosophila melanogaster photobehavior during larval development.
791    *Journal of Neurogenetics*, *10*(2), 119–135. https://doi.org/10.3109/01677069509083459
792  Schlenke, T. A., Morales, J., Govind, S., & Clark, A. G. (2007). Contrasting Infection Strategies in
793    Generalist and Specialist Wasp Parasitoids of Drosophila melanogaster. *PLoS Pathogens*,
794    *3*(10), e158. https://doi.org/10.1371/journal.ppat.0030158
795  Sendoya, S. F., Freitas, A. V. L., & Oliveira, P. S. (2009). Egg-laying butterflies distinguish
796    predaceous ants by sight. *American Naturalist*, *174*(1), 134–140.
797    https://doi.org/10.1086/599302
798  Sidhu, C. S., & Rankin, E. E. W. (2016). Honey Bees Avoiding Ant Harassment at Flowers Using
799    Scent Cues. *Environmental Entomology*, *45*(2), 420–426.

https://doi.org/10.1093/EE/NVV230

Soto-Yéber, L., Soto-Ortiz, J., Godoy, P., & Godoy-Herrera, R. (2018). The behavior of adult Drosophila in the wild. *PLoS ONE*, *13*(12), 1–26. https://doi.org/10.1371/journal.pone.0209917

Stensmyr, M. C., Dweck, H. K. M., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J., Grabe, V., Steck, K., Lavista-Llanos, S., Wicher, D., Sachse, S., Knaden, M., Becher, P. G., Seki, Y., & Hansson, B. S. (2012). A conserved dedicated olfactory circuit for detecting harmful microbes in drosophila. *Cell*, *151*(6), 1345–1357. https://doi.org/10.1016/j.cell.2012.09.046

Taylor, A. J., Muller, C. B., & Godfray, H. C. J. (1998). Effect of Aphid Predators on Oviposition Behavior of Aphid Parasitoids. *Journal of Insect Behavior*, *11*(2), 297–302.

Teal, P. E. A., & Tumlinson, J. H. (1989). Isolation, identification, and biosynthesis of compounds produced by male hairpencil glands of Heliothis virescens (F.) (Lepidoptera: noctuidae). *Journal of Chemical Ecology*, *15*(1).

Traniello, J. (2010). Pheidole: Sociobiology of a Highly Diverse Genus. In *Encyclopedia of Animal Behavior* (Vol. 2, pp. 699–706).

Van Mele, P., Vayssieres, J. F., Adandonon, A., & Sinzogan, A. (2009). Ant cues affect the oviposition behaviour of fruit flies (Diptera: Tephritidae) in Africa. *Physiological Entomology*, *34*(3), 256–261. https://doi.org/10.1111/j.1365-3032.2009.00685.x

Venken, K. J. T., Simpson, J. H., & Bellen, H. J. (2011). Genetic manipulation of genes and cells in the nervous system of the fruit fly. *Neuron*, *72*(2), 202–230. https://doi.org/10.1016/j.neuron.2011.09.021

Vijendravarma, R. K., Narasimha, S., & Kawecki, T. J. (2013). Predatory cannibalism in Drosophila melanogaster larvae. *Nature Communications 2013 4:1*, *4*(1), 1–8. https://doi.org/10.1038/ncomms2744

Wang, F., Wang, K., Forknall, N., Patrick, C., Yang, T., Parekh, R., Bock, D., & Dickson, B. J. (2020). Neural circuitry linking mating and egg laying in Drosophila females. *Nature*, *579*(7797), 101–105. https://doi.org/10.1038/s41586-020-2055-9

Worthen, W. B., Natasha Hipp, M., & Twardokus, C. T. (1993). Effects of Ant Predation and Larval Density on Mycophagous Fly Communities. *Oikos*, *66*(3), 526–532. Retrieved from https://www.jstor.org/stable/3544948

Yang, C.-H., Belawat, P., Hafen, E., Jan, L. Y., & Jan, Y.-N. N. (2008). Drosophila egg-laying site selection as a system to study simple decision-making processes. *Science*, *319*(5870), 1679–1683. https://doi.org/10.1126/science.1151842

Yang, Fa-zhong, Li, Y., & Yang, B. (2013). The inhibitory effects of rose powdery mildew infection on the oviposition behaviour and performance of beet armyworms. *Entomologia Experimentalis et Applicata*, *148*(1), 39–47. https://doi.org/10.1111/EEA.12069

Yang, Fazhong, Dong, W., Zhang, X., Li, Y., Zhou, S., Zhu, G., & Xiao, C. (2019). Volatile-organic compound changes in rose twigs consequent to infection with rose powdery mildew. *CHILEAN JOURNAL OF AGRICULTURAL RESEARCH*, *79*(4). https://doi.org/10.4067/S0718-58392019000400596

Zhang, L., Yu, J., Guo, X., Wei, J., Liu, T., & Zhang, W. (2020). Parallel Mechanosensory Pathways Direct Oviposition Decision-Making in Drosophila. *Current Biology*, *30*(16), 3075-3088.e4. https://doi.org/10.1016/j.cub.2020.05.076

Fig. 1: Flies alter egg depth when exposed to ants. *D. melanogaster* flies exposed to either
wild-caught *Forelius mccooki* (**A**) or lab-reared *Pheidole hyatti* (**B**) ants did not reduce
oviposition rates (unpaired, Welch's t-test; *F. mccooki*: *P*=0.3551; *P. hyatti*: *P*=0.0713). The
lower, but non-significant, exposed egg numbers in (B) are likely due to some flies being caught
by the ants (**C**). The arrowhead indicates a leg that was severed. We did not observe any flies
being caught by *F. mccooki* ants. (**D**) Photo of eggs in an exposure experiment, indicating the
three different egg depth categories. (**E**) Schematic of oviposition depth categories. (**F**) Ant
exposure significantly alters the distribution of eggs in each category (χ^2=28.93, *P*<0.0001, bars
represent mean ± S.E.M). (**G** and **H**) Both *F. mccooki* and *P. hyatti* ants significantly altered the
egg position index towards a preference for buried eggs rather than surface eggs (unpaired,
Welch's t-test: *F. mccooki*: *P*<0.0001; *P. hyatti*: *P*=0.0002).

Fig. 2: Fly eggs positioned deeply are protected from foraging ants. (A) Photo of ant foraging experiment with *P. hyatti* ants. (B) Buried fly eggs were significantly more likely to remain after ant foraging, while surface and partially buried eggs were removed by the ants (Surface vs. Partial: *P*=0.0135; Surface vs Buried: *P*<0.0001; Partial vs Buried: *P*=0.0001). (C) There was no significant difference in egg hatching rates across all three categories of the remaining eggs after ant foraging (Surface vs. Partial: *P*=0.6613; Surface vs Buried: *P*=0.2665; Partial vs Buried: *P*=0.8782). (Kruskal-Wallis test with Dunn's multiple comparison test). Bars represent mean ± S.E.M.

Fig. 3: The induced egg burying behavior is conserved across Drosophilids. (**A**)Three other fly
species did not reduce oviposition rates when exposed to *F. mccooki* ants (*D. simulans*:
P=0.9729; *D. yakuba*: *P*=0.4829; *D. virilis*: *P*=0.6040). (**B**) All three species altered their
oviposition depth when exposed to ants (*P*<0.0001 for all comparisons). (Two-way ANOVA with
Bonferroni multiple comparison test).

Fig. 4: Flies detect ants through olfactory cues. (**A**) Blind flies continued to bury their eggs
when exposed to *F. mccooki* ants, similar to control OreR flies (*P*<0.0001 for both comparisons).
(**B**) Both control heterozygous *orco* mutant flies and homozygous *orco* mutants buried their
eggs when exposed to ants, although the magnitude of the effect was much weaker for the
homozygous mutants (*orco²*/+: *P*<0.0001; *orco²*: *P*=0.0210; two-way ANOVA with Bonferroni
multiple comparison test for A and B). (**C**) Ant (*P. hyatt*i) odors extracted in hexane significantly
altered fly oviposition depth behavior, although not to the same extent as direct ant exposure
(Hex vs live ants: *P*<0.0001; Hex vs ant odors: *P*=0.0166; Live ants vs ant odors: *P*=0.0150;
Brown-Forsythe and Welch's ANOVA with Dunnett's multiple comparison test).

Fig. 5: **The ant pheromone 1-octadecanol induces fly egg burying.** Whole body-wash extract from *F. mccooki* ants was fractioned, resulting in 11 subfractions. (**A**) Flies did not alter oviposition rate after exposure to any of the odor subfractions, as compared to hexane-only exposed flies. (**B**) However, fraction 9 did significantly alter oviposition depth behavior (Hex vs fraction 9: P=0.0003). (**C**) The chromatogram from GC-MS of the whole-body wash extract revealed many peaks, including iridomyrmecin (peak 1). (**D**) Fraction 9 contained a single large peak that was identified as 1-octadecanol, which was also present in the whole-body wash extract (peak 2). (**E**) Flies exposed to *F. mccooki* (Fmcc) ant body wash, or higher concentrations of 1-octadecanol, significantly altered oviposition depth, as compared to hexane-exposed flies (Hex vs Fmcc: P<0.0001; Hex vs 10^{-6} M: P=0.0005; Hex vs 10^{-8} M: P=0.0010; Hex vs 10^{-10} M: P=0.0271; Hex vs 10^{-12} M: P>0.9999). (A, B, and E: Brown-Forsythe and Welch's ANOVA with Dunnett's multiple comparison test).

Fig. 6: Hymenopteran odors trigger the fly egg burying behavior. A phylogeny is shown of the different insect species whose odors were tested for the ability to induce the fly egg burying behavior. The number of independent replicates run for each insect species is shown, while the percentage of replicates resulting in significant fly egg burying is colored in red. Flies responded to odors from a diversity of hymenopterans, but rarely to odors from non-hymenopterans. * live insect exposure rather than body wash exposures; † odor extracted from larval stage; ‡ unlike the ant replicates that were derived from independent ant colonies, the *Leptoglossus zonatus* replicates were derived from the same colony, although the odor extractions were independent.

Supp. Fig. 1: Exposure to parasitoid wasps also alters fly oviposition behaviors. Unlike their
response to ant exposure, flies reduced oviposition rates when exposed to *L. heterotoma* (**A**,
P=0.0001) or *L. boulardi* (**C**, *P*=0.0025) parasitoid wasps. However, like their response to ant
exposure, flies oviposited eggs more deeply when exposed to *L. heterotoma* (**B**, *P*=0.0022) and
L. boulardi (**D**, *P*=0.0002) wasps. (Unpaired, Welch's t-test).

Supp. Fig. 2: Any cost to deeper oviposition remains unknown. Egg depth had no effect on
adult eclosion rates under normal conditions (**A**, *P*=0.2230) or when other survival risks were
present, such as the presence of older larvae (**B**, *P*=0.7234), cold overnight temperatures (**C**,
P=0.7577), or simulated rain (**D**, *P*=0.6774) (unpaired, Welch's t-test for A, B, and D; Mann-
Whitney test for C). (**E**) Maternal oviposition duration was significantly increased for partially
inserted eggs, but no timing difference was found between surface and buried eggs (Surface vs.
Partial: *P*<0.0001; Surface vs Buried: *P*<0.3475; Partial vs Buried: *P*=0.0386). (**F**) After
ovipositing, the female flies showed no difference in the length of their clean-and-rest phase
across the three egg position categories (Surface vs. Partial: *P*=0.; Surface vs Buried: *P*<0.;
Partial vs Buried: *P*=0.). (Kruskal-Wallis with Dunn's multiple comparison test in E and F). Bars
represent mean ± S.E.M.

Supp. Fig. 3: Olfaction is required to detect ants and wasps. (**A** and **B**) Blind flies exposed to *L.*
heterotoma parasitoid wasps maintained normal oviposition depth change similar to control
OreR flies or heterozygous controls (*P*<0.0001 for all comparisons in A and B). (**C**) Anosmic *orco*

mutant flies failed to alter oviposition depth behavior when exposed to *P. hyatti* ants (*orco²*/+:
P<0.0001; *orco²*: *P*=0.4805). (**D**) Similarly, anosmic flies also failed to respond to *L. heterotoma*
wasps (OreR: *P*<0.0001; *orco²*/+: *P*<0.0001; *orco²*: *P*>0.9999). (**E**) Control OreR flies with ablated
antenna had a significantly higher egg position index when exposed to *F. mccooki* ants than
fully intact flies. Ant-exposed flies with only maxillary palp ablations showed no significant
difference in oviposition depth compared to intact flies (intact exposed vs intact unexposed:
P<0.0001; intact exposed vs antenna-ablated exposed: *P*<0.0001; intact exposed vs maxillary
palp-ablated exposed: *P*=0.6090; intact exposed vs both-ablated exposed: *P*=0.0138). (**F** and **G**)
Flies responded to *L. heterotoma* wasp presence by burying eggs even with the silencing
(*Kir2.1*) or ablation (*hid*) of wasp odor-detecting Or49a sensory neurons (*P*<0.0001 for all
comparisons in F and G). (A-D, F, G: Two-way ANOVA with Bonferroni multiple comparison test;
E: Brown-Forsythe and Welch's ANOVA with Dunnett's multiple comparison test).

Supp. Fig. 4: Body wash solvent specificity in fly detection of ant and wasp odors. Different
solvents used to extract body odors varied in their success at inducing the fly egg burying
behavior. Flies responded to *L. heterotoma* parasitoid wasp body washes when either hexane
(Hex) (**A**, *P*=0.0110) or dichloromethane (DCM) (**B**, *P*=0.0099) were used. Body washes using
other solvents, such as acetonitrile (ACN) (**C**, *P*=0.2802) and water (**D**, *P*=0.0899), did not induce
the fly behavior. *P. hyatti* ant body washes using the ACN solvent also showed no significant
change in oviposition behavior (**E**, *P*=0.1249). (Unpaired, Welch's t-test).