1   **Prime Editing Efficiency and Fidelity are Enhanced in the Absence of**
2   **Mismatch Repair**
3

4   Ferreira da Silva J[1,2#], Oliveira GP[1#], Arasa-Verge EA[1], Moretton A[1,2], Timelthaler G[1],
5   Jiricny J[4,5], Loizou JI[1,2*]
6
[1]Institute of Cancer Research, Department of Medicine I, Comprehensive Cancer Centre,
Medical University of Vienna, Borschkegasse 8a, 1090 Vienna, Austria
[2]CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences,
Vienna, Austria
[3]Institute of Life Sciences, University of Zurich, Winterthurerstrasse 190, 8057 Zurich,
Switzerland
[4]Institute of Biochemistry of the ETH Zurich, Otto-Stern-Weg 3, 8093 Zurich, Switzerland.
[#]Equal contribution
*Correspondence
16
17   Correspondence: joanna.loizou@meduniwien.ac.at (JIL)

19   **Running title**
20
21

22   **Keywords**
23   Prime editing; mismatch repair; genome engineering; DNA repair
24

Abstract

Prime editing is a powerful genome engineering approach that enables the introduction of
base substitutions, insertions and deletions, into any given genomic locus. But prime editing,
at even the same locus, can exhibit wildly different efficiencies in various cell backgrounds. It
is unclear what determines these variations in efficiencies in a given cellular context. Through
a focused genetic screen targeting DNA repair factors, we show that the efficiency of prime
editing is attenuated by the mismatch repair pathway. The accumulation of the mismatch
repair protein MLH1 at sites of prime editing, indicates that mismatch repair acts at these
regions to directly counteract the insertion of the edit. Consequently, ablation of mismatch
repair yields an up to 17-fold increase in prime editing efficiency across different human cell
lines, several types of edits and multiple genomic loci. Our results shed new light on the cellular

requirements for prime editing and identify that ablation of mismatch repair increases editing
efficiency and fidelity.
38
Introduction
Genome editing technologies, using CRISPR-Cas, are powerful approaches that can be used
for functional genomics, as well as for therapeutic applications (Jinek et al., 2012). Their
optimal implementation, however, relies on a thorough understanding of the DNA metabolic
pathways and mechanisms that are responsible for editing outcomes, as well as the activity
of these pathways in different tissue contexts (Richardson et al., 2018; Yeh et al., 2019;
Ferreira da Silva et al., 2021; Hussmann et al., 2021). Amongst the toolbox of Cas9 proteins
that are used for genome engineering are Cas9 nucleases that can be directed to a specific
region of the genome by a single-guide RNA (sgRNA). Following the generation of a DNA
double-strand break (DSB), the endogenous cellular DNA end-joining pathways can repair the
lesion in an error prone manner, leading to insertions and deletions (indels) that culminate in
loss-of-function alleles (Bothmer et al., 2017). This approach can be further adapted to include
a donor template, either single-stranded or double-stranded, containing a desired edit. Here,
following the generation of a DNA DSB, homology directed repair (HDR) mediates the
insertion of the desired template, thus editing the genome. This approach can be used to
introduce or correct mutations with high precision. However, since HDR is an inefficient
mechanism, depends on potentially deleterious DSBs and requires cell division, other
approaches have been developed.
57
Amongst these are base editors (BE) which use nucleobase modification chemistry to
efficiently and precisely incorporate single nucleotide variants into the genome of living cells
(Komor et al., 2016; Gaudelli et al., 2017; Gu et al., 2021). BEs are however limited to single-
base substitutions. Thus, more recently, prime editing (PE) was developed as a highly
versatile genome editing approach that allows for targeted insertions, deletions and all
possible combinations of base substitutions in a precise manner, independent on cell division
(Anzalone et al., 2019). PE utilises a fusion of a Cas9(H840A) nickase (Jinek et al., 2012) and
reverse transcriptase (RT) that is targeted to a precise genomic region by a PE guide RNA
(pegRNA). The pegRNA encodes, in its 3' extension region, the genetic information that is
copied to the targeted genomic locus by the RT. The installation of the desired edit is
dependent on the resolution of a DNA intermediate that contains redundant single-stranded
DNA (ssDNA) flaps. Preferential 5' flap excision and 3' flap ligation drives the integration of
the edit, resulting in a DNA heteroduplex containing one edited- and one non-edited strand
(PE2). The editing outcome depends on the resolution of this heteroduplex. Utilising a sgRNA
that directs the nicking of the non-edited strand, either concurrently to the edit installation

(PE3), or afterwards (PE3b), has been shown to increase PE efficiencies (Anzalone et al.,

2019).

A notable feature of prime editing is its highly variable rates of efficiency across different

genetic backgrounds, even within the same genomic region and using the same pegRNA

(Anzalone et al., 2019). To address whether this could be explained by different DNA repair

capacities, we performed a targeted genetic screen aimed at identifying DNA repair factors

involved in PE. Here, we uncover an unexpected inhibitory role for the mismatch repair (MMR)

pathway in PE and show that MMR localises to sites of PE to counteract edit installation. Thus,

MMR depletion increases PE efficiency and fidelity across different edit sites, types and cell

lines.

Results

A targeted genetic screen identifies the mismatch repair pathway as inhibitory for prime

editing

To investigate the DNA repair requirements for PE, we conducted a targeted genetic screen,

utilising a collection of isogenic knockout cell lines in the human near-haploid cell line HAP1

(**Supplementary Table 1**). The 32 targeted genes were selected to represent divergent

functions within all known human DNA repair pathways. The library thus provided a

comprehensive coverage of the DNA damage response. The cell lines received the PE

machinery, including the Cas9(H840)-RT and a pegRNA encoding a 5-base pair (bp) deletion

in the *HEK3* locus. Editing efficiency was then determined by amplicon sequencing of the

genomic locus.

Wild-type HAP1 cells were remarkably inefficient at prime editing (<1% alleles edited). In

contrast, isogenic HAP1 cell lines mutated at the *MLH*1, *PSM2*, *MSH2*, *EXO1* and *MSH3* loci

displayed higher levels of PE (ranging from 2.3 to 7.9-fold) (**Figure 1A**). Deficiencies in other

DNA repair pathways had little impact on the efficacy of prime editing. This finding clearly

indicates that MMR plays an inhibitory role in PE. Of all MMR genes targeted in the screen,

only the loss of *MSH6* failed to increase PE efficiency.

The MMR pathway evolved to correct base/base mispairs and indels arising in DNA during

replication. Base/base mismatches and small indels of 1-3 nucleotides are recognised by the

heterodimer MSH2-MSH6, whereas indels are generally recognised by the MSH2-MSH3

heterodimer (Drummond et al., 1995; Palombo et al., 1995, 1996; Acharya et al., 1996; Gradia

et al., 1997). Detection of the lesion is followed by an ATP-dependent conformational change

of these complexes and recruitment of the MLH1-PMS2 (Li and Modrich, 1995) or MLH1-

MLH3 (Lipkin et al., 2000) heterodimers. Assembly of this complex together with RFC and

PCNA (Pluciennik et al., 2010) activates cryptic endonucleases of the PMS or MLH3 proteins,

which then introduce DNA single-strand breaks (SSBs) into the nascent DNA strand in the

vicinity of the mismatch. These SSBs act as entry points for EXO1, which degrades the

nascent strand up to, and some distance past, the misincorporated nucleotide(s) (Kadyrov et

al., 2006). The resulting gap is filled-in by polymerase δ and the remaining nick is ligated by

DNA ligase I (Stojic et al., 2004; Iyer et al., 2006; Fishel, 2015). The edit introduced in our

screen, a 5bp deletion, is a preferential target of the MSH2-MSH3 heterodimer, but not the

MSH2-MSH6 heterodimer (Palombo et al., 1996), thus explaining the lack of an effect on

editing upon loss of MSH6 (**Figure 1A**). This result highlights the highly specialised nature of

the DNA damage response and shows that different edits might be addressed by distinct

factors.

Figure 1: Mismatch repair inhibits prime editing in human cells. A) Genetic screen in a
panel of 32 HAP1 isogenic knockout cell lines covering different DNA damage repair

pathways, as well as their wild-type (WT) counterpart, showing the efficiency of installation of
a 5 base pair (bp) deletion in the *HEK3* locus, using PE2. Values correspond to editing
efficiency, measured by amplicon sequencing analysis in two independent biological
replicates, with two technical replicates each. Each radial line represents an increment of 1%.
B) PE efficiency of a 5 bp deletion in the *HEK3* locus in different mismatch repair-deficient
human cell lines, along with their respective complemented counterparts. Values correspond
to editing efficiency, measured by amplicon sequencing analysis in three independent
biological replicates, with two technical replicates each. **C)** PE efficiency of different types of
edits (*RNF2* locus) in the HEC59 cell line complemented with chromosome 2 (MMR proficient;
MMR+) and HEC59 cell line (MMR deficient; MMR-). Values correspond to editing efficiency,
measured by amplicon sequencing analysis in three independent biological replicates, with
two technical replicates each. **D)** Comparison of different types of PE efficiency (PE2, PE3
and PE3b) upon installation of a G>T substitution in the *FANCF* locus, in HEC59 cells
complemented with chromosome 2 (MMR+) and HEC59 cells (MMR-). Values correspond to
editing efficiency, measured by amplicon sequencing analysis in three independent biological
replicates, with two technical replicates each. Ns, p-value non-significant; *, p-value < 0.05; **,
p-value < 0.01; ***, p-value<0.001; ****, p-value < 0.0001.

Mismatch repair hinders PE2 and PE3 across several human cell lines, genomic loci
and edits

To further explore the inhibitory role of MMR in PE, we measured editing efficiency and fidelity

of the *HEK3* locus in a panel of MMR-deficient human cell lines, alongside their complemented

counterparts. We used the colorectal cancer line HCT116, which is deficient for both MSH3

and MLH1, along with an HCT116 MMR-proficient cell line complemented with both

chromosomes 5 and 3 (**Supplementary Figure 1A)** (Koi et al., 1994; Haugen et al., 2008).

Additionally, the endometrial adenocarcinoma cell line HEC59, which is deficient for MSH2,

was used together with its MMR-proficient counterpart that was complemented with

chromosome 2 (**Supplementary Figure 1B**) (Umar et al., 1997). Finally, we used a

doxycycline-inducible model of MLH1 deficiency in the embryonic kidney cell line HEK293T

(293T-Lα) (**Supplementary Figure 1C**) (Cejka et al., 2003). All cell lines showed significantly

increased PE2 editing (1.7 to 5-fold) when MMR was ablated, compared to their MMR-

proficient counterparts (**Figure 1B**). Importantly, even though PE efficiencies were increased

by MMR deficiency, this did not come at the expense of higher indel frequencies at the

amplicon region (**Supplementary Figure 1D**). Indeed, we observed that loss of MMR

prevented unwanted indels at the *HEK3* locus in the 293T-Lα cell line (**Supplementary Figure**

1D).

To investigate further the substrates of MMR in PE, we measured the editing efficiencies of a

transition (G>A), two transversions (C>G and C>T), a 1 bp insertion and a 3 bp deletion, all

within a different endogenous locus. We found that active MMR significantly diminished the

efficiency of all these edits in HEC59 cells (ranging from 2.8 to 17-fold) (**Figure 1C**). To test

the inhibitory role of MMR on different types of PE, we measured the efficiency of PE2, PE3

and PE3b in the *FANCF* locus, via the installation of a G>T substitution. Editing efficiency was

improved by loss of MMR for all types of PE, albeit to a lesser extent for PE3b (**Figure 1D**). In

summary, MMR counteracts PE efficiency across different edits and different genomic loci.

The mismatch repair factor MLH1 is directly recruited to sites of prime editing

We hypothesized that MMR could reduce PE efficiency either directly at the edit site, or

indirectly by other mechanisms. To address this, we determined if the MMR factor MLH1 was

recruited to sites of active and ongoing editing, through co-localization with Cas9(H840)-RT.

Cas9(H840)-RT was directed to human repetitive telomeric regions, a strategy that has proven

efficient for CRISPR imaging (Chen et al., 2013) (**Figure 2A**). With this experimental

approach, Cas9(H840)-RT was shown to co-localise with TRF1, a major telomere-binding

protein (Wang et al., 2008) (**Supplementary Figures 2A-B**). Similarly, endogenous MLH1 co-

localised with Cas9(H840)-RT-foci, in a pegRNA-dependent manner (**Figures 2B-C**). Based

on these results, we postulate that PE inhibition by MMR occurs through recruitment of MMR

factors to sites of editing, where the heteroduplex DNA is recognised as a mismatch and

removed, using the non-edited strand as a template, and thus restoring the non-edited

sequence of nucleotides.

Figure 2: MLH1 is enriched at prime editing sites and its ablation can be exploited to
increase prime editing efficiency. A) Schematic representation of the approach used to co-
localise MLH1 at PE sites. The PE machinery is directed to human telomeric repetitive regions,

where the telomeric protein, TRF1, also binds. **B)** Representative super resolution images of
Cas9(H490)-RT, MLH1 and both merged, in U2OS cells. Images were taken 24 hours
following reverse transfection of Cas9(H480)-RT and in the presence or absence of a pegRNA
targeting telomeric repeats. Scale bars = 5 μm **C)** Quantification of Cas9(H480)-RT foci that
co-localise with MLH1. Values correspond to quantification of 13 cells in three independent
biological replicates. **D)** Immunoblot for MLH1 and tubulin in HEK293 cell extracts three days
post transfection with non-targeting and MLH1 siRNA pools (of four siRNAs). **E)** PE efficiency
of a 5 bp deletion in the *HEK3* locus in HEK293 cells transfected with non-targeting and MLH1
siRNA pools, two days prior to PE vector delivery. Values correspond to editing efficiency,
measured by Sanger sequencing and analysed by TIDE analysis (Brinkman et al., 2014), in
three independent biological replicates, with two technical replicates each. **F)** Immunoblot for
MLH1 and tubulin in HAP1 cell extracts. Expression of dTAG-MLH1 is abrogated upon 1-hour
treatment with 500nM dTAG-7 ligand ('+' condition). Recovery of dTAG-MLH1 expression is
visible 24 hours after ligand removal ('24h' condition) **G)** PE efficiency of a 5 bp deletion in
the *HEK3* locus in HAP1 dTAG-MLH1 cells in the presence and absence of 500 nM dTAG-7.
Values correspond to editing efficiency, measured by Sanger sequencing and analysed by
TIDE analysis (Brinkman et al., 2014), in three independent biological replicates, with two
technical replicates each. Ns, p-value non-significant; *, p-value < 0.05; **, p-value < 0.01;
****, p-value < 0.0001.
208
209

Mismatch repair inhibition can be exploited to increase prime editing efficiency

We next sought to inhibit MMR as a strategy to improve PE efficiency. Since MMR loss leads

to increased mutational burden and genome instability (Jiricny, 2006), long-term inhibition of

MMR is not desirable. Thus, to achieve transient MMR ablation, we depleted MLH1 in HEK293

cells with a pool of siRNAs (**Figure 2D**), and subsequently showed that this effectively

increased PE efficiency in the *HEK3* locus (**Figure 2E**).

An alternative way to achieve transient loss of function is through targeted protein degradation.

The degradation tag (dTAG) system has proved to be an efficient strategy for rapid and

transient ligand-induced targeted protein degradation (Winter et al., 2015). Using CRISPR-

mediated knock-in, we introduced the dTAG system into the *MLH1* locus of HAP1 cells which

allowed for the targeted degradation of MLH1, after ligand treatment (**Figure 2F**). Upon MLH1

degradation, the efficiency of PE was increased by 2-fold (**Figure 2G**). Taken together, these

results indicate that transient inhibition of MMR is a promising strategy to increase the

efficiency of PE.

Discussion

Here, we describe an inhibitory role of the MMR pathway during PE, across different human

cell lines, genomic loci and types of edits, thus challenging the current view that this is a

pathway that is required for edit installation (Petri et al., 2021). We propose that MMR proteins

recognise the heteroduplex DNA, containing the desired edit introduced by the pegRNA, as a

mismatch. The mismatch is promptly removed, and the non-edited strand is used as a

template to restore the original sequence of nucleotides. Hence, suppression of MMR
increases PE efficiency.

Our findings suggest that the improvement in PE efficiencies upon loss of MMR activity does
not come at the cost of indel generation around the edit site, which would be an undesirable
outcome of editing. Much effort has gone into improving genome editing technologies through
both computational and experimental approaches aimed at improving editing efficiencies and
fidelity. While globally manipulating MMR in dividing cells would come at the expense of
mutations, precisely interfering with MMR at the edit site might offer an approach to specifically
perturb this pathway only at the sites of PE. Similar approaches have proved useful for
improving the efficiency of HDR (Charpentier et al., 2018; Rees et al., 2019), suggesting that
their development for PE might also be valuable.

The highly specialised nature of the DNA damage response suggests that different edits might
be dealt with by different pathways, and even different factors within the same pathway. Our
data on the lack of involvement of MSH6 in the repair of a 5bp deletion supports this
hypothesis. While it is known that MMR functions to repair base substitutions and small indels,
it remains to be seen what the size limitation of an edit is to be removed by this pathway, when
generated by PE. Future directions might provide a deeper understanding of the specificity
and size of MMR substrates in PE to determine if perturbing MMR could be used to enhance
the insertion of small tags such as histidine or hemagglutinin tags.

Overall, our data shed new light on the DNA repair mechanisms of a new and highly promising
genome editing technology. This knowledge can be applied in further development of prime
editors, as well as in the design of therapeutic strategies. Finally, taking into consideration the
expression levels of MMR factors in particular tissue or cell types might predict PE efficiencies.

Methods
Plasmids and oligos
DNA oligos were obtained from Integrated DNA Technologies (IDT) unless otherwise noted.
All plasmids used for prime editing experiments were obtained from Addgene (pCMV-PE2
no.132775, pU6-pegRNA-GG-acceptor no.132777). PegRNAs were cloned using BsaI
Golden Gate assembly (NEB), as described previously (Anzalone et al., 2019). The pegRNA
targeting telomeres included a stem loop extension as described in. SgRNAs utilized in PE3
and PE3b experiments were cloned in the plenti-sgRNA puro vector (Addgene no. 104990),

using BsmBI Golden Gate assembly (NEB), following the manufacturer's instructions. The

pegRNA targeting telomeres included a stem loop extension as described in (Chen et al.,

2013). Sequences of sgRNA and pegRNA constructs are listed in **Supplementary Table 2**.

All plasmids for mammalian cell experiments were purified using the Plasmid Plus Midi Kit

(Qiagen) or the Spin Miniprep Kit (Qiagen), both including endotoxin removal steps.

Mammalian cell culture

All cells were grown at 3% oxygen at 37°C and routinely checked for possible mycoplasma

contamination. Human HAP1 cells were obtained from Horizon Discovery and were grown in

Iscove's Modified Dulbecco's Medium (IMDM) (Gibco), containing L-glutamine and 25nM

HEPES and supplemented with 10% Fetal Bovine Serum (FBS) (Gibco) and 1%

Penincillin/Streptomycin (P/S) (Sigma-Aldrich). U2OS cells were purchased from ATCC cell

repository and cultured in DMEM (Gibco), supplemented with 10% FBS and 1% P/S. HEC59,

wild-type and complemented with chromosome 2, were provided by Josef Jiricny (ETH Zurich,

Switzerland) and cultured in F12 DMEM with 10% FBS and 1% P/S. HEC59 complemented

cells were cultured with 400 µg/mL of geneticin (G418, Gibco). HCT116 cells, wild-type and

complemented with both chromosomes 3 and 5, were provided by Josef Jiricny (ETH Zurich,

Switzerland) and cultured with McCoy's 5A medium (Gibco), with 10% FBS and 1% P/S.

HCT116 cells complemented with chromosomes 3 and 5 were cultured with 400 µg/mL

geneticin (G418, Gibco) and 6 ug/mL blasticidine (Invivogen). 293T-Lα were provided by Josef

Jiricny (ETH Zurich, Switzerland) and cultured in DMEM medium (Gibco) supplemented with

10% FBS or Tet-system approved FBS (Takara Bio), 1% P/S, 100 µg/mL zeocin (Gibco) and

300 µg/mL hygromycin (Gibco). 293T-Lα were grown in doxycycline (1µg/mL) for 7 days

before any experiment, to completely deplete MLH1 expression. Doxycycline was replenished

in the medium every 2 days.

Focused DNA repair genetic screen

CRISPR-Cas9 knockouts of DNA repair genes were generated in collaboration with Horizon

Genomics. Sequences of sgRNAs were designed by Horizon Genomics or with the use of

http://chopchop.cbu.uib.no/. sgRNA sequences and frameshift mutations can be found in

Supplementary Table 1.

For the genetic screen, 80,000 cells were seeded in technical duplicates in 12-well plates.

Cells were transfected the day after with 636ng of pCMV-PE2 and 159ng of the *HEK3* pegRNA

inducing a 5bp deletion, per well. 1.6uL Lipofectamine 2000 (ThermoFisher Scientific) were

used per well, following the manufacturer's instructions. A separate transfection control was

performed using 795ng of the pmaxGFP™ vector (Lonza). Medium containing transfection

reagents was removed 16h post-transfection. Transfection efficiency was measured 48h after

transfection, by determining the percentage of GFP+ cells by flow cytometry. Genomic DNA

was harvested 96h post-transfection, using the QUIAmp DNA Blood Mini kit (Quiagen),

following the manufacturer's instructions.

Transfection and genomic DNA preparation of mismatch repair-deficient human cell

lines

HEC59, HCT116 and HEK293Tα were seeded in 48-well plates in duplicates (50,000

cells/well). Transfections were performed the next day, using 1µL Lipofectamine 2000

(ThermoFisher Scientific) per well, following the manufacturer's instructions. Cells were

transfected with 320 ng of the pCMV-PE2 vector, 80 ng of the respective pegRNA and, for

PE3 and PE3b, 33.2 ng of the nicking sgRNA, per well. A transfection control was performed

in parallel, by transfecting 400 ng per well of the pmaxGFP^{TM} vector (Lonza).

Genomic DNA was extracted 96 hours after transfection, by removing the medium,

resuspending the cells in a lysis solution (100 µL DirectPCR Lysis Reagent (Cell) (Viagen

Biotech), 76 µL of water and 4 µL Proteinase K) and incubating 45 minutes at 55 ^{o}C and 45

minutes at 85 ^{o}C.

High-throughput DNA sequencing of genomic samples

Genomic sites of interest were amplified from genomic DNA samples and sequenced on an

Illumina Miseq or NextSeq, depending on the number of pooled samples. Amplification primers

containing Illumina forward and reverse primers **(Supplementary Table 2)** were used for a

first round of PCR (PCR1) to amplify the genomic region of interest. A mixture of staggered

forward primers was used to create complexity. PCR1 reactions were performed in a final

volume of 25 µL, using 0.5µM of each forward and reverse primers, 1 µL genomic DNA and

12.5 µL of Phusion U Multiplex PCR 2x Master Mix (ThermoFisher Scientific). PCR1 was

carried as following: 98 ^{o}C 2 min, 30 cycles [98^{o}C 10 s, 61 ^{o}C 20 s, 72 ^{o}C 30 s], followed by a

final extension of 72 ^{o}C for 7 min. Unique Illumina dual index barcode primer pairs were added

to each sample in a secondary PCR reaction (PCR2). PCR2 was performed in a final volume

of 25 µL, using 0.5 µM of each unique forward and reverse Illumina barcoding primer pair,1

µL of unpurified PCR1 reaction and 12.5 µL of of Phusion U Multiplex PCR 2x Master Mix.

PCR2 was carried as following: 98 ^{o}C 2 min, 12 cycles [98^{o}C 10 s, 61 ^{o}C 20 s, 72 ^{o}C 30 s],

followed by a final extension of 72 ^{o}C for 7 min. PCR products were analysed by

electrophoresis in a 1% (w/v) agarose gel and purified using magnetic AMPure XP beads

(Beckman Coulter), using a ratio of beads:PCR product of 2:1. DNA concentration was

measured by fluorometric quantification (Qubit, ThermoFisher Scientific) and sequenced on
an Illumina instrument, according to manufacturer's instructions.
343
Sequencing reads were demultiplexed using MiSeq Reporter (Illumina) and alignment of
amplicon sequences to a reference sequence was performed using CRISPResso 2 (Clement
et al., 2019). CRISPResso2 was ran in standard mode and prime editing yield was calculated
as: number of aligned reads containing the desired edit/total aligned reads. Percentage of
indels was calculated as: number of aligned reads containing indels that are not the desired
edit/ total number of aligned reads.
350

siRNA transfections
The following siRNAs from Dhamacon (used at a final concentration of 100nM) were used in
this study: MLH1 SMARTpool ON-TARGETplus (L-003906-00-0005) and Non-targeting
control SMARTpool ON-TARGETplus (D-001810-10-05). siRNA transfections in HEK293 cells
were performed using Dharmafect 1 following manufacturer´s instructions. siRNA delivery was
performed 48 hours prior to transfection of prime editing vectors.
357

Generation of dTAG-MLH1 HAP1 cell line
A targeting vector encoding for the BSD-dTAG sequence (amplified from pCRIS-PITChv2-
BSD-dTAG (BRD4) Addgene no.91792) surrounded by two 1kb-long homology arms
upstream and downstream of the start codon of MLH1 was generated using Gibson assembly
(NEB). An in-vitro transcribed sgRNA targeting the region spanning the start codon of MLH1
was generated as previously described (Richardson et al., 2018). Cas9 protein (IDT) together
with the targeting vector and in-vitro transcribed sgRNA were nucleofected into 200,000
haploid cells in 16-well strips, using a 4D Nucleofector (Lonza) and the program DS-118.
Three days after nucleofection, 10 µg/ml blasticidin (Invivogen) were added to the culture
medium for one week after which single and haploid clones were sorted into 96-well plates.
Clonal haploid populations were grown and validated for correct homology-directed repair by
LR-PCR and immuno blot analysis. dTAG-7 (R&D Systems), at the final concentration of 500
nM, was used to test target degradation in the generated clones and all further targeted protein
degradation experiments.
372

Immunoblotting
Cell extracts were prepared in RIPA lysis buffer (NEB) supplemented with protease inhibitors
(Sigma) and phosphatase inhibitors (Sigma, NEB). Immunoblots were performed using
standard procedures. Protein samples were separated by sodium dodecyl sulfate-

polyacrylamide gel electrophoresis (SDS-PAGE) (3–8% gradient gels, Invitrogen) and
subsequently transferred onto nitrocellulose membranes. Primary antibodies for MLH1
(554073, BD Pharmigen), MSH2 (ab52266, Abcam), MSH3 (ab69619, Abcam), Tubulin (3873,
Cell Signaling) and ß-Actin (A5060, Sigma) were used at 1:1,000. Secondary antibodies were
used at 1:5,000 (HRP-conjugated goat anti-mouse or anti-rabbit IgG from Jackson
Immunochemicals). Immunoblots were imaged using a Curix 60 (AGFA) table-top processor.
384

Immunofluorescence
U2OS cells were reverse transfected using PEI together with pCMV-PE2 in the presence and
absence of pegRNA targeting telomeres. 50.000 cells were seeded per well of μ-Slide 8 well
(Ibidi) chambered coverslip plates. Pre-extraction was performed using 0.1% Tween in PBS
24 hours after reverse transfection. Cells were then fixed with 4% para-formaldehyde and fixed
cells were processed for immunofluorescence using the following antibodies: anti-Cas9 (Cell
Signalling, 14697), anti-TRF1 (Abcam, ab1423), anti-MLH1 (ThermoFisher, A300-015A), anti-
MSH2 (Bethyl, A300-452A). Primary antibodies were diluted 1:500 and secondary antibodies
(Alexa Fluor® 568 goat anti-mouse and Alexa Fluor® 488 goat anti-rabbit, LifeTechnologies)
were diluted 1:2000.
395

Imaging
16-bit fluorescence images were acquired using an Olympus IXplore spinning disk confocal
microscope (equipped with the Yokogawa CSU-W1 with 50 μm pinhole disk and a Hamamatsu
ORCA Fusion CMOS camera). A 60X oil immersion objective (NA 1,42) in combination with a
3,2X magnification lens (equaling 192X total magnification) was used for superresolution
imaging of fixed cells and z-stacks with a 0,24μm slice interval were acquired. These z-stacks
were then processed using the Olympus 3D deconvolution software (constrained iterative
deconvolution, using automatic background removal and noise reduction, filter using
advanced maximum likelihood algorithm and 5 iterations). Finally "maximum-z" projection
images of the deconvoluted z-stacks were generated. For further data analysis, the ImageJ
(NIH) distribution FIJI was used. Nuclear foci were counted manually and at least 15 cells per
condition were imaged in each experiment. Quantification of the foci was performed manually
based on maximum intensity projections.
409

References
Acharya, S., Wilson, T., Gradia, T., Kane, M., Guerrette, S., Marsischky, G., et al. (1996).
hMSH2 forms specific mispair-binding complexes with hMSH3 and hMSH6. *Proc. Natl.*

Acad. Sci. U. S. A. 93, 13629–13634. doi:10.1073/PNAS.93.24.13629.

Anzalone, A. V, Randolph, P. B., Davis, J. R., Sousa, A. A., Koblan, L. W., Levy, J. M., et al.

(2019). *Search-and-replace genome editing without double-strand breaks or donor*

DNA. doi:10.1038/s41586-019-1711-4.

Bothmer, A., Phadke, T., Barrera, L. A., Margulies, C. M., Lee, C. S., Buquicchio, F., et al.

(2017). Characterization of the interplay between DNA repair and CRISPR/Cas9-

induced DNA lesions at an endogenous locus. *Nat. Commun.* 8, 13905.

doi:10.1038/ncomms13905.

Brinkman, E. K., Chen, T., Amendola, M., and van Steensel, B. (2014). Easy quantitative

assessment of genome editing by sequence trace decomposition. *Nucleic Acids Res.*

42, e168–e168. doi:10.1093/NAR/GKU936.

Cejka, P., Stojic, L., Mojas, N., Russell, A. M., Heinimann, K., Cannavó, E., et al. (2003).

Methylation-induced G2/M arrest requires a full complement of the mismatch repair

protein hMLH1. *EMBO J.* 22, 2245. doi:10.1093/EMBOJ/CDG216.

Charpentier, M., Khedher, A. H. Y., Menoret, S., Brion, A., Lamribet, K., Dardillac, E., et al.

(2018). CtIP fusion to Cas9 enhances transgene integration by homology-dependent

repair. *Nat. Commun.* 9, 1–11. doi:10.1038/s41467-018-03475-7.

Chen, B., Gilbert, L., Cimini, B., Schnitzbauer, J., Zhang, W., Li, G., et al. (2013). Dynamic

imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. *Cell*

155, 1479–1491. doi:10.1016/J.CELL.2013.12.001.

Clement, K., Rees, H., Canver, M. C., Gehrke, J. M., Farouni, R., Hsu, J. Y., et al. (2019).

CRISPResso2 provides accurate and rapid genome editing sequence analysis. *Nat.*

Biotechnol. 2019 373 37, 224–226. doi:10.1038/s41587-019-0032-3.

Drummond, J., Li, G., Longley, M., and Modrich, P. (1995). Isolation of an hMSH2-p160

heterodimer that restores DNA mismatch repair to tumor cells. *Science* 268, 1909–

1912. doi:10.1126/SCIENCE.7604264.

Ferreira da Silva, J., Meyenberg, M., and Loizou, J. I. (2021). Tissue specificity of DNA

repair: the CRISPR compass. *Trends Genet.* 0. doi:10.1016/J.TIG.2021.07.010.

Fishel, R. (2015). Mismatch repair. *J. Biol. Chem.* 290, 26395–26403.

doi:10.1074/JBC.R115.660142.

Gaudelli, N. M., Komor, A. C., Rees, H. A., Packer, M. S., Badran, A. H., Bryson, D. I., et al.

(2017). Programmable base editing of T to G C in genomic DNA without DNA cleavage.

Nature 551, 464–471. doi:10.1038/nature24644.

Gradia, S., Acharya, S., and Fishel, R. (1997). The Human Mismatch Recognition Complex

hMSH2-hMSH6 Functions as a Novel Molecular Switch. *Cell* 91, 995–1005.

doi:10.1016/S0092-8674(00)80490-0.

Gu, S., Bodai, Z., Cowan, Q. T., and Komor, A. C. (2021). Base editors: Expanding the types

of DNA damage products harnessed for genome editing. *Gene Genome Ed.* 1, 100005.

doi:10.1016/J.GGEDIT.2021.100005.

Haugen, A. C., Goel, A., Yamada, K., Marra, G., Nguyen, T.-P., Nagasaka, T., et al. (2008).

Genetic Instability Caused by Loss of MutS Homologue 3 in Human Colorectal Cancer.

Cancer Res. 68, 8465–8472. doi:10.1158/0008-5472.CAN-08-0002.

Hussmann, J. A., Ling, J., Ravisankar, P., Yan, J., Cirincione, A., Xu, A., et al. (2021).

Mapping the Genetic Landscape of DNA Double-strand Break Repair. *bioRxiv*,

2021.06.14.448344. doi:10.1101/2021.06.14.448344.

Iyer, R., Pluciennik, A., Burdett, V., and Modrich, P. (2006). DNA mismatch repair: functions

and mechanisms. *Chem. Rev.* 106, 302–323. doi:10.1021/CR0404794.

Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., and Charpentier, E. (2012). A

programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity.

Science (80-.). 337, 816–821. doi:10.1126/science.1225829.

Jiricny, J. (2006). The multifaceted mismatch-repair system. *Nat. Rev. Mol. Cell Biol. 2006*

75 7, 335–346. doi:10.1038/nrm1907.

Kadyrov, F., Dzantiev, L., Constantin, N., and Modrich, P. (2006). Endonucleolytic function of

MutLalpha in human mismatch repair. *Cell* 126, 297–308.

doi:10.1016/J.CELL.2006.05.039.

Koi, M., Umar, A., Chauhan, D. P., Cherian, S. P., Carethers, J. M., Kunkei, T. A., et al.

(1994). Human Chromosome 3 Corrects Mismatch Repair Deficiency and Microsatellite

Instability and Reduces N-Methyl-N′-nitro-N-nitrosoguanidine Tolerance in Colon Tumor

Cells with Homozygous hMLH1 Mutation. *Cancer Res.* 54.

Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A., and Liu, D. R. (2016). Programmable

editing of a target base in genomic DNA without double-stranded DNA cleavage.

Nature 533, 420–424. doi:10.1038/nature17946.

Li, G., and Modrich, P. (1995). Restoration of mismatch repair to nuclear extracts of H6

colorectal tumor cells by a heterodimer of human MutL homologs. *Proc. Natl. Acad. Sci.*

U. S. A. 92, 1950–1954. doi:10.1073/PNAS.92.6.1950.

Lipkin, S., Wang, V., Jacoby, R., Banerjee-Basu, S., Baxevanis, A., Lynch, H., et al. (2000).

MLH3: a DNA mismatch repair gene associated with mammalian microsatellite

instability. *Nat. Genet.* 24, 27–35. doi:10.1038/71643.

Palombo, F., Gallinari, P., Iaccarino, I., Lettieri, T., Hughes, M., D'Arrigo, A., et al. (1995).

GTBP, a 160-kilodalton protein essential for mismatch-binding activity in human cells.

Science (80-.). 268, 1912–1914. doi:10.1126/SCIENCE.7604265.

Palombo, F., Iaccarino, I., Nakajima, E., Ikejima, M., Shimada, T., and Jiricny, J. (1996).

hMutSβ, a heterodimer of hMSH2 and hMSH3, binds to insertion/deletion loops in DNA.

Curr. Biol. 6, 1181–1184. doi:10.1016/S0960-9822(02)70685-4.

Petri, K., Zhang, W., Ma, J., Schmidts, A., Lee, H., Horng, J. E., et al. (2021). CRISPR prime

editing with ribonucleoprotein complexes in zebrafish and primary human cells. *Nat.*

Biotechnol. 2021, 1–5. doi:10.1038/s41587-021-00901-y.

Pluciennik, A., Dzantiev, L., Iyer, R. R., Constantin, N., Kadyrov, F. A., and Modrich, P.

(2010). PCNA function in the activation and strand direction of MutLα endonuclease in

mismatch repair. *Proc. Natl. Acad. Sci.* 107, 16066–16071.

doi:10.1073/PNAS.1010662107.

Rees, H. A., Yeh, W.-H., and Liu, D. R. (2019). Development of hRad51–Cas9 nickase

fusions that mediate HDR without double-stranded breaks. *Nat. Commun. 2019 101* 10,

1–12. doi:10.1038/s41467-019-09983-4.

Richardson, C. D., Kazane, K. R., Feng, S. J., Zelin, E., Bray, N. L., Schäfer, A. J., et al.

(2018). CRISPR–Cas9 genome editing in human cells occurs via the Fanconi anemia

pathway. *Nat. Genet.* doi:10.1038/s41588-018-0174-0.

Stojic, L., Brun, R., and Jiricny, J. (2004). Mismatch repair and DNA damage signalling. *DNA*

Repair (Amst). 3, 1091–1101. doi:10.1016/J.DNAREP.2004.06.006.

Umar, A., Koi, M., Risinger, J. I., Glaab, W. E., Tindall, K. R., Kolodner, R. D., et al. (1997).

Correction of Hypermutability, N-Methyl-N′-nitro-N-nitrosoguanidine Resistance, and

Defective DNA Mismatch Repair by Introducing Chromosome 2 into Human Tumor

Cells with Mutations in MSH2 and MSH6. *Cancer Res.* 57.

Wang, X., Kam, Z., Carlton, P. M., Xu, L., Sedat, J. W., and Blackburn, E. H. (2008). Rapid

telomere motions in live human cells analyzed by highly time-resolved microscopy.

Epigenetics Chromatin 2008 11 1, 1–19. doi:10.1186/1756-8935-1-4.

Winter, G. E., Buckley, D. L., Paulk, J., Roberts, J. M., Souza, A., Dhe-Paganon, S., et al.

(2015). Phthalimide conjugation as a strategy for in vivo target protein degradation.

Science (80-.). 348, 1376–1381. doi:10.1126/SCIENCE.AAB1433.

Yeh, C. D., Richardson, C. D., and Corn, J. E. (2019). Advances in genome editing through

control of DNA repair pathways. *Nat. Cell Biol.* 21(12), 1468–1478.

doi:10.1038/s41556-019-0425-z.

Data availability

All sequencing data will be deposited in the European Nucleotide Archive (EMBL-EBI).

Competing Interests

The authors declare no commercial or financial relationships that could be construed as

potential conflict of interest.

Author Contributions

526 JFdaS, GO, JJ and JIL conceptualized the study. JFdaS and JIL obtained funding. JFdaS, GO, EA and AM carried out investigations. GT performed the confocal microscopy. JFdaS performed analysis and visualization. JIL supervised the study. JFsaS with input from JIL wrote the original draft and all authors reviewed and edited the final manuscript.

Funding

532 JFdaS is supported by a DOC fellowship from the Austrian Academy of Sciences (ÖAW25035). A.M. is funded by the Austrian Science Fund, grant number P 33024 awarded to J.I.L. The Loizou lab is funded by an ERC Synergy Grant (DDREAMM Grant agreement ID: 855741). CeMM is funded by the Austrian Academy of Sciences.

Acknowledgments

538 We would like to thank the VBCF (Vienna, Austria) for all next generation sequencing. We would like to acknowledge Dr Jacob Corn (ETHZ, Zurich, Switzerland) and Dr Richard Sherwood (Harvard Medical School, Boston, USA) for critically reading the manuscript. We would like to thank Mr Chris Fell (Medical University of Vienna, Vienna, Austria) and Mr Marc Wiedner (CeMM, Vienna, Austria) for technical support. We are thankful to Prof KJ Patel (Weatherall Institute of Molecular Medicine, Oxford, UK) for providing the FANCD2 deficient HAP1 cell line. The Winter Lab (CeMM, Vienna, Austria) provided valuable tools for protein degradation. We are grateful to members of the Loizou lab for helpful discussions and feedback.