Greater spear nosed bats commute long distances alone, rest together, but forage apart

M. Teague O'Mara[1-4], Dina K.N. Dechmann[3-6]

1. Bat Conservation International
2. Department of Biological Sciences, Southeastern Louisiana University
3. Department of Migration, Max Planck Institute of Animal Behavior
4. Smithsonian Tropical Research Institute
5. Department of Biology, University of Konstanz
6. Centre for the Advanced Study of Collective Behaviour, University of Konstanz

Abstract

Animals frequently forage in groups on ephemeral resources to profit from social information and increase efficiency. Greater spear-nosed bats (*Phyllostomus hastatus*) develop group-specific social calls, which are hypothesized to coordinate social foraging to feed on patchily-distributed balsa flowers. We tracked three social groups of *P. hastatus* on Isla Colón, Panamá, using high frequency GPS. We found commuting distances of 20-30 km to foraging sites, more than double of what has been previously reported. In contrast to our expectations, we found that individuals commuted alone. They then joined group members in small foraging patches with high densities of flowering balsas on the mainland. Close proximity to group members did not influence foraging efficiency or energy expenditure, but it positively influenced time outside of the cave, and increased the duration and synchrony of rest. These results show that stronger social proximity associations were more closely tied to resting behaviour, and indicate that factors other than increased feeding efficiency may structure social relationships of group members while foraging. It appears that depending on the local resource landscape these bats have an excellent map even of distant resources and may use social information only for current patch discovery. They then may no longer rely on social information during daily foraging.

Keywords: GPS tracking, movement ecology, energy expenditure, foraging, social behaviour

Introduction

Animals must respond to changes in the spatiotemporal distribution of resources, and their movement decisions to search for and exploit food resources directly impact the ability to satisfy dietary requirements and affects their fitness (Bell, 1990). When resources are ephemeral (e.g., spatially patchy, temporally unpredictable), using social information while moving with others may help them find resources more efficiently (Bhattacharya & Vicsek, 2014). For example, fish track moving refugia by matching speed to group mates (Berdahl, Torney, Ioannou, Faria, & Couzin, 2013), insect- and fish-eating bats converge on the feeding calls of conspecifics (Dechmann et al., 2009; Egert-Berg et al., 2018), seabirds follow the white plumage of foraging flocks (Beauchamp & Heeb, 2001), and penguins are able to capture more fish when foraging together (McInnes, McGeorge, Ginsberg, Pichegru, & Pistorius, 2017).

Foraging in groups can convey energetic benefits by increasing foraging success and make energy intake more reliable (Giraldeau & Beauchamp, 1999; McInnes et al., 2017; Snijders et al., 2021). Individuals are required to maintain cohesion and spatiotemporal coordination to benefit from interactions with conspecifics (Conradt & Roper, 2005). While maintaining strong social bonds can provide long-term fitness benefits (Bohn, Moss, & Wilkinson, 2009; Silk et al., 2010), moving with group members can increase immediate costs of transport, and feeding competition. Less time and energy spent on finding food patches due to information provided by group members may be especially important for species foraging on ephemeral resources, but little evidence is available.

Bats are an excellent group to test how resource ephemerality directs group foraging. They spend large proportions of their energy budget on locomotion that is fueled by the food of the day (O'Mara et al., 2017), and feed on resources that are often widely dispersed and unpredictable. Bat species that forage for ephemeral insect swarms eavesdrop on the

echolocation buzzes that reveal a capture attempt (Dechmann et al., 2009; Dechmann,

Kranstauber, Gibbs, & Wikelski, 2010; Egert-Berg et al., 2018). Beyond this often

opportunistic behaviour, some species will also search food near group members to

maximize the discovery of feeding patches patches (Dechmann et al., 2009; Dechmann et

al., 2010; Egert-Berg et al., 2018). Bats of many species readily incorporate social

information about food across a range of cues and (O'Mara, Dechmann, & Page, 2014;

Page & Ryan, 2006; Ramakers, Dechmann, Page, & O'Mara, 2016; Ratcliffe & ter Hofstede,

2005; Wright, 2016) and the nature of the resource they feed on as well as how tightly they

depend on it can be used to predict if and when social information should be used during

foraging (Kohles, O'Mara, & Dechmann, 2022).

Greater spear-nosed bats (*Phyllostomus hastatus*) seasonally feed on an ephemeral

resource, the nectar of balsa trees (*Ochroma pyramidale*). In a well-studied population on

Trinidad, female *P. hastatus* form stable life-long groups of unrelated females (McCracken &

Bradbury, 1981; Wilkinson, Carter, Bohn, & Adams, 2016). They synchronize reproduction,

converge on a group-specific call that requires extended learning, and perform several

cooperative behaviours at the group level, such as babysitting and pup guarding from the

infanticide attempts of neighbouring groups (Boughman & Wilkinson, 1998; Boughman,

1998; Wilkinson & Boughman, 1998; Wilkinson et al., 2016). They are omnivorous, but

during the dry season, they feed nearly exclusively on balsa nectar. These pioneer trees are

a rare and patchy resource and the group-specific social calls are hypothesized to recruit

group members to flowering trees to exploit or defend them collectively (Wilkinson &

Boughman, 1998). However, the number of flowers available on a given tree is limited

(Kays, Rodríguez, Valencia, & Horan…, 2012), many other animals feed on them, and the

energy requirements of these bats are large (Kunz, Robson, & Nagy, 1998). Thus, the

potential reasons for recruiting others to these flowers warrant further investigation. In

addition, the availability of balsa and thus the value of social information (Kohles et al.,

2022) may vary locally and it is unclear if group foraging and resource defence occur across

the species' range.

We used high frequency GPS loggers to track three groups of *P. hastatus* in Panamá and

recorded complete foraging trips. We used these GPS data to construct proximity-based

social networks to test how social associations influence foraging performance and

behaviour. We hypothesized that, like in Trinidad, *P. hastatus* forage in groups during the

dry season. We thus expected them to commute to a food source and exploit it together.

We also hypothesized that social foraging increases foraging success, and that social

proximity increases foraging efficiency despite potential competition trade-offs. With this

study we make an important contribution to how foraging behaviour may vary

intraspecifically, and thus the intricate link between a local resource landscape and the

resulting social behaviour.

Methods

Data were derived from 40 adult *Phyllostomus hastatus* (38 F / 2 M) that were captured

from three roosting groups in a cave ("La Gruta") on Isla Colón, Bocas del Toro, Panamá

using a bucket trap. Groups were captured sequentially and there was no overlap among

groups in the nights they were tracked. Roosting groups were individuals co-roosting in

small depressions on the cave ceiling, consistent with previous work in Trinidad. Both

males were adult harem males, and the females comprised 15 nulliparous young females,

and 19 postlactating females. Bats from a single social group were placed into a wire mesh

cage covered with a breathable cotton cloth where they roosted together calmly until

removed for processing. Bat mass was recorded to the nearest 0.5 g, forearm length

measured to the nearest 0.1 mm, and each bat received a subcutaneous PIT tag (ID100;

Euro ID, Frechen, Germany). To measure wing dimensions for flight power estimates, we

took photos of one fully outstretched wing placed flat over mm graph paper. A subset of bats from each group were fitted with a GPS data logger (Gypsy-5 GPS, TechnoSmart, Rome, Italy (O'Mara et al., 2021)) that was wrapped in clear shrink tube. The logger was mounted on a silk collar (0.8 cm wide) and closed with Safil-C degradable suture (Aesculap/B. Braun, Co, Tuttlingen, Germany, (O'Mara, Wikelski, & Dechmann, 2014)). Total collar + GPS weight was 6.8 ± 0.51 g, which represented 5.7 ± 0.4 % of body mass (range: 4.5 - 6.6 %).

GPS tags collected location fixes from 18 h – 06 h local time at 0.5 or 1 Hz. When there was not adequate GPS reception, tags went into a low energy sleep state for five mins and then restarted to search for satellites for 90 s. Tag function varied due to the deep cave roost used by the bats and resting under presumably dense foliage while foraging. We retrieved 18 tags with analyzable data: five females and one male from group one, three females from group two, and eight females and one male from group three (Table S1). The 18 recovered tags collected one to four nights of data for a total of 34 bat nights. We removed from analysis five nights from five different bats where fewer than 30 mins were tracked for various reasons (e.g., the bat remained in the cave for most of the night draining the battery), leaving 30 bat nights from 16 females and 2 males with a range of 75.5 – 307.5 min of data collected per night (mean ± sd: 197.31 ± 60.35, Table S1).

To estimate flight airspeed and subsequent energy expenditure, wind data were collected at an automated weather station (9.351, -82.258) at 15-min intervals by the Physical Monitoring Program at the Smithsonian Tropical Research Institute for their Bocas del Toro field station and downloaded from http://biogeodb.stri.si.edu/physical_monitoring/research/bocas. Wind speed and direction were collected every 10 s with a RM Young Wind Monitor Model 05103. Mean wind speed and wind direction were then calculated at the end of every 15-min interval.

141   **Analysis**

142   All analyses were conducted in R 4.02 (R Core Team, 2020).
143

Ground speed, wind speed, and wind accommodation.

Ground speed (speed of movement relative to the ground) and bearing were calculated for

successive time points in the *move* package (Kranstauber, Smolla, & Scharf, 2018). To

calculate airspeed (speed of movement relative to the moving air column), wind support

and crosswind for were annotated for each GPS location using a weighted interpolation of

the U and V components of wind to match the resolution of the GPS sampling

sampling(O'Mara et al., 2021; O'Mara et al., 2019; Safi et al., 2013). Wind support was

calculated as the length of the wind vector in the direction of the bat's flight where positive

values represent tailwind and negative values headwind and are given as total support in m

s^{-1}. Crosswind was calculated as the absolute value of the speed of the wind vector

perpendicular to the travel direction, and airspeed was calculated as the square root of

[(ground speed – wind support)^2 + crosswind^2].

Behavioural segmentation

To identify behavioural states of resting, slow foraging flight (i.e., feeding), moving between

patches, and commuting, we applied a four state hidden Markov model in *momentuHMM*

(McClintock, Michelot, & Goslee, 2018)*.* These behaviours were entered into the model in

order of increasing speed and decreasing turning angular mean (i.e., slow flight had larger

turning angles, commuting flight was fast with high concentrated turning angles near zero),

with step lengths modelled with a gamma error distribution and turning angles with a

wrapped Cauchy distribution. Models were fit for each bat on each bat night, and each

resulting model was visually inspected to ensure reasonable classification. The track for

each bat night was first regularized to one-second intervals using a correlated random walk

procedure in function *momentuHMM::crawlWrap* and then passed to the hidden Markov

model. We used simulations to target the number of identifiable states and identify the

starting values for each state (McClintock et al., 2018; Michelot et al., 2017), and these

simulations showed that four state model always performed better (had lower AIC values)

than three- or two-state models. On occasion, a five-state model had better fit although it

was often difficult to discern biological meaning between the additional state that was

placed very close to the low speed and higher turning angle behaviours of roosting and

slow foraging flight.

We used a patch approach to identify foraging and resting (i.e., night roost) areas since

aggregations of GPS locations should indicate a site of behavioural interest. For each bat

night a patch was defined as a cluster of GPS locations that were classified as foraging

(slow flight or moving) or as rest. These clusters were identified using density-based spatial

clustering of applications with noise (DBSCAN) using function *fpc::dbscan* (Hennig, 2020;

Schubert, Sander, Ester, Kriegel, & Xu, 2017) with a minimum of 15 points per cluster at a

maximum spatial distance (eps) of 10 m among nearest neighbours. This distance was

chosen through visual inspection of diagnostic plots in function *dbscan*::*kNNdistplot*

(Hahsler, Piekenbrock, & Doran, 2019). To facilitate spatial comparisons across all

individuals in the sequentially tracked groups, we labelled patches with centroids that were

less than 30 m apart as a single patch regardless of the night on which they were used.

This distance was chosen as there was a clear break in the distribution of pairwise

distances among patch centroids and 30 m is slightly larger than the approximate diameter

of a balsa crown. These patches were also ground-truthed to evaluate potential plant food

composition and presence.

Once foraging patches were identified, we then further classified feeding locations that

were likely flower clusters. We used the same DBSCAN procedure on the GPS locations

within each foraging patch per bat night, to identify a flower cluster as a position with a

minimum of six points per cluster within a maximum spatial distance of 0.8 m. This distance

was chosen based on the spatial distribution of flowers observed within *O. pyramidale*

(personal observation). We used these likely feeding clusters to define foraging efficiency as

feeding clusters divided by the total time tracked (in mins) per night.

Energy expenditure

The speed that an animal flies in an air column (airspeed) is the most important predictor of

its mechanical power output and subsequent total metabolic power output. We estimated

the mechanical power of flight in Watts (P_{mech}) following Pennycuick (2008) using calculated

airspeeds, the capture mass of the animals, and wing length taken from each bat's wing

photo multiplied by two. An individual's average power curve was generated for each bat at

the mean flight altitude (50 m) and 25°C (Figure S1) and returned estimates across all

individuals for the minimum power speed of 6.81 ± 0.21 ms^{-1} and maximum range speed of

11.0 ± 0.35 ms^{-1}. Minimum power speed represents the most efficient instantaneous flight

speed and maximum range speed maximizes the range covered over ground per unit

energy expended. In general, bats should fly at their minimum power speed when moving

short distances and at the maximum range speed when moving long distances

(Hedenström, 2003). To estimate energy expenditure for each night, the instantaneous

power output during flight was calculated for each GPS location at the observed altitude

and airspeed. Mechanical power output alone underestimates metabolic power

requirements (Pennycuick, 2008; von Busse, Swartz, & Voigt, 2013; Ward et al., 2001)   . To

estimate total metabolic power required, we estimated the metabolic power of flight (P_{met})

following Ward et al (2001) using the mean estimated flight muscle partial efficiency (E_{FM}) for

P. hastatus in a wind tunnel (0.24667, range: 0.13 – 0.34, (Thomas, 1975)). Total metabolic

power was then calculated as: $P_{met} = 1.1[(P_{mech} / E_{FM}) + P_{BMR}]$. For locations where the bat

was at rest (including while in the cave), we substituted the resting metabolic rate (P_{BMR}) of

23.8 J g^{-1} h^{-1} (McNab, 1969) and converted this to 0.0661 W g^{-1}. These values were

summed to daily energy expenditure (DEE) and compared to DEE values from smaller *P.*

hastatus in Trinidad measured through doubly labelled water (Kunz et al., 1998).

To estimate energy returns from foraging, we used the energy content per ml of balsa nectar

and the likely flower density per tree (Kays et al., 2012). Total nectar produced by a flower is

estimated at 25.5 ml, and balsa nectar sugar concentration decreases over the night from

13.3% at 18 h to 7.9% at 06 h, with an average concentration of 12.4% total sugars (Kays

et al., 2012). This is 0.124 g sucrose ml^{-1} nectar that yields 0.47988 kcal ml^{-1} (3.87 kcal g^{-1}

sugar *0.124 g ml^{-1}). Balsa nectar then has an energy density of 2.007818 kJ ml^{-1}. Flowers

open with 4.9 ± 1.3 ml of nectar, and there is a sharp decline in nectar production over the

night. We assumed that bats drink the full nectar volume present when flowers open (5 ml),

which likely over-estimates the amount of nectar truly ingested during feeding events. Peak

flower availability is approximately 60 flowers per patch or 5 flowers per m^2. However, the

mean is 20 flowers in a patch and 2.5 flowers per m^2, with flower density following a normal

distribution over the season (Kays et al., 2012).

Social Proximity Effects on Behaviour

To test the effects of distance to a nearest neighbour on behaviour we used the distance

among individuals as a dynamic metric that could change with every second of tracking.

We excluded commuting behaviour from this analysis after inspection of the data showed

that individuals did not commute together with group mates. We used this pairwise

distance to further identify if the changing proximity between individuals affects movement

decisions. We limited the potential distance that behaviour could be affected by another

individual to 500 m, which is twice the potential perceptual distance of *P. hastatus* social

calls at a peak call frequency of 6725 ± 36.3 Hz at ambient weather conditions (Stilz &

Schnitzler, 2012).

Generalized linear mixed effects models were fit in *lme4* with individual as a random

intercept nested within social group (Bates, Mächler, Bolker, & Walker, 2015). For models

evaluating proportional activity budgets as a response, a binomial family was specified and

for all others Gaussian models were used. When a non-linear relationship seemed likely, we

fit second-order polynomial models and tested if they fit the data more efficiently than a

first order model using the second-order Akaike Information Criterion calculated using

MuMIn::AICc. The most efficient model was the model with an AICc value at least three

units lower than competing models. To evaluate the significance of the fixed effects, we

calculated type II p-values using Satterthwaite degrees of freedom method with

lmerTest::anova for Gaussian models (Kuznetsova, Brockhoff, & Christensen, 2017) and

with *car::Anova* for binomial models. To measure the effect size of each Gaussian model, R^2

was calculated for both the marginal (fixed effects, R^2_m) and conditional (fixed and random

effects, R^2_c) in *MuMIn* (Bartoń, 2016). The full GPS data set and code are available from the

Movebank Data Repository (doi released on acceptance, code and data with environmental

and behavioural annotation is available at osf.io for review:

https://osf.io/d8gaz/?view_only=324382133cfb44c28ada3120b49ea42e).

266   Results

Tracking Summary

The 18 tracked bats mostly foraged in sites that were 20 – 30 km away from their cave

roost across the sea (Fig 1), however one of the harem males foraged close to the cave and

most individuals showed some indications of quick foraging stops on their return flights to

the cave. While GPS tags had the same programmed on/off time, because of low satellite

coverage and late cave emergence, the first GPS record of each tag was 149 ± 49 min after

sunset. At this time bats were already commuting and 5.2 ± 6.2 km from the roost when

first locations were recorded (range: 350 m – 24.2 km). Each night, bats spent 197 ± 60

minutes outside the roost and travelled 59.2 ± 16.2 km. Bats commuted to their foraging

areas with slight headwinds at ground speeds of 8.63 ± 2.63 m s^{-1} (airspeed: 9.12 ± 2.69 m

s^{-1}) and returned to the cave at 7.89 ± 4.08 m s^{-1} (airspeed: 7.76 ± 3.82 m s^{-1}) flying with

tailwinds, airspeeds that are between their minimum power speed (6.81 ± 0.21 m s^{-1}) and

maximum range speed (11.0 ± 0.35 m s^{-1}). Wind speeds were generally low during the

tracking period, with prevailing offshore winds blowing eastward. Bats foraged at ground

speeds of 3.95 ± 3.43 m s^{-1} (airspeed: 4.15 ± 3.38 m s^{-1}).

Figure 1. A) Tracking overview with individuals colored by group membership. B) The island

in the Changuinola river (Isla Changuinola) shown with three individuals from group 3 on

2016-03-04, and C) the pairwise distances between these three individuals across that

night. Note that in C both axes are repeated to show the simultaneous distance between

pairs of individuals and the large distances among individuals at the beginning and end of

the nights when commuting.

Behavior & foraging patch use

We ground-truthed the foraging patches on the mainland and found flowering *O.*

pyramidale trees in each of them. There were no flowering *O. pyramidale* on Isla Colón, but

foraging patches always included flowering *Luehea seemannii*. It is unknown if bats fed on

the nectar of *L. seemannii* flowers, or on the animals attracted to this resource. All

individuals completed the ca. 25 km commute from the roost to the foraging areas alone

(Fig 1C), and individuals were 9.1 ± 5.8 km (mean ± sd) away from one another when

commuting. Individuals then converged in the same foraging areas, mostly on the mainland.

Individuals used a mix of slow flight, moving and resting during the approximately 200 mins

they were outside of the cave, and this did not differ across the three groups (Fig 2A). Bats

spent 24.0 ± 19.3% of their time in rest, 24.6 ± 14.6% of time in slow foraging/feeding

flights, 31.3 ± 23.6% in faster foraging movements between feeding sites, and 25.1 ±

16.8% of time commuting. To examine social effects on foraging, we only further analysed

behaviours other than commuting.

Bats used 7.2 ± 4.2 foraging patches (i.e., trees or groups of trees) per night, and patches

were 4.0 ± 3.8 km apart. These patches were 23.35 ± 1.4 km from the roost, and only a few

foraging patches were used near the roost during the return flight to the cave. As bats

increased their total nightly flight distance, they used more patches (estimate: 0.886 ±

0.034 per additional km, $F_{1,\ 20.174}$ = 6.861, p = 0.016, R^2_m = 0.13, R^2_c = 0.70; Fig S2). Over the

course of the night, bats used 20.72 ± 19.91 feeding locations (flower clusters) (range: 3 -

82). However, there was no relationship between the number of feeding locations they

visited within the patches and the number of patches used, or with the time they spent

outside the roost (Fig S2).

Figure 2. Bat nightly activity budgets and the synchronization of behaviour differ when nearest neighbours are in the same patch or a different patch. A) Bat activity budgets when nearest neighbours occupied the same and different patches, but were within 500 m of one another. B) Behavioral synchrony of nearest neighbours in and out of the same patch. Box plots show proportion of time engaged in behaviors for each patch category, and points show the proportions for each individual bat per night.

Activity budgets differed depending on whether individuals were in the same patch as their neighbours or not (Fig 2A-B, patch estimate $F_{1,3}$ = 0.029; Fig 2). When in the same patch, individuals were 24.3 ± 21.4 m apart (0.25 – 133 m). The proportion of time spent in rest was larger when an individual's nearest neighbour was in the same patch as their nearest neighbour than when individuals occupied different patches that were within 500 m of one another (same patch (median ± MAD): 85.2 ± 18.6%, different patch: 32.2% ± 44.3%, X^2_3 = 10.12, p = 0.018, Fig 2A). The proportion of each behaviour that was synchronized with a nearest neighbour was not evenly distributed (X^2_1 = 4.08, p = 0.043, Fig 2B). Individuals were more likely to synchronize resting than other behaviours when in the same patch as their nearest neighbour (Fig 2B). Bats rested 19.3 ± 14.3 m away from each other, and resting bouts were longer when they were in closer proximity to another individual, with the

strongest effects less than three meters away (power curve / Freundlich equation intercept

= 2.95 ± 0.24 m SE, p < 0.001, Fig 3).

Figure 3. Duration of continuous resting bouts relative to the distance to a bat's nearest

neighbour.

Despite groups being tracked on different nights, the same resting areas tended to be used

by bats across all social groups, regardless of the tracking night. We identified 43 resting

areas, and one of these resting locations on Isla Changuinola was used by 11 bats over five

different nights (Fig S3). Three other locations on Isla Colón were used repeatedly by a

single bat over two nights. The remaining 39 sites were used by one bat on one night each.

Energy costs & feeding requirements

The mean minimum power speed of these bats was 6.81 ± 0.21 m s^{-1} and maximum range

speed was 11.03 ± 0.34 m s^{-1}, with the airspeeds used by bats (7 – 9 m s^{-1}) falling well

within these estimates for energy-efficient flight. The power curve for each individual bat

was used to estimate total metabolic power required for the night. Estimated daily energy

expenditure (DEE) was 198.47 ± 69.44 kJ day^{-1} and increased with tracking time (estimate:

0.729 ± 0.174, $F_{1,25.61} = 17.462$, $p < 0.001$, $R^2_m = 0.383$, $R^2_c = 0.448$, Fig 4A). This is similar to

DEE estimates based on allometric estimates from body mass alone of 153.02 ± 5.96 kJ

day^{-1} (range: $143.07 - 164.89$ (Speakman, 2005)), and to DEE derived from doubly-labelled

water measurements in *P. hastatus* that were 36% smaller than those at our study site (76.8

± 6.6 g vs 121.5 ± 7.6 g in this study, estimates from Kunz et al (1998) are shown in blue in

Fig 4A). We converted the estimated energetic needs for each individual on each tracking

night into the number of full *O. pyramidale* flowers required to support that estimated

energy expenditure. We found that individuals fed from an increasing number of flower

clusters as their total estimated energy needs increased (second order polynomial estimate

$= 55.45 \pm 14.89$, 27.73 ± 14.86, $F_{2,26.45} = 8.84$, $p = 0.001$, $R^2_m = 0.376$, $R^2_c = 0.494$, Fig 4B).

Estimated energy expenditure, as measured through movement, more strongly depended

on the total time tracked per night than on the proximity relationships among individuals in

a social group.

Figure 4. Estimated daily energy expenditure and the estimated energetic returns from
foraging. A) Daily energy expenditure and the minutes tracked per night, with comparative
data from smaller *P. hastatus* in Trinidad shown in blue (Kunz et al., 1998). The line shows
the relationship derived from the individuals tracked in this study. B) The number of flowers
needed, based on average energy content, to support the energy requirements of an
individual's estimated energy expenditure predict the number of feeding clusters used per
night.

Discussion

Based on previous work (Wilkinson & Boughman, 1998), we predicted that *Phyllostomus*
hastatus would commute to foraging patches together and forage with their social group to
feed on *O. pyramidale* flowers. Instead, we found that individuals commuted alone to the
foraging patches over long distances that included the ocean and large commercial banana
plantations – landscapes that have few available resources to these bats. Tracked
individuals then used the same foraging patches, but on different feeding locations within
the patch. When individuals were near group members, they tended to rest, and individuals
that rested close to one another tended to rest for longer. It appears that while the main
benefit of social foraging is often assumed to be increased foraging success, resting in

these *P. hastatus* groups may reinforce social bonds or has benefits outside of foraging

efficiency such as predator detection.

Social foraging should increase foraging efficiency, either because food patches are

detected more efficiently or because of social facilitation increasing feeding rates. With

increasing group size, Trinidad guppies decrease the time it takes to locate food patches

and increase their intake rates (Snijders et al., 2021). Larger guppy groups act as more

efficient sensors, but this comes at a cost of perceived feeding competition that drives

increased bite rates. Individuals of many bat species forage socially to eavesdrop on

feeding calls when resources are ephemeral and searching costs are high (Egert-Berg et al.,

2018; Fenton, 2003), and some species show extraordinarily coordinated group foraging for

these ephemeral resources (Dechmann et al., 2009; Dechmann et al., 2010; Kohles et al.,

2022). The time spent foraging, the number of foraging patches or the number of feeding

locations used were independent of the proximity of group members.

Astonishingly, we found that individual *P. hastatus* from multiple social groups used the

same patches of *O. pyramidale*, but that they commuted over 25 km from their roost

without other tagged group members. *Phyllostomus hastatus* were previously found to

forage within 10 km of their roost (McCracken & Bradbury, 1981; Williams & Williams,

1970). Displacement studies showed that most *P. hastatus* individuals would successfully

navigate back to their home roost after displacement of up to 20 km. At 30 km or more,

bats often failed completely to return home (Williams & Williams, 1970). This indicates that

the distance at which our study animals were foraging from their cave was at the edge or

even outside the range they were familiar with. They had to cross the sea and large

commercial banana farms to reach the foraging locations. All of our tracked bats foraged in

mainland areas with *O. pyramidale*. During our resource ground-truthing we found that on

their home island and elsewhere on the mainland *O. pyramidale* was not yet available. This

suggests that once an isolated *O. pyramidale* patch is discovered, especially early in the

flowering season, social information about this quickly moves through a colony. Once they

have located it, group members commute to the patch alone. *Phyllostomus hastatus* feeds

on various food sources throughout most of the year, but almost exclusively switches to *O.*

pyramidale when it is available during the dry season (Wilkinson & Boughman, 1998). It is

unknown if this is due to a shortage of other food sources, or to a distinct preference for

nectar. However, locating this widely distributed food source may rely on social information

transfer (Kohles et al., 2022). This may be a rare, but crucial resource during the transition

period when few trees are flowering, and finding food is more unpredictable. Further

tracking, mapping of the resource landscape, and detailed dietary analysis are needed to

identify how and when bats switch between food resources and the behavioural and

energetic correlates of this change.

While they commuted alone, we found that some individuals would reunite with social

group members in these patches, while others tended to forage completely on their own.

This is similar to vampire bats (*Desmodus rotundus*) that fly individually to cattle

approximately 300 m away where they then feed with partners they have close social

relationships with (Ripperger & Carter, 2021). Previous radio tracking of *P. hastatus* in

Trinidad found that during most of the year, social group members foraged alone and did

not depart or arrive at the cave together, but on rare occasions they joined a group mate in

a nearby foraging area (McCracken & Bradbury, 1981). Members of social groups occupied

adjacent foraging ranges and social groups were segregated across the landscape

(McCracken & Bradbury, 1981), unlike the large overlap we found in our data. In Trinidad

groups were more likely to depart together, and females from the same social group were

captured around a *O. pyramidale* feeding site more often than randomly expected during

the dry season (Wilkinson & Boughman, 1998). There also appears to be strong social

attraction among Trinidad *P. hastatus* groups. Social calls broadcast at flowering *O.*

pyramidale attracted bats, and social calls from flying *P. hastatus* were most often noted 3-

4 hours after sunset during foraging (Wilkinson & Boughman, 1998). The higher spatial and

temporal resolution of our GPS tracks now indicates they are at least not only attracted to

the possibility of food resources, but that they may form resting associations while outside

of their roost.

There could be regional or population differences in the main drivers on social group

formations depending on the resource landscape. In Trinidad it has been hypothesized that

recruitment of group members to flowering trees may predominantly help bats defend trees

against competitors (Wilkinson & Boughman, 1998). However, in Panamá large animals that

bats cannot defend against, such as kinkajous and opossums are the main visitors of *O.*

pyramidale flowers (Kays et al., 2012). A flowering *O. pyramidale* with mean peak flower

availability of 60 flowers per night provides approximately 600 kJ of energy at the beginning

of the night. This is before nectar pools are depleted and trees begin producing less

energy-dense nectar (Kays et al., 2012). We estimated that our tracked bats expended 198

± 69 kJ / day, indicating that a single *O. pyramidale* crown could support the daily energy

needs of only 3 - 7 bats. Such a limited resource may be worth defending from

conspecifics if all available flowers could be fully exploited (Wilkinson & Boughman, 1998),

but would not support the needs of an entire social group of bats, suggesting that a

collective resource defence is not the likely explanation for foraging near group mates for

this population in Panamá.

It is possible that at this time of year or within this population in Panamá there is more

pressure to maintain social associations than to group forage. Within social groups that are

structured by kin relationship, strong social bonds between related individuals have

numerous life history advantages (Silk, 2007), extending lifespans (Barocas, Ilany, Koren,

Kam, & Geffen, 2011; Silk et al., 2010), success in group conflicts (Samuni, Crockford, &

Wittig, 2021), and individual reproductive success (Frere et al., 2010; Schülke, Bhagavatula,

Vigilant, & Ostner, 2010; Silk, Alberts, & Altmann, 2003). On Trinidad, however, female *P.*

hastatus form stable, relatively closed groups of unrelated females that stay together for

their lifetime and show highly developed social bonds (McCracken & Bradbury, 1981;

Wilkinson et al., 2016). They develop group-specific social calls (Boughman, 1997;

Wilkinson & Boughman, 1998), and recognize and guard group-members' offspring (Bohn

et al., 2009; Bohn, Wilkinson, & Moss, 2007). Low reproductive rates and high infant

mortality in this species (Stern & Kunz, 1998) are strong selective pressures on potential

cooperation among unrelated females. The ecological and physiological conditions

structuring *P. hastatus* social groups may be similar to groups of unrelated females in wild

equids (Cameron, Setsaas, & Linklater, 2009) and cooperative breeding birds (Riehl &

Strong, 2018). Close rest while foraging, rather than feeding together, may be one way for

the bats to reinforce their social bonds since *P. hastatus* has not been observed to engage

in the extended grooming that characterizes social bond maintenance in other mammals

(Kern & Radford, 2021; Silk et al., 2003; Wilkinson, 1986), or to potentially guard against

predation.

Social information may mainly be used for the discovery of feeding areas. Commuting to

repeatedly used foraging patches 25 km or more from a central roost may not be unusual

for some bats (Calderón-Capote et al., 2020; Goldshtein et al., 2020; Harten, Katz,

Goldshtein, Handel, & Yovel, 2020; O'Mara et al., 2021), but still presents navigational and

energetic risks due to the long distances travelled and potential weather hazards. Nectar

and fruit-feeding fruit bats appear to have large and robust cognitive maps of their (Harten

et al., 2020; Toledo et al., 2020). These species will all commute from a common roost to

distant foraging patches, but there is little to no overlap among individual foraging ranges in

animals from these roosting aggregations, and these non-overlapping ranges may be

developed through reinforcement learning that minimizes competition (Goldshtein et al.,

2020).

Further work mapping resources as seasons change and social groups reach decision

points to alter movements will help us further elucidate the intricate relationships between

social and foraging behaviour and its energetic context. This may have strong variation

among populations that differ in the resource landscapes they encounter, but may allow us

to better understand the links between species ecological niche and how sociality responds

to resource environments and the need for information use.

Data statement

The full GPS data set and code are available from the Movebank Data Repository (doi

released on acceptance, code and data with environmental and behavioural annotation is

available at osf.io for review:

https://osf.io/d8gaz/?view_only=324382133cfb44c28ada3120b49ea42e).

Acknowledgements

We would like to thank the owners of Cueva La Gruta for access to the research site.

Maurice Thomas, and the Smithsonian Tropical Research Institute assisted with logistics

and support, particularly Rachel Page, Rachel Collin, Plinio Gondola, and Urania González.

We would also like to thank Gary McCracken, Camila Calderón, Jenna Kohles, Anne

Scharf, Mariëlle van Toor, and Kamran Safi for suggestions and insight, and Martin Wikelski

and the Max Planck Institute of Animal Behavior (formerly Ornithology) for exceptional

support. This work was approved by the Ministerio del Ambiente, Panamá (SE/A-96-15),

and the Animal Care and Use Committee at the Smithsonian Tropical Research Institute

(2014-0701-2017).

Funding

This was funded by funded in part by the University of Konstanz Young Scholars Fund to

MTO, the Max Planck Institute of Ornithology, and the US National Science Foundation

(BIO-2217920 to MTO).

References

Barocas, A., Ilany, A., Koren, L., Kam, M., & Geffen, E. (2011). Variance in centrality within rock hyrax social networks predicts adult longevity. *PLoS One, 6*(7), e22375. doi: 10.1371/journal.pone.0022375

Bartoń, K. (2016). MuMIn: Multi-Model Inference. Retrieved from https://CRAN.R-project.org/package=MuMIn

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw, 67*(1), 1-48. doi: 10.18637/jss.v067.i01

Beauchamp, G., & Heeb, P. (2001). Social foraging and the evolution of white plumage. *Evol Ecol Res, 3*(6), 703-720.

Bell, W. J. (1990). *Searching Behaviour: The behavioural ecology of finding resources*. London: Chapman and Hall.

Berdahl, A., Torney, C., Ioannou, C., Faria, J., & Couzin, I. (2013). Emergent sensing of complex environments by mobile animal groups. *Science, 339*(6119), 574-576. doi: 10.1126/science.1225883

Bhattacharya, K., & Vicsek, T. (2014). Collective foraging in heterogeneous landscapes. *J R Soc Interface, 11*(100), 20140674. doi: 10.1098/rsif.2014.0674

Bohn, K. M., Moss, C. F., & Wilkinson, G. S. (2009). Pup guarding by greater spear-nosed bats. *Behav Ecol Sociobiol, 63*(12), 1693-1703. doi: 10.1007/s00265-009-0776-8

Bohn, K. M., Wilkinson, G. S., & Moss, C. F. (2007). Discrimination of infant isolation calls by female greater spear-nosed bats, *Phyllostomus hastatus*. *Anim Behav, 73*(Part 3), 423-432. doi: 10.1016/j.anbehav.2006.09.003

Boughman, J., & Wilkinson, G. (1998). Greater spear-nosed bats discriminate group mates by vocalizations. *Anim Behav, 55*(6), 1717-1732. doi: 10.1006/anbe.1997.0721

Boughman, J. W. (1997). Greater spear-nosed bats give group-distinctive calls. *Behav Ecol Sociobiol, 40*(1), 61-70. doi: 10.1007/s002650050316

Boughman, J. W. (1998). Vocal learning by greater spear-nosed bats. *Proc R Soc B, 265*(1392), 227-233. doi: 10.1098/rspb.1998.0286

Calderón-Capote, M. C., Dechmann, D. K. N., Fahr, J., Wikelski, M., Kays, R., & O'Mara, M. T. (2020). Foraging movements are density-independent among straw-coloured fruit bats. *R Soc Open Sci, 7*(5). doi: 10.1098/rsos.200274

Cameron, E. Z., Setsaas, T. H., & Linklater, W. L. (2009). Social bonds between unrelated females increase reproductive success in feral horses. *Proc Natl Acad Sci U S A, 106*(33), 13850-13853. doi: 10.1073/pnas.0900639106

Conradt, L., & Roper, T. J. (2005). Consensus decision making in animals. *Trends Ecol Evol, 20*(8), 449-456.

Dechmann, D., Heucke, S., Giuggioli, L., Safi, K., Voigt, C., & Wikelski, M. (2009). Experimental evidence for group hunting via eavesdropping in echolocating bats. *Proc R Soc B, 276*(1668), 2721-2728. doi: 10.1098/rspb.2009.0473

Dechmann, D., Kranstauber, B., Gibbs, D., & Wikelski, M. (2010). Group hunting-a reason for sociality in molossid bats? *PLoS One, 5*(2), e9012. doi: 10.1371/journal.pone.0009012

Egert-Berg, K., Hurme, E. R., Greif, S., Goldstein, A., Harten, L., Herrera, M. L., . . . Yovel, Y. (2018). Resource ephemerality drives social foraging in bats. *Curr Biol, 28*(22), 3667-3673. doi: 10.1016/j.cub.2018.09.064

Fenton, M. B. (2003). Eavesdropping on the echolocation and social calls of bats. *Mammal Rev, 33*(3-4), 193-204. doi: 10.1046/j.1365-2907.2003.00019.x

Frere, C. H., Krutzen, M., Mann, J., Connor, R. C., Bejder, L., & Sherwin, W. B. (2010). Social and genetic interactions drive fitness variation in a free-living dolphin population. *Proc Natl Acad Sci USA, 107*(46), 19949-19954. doi: 10.1073/pnas.1007997107

Giraldeau, L., & Beauchamp, G. (1999). Food exploitation: searching for the optimal joining policy. *Trends Ecol Evol, 14*(3), 102-106.

Goldshtein, A., Handel, M., Eitan, O., Bonstein, A., Shaler, T., Collet, S., . . . Yovel, Y.
(2020). Reinforcement learning enables resource partitioning in foraging bats. *Curr*
Biol. doi: 10.1016/j.cub.2020.07.079
Hahsler, M., Piekenbrock, M., & Doran, D. (2019). dbscan: Fast Density-Based Clustering
with R. *J Stat Softw, 91*(1). doi: 10.18637/jss.v091.i01
Harten, L., Katz, A., Goldshtein, A., Handel, M., & Yovel, Y. (2020). The ontogeny of a
mammalian cognitive map in the real world. *Science, 369*(6500), 194-197. doi:
10.1126/science.aay3354
Hedenström, A. (2003). Twenty-Three Testable Predictions About Bird Flight. In P. Berthold,
E. Gwinner & E. Sonnenschein (Eds.), *Avian Migration* (pp. 563-582). Berlin:
Springer.
Hennig, c. (2020). fpc: Flexible Procedures for Clustering (Version R package version 2.2-5).
Retrieved from https://CRAN.R-project.org/package=fpc\
Kays, R., Rodríguez, M., Valencia, L., & Horan…, R. (2012). Animal visitation and pollination
of flowering balsa trees (*Ochroma pyramidale*) in Panama. *Mesoamericana, 16*, 54-
70.
Kern, J. M., & Radford, A. N. (2021). Strongly bonded individuals prefer to forage together
in cooperatively breeding dwarf mongoose groups. *Behav Ecol Sociobiol, 75*(5). doi:
10.1007/s00265-021-03025-0
Kohles, J. E., O'Mara, M. T., & Dechmann, D. K. N. (2022). A conceptual framework to
predict social information use based on food ephemerality and individual resource
requirements. *Biol Rev Camb Philos Soc*. doi: 10.1111/brv.12881
Kranstauber, B., Smolla, M., & Scharf, A. K. (2018). move: Visualizing and analyzing animal
track data.
Kunz, T. H., Robson, S. K., & Nagy, K. A. (1998). Economy of harem maintenance in the
greater spear-nosed bat, *Phyllostomus hastatus*. *J Mammal*, 631-642.
Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in
Linear Mixed Effects Models. *J Stat Softw, 82*(13). doi: 10.18637/jss.v082.i13
McClintock, B. T., Michelot, T., & Goslee, S. (2018). momentuHMM: R package for
generalized hidden Markov models of animal movement. *Methods Ecol Evol, 9*(6),
1518-1530. doi: 10.1111/2041-210x.12995
McCracken, G. F., & Bradbury, J. W. (1981). Social organization and kinship in the
polygynous bat *Phyllostomus hastatus*. *Behav Ecol Sociobiol, 8*(1), 11-34. doi:
10.1007/Bf00302840
McInnes, A. M., McGeorge, C., Ginsberg, S., Pichegru, L., & Pistorius, P. A. (2017). Group
foraging increases foraging efficiency in a piscivorous diver, the African penguin. *R*
Soc Open Sci, 4(9), 170918. doi: 10.1098/rsos.170918
McNab, B. K. (1969). The economics of temperature regulation in neotropical bats. *Comp*
Biochem Physiol, 31(2), 227-268. doi: 10.1016/0010-406X(69)91651-X
Michelot, T., Langrock, R., Bestley, S., Jonsen, I. D., Photopoulou, T., & Patterson, T. A.
(2017). Estimation and simulation of foraging trips in land-based marine predators.
Ecol, 98(7), 1932-1944. doi: 10.1002/ecy.1880
O'Mara, M., Dechmann, D., & Page, R. (2014). Frugivorous bats evaluate the quality of
social information when choosing novel foods. *Behav Ecol, 25*, 1233-1239. doi:
10.1093/beheco/aru120
O'Mara, M. T., Amorim, F., Scacco, M., McCracken, G. F., Safi, K., Mata, V., . . .
Dechmann, D. K. N. (2021). Bats use topography and nocturnal updrafts to fly high
and fast. *Curr Biol, 31*(6), 1311-1316 e1314. doi: 10.1016/j.cub.2020.12.042
O'Mara, M. T., Scharf, A. K., Fahr, J., Abedi-Lartey, M., Wikelski, M., Dechmann, D. K. N., &
Safi, K. (2019). Overall dynamic body acceleration in straw-colored fruit bats
increases in headwinds but not with airspeed. *Front Ecol Evol, 7*, 200. doi:
10.3389/fevo.2019.00200

O'Mara, M. T., Wikelski, M., & Dechmann, D. K. N. (2014). 50 years of bat tracking: device
attachment and future directions. *Methods Ecol Evol, 5*(4), 311-319. doi:
10.1111/2041-210x.12172
O'Mara, M. T., Wikelski, M., Voigt, C. C., Ter Maat, A., Pollock, H. S., Burness, G., . . .
Dechmann, D. K. (2017). Cyclic bouts of extreme bradycardia counteract the high
metabolism of frugivorous bats. *eLife, 6*, e26686. doi: 10.7554/eLife.26686
Page, R., & Ryan, M. (2006). Social transmission of novel foraging behavior in bats: frog
calls and their referents. *Curr Biol, 16*(12), 1201-1205. doi:
10.1016/j.cub.2006.04.038
Pennycuick, C. J. (2008). *Modelling the flying bird* (Vol. 5). London: Academic Press.
R Core Team. (2020). R: A language and environment for statistical computing: R
Foundation for Statistical Computing, Vienna Austria. Retrieved from http://www.R-
project.org
Ramakers, J. J. C., Dechmann, D. K. N., Page, R. A., & O'Mara, M. T. (2016). Frugivorous
bats prefer information from novel social partners. *Anim Behav, 116*, 83-87. doi:
10.1016/j.anbehav.2016.03.021
Ratcliffe, J., & ter Hofstede, H. (2005). Roosts as information centres: social learning of food
preferences in bats. *Biol Lett, 1*(1), 72-74. doi: 10.1098/rsbl.2004.0252
Riehl, C., & Strong, M. J. (2018). Stable social relationships between unrelated females
increase individual fitness in a cooperative bird. *Proc R Soc B, 285*(1876). doi:
10.1098/rspb.2018.0130
Ripperger, S. P., & Carter, G. G. (2021). Social foraging in vampire bats is predicted by
long-term cooperative relationships. *PLoS Biol, 19*(9), e3001366. doi:
10.1371/journal.pbio.3001366
Safi, K., Kranstauber, B., Weinzierl, R., Griffin, L., Rees, E. C., Cabot, D., . . . Bohrer, G.
(2013). Flying with the wind: scale dependency of speed and direction
measurements in modelling wind support in avian flight. *Mov Ecol, 1*(1), 4. doi:
10.1186/2051-3933-1-4
Samuni, L., Crockford, C., & Wittig, R. M. (2021). Group-level cooperation in chimpanzees
is shaped by strong social ties. *Nat Commun, 12*(1), 539. doi: 10.1038/s41467-020-
20709-9
Schubert, E., Sander, J., Ester, M., Kriegel, H. P., & Xu, X. (2017). DBSCAN Revisited,
Revisited. *ACM Transactions on Database Systems, 42*(3), 1-21. doi:
10.1145/3068335
Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance
reproductive success in male macaques. *Curr Biol, 20*(24), 2207-2210. doi:
10.1016/j.cub.2010.10.058
Silk, J. B. (2007). The adaptive value of sociality in mammalian groups. *Philos Trans R Soc
B, 362*(1480), 539-559. doi: 10.1098/rstb.2006.1994
Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance
infant survival. *Science, 302*(5648), 1231-1234.
Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., . . .
Cheney, D. L. (2010). Strong and consistent social bonds enhance the longevity of
female baboons. *Curr Biol, 20*(15), 1359-1361. doi: 10.1016/j.cub.2010.05.067
Snijders, L., Krause, S., Tump, A. N., Breuker, M., Ortiz, C., Rizzi, S., . . . Kurvers, R. (2021).
Causal evidence for the adaptive benefits of social foraging in the wild. *Commun
Biol, 4*(1), 94. doi: 10.1038/s42003-020-01597-7
Speakman, J. (2005). Body size, energy metabolism and lifespan. *J Exp Biol, 208*(Pt 9),
1717-1730. doi: 10.1242/jeb.01556
Stern, A. A., & Kunz, T. H. (1998). Intraspecific variation in postnatal growth in the greater
spear-nosed bat. *J Mammal, 79*(3), 755. doi: 10.2307/1383086
Stilz, W. P., & Schnitzler, H. U. (2012). Estimation of the acoustic range of bat echolocation
for extended targets. *J Acoust Soc Am, 132*(3), 1765-1775. doi: 10.1121/1.4733537

Thomas, S. P. (1975). Metabolism during flight in two species of bats, *Phyllostomus*
hastatus and *Pteropus gouldii*. *J Exp Biol, 63*(1), 273-293.
Toledo, S., Shohami, D., Schiffner, I., Lourie, E., Orchan, Y., Bartan, Y., & Nathan, R. (2020).
Cognitive map–based navigation in wild bats revealed by a new high-throughput
tracking system. *Science, 369*(6500), 188-193. doi: 10.1126/science.aax6904
von Busse, R., Swartz, S. M., & Voigt, C. C. (2013). Flight metabolism in relation to speed in
Chiroptera: testing the U-shape paradigm in the short-tailed fruit bat *Carollia*
perspicillata. *J Exp Biol, 216*(Pt 11), 2073-2080. doi: 10.1242/jeb.081760
Ward, S., Moller, U., Rayner, J. M., Jackson, D. M., Bilo, D., Nachtigall, W., & Speakman, J.
R. (2001). Metabolic power, mechanical power and efficiency during wind tunnel
flight by the European starling *Sturnus vulgaris*. *J Exp Biol, 204*(Pt 19), 3311-3322.
Wilkinson, G., & Boughman, J. (1998). Social calls coordinate foraging in greater spear-
nosed bats. *Anim Behav, 55*, 337-350.
Wilkinson, G. S. (1986). Social grooming in the common vampire bat, *Desmodus rotundus*.
Anim Behav, 34(6), 1880-1889.
Wilkinson, G. S., Carter, G. G., Bohn, K. M., & Adams, D. M. (2016). Non-kin cooperation in
bats. *Philos Trans R Soc B, 371*(1687). doi: 10.1098/rstb.2015.0095
Williams, T. C., & Williams, J. M. (1970). Radio tracking of homing and feeding flights of a
neotropical bat, *Phyllostomus hastatus*. *Anim Behav, 18*, 302-309. doi:
10.1016/s0003-3472(70)80042-2
Wright, G. S. (2016). Social Learning and Information Transfer in Bats: Conspecific
Influence Regarding Roosts, Calls, and Food. In J. Ortega (Ed.), *Sociality in Bats* (pp.
211-230): Springer.
706

 Supplementary Material

Table S1. Biometric and tracking information for the individuals analyzed in this study. Reproductive state is coded as nr: non-reproductive, plac: post-lactation, and nulli: nulliparous. Nightly values for the minutes of tracking and number of locations used are given for means ± sd when more than one night was recorded.

Bat ID	Group ID	Sex	Reproductive State	Mass (g)	Forearm length (mm)	Wing length (mm)	Wing area (mm²)	Nights Analyzed	Time Analyzed (min)	Number of Locations Used
2016030703	one	m	nr	134	93.5	271.96	22966.52	2	133.99 ± 82.68	7296.5 ± 3939.29
2016030705	one	f	plac	124	87.8	280.82	24543.47	1	156.67	7694
71A0D95	one	f	nulli	122	89.9	282.74	24088.56	2	224.89 ± 4.82	12371.5 ± 1289.06
71A111A	one	f	nulli	109	90.8	296.86	25587.39	1	279.52	16752
74DDA80	one	f	nulli	123	91.2	278.44	23724.58	1	151.88	6268
74DE9A7	one	f	plac	122	91.5	274.95	23752.23	1	236.13	13488
74D8954	two	f	plac	128	93.02	317.17	29422.44	1	161.88	6211
74DA92F	two	f	plac	109	87.6	250.07	19667.46	2	192.87 ± 25.53	6767 ± 1236.02
74DCA83	two	f	nulli	123	92.1	300.31	29703.42	2	131.85 ± 51.45	5028 ± 2224.56
74D8C25	three	f	nulli	131	93.2	308.25	28074.65	1	118.15	5158
74D932E	three	f	nulli	116	93.6	304.29	28397.6	2	207.63 ± 2.4	8783 ± 83.44
74DAF9C	three	f	nulli	126	94.5	296.29	26317.99	1	95.27	5628
74DCBCC	three	f	plac	114	89.7	289.26	22936.64	2	278.75 ± 18.27	11910 ± 1578.26
74DE4E7	three	f	plac	124	94.2	296.11	26383.96	1	197.03	6618
74F7D4C	three	m	nr	137	93.2	277.12	23586.97	1	152.63	3748
74F8E19	three	f	plac	113	91.7	289.38	25321.02	3	201.26 ± 61.87	7702.67 ± 2597.55
74F9F83	three	f	plac	116	92	284.51	24720.16	3	264.17 ± 54.4	7866.33 ± 1690.4
74FE24E	three	f	plac	113	91.7	270.5	22057.56	2	218.35 ± 3.15	7528 ± 622.25

Figure S1. Mechanical power requirements at airspeeds for each bat in this study. Power

output was calculated following Pennycuick (2008) at 50 m and 25°C using each

individual's body mass and wing dimensions.

Figure S2. Individual nightly summaries for foraging and feeding. The number of foraging

patches used is predicted by A) the distance that individuals moved each night and B) the

total time that individuals were tracked each night. The number of feeding clusters

identified was not predicted by either C) the distance moved per night or D) the number of

foraging patches used. Statistically significant slopes that differed from zero are shown by

solid lines, non-significant slopes by dashed lines, and shaded areas indicate 95%

confidence intervals.

Figure S3. Resting patches clustered from GPS locations. A) Resting patches are shown

across the study area, scaled by the number of individual bat days that the sites were used

and colored by the number of individual bats that used the patch. Resting patches are

shown for B) Isla Changuinola, and C) Isla Colón.