Lay Summary
Social foraging can increase foraging efficiency on hard-to-find food sources. Miniaturized

high-frequency GPS allowed us to test this on groups of greater spear-nosed bats. We

estimated food intake rates, energy expenditure, and the strength of bat social

relationships. In contrast to our expectations, we did not find a relationship between bat

social relationships and energy expenditure or foraging efficiency. However, bats took

longer resting breaks when group members were close. Why these bats commute 25 km

individually across the ocean to then forage near each other thus requires further

investigation.

10
Greater spear nosed bats in Panamá do not use social proximity to improve foraging
efficiency

Abstract

Group foraging animals may change their spatial relationships in response to the resource

landscape. The strength of social network ties across foraging groups could then reflect the

energy balance of individual group members through their effect on foraging efficiency.

Greater spear-nosed bats (*Phyllostomus hastatus*) live in social groups where unrelated

females develop group-specific social calls, defend each other's offspring, and are

hypothesized to coordinate social foraging to feed on balsa flowers. To test if social groups

forage together, and if this increases foraging efficiency, we tracked 18 individuals from

three social groups of *P. hastatus* on Isla Colón, Panamá using high frequency GPS for 1 –

3 days each. We found that individuals commuted alone but then joined group members in

small foraging patches with high densities of flowering balsas, 25 km from their roost on the

mainland. Proximity-based social network measures did not predict foraging efficiency or

energy expenditure, but individuals with higher social associations spent more time outside

of the cave. Individuals did not coordinate movement, but close proximity among

individuals increased resting bout duration and synchrony, particularly when in the same

patch. These results show that stronger social proximity associations were more closely

tied to resting behavior, and indicate that factors other than increased feeding efficiency

may structure network relationships of group members while foraging.

Keywords: GPS tracking, movement ecology, energy expenditure, foraging, social behavior

Introduction

Animals must respond to changes in the spatiotemporal distribution of resources, and their

movement decisions to search for and exploit food resources directly impacts the ability to

satisfy dietary requirements and affects their fitness (Bell, 1990). When resources are

ephemeral (e.g spatially patchy, temporally unpredictable), using social information while

moving with others may help them find resources more efficiently (Bhattacharya and

Vicsek, 2014; Valone, 1989). For example, fish track moving refugia by matching speed to

group mates (Berdahl et al., 2013), insect- and fish-eating bats converge on the feeding

calls of conspecifics (Dechmann et al., 2009; Egert-Berg et al., 2018), seabirds follow the

white plumage of foraging flocks (Beauchamp and Heeb, 2001), and penguins are able to

capture more fish when foraging together (McInnes et al., 2017).

Foraging in groups can convey energetic benefits by increasing foraging success and make

energy intake more (Giraldeau and Beauchamp, 1999; McInnes et al., 2017; Snijders et al.,

2021). To benefit from interactions with conspecifics, individuals are required to maintain a

degree of cohesion and spatial and temporal coordination in their activities (Conradt and

Roper, 2005). While maintaining strong social bonds can provide long-term fitness benefits

(Bohn et al., 2009; Silk et al., 2003), moving with group members can increase immediate

costs of transport, and feeding competition. However, few studies have directly addressed

the energetic trade-offs of social foraging or changes in the variance in food intake (Snijders

et al., 2021). Less time and energy spent on finding food patches due to information

provided by group members may be especially important for species foraging on

ephemeral resources, but little evidence is available.

Bats are an excellent group to test how resource ephemerality directs group foraging. They spend large proportions of their energy budget on locomotion that is fueled by the food of the day (O'Mara et al., 2017), and feed on resources that are often widely dispersed and unpredictable. Bat species that forage for ephemeral insect swarms eavesdrop on the echolocation buzzes that reveal where an individual is attempting to capture (Dechmann et al., 2009; Egert-Berg et al., 2018; Fenton, 2003). Beyond this potentially opportunistic attraction to a resource, social group members of some species will also forage near each other to maximize the discovery of these feeding patches (Dechmann et al., 2009; Dechmann et al., 2010). Bats of many species readily incorporate social information about food across a range of cues and modalities (O'Mara et al., 2014a; Page and Ryan, 2006; Ramakers et al., 2016; Ratcliffe and ter Hofstede, 2005; Wright, 2016) reinforcing the inherently social nature of their foraging ecology.

To test the potential energetic benefits of foraging together, we tracked social groups of greater spear-nosed bats (*Phyllostomus hastatus*) in Panamá. Due to bat nocturnal lifestyles, acoustic cues are necessary for the coordination of group movement, and since sound has a relatively short transmission distance, bat social groups should move close to one another in a coordinated fashion. In a well-studied population on Trinidad, female *P. hastatus* form stable closed groups of unrelated females that stay together for their lifetime (McCracken and Bradbury, 1981; Wilkinson et al., 2016). Females in these social groups synchronize their reproduction, converge on a group-specific call that requires extended learning, and perform several cooperative behaviors at the group level, such as babysitting and pup guarding from the infanticide attempts of neighboring groups (Boughman and Wilkinson, 1998; Boughman, 1998; Boughman, 2006; Wilkinson and Boughman, 1998; Wilkinson et al., 2016). These omnivorous bats have diverse diets, but during the dry season, they feed nearly exclusively on the nectar of balsa trees (*Ochroma pyramidale*).

These pioneer trees are a rare and patchy resource and females are hypothesized to use

group-specific social calls to recruit group members to flowering trees and exploit or

defend them collectively (Wilkinson and Boughman, 1998). However, the number of flowers

available on these trees is limited at any given time (Kays et al., 2012), and the energy

requirements of these bats are large (Kunz et al., 1998) which may increase competition. In

addition, the availability of balsa may vary locally and it is unclear if group foraging and

resource defense occurs across the species' range.

We used high frequency GPS loggers to track three groups of *P. hastatus* in Panamá and

recorded complete foraging trips. We used these GPS data to construct proximity-based

social networks to test how social associations influence foraging performance and

behavior. Based on the previous literature we hypothesized that (H1) *P. hastatus* forage in

groups during the dry season and (H2) group foraging increases foraging efficiency despite

competition trade-offs. We predicted that (P1) individuals with stronger proximity networks

forage in the same food patches, and (P2) network strength centrality is positively

associated with foraging efficiency and negatively associated with nightly energy

expenditure. However, we found no clear evidence for social foraging or on energy

expenditure. *Phyllostomus hastatus* at our study site in Panamá seem to behave profoundly

differently from what was described previously.

Methods

We analyzed data of females from three social groups: five females and one male from

group one, three females from group two, and eight females and one male from group three

(Table S1). The 18 bats collected one to three nights of data, and we analyzed 30 complete

bat nights (with a range of 75.5 – 307.5 min of data collected per night (mean ± sd: 197.31

± 60.35). Groups were captured and tracked sequentially, with no overlap in tracking

among groups.

These data were derived from 40 adult *Phyllostomus hastatus* (38 F / 2 M) that were

captured from three social groups in a cave ("La Gruta") on Isla Colón, Bocas del Toro,

Panamá using a bucket trap. Social groups were defined as individuals co-roosting in small

depressions on the cave ceiling, consistent with previous work in Trinidad. Both males were

adult harem males, and the females comprised 15 nulliparous young females, and 19

postlactating females. Bats from a single social group were placed into a wire mesh cage

covered with a breathable cotton cloth where they roosted together calmly until removed

for processing. Bat mass was recorded to the nearest 0.5 g, forearm length measured to

the nearest 0.1 mm, and each bat received a subcutaneous PIT tag (ID100; Euro ID,

Frechen, Germany). To measure wing dimensions for flight power estimates, wing photos

were taken of one fully outstretched wing placed flat over mm graph paper. A subset of

bats from each group were fitted with a GPS data logger (37 tags, Gypsy-5 GPS,

TechnoSmArt, Rome, Italy (O'Mara et al., 2021)) that was wrapped in clear shrink tube. The

logger was mounted on a silk collar (0.8 cm wide) and closed with Safil-C degradable

suture (Aesculap/B. Braun, Co, Tuttlingen, Germany, (O'Mara et al., 2014b)). Total collar +

GPS weight was 6.8 ± 0.51 g, which represented 5.7 ± 0.4 % of body mass (range: 4.5 -

6.6 %).

GPS tags were programmed to collect location fixes from 18 h – 06 h local time at 0.5 or 1

Hz. When adequate GPS reception was not received, tags went into a low energy sleep

state for five minutes and then restarted to search for satellites for 90 seconds. The GPS

tags had variable functionality due to both the deep cave roost used by the bats and

roosting under presumably dense foliage while foraging. The 18 recovered tags collected

one to four nights of data for a total of 34 bat nights. We removed from analysis five nights

of tracking from five different bats that were less than 30 mins for various reasons (e.g., the

bat remained in the cave for most of the night draining the battery), leaving 30 bat nights

from 16 females and 2 males.

Wind data were collected at an automated weather station (9.351, -82.258) at 15-min

intervals by the Physical Monitoring Program at the Smithsonian Tropical Research Institute

for their Bocas del Toro field station and downloaded from

http://biogeodb.stri.si.edu/physical_monitoring/research/bocas. Wind speed and direction

were collected every 10 seconds with a RM Young Wind Monitor Model 05103. Mean wind

speed and wind direction were then calculated at the end of every 15-minute interval.

Analysis

All analyses were conducted in R 4.02 (R Core Team, 2020).
145

Ground speed, wind speed, and wind accommodation.

Ground speed (speed of movement relative to the ground) and bearing were calculated for

successive time points in the *move* package (Kranstauber et al., 2012). To calculate

airspeed (speed of movement relative to the moving air column) wind support and

crosswind for were annotated for each GPS location using a weighted interpolation of the U

and V components of wind to match the resolution of the GPS sampling (O'Mara et al.,

2021; O'Mara et al., 2019; Safi et al., 2013). Wind support was calculated as the length of

the wind vector in the direction of the bat's flight where positive values represent tailwind

and negative values headwind and are given as total support in ms^{-1}. Crosswind was

calculated as the absolute value of the speed of the wind vector perpendicular to the travel

direction, and airspeed was calculated as the square root of [(ground speed – wind

support)^2 + crosswind^2].

Behavioral segmentation

To identify behavioral states of resting, slow foraging flight (or feeding), moving between

patches, and commuting, we applied a four state hidden Markov model in *momentuHMM*

(McClintock et al., 2018). These behaviors were entered into the model in order of

increasing speed and decreasing turning angular mean (i.e., slow flight had larger turning

angles, commuting flight was fast with high concentrated turning angles near zero), with

step lengths modeled with a gamma error distribution and turning angles with a wrapped

Cauchy distribution. Models were fit for each bat on each bat night, and each resulting

model was visually inspected to ensure reasonable classification. The track for each bat

night was regularized to one-second intervals using a correlated random walk procedure in

function *momentuHMM::crawlWrap* and then passed to the hidden Markov model. We used

simulations to target the number of identifiable states and identify the starting values for

each state (McClintock et al., 2018; Michelot et al., 2017), and these simulations showed

that four state model always performed better (had lower AIC values) than three or two

state models. On occasion a five-state model had better fit although it was often difficult to

discern biological meaning between the additional state that was placed very close to the

low speed and higher turning angle behaviors of roosting and slow foraging flight.

We used a patch approach to identify foraging and resting (night roost) areas since

aggregations of GPS locations should indicate a site of behavioral interest. For each bat

night a patch was defined as a cluster of GPS locations that were classified as foraging

(slow flight or moving) or as rest. These clusters were identified using density-based spatial

clustering of applications with noise (DBSCAN) using function *fpc::dbscan* (Hennig, 2020;

Schubert et al., 2017) with a minimum of 15 points per cluster at a maximum spatial

distance (eps) of 10 m among nearest neighbors. This distance was chosen through visual

inspection of diagnostic plots in function *dbscan::kNNdistplot* (Hahsler et al., 2019). To

facilitate spatial comparisons across all individuals in the sequentially tracked groups, we

labeled patches with centroids that were less than 30 m apart as a single patch regardless

of the night on which it was used. This distance was chosen as there was a clear break in

the distribution of pairwise distances among patch centroids and 30 m is slightly larger than

the approximate diameter of a balsa crown. These patches were also ground-truthed to

evaluate potential food composition and presence.

Once foraging patches were identified, we then further classified feeding locations that

were likely flower clusters. We used the same DBSCAN procedure on the GPS locations

within each foraging patch per bat night, to identify a flower cluster as a position with a

minimum of six points per cluster within a maximum spatial distance of 0.8 m. This distance

was chosen based on the spatial distribution of flowers observed within *O. pyrimidale*. We

used these likely feeding clusters to define foraging efficiency as feeding clusters divided

by the total time tracked (in minutes) per night.

Energy expenditure

The speed that an animal flies in an air column (airspeed) is the most important predictor of

its mechanical power output. We estimated the mechanical power of flight in Watts (P_{mech})

following Pennycuick (2008) using calculated airspeeds, the capture mass of the animals,

and wing length taken from a photo of the bat's fully outstretched wing over graph paper

multiplied by two. An individual's average power curve was generated for each bat at the

mean flight altitude (50 m) and 25°C (Figure S1) and returned estimates across all

individuals for the minimum power speed of 6.81 ± 0.21 m s^{-1} and maximum range speed of

11.0 ± 0.35 m s^{-1}. Minimum power speed represents the most efficient instantaneous flight

speed and maximum range speed maximizes the range covered over ground per unit

energy expended. In general, bats should fly at their minimum power speed when moving

short distances and at the maximum range speed when moving long distances

(Hedenström, 2003). To estimate energy expenditure for each night, the instantaneous

power output during flight was calculated for each GPS location at the observed altitude

and airspeed. Mechanical power output alone underestimates metabolic power

requirements (Pennycuick, 2008; von Busse et al., 2013; Ward et al., 2001). To estimate

total metabolic power required, we estimated the metabolic power of flight (P_{met}) following

Ward et al (2001) using the mean estimated flight muscle partial efficiency (E_{FM}) for *P.*

hastatus in a wind tunnel (0.24667, range: 0.13 – 0.34, (Thomas, 1975)). Total metabolic

power was then calculated as: $P_{met} = 1.1[(P_{mech} / E_{FM}) + P_{BMR}]$. For locations where the bat

was at rest (including while in the cave), we substituted the resting metabolic rate (P_{BMR}) of

23.8 J g^{-1} h^{-1} (McNab, 1969) and converted this to 0.0661 W g^{-1}. These values were

summed to daily energy expenditure (DEE) and compared to DEE values from smaller *P.*

hastatus in Trinidad measured through doubly labeled water (Kunz et al., 1998).

To estimate energetic returns from foraging, we used the energy content per ml of balsa

nectar and the likely flower density per tree (Kays et al., 2012). Total nectar produced by a

flower is estimated at 25.5 ml, and balsa nectar sugar concentration decreases over the

night from 13.3% at 18 h to 7.9 % at 06 h, with an average concentration of 12.4% total

sugars. This is 0.124 g sucrose per ml nectar that yields 0.47988 kcal per ml (3.87 kcal/g

sugar *0.124 g ml^{-1}). Balsa nectar then has an energy density of 2.007818 kJ ml^{-1}. Flowers

open with 4.9 ± 1.3 ml of nectar, and there is a sharp decline in nectar production over the

night. We assumed that bats drink the full nectar volume present when flowers open (5 ml).

Peak flower availability is approximately 60 flowers per patch or 5 flowers per m^2. However,

the mean is 20 flowers per patch and 2.5 flowers per m^2, with flower density following a

normal distribution over the season.

Social Proximity Effects on Behavior

To test the effects of distance to a nearest neighbor on behavior we used the distance

among individuals as a dynamic metric that could change with every second of tracking.

We also used this distance and if the focal individual's nearest neighbor was in the same

patch to further identify if the changing proximity between individuals affects movement

decisions. We limited the potential distance that behavior could be affected by another

individual to 500 m, which is twice the potential perceptual distance of *P. hastatus* calls at a

peak call frequency of 6725 ± 36.3 Hz at ambient weather conditions (Stilz and Schnitzler,

2012).

To test if cumulative proximity relationships influence behavior, we constructed individual-

level proximity-based social networks across all tracking nights in *spatsoc* (Robitaille et al.,

2019). GPS fixes were grouped to one-minute intervals, and proximity was assessed as a

binary variable at a distance threshold of less than 30 m. This is twice the spatial error of a

stationary tag (14.22 ± 12.05 m (O'Mara et al., 2021)), although error decreases dramatically

when in flight (Ranacher et al., 2016), and is slightly larger than the approximate diameter of

a balsa crown. We then calculated single values for strength centrality and degree for each

individuals in *asnipe* (Farine, 2013), with edge weights as a simple ratio index of individuals

within 30 m of each other. We used strength centrality of each individual in subsequent

analyses as it reflects how much the focal bat was within 30 m of other individuals (Wyman

et al., 2021), as well as degree (the number of different individuals that came within 30 m,

Supplemental Table 1). We also wanted to identify if there were particular spatial locations

that were used by bats across the social groups since groups were tracked sequentially

and did not overlap in time. To do so we created an undirected patch use network with

edge weights as the simple ratio index of the presence or absence of the use of a foraging

patch by an individual. We used fast-greedy community detection in *igraph* to identify if

there were foraging patches that were commonly used among individuals from multiple

social groups at different times (Clauset et al., 2004). These communities then show how

spatial location can connect individuals and groups across time.

Generalized linear mixed effects models were fit in *lme4* with individual as a random

intercept nested within group (Bates et al., 2015). For models evaluating on proportional

activity budgets as a response, a binomial family was specified and for all others Gaussian

models were used. When a non-linear relationship seemed likely, we fit second order

polynomial models and tested if they fit the data more efficiently than a first order model

using the second-order Akaike Information Criterion calculated using *MuMln::AICc.* The

most efficient model was the model with an AICc value at least 3 units lower than

competing models. To evaluate the significance of the fixed effects in the models, we

calculated type II p-values using Satterthwaite degrees of freedom method with

lmerTest::anova for gaussian models (Kuznetsova et al., 2017) and with *car::Anova* for

binomial models. To measure the effect size of each Gaussian model, R^2 was calculated for

both the marginal (fixed effects, R^2_m) and conditional (fixed and random effects, R^2_c) in

MuMln (Bartoń, 2016). The full GPS data set and code are available from the Movebank

Data Repository (doi released on acceptance, code and data with environmental and

behavioral annotation is available at Dryad for review:

https://datadryad.org/stash/share/DBJdu_DVmEVBp6Yeq5x2RwupCy6W9NOaprjDtUZIH3E or at

https://github.com/teague0/hastatusForaging).

Results

Tracking Summary

The 18 tracked bats mostly foraged in sites that were 20 – 30 km away from their cave

roost across the sea (Fig 1). While GPS tags had the same programmed on / off time,

because of low satellite coverage and late cave emergence, the first GPS record of each

tag was 149 ± 49 min after sunset. At this time bats were already commuting and 5.2 ± 6.2

km from the roost when first locations were recorded (range: 350 m – 24.2 km). Each night,

bats spent 197 ± 60 minutes outside the roost and traveled 59.2 ± 16.2 km. Bats

commuted to their foraging areas with slight headwinds at ground speeds of 8.63 ± 2.63 m

s^{-1} (airspeed: 9.12 ± 2.69 m s^{-1}) and returned to the cave at 7.89 ± 4.08 m s^{-1} (airspeed:

7.76 ± 3.82 m s^{-1}) flying with tailwinds, airspeeds that are between their minimum power

speed (6.81 ± 0.21 m s⁻¹) and maximum range speed (11.0 ± 0.35 m s⁻¹). Wind speeds were

generally low during the tracking period, with prevailing offshore winds blowing eastward.

Bats foraged at ground speeds of 3.95 ± 3.43 m s⁻¹ (airspeed: 4.15 ± 3.38 m s⁻¹).

Figure 1. A) Tracking overview with individuals colored by group membership. B) The island

in the Changuinola river (Isla Changuinola) shown with three individuals from group 3 on

2016-03-04, and C) the pairwise distances between these three individuals across that

night. Note that in C both axes are repeated to show the simultaneous distance between

pairs of individuals and the large distances among individuals at the beginning and end of

the nights when commuting.

Behavior & foraging patch use

All individuals completed the 25 km commute from the roost to the foraging areas alone

(Fig 1C), and individuals were 9.1 ± 5.8 km (mean ± sd) away from one another when

commuting. Individuals then converged in the same foraging areas, mostly on the mainland.

We only analyzed behaviors other than commuting to look at social influences on them.

Individuals used a mix of moving and resting during the approximately 200 minutes they

were outside of the cave, and this did not differ across the three groups (Fig 2A). Bats spent

24.0 ± 19.3% of their time in rest, 24.6 ± 14.6% of time in slow foraging/feeding flights,

31.3 ± 23.6% in faster foraging movements between feeding sites, and 25.1 ± 16.8% of

time commuting. We ground-truthed the mainland foraging patches and found flowering *O.*

pyramidale trees in each of them. There were no flowering *O. pyramidale* on Isla Colón, but

foraging patches always included flowering *Luehea seemannii*. It is unknown if bats fed on

the nectar of *Luehea* flowers, or on the animals attracted to this resource.

Bats used 7.2 ± 4.2 foraging patches (i.e. trees or groups of trees) per night, and patches

were 4.0 ± 3.8 km apart. These patches were 23.35 ± 1.4 km from the roost, and only a few

foraging patches were used near the roost during the return flight to the cave. As bats

increased their total nightly flight distance, they used more patches (Estimate: 0.886 ±

0.034 per additional km, $F_{1,\,20.174}$ = 6.861, p = 0.016, R^2_m = 0.13, R^2_c = 0.70; Fig S2). Over the

course of the night, bats used 20.72 ± 19.91 feeding locations (flower clusters) (range: 3 -

82). However, there was no relationship between the number of feeding clusters they

visited within the patches and the number of patches used, or with the time they spent

outside the roost (Fig S2).

Figure 2. Effects of proximity on bat nightly activity budgets. A) Bat activity budgets when nearest neighbors occupied the same and different patches, but are within 500 m of one another. B) Behavioral synchrony of nearest neighbors in and out of the same patch. Box plots show proportion of time engaged in behaviors for each patch category, and points show the proportions for each individual bat per night.

Activity budgets differed depending on whether individuals were in the same patch as their neighbors or not (Fig 2A-B, Patch Estimate $F_{1,3} = 0.029$; Fig 2). When in the same patch, individuals were 24.3 ± 21.4 m apart ($0.25 – 133$ m). The proportion of time spent in rest was larger when an individual's nearest neighbor was were in the same patch as their nearest neighbor than when individuals occupied different patches that were within 500 m of one another (same patch (median $\pm$ MAD): $85.2 \pm 18.6\%$, different patch: $32.2\%\pm$ 44.3%, $X^2 = 10.12$, df = 3, p = 0.018, Fig 2A). The proportion within a behavior that was synchronized with a nearest neighbor was not evenly distributed ($X^2 = 4.08$, df = 1, p = 0.043, Fig 2B). Individuals were more likely to be simultaneously engaged in rest when in the same patch as their nearest neighbor that to be synchronized in other behaviors (Fig 2B). Bats rested 19.3 ± 14.3 m away from each other, and resting bouts were longer when they were in closer proximity to another individual, with the strongest effects less than three

meters away (Power Curve / Freundlich Equation intercept = 2.95 ± 0.24 m SE, p < 0.001,

Fig 3).

Figure 3. Duration of continuous resting bouts relative to the distance to a bat's nearest

neighbor.

The same resting areas tended to be used by bats across all social groups, regardless of

the tracking night. We identified 43 resting areas, and one of these resting locations on Isla

Changuinola was used by 11 bats over five different nights (Fig S3). Three other locations

on Isla Colón were used repeatedly by a single bat over two nights. The remaining 39 sites

were used by one bat on one night each.

Social effects on foraging and resting

Proximity networks of simultaneous GPS locations showed that entire social groups were

never in close contact while tracked. There were strongly differentiated spatial relationships

among some group members, and the strength centrality and degree of individuals within a

social group was variable among groups (Fig S4). The males tracked as part of this study

were never in close proximity to the females from their group, despite similar long-distance

commutes. While close proximity between individuals at rest increased the likelihood they

would continue resting, there were no overall effects of an individual's strength centrality on

overall nightly activity budgets (Fig S5). However, individual strength centrality was

positively related to the amount of time that bats were tracked outside of their roost per

night (Estimate: 274.58 ± 103.38, $F_{1, 15.62}$ = 7.054, p = 0.018, R^2_m = 0.220, R^2_c = 0.393, Fig

3A). There were no effects of strength centrality on foraging efficiency when evaluated

using a second order mixed effects model ($F_{2,16.38}$ = 0.678, p = 0.521, Fig 3B), or on

estimated daily energy expenditure ($F_{1,18.18}$ = 1.52, p = 0.245, Fig 3C).

Figure 3. Effects of individual strength centrality on nightly measures of A) duration tracked,

B) foraging efficiency, and C) estimated daily energy expenditure. Solid lines indicate

statistically significant relationships using a mixed effects model, and dashed lines show

relationships that were not significant. Shaded regions show the 95% confidence interval.

Because we tracked social groups sequentially, we were unable to assess if multiple

groups are found in the same sites at the same time. However, there was extensive spatial

overlap of individuals across the nights of tracking. Proximity networks of GPS locations

that were within 30 m of each other, regardless of time, show overlapping use of foraging

and resting locations. We detected three communities from the proximity network created

across all tracking nights. These communities link together bats that foraged in the same

patches, but at different times and highlights that the same foraging locations host many

social groups. Individuals who shared close space at the same time did not necessarily use

the same number and constellation of foraging patches. Some individuals did not share

space with any other tracked bats, either from their social group or from the other tracked

groups (Fig S4).

Energy costs & feeding requirements

The mean minimum power speed of these bats was 6.81 ± 0.21 m s^{-1} and maximum range

speed was 11.03 ± 0.34 m s^{-1}. The power curve for each individual bat was used to

estimate total metabolic power required for the night. Bat estimated daily energy

expenditure (DEE) was 198.47 ± 69.44 kJ per day and increased with tracking time

(Estimate: 0.729 ± 0.174, $F_{1,25.61} = 17.462$, $p < 0.001$, $R^2_m = 0.383$, $R^2_c = 0.448$, Fig 4A). This

is similar to DEE estimates based on body mass alone of 153.02 ± 5.96 kJ per day (range:

$143.07 - 164.89$ (Speakman, 2005)) and to doubly-labeled water measurements in *P.*

hastatus that were 36% smaller than those at our study site (76.8 ± 6.6 g vs 121.5 ± 7.6 g in

this study, estimates from (Kunz et al., 1998) are shown in blue in Fig 4A). We converted the

estimated energetic needs for each individual on each tracking night into the number of full

O. pyramidale flowers required to support that estimated energy expenditure. We found

that individuals fed from an increasing number of flower clusters as their total estimated

energy needs increased (second order polynomial estimate = 55.45 ± 14.89, 27.73 ± 14.86,

$F_{2,26.45} = 8.84$, $p = 0.001$, $R^2_m = 0.376$, $R^2_c = 0.494$, Fig 4B). Estimated energy expenditure, as

measured through movement, more strongly depended on the total time tracked per night

than on the proximity relationships among individuals in a social group.

Figure 4. Estimated daily energy expenditure and the estimated energetic returns from foraging. A) Daily energy expenditure and the minutes tracked per night, with comparative data from smaller *P. hastatus* in Trinidad as comparison (Kunz et al., 1998). The line shows the relationship derived from the individuals tracked in this study. B) The number of flowers needed based on average energy content to support the energy requirements of an individual's estimated energy expenditure predict the number of feeding clusters used per night.

Discussion

Based on previous work (Wilkinson and Boughman, 1998), we predicted that *Phyllostomus hastatus* would forage with their social group to feed on *O. pyramidale* flowers. While tracked individuals did use the same foraging areas, most individuals flew to these foraging patches alone instead of commuting with social group members. When individuals were near group members, they tended to rest, and individuals that rested close to one another tended to rest for longer. The strength of their proximity-based networks was not related to measures of foraging success or energy conservation, and it appears that while the main benefit of social foraging is often assumed to be increased foraging success, the bats in

our study rely on social partners during rest. It appears, unlike in the foraging associations

of other bat species, resting in *P. hastatus* reinforces social associations or has benefits

outside of foraging efficiency such as predator detection.

Social foraging should increase foraging efficiency, either through effects of food detection

or social facilitation that increase feeding rates. Individuals of many bat species forage

socially to eavesdrop on feeding calls when resources are ephemeral and searching costs

are high (Egert-Berg et al., 2018; Fenton, 2003), and some species show extraordinarily

coordinated group foraging for these ephemeral resources (Dechmann et al., 2009;

Dechmann et al., 2010). With increasing group size, Trinidad guppies decrease the time it

takes to locate food patches and increase their intake rates (Snijders et al., 2021). Larger

guppy groups act as more efficient sensors, but come at a cost of perceived feeding

competition that drives increased bite rates. While we did not directly measure intake rate,

we did not find any association between proximity network strength and any foraging

metrics. The *P. hastatus* in our study did not modify their foraging, either in the time spent

foraging, or the number of foraging patches or feeding locations used in relation to group

members.

We found that individual *P. hastatus* from multiple social groups used the same patches of

O. pyramidale, but arrived at these patches over 25 km from their roost alone. We found

that some individuals would reunite with social group members in these patches, while

other tended to forage completely on their own and show no spatial or temporal overlap

with social group members. This is similar to vampire bats (*Desmodus rotundus*) that fly

individually to cattle approximately 300 m away and feed with partners with close social

relationships (Ripperger and Carter, 2021). Previous radio tracking work in Trinidad found

that during most of the year, social group members did not depart or arrive at the cave

together, and tended to forage alone, but on rare occasions joined a group mate in a

nearby foraging area (McCracken and Bradbury, 1981). Members of social groups occupied

adjacent foraging ranges and social groups were segregated across the landscape

(McCracken and Bradbury, 1981), unlike the large overlap we found in our data. In Trinidad

groups were more likely to depart together, and females from the same social group were

captured around a *O. pyramidale* feeding site more often than randomly expected during

the dry season (Wilkinson and Boughman, 1998). There also appears to be strong social

attraction among *P. hastatus* social groups. Social calls broadcast at flowering *O.*

pyramidale attracted bats, and social calls from flying *P. hastatus* were most often noted 3-

4 hours after sunset when bats were already out foraging (Wilkinson and Boughman, 1998).

The higher spatial and temporal resolution of our GPS tracks now indicates they are at least

not only attracted to the possibility of food resources, but that they may form resting

associations while outside of their roost.

It is possible that at this time of year or within this population in Panama there is more

pressure to maintain social associations than to group forage. Within social groups that are

structured by kin relationship, strong social bonds between related individuals have

numerous life history advantages (Silk, 2007), extending lifespans (Barocas et al., 2011; Silk

et al., 2010), success in group conflicts (Samuni et al., 2021), and individual reproductive

success (Frere et al., 2010; Schülke et al., 2010; Silk et al., 2003). However, on Trinidad,

female *P. hastatus* form stable, relatively closed groups of unrelated females that stay

together for their lifetime and show highly developed social bonds (McCracken and

Bradbury, 1981; Wilkinson et al., 2016). They develop group-specific social calls

(Boughman, 1997; Wilkinson and Boughman, 1998), and recognize and guard group-

members' offspring (Bohn et al., 2009; Bohn et al., 2007). Low reproductive rates and high

infant mortality in this species (Stern and Kunz, 1998) are strong selective pressures on

potential cooperation among unrelated females. The conditions structuring *P. hastatus*

social groups may be similar to groups of unrelated females in wild equids (Cameron et al.,

2009) and cooperative breeding birds (Riehl and Strong, 2018). Close rest while foraging,

rather than feeding together, may be one way for the bats to reinforce their social bonds

since *P. hastatus* has not been observed to engage in the extended grooming that

characterizes social bond maintenance in other mammals (Kern and Radford, 2021; Silk et

al., 2003; Wilkinson, 1986), or to potentially guard against predation.

Alternatively, there could be regional differences in the main drivers on social group

formations depending on the resource landscape. On Trinidad it has been hypothesized

that recruitment of group members to flowering trees may predominantly help bats defend

trees against competitors (Wilkinson and Boughman, 1998). However, in Panamá large

animals that bats cannot defend against, such as kinkajous and opossums are the main

visitors of *O. pyramidale* flowers (Kays et al., 2012). A flowering *O. pyramidale* with mean

peak flower availability of 60 flowers per night provides approximately 600 kJ of energy at

the beginning of the night before nectar pools are depleted and trees begin producing less

energy-dense nectar (Kays et al., 2012). We estimated that our tracked bats expended 198

± 69 kJ per day, indicating that a single *O. pyramidale* crown could support the daily energy

needs of 3 - 7 bats. Such a limited resource may be worth defending from conspecifics if all

available flowers could be fully exploited (Wilkinson and Boughman, 1998), but would not

support the needs of an entire social group of bats without moving to several patches to

feed. The closest spatial associations among foraging group members did not occur in the

feeding patches, suggesting that a collective resource defense is not the likely explanation

for foraging and resting near group mates for this population in Panamá.

Instead, social information may mainly be used for the discovery of feeding areas.

Commuting to repeatedly used foraging patches 25 km or more from a central roost may

not be unusual for some bats (Calderón-Capote et al., 2020; Goldshtein et al., 2020; Harten

et al., 2020; O'Mara et al., 2021), but still presents navigational and energetic risks due to

the long distances traveled and potential weather hazards. Nectar and fruit-feeding fruit

bats appear to have large and robust cognitive maps of their (Harten et al., 2020; Toledo et

al., 2020). These species will all commute from a common roost to distant foraging

patches, but there is little to no overlap among individual foraging ranges in animals from

these roosting aggregations, and these non-overlapping ranges may be developed through

reinforcement learning that minimizes competition (Goldshtein et al., 2020).

On the other hand, social groups of *P. hastatus* seem to seek out one another while

foraging away from their cave roosting site. Individuals commuted separately to a foraging

area, and tended to only synchronize their behavior when resting. These individuals were

exploiting the same resources, but did not seem to do so in a coordinated way.

Phyllostomus hastatus were previously found to forage within 10 km of their (McCracken

and Bradbury, 1981; William and Williams, 1970), and most will successfully navigate back

to their home roost after displacement of up to 20-25 km. At 30 km, however, bats

searched seemingly randomly, then returned to the release point, repeating this process up

to four times, and rarely returned when displaced 50 km or more (William and Williams,

1970). This indicates that the distance at which our study animals were foraging from their

cave was at the edge or even outside the range they were familiar with. This is particularly

true since they had to cross the sea and large commercial banana farms to reach these

areas. All of our tracked bats foraged in areas with *O. pyramidale*, where on their home

island and other nearby mainland locations *O. pyramidale* was not yet available. This far-

removed distance suggests that once an *O. pyramidale* patch is discovered, the social

information quickly moves through a colony bringing social group members together in

foraging areas. *Phyllostomus hastatus* feeds on various food sources throughout most of

the year, but almost exclusively switches to *O. pyramidale* when it is available during the

dry season (Wilkinson and Boughman, 1998). It is unknown if this is due to shortage of

other food sources or to a preference for nectar, but locating this widely distributed food

source may rely on social information transfer. This may be a rare, but crucial resource

during the transition period when few trees are flowering or in fruit, and finding food is more

unpredictable. Further tracking and detailed dietary analysis are needed to identify how and

when bats switch between food resources and the behavioral and energetic correlates of

this change.

Data availability

The full GPS data set and code is available from the Movebank Data Repository (doi

released on acceptance, code and data with environmental and behavioral annotation is

available at Dryad for review

(https://datadryad.org/stash/share/DBJdu_DVmEVBp6Yeq5x2RwupCy6W9NOaprjDtUZIH3E).

References

Barocas A, Ilany A, Koren L, Kam M, Geffen E, 2011. Variance in centrality within rock hyrax
social networks predicts adult longevity. PLoS One 6:e22375. doi:
10.1371/journal.pone.0022375.
Author. 2016. MuMIn: Multi-Model Inference.
Bates D, Mächler M, Bolker B, Walker S, 2015. Fitting Linear Mixed-Effects Models Using
lme4. J Stat Softw 67:1-48. doi: 10.18637/jss.v067.i01.
Beauchamp G, Heeb P, 2001. Social foraging and the evolution of white plumage. Evol Ecol
Res 3:703-720.
Bell WJ, 1990. Searching Behaviour: The behavioural ecology of finding resources. London:
Chapman and Hall.
Berdahl A, Torney C, Ioannou C, Faria J, Couzin I, 2013. Emergent sensing of complex
environments by mobile animal groups. Science 339:574-576. doi:
10.1126/science.1225883.
Bhattacharya K, Vicsek T, 2014. Collective foraging in heterogeneous landscapes. J R Soc
Interface 11:20140674. doi: 10.1098/rsif.2014.0674.
Bohn KM, Moss CF, Wilkinson GS, 2009. Pup guarding by greater spear-nosed bats. Behav
Ecol Sociobiol 63:1693-1703. doi: 10.1007/s00265-009-0776-8.
Bohn KM, Wilkinson GS, Moss CF, 2007. Discrimination of infant isolation calls by female
greater spear-nosed bats, *Phyllostomus hastatus*. Anim Behav 73:423-432. doi:
10.1016/j.anbehav.2006.09.003.
Boughman J, Wilkinson G, 1998. Greater spear-nosed bats discriminate group mates by
vocalizations. Anim Behav 55:1717-1732. doi: 10.1006/anbe.1997.0721.
Boughman JW, 1997. Greater spear-nosed bats give group-distinctive calls. Behav Ecol
Sociobiol 40:61-70. doi: 10.1007/s002650050316.
Boughman JW, 1998. Vocal learning by greater spear-nosed bats. Proc R Soc B 265:227-
233. doi: 10.1098/rspb.1998.0286.

Boughman JW, 2006. Selection on social traits in greater spear-nosed bats, Phyllostomus
hastatus. Behav Ecol Sociobiol 60:766-777. doi: 10.1007/s00265-006-0220-2.
Calderón-Capote MC, Dechmann DKN, Fahr J, Wikelski M, Kays R, O'Mara MT, 2020.
Foraging movements are density-independent among straw-coloured fruit bats. R
Soc Open Sci 7. doi: 10.1098/rsos.200274.
Cameron EZ, Setsaas TH, Linklater WL, 2009. Social bonds between unrelated females
increase reproductive success in feral horses. Proc Natl Acad Sci U S A 106:13850-
13853. doi: 10.1073/pnas.0900639106.
Clauset A, Newman ME, Moore C, 2004. Finding community structure in very large
networks. Phys Rev E Stat Nonlin Soft Matter Phys 70:066111. doi:
10.1103/PhysRevE.70.066111.
Conradt L, Roper TJ, 2005. Consensus decision making in animals. Trends Ecol Evol
20:449-456.
Dechmann D, Heucke S, Giuggioli L, Safi K, Voigt C, Wikelski M, 2009. Experimental
evidence for group hunting via eavesdropping in echolocating bats. Proc R Soc B
276:2721-2728. doi: 10.1098/rspb.2009.0473.
Dechmann D, Kranstauber B, Gibbs D, Wikelski M, 2010. Group hunting-a reason for
sociality in molossid bats? PLoS One 5:e9012. doi: 10.1371/journal.pone.0009012.
Egert-Berg K, Hurme ER, Greif S, Goldstein A, Harten L, Herrera ML, Flores-Martinez JJ,
Valdes AT, Johnston DS, Eitan O, Borissov I, Shipley JR, Medellin RA, Wilkinson
GS, Goerlitz HR, Yovel Y, 2018. Resource ephemerality drives social foraging in
bats. Curr Biol 28:3667-3673. doi: 10.1016/j.cub.2018.09.064.
Farine DR, 2013. Animal social network inference and permutations for ecologists in R using
asnipe. Methods Ecol Evol 4:1187-1194.
Fenton MB, 2003. Eavesdropping on the echolocation and social calls of bats. Mammal Rev
33:193-204. doi: 10.1046/j.1365-2907.2003.00019.x.
Frere CH, Krutzen M, Mann J, Connor RC, Bejder L, Sherwin WB, 2010. Social and genetic
interactions drive fitness variation in a free-living dolphin population. Proc Natl Acad
Sci USA 107:19949-19954. doi: 10.1073/pnas.1007997107.
Giraldeau L, Beauchamp G, 1999. Food exploitation: searching for the optimal joining policy.
Trends Ecol Evol 14:102-106.
Goldshtein A, Handel M, Eitan O, Bonstein A, Shaler T, Collet S, Greif S, Medellín RA,
Emek Y, Korman A, Yovel Y, 2020. Reinforcement learning enables resource
partitioning in foraging bats. Curr Biol. doi: 10.1016/j.cub.2020.07.079.
Hahsler M, Piekenbrock M, Doran D, 2019. dbscan: Fast Density-Based Clustering with R. J
Stat Softw 91. doi: 10.18637/jss.v091.i01.
Harten L, Katz A, Goldshtein A, Handel M, Yovel Y, 2020. The ontogeny of a mammalian
cognitive map in the real world. Science 369:194-197. doi: 10.1126/science.aay3354.
Hedenström A, 2003. Twenty-Three Testable Predictions About Bird Flight. In: Berthold P,
Gwinner E, Sonnenschein E, editors. Avian Migration Berlin: Springer. p. 563-582.
Author. 2020. fpc: Flexible Procedures for Clustering. Version R package version 2.2-5.
Kays R, Rodríguez M, Valencia L, Horan… R, 2012. Animal visitation and pollination of
flowering balsa trees (*Ochroma pyramidale*) in Panama. Mesoamericana 16:54-70.
Kern JM, Radford AN, 2021. Strongly bonded individuals prefer to forage together in
cooperatively breeding dwarf mongoose groups. Behav Ecol Sociobiol 75. doi:
10.1007/s00265-021-03025-0.
Kranstauber B, Kays R, Lapoint S, Wikelski M, Safi K, 2012. A dynamic Brownian bridge
movement model to estimate utilization distributions for heterogeneous animal
movement. J Anim Ecol 81:738-746. doi: 10.1111/j.1365-2656.2012.01955.x.
Kunz TH, Robson SK, Nagy KA, 1998. Economy of harem maintenance in the greater
spear-nosed bat, *Phyllostomus hastatus*. J Mammal:631-642.
Kuznetsova A, Brockhoff PB, Christensen RHB, 2017. lmerTest Package: Tests in Linear
Mixed Effects Models. J Stat Softw 82. doi: 10.18637/jss.v082.i13.

McClintock BT, Michelot T, Goslee S, 2018. momentuHMM: R package for generalized
hidden Markov models of animal movement. Methods Ecol Evol 9:1518-1530. doi:
10.1111/2041-210x.12995.
McCracken GF, Bradbury JW, 1981. Social organization and kinship in the polygynous bat
Phyllostomus hastatus. Behav Ecol Sociobiol 8:11-34. doi: 10.1007/Bf00302840.
McInnes AM, McGeorge C, Ginsberg S, Pichegru L, Pistorius PA, 2017. Group foraging
increases foraging efficiency in a piscivorous diver, the African penguin. R Soc Open
Sci 4:170918. doi: 10.1098/rsos.170918.
McNab BK, 1969. The economics of temperature regulation in neotropical bats. Comp
Biochem Physiol 31:227-268. doi: 10.1016/0010-406X(69)91651-X.
Michelot T, Langrock R, Bestley S, Jonsen ID, Photopoulou T, Patterson TA, 2017.
Estimation and simulation of foraging trips in land-based marine predators. Ecol
98:1932-1944. doi: 10.1002/ecy.1880.
O'Mara M, Dechmann D, Page R, 2014a. Frugivorous bats evaluate the quality of social
information when choosing novel foods. Behavioral Ecology 25:1233-1239. doi:
10.1093/beheco/aru120.
O'Mara MT, Amorim F, Scacco M, McCracken GF, Safi K, Mata V, Tome R, Swartz S,
Wikelski M, Beja P, Rebelo H, Dechmann DKN, 2021. Bats use topography and
nocturnal updrafts to fly high and fast. Curr Biol 31:1311-1316 e1314. doi:
10.1016/j.cub.2020.12.042.
O'Mara MT, Scharf AK, Fahr J, Abedi-Lartey M, Wikelski M, Dechmann DKN, Safi K, 2019.
Overall dynamic body acceleration in straw-colored fruit bats increases in headwinds
but not with airspeed. Front Ecol Evol 7:200. doi: 10.3389/fevo.2019.00200.
O'Mara MT, Wikelski M, Dechmann DKN, 2014b. 50 years of bat tracking: device
attachment and future directions. Methods Ecol Evol 5:311-319. doi: 10.1111/2041-
210x.12172.
O'Mara MT, Wikelski M, Voigt CC, Ter Maat A, Pollock HS, Burness G, Desantis LM,
Dechmann DK, 2017. Cyclic bouts of extreme bradycardia counteract the high
metabolism of frugivorous bats. eLife 6:e26686. doi: 10.7554/eLife.26686.
Page R, Ryan M, 2006. Social transmission of novel foraging behavior in bats: frog calls and
their referents. Curr Biol 16:1201-1205. doi: 10.1016/j.cub.2006.04.038.
Pennycuick CJ, 2008. Modelling the flying bird. London: Academic Press.
Author. 2020. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna Austria.
Ramakers JJC, Dechmann DKN, Page RA, O'Mara MT, 2016. Frugivorous bats prefer
information from novel social partners. Anim Behav 116:83-87. doi:
10.1016/j.anbehav.2016.03.021.
Ranacher P, Brunauer R, Trutschnig W, Van der Spek S, Reich S, 2016. Why GPS makes
distances bigger than they are. Int J Geogr Inf Sci 30:316-333. doi:
10.1080/13658816.2015.1086924.
Ratcliffe J, ter Hofstede H, 2005. Roosts as information centres: social learning of food
preferences in bats. Biol Lett 1:72-74. doi: 10.1098/rsbl.2004.0252.
Riehl C, Strong MJ, 2018. Stable social relationships between unrelated females increase
individual fitness in a cooperative bird. Proc R Soc B 285. doi:
10.1098/rspb.2018.0130.
Ripperger SP, Carter GG, 2021. Social foraging in vampire bats is predicted by long-term
cooperative relationships. PLoS Biol 19:e3001366. doi:
10.1371/journal.pbio.3001366.
Robitaille AL, Webber QMR, Vander Wal E, Golding N, 2019. Conducting social network
analysis with animal telemetry data: Applications and methods using spatsoc.
Methods Ecol Evol 10:1203-1211. doi: 10.1111/2041-210x.13215.
Safi K, Kranstauber B, Weinzierl R, Griffin L, Rees EC, Cabot D, Cruz S, Proano C,
Takekawa JY, Newman SH, Waldenstrom J, Bengtsson D, Kays R, Wikelski M,

687    Bohrer G, 2013. Flying with the wind: scale dependency of speed and direction
688    measurements in modelling wind support in avian flight. Mov Ecol 1:4. doi:
689    10.1186/2051-3933-1-4.
690 Samuni L, Crockford C, Wittig RM, 2021. Group-level cooperation in chimpanzees is shaped
691    by strong social ties. Nat Commun 12:539. doi: 10.1038/s41467-020-20709-9.
692 Schubert E, Sander J, Ester M, Kriegel HP, Xu X, 2017. DBSCAN Revisited, Revisited. ACM
693    Transactions on Database Systems 42:1-21. doi: 10.1145/3068335.
694 Schülke O, Bhagavatula J, Vigilant L, Ostner J, 2010. Social bonds enhance reproductive
695    success in male macaques. Curr Biol 20:2207-2210. doi: 10.1016/j.cub.2010.10.058.
696 Silk JB, 2007. The adaptive value of sociality in mammalian groups. Philos Trans R Soc B
697    362:539-559. doi: 10.1098/rstb.2006.1994.
698 Silk JB, Alberts SC, Altmann J, 2003. Social bonds of female baboons enhance infant
699    survival. Science 302:1231-1234.
700 Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM,
701    Seyfarth RM, Cheney DL, 2010. Strong and consistent social bonds enhance the
702    longevity of female baboons. Curr Biol 20:1359-1361. doi:
703    10.1016/j.cub.2010.05.067.
704 Snijders L, Krause S, Tump AN, Breuker M, Ortiz C, Rizzi S, Ramnarine IW, Krause J,
705    Kurvers R, 2021. Causal evidence for the adaptive benefits of social foraging in the
706    wild. Commun Biol 4:94. doi: 10.1038/s42003-020-01597-7.
707 Speakman J, 2005. Body size, energy metabolism and lifespan. Journal of Experimental
708    Biology 208:1717-1730. doi: 10.1242/jeb.01556.
709 Stern AA, Kunz TH, 1998. Intraspecific variation in postnatal growth in the greater spear-
710    nosed bat. J Mammal 79:755. doi: 10.2307/1383086.
711 Stilz WP, Schnitzler HU, 2012. Estimation of the acoustic range of bat echolocation for
712    extended targets. J Acoust Soc Am 132:1765-1775. doi: 10.1121/1.4733537.
713 Thomas SP, 1975. Metabolism during flight in two species of bats, *Phyllostomus hastatus*
714    and *Pteropus gouldii*. J Exp Biol 63:273-293.
715 Toledo S, Shohami D, Schiffner I, Lourie E, Orchan Y, Bartan Y, Nathan R, 2020. Cognitive
716    map–based navigation in wild bats revealed by a new high-throughput tracking
717    system. Science 369:188-193. doi: 10.1126/science.aax6904.
718 Valone TJ, 1989. Group foraging, public information, and patch estimation. Oikos:357-363.
719 von Busse R, Swartz SM, Voigt CC, 2013. Flight metabolism in relation to speed in
720    Chiroptera: testing the U-shape paradigm in the short-tailed fruit bat *Carollia*
721    *perspicillata*. J Exp Biol 216:2073-2080. doi: 10.1242/jeb.081760.
722 Ward S, Moller U, Rayner JM, Jackson DM, Bilo D, Nachtigall W, Speakman JR, 2001.
723    Metabolic power, mechanical power and efficiency during wind tunnel flight by the
724    European starling *Sturnus vulgaris*. J Exp Biol 204:3311-3322.
725 Wilkinson G, Boughman J, 1998. Social calls coordinate foraging in greater spear-nosed
726    bats. Anim Behav 55:337-350.
727 Wilkinson GS, 1986. Social grooming in the common vampire bat, *Desmodus rotundus*.
728    Anim Behav 34:1880-1889.
729 Wilkinson GS, Carter GG, Bohn KM, Adams DM, 2016. Non-kin cooperation in bats. Philos
730    Trans R Soc B 371. doi: 10.1098/rstb.2015.0095.
731 William TC, Williams JM, 1970. Radio tracking of homing and feeding flights of a neotropical
732    bat, *Phyllostomus hastatus*. Anim Behav 18:302-309. doi: 10.1016/s0003-
733    3472(70)80042-2.
734 Wright GS, 2016. Social Learning and Information Transfer in Bats: Conspecific Influence
735    Regarding Roosts, Calls, and Food. In: Ortega J, editor. Sociality in Bats: Springer.
736    p. 211-230.
737 Wyman MT, Pinter-Wollman N, Mooring MS, Hayes L, 2021. Trade-offs between fighting
738    and breeding: a social network analysis of bison male interactions. J Mammal. doi:
739    10.1093/jmammal/gyaa172.

741   **Supplemental Information**

Table S1. Biometric and tracking information for the individuals analyzed in this study. Reproductive state is coded as nr: non-reproductive, plac: post-lactation, and nulli: nulliparous. Nightly values for the minutes of tracking and number of locations used are given for means ± sd when more than one night was recorded.

Bat ID	Network Label ID	Group ID	Sex	Reproductive State	Mass (g)	Forearm length (mm)	Wing length (mm)	Wing area (mm2)	Nights Analyzed	Time Analyzed (min)	Number of Locations Used	Strength Centrality	Degree
2016030703	1	one	m	nr	134	93.5	271.96	22966.52	2	133.99 ± 82.68	7296.5 ± 3939.29	0	0
2016030705	2	one	f	plac	124	87.8	280.82	24543.47	1	156.67	7694	0	0
71A0D95	3	one	f	nulli	122	89.9	282.74	24088.56	2	224.89 ± 4.82	12371.5 ± 1289.06	0.13	1
71A111A	4	one	f	nulli	109	90.8	296.86	25587.39	1	279.52	16752	0.13	1
74DDA80	5	one	f	nulli	123	91.2	278.44	23724.58	1	151.88	6268	0	0
74DE9A7	6	one	f	plac	122	91.5	274.95	23752.23	1	236.13	13488	0	0
74D8954	7	two	f	plac	128	93.02	317.17	29422.44	1	161.88	6211	0.178	3
74DA92F	8	two	f	plac	109	87.6	250.07	19667.46	2	192.87 ± 25.53	6767 ± 1236.02	0.24	5
74DCA83	10	two	f	nulli	123	92.1	300.31	29703.42	2	131.85 ± 51.45	5028 ± 2224.56	0.004	1
74D8C25	11	three	f	nulli	131	93.2	308.25	28074.65	1	118.15	5158	0.288	5
74D932E	12	three	f	nulli	116	93.6	304.29	28397.6	2	207.63 ± 2.4	8783 ± 83.44	0.317	5
74DAF9C	9	three	f	nulli	126	94.5	296.29	26317.99	1	95.27	5628	0	0
74DCBCC	13	three	f	plac	114	89.7	289.26	22936.64	2	278.75 ± 18.27	11910 ± 1578.26	0	0
74DE4E7	14	three	f	plac	124	94.2	296.11	26383.96	1	197.03	6618	0.144	4
74F7D4C	15	three	m	nr	137	93.2	277.12	23586.97	1	152.63	3748	0.137	4
74F8E19	16	three	f	plac	113	91.7	289.38	25321.02	3	201.26 ± 61.87	7702.67 ± 2597.55	0.013	2
74F9F83	17	three	f	plac	116	92	284.51	24720.16	3	264.17 ± 54.4	7866.33 ± 1690.4	0.035	2
74FE24E	18	three	f	plac	113	91.7	270.5	22057.56	2	218.35 ± 3.15	7528 ± 622.25	0.026	1

Figure S1. Mechanical power requirements at airspeeds for each bat in this study. Power

output was calculated following Pennycuick (2008) at 50 m and 25°C using each

individual's body mass and wing dimensions.

Figure S2. The number of foraging patches used is predicted by A) the distance that

individuals moved each night and B) the total time that individuals were tracked each night.

The number of feeding clusters identified was not predicted by either C) the distance

moved per night or D) the number of foraging patches used. Statistically significant slopes

that differed from zero are shown by solid lines, non-significant slopes by dashed lines, and

shaded areas indicate 95% confidence intervals.

756
757 Figure S3. Resting patches clustered from GPS locations. A) Resting patches are shown
758 across the study area, scaled by the number of individual bat days that the sites were used
759 and colored by the number of individual bats that used the patch. Resting patches are
760 shown for B) Isla Changuinola, and C) Isla Colón.
761
762
763

A   B

Figure S4. A) Proximity networks of simultaneous spatial associations (same space, same

time). Edge weights show strength centrality, and note that individuals 1 and 15 are males.

B) networks of across-night foraging patch associations (same space, different time).

Individual bats are shown as part of their social group (group 1: blue square, group 2: green

circle, group 3: yellow triangle. Foraging patch communities are highlighted by the size of

their membership (community 1: red, community 2: green, community 3: purple).

Figure S5. Relationships between proximity network strength of individual bats and the

proportion of time that they spend per night while A) Resting, B) Moving between patch, C)

slow foraging and feeding flight, and D) Commuting.