Pharyngeal timing and particle transport defects in *Caenorhabditis elegans* feeding mutants

Isaac Ravi Brenner[1], David M. Raizen[2], and Christopher Fang-Yen[2,3,*]

[1]Department of Bioengineering, School of Engineering and Applied Science, University of Pennsylvania, Philadelphia PA, 19104

[2]Department of Neurology, Perelman School of Medicine, University of Pennsylvania, Philadelphia PA, 19104

[3]Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania, Philadelphia PA, 19104

*Corresponding Author. Email: fangyen@seas.upenn.edu

Mailing address: 210 S 33rd St., Skirkanich Hall 240, Philadelphia, PA 19104

Abstract: The nematode *C. elegans* uses rhythmic muscle contractions and relaxations called pumps to filter, transport, and crush food particles. A number of feeding mutants have been identified, including those with slow pharyngeal pumping rate, weak muscle contraction, defective muscle relaxation, and defective grinding of bacteria. Many aspects of these pharyngeal behavioral defects and how they affect pharyngeal function are not well understood. For example, the behavioral deficits underlying inefficient particle transport in 'slippery' mutants have been unclear. Here we use high speed video microscopy to describe pharyngeal pumping behaviors and particle transport in wild-type animals and in feeding mutants. Different 'slippery' mutants exhibit distinct defects including weak isthmus contraction, failure to trap particles in the anterior

isthmus, and abnormal timing of contraction and relaxation in pharyngeal

compartments. Our results show that multiple deficits in pharyngeal timing or

contraction can cause defects in particle transport.

Abbreviated title: Timing and transport defects in *C. elegans* feeding mutants

Introduction

The nematode *C. elegans* consumes food bacteria using movements of the pharynx, a

neuromuscular feeding organ that connects to the buccal cavity and to the intestine.

The pharynx displays two behaviors: pharyngeal pumping and isthmus peristalsis.

Pharyngeal pumping consists of rhythmic cycles of contraction and relaxation

throughout most of the pharynx. During contraction, the radially oriented muscle fibers

open the pharyngeal lumen; during relaxation the lumen closes. Isthmus peristalsis is

an anterior-to-posterior contraction in the isthmus that occurs after a fraction of pumps

(Avery and Horvitz, 1987). Together, these movements filter food particles from

surrounding fluids, transports the particles posteriorly, and crushes them before they

enter the intestine.

The pharynx is divided into several compartments including the anterior and posterior

(terminal) bulbs. Detailed studies of pharyngeal pumping have revealed differences in

timing of contractions and relaxations in these compartments (Avery, 1993a, 1993b;

Fang-Yen et al., 2009). The corpus, which extends from the buccal cavity to the

anterior bulb, is the first to contract, drawing particles from the buccal cavity into the

pharyngeal lumen. Contraction of the isthmus and then the terminal bulb follow

contraction of the corpus with short delays. Pharyngeal relaxation of the various

compartments proceeds in the same order as their contraction but several times more

rapidly (Fang-Yen 2009).

The sequence of relaxations in pharyngeal compartments is closely tied to the

mechanism of particle trapping. The anterior tip of the corpus relaxes slightly before the

rest of the corpus; this allows food particles to become trapped at the anterior corpus.

Similarly, relaxation of the metacorpus before the anterior isthmus allows food particles

to become trapped at the anterior isthmus. At each of the two trap sites (anterior corpus

and anterior isthmus), earlier relaxation of the structures immediately anterior to the trap

prevents anteriorward transport of particles while fluids are expelled (Fang-Yen et al.,

2009). Under some conditions such as violet light exposure, the trap at the anterior

procorpus remains contracted, causing particles to be ejected from the anterior pharynx

rather than trapped at the end of the pump (Bhatla et al., 2015; Sando et al., 2021).

Particles trapped in the anterior isthmus are transported to the pharyngeal grinder via

isthmus peristalsis.

A number of 'eat' mutants with defective feeding behavior were identified by Avery

(Avery, 1993a).

Feeding phenotypes included slow pumping, defective timing of relaxation, and a

"slippery pharynx" phenotype characterized by inefficient transport of particles. These

feeding mutants were described using video recordings at 60 frames per second (fps), a

frame rate insufficient to resolve the rapid movements of the pharynx and food particles

during muscle relaxation. As a result, aspects of pharyngeal behavior in feeding

mutants have remained unclear. In particular, the mechanisms by which 'slippery'

mutants are defective in anterior-to-posterior particle transport are poorly understood.

Here, we revisit the analysis of feeding mutants using high speed video microscopy at

1000 fps. We imaged worms feeding on bacteria and on polystyrene beads. By

tracking the motions of pharyngeal components and particles within the pharyngeal

lumen, we quantified the timing of contraction and relaxation, motion of particles in the

pharyngeal lumen, and efficiency by which particles were transported and trapped. We

also measured the pumping rate and the frequency of posterior isthmus peristalsis, by

which particles trapped in the anterior isthmus are transported to the terminal bulb.

Different slippery mutants showed a variety of behavioral defects, including weak

isthmus contraction, failure to trap particles in the anterior isthmus, and abnormal timing

of contractions and relaxations in different pharyngeal compartments. We also describe

behavioral differences in slow-feeding and relaxation-defective mutants.

Results

The class of slippery mutants were labeled as such by Avery because in these animals,

bacteria were observed to slip toward the anterior instead of being trapped during

pharyngeal relaxation (Avery, 1993a). We asked what differences in pharyngeal

behavior relative to wild-type animals led to the slippery phenotype.

We first used our assay to describe wild-type pharyngeal behavior. In wild-type

animals, pump timing and particle movement were consistent with previous findings

(Fang-Yen et al., 2009). Each pump began with the onset of contraction of the corpus,

followed 28.5 ± 14.8 ms (mean ± SD) later by onset of contraction of the anterior

isthmus, followed 9.6 ± 11.7 ms (mean ± SD) later by onset of contraction of the

terminal bulb (Fig, 1b, Video S1, Supplemental Table A3). Onset of relaxation occurred

in the same order but faster, beginning with the corpus, followed 13.4 ± 9.9 ms later by

the anterior isthmus, and then 6.4 ± 10.7 ms later by the terminal bulb (Fig. 1b). The

pumping rate was 4.4 Hz ± 0.48 (mean ± SD). Isthmus peristalsis occurred after 22.6%

of pumps, consistent with prior studies (Avery, 1993a; Song and Avery, 2012).

To assay the efficiency of particle transport, we reviewed video sequences to manually

track trajectories of particles as they moved through the pharynx (see Methods). We

found that pharynxes in wild-type animals transported particles efficiently: 79.5% (35 out of 44) of tracked particles in the corpus at the beginning of contraction reached the anterior isthmus and were trapped there at the end of the relaxation phase (Supplemental Table A4).

Various behavioral defects contribute to inefficient transport in slippery mutants

We analyzed four slippery mutants: *eat-10*, *eat-13*, *eat-15*, and *eat-17*. By reviewing video recordings, we examined the patterns and timing of contraction and relaxation of pharyngeal compartments in each mutant. We also analyzed transport of particles (bacteria and/or polystyrene beads).

In wild-type animals, robust contraction of the anterior isthmus causes particles in the corpus to be drawn into the isthmus (Fig. 1a,b) where they can be trapped. By contrast, in *eat-10* mutants the anterior isthmus frequently either failed to contract (18/49 pumps, 36%) or contracted weakly (14/49 pumps, 29%). Neither weak nor absent contractions were ever observed in wild-type worms. During weak contractions in *eat-10* worms, there was very limited (<20% of normal) opening of the pharyngeal lumen (Fig. 1c, Video S2 worms 3-4). In cases of weak or absent anterior isthmus contraction, particles were not drawn into the anterior isthmus (Fig. 1c, Video S2, Worm 3-4). As a result, in

eat-10 mutants, only 33% (16 out of 49) of the particles starting in the corpus at the

beginning of a pump became trapped in the anterior isthmus at the end of the pump. In

comparison, in wild-type animals, 79.5% moved from the corpus to isthmus at the end

of the pump (Fig. 4). This finding is consistent with the original description of slippery

pumping in which many particles reach the metacorpus at the peak of contraction but

then return to the anterior procorpus during relaxation (Avery 1993a).

*Figure 1: **Particle tracking and pharyngeal subcomponent timing of wild type***

worms and slippery mutants. *(A) The pharynx and pharyngeal compartments in a*

wild type animal. Scale bar: 25 μm. (B-F) Pharyngeal muscle dynamics and particle

tracking data from high-speed video microscopy of a single representative pump cycle

from wild type (N2) and mutant worms. Colored traces indicate trajectories of four

 particles. Colored areas show timings of contraction and relaxation of the corpus,

 anterior isthmus, and terminal bulb, with red representing contraction and blue

 representing relaxation. The x axis corresponds to the longitudinal positions shown in

 (A). For eat-10, no isthmus contraction or relaxation was visible during this pump

 (Panel C, Supplemental Video S2 worm 4). Details of particle tracking data are given in

 Supplemental Table A4.

As in *eat-10* mutants, we found in *eat-15* worms a low (20/42 = 47.6%) rate of particle

transport from the corpus to isthmus compared to *wild type* (Fig. 4). In some pumps,

isthmus contraction was weak (Video S4 worm 3). Particle tracking showed that

particles initially located in the corpus often did not enter the isthmus, instead stalling

upon reaching the metacorpus (Fig. 1e). In some pumps, the particles moved back and

forth along the lumen during relaxation, perhaps due to influence from movements in

the terminal bulb (e.g. Video S4 worm 4).

In *eat-13* mutants, efficiency of transport from the corpus to isthmus (24/38 = 63%) was

only slightly lower than that of wild-type animals (79.5%). Pharynxes in *eat-13* mutants

appeared physically distorted: the ratio of the isthmus width and the terminal bulb width

to the length of the pharynx in *eat-13* mutants was greater than that ratio in wild-type

worms (Video S3, Fig. 2a-e). This difference in morphology is unlikely to be explained

by a nutritional defect alone since *eat-2* mutants, which have a defect in cholinergic

excitation of the pharynx and therefore pump slowly (McKay et al., 2004; Raizen, Lee,

and Avery, 1995), had a ratio no different from that of wild-type worms.

Figure 2: Pharynxes of *eat-13* animals are distorted. (A) Pharynx length for N2, *eat-13*, and *eat-2* animals. (B) Ratio of procorpus width to pharynx length. (C) Ratio of isthmus width to pharynx length. (D). Ratio of TB width to pharynx length. Significance threshold from t-test: * indicates $p < 0.05$, ** indicates $p < 10^{-5}$, *** indicates $p < 10^{-8}$. Note that the *eat-13* pharynx is generally shorter and wider than N2. See Supplemental Table A5

for detailed measurements. (E) Deformed *eat-13* pharynx. (F) Pharynx of an *eat-17*

worm has a deformed grinder. See also Video S5. Scale bar: 25 µm for both images.

The gene *eat-17* encodes a GTPase-activating protein required for pharyngeal cuticle

structure (Straud et al., 2013). In *eat-17* mutants, we observed a high percentage of

particles failing to make posterior progress along the pharyngeal lumen (29% compared

to 14% in wild-type, Supplemental Table A4). However, unlike *eat-13* mutants, *eat-17*

mutants were highly efficient in corpus-to-isthmus transport, suggesting that most of the

transport inefficiency in *eat-17* mutants was occurring in the isthmus. The grinder in the

terminal bulb of *eat-17* mutant pharynxes appeared deformed (Video S5, Fig. 2f), as

previously reported (Straud et al., 2013). Overall, *eat-17* mutants were able to transport

particles from the corpus to the isthmus efficiently, but had defective transport within the

isthmus (Video S5).

Slippery mutants display altered isthmus and terminal bulb contraction timings

In wild-type worms, the onset of contraction of the terminal bulb nearly always occurred

after onset of contraction in the isthmus, and the onset of relaxation of the terminal bulb

occurred predominantly after onset of relaxation in the isthmus (Fig. 3, Supplemental

Table A3). In *eat-10, eat-13,* and *eat-15* animals, the terminal bulb began contracting

earlier on average than the isthmus (Fig 3a). Similarly, on average, the terminal bulb

also began relaxation before the isthmus (Fig. 3b), although the difference was smaller

than that of the contraction delay.

Figure 3: Delay between the onset of (A) contraction and (B) relaxation of the terminal

blub (TB) relative to the isthmus in wild-type worms and slippery mutants. Negative

delay indicates that terminal bulb contracted/relaxed prior to isthmus. Each point

represents a single pump. For each genotype, data is from 1 pump from each of N =

10-19 animals (See Supplemental Table A3). Error bars indicate +/- SEM. Results of

two-tailed Student's t-test comparing each strain to N2; * indicates p<0.05, ** indicates

p<0.005, *** indicates p<0.0005.

Relaxation defective mutants show defective transport

The pharynxes of loss-of-function mutants e*at-6* fail to relax completely during the pump cycle (Avery, 1993a). The gene *eat-6* encodes a sodium/potassium ATPase, a pump required to maintain the resting membrane potential (Davis et al., 1995). In *eat-6* mutants, we observed that the terminal bulb only occasionally came back to its fully relaxed position, usually appearing to only partially relax (Video S6). In addition to this failure of the terminal bulb to fully relax, the relaxation phases were proportionally longer in *eat-6* mutants than in the wild type. Corpus (p=0.001) and isthmus (p=0.02) contraction phases were significantly shorter in *eat-6* after normalizing for pump period, indicating that relaxation made up a greater percentage of the overall pump. Like *eat-10* animals, *eat-6* worms frequently showed weak isthmus contractions, precluding effective transport (Fig. 5c, Video S6 worms 2-4). This resulted in a lower rate of corpus to isthmus transport (23.2% compared to 79.5% in N2; Fig. 4).

Like *eat-6* mutants, gain-of-function mutants of *egl-19 (*an L-type voltage-gated calcium channel (Lee et al 1997) fail to relax completely during the pump cycle (Avery, 1993a). In an *egl-19* gain-of-function mutant, the most notable behavior was occasional (3/46 pumps) isthmus and terminal bulb "yawns" in which those parts of the pharynx would remain contracted while the corpus went through its normal pump cycle (Video S7 worm 1-2). These terminal bulb contractions would span two pumps of the corpus. Even when the contraction ended prior to the next pump (i.e., no discernable "yawn" occurred), the isthmus and terminal bulb remained contracted longer than observed in wild-type worms (Supplemental Table A2). Consistent with this, we found a longer delay between onset

of corpus relaxation, and that of the isthmus, and subsequently the terminal bulb

(Supplemental Table A3). We observed a lower rate of corpus to isthmus transport

(18/40, 45%), as well as an increase in particles ending in the metacorpus (21%

compared to 0% in wild type (Fig. 4, Supplemental Table A4).

Figure 4: (A) Particle transport efficiency, defined as the fraction of particles starting in

the corpus that are transported to the isthmus during a pump. Error bars show ±

standard deviation assuming a binomial distribution. (B) Fraction of pumps that were

followed by isthmus peristalsis. Error bars show standard deviation assuming a

binomial distribution.   (C) Pump rate, with black bars indicating mean +/- SD. Additional

data can be found in Supplemental Tables A1 and A4.

Figure 5. **Particle tracking and pharyngeal subcomponent timing of N2, eat-6, egl-**

19, and eat-2 mutants. *(A-E) Representative particle trajectories and*

contraction/relaxation timings, as in Fig. 1.

Slow pumping mutant *eat-2* shows normal pharyngeal dynamics and particle transport

We next analyzed animals mutant for the gene *eat-2*, which encodes a non-alpha subunit of a nicotinic acetylcholine receptor (McKay et al., 2004). Because EAT-2 is required for excitation of the pharynx by the motor neuron MC (Raizen et al., 1995) animals mutant for *eat-2* have a pump rate much lower than that of wild-type (1.3 ± 0.5 Hz compared to 4.4 ± 0.48 Hz for N2). To assess differences in pump timing, we normalized timing measurements by pump duration. After normalizing for pump duration, we found that the duration of the contraction and relaxation phases were nearly identical to that of N2. Consistent with previously reported results (Kozlova et al., 2019), we observed a higher rate of peristalsis in *eat-2,* with peristalsis occurring after 45.5% of pumps, compared to 22.6% in N2. Aside from slow pumping and increased peristalsis, *eat-2* mutants displayed normal pharyngeal muscle dynamics and particle trapping (Fig. 5E, Video S8). Transport efficiency was similar to that of wild-type worms, with about 65% of particles in the corpus becoming trapped in the isthmus.

Discussion

In this work we used high speed video microscopy to observe pharyngeal muscle movements and particle transport in feeding mutants. We found that three of the four

mutants described as slippery had defects in posteriorward transport, consistent with

previous observations (Avery, 1993a).

The mechanisms by which differences in pharyngeal movements create particle

transport defects were clear only in a few cases. In *eat-10*, and possibly *eat-15*,

inefficiency of posterior transport could be explained by a deficiency or absence of

anterior isthmus contraction. Isthmus contraction is required for particles to be drawn

from the metacorpus into the isthmus, prior to being trapped there.

Though pharyngeal muscles are electrically coupled (Starich et al., 1996) the pharynx

exhibits highly compartmentalized muscle dynamics, between and even within individual

muscle cells. For example, the anterior tips of the pm3 muscle cells, which span the

procorpus, relax earlier than the rest of the corpus during pumping (Fang-Yen et al.,

2009), a behavior that is antagonized by activity of the M1 neuron (Bhatla et al., 2015;

Sando et al., 2021). Similarly, though the isthmus is composed of just one muscle cell

type (pm5), the anterior and posterior compartments of the isthmus display different

behaviors: the anterior isthmus contracts during pumps whereas the posterior isthmus

displays a peristaltic movement after every fourth or five pumps. The finding of weak

isthmus pumping but normal posterior isthmus peristalsis in *eat-10* and possibly *eat-15*

mutants suggests that muscle contractions in subcellular compartments of the

pharyngeal isthmus are mediated by distinct mechanisms.

In discussing reasons for isthmus peristalsis, Avery proposed that normal corpus and

terminal bulb function require them to be hydrodynamically isolated from each other

during pumping (Avery, 1993c). During corpus contraction, pharyngeal lumen pressure

decreases, and the bacteria suspension is drawn into the pharyngeal lumen from the

stoma. The posterior isthmus remaining closed (relaxed) at this time may serve to

prevent gastro-pharyngeal reflux, i.e. fluids and/or particles drawn into the pharynx from

the intestine. However, if the relaxed position of the posterior isthmus mechanically

isolates the terminal bulb from the corpus, then the timing of terminal bulb contraction

should not affect corpus to anterior isthmus transport. In all four slippery mutants

studied, we observed a premature terminal bulb relaxation (Fig 3). This observation

suggests that there is a connection between terminal bulb contraction and particle

transport efficiency, and that the TB is not fully isolated from the corpus and anterior

isthmus. It is possible that a delay between terminal bulb and corpus contraction

reduces the fluid resistance of the posterior pharynx, and so desynchronization of the

terminal bulb contraction relative to that of the corpus and isthmus normally helps to

maintain posteriorward flow.

We observed poor pharyngeal transport in animals mutant for *eat-17*, which encodes a

GTPase-activating protein required for pharyngeal cuticle structure (Straud et al., 2013).

The observation of poor particle transport despite no detectable defect in pharyngeal

motion timing suggests that the pharyngeal cuticle may play a role in transport, perhaps

via mechanical interactions with food particles. Alternatively, poorly macerated bacteria

in the terminal bulb due to *eat-17* grinder dysfunction may affect the mechanics of

bacterial transport in the entire pharynx, again perhaps because the terminal bulb is not

fully mechanically isolated from the corpus and anterior isthmus.

We found that *eat-6* and *egl-19* mutants, previously described as defective in relaxation

(Avery 1993a), were most notable for their extended contraction phases. We also found

that both of these mutants were deficient in posteriorward particle transport. Therefore,

these relaxation-defective mutants can also be described as slippery. Considering the

importance of the relaxation phase for trapping and transport, this result is not

surprising.

In this work we have focused on describing the behavioral defects in feeding mutants.

Future efforts will aim to connect the molecular identity of specific genes to specific

behavioral defects. The future use of genetically encoded calcium and/or voltage

sensors could allow for an assessment of excitation in different muscle compartments.

The combination of genetic analysis and high-resolution phenotyping of pharyngeal

behavior may lead to new insights into excitable cells and organs, in particular regarding

the mechanisms of subcellular compartmentalization of muscle contraction.

Material and Methods

Strains used in this study are shown in Figures 1 and 5. *C. elegans* were cultured

using standard methods (Stiernagle, 2006). For experiments, 5-10 young adult animals

were transferred to a 1 mm thick pad composed of 2% agarose in NGM buffer on a

microscope slide, in addition to 1-2 µL of either an OP50 *E. coli* suspension in LB

medium or a 1% (vol/vol) suspension of 1 µm diameter polystyrene beads

(Polysciences 431), diluted in NGM buffer to 0.1% vol/vol. NGM buffer consisted of the

same components as NGM media (Stiernagle, 2006) without agar, peptone, or

cholesterol. The pad was covered with a #1 cover slip and worms were given 15-45

minutes to acclimate to the pad before imaging. We mounted slides on an inverted

microscope (Nikon TE-2000S) with a 60X Plan Apo (NA 1.4) oil immersion objective

lens under differential interference contrast optics. For each animal we saved 1-3

recordings, each 4-5 s in duration, at 1000 fps using a Vision Research Phantom V9.1

high speed camera.

To record muscle timing data, we used a custom MATLAB script to review videos and

manually recorded the frame numbers corresponding to 10 events: the start of corpus

contraction, the start of corpus relaxation, the end of corpus relaxation, the same for

both the isthmus and the terminal bulb, and the beginning of the next pump. This was

done only for recording periods in which all parts of the pump cycle were clearly visible.

We manually scored each pump for whether or not it was followed by isthmus

peristalsis.

To perform particle tracking, we used the same MATLAB program to analyze videos

frame by frame and recorded the approximate locations of several particles in each

pump, as well as the timings of each particle's beginning, peak, and end of movement.

To approximate particle locations, we established several guideposts along the pharynx

(tip of the corpus, both ends of the metacorpus, the isthmus trap, the end of the anterior

isthmus, and the grinder) and assigned them approximate distances relative to the tip of

the stoma. We tracked at least one particle in the corpus and isthmus on every pump in

which there were particles in those compartments.

All statistical analyses were performed in Python. Because duration of components of

the pharyngeal pump is affected by the total pump time (Fang-Yen et al., 2009), we

normalized time measurements by the pump period for each trial before making

statistical comparisons between strains. To assess the significance of these particle

tracking samples, we used the 2x2 Fisher test, comparing each mutant strain with N2.

All pump timing data can be found in Supplemental Tables A1-3, and all particle

tracking data can be found in Supplemental Table A4.

Acknowledgements

Some strains were provided by the *Caenorhabditis* Genetics Center (CGC), which is

funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We thank

Levi Kanu for conducting preliminary experiments in this project.

Video captions

Video S1. High speed video of pharyngeal pumping in N2 animals. Pump cycles from 4 wild type worms feeding on 1 μm beads. The beginning of each animal's recording can be found at the following time points (m:s): Worm 1: 0:00; Worm 2: 0:13; Worm 3: 0:27; Worm 4: 0:42. Scale bar = 25 μm in all videos. Worm 1 corresponds to tracking plot in Fig. 1b.

Video S2. High speed video of pharyngeal pumping in *eat-10* animals. The beginning of each animal's recording can be found at the following time points (m:s): Worm 1: 0:00; Worm 2: 0:15; Worm 3: 0:30; Worm 4: 0:45. Note the weak or absent isthmus contractions in worms 3 and 4. Worm 4 corresponds to tracking plot in Fig. 1c.

Video S3. High speed video of pharyngeal pumping in *eat-13* animals. The beginning of each animal's recording can be found at the following time points (m:s): Worm 1: 0:00; Worm 2: 0:12; Worm 3: 0:28; Worm 4: 0:41. Note the deformed pharynx in these worms. Worm 2 corresponds to tracking plot in Fig. 1d.

Video S4. High speed video of pharyngeal pumping in *eat-15* animals. The beginning of each animal's recording can be found at the following time points (m:s): Worm 1: 0:00; Worm 2: 0:16; Worm 3: 0:29; Worm 4: 0:40. Worm 3 has weak isthmus contraction. Worm 4 shows particles moving back and forth along the lumen during

relaxation, perhaps due to the influence of the terminal bulb. Worm 2 corresponds to

tracking plot in Fig. 1e.

Video S5. High speed video of pharyngeal pumping in *eat-17* animals. The

beginning of each animal's recording can be found at the following time points (m:s):

Worm 1: 0:00; Worm 2: 0:12; Worm 3: 0:27; Worm 4: 0:44. Note the deformed

pharynxes in these worms. The grinder flaps are hard to resolve, indicating a potential

defect in grinder formation. Worm 3 corresponds to tracking plot in Fig. 1f.

Video S6. High speed video of pharyngeal pumping in *eat-6* animals. The

beginning of each animal's recording can be found at the following time points (m:s):

Worm 1: 0:00; Worm 2: 0:20; Worm 3: 0:48; Worm 4: 1:12. Note that the terminal bulb

does not come back to the fully relaxed position in these worms, but remains slightly

contracted even when the other compartments have closed. Worms 2-4 also have weak

isthmus contraction. Worm 2 corresponds to tracking plot in Fig. 5c.

Video S7. High speed video of pharyngeal pumping in *egl-19* animals. The

beginning of each animal's recording can be found at the following time points (m:s):

Worm 1: 0:00; Worm 2: 0:16; Worm 3: 0:40; Worm 4: 0:58. Note the unique "yawns" of

the terminal bulb in worms 1 and 2. In those pumps, the terminal bulb remained fully

contracted for two pump cycles. Worm 3 corresponds to tracking plot in Fig. 5d.

Video S8. High speed video of pharyngeal pumping in *eat-2* animals. The beginning of each animal's recording can be found at the following time points (m:s): Worm 1: 0:00; Worm 2: 0:27; Worm 3: 0:47; Worm 4: 1:15. Note the extended delay between pump cycles in these worms compared to N2. Worm 2 corresponds to tracking plot in Fig. 5e.

References

Avery, L. and Horvitz, H.R., 1987. A cell that dies during wild-type C. elegans development can function as a neuron in a ced-3 mutant. Cell, 51(6), pp.1071-1078.

Altun, Z.F., Hall, D.H., 2003. WormAtlas Hermaphrodite Handbook - Alimentary System - Pharynx. WormAtlas. https://doi.org/10.3908/wormatlas.1.3

Avery, L., 1993a. The genetics of feeding in Caenorhabditis elegans. Genetics 133, 897–917.

Avery, L., 1993b. Motor neuron M3 controls pharyngeal muscle relaxation timing in Caenorhabditis elegans. J. Exp. Biol. 175, 283–297.

Avery, L., 1993c. Motor neuron M3 controls pharyngeal muscle relaxation timing in Caenorhabditis elegans. J. Exp. Biol. 175, 283–297.

Bhatla, N., Droste, R., Sando, S.R., Huang, A., Horvitz, H.R., 2015. Distinct Neural Circuits Control Rhythm Inhibition and Spitting by the Myogenic Pharynx of C. elegans. Curr. Biol. CB 25, 2075–2089. https://doi.org/10.1016/j.cub.2015.06.052

Davis, M.W., Somerville, D., Lee, R.Y., Lockery, S., Avery, L., Fambrough, D.M., 1995.
Mutations in the Caenorhabditis elegans Na,K-ATPase alpha-subunit gene, eat-
6, disrupt excitable cell function. J. Neurosci. Off. J. Soc. Neurosci. 15, 8408–
8418.

Fang-Yen, C., Avery, L., Samuel, A.D.T., 2009. Two size-selective mechanisms
specifically trap bacteria-sized food particles in Caenorhabditis elegans. Proc.
Natl. Acad. Sci. U. S. A. 106, 20093–20096.
https://doi.org/10.1073/pnas.0904036106

Kozlova, A.A., Lotfi, M., Okkema, P.G., 2019. Cross Talk with the GAR-3 Receptor
Contributes to Feeding Defects in Caenorhabditis elegans eat-2 Mutants.
Genetics 212, 231–243. https://doi.org/10.1534/genetics.119.302053

McKay, J.P., Raizen, D.M., Gottschalk, A., Schafer, W.R., Avery, L., 2004. eat-2 and
eat-18 are required for nicotinic neurotransmission in the Caenorhabditis elegans
pharynx. Genetics 166, 161–169.

Raizen, D.M., Lee, R.Y., Avery, L., 1995. Interacting genes required for pharyngeal
excitation by motor neuron MC in Caenorhabditis elegans. Genetics 141, 1365–
1382.

Sando, S.R., Bhatla, N., Lee, E.L., Horvitz, H.R., 2021. An hourglass circuit motif
transforms a motor program via subcellularly localized muscle calcium signaling
and contraction. eLife 10, e59341. https://doi.org/10.7554/eLife.59341

Song, B., Avery, L., 2012. Serotonin activates overall feeding by activating two separate
neural pathways in Caenorhabditis elegans. J. Neurosci. Off. J. Soc. Neurosci.
32, 1920–1931. https://doi.org/10.1523/JNEUROSCI.2064-11.2012

476   Stiernagle, T., 2006. Maintenance of C. elegans. WormBook Online Rev. C Elegans

477   Biol. 1–11. https://doi.org/10.1895/wormbook.1.101.1

478   Straud, S., Lee, I., Song, B., Avery, L., You, Y.-J., 2013. The jaw of the worm: GTPase-

479   activating protein EAT-17 regulates grinder formation in Caenorhabditis elegans.

480   Genetics 195, 115–125. https://doi.org/10.1534/genetics.113.152538