Competitive evolution of H1N1 and H3N2 influenza viruses in the United States

Chaiwat Wilasang[1], Pikkanet Suttirat[1], Anuwat Wiratsudakul[2], Sudarat Chadsuthi[3,4], Charin Modchang[1,5,6,*]

[1]Biophysics Group, Department of Physics, Faculty of Science, Mahidol University, Bangkok 10400, Thailand.

[2]Department of Clinical Sciences and Public Health, and the Monitoring and Surveillance Center for Zoonotic Diseases in Wildlife and Exotic Animals, Faculty of Veterinary Science, Mahidol University, Nakhon Pathom 73170, Thailand.

[3]Department of Physics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand.

[4]Research Center for Academic Excellence in Applied Physics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

[5]Centre of Excellence in Mathematics, CHE, Bangkok 10400, Thailand.

[6]Thailand Center of Excellence in Physics, Commission on Higher Education, Bangkok 10400, Thailand.

* Corresponding author

Email: charin.mod@mahidol.edu

Abstract

Seasonal influenza causes vast public health and economic impact globally. The prevention and control of the annual epidemics remain a challenge due to the antigenic evolution of the viruses. Here, we presented a novel modeling framework based on changes in amino acid sequences and relevant epidemiological data to retrospectively investigate the competitive evolution and transmission of H1N1 and H3N2 influenza viruses in the United States during October 2002 and April 2019. To do so, we estimated the time-varying disease transmission rate from the reported influenza cases and the time-varying evolutionary rate of the viruses from the changes in amino acid sequences. By incorporating these time-varying rates into the transmission models, we found that the models could accurately capture the evolutionary transmission dynamics of influenza viruses in the United States. Our modeling results also showed that models incorporating evolutionary change of the virus could provide better modeling performance suggesting the critical role of the evolutionary change of virus on the disease transmission.

Keywords: H1N1, H3N2, Influenza virus, Evolutionary dynamics, Transmission dynamics.

Introduction

The spread of the influenza virus is a major public health threat that leads to mortality, hospitalization, and an economic impact. Each year, there are one billion influenza cases globally, including 3 to 5 million cases of severe illness[1,2] and at least 200,000 to 600,000 recorded deaths[3-6]. In the US, seasonal influenza accounts for more than tens of thousands of deaths during annual epidemics[7]. Moreover, the economic burden of seasonal influenza in the US has been estimated at more than ten billion dollars in healthcare and social costs[7]. The influenza epidemics recur annually and are very challenging to control. These recurrent epidemics are usually governed by a complex interaction of evolutionary and transmission dynamics of influenza viruses[1,6]. A better understanding of the evolutionary transmission dynamics of the virus could provide insights to mitigate the burden of seasonal influenza.

Influenza is caused by a rapidly mutating virus, which is an enveloped negative-strand RNA virus belonging to the family of *Orthomyxoviridae*[6,8,9]. The influenza viruses are classified into types A–D, where influenza type A is the most pathogenic[8]. Among influenza type A, different viral subtypes have been continuously reported. However, the subtype A(H3N2) and A(H1N1) are commonly found during epidemics each year[1]. In the US, these two influenza subtypes are reported almost every winter, usually from October to April. Seasonality is known to affect the transmission dynamics of the influenza virus[10-14]. A change in temperature and humidity throughout the year could profoundly affect the survival of these viruses. For example, in winter, the cold and dry weather can accelerate the growth of the virus and facilitate viral transmisison[15].

Vaccination is one of the most effective interventions used to mitigate the seasonal influenza epidemics[16,17]. Nonetheless, the annual vaccination may fail to fully protect vaccinated individuals due to antigenic changes of the viruses. Influenza viruses exhibit high genetic variability in which the viruses continually undergo an antigenic change through natural selection. For instance, the changes in the surface protein, called hemagglutinin (HA), help the viruses to escape from immune attacks[6,18,19]. The primary target of immunity against influenza is the HA proteins[7,20]. By changing the properties of antigenic sites in the HA proteins, the immune system may not recognize the virus and allows it to infect host cells[6,18,20,21]. Moreover, a high mutation rate has been observed in several studies due to the short generation times and large population

sizes of the viruses [22-24]. These evolutionary processes are believed to be a major driver of seasonal
influenza transmission[20].

In the US, influenza subtype A(H3N2) emerges and spreads more frequently than
A(H1N1). Nevertheless, the two subtypes seem to be competitive for the susceptible hosts. Each
year, the subtype virus that gets the highest fitness from mutation may spread predominantly in
the population. In general, individuals infected with a single influenza strain will acquire immunity
against that certain strain[16,22]. Partial or cross-immunity to other subtypes is also observed[15,25-29].
However, the level of cross-immunity is weakening with an increasing antigenic change of the
influenza virus[2]. Therefore, co-circulation of the two influenza subtypes may result in cross-
immunity.

Mathematical models, specifically the compartmental frameworks, have been widely used
to investigate the transmission dynamics of influenza viruses. Most of the previous studies focused
on the impact of seasonality on influenza transmission dynamics[12,13,30-33]. Subsequently, the
evolution of these viruses has increasingly been studied as an important source of disease burden[6].
For instance, in 2014, J. B. Axelsen and co-workers developed a model to examine the role of
climatic drivers and viral evolution on influenza dynamics[30]. The model also integrated an
antigenic evolution process of the influenza virus by adding a new strain pulse to the year that
drastic antigenic change occurs. However, the model did not consider the evolutionary dynamics
of the influenza virus. Later, in 2018, X. Du et al. developed advances in mathematical modeling
to incorporate such evolutionary changes. Their proposed model was designed specifically for
seasonal H3N2 influenza to predict influenza incidence in the US[1]. Incorporating information on
evolutionary change into transmission models can accurately improve epidemiological forecasts.
Although such models may fit data accurately, their model does not consider the competition
between different subtypes of the virus. Indeed, the co-circulation and co-transmission of influenza
subtypes may result in different transmission characteristics[15,34].

In this study, we proposed epidemiological models for seasonal influenza A(H3N2) and
A(H1N1) viruses that incorporate the evolutionary change of the viruses. The models integrated
the data on the changes of amino acid sequences of HA proteins in epitope sites to the transmission
models. To do so, we firstly estimated the time-varying evolutionary rate of each influenza subtype
using the sequence data. We then incorporated this evolutionary rate into the transmission models

to investigate the transmission dynamics of seasonal influenza in the US. We utilized long-term

influenza surveillance data[35] from October 2002 to April 2019 in the US to fit the models. Finally,

we demonstrated that evolutionary changes of the virus could play an important role in changing

seasonal influenza transmission dynamics at the population level.

Materials and methods

Influenza surveillance data

The influenza surveillance data used in this study were obtained from the FluView of the

US Centers for Disease Control and Prevention (CDC)[35]. The US CDC collects seasonal influenza

activity data from both the US World Health Organization (WHO) Collaborating Laboratories and

National Respiratory and Enteric Virus Surveillance System (NREVSS) laboratories. The US

WHO and NREVSS laboratories report the total number of weekly respiratory specimens tested

and the number of weekly influenza-like illness (ILI) cases that tested positive for influenza types

A and B to the US CDC. Besides, the US WHO collaborating laboratories also report influenza

subtype A(H1N2) and A(H3N2). An ILI case was defined as a patient with a fever greater than 38

C° and a cough or sore throat[36,37]. Following X. Du et al.,[1] the weekly influenza cases were

computed as the product of the ILI incidence rate, the proportion of ILI samples that tested positive

for influenza, the subtype proportion, and the US population size. The population data were

downloaded from the US Census Bureau[38]. The monthly influenza cases from October 2002 to

April 2019 were obtained by aggregating the weekly data.

Influenza amino acid sequences data

Amino acid sequences of influenza A(H3N2) and A(H1N1) were downloaded from the

Global Initiative on Sharing Avian Influenza Data (GISAID)[39,40]. To analyze the HA protein

sequences, whole sequences were aligned using Multiple Sequence Comparison by Log-

Expectation (MUSCLE)[41,42]. The MUSCLE program uses FASTA files containing amino acid

sequences as an input and provides the aligned amino acid sequences. Unknown amino acids were

replaced by gaps. Aligned sequences were then edited using ClustalW, and the incomplete

sequences were manually removed[43,44]. Since the major target of immunity against influenza is

located in the epitope sites of the HA proteins[6,7,20], we used only the amino acid sequences

encoding epitope sites of these proteins to analyze the antigenic change of the virus.

Evolutionary transmission model with no competition

To investigate the evolutionary transmission dynamics of influenza, we analyzed two

transmission models that explicitly integrate the virus evolutionary dynamics (**Figure 1**). **Figure**

1(A) illustrates the evolutionary transmission model with no competition. In this model, the

population was divided into three epidemiological classes, namely, susceptible, infectious, and

recovered. The susceptible individuals (S) can be infected if they contact an infectious individual.

After being infected, susceptible individuals move to the infectious class at rate $\beta(t)I/N$, where

$\beta(t)$ is a time-dependent transmission rate, and N is the total population size. The infectious

individuals (I) recover from the disease after the infectious period ($1/\gamma$) and then are moved to

the recovered class. Individuals in the recovered class are immune to the virus and cannot be

infected. The model also integrates an effect of the virus antigenic change represented by an

evolutionary rate ($\varepsilon(t)$). We assumed that recovered individuals (R) could become susceptible

again at rate $\varepsilon(t)R$. The evolutionary transmission dynamics of influenza are described by the

following ordinary differential equations (ODEs):

$$\frac{dS}{dt} = \mu N - \beta(t)S\left(\frac{I}{N}\right) + \varepsilon(t)R - \mu S, \tag{1}$$

$$\frac{dI}{dt} = \beta(t)S\left(\frac{I}{N}\right) - \gamma I - \mu I, \tag{2}$$

$$\frac{dR}{dt} = \gamma I - \varepsilon(t)R - \mu R, \tag{3}$$

$$\frac{dC}{dt} = \beta(t)S\left(\frac{I}{N}\right), \tag{4}$$

149   where S, I, R, and C represent the number of susceptible, infectious, recovered, and cumulative

150   infected individuals, respectively. In this model, we assumed that the newborns are susceptible,

151   and the number of natural births and deaths are balanced at a rate μ so that the total population size

152 is constant. The parameters descriptions and the values used in the model are summarized in **Table**
153 **1**.

Figure 1. Schematic of the evolutionary transmission model. (A) The flow diagram of the evolutionary transmission model of the A(H1N1) and A(H3N2) influenza viruses with no competition and **(B)** with the selective competition.

161 **Table 1. Model parameters and their values.**

Parameter	Definition	Value	References
$\beta_i(t)$	Monthly transmission rate of influenza subtype i	[0.001-10]	Estimated
γ	Recovery rate	1/5 per day	[45-47]
μ	Natural birth and death rate	1/67 per year	[3]
ε_0	Waning rate of immunity	1/20 per year	Estimated
Ψ	Cross-immunity induced by primary infection	0.7	[15,48,49]
α	The steepness of antigenic change	2	Assumed
N	Population size	3.25×10^8	[38]

Evolutionary transmission model with the selective competition

Several influenza virus strains can co-circulate in a population leading to selective
competition in susceptible hosts[7,22,25,50,51]. The co-circulation of influenza viruses could result in
cross-immunity and mediate the ecological competition between different strains. Consequently,
for the modeling pursued here, we constructed an evolutionary transmission model considering
cross-immunity and selective competition of the A(H3N2) and A(H1N1) influenza viruses. The
schematic for the model with selective competition is shown in **Figure 1(B)**. Similar to the model
with no competition, the population was categorized into three different groups, namely. However,
each group of individuals here can be infected with either A(H1N1) or A(H3N2), indicated by the
subscripts H1 and H3, respectively. Individuals recently recovered from the infection of one stain
develop partial protection and hence are less susceptible to another strain. In our equations, we use
the order of subscripts to indicate the order of infection. For example, I_{H3H1} represents a group of
individuals who had recently recovered from the infection of A(H3N2) but is currently infected
with A(H1N1). R_d represents a group of individuals that are immune to the two virus strains. The
evolutionary transmission model with selective competition can be described by the following
system of ODEs:

$\frac{dS}{dt} = \mu N - \beta_{H1}(t)S(t)(I_{H1} + I_{H3H1})/N - \beta_{H3}(t)S(I_{H3} + I_{H1H3}) + \varepsilon_{H1}(t)R_{H1} + \varepsilon_{H3}(t)R_{H3} - \mu S,$ (5)

$$181 \qquad \frac{dI_{H1}}{dt} = \beta_{H1}(t)S(I_{H1} + I_{H3H1})/N - \gamma I_{H1} - \mu I_{H1}, \qquad (6)$$

$$182 \qquad \frac{dI_{H3}}{dt} = \beta_{H3}(t)S(I_{H3} + I_{H1H3})/N - \gamma I_{H3} - \mu I_{H3}, \qquad (7)$$

$$183 \qquad \frac{dR_{H1}}{dt} = \gamma I_{H1} - \varepsilon_{H1}(t)R_{H1} - \beta_{H3}(t)(1-\Psi)R_{H1}(I_{H3} + I_{H1H3})/N + \varepsilon_{H3}(t)R_d - \mu R_{H1}, \quad (8)$$

$$184 \qquad \frac{dR_{H3}}{dt} = \gamma I_{H3} - \varepsilon_{H3}(t)R_{H3} - \beta_{H1}(t)(1-\Psi)R_{H3}(I_{H1} + I_{H3H1})/N + \varepsilon_{H1}(t)R_d - \mu R_{H3}, \quad (9)$$

$$185 \qquad \frac{dI_{H1H3}}{dt} = \beta_{H3}(t)(1-\Psi)R_{H1}(I_{H3} + I_{H1H3})/N - \gamma I_{H1H3} - \mu I_{H1H3}, \qquad (10)$$

$$186 \qquad \frac{dI_{H3H1}}{dt} = \beta_{H1}(t)(1-\Psi)R_{H3}(I_{H1} + I_{H3H1})/N - \gamma I_{H3H1} - \mu I_{H3H1}, \qquad (11)$$

$$187 \qquad \frac{dR_d}{dt} = \gamma I_{H1H3} + \gamma I_{H3H1} - \varepsilon_{H3}(t)R_d - \varepsilon_{H1}(t)R_d - \mu R_d, \qquad (12)$$

where $\beta_{H1}(t)$ and $\beta_{H3}(t)$ is the time-varying transmission rate of infectious individuals who are
currently infected with A(H1N1) and A(H3N2), respectively. Ψ is the cross-immunity between the
two influenza subtypes. ε_{H1} and ε_{H3} are the evolutionary rate of the A(H1N1) and A(H3N2) strain,
respectively.

To generate the long-term seasonal influenza dynamics, the infection was initiated by
randomly selected seed infections and run for a period of 100 years to allow the system to reach
equilibrium. We then used the averaged population in each compartment as the initial population
of the models. For the simulation process, we used the Euler method to solve the sets of ODEs
implemented in MATLAB R2019b. The model parameters are summarized in **Table 1**.

198 Estimating the time-varying evolutionary rate

Seasonal influenza viruses can better survive and spread in a specific environment,
especially in winter[30,52]. The recurrence of seasonal influenza epidemics thus frequently occurs
between October and March. To consider the effects of seasonality, the period from October 1 to
March 31 was assigned as the high transmission season (HTS), while the remaining period was
the low transmission season (LTS)[1]. We compared the amino acid sequences of both influenza
subtypes between two consecutive seasons to estimate the evolutionary rate.

To investigate the evolutionary dynamics of the seasonal influenza virus, we used amino
acid changes in the epitope as a measure of the antigenic distance between the two strains. Once
the virus infects the host cells, the antibody attacks the antigen by binding to HA proteins located
on the viral surface[18]. This part of an antigen memorized by the immune system is called
epitope[3,53]. The mutation within epitope regions could, thus, weaken the immune system[19].
Consequently, the virus can enter and infect the host cell. Epitope regions are categorized
differently according to influenza virus subtypes[6,54-57]. The amino acid residues of epitope regions
are summarized in **Table S1** and **Table S2** in the Supplementary Information.

To estimate the antigenic change among influenza viruses, we considered aligned amino
acid sequences as character strings and computed the number of amino acid differences within all
five epitopes, known as the Hamming distance[58]. The Hamming distances were calculated by
comparing the viral sequences in the current and previous seasons. However, as the numbers of
viral sequences recorded in each season are different, we first sampled 10^6 random pairs of
sequences between seasons to avoid sampling biases. Subsequently, we calculated the Hamming
distance for each pair of amino acid sequences. We then computed $P_{all-epitopes}$, which is a fraction
of the number of amino acid differences within all five epitopes and the total number of amino
acids presented[59].

We incorporated the evolutionary rate into the transmission model to investigate how
antigenic change affects transmission dynamics. The time-varying evolutionary rate, $\varepsilon(t)$, is given
by:

$$\varepsilon(t) = \varepsilon_0 + c \left[\frac{P_{all-epitope}}{1 - P_{all-epitope}} \right]^{\alpha}, \tag{13}$$

where ε_0 represents the naturally waning rate of immunity. Parameters α and c were estimated
such that the resulting cumulative cases derived from the simulations match the influenza
surveillance data. We varied the evolutionary rate and simulated the H1N1 influenza model before
the pandemic season of 2009–2010. The values of $\varepsilon(t)$ that the model can search for are uniformly
distributed in the interval [0, 1] with 0.001 resolution. We recorded only the values of the
evolutionary rate that yield the best simulation results. Subsequently, we calculated the constant
parameter c according to Eq. (13). Simultaneously, we varied α; the value of steepness changes.

The estimated parameters are $\alpha = 2$ and $c = 5.9911 \times 10^{-2}$ as they minimize the difference between

simulated and reported data.

Estimating the time-varying transmission rate

In this study, we employed the parameter estimation method developed by P. Suparit et

al.[60] to estimate the time series of monthly transmission rates. The monthly influenza cases in the

US starting from October 2002 to April 2019 were used to estimate the time series of monthly

transmission rates. Briefly, we divided the model simulation into several consecutive monthly time

window simulations. We varied $\beta(t)$ ranging from 0.001 to 10 with the resolution of 0.01 and

searched for the transmission rate that provides the best fit between the number of monthly

simulated cases (C_{sim}) and the number of monthly reported cases (C_{rep}). The values of the monthly

transmission rates that minimizes $|C_{sim} - C_{rep}|$ were recorded in the time series of monthly

transmission rates. We also recorded the number of individuals in each epidemiological class for

the current monthly time window simulation. Subsequently, we used the number of individuals in

the previous monthly time window as an initial condition for the next monthly time window

simulation. The process was repeated until the complete time series of the monthly transmission

rates were obtained.

Results

Seasonal patterns of influenza epidemics in the US

The monthly influenza cases from October 2002 to April 2019 (~17 years) reported in the

United States are shown in **Figure 2** (see also **Figure S1**). The data shows the incidence of monthly

influenza of two subtypes: subtype A(H3N2) and subtype A(H1N1). The outbreaks of influenza

A (H3N2) and A (H1N1) usually appear between October and March. The numbers of cases are

usually lower in the summer seasons (April to September). We, therefore, classified the period

from 1st October to 31st March as a high transmission season (HTS) and the period from 1st April

to 30th September as a low transmission season (LTS). Thus, there were 17 HTSs and 16 LTSs in

the time series. As illustrated in **Figure 2**, both subtypes of influenza viruses have co-circulated

throughout the study period. Aside from the apparent annual cycles, epidemics of A(H3N2) and
A(H1N1) usually occur with different magnitudes, especially during the year 2009 when the novel
2009 A(H1N1) influenza emerged [47,61,62].

Figure 2. Influenza epidemics in the US. The monthly cases of influenza A(H1N1) and A(H3N2)
in the US between October 2002 and April 2019.

Evolutionary rates of influenza virus

The recurrent influenza epidemics are a combined outcome of viral antigenic changes,
interactions among co-circulating influenza viruses, transmission, and host immune recognition
against the virus[6]. When influenza viruses undergo evolution, the immune system may not be able
to recognize the viruses, allowing the viruses to infect host cells. Thus, previously infected
individuals may eventually become susceptible to the viruses again[19]. To quantify the evolutionary
rate, we calculated $P_{all-epitopes}$, which is a measure of the antigenic change between two influenza
virus strains[4,59] (see Material and Methods for more details). The average $P_{all-epitopes}$ for both
influenza A(H3N2) and A(H1N1) are shown in **Figure 3 (A)** and **(B)**, respectively. $P_{all-epitopes}$
were also compared with the monthly influenza cases. According to the results shown in **Figure**

3(A) and **(B)**, we found that the large epidemic curves usually occur after a significant evolutionary change of influenza viruses. Specifically, the antigenic change of influenza A(H1N1) was around 80% in the 2008-2009 HTS, which resulted in the subsequent surge of the 2009 influenza pandemic (**Figure 3(B)**). This result suggested that $P_{all-epitopes}$ might be a potential early warning for a pandemic. We also found that the average $P_{all-epitopes}$ of A(H3N2) is less fluctuate than that of A(H1N1). The corresponding evolutionary rates of influenza A(H1N1) and A(H3N2) are illustrated in **Figure 3(C)** and **(D)**.

Figure 3. Average monthly $P_{all-epitopes}$ and evolutionary rates. $P_{all-epitopes}$ of influenza **(A)** A(H3N2) virus, and **(B)** influenza A(H1N1) virus, **(C)** the evolutionary rate of influenza A(H3N2) virus, and **(D)** influenza A(H1N1) virus.

Seasonal influenza transmission dynamics in the US

We found that the trends of monthly influenza cases generated from the model agree well with the surveillance data with R-square = 0.9989 (**Figure 4**). During the pandemic season 2008-2009, for example, the total number of influenza cases was predicted at 39.03 million, which is close to the number of reported cases at 38.3 million. We also employed the model to reconstruct

297 the transmission dynamics of influenza A(H3N2) during the same period in the US. We found that

298 the model also provided a good fit to the observed data with R-square = 0.9989 (**Figure S2**).

300 **Figure 4. The influenza A(H1N1) transmission dynamics**. The model provided a good fit to the

301 16-year observed data with R-square = 0.9989.

Effect of the evolutionary change of the influenza virus

We found that the estimated transmission rates from the two models are different. The

estimated monthly transmission rates that provide the best fit of the two models are shown in

Figure 5. The results illustrate that the transmission rates estimated from the model without the

evolutionary dynamics increased after the major changes of influenza viruses in 2009. In contrast,

there is no change in the post-2009 trend of the transmission rates estimated from the model

incorporating the evolutionary dynamics. The transmission rate fluctuates between 0.11 and 0.31

per month during the years 2009 and 2019. There is also an approximately 2-fold increase in

oscillation amplitude of transmission rates when excluding the evolutionary dynamics in the

model. These suggested that the evolutionary change of the virus could affect their transmission

dynamics. This effect was also observed in the transmission dynamics of the influenza A(H3N2)

virus (see **Figure S3** in the supplementary results).

Figure 5. Comparison of H1N1 monthly transmission rates. The time series of monthly

transmission rates during October 2002 and April 2019. The blue line shows the monthly

transmission rate of the A(H1N1) influenza virus estimated using the model without evolutionary

change. Redline denotes the estimated monthly transmission rate of the A(H1N1) influenza virus

from the model with the evolutionary change of the virus.

Competitive evolutionary transmission dynamics of the A(H3N2) and A(H1N1) influenza viruses

The infection of a certain influenza subtype in humans results in partial immunity to other

subtypes of influenza viruses[22,63]. The empirical evidence suggests that the co-circulation of

influenza viruses not only results in cross-immunity but also appears to mediate the ecological

competition between the subtypes[7,22,25,50,51]. A competitive model was then constructed to

investigate the combined co-evolution and co-transmission dynamics of the A(H3N2) and

A(H1N1) influenza viruses. We also incorporated the cross-immunity between the two subtypes

into our competitive model. The results illustrated that the models could faithfully reconstruct the

influenza epidemics (**Figure S4** and **Figure S5** in the supplementary results). Nevertheless, the

competitive model that incorporates the cross-immunity is slightly more accurate in our model

validation. For the competitive model incorporating the cross-immunity, the model fit correlated

to the data with R-square = 0.9849 for H3N2 influenza cases, and it is 0.9468 in the case of H1N1

influenza cases. For the competitive model without the cross-immunity, we observed a correlation

of R-square = 0.9723 and R-square = 0.8905 for H3N2 and H1N1 influenza viruses, respectively.

The correlations were driven down partly due to the pandemic season of 2009-2010, in which

antigenic evolution of the A(H1N1) influenza virus occurred.

We then further investigated the effects of cross-immunity between the two subtypes. The

strength of cross-immunity between the two influenza subtypes is represented by the parameter Ψ.

If $\Psi = 0$, the influenza subtypes are completely different antigenically. In contrast, if $\Psi = 1$, the

two subtypes are antigenically indistinguishable[25]. **Figure 6** illustrates the effect of the cross-

immunity on the transmission rates estimated by the three models. The transmission rate estimated

from the competitive model with the cross-immunity $\Psi = 0.7$ was found in the range of $0.01–1.60$

per month with an average value of 0.59. In contrast, the transmission rates estimated from the

competitive model without cross-immunity ($\Psi = 0.0$) fluctuates in the range of $0.01-0.60$ with the

average value of 0.20 per month, which is close to the transmission rates estimated from the single

strain model (H1N1 model).

Figure 6. The time series of transmission rate estimated from three different models. The red

line shows the monthly transmission rate of the H1N1 influenza virus estimated from the model

without evolutionary change of the virus. The blue and green lines display the monthly

transmission rate of the H1N1 influenza virus estimated from the competitive model without cross-

immunity ($\Psi = 0.0$) and with the cross-immunity $\Psi = 0.7$, respectively.

Comparing the proportion of immune individuals predicted by each model, we found the

fraction of the immune population obtained from the non-competitive model is lower than that of

the competitive one (**Figure 7** and **Figure S6(B)** in the supplementary results). Noticeably, at the

end of 2009, the immunity levels of influenza A(H1N1) from the three modeling scenarios were

predicted at 7.94%, 12.82%, and 20.10%, respectively. Regarding the seroprevalence of influenza-

specific antibodies, we found that the competitive model with the cross-immunity provides the

proportion of immune population closest to the observed data at 21.0%[64,65].

Figure 7. Immunity level of influenza A(H1N1). The immunity levels of influenza A(H1N1)

estimated from the H1N1 model, the competitive model without cross-immunity, and the

competitive model with Ψ=0.7 are 7.94%, 12.82%, and 20.10%, respectively. The seroprevalence

of influenza observed by S. M. Zimmer et al.[64] was approximately 21.0%.

Discussion

Influenza epidemics have been observed annually, resulting in significant morbidity and mortality rates. The recurrence of the influenza epidemic stems from several possibilities[6]. Seasonal factors are known to affect the transmission dynamics of the influenza virus[10-14]. However, one of the most significant factors that drive the annual influenza epidemic is the evolution of the virus[1,19,58]. Thus, the integration of the evolutionary change of the virus into the epidemic model is necessary. To capture the transmission dynamics of the seasonal influenza virus in the US, we proposed mathematical models incorporating the evolutionary change of the virus. The models were employed to reconstruct the long-term influenza epidemics in the US from October 2002 to April 2019. We also compared competitive evolutionary transmission models of influenza A(H1N1) and A(H3N2) viruses with and without cross-immunity.

To investigate the interplay between the epidemic dynamics and the evolutionary dynamics of the influenza virus, we compared the virus evolutionary rate with the monthly influenza incidence (**Figure 3**). The evolutionary rate of influenza A(H3N2) increased a little during the season 2013 (October 2012 to April 2013), which might contribute to the large outbreak in the US in that season, resulting in approximately 30 million influenza illnesses[66]. For influenza A(H1N1), the evolutionary rate drastically increased during October 2008 and April 2009, consequently causing the 2009 A(H1N1) influenza pandemic[67]. Over this period, we observed that the evolutionary rate of influenza A(H1N1) considerably increased from 0.01 to 0.80. Thus, the emerging strain could replace the existing stains rapidly[68].

Several studies have examined the impacts of seasonal factors (e.g., temperature and humidity) on the transmission of seasonal influenza[11-14,25]. Few studies have attempted to bridge the gap between the evolutionary change of the virus and the transmission model[1]. Previously, this was modeled by adding a constant rate to the year that drastic antigenic change occurs to represent the effect of the influenza virus evolution[30]. However, in this work, we intended to link antigenic changes of the influenza virus to disease transmission models. Specifically, we investigated the effect of evolutionary change on seasonal influenza epidemics by incorporating evolutionary rate into the transmission model. We also validated the models by comparing the modeling results with the reported data. The model-fitting results show that the proposed influenza transmission model incorporating evolutionary change provides the best fit between the simulation results and the

reported data (R-square=0.9989) (**Figure 5**). Also, our models could reconstruct the observed

epidemic dynamics for both high transmission and low transmission seasons for the two subtypes.

Furthermore, the evolutionary rate included in the model can describe the transmission

dynamics of seasonal influenza. To interpret these findings, we compared the monthly

transmission rates estimated from the model with and without evolutionary change of the influenza

virus. We then observed significant differences in the transmission rate. The transmission rate

estimating from the model incorporating evolutionary rate fluctuates around a certain constant

even though there is the drastic change in the genetic sequence of the influenza virus in 2009. In

contrast, in the model without evolutionary rate, the fluctuation of the estimated transmission rate

is shifted up after the year of pandemics. The increase in transmission rate may result from the

large outbreaks in 2009, which drastically depleted the susceptible pool. Consequently, the model

compensates by increasing the transmission rate to generate the same number of reported cases as

a model incorporating evolutionary change. This means that either human behavior or the viral

characteristic has abruptly changed. However, there is no obvious evidence that those factors

occurred, unlike the current COVID-19 pandemic[69,70]. Since the transmission rate depends on

human behavior, it is unlikely to change after the 2009 pandemic. These results highlight the

importance of incorporating the evolutionary rate into the model. The models incorporating

evolutionary information of seasonal influenza virus are significantly better than the model that

does not. Thus, to accurately reconstruct the dynamics of seasonal influenza transmission, the

evolutionary change of the virus should be considered in transmission models.

Indeed, people can be infected by multiple strains of the influenza virus. We, therefore,

constructed competitive transmission models to investigate the combined co-evolution and co-

transmission dynamics of the A(H3N2) and A(H1N1) influenza viruses. Our findings indicate that

population susceptibility is associated with the evolutionary change of the virus and the strength

of cross-immunity. The evolutionary change of the virus replenishes the pool of susceptible

individuals, which increases population susceptibility, whereas cross-immunity can reduce

population susceptibility[71]. The antigenic variation of the influenza virus may allow the virus to

escape the existing protective immunity and cause new epidemics. This evolutionary change can

accommodate the system dynamics, e.g., increased population susceptibility, and allows abrupt

increases in population susceptibility in response to epidemic surges. In addition, we also found

that there was a competition between influenza subtypes via cross-immunity. Different values of

estimated transmission in **Figure 6** illustrate the effect of cross-immunity. These indicate that

cross-immunity could modulate the ecological interactions between co-circulating influenza

viruses by increasing the transmission rate of the virus. If the cross-immunity level is high, the

competition between the two subtypes becomes extensively strict[25]. This finding agrees with the

conclusions from other studies[2,25,50,51], which found that cross-immunity among influenza viruses

would naturally lead to competition between subtypes. Besides, we also validated our competitive

transmission models by comparing the level of immunity from model predictions with

seroprevalence data. The competitive transmission model with cross-immunity yields the best

simulation result. The model predicted the immunity level at the end of 2009 to be 20.10%, which

is similar to other studies[64,65,72-74].

Our models made several simplifying assumptions: firstly, we assumed the population is

homogeneous mixing, in which all individuals have an equal probability of transmitting the

disease. Indeed, as observed in other works[75], models with a homogeneously mixed population

usually present a higher attack rate than heterogeneous models, even for the same R_0. Second, we

ignore heterogeneity in disease transmissions such as sex, age structure, and geographical region.

The individual heterogeneity in infectiousness, e.g., superspreading events, is known to affect the

dynamics of disease transmission[76,77]. Moreover, individual heterogeneity in disease transmission

can also result in a higher probability of stochastic extinction[77]. Finally, we did not consider human

mobility in our model; incorporating these data into the model could improve a more realistic

model[78,79]. It is worth stressing that the knowledge of the evolutionary change of the influenza

virus is necessary to be investigated. Furthermore, it could also inform vaccine strain selection to

prevent yearly epidemics[80,81].

In summary, we constructed evolutionary transmission models for investigating seasonal

influenza transmission in the US. The models integrate the changes of amino acid sequences of

HA proteins in epitope sites and the time-varying disease transmission. We found that the models

with information on evolutionary change outperformed the models without it. We also constructed

the competitive model to investigate the competitive evolutionary transmission dynamics of the

A(H3N2) and A(H1N1) influenza viruses. Our result suggested that the competitive model with

cross-immunity is better at capturing the transmission dynamics of seasonal influenza.

Acknowledgments

Chaiwat Wilasang is supported by the Science Achievement Scholarship of Thailand (SAST).

References

Du, X., King, A. A., Woods, R. J. & Pascual, M. Evolution-informed forecasting of seasonal influenza A (H3N2). *Science translational medicine* **9** (2017).

Yang, W., Lau, E. H. & Cowling, B. J. Dynamic interactions of influenza viruses in Hong Kong during 1998-2018. *PLoS computational biology* **16**, e1007989 (2020).

Xu, R. *et al.* Structural basis of preexisting immunity to the 2009 H1N1 pandemic influenza virus. *Science* **328**, 357-360 (2010).

Gupta, V., Earl, D. J. & Deem, M. W. Quantifying influenza vaccine efficacy and antigenic distance. *Vaccine* **24**, 3881-3888 (2006).

Hill, E. M., Petrou, S., De Lusignan, S., Yonova, I. & Keeling, M. J. Seasonal influenza: Modelling approaches to capture immunity propagation. *PLoS computational biology* **15**, e1007096 (2019).

Petrova, V. N. & Russell, C. A. The evolution of seasonal influenza viruses. *Nature Reviews Microbiology* **16**, 47 (2018).

Zinder, D., Bedford, T., Gupta, S. & Pascual, M. The roles of competition and mutation in shaping antigenic and genetic diversity in influenza. *PLoS Pathog* **9**, e1003104 (2013).

Long, J. S., Mistry, B., Haslam, S. M. & Barclay, W. S. Host and viral determinants of influenza A virus species specificity. *Nature Reviews Microbiology* **17**, 67-81 (2019).

Maclachlan, N. J. & Dubovi, E. J. *Fenner's veterinary virology*. (Academic press, 2010).

Tamerius, J. D. *et al.* Environmental predictors of seasonal influenza epidemics across temperate and tropical climates. *PLoS pathogens* **9**, e1003194 (2013).

Steel, J., Palese, P. & Lowen, A. C. Transmission of a 2009 pandemic influenza virus shows a sensitivity to temperature and humidity similar to that of an H3N2 seasonal strain. *Journal of virology* **85**, 1400-1402 (2011).

12   Lowen, A. C., Mubareka, S., Steel, J. & Palese, P. Influenza virus transmission is
dependent on relative humidity and temperature. *PLoS Pathog* **3**, e151 (2007).
13   Lowen, A. C. & Steel, J. Roles of humidity and temperature in shaping influenza
seasonality. *Journal of virology* **88**, 7692-7695 (2014).
14   Chadsuthi, S., Iamsirithaworn, S., Triampo, W. & Modchang, C. Modeling seasonal
influenza transmission and its association with climate factors in Thailand using time-series
and ARIMAX analyses. *Computational and mathematical methods in medicine* **2015**
(2015).
15   Zhang, X.-S. Strain interactions as a mechanism for dominant strain alternation and
incidence oscillation in infectious diseases: seasonal influenza as a case study. *PloS one*
10, e0142170 (2015).
16   Ahn, K. W., Kosoy, M. & Chan, K.-S. An approach for modeling cross-immunity of two
strains, with application to variants of Bartonella in terms of genetic similarity. *Epidemics*
7, 7-12 (2014).
17   Weir, J. P. & Gruber, M. F. An overview of the regulation of influenza vaccines in the
United States. *Influenza and other respiratory viruses* **10**, 354-360 (2016).
18   Wiley, D., Wilson, I. & Skehel, J. Structural identification of the antibody-binding sites of
Hong Kong influenza haemagglutinin and their involvement in antigenic variation. *Nature*
289, 373 (1981).
19   Ferguson, N. M., Galvani, A. P. & Bush, R. M. Ecological and immunological
determinants of influenza evolution. *Nature* **422**, 428 (2003).
20   Huang, J.-W. & Yang, J.-M. Changed epitopes drive the antigenic drift for influenza A
(H3N2) viruses. *BMC bioinformatics* **12**, 1-10 (2011).
21   van de Sandt, C. E. *et al.* Differential recognition of influenza a viruses by M158–66
epitope-specific CD8+ T cells is determined by extraepitopic amino acid residues. *Journal*
of virology **90**, 1009-1022 (2016).
22   Zhang, X.-S. *et al.* Co-circulation of influenza A virus strains and emergence of pandemic
via reassortment: the role of cross-immunity. *Epidemics* **5**, 20-33 (2013).
23   Stern, A. & Andino, R. in *Viral Pathogenesis*   233-240 (Elsevier, 2016).
24   Peck, K. M. & Lauring, A. S. Complexities of viral mutation rates. *Journal of virology* **92**
(2018).

Kamo, M. & Sasaki, A. The effect of cross-immunity and seasonal forcing in a multi-strain epidemic model. *Physica D: Nonlinear Phenomena* **165**, 228-241 (2002).

Laurie, K. L. *et al.* Multiple infections with seasonal influenza A virus induce cross-protective immunity against A (H1N1) pandemic influenza virus in a ferret model. *The Journal of infectious diseases* **202**, 1011-1020 (2010).

Barry, J. M., Viboud, C. & Simonsen, L. Cross-protection between successive waves of the 1918–1919 influenza pandemic: epidemiological evidence from US Army camps and from Britain. *The Journal of infectious diseases* **198**, 1427-1434 (2008).

Epstein, S. L. & Price, G. E. Cross-protective immunity to influenza A viruses. *Expert review of vaccines* **9**, 1325-1341 (2010).

Bodewes, R. *et al.* Infection of the upper respiratory tract with seasonal influenza A (H3N2) virus induces protective immunity in ferrets against infection with A (H1N1) pdm09 virus after intranasal, but not intratracheal, inoculation. *Journal of virology* **87**, 4293-4301 (2013).

Axelsen, J. B., Yaari, R., Grenfell, B. T. & Stone, L. Multiannual forecasting of seasonal influenza dynamics reveals climatic and evolutionary drivers. *Proceedings of the National Academy of Sciences* **111**, 9538-9542 (2014).

Yaari, R., Katriel, G., Huppert, A., Axelsen, J. & Stone, L. Modelling seasonal influenza: the role of weather and punctuated antigenic drift. *Journal of The Royal Society Interface* **10**, 20130298 (2013).

Chowell, G., Miller, M. & Viboud, C. Seasonal influenza in the United States, France, and Australia: transmission and prospects for control. *Epidemiology & Infection* **136**, 852-864 (2008).

Shaman, J. & Karspeck, A. Forecasting seasonal outbreaks of influenza. *Proceedings of the National Academy of Sciences* **109**, 20425-20430 (2012).

Zhang, X.-S. & De Angelis, D. Construction of the influenza A virus transmission tree in a college-based population: co-transmission and interactions between influenza A viruses. *BMC infectious diseases* **16**, 1-8 (2015).

(CDC), C. f. D. C. a. P. *FluView*, <https://gis.cdc.gov/grasp/fluview/fluportaldashboard.html> (2019).

36 Fitzner, J. *et al.* Revision of clinical case definitions: influenza-like illness and severe acute
respiratory infection. *Bulletin of the World Health Organization* **96**, 122 (2018).

37 Kasper, M. R., Wierzba, T. F., Sovann, L., Blair, P. J. & Putnam, S. D. Evaluation of an
influenza-like illness case definition in the diagnosis of influenza among patients with
acute febrile illness in Cambodia. *BMC infectious diseases* **10**, 1-5 (2010).

38 Bureau, U. S. C. *Population and housing unit estimates*,
<https://www.census.gov/programs-surveys/popest.html> (2020).

39 Shu, Y. & McCauley, J. GISAID: Global initiative on sharing all influenza data–from
vision to reality. *Eurosurveillance* **22**, 30494 (2017).

40 Bogner, P., Capua, I., Lipman, D. J. & Cox, N. J. A global initiative on sharing avian flu
data. *Nature* **442**, 981-981 (2006).

41 Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic acids research* **32**, 1792-1797 (2004).

42 Edgar, R. C. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity. *BMC bioinformatics* **5**, 1-19 (2004).

43 Thompson, J. D., Gibson, T. J. & Higgins, D. G. Multiple sequence alignment using
ClustalW and ClustalX. *Current protocols in bioinformatics*, 2.3. 1-2.3. 22 (2003).

44 Larkin, M. A. *et al.* Clustal W and Clustal X version 2.0. *bioinformatics* **23**, 2947-2948
(2007).

45 Carrat, F. *et al.* Time lines of infection and disease in human influenza: a review of
volunteer challenge studies. *American journal of epidemiology* **167**, 775-785 (2008).

46 Nsoesie, E. O., Beckman, R. J. & Marathe, M. V. Sensitivity analysis of an individual-
based model for simulation of influenza epidemics. *PloS one* **7**, e45414 (2012).

47 Modchang, C., Iamsirithaworn, S., Auewarakul, P. & Triampo, W. A modeling study of
school closure to reduce influenza transmission: A case study of an influenza A (H1N1)
outbreak in a private Thai school. *Mathematical and Computer Modelling* **55**, 1021-1033
(2012).

48 Truscott, J. *et al.* Essential epidemiological mechanisms underpinning the transmission
dynamics of seasonal influenza. *Journal of The Royal Society Interface* **9**, 304-312 (2012).

49   Furuse, Y. & Oshitani, H. Mechanisms of replacement of circulating viruses by seasonal
and pandemic influenza A viruses. *International Journal of Infectious Diseases* **51**, 6-14
(2016).

50   Latorre-Margalef, N. *et al.* Competition between influenza A virus subtypes through
heterosubtypic immunity modulates re-infection and antibody dynamics in the mallard
duck. *PLoS pathogens* **13**, e1006419 (2017).

51   Andreasen, V., Lin, J. & Levin, S. A. The dynamics of cocirculating influenza strains
conferring partial cross-immunity. *Journal of mathematical biology* **35**, 825-842 (1997).

52   Tang, J. W. The effect of environmental parameters on the survival of airborne infectious
agents. *Journal of the Royal Society Interface* **6**, S737-S746 (2009).

53   Holmes, E. C. *et al.* Whole-genome analysis of human influenza A virus reveals multiple
persistent lineages and reassortment among recent H3N2 viruses. *PLoS biology* **3**, e300
(2005).

54   Skowronski, D. M. *et al.* Low 2012–13 influenza vaccine effectiveness associated with
mutation in the egg-adapted H3N2 vaccine strain not antigenic drift in circulating viruses.
PLoS one **9**, e92153 (2014).

55   Munoz, E. T. & Deem, M. W. Epitope analysis for influenza vaccine design. *Vaccine* **23**,
1144-1148 (2005).

56   Skowronski, D. M. *et al.* Influenza A/subtype and B/lineage effectiveness estimates for the
2011–2012 trivalent vaccine: cross-season and cross-lineage protection with unchanged
vaccine. *The Journal of infectious diseases* **210**, 126-137 (2014).

57   Skowronski, D. M. *et al.* A sentinel platform to evaluate influenza vaccine effectiveness
and new variant circulation, Canada 2010–2011 season. *Clinical infectious diseases* **55**,
332-342 (2012).

58   Klein, E. Y., Serohijos, A. W., Choi, J.-M., Shakhnovich, E. I. & Pekosz, A. Influenza A
H1N1 pandemic strain evolution–divergence and the potential for antigenic drift variants.
PLoS one **9**, e93632 (2014).

59   Pan, K., Subieta, K. C. & Deem, M. W. A novel sequence-based antigenic distance measure
for H1N1, with application to vaccine effectiveness and the selection of vaccine strains.
Protein Engineering, Design & Selection **24**, 291-299 (2011).

60 Suparit, P., Wiratsudakul, A. & Modchang, C. A mathematical model for Zika virus transmission dynamics with a time-dependent mosquito biting rate. *Theoretical Biology and Medical Modelling* **15**, 1-11 (2018).

61 Shrestha, S. S. *et al.* Estimating the burden of 2009 pandemic influenza A (H1N1) in the United States (April 2009–April 2010). *Clinical Infectious Diseases* **52**, S75-S82 (2011).

62 Fraser, C. *et al.* Pandemic potential of a strain of influenza A (H1N1): early findings. *science* **324**, 1557-1561 (2009).

63 Chung, K. & Lui, R. Dynamics of two-strain influenza model with cross-immunity and no quarantine class. *Journal of mathematical biology* **73**, 1467-1489 (2016).

64 Zimmer, S. M. *et al.* Seroprevalence following the second wave of Pandemic 2009 H1N1 influenza in Pittsburgh, PA, USA. *PLoS one* **5**, e11601 (2010).

65 Reed, C., Katz, J. M., Hancock, K., Balish, A. & Fry, A. M. Prevalence of seropositivity to pandemic influenza A/H1N1 virus in the United States following the 2009 pandemic. *PloS one* **7**, e48187 (2012).

66 Control, C. f. D. & Prevention. Update: influenza activity-United States, 2011-12 season and composition of the 2012-13 influenza vaccine. *MMWR. Morbidity and mortality weekly report* **61**, 414-420 (2012).

67 Control, C. f. D. & Prevention. Update: influenza activity-United States, 2009-10 season. *MMWR. Morbidity and mortality weekly report* **59**, 901-908 (2010).

68 Felsenstein, J. The evolutionary advantage of recombination. *Genetics* **78**, 737-756 (1974).

69 Jarvis, C. I. *et al.* Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. *BMC medicine* **18**, 1-10 (2020).

70 Schlosser, F. *et al.* COVID-19 lockdown induces disease-mitigating structural changes in mobility networks. *Proceedings of the National Academy of Sciences* **117**, 32883-32890 (2020).

71 Sonoguchi, T., Naito, H., Hara, M., Takeuchi, Y. & Fukumi, H. Cross-subtype protection in humans during sequential, overlapping, and/or concurrent epidemics caused by H3N2 and H1N1 influenza viruses. *Journal of Infectious Diseases* **151**, 81-88 (1985).

72 Achonu, C. *et al.* Seroprevalence of pandemic influenza H1N1 in Ontario from January 2009–May 2010. *PLoS one* **6**, e26427 (2011).

73 Broberg, E., Nicoll, A. & Amato-Gauci, A. Seroprevalence to influenza A (H1N1) 2009
virus—where are we? *Clinical and vaccine immunology* **18**, 1205-1212 (2011).
74 Tsai, T. *et al.* Regional and age-specific patterns of pandemic H1N1 influenza virus
seroprevalence inferred from vaccine clinical trials, August-October 2009.
Eurosurveillance **15**, 19624 (2010).
75 Fox, S. J., Miller, J. C. & Meyers, L. A. Seasonality in risk of pandemic influenza
emergence. *PLoS computational biology* **13**, e1005749 (2017).
76 Opatowski, L. *et al.* Transmission characteristics of the 2009 H1N1 influenza pandemic:
comparison of 8 Southern hemisphere countries. *PLoS Pathog* **7**, e1002225 (2011).
77 Lloyd-Smith, J. O., Schreiber, S. J., Kopp, P. E. & Getz, W. M. Superspreading and the
effect of individual variation on disease emergence. *Nature* **438**, 355-359 (2005).
78 Barbosa, H. *et al.* Human mobility: Models and applications. *Physics Reports* **734**, 1-74
(2018).
79 Changruenngam, S., Bicout, D. J. & Modchang, C. How the individual human mobility
spatio-temporally shapes the disease transmission dynamics. *Scientific Reports* **10**, 1-13
(2020).
80 Łuksza, M. & Lässig, M. A predictive fitness model for influenza. *Nature* **507**, 57 (2014).
81 Agor, J. K. & Özaltın, O. Y. Models for predicting the evolution of influenza to inform
vaccine strain selection. *Human vaccines & immunotherapeutics* **14**, 678-683 (2018).

Supporting information captions

S1 Text. Supplementary Information

665