Bacillus subtilis with a 1% population enhances the activity of *Enterococcus*

faecalis with a 99% population

Tsukasa Ito,[a]# and Yu Yamanashi,[a]

[a]Department of Environmental Engineering Science, Gunma University, Kiryu,

Gunma, Japan

Running title: 1% *Bacillu*s activated 99% *Enterococcus*

Keywords: microbial interaction; dye decolorization assay; minority; redox

potential (ORP); amino acids, metabolomics

#Address corresponding to Tsukasa Ito, t.ito@gunma-u.ac.jp

Mailing address: Department of Environmental Engineering Science, Gunma

University, 1-5-1 Tenjin-cho, Kiryu, Gunma 376-8515, Japan.

Phone & Fax: +81-(0)277-30-1632

ABSTRACT

Microbes are present as communities in the environment. However, the importance of minor populations has not been well studied experimentally. In this study, we evaluated the role of *Bacillus subtilis* with a 1% population and its effect on co-incubated *Enterococcus faecalis* with a 99% population. Here we used an azo dye-decolorizing *Enterococcus faecalis* strain T6a1 and non-dye-decolorizing *Bacillus subtilis* strain S4ga. The dye decolorization assay enabled the investigation of the effects of *B. subtilis* S4ga on the activity of *E. faecalis* T6a1, even when *B. subtilis* S4ga was present at only 1% relative abundance or lower. We found that non-decolorizing *B. subtilis* S4ga enhanced the dye decolorization activity of *E. faecalis* T6a1, shortened the lag time of *E. faecalis* T6a1 to start decreasing the dye concentration, and increased the time for *E. faecalis* T6a1 to continue dye decolorization. These effects were correlated with redox potential values. We compared the extracellular amino acids between each incubation culture of *E. faecalis* T6a1 and *B. subtilis* S4ga, which revealed their mutual relationship by cross-feeding of specific amino acids. We also compared the intracellular primary metabolites between co-incubation and sole incubation of *E. faecalis* T6a1. The arginine deiminase (ADI) pathway in the co-incubated *E. faecalis* T6a1 was activated compared to that of *E. faecalis* T6a1 incubated solely. These findings explained that co-incubation with *B. subtilis* S4ga promoted ATP production in *E. faecalis* T6a1 cells to a greater extent and enhanced dye-decolorization activity.

IMPORTANCE

This study highlights the importance of minor bacterial populations and their effects on major populations. We used *Enterococcus faecalis* as the major population and *Bacillus subtilis* as the minor population. Both species are becoming increasingly important. Some strains of *E. faecalis* are antibiotic-resistant pathogens, show probiotic effects, or are applicable for textile wastewater treatment. Some strains of *B. subtilis* are known to produce antimicrobial agents, reduce intestinal inflammation, or restore gut microbiota. We demonstrated that a low abundance of *B. subtilis* with 1% population increased the amount of energy produced by *E. faecalis* with 99% population, which appeared as enhanced dye decolorization activity of dye-decolorizing *E. faecalis*. Metabolomic analysis suggested that *E. faecalis* and *B. subtilis* had a mutual relationship by feeding specific amino acids to each other. These results provide new insights into co-existing minor populations in microbial communities and will improve our understanding of bacterial control.

INTRODUCTION

Enterococcus faecalis is becoming an enormously important bacterium in various aspects. Some strains are pathogens and are nosocomial antibiotic-resistant bacteria known as vancomycin-resistant enterococci (**1, 2**), while some strains produce an antimicrobial agent, bacteriocin, against spoilage bacteria (**3, 4**) or add flavoring (**4**) in the food industry. Probiotics for humans and animals have also been reported (**4, 5**). In the environment, *E. faecalis* is a fecal indicator and is used for microbial source tracking (**6**), and other strains are dye-decolorizing bacteria applicable for wastewater treatment discharged from the dye/textile industry (**7, 8**). Since *E. faecalis* is present in communities with other bacteria in the above situations, investigating how *E. faecalis* is associated with other bacteria is the key to establishing a way to control *E. faecalis*.

Bacillus subtilis is known for its beneficial effects by producing antimicrobial agents and surfactants (**9, 10**), by reducing intestinal inflammation (**9, 11, 12**), and by modulating or restoring the gut microbiota (**13, 14**), despite its lower abundance in the microbial communities. Here we focused on the combination of *B. subtilis* and *E. faecalis* and on their relationship when *B. subtilis* is a minor population, such as an additive.

Although next-generation sequencing technology currently enables the detection of microbial community members with very low abundances, their functional roles in microbial communities are poorly understood. To investigate the role of the minor population, it is necessary to establish a stable community composition during the investigation and to quantify the population composition. Furthermore, it is necessary to assay the functional activity of the major

population affected by minor populations. Some assays may be required to

separate or distinguish minor populations from mixed communities.

In this study, we used a dye-decolorizing *E. faecalis* strain as a reporter

strain, such that when the strain is activated its decolorization is enhanced, and a

non-dye-decolorizing *B. subtilis* strain as the partner strain with a minor

population. We evaluated the activities of *E. faecalis* when incubated solely and

when co-incubated with *B. subtilis* using the dye decolorization assay. The dye

decolorization assay, though a low-tech method, visualized the activity of *E.*

faecalis as the color change of test tubes within a day with high throughput.

Although green fluorescent protein (GFP) has been used in many reporter assays,

GFP-generating fluorophores require oxygen (**15**). The GFP-independent activity

assay (dye decolorization assay) is suitable for targeting the dye-decolorizing

enzyme azoreductase of *E. faecalis*, which is activated in reduced environments

(**16**). Another benefit is that *E. faecalis* and *B. subtilis* are easily culturable,

distinguishable, and therefore quantifiable on the same agar plate even though

their population ratio is 100:1 or more. Furthermore, we were able to separate

each strain from the co-incubated culture by using a 0.8-µm filter for analyzing

intracellular and extracellular metabolites. In the dye decolorization assay, we

prepared the initial *E. faecalis* population to be approximately 7–8 log CFU/mL

and incubated in 10% Luria-Bertani (LB) medium, which controlled the growth of

E. faecalis and *B. subtilis* during co-incubation. For example, although *B. subtilis*

can grow from 5 log CFU/mL to 8 log CFU/mL within a day, it was restrained to

6 CFU/mL when co-incubated with 8 log CFU/mL of *E. faecalis* in 10% LB

medium. Thus, we controlled the population ratio of the two strains (*B. subtilis*/*E.*

faecalis) between 0.04 and 0.4%. Therefore, we successfully evaluated the effects

of non-dye-decolorizing *B. subtilis* with a 1 % population on the enhancement of

the activity of dye-decolorizing *E. faecalis*.

RESULTS AND DISCUSSION

Effects of *B. subtilis* on the activity of *E. faecalis*

We investigated the activity of *E. faecalis* T6a1 using the dye decolorization

assay at a dye concentration of 60 mg/L. **Fig. 1A** shows typical changes in dye

concentrations during the co-incubation of *E. faecalis* T6a1 and *B. subtilis* S4ga

and the sole incubation of *E. faecalis* T6a1. The *E. faecalis* T6a1 population

during the co-incubation was approximately one-fifth that of the sole incubation

at 48 h: 1.8×10^8 CFU/mL for the co-incubation, 8.4×10^8 CFU/mL for the sole

incubation. However, the dye concentration decreased earlier and faster during

the co-incubation, and at 48 h, the amount of dye decolorized during the co-

incubation was almost twice that of the sole incubation. In contrast, the sole

incubation did not show additional decolorization after 48 h. In the co-incubation,

the population of *B. subtilis* S4ga was 1.4×10^5 CFU/mL, and thus, the

population ratio of *B. subtilis* S4ga to *E. faecalis* T6a1 was 0.08%; this low

ratio resulted in differences in the dye decolorization activity of *E. faecalis* T6a1.

Furthermore, the maximum dye decolorization rate of the co-incubation was

twice as high as that of the sole incubation (**Fig. 1B**). We also repeatedly

observed that the co-incubation doubled the amount of dye decolorized and

halved the lag time to start decreasing the dye concentration compared with those

of the sole incubation (**Fig. 1C, D**). Additionally, co-incubation increased the

duration of dye decolorization by approximately 2.2 times compared with sole

incubation of *E. faecalis* T6a1 (**Fig. 1E**). In other words, dye decolorization by

the co-incubated strain started earlier and lasted longer compared with that of the

sole incubated strain, resulting in decolorization of a large amount of dye. The

population ratio of *B. subtilis* S4ga to *E. faecalis* T6a1 was only 0.1%.

Co-incubation enhanced the decolorization effects of *E. faecalis* T6a1. For

instance, shortening the lag time to start decreasing the dye concentration

indicated that *B. subtilis* S4ga affected the activity of *E. faecalis* T6a1 before or

at the same time that dye degradation proceeded. Therefore, we measured the

intracellular ATP levels of co-incubated *E. faecalis* T6a1 and that of the sole

incubated *E. faecalis* T6a1. The intracellular ATP level of co-incubated *E.*

faecalis T6a1 was 16 times higher than that of the sole-incubated *E. faecalis* T6a1

(**Fig. 1F**). On the other hand, the intracellular ATP of co-incubated *B. subtilis*

S4ga was 2.5 times higher than that of the sole-incubated *B. subtilis* S4ga (**Fig.**

1G). These results suggest that both *E. faecalis* T6a1 and *B. subtilis* S4ga were

activated by co-incubation.

Population ratio of *B. subtilis* to *E. faecalis*

We evaluated the effect of various population ratios of *B. subtilis* S4ga to *E.*

faecalis T6a1 on dye decolorization activity of *E. faecalis* T6a1 by changing the

amount of *B. subtilis* S4ga added to the same amount of *E. faecalis* T6a1. The

effective population ratios ranged between 0.04% and 40%. There were no

significant differences in the decolorization rates of the cocultures within this

wide range of *B. subtilis* S4ga population ratios (**Fig. 1H**). However, the

decolorization rates of the co-incubations were higher than those of the sole

incubation at each time point (**Fig. 1B, H**). We created a box and whisker plot to

show the distribution of the effective population ratio with data lower than 4%

and demonstrated that 0.4–0.8% *B. subtilis* S4ga (less than 1%) enhanced the dye

decolorization activity of *E. faecalis* T6a1 (**Fig. 1I**).

Effects of *B. subtilis* on the oxidation state

The dye-decolorizing enzyme azoreductase, which is expressed by *E. faecalis*, is

activated under reduced environments (**16**). Therefore, we investigated dissolved

oxygen concentration (DO) and decolorization rate of cocultures of *E. faecalis*

T6a1 and *B. subtilis* S4ga under oxic and anoxic conditions for 23 h (**Fig. 2**). Sole

incubation of each strain was performed for comparison. The sole incubation of

E. faecalis T6a1 showed approximately 75% decolorization under anoxic

conditions at 23 h, while the sole incubation of *E. faecalis* T6a1 showed a much

lower decolorization rate (20–45%) under oxic conditions at 23 h, mainly

between 3.5 and 4.2 mg/L DO (**Fig. 2A**). On the other hand, the co-incubation of

E. faecalis T6a1 and *B. subtilis* S4ga resulted in higher dye decolorization rates

of 80–95% at 8 h under oxic conditions, mainly between 1.5 and 3.2 mg/L DO,

and further increased to 90–100% at 23 h. Under anoxic conditions, the co-

incubation showed incomplete decolorization of approximately 75–90% at 23 h

(**Fig. 2B**). We found that under oxic conditions, co-incubation of *E. faecalis* T6a1

and *B. subtilis* S4ga lowered DO below 3 mg/L, whereas sole incubation of *E.*

faecalis T6a1 did not decrease DO as much.

Next, we examined the effects of *B. subtilis* S4ga on ORP values for

dye decolorization of *E. faecalis* T6a1, both sole incubation of *E. faecalis* T6a1

and co-incubation with *B. subtilis* S4ga were carried out under oxic conditions

below approximately 3 mg/L DO (**Fig. 2D**). The initial oxidation-reduction

potential (ORP) was +163 ± 11 mV in the presence of 1.9 mg/L DO. The ORP

values of the sole incubation of *E. faecalis* T6a1 were in the range of +90 mV to

+140 mV at 20 h. In contrast, the ORP values of the co-incubation were in the

range of −30 mV to +80 mV (**Fig. 2E**).

To confirm the effects of *B. subtilis* S4ga on lowering ORP, we

monitored the changes in ORP during the sole incubation of *B. subtilis* S4ga

under oxic and microaerophilic conditions (**Fig. 3**). Under oxic condition, *B.*

subtilis S4ga lowered ORP to approximately 0 mV after 8-h incubation after

decreasing DO below 3 mg/L. Under microaerophilic condition, *B. subtilis* S4ga

lowered ORP to approximately + 60 mV after 8-h incubation with decreasing DO

below 1 mg/L. *B. subtilis* S4ga created a moderately reduced environment when

DO was present and the ORP level under oxic condition was lower than that

under microaerophilic condition. These results demonstrate that *B. subtilis* S4ga

lowered the ORP to a condition that enhanced dye decolorization of *E. faecalis*

T6a1 under oxic conditions and that oxygen was required to enhance dye

decolorization of *E. faecalis* T6a1 when co-incubated with *B. subtilis* S4ga. *B.*

subtilis has been widely used to produce amino acids (**17**) and vitamins (**18**).

Since oxygen is required for the production of these compounds (**19, 20**), *B.*

subtilis probably produced such reducing agents under oxic conditions, resulting

in the lower ORP values observed in this study.

Extracellular metabolites of co-incubation and sole incubations

Extracellular metabolites of individual incubation of *E. faecalis* T6a1 and *B.*

subtilis S4ga and co-incubated culture revealed characteristics of amino acid

production and consumption from the LB 10% medium (**Fig. 4**). Amino acids

produced by *B. subtilis* S4ga were tyrosine, methionine, tryptophan,

phenylalanine, valine, and leucine ($P<0.05$). Solely incubated *E. faecalis* T61a

utilized all of these amino acids. However, the decreased amounts of methionine,

tryptophan, phenylalanine, valine, and leucine in co-culture were not as high as

those in the sole incubation of *E. faecalis* T6a1, which is probably because *B.*

subtilis S4ga provided the co-culture with these amino acids. Although *B. subtilis*

S4ga sole incubation produced tyrosine twice as much as the amount of LB 10%

medium, the co-incubation of *E. faecalis* T6a1 with 1% *B. subtilis* S4ga

consumed tyrosine and arginine, both of which were possibly rate-limiting

substrates. Interestingly, all the amino acids produced by *B. subtilis* S4ga

(methionine, tryptophan, phenylalanine, valine, and leucine) were hydrophobic.

Since hydrophobic amino acids are more permeable through the cell membrane

than hydrophilic amino acids, *E. faecalis* T6a1 can readily acquire these amino

acids. In turn, amino acids produced by *E. faecalis* T6a1 were glutamine,

histidine, aspartic acid, and proline ($P<0.05$). Solely incubated *B. subtilis* S4ga

utilized all of these amino acids. The production and utilization of the amino

acids by both strains demonstrated cross-feeding of amino acids between *E.*

faecalis T6a1 and *B. subtilis* S4ga, indicating a synergistic effect on the activities

of each strain.

Intracellular metabolites of *E. faecalis* incubated with *B. subtilis*

Among the 97 water-soluble primary metabolites, citrulline was the most

significantly altered intracellular metabolite, with higher levels in *E. faecalis*

T6a1 cells co-incubated with *B. subtilis* S4ga than in solely incubated *E. faecalis*

T6a1 cells (**Table S1**). Citrulline is an amino acid produced during arginine

degradation by the ADI pathway of *E. faecalis* (**21**). The ADI pathway is an ATP-

producing energy-generating process for *E. faecalis* (**22–24**). The ADI pathway

catalyzes the conversion of arginine to citrulline and ammonia and further to

ornithine, carbon dioxide, and ammonia, with the generation of 2 mol ATP for 1

mol arginine (EC 3.5.3.6, EC 6.3.4.16). In fact, we demonstrated more than 10

times higher amounts of ATP in the co-incubated *E. faecalis* T6a1 cells than in

the solely incubated *E. faecalis* T6a1 cells (**Fig. 1F**). Although low oxygen

tension is required to induce the ADI pathway in *Pseudomonas aeruginosa* (**25**),

in *E. faecalis*, oxygen tension has a weak effect on the ADI pathway enzymes,

and arginine is a more important inducer (**23, 24**). Similarly, higher dye-

decolorization activities of *E. faecalis* T6a1 were obtained under oxic conditions

(**Fig. 2B**), and extracellular arginine was extensively consumed in both solely

incubated *E. faecalis* T6a1 and co-incubated *E. faecalis* T6a1 (**Fig. 4**). In

addition, among the intracellular metabolites of *E. faecalis* T6a1, the detected

amount of arginine was higher in the co-incubation than in the sole incubation of

E. faecalis T6a1 (**Fig. 5, Table S1**). Together with citrulline and arginine, the

increase in ornithine is also explained by the observation that the ADI pathway

excretes abundant ornithine (**21**). Accordingly, we concluded that co-incubation

with *B. subtilis* S4ga activated the ADI pathway and promoted energy generation

in *E. faecalis* T6a1 cells.

Proline levels were also significantly higher in cocultured *E. faecalis* T6a1

cells. Proline is an abundant amino acid in *E. faecalis* enterolysin A (**26**), which

breaks down bacterial cell walls. Because some strains of *Bacillus*, including *B.*

subtilis, show weak sensitivity to enterolysin A (**26**), the presence of *B. subtilis*

S4ga may stimulate enterolysin A production by *E. faecalis* T6a1. Moreover, we

assumed that the presence of *B. subtilis* S4ga affected the oxidative stress

response of *E. faecalis* T6a1 and therefore focused on metabolites related to

oxidative stress responses, such as glutathione and oxidized glutathione (**27, 28**).

It seems that the amount of glutathione detected in solely incubated *E. faecalis*

T6a1 was higher than that in the co-incubation, although the difference was not

statistically significant. Further studies are needed to determine whether the

oxidative stress level of *E. faecalis* T6a1 was decreased by co-incubation with *B.*

subtilis S4ga.

In conclusion, we report here that *B. subtilis* S4ga enhanced the activity

of *E. faecalis* T6a1, even when *B. subtilis* S4ga was present at only 0.1–1% of the

total population, as demonstrated by the dye decolorization assay using non-dye-

decolorizing *B. subtilis* S4ga and dye-decolorizing *E. faecalis* T6a1. The effects

of *B. subtilis* S4ga on increasing the activity of *E. faecalis* T6a1 appeared not

only as a high maximum dye decolorization rate, but also as a shortened lag time

for *E. faecalis* T6a1 to start decreasing the dye concentration, as an increased

duration of dye decolorization, and the amount of dye decolorization was doubled.

During co-incubation, *B. subtilis* S4ga lowered the ORP to approximately 0 mV

by consuming oxygen and/or producing reducing agents such as amino acids. The

amino acids produced by *B. subtilis* S4ga were the same as those consumed by *E.*

faecalis T6a1, while the amino acids produced by *E. faecalis* T6a1 were the same

as those consumed by *B. subtilis* S4ga. In other words, these strains had a mutual

supply relationship with the amino acids. This elevated the intracellular ATP

content of *E. faecalis* T6a1 and *B. subtilis* S4ga, partly via activation of the ADI

pathway of *E. faecalis* T6a1. In the future, we will clarify whether the

combination of these species or the presence of *B. subtilis* is important for

enhancing the whole-cell activity of coexisting bacteria.

MATERIALS AND METHODS

Strains used in this study and identification

One bacterial strain (*E. faecalis* strain T6a1) with dye-decolorizing ability was

isolated from the fingertips, as described by Ito et al. (**29, 30**). Seven non-dye-

decolorizing *Bacillus*-like strains (S4ga, S4gb, C16caA, C47ea, A14d3B, D3da,

and C8ba) were isolated in the same manner (**Table S2**). Full-length 16S rRNA

gene sequences of these strains were analyzed by colony polymerase chain

reaction, followed by Sanger sequencing, which were carried out at GENEWIZ

Japan Corp. (Tokyo, Japan). Identification of the strains was based on an NCBI

BLAST search for 16S rRNA gene sequences, colony morphology, and

microscopy.

Screening of non-decolorizing *Bacillus* strains that enhanced dye

decolorization of *E. faecalis* T6a1

The isolated strains were pre-incubated under two conditions: sole incubation of

each strain and co-incubation of *E. faecalis* T6a1 with one *Bacillus*-like strain.

Each strain was inoculated into 5 mL of LB liquid medium in a test tube with a

lid. After culturing statically for 17 h at 37°C, 30 μL of 10 g/L Congo red dye

(Wako Pure Chemicals, Tokyo, Japan; cat. no. 032-03922) was added to each test

tube to achieve an initial dye concentration of 60 mg/L. After the addition of

Congo red dye, the samples were incubated statically at 37°C under oxic

conditions by leaving the gas phase of the test tubes as air. The color of the liquid

in the test tubes was regularly observed. The decreased dye concentration was

determined by measuring the absorbance of the liquid in the test tubes at a

wavelength of 490 nm. When measuring the absorbance, the liquid containing the

cells and dye was centrifuged at $14000 \times g$ for 5 min, and the supernatant was

analyzed using a spectrophotometer. The dye decolorization rate was calculated

from the maximum slope of the line on each dye concentration-time graph. The

dye decolorization assay was performed in triplicate.

Among the seven *Bacillus* strains, strains S4ga and S4gb were closely

related to *B. subtilis* (100% similarity for the full-length 16S rRNA gene). In

addition, strain S4ga showed the highest acceleration of dye decolorization of *E.*

faecalis T6a1, resulting in complete decolorization of 60 mg/L Congo red to 0

mg/L (**Table S2**). Sole incubation of *E. faecalis* T6a1 decreased the dye

concentration, but at a decolorization rate of two-thirds that of the co-incubation

with strain S4ga. Thus, strain S4ga was selected for subsequent analyses.

Population ratio setting

To test different population ratios of *E. faecalis* T6a1 and *B. subtilis* S4ga, we

prepared pre-incubated *E. faecalis* T6a1 cells that were suspended in 10% LB

medium with high turbidity and pre-incubated *B. subtilis* S4ga cells as follows. *E.*

faecalis T6a1 and *B. subtilis* S4ga were separately pre-incubated in 100% LB

medium at 37°C for 14 h. The cells were collected and washed with 10% LB

medium, and each strain was then suspended in 10% LB medium. The cell

turbidity of *E. faecalis* T6a1 was adjusted to approximately 3 O.D. in 10% LB

medium, which suppressed the growth of the strains by 2–8 times compared with

their initial cell numbers during subsequent incubation with dye. This enabled the

regulation of different population ratios of *B. subtilis* S4ga to *E. faecalis* T6a1

during incubation. The population ratios were set to be approximately 1% and in

the range between 0.01% and 50%, as estimated based on the cell turbidity after

pre-incubation and confirmed by CFU counting. The cell number of *E. faecalis*

T6a1 in sole incubation was adjusted to be similar to that of *E. faecalis* T6a1 in

co-incubation with *B. subtilis* S4ga.

Dye decolorization assay

The dye-decolorizing *E. faecalis* T6a1 was co-incubated with the non-dye-

decolorizing *B. subtilis* S4ga and Congo red dye at 37°C. In parallel, *E. faecalis*

T6a1 sole incubation was used as a control. Each strain was inoculated into 5 mL

of 10% LB liquid medium in a test tube with a lid. To assay dye decolorization

under oxic conditions, oxic conditions were assessed by leaving the gas phase as

air. The initial DO under oxic conditions was between 6 and 7 mg/L. To assay

dye decolorization under anoxic conditions, anoxic conditions were created by

nitrogen purging of the gas phase in the tube for 60 s. After nitrogen purging, the

DO in the liquid in the test tube was less than 0.5 mg/L. The concentration of

Congo red dye was set to 60 mg/L. Immediately after the addition of the dye, the

DO was measured routinely using a noncontact-type DO sensor, Fibox3-trace

(PreSens Precision Sensing, Regensburg, Germany) until the liquid in the test

tube was decolorized or the incubation was terminated. The ORP value of the

liquid was measured when the incubation was terminated or when the dye in the

liquid was completely decolorized. The initial ORP values were in the range of

+155 mV to +175 mV. The pH did not change significantly before and after the

incubation, which was between 7.2 and 7.6. The viable cell count as CFU was

evaluated after decolorization. The dye concentration during incubation was

determined from the RGB of the routinely acquired images (**Fig. S2**). The final

dye concentration was determined by measuring the absorbance at 490 nm

wavelength.

Viable cell counts of *E. faecalis* T6a1 and *B. subtilis* S4ga

The amount of dye decolorized in the test tube was almost proportional to the

population of dye-decolorizing *E. faecalis* T6a1. Therefore, *E. faecalis* T6a1 cell

numbers were enumerated by dilution plating before and after incubation. The

incubated solutions collected from sole incubation of *E. faecalis* T6a1 and co-

incubation of *E. faecalis* T6a1 and *B. subtilis* S4ga were spread on 10% LB agar

medium with serial dilutions and then incubated at 37°C for 12 h, followed by

colony counting. Although *B. subtilis* S4ga colonies appeared with *E. faecalis*

T6a1 colonies on the same agar plate, they were easily discriminated based on

their colony sizes (**Fig. S1**). *E. faecalis* T6a1 lacked motility and was obviously

smaller in size (colonies measured approximately 0.5 mm in diameter), whereas

B. subtilis S4ga was motile and had larger colony sizes (1.5–3 mm in diameter)

after incubation for 12 h. *B. subtilis* S4ga colonies were further confirmed by

their *Bacillus*-like wrinkled appearance after incubation for 24–48 h.

Dye concentration estimation by measuring RGB values in photos of

incubating tubes

Time-course monitoring of dye concentration was performed by measuring the

RGB values in the photographs of the test tubes during incubation. First, standard

solutions of Congo red dye were prepared in 10% LB medium at concentrations

ranging from 0 to 300 mg/L to obtain a calibration equation. Five milliliters of

each standard solution was placed in a flat-bottom test tube for spectrophotometer

analysis (Spectroquant Empty cells 16 mm; Merck KGaA, Darmstadt, Germany),

which was also used for incubation. All test tubes of the standard solutions were

placed in a photography box equipped with an LED backlight, and photographs of

the test tubes were obtained using a camera with the following settings: shutter

speed, 1/100 s; aperture, f/2.8; focal length, 4.5 mm; ISO speed, 100 (**Fig. S2A**).

After loading the images on a PC, the values of R, G, and B at the center of each

test tube were measured using the Eyedropper tool in PowerPoint (Microsoft

Corporation, Redmond, WA, USA). The total RGB value was calculated for each

standard solution to create calibration curves of the total RGB value versus dye

concentration, yielding a linear approximation for concentrations below 40 mg/L

and an exponential approximation between 40 and 300 mg/L (**Fig. S2B**). When

determining the dye concentration in the incubation tubes, photographs of the

tubes were taken along with test tubes of the standard solutions with initial dye

concentrations of 60 and 300 mg/L as internal standards.

This method is a non-contact type measurement, which does not require

collecting the liquid from the test tube and thus does not disturb the incubation

solution in the test tube and the gas phase of the tube. Furthermore, the

measurement can be performed quickly and repeatedly by taking photographs of

the test tubes.

ORP measurements during *B. subtilis* S4ga incubation

A part of the *B. subtilis* S4ga colony on an agar plate was suspended in 50 mL of

LB liquid medium and placed in an incubator at 37°C for 1 h. The cell suspension

was centrifuged at 12000 rpm for 5 min and the supernatant was replaced with

10% LB liquid medium. The centrifuged cells were re-suspended in 10% LB and

thus a *B. subtilis* S4ga cell suspension (8 log CFU/mL) was prepared in a 50 mL

tube. For aerobic incubation, 200 mL of 10% LB liquid medium was prepared in

a 500 mL vial, and 240 μL of *B. subtilis* cell suspension was added to the

medium. Congo red (10 g/L) was then added at a concentration of 60 mg/L. Three

sets of vials were shaken at 60 rpm. For microaerophilic incubation, 5 mL of 10%

LB liquid medium was prepared in a 16 mL test tube with a rubber stopper, and

30 μL of *B. subtilis* S4ga cell suspension was added to the medium. Congo red

(10 g/L) was added at a final concentration of 60 mg/L. After the tube was

capped with a rubber stopper, nitrogen gas was injected into the test tube with a

needle for 3 s. DO was 1 mg/L at 5 h and thereafter remained lower than 1 mg/L.

During incubation at 37°C, the ORP and DO of the incubation medium of the

vials and test tubes were monitored for 27 h. Since it is necessary to open the

caps of the vials and tubes for measuring ORP, DO was measured using a non-

contact-type DO sensor before opening them. To measure the ORP of the

incubation solution under microaerophilic conditions, unopened test tubes were

opened each time. Therefore, 30 test tubes were initially prepared, and four to six

test tubes were measured at each time point.

Metabolomic analysis

(i) Incubation conditions. To investigate whether *B. subtilis* S4ga had some

effects on the metabolism of *E. faecalis* T6a1, metabolomic analysis of the

intracellular and extracellular water-soluble primary metabolites of strains T6a1

and S4ga was carried out for coculture and sole incubation cultures of each strain.

The incubations were conducted as described above (dye decolorization assay).

Briefly, after pre-incubation of each strain, co-incubation of *E. faecalis* T6a1 and

B. subtilis S4ga and sole incubation of each strain were performed at an initial

Congo red dye concentration of 60 mg/L in 10% LB medium at 37°C. The

incubations were terminated at 10 h during decolorization. The use of 10% LB

medium reduced background amino acids and vitamins derived from tryptone and

yeast extract of the incubation medium when intracellular and extracellular

metabolites were extracted.

(ii) Extraction and analysis of intracellular water-soluble primary

metabolites of *E. faecalis* T6a1. To compare intracellular metabolites of the co-

incubated *E. faecalis* T6a1 with those of the sole incubated *E. faecalis* T6a1, it

was necessary to remove *B. subtilis* S4ga cells from the co-incubation solution.

To do this, we focused on cell sizes; *E. faecalis* T6a1 cells were spherical,

approximately 0.8 μm in width and 1.3 μm in length, whereas *B. subtilis* S4ga

cells were elongated, approximately 0.8 μm in width and 2.6 μm or more in

length (**Fig. S3A, B**). Therefore, we expected that *B. subtilis* S4ga cells would not

pass through a 0.8-μm pore size filter compared with *E. faecalis* T6a1 cells.

Accordingly, a 0.8-μm filter was used to reduce the number of *B. subtilis* S4ga

cells in the co-incubation solution. After 10-h incubation with dye, the co-

incubation solution was filtered through a 0.8-μm pore size membrane filter

(DISMIC 25CS080AS; Advantec, Tokyo, Japan). In parallel, the sole incubation

solution of *E. faecalis* T6a1 was also filtered to ensure that the filtration

procedure did not affect the metabolites in *E. faecalis* T6a1 cells. The filtered

solutions were further filtered twice more to minimize contamination by *B.

subtilis* S4ga cells. Although *B. subtilis* S4ga accounted for 2% of the total

population after the 10-h incubation (**Fig. S3C, D**), the filtration treatment did not

allow us to find vegetative cells of *B. subtilis* S4ga, despite observing

approximately 10,000 cells of *E. faecalis* T6a1 by microscopy. We observed only

some endospores or non-vegetative thin cells (**Fig. S3E, F**), and *B. subtilis* S4ga

cells accounted for 0.07% of the total population. Such a small population and the

remaining non-vegetative cells would not affect the metabolomic analysis of *E.

faecalis* T6a1.

E. faecalis T6a1 cells prepared by filtration after co-incubation and sole

incubation were washed with phosphate-buffered saline once and then suspended

in sterilized ultrapure water. The turbidity of the cell suspension of *E. faecalis*

T6a1 under co-incubation and sole incubation was adjusted to be the same among

all samples by adding ultrapure water. Methanol was then added to the

suspensions at a concentration of 80% to extract the intracellular metabolites.

Next, 2-morpholinoethanesulfonic acid (2-MES) and methionine sulfone as

internal standards were added to the cell suspensions at a final concentration of

10 μM. After thorough mixing and centrifugation at $20000 \times g$ for 5 min, the

supernatants of the cell suspensions were collected for metabolomic analysis.

Two sets of samples were prepared for *E. faecalis* T6a1 co-incubation and sole

incubation.

Ninety-seven water-soluble primary metabolites (e.g., amino acids,

organic acids, and coenzymes) were evaluated in intracellular extraction solutions

of strain T6a1 by liquid chromatography tandem mass spectrometry (Shimadzu

LCMS-8050; Kyoto, Japan) equipped with a pentafluorophenylpropyl column.

The resulting intensity was normalized as the height ratio (peak height of each

metabolite divided by the peak height of 2-MES). In cases in which a small peak

was overlaid with the next peak or in which small peaks were considered noise,

such peaks were regarded as undetected metabolites. Metabolic pathways of *E.*

faecalis were analyzed using the Kyoto Encyclopedia of Genes and Genomes

(KEGG) database (**31**).

(iii) Extraction and analysis of extracellular water-soluble primary

metabolites of co-culture and sole incubation cultures of *E. faecalis* T6a1 and

B. subtilis **S4ga.** The incubation solutions of the co-culture and sole incubation

cultures of *E. faecalis* T6a1 and *B. subtilis* S4ga were centrifuged at $20000 \times g$

for 7 min, and the supernatants were filtered using a 0.2-μm pore size sterile

membrane filter (ASFIL 033022SO-SFCA; AS ONE, Osaka, Japan). In addition,

10% LB medium was used as a reference to determine the relative concentration

of amino acids from the components of tryptone and yeast extract (**32**). Methanol

was then added to the filtered supernatants to a concentration of 80% to extract

the extracellular metabolites. Next, 2-MES and methionine sulfone as internal

standards were added to the supernatants at a final concentration of 10 μM. Three

sets of samples were prepared for the co-culture and sole incubation cultures of

E. faecalis T6a1 and *B. subtilis* S4ga, and 10% LB medium. In the same manner

as for the analysis of intracellular metabolites, ninety-seven water-soluble

primary metabolites were evaluated in the extracellular extraction solutions.

Measurement of intracellular ATP of *E. faecalis* T6a1 and *B. subtilis* S4ga

The amount of intracellular ATP in *E. faecalis* T6a1 in sole culture was

determined by subtracting the amount of extracellular ATP of the 0.2 μm

filtration permeate from the amount of the total ATP of the culture medium

containing cells: $ATP_{E.faecalis} = ATP_{total} - ATP_{0.2 \ \mu m\text{-extracellular}}$. The amount of

intracellular ATP of *B. subtilis* S4ga in sole culture was determined in the same

manner: $ATP_{B.subtilis} = ATP_{total} - ATP_{0.2 \ \mu m\text{-extracellular}}$. For co-culture of *E. faecalis*

T6a1 and *B. subtilis* S4ga, the amount of intracellular ATP of *E. faecalis* T6a1 in

the co-culture was determined by subtracting the amount of extracellular ATP in

the 0.2 μm filtration permeate from the amount of ATP in the 0.8 μm filtration

permeate containing extracellular ATP and *E. faecalis* T6a1 cells (but not strain

S4ga cells): $\text{ATP}_{\text{E.faecalis}} = (\text{ATP}_{0.8\ \mu\text{m-extracellular}} + \text{ATP}_{0.8\ \mu\text{m-E.faecalis}}) - \text{ATP}_{0.2\ \mu\text{m-}}$

$_{\text{extracellular}}$.

The amount of intracellular ATP in *B. subtilis* S4ga in the co-culture was

determined by subtracting the amount of ATP in the 0.8 μm filtration permeate

containing extracellular ATP and *E. faecalis* T6a1 cells (but not *B. subtilis* S4ga

cells) from the amount of total ATP in the culture medium containing

extracellular ATP and intracellular ATP of both strain cells: $\text{ATP}_{\text{B.subtilis}} =$

$\text{ATP}_{\text{total}} - (\text{ATP}_{0.8\ \mu\text{m-extracellular}} + \text{ATP}_{0.8\ \mu\text{m-E.faecalis}})$. ATP measurement was

performed with BacTiter Glo ATP measurement reagent (Promega, WI, USA) and

a luminometer (Lu-mini, Viti Life Science Solutions, Shanghai, China).

Statistical analysis

Statistical significance was determined by two-tailed paired *t*-tests to compare the

co-incubation with *B. subtilis* S4ga with the sole-incubations of *E. faecalis* T6a1.

Data availability

The 16S rRNA gene sequences of the isolated strains are available in the

DDBJ/EMBL/GenBank databases under the accession numbers LC557810–

LC557817.

ACKNOWLEDGMENTS

This study was supported in part by JSPS KAKENHI (grant number 15K12369 to

T.I.) and by a Grant-in-Aid for JSPS Research Fellow (grant number JP17J11210

to Y.Y.). Metabolomic analysis was performed at the Laboratory for Analytical

Instruments, Education and Research Support Center, Gunma University.

REFERENCES

1. Fisher K, Phillips C. 2009. The ecology, epidemiology and virulence of

Enterococcus. Microbiology 155:1749–1757.

2. Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen

GS, Colomb-Cotinat M, Kretzschmar ME, Devleesschauwer B, Cecchini M,

Ouakrim DA, Oliveira TC, Struelens MJ, Suetens C, Monnet DL. 2019.

Burden of AMR Collaborative Group. Attributable deaths and disability-

adjusted life-years caused by infections with antibiotic-resistant bacteria in

the EU and the European Economic Area in 2015: a population-level

modelling analysis. Lancet Infect Dis 19:56–66.

3. Silva CCG, Silva SPM, Ribeiro SC. 2018. Application of bacteriocins and

protective cultures in dairy food preservation. Front Microbiol 9:594.

4. Foulquié Moreno MR, Sarantinopoulos P, Tsakalidou E, De Vuyst L. 2006.

The role and application of enterococci in food and health. Int J Food

Microbiol 106:1–24.

5. Klein G. 2003. Taxonomy, ecology and antibiotic resistance of enterococci

from food and the gastro-intestinal tract. Int J Food Microbiol 88:123–131.

6. Byappanahalli MN, Nevers MB, Korajkic A, Staley ZR, Harwood VJ. 2012.

Enterococci in the environment. Microbiol Mol Biol Rev 76:685–706.

7. Walker R, Gingell R, Murrells DF. 1971. Mechanisms of azo reduction by
Streptococcus faecalis I. Optimization of assay conditions. Xenobiotica
1:221–229.

8. Zahran SA, Ali-Tammam M, Hashem AM, Aziz RK, Ali AE.
2019. Azoreductase activity of dye-decolorizing bacteria isolated from the
human gut microbiota. Sci Rep 9:5508

9. Ilinskaya ON, Ulyanova VV, Yarullina DR, Gataullin IG. 2017. Secretome of
intestinal Bacilli: a natural guard against pathologies. Front Microbiol
8:1666.

10. Kaspar F, Neubauer P, Gimpel M. 2019. Bioactive secondary metabolites
from *Bacillus subtilis*: A Comprehensive Review. J Nat Prod 82:2038–2053.

11. Okamoto, K., Fujiya, M., Nata, T., Ueno, N., Inaba, Y., Ishikawa, C., et al.
2012. Competence and sporulation factor derived from *Bacillus subtilis*
improves epithelial cell injury in intestinal inflammation via
immunomodulation and cytoprotection. Int J Colorectal Dis 27:1039–1046.

12. Zhang HL, Li WS, Xu DN, Zheng WW, Liu Y, Chen J, Qiu ZB, Dorfman RG,
Zhang J, Liu J. 2016. Mucosa-reparing and microbiota-balancing therapeutic
effect of Bacillus subtilis alleviates dextrate sulfate sodium-induced
ulcerative colitis in mice. Exp Ther Med 12:2554-2562.

13. Hadieva G, Lutfullin M, Pudova D, Akosah Y, Shagimardanova E, Gogoleva
N, Sharipova M, Mardanova A. 2021. Supplementation of *Bacillus subtilis*
GM5 enhances broiler body weight gain and modulates cecal microbiota. 3
Biotech 11(3):126.

14. Ma Y, Wang W, Zhang H, Wang J, Zhang W, Gao J, Wu S, Qi G. 2018.

Supplemental Bacillus subtilis DSM 32315 manipulates intestinal structure

and microbial composition in broiler chickens. Sci Rep. 8:15358.

15. Cubitt AB, Heim R, Adams SR, Boyd AE, Gross LA, Tsien RY. 1995.

Understanding, improving and using green fluorescent proteins. Trends

Biochem Sci 20:448–455.

16. Chung KT, Stevens Jr SE, Cerniglia CE. 1992. The reduction of azo dyes by

the intestinal microflora. Crit Rev Microbiol 18:175–190.

17. Su Y, Liu C, Fang H, Zhang D. 2020. *Bacillus subtilis*: a universal cell

factory for industry, agriculture, biomaterials and medicine. Microb Cell

Fact 19:173.

18. Abbas CA, Sibirny AA. 2011. Genetic control of biosynthesis and transport

of riboflavin and flavin nucleotides and construction of robust

biotechnological producers. Microbiol Mol Biol Rev 75:321–360.

19. Hu J, Lei P, Mohsin A, Liu X, Huang M, Li L, Hu J, Hang H, Zhuang Y, Guo

M. 2017. Mixomics analysis of *Bacillus subtilis*: effect of oxygen availability

on riboflavin production. Microb Cell Fact 16:150.

20. Utagawa T, 2004. Production of arginine by fermentation. J Nutr 134:2854S–

857S.

21. Cunin R, Glansdorff N, Piérard LA, Stalon V. 1986. Biosynthesis and

metabolism of arginine in bacteria. Microbiol Rev 50:314–352.

22. Slade HD, Slamp WC. 1952. The formation of the arginine dihydrolase by

Streptococci and some properties of the enzyme system. J Bacteriol 64:455–

466.

23. Simon JP, Wargnies B, Stalon V. 1982. Control of enzyme synthesis in the

arginine deiminase pathway of *Streptococcus faecalis*. J Bacteriol 150:1085–

1090.

24. Barcelona-Andrés B, Marina A, Rubio V. 2002. Gene structure, organization,

expression, and potential regulatory mechanisms of arginine catabolism

in *Enterococcus faecalis*. J Bacteriol 184:6289–6300.

25. Lu CD, Winteler H, Abdelal A, Haas D. 1999. The ArgR regulatory protein, a

helper to the anaerobic regulator ANR during transcriptional activation of the

arcD promoter in *Pseudomonas aeruginosa*. J Bacteriol 181:2459–2464.

26. Nilsen T, Nes IF, Holo H. 2003. Enterolysin A, a cell wall-degrading

bacteriocin from *Enterococcus faecalis* LMG2333. Appl Environ Microbiol

69:2975–2984.

27. Patel MP, Marcinkeviciene J, Blanchard JS. 1998. *Enterococcus faecalis*

glutathione reductase: purification, characterization and expression under

normal and hyperbaric O_2 conditions. FEMS Microbiol Lett 166:155–163.

28. Portela CA, Smart KF, Tumanov S, Cook GM, Villas-Bôas SG. 2014. Global

metabolic response of *Enterococcus faecalis* to oxygen. J Bacteriol

196(11):2012–2022.

29. Ito T, Shimada Y, Suto T. 2018. Potential use of bacteria collected from

human hands for textile dye decolorization. Water Resour Ind 20:46–53.

30. Ito T. 2013. Color-removal by microorganisms isolated from human hands. J

Microbiol Biol Educ 14:244–247.

31. Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2014.

Data, information, knowledge and principle: back to metabolism in KEGG.

Nucleic Acids Res 42 D199–D205.

32. Becton, Dickinson and Company. 2009. Difco & BBL manual of

microbiological culture media. 2nd ed., p 686–687.

 FIGURE LEGENDS

Figure 1. Effects of *B. subtilis* S4ga on dye decolorization of *E. faecalis* T6a1.

(A) Dye decolorization over time during co-incubation of *B. subtilis* S4ga and *E.*

faecalis T6a1 or sole incubation of *E. faecalis* T6a1. (B) Dye decolorization rates

of solely incubated *E. faecalis* T6a1 and co-incubated *E. faecalis* T6a1 with *B.*

subtilis S4ga for 11 h. (C) Ratio of dye amount decolorized by the co-incubation

to dye amount decolorized by the sole incubation. (D) Ratio of the time when the

dye concentration started to decrease during the co-incubation to the time when

the dye concentration started decreasing in the sole incubation. (E) Ratio of the

time when the dye concentration continued to decrease in the co-incubation to the

time when the dye concentration continued to decrease in the sole incubation. (F)

Intracellular ATP of solely incubated *E. faecalis* T6a1 or co-incubated *E. faecalis*

T6a1 after removal of *B. subtilis* S4ga at 24 h. (G) Intracellular ATP of solely

incubated *B. subtilis* S4ga or co-incubated *B. subtilis* S4ga after removal of *E.*

faecalis T6a1 at 24 h. (H) Dye decolorization rates of *E. faecalis* T6a1 co-

incubated with *B. subtilis* S4ga at the different population ratios ranging from

0.04% to 40%. (I) Distributions of the population ratios of *B. subtilis* S4ga to *E.*

faecalis T6a1 that enhanced the dye-decolorization activity of *E. faecalis* T6a1. A

box and whisker plot was created with data lower than 4% to focus on the

effectiveness of the low ratio range. Error bars represent standard deviations (B–

I); $*P < 0.05$; $**P < 0.01$; n = 3 (B); n = 9 (C–E); n = 3 (F, G); n = 8 (I). a.u.,

arbitrary units.

Figure 2. Impact of *B. subtilis* S4ga on dissolved oxygen (DO) concentration and

redox potential (ORP) during co-incubation with *E. faecalis* T6a1 and during

individual incubation with dye under oxic and anoxic conditions. Correlation

between final DO and dye decolorization rates of the incubations of (A) *E.*

faecalis T6a1, (B) *E. faecalis* T6a1 with *B. subtilis* S4ga, and (C) *B. subtilis*

S4ga, as determined at 8 h and 23 h. (D) Correlation between final ORP and DO

values after the incubations of *E. faecalis* T6a, *E. faecalis* T6a1 with *B. subtilis*

S4ga, and *B. subtilis* S4ga for 20 h. Populations of *E. faecalis* T6a1 and *B.*

subtilis S4ga at 20 h (CFU/mL), Co-1: T6a1, 2×10^7; S4ga, 7×10^5 (S4ga-to-

T6a1 population ratio, 3%). Co-2: T6a1, 2×10^8; S4ga, 1×10^6 (S4ga-to-T6a1,

0.6%). Co-3: T6a1, 7×10^7; S4ga, 5×10^5 (S4ga-to-T6a1, 0.7%). T-1: T6a1, $3 \times$

10^6. T-2: T6a1, 2×10^7. T-3: T6a1, 3×10^7. (E) A box and whisker plot showing

ORP values after incubation of *E. faecalis* T6a1 or *E. faecalis* T6a1 with *B.*

subtilis S4ga. ***$P < 0.001$; n = 11 (T6a1), n = 11 (T6a1 with S4ga). The initial

DO under oxic conditions was between 6 and 7 mg/L (A–E). The initial

concentration of Congo red dye was 60 mg/L (A–E).

Figure 3. Effects of *B. subtilis* S4ga on redox potential (ORP) under oxic and

microaerophilic conditions. ORP (A) and dissolved oxygen (DO) concentration

(B) over time during sole incubation of *B. subtilis* S4ga under shaking and static

conditions. Rotational shaking at 60 rpm created oxic conditioned. Static

incubation after nitrogen purging for nine seconds created microaerophilic

condition. Population of *B. subtilis* S4ga at 14 h: oxic, 1×10^8 (CFU/mL);

microaerophilic, 0.5×10^8 (CFU/mL). The initial *B. subtilis* S4ga population

was 3×10^5 (CFU/mL). Error bars represent standard deviations; n = 3.

Figure 4. Extracellular amino acids produced or consumed by *B. subtilis* S4ga, *E.*

faecalis T6a1 and co-culture of *E. faecalis* T6a1 and *B. subtilis* S4ga. Data

represent ratios of amino acids in the supernatants of the incubated cultures at 14

h to amino acids in 10% LB medium at 0 h. Co-culture: *E. faecalis* T6a1, 5×10^9

(CFU/mL); *B. subtilis* S4ga, 1×10^8 (CFU/mL) (S4ga-to-T6a1 population ratio,

2%). n = 3. a.u., arbitrary units.

Figure 5. Metabolites of the arginine deaminase (ADI) pathway produced by *E.*

faecalis T6a1 during co-incubation with *B. subtilis* S4ga and dye decolorization.

Error bars represent standard deviations; $**P < 0.01$; n = 2. Data represent

normalized intensities of the metabolites. a.u., arbitrary unit.

Figure 1. Effects of *B. subtilis* S4ga on dye decolorization of *E. faecalis* T6a1. (A) Dye decolorization over time during co-incubation of *B. subtilis* S4ga and *E. faecalis* T6a1 and sole incubation of *E. faecalis* T6a1. (B) Dye-decolorization rates of solely incubated *E. faecalis* T6a1 or co-incubated *E. faecalis* T6a1 with *B. subtilis* S4ga for 11 h.) (C) Ratio of dye amount decolorized by the co-incubation to dye amount decolorized by the sole incubation. (D) Ratio of the time when the dye concentration started to decrease during the co-incubation to the time when the dye concentration started decreasing in the sole incubation. (E) Ratio of the time when the dye concentration continued to decrease in the co-incubation to the time when the dye concentration continued to decrease in the sole incubation. (F) Intracellular ATP of solely incubated *E. faecalis* T6a1 or co-incubated *E. faecalis* T6a1 after removal of *B. subtilis* S4ga at 24 h. (G) Intracellular ATP of solely incubated *B. subtilis* S4ga or co-incubated *B. subtilis* S4ga after removal of *E. faecalis* T6a1 at 24 h. (H) Dye-decolorization rates of *E. faecalis* T6a1 co-incubated with *B. subtilis* S4ga at the different population ratios ranging from 0.04% to 40%. (I) Distributions of the population ratios of *B. subtilis* S4ga to *E. faecalis* T6a1 that enhanced the dye-decolorization activity of *E. faecalis* T6a1. A box and whisker plot was created with data lower than 4% to focus on the effectiveness of the low ratio range. Error bars represent standard deviations (B–I); *$P < 0.05$; **$P < 0.01$; $n = 3$ (B); $n = 9$ (C–E); $n = 3$ (F, G); $n = 8$ (I). a.u., arbitrary units.

Figure 1. Ito & Yamanashi

Figure 2. Impact of *B. subtilis* S4ga on dissolved oxygen (DO) concentration and redox potential (ORP) during co-incubation with *E. faecalis* T6a1 and during individual incubation with dye under oxic and anoxic conditions. Correlation between final DO and dye-decolorization rates of the incubations of (A) *E. faecalis* T6a1, (B) *E. faecalis* T6a1 with *B. subtilis* S4ga, and (C) *B. subtilis* S4ga. (D) Correlation between final ORP and DO values after the incubations of *E. faecalis* T6a, *E. faecalis* T6a1 with *B. subtilis* S4ga, and *B. subtilis* S4ga. The incubation period was 20 h. Populations of *E. faecalis* T6a1 and *B. subtilis* S4ga at 20 h (CFU/mL), Co-1: T6a1, 2×10^7; S4ga, 7×10^5 (S4ga-to-T6a1 population ratio, 3%). Co-2: T6a1, 2×10^8; S4ga, 1×10^6 (S4ga-to-T6a1, 0.6%). Co-3: T6a1, 7×10^7; S4ga, 5×10^5 (S4ga-to-T6a1, 0.7%). T-1: T6a1, 3×10^6. T-2: T6a1, 2×10^7. T-3: T6a1, 3×10^7. (E) ORP values after incubation of *E. faecalis* T6a1 or *E. faecalis* T6a1 with *B. subtilis* S4ga. ***$P < 0.001$; n = 11 (T6a1), n = 11 (T6a1 with S4ga). The initial DO under oxic conditions was between 6 and 7 mg/L (A–E). The initial concentration of Congo red dye was 60 mg/L (A–E).

Figure 2. Ito & Yamanashi

Figure 3. Effects of *B. subtilis* S4ga on redox potential (ORP) under oxic and microaerophilic conditions. ORP (A) and dissolved oxygen (DO) concentration (B) over time during sole incubation of *B. subtilis* S4ga under shaking and static conditions. Rotational shaking at 60 rpm created oxic conditioned. Static incubation after nitrogen purging for nine seconds created microaerophilic condition. Population of *B. subtilis* S4ga at 14 h: oxic, 1×10^8 (CFU/mL); microaerophilic, 0.5×10^8 (CFU/mL). The initial *B. subtilis* S4ga population was 3×10^5 (CFU/mL). Error bars represent standard deviations; n = 3.

Figure 3. Ito & Yamanashi

Figure 4. Extracellular amino acids produced or consumed by *B. subtilis* S4ga, *E. faecalis* T6a1 and co-culture of *E. faecalis* T6a1 and *B. subtilis* S4ga. Data represent ratios of amino acids in the supernatants of the incubated cultures at 14 h to amino acids in 10% LB medium at 0 h. Co-culture: *E. faecalis* T6a1, 5 × 10^9; *B. subtilis* S4ga, 1 × 10^8 (CFU/mL) (S4ga-to-T6a1 population ratio, 2%). n = 3. a.u., arbitrary units.

Figure 5. Metabolites of the arginine deaminase (ADI) pathway produced by *E. faecalis* T6a1 during co-incubation with *B. subtilis* S4ga and dye decolorization. Error bars represent standard deviations; **$P < 0.01$; n = 2. Data represent normalized intensities of the metabolites. a.u., arbitrary unit.