Natural and designed proteins inspired by extremotolerant organisms can form

condensates and attenuate apoptosis in human cells

Mike T. Veling[1,2,8], Dan T. Nguyen[1,2,8], Nicole N. Thadani[1], Michela E. Oster[1,2], Nathan J.

Rollins[1,2], Kelly P. Brock[1], Neville P. Bethel[3,4], Samuel Lim[1,2], David Baker[3,4,5], Jeffrey C.

Way[1,2], Debora S. Marks[1,6], Roger L. Chang[1,2,7,*], Pamela A. Silver[1,2,*]

Affiliations

1. Department of Systems Biology, Harvard Medical School, Boston, MA, 02115, USA;

2. Wyss Institute for Biologically Inspired Engineering, Boston, MA, 02115, USA;

3. Institute for Protein Design, University of Washington, Seattle, WA, 98195, USA;

4. Department of Biochemistry, University of Washington, Seattle, WA, 98195, USA;

5. Howard Hughes Medical Institute, University of Washington, Seattle, WA, 98195, USA;

6. Broad Institute of Harvard and MIT, Cambridge, MA, 02142, USA;

7. Department of Systems & Computational Biology, Albert Einstein College of Medicine,

Bronx, NY, 10461, USA;

8. Co-first authors

*. Co-corresponding authors (pamela_silver@hms.harvard.edu,

roger.chang@einsteinmed.edu)

ABSTRACT

Many organisms can survive extreme conditions and successfully recover to normal life. This extremotolerant behavior has been attributed in part to repetitive, amphipathic, and intrinsically disordered proteins that are upregulated in the protected state. Here, we assemble a library of approximately 300 naturally-occurring and designed extremotolerance-associated proteins to assess their ability to protect human cells from chemically-induced apoptosis. We show that several proteins from tardigrades, nematodes, and the Chinese giant salamander are apoptosis protective. Notably, we identify a region of the human ApoE protein with similarity to extremotolerance-associated proteins that also protects against apoptosis. This region mirrors the phase separation behavior seen with such proteins, like the tardigrade protein CAHS2. Moreover, we identify a synthetic protein, DHR81, that shares this combination of elevated phase separation propensity and apoptosis protection. Finally, we demonstrate that driving protective proteins into the condensate state increases apoptosis protection, and highlight the ability for DHR81 condensates to sequester caspase-7. Taken together, this work draws a link between extremotolerance-associated proteins, condensate formation, and designing human cellular protection.

INTRODUCTION

When exposed to extreme environmental stress, extremotolerance-associated (ExTol) organisms enter a state of reduced cellular metabolic activity that promotes survival in harsh conditions. This state has been shown to protect these organisms from high salinity, desiccation, and even the vacuum of outer space [1]. To acclimate to these conditions,

ExTol organisms significantly upregulate the production of protective metabolites, like

trehalose [2], as well as tolerance-conferring proteins [3], many of which are intrinsically

disordered proteins (IDPs) with repetitive sequence and predicted amphipathic helices.

While the mechanism of protection is unknown, one model suggests that at least some

of these proteins mediate formation of a gel- or glass-like material state within an entire

cell to protect it from damage [3]. The model posits that such proteins promiscuously

interact with membranes and other proteins, increasing intracellular viscosity and

protecting vulnerable structures [4-6] (Figure 1A top). Such interactions would also work to

prevent stress-induced protein unfolding and protein aggregation [7]. This vitrification

hypothesis is the subject of active investigation at the time of this writing, with evidence

both for and against its validity [8-10]. An alternative model suggests that these proteins

form intracellular phase-separated membraneless compartments with distinct physical

parameters (referred to here as condensates) that partition key stress-sensitive cellular

components [11] (Figure 1A bottom). Within these dense protein structures, essential

components are sequestered and protected from damage. The models are not mutually

exclusive; for example, in the case of desiccation tolerance, continued dehydration could

lead phase-separated compartments to fuse, comprise much (if not all) of the cell, and

promote a glassy material state.

Figure 1: Selection of candidate extremotolerance-associated proteins (A) Illustration of different protective models for cellular stress tolerance. (B) Illustration of amphipathic membrane interaction for rotifer LEA4 protein and Alpha-synuclein, thus highlighting a feature indicative of extremotolerance-associated proteins. (C) Hydrophobic moment analysis of rotifer LEA4, showing the repetitive nature of the amphipathic signature. Highlighted green segment corresponds to the LEA4 helix in (B). (D) Venn diagram of the lines of evidence used to select targets in the final library. Abbreviations: KD, knockdown; OE, overexpression; Δ, change in expression levels. (E) Domain analysis of included targets reveals LEA and ApoE domains in almost half of the targets tested. The other half contains proteins with other PFAM domains (19%) or without any affiliated PFAM entry (32%). Of the proteins lacking a known PFAM annotation, many exhibit repetitive sequence (20% of all targets).

The resurgence of interest in biological phase separation, extensively reviewed in [12-15],

has shown that human cells can employ similar strategies to survive stressful conditions.

While the reversible formation of stress granules in healthy cells is the archetype of stress

tolerance, recent studies have revealed phase separation as a key adaptation that cancer

cells use to avoid apoptosis and cell death [16-17]. Phase-separated condensates have been

implicated in sequestering key tumor repressors[17], enhancing or repressing gene

expression in response to stress [18], and regulating mRNA or protein function through

sequestration [19]. These processes parallel the mechanisms that ExTol proteins are

thought to use to protect complex biological systems, suggesting there could be some

functional overlap. That is, sequence features and properties within ExTol proteins could

be implicated in mammalian stress tolerance pathways, including apoptosis resistance.

A key sequence feature of some ExTol proteins is the presence of amphipathic helices

[20]. While disordered in most contexts, ExTol proteins can transiently form amphipathic

helices in certain molecular environments, notably when interacting with and stabilizing

membranes [6]. The helices often exhibit a hydrophobic face, which interacts with core

lipids within the membrane, flanked by positively charged residues that might interact with

the negatively charged lipid head groups (Figure 1B). This amphipathic moment often

repeats across the length of the protein (Figure 1C). This feature is a hallmark of the late

embryonic abundant (LEA) class of ExTol proteins [21]. Human membrane interacting

proteins, like alpha-synuclein, also exhibit a similar repeat motif [22]. Such parallels in

sequence features suggest that orthogonal expression of ExTol proteins may confer their

protective properties to other biological systems, like human cells.

Indeed, previous reports have shown that some of these ExTol proteins provide similar

protective properties when expressed heterologously. Expressing tardigrade-specific

IDPs (TDPs) in bacteria provides protection against desiccation [3]. Anhydrin, an IDP

upregulated by Aphelenchus avenae during dehydration, limits PABPN1-A17

aggregation, a hallmark of oculopharyngeal muscular dystrophy, when expressed in a

human cell line [7]. Similar reports demonstrate protection of human cells provided by

expression of LEA proteins originating from the African midge [23] and by TDPs from

tardigrades [24]. These findings reinforce the promise of transferring protective phenotypes

from their native contexts to human cells.

Taken together, these data indicate a potential avenue for using this class of proteins to

protect biological systems in a controlled manner. To test this hypothesis, we generated

approximately 300 sequences inspired by ExTol proteins and examined their ability to

protect against apoptotic cell death. Our results show that human proteins containing

sequence hallmarks of ExTol proteins as well as de novo designed condensate forming

proteins can protect human cells from chemically-induced apoptosis. We also

demonstrate that condensate induction using synthetic modifications can strengthen

apoptotic protection. Finally, we show that key components of the apoptotic pathway can

localize to inducible condensates, suggesting enzyme sequestration could be a

mechanism of action.

RESULTS

Assembling a library of stress tolerance proteins

Based on the hallmark sequence features of ExTol proteins, we searched the published literature to identify proteins that not only contain these features but also exhibit some evidence for protection of biological structures. The library was generated based on over one hundred naturally-occurring proteins based on five lines of evidence for stress tolerance association, including: change in expression during extremotolerance; knockdown or overexpression changing the tolerant state; in vitro protection; orthogonal expression transferring tolerance; and sequence similarity to other proteins with these features (Figure 1D and Supplemental Table S1). We also examined our sequences for the presence of repetitive elements and intrinsic disorder (Supplemental Figure 1).

A PFAM domain analysis of the identified proteins reveals almost half of the identified targets have two key domains: Late Embryonic Abundant (LEA) and Apolipoprotein E (ApoE) domains (Figure 1E, Supplemental Table S1). LEA proteins are well known to be upregulated during the last stages of plant seed maturation [25]. This domain is thought to protect the seed embryo during transport to its new destination. There is also evidence that this domain has been co-opted by bacteria as protection from desiccation and radiation [26]. ApoE, in contrast, is not commonly discussed in the context of extremotolerance. ApoE proteins interact with lipid particles to allow for lipid reuptake into cells. Mutations within human ApoE have been identified as strong genetic predictors

of Alzheimer's disease in human populations [27].

While seemingly unrelated to other proteins in our

study, we chose to include human ApoE and its

various mutations as ExTol proteins due to the

prominence of the ApoE domain in the PFAM

analysis and since they exhibit similar features as

other selected proteins (e.g. their lipid interaction

properties and phase separation tendencies [28]).

Designing novel extremotolerance-associated

proteins

To test hypotheses about sequence features and

their relationship to phase separation and

apoptosis, we designed novel proteins based on

the set of ExTol proteins. These new designs take

3 approaches (Figure 2): truncation of proteins to

focus on repetitive, intrinsically disordered, and

amphipathic regions; isolation of individual helices

with desired sequence features; and de novo

design of alpha helix-containing proteins that self-

assemble into higher order structures. Through

these designs, we sought to identify the minimal

Figure 2: Design based on candidate extremotolerance-associated proteins (A) Truncations were made to isolate regions of repetitive and amphipathic sequence. (B) Small helices were selected and altered based on their hydrophobic moment. (C) Synthetic alpha helix-containing proteins designed to oligomerize. (D) Venn diagram highlighting the library that was tested.

elements required for protection and phase separation.

First, we identified regions within natural proteins that contain intrinsic disorder [29],

repetitive sequences [30], and amphipathic helices [31]. Figure 2A illustrates the use of

HHRepID on the sequence of full length human ApoE. Based on this analysis, we

identified the repeating domain within ApoE for further testing. Other ExTol proteins were

similarly modified and examined.

We also included short peptides based on the consensus sequence of repetitive regions

that have previously been shown to be protective [32]. To test if any of the sequence

features we identified are important for apoptosis protection, we altered these features

within these short peptides. For example, we altered the amphipathic character of these

helices (Figure 2B) to determine if apoptosis protection was changed.

To examine whether generic condensate formation is sufficient to confer apoptosis

protection, we also examined completely synthetic alpha helix-containing proteins that

were designed to form repetitive structure with sequences orthogonal to naturally-

occurring proteins [33]. These designed helical repeat (DHR) proteins interact along the

axis of the helix, forming long spiraling fibrils, but may also potentially self-interact along

the face of the protein based on a pattern of positive and negative charges (Figure 2C).

This ability to oligomerize and form multivalent complexes, which may then interact via

weak non-covalent interactions, is an established determinant of condensate formation

[15].

Together, our library consists of 104 naturally-occurring proteins, 166 truncations or isolations of helical regions, and 19 de novo designs (Figure 2D). These proteins interrogate several hypotheses about how disordered proteins may protect human cells against apoptosis.

Apoptosis assay identifies extremotolerance-associated proteins that protect human cells

To model stress in a mammalian cell system, we induced cellular apoptosis by treatment with the chemotherapeutic camptothecin (CPT), a DNA topoisomerase inhibitor well known to trigger apoptosis [34-35]. Under these conditions, protection against apoptosis could be achieved through a number of avenues, including protection against the DNA damage caused by CPT-induced topoisomerase inhibition [34], interruption of the signaling cascade, or even protection of the mitochondrial membrane whose rupture ultimately triggers apoptosis [36].

To test our designs for protection against apoptosis, we used a human cell-based apoptosis assay (Figure 3A). This assay uses HT-1080 fibrosarcoma cells from ATCC [37]. These cells are seeded in a 96-well plate and allowed to adhere overnight. The next day, the cells are co-transfected with a plasmid encoding the ExTol candidate proteins of interest as well as a transfection marker. The following day, the media is swapped to one containing 13.2 μM CPT. After addition of CPT, caspase-3 and -7 activation is measured by microscopy every 45 minutes for 1 day using a fluorophore quencher pair attached to

either end of a peptide with the sequence "DEVD". As caspase-3 and -7 are activated,

the DEVD peptide is cleaved, releasing the fluorophore from the quencher. The

unquenched fluorophore localizes to the nucleus where it can be quantified by imaging

on a single-cell level. The single-cell resolution allows for the separation of transfected

and non-transfected cells based on the transfection marker. Only single cells with the

transfection marker were subjected to further analysis; Figure 3B shows sample images

highlighting the type of data being analyzed.

On every plate, an empty vector negative control is run as well as a positive control

containing the X-linked inhibitor of apoptosis (XIAP). This protein is known to bind to

caspase-3 and -7 to block their activity [38] . Figure 3C shows example data comparing

median DEVD peptide signal intensity over time after induction with CPT of four wells on

the same plate transfected with XIAP and empty vector. The results show a reduction in

DEVD peptide signal intensity when the XIAP-containing vector is transfected,

demonstrating that the assay can identify modulators of apoptosis.

To convert these time-resolved data to a measure of protection against apoptosis, the

maximum DEVD peptide signal intensity is calculated and normalized to a plate-matched

empty vector control (see methods for further details). The normalized apoptosis induction

intensity is shown in Figure 3D left. The data are slightly skewed towards apoptosis-

protection, which supports our hypothesis that the targeted proteins can protect against

apoptosis.

Figure 3: Screen for CPT resistance by extremotolerance-associated proteins (A) Flow diagram of the assay. (B) Representative images showing cells during the assay. Cells are identified using SiR-Hoechst marking the nucleus (blue). Transfected cells are identified by the presence of the mCherry marker (green). Finally, the degree of apoptosis is identified by the presence of the cleaved DEVD peptide (red). (C) Positive control showing apoptosis reduction upon expression of XIAP. (D) Bar graph highlighting the apoptosis resistance provided by different constructs, including zoomed in area to highlight the top hits in the assay. Source species of a sequence are abbreviated and shown in italics; abbreviations are drawn from Uniprot, including RAMVA (Ramazzottius varieornatus), HYPDU (Hypsibius dujardini or Hypsibius exemplaris historical naming), PARRC (Paramacrobiotus richtersi), ANDDA (Andrias davidianus), APHAV (Aphelenchus avenae), ARTSF (Artemia franciscana), CRAPL (Craterostigma plantagineum), and DEIRA (Deinococcus radiodurans).

To highlight the top-performing designs, the top 10% of the targets are shown in Figure

3D right. As expected, the XIAP positive control showed the strongest anti-apoptotic

effect. The two top-performing hits were ApoE designs. Human alpha-synuclein is the 5th

hit, which is expected as it is known to be both intrinsically disordered and anti-apoptotic

[39]. Additionally, several ApoE designs and short LEA helices are top hits. The data used

to generate these figures is included in Supplemental Table S2.

Known extremotolerance-associated proteins affect apoptosis

Among the top hits in apoptosis reduction are an IGKV protein highly expressed in mucus

of the Chinese giant salamander Andrias davidianus [40] that is used as a medical glue in

traditional Chinese medicine, an LEA protein from the nematode Aphelenchus avenae

previously shown to have the ability to reduce a form of protein aggregation associated

with oculopharyngeal muscular dystrophy in human cells [7], and a mitochondrial LEA

protein from the brine shrimp Artemia franciscana shown to protect human cells during

desiccation [23] (Figure 3D). The prominence of these ExTol proteins suggests the assay

can help identify other proteins that can protect cells from more general stresses beyond

CPT-induced apoptosis.

Short repetitive peptides and de novo designs modulate apoptosis

LEA repeat-containing peptides are among the top-performing hits in the screen. While

the original short LEA designs from Furuki et al. 2019 had a minor effect reducing

apoptosis, several of the designs tested here performed better (Supplemental Figure 2A)

[32]. As 22 amino acids is small and could lead to peptide stability issues, longer 50-amino

acid designs inspired by the same LEA proteins utilized by Furuki et al. were also

developed (Supplemental Figure 2B). Some of these designs were also able to reduce

apoptosis, including the PvLEA4 with extended repeats (notably PvLEA4 repeats

extended #2), which fell into the top hits of the entire library (Figure 3D and Supplemental

Figure 2B). These results suggest that optimization of synthetic designs yields proteins

with stronger function than naturally-occurring proteins.

While these proteins were designed to interrogate the relationship between sequence

and apoptosis protection in short peptides, little relationship was found with respect to the

sequence features we analyzed (Supplemental Figure 2C). Several designs were made

that would alter the hydrophobic moments of putative helices these peptides are predicted

to form. To examine if this sequence property relates to the stress-tolerance function of

these proteins, we attempted to correlate this property with assay performance. However,

no significant correlation was found, suggesting (at least in these cases and experimental

conditions) that degree of amphipathicity on its own is not the key determinant of stress-

tolerance function of these proteins.

The repeat region of ApoE comprises its minimal anti-apoptotic unit

The top two hits in the assay were ApoE4 (47-300) and ApoE2 (47-300). These truncated

forms of ApoE maintain the repeat domains shown in Figure 4A, while removing the signal

peptide and terminal regions. This suggests that the repeat region of ApoE accounts for

its core anti-apoptotic functionality. Interestingly, the full length ApoE2 and E4 do not

show up as top hits, suggesting that these truncated forms improve on the apoptosis

protection provided by the full-length variants (Figure 4B).

Based on the location of the signal peptide, it is expected that ApoE4 should localize to

the secretory pathway whereas the truncated form may be cytoplasmic. To test this

hypothesis, cells were subjected to Western blot and immunofluorescence analysis using

commercially available antibodies (Abcam ab24139; Figure 4C and Supplemental Figure

3A). The Western blot results show that both the full-length and truncated forms of ApoE4

are expressed at the expected size (Supplemental Figure 3A). The immunofluorescence

shows co-localization between the ApoE4 full length protein and the endoplasmic

reticulum (marked by Thermo's PDI antibody MA3-019, Figure 4C). However, the

truncated form did not show the same co-localization, suggesting as expected that it may

not enter the secretory pathway as the full-length version does.

Strikingly, the truncated version appeared to form more pronounced puncta throughout

the cytoplasm of the cell, suggesting the formation of potentially phase-separated

condensates. In screening through the point mutants of ApoE also tested here, we found

a similar condensate phenotype with the full length ApoE3 D125I mutation (Supplemental

Figure 3B). Together, these data suggest that ApoE may be able to support condensate

formation and confer apoptosis protection in part via the protective compartment

mechanism (Figure 1A bottom).

Figure 4. ApoE truncations and synthetic condensates further reduce apoptosis. (A) Gene diagram highlighting the repetitive region of ApoE. (B) Boxplots highlighting the apoptosis assay performance of the full length and truncated forms of ApoE2 and E4. (C) Immunofluorescence of ApoE4, full length and truncated forms. White indicates colocalization, which is more apparent with the full-length protein. The truncated form appears to exhibit larger and more distinguishable puncta. (D) Diagram of the induced multi-valency systems implemented here, including the light-inducible system (top) and the chemically-inducible system (bottom). For the chemically-induced system, we used two FKBP domains. (E) Condensate formation assay using the light-inducible system, including the PARRC CAHS2 positive control showing condensates only upon light-induction and several ApoE examples that form condensates both before and after induction. (F) Condensate formation assay using the chemically-inducible system, including the PARRC CAHS2 positive control showing condensates only upon added dimerizer and ApoE4 (47-300) exhibiting condensates both before and after induction. (G) Apoptosis data showing reduction in apoptosis upon condensate induction with dimerizer. Data are normalized to mean apoptosis level of uninduced (i.e. no dimerizer) sample for each construct.

Induced multivalency to probe the relation between condensates and apoptosis

We made use of established condensate engineering tools [41] to test for latent phase

separation propensity among a selection of ExTol proteins representing the range of

apoptosis effects. We used both light- and chemically-inducible systems of multivalency.

In the light system, a ferritin core (comprised of self-assembled monomers, FTH1 domain-

eGFP-iLID) becomes decorated with SspB-modified proteins upon exposure to blue light

[42] (Figure 4D top). In the chemical system, multivalency (i.e. multiple arms of the ExTol

protein of interest) is induced using a dimerization agent [43] that brings together FKBP-

modified proteins [41] (Figure 4D bottom). These inducible methods increase the valency

of a system, and thus work to favor phase separation. In this way, they can help reveal

any latent ability to form condensates that is not immediately observable without

modification.

Condensates were indeed observed in a number of ExTol proteins (Supplemental Figure

4), including ApoE, ARTSF LEA3m, DHR81, HYPDU CAHS3, and PARRC CAHS2 using

the light-inducible system. For those proteins with no visible condensates, the

experimental conditions may be insufficient to support phase separation. PARRC CAHS2

exhibited the most dramatic phenotype, which is consistent with recent work describing

the gel forming capabilities of CAHS proteins [44-45] (Figure 4E&F). ApoE variants were of

interest, based on the puncta observed in the top performing ApoE constructs. All

versions of ApoE tested formed puncta without activation, consistent with our previous

observation using untagged native ApoE (Figure 4C). Upon light- or chemical-activation,

there is a noticeable increase in condensates, exemplifying how phase separation can be

enhanced with increased multivalency (Figure 4E&F).

To examine how enhanced phase separation may affect apoptosis, we performed the

apoptosis assay with ExTol proteins that were modified for chemically-induced

multivalency. Chemical induction was more compatible with the apoptosis assay, while

enabling the same increase in condensates seen with the light-induced system (Figure

4F). We observed that the presence of the chemical dimerizer had little to no effect on

apoptosis in cells transfected with the empty vector, with a construct encoding the

dimerization domains alone, or with constructs encoding an ExTol protein that was not

strongly anti-apoptotic (e.g. PARRC CAHS2) (Figure 4G; Supplemental Figure 5). In

contrast, with constructs encoding a strongly anti-apoptotic ExTol protein – including

ApoE, ARTSF LEA3m, and DHR81 – added dimerizer corresponded with reduced

apoptosis in transfected cells (Figure 4G; Supplemental Figure 5). These data suggest

that condensate formation plays a role in the anti-apoptotic effect of these proteins.

Synthetic DHR81 condensates sequester caspase-7

To begin elucidating the mechanism by which ExTol proteins protect against apoptosis,

particularly the hypothesis of membraneless compartments that sequester essential

effectors, we examined the ability for the various condensates to partition apoptotic

signaling molecules, identifying the synthetic protein DHR81 as the only ExTol protein

capable of doing so. DHR81 is a synthetic alpha helical-rich protein designed to promote

self-interaction. It is formed from four helical hairpin repeat regions; the core of the protein

is hydrophobic to ensure consistent folding, while

its outside is decorated with checkerboard positive

and negative charges (Figure 5A) [33]. This structure

ensures the protein is well folded, while also

enabling the potential for higher order structure

formation via electrostatic interactions. Not only

did DHR81 support condensates in the

chemically-inducible condensate system (Figure

5B), it also exhibited a strong increase in apoptotic

protection after induced-multivalency (Figure 5C;

Supplemental Figure 5&6A).

As such, we chose DHR81, along with a few other

ExTol proteins with notable phase separation

phenotypes (e.g. ApoE4 and PARRC CAHS2), to

further explore how their condensates may confer

an anti-apoptotic effect. Specifically, we used

immunofluorescence to examine whether they

sequester major components of apoptotic

signaling, namely caspase-3 and -7. The results

showed that activated caspase-7 localized to

DHR81 condensates but not PARRC CAHS2 or

ApoE4 condensates (Figure 5D; Supplemental

Figure 5. DHR81 condensate formation and apoptosis reduction. (A) Diagram of DHR81, illustrating a charge distribution that can support oligomerization. (B) Sample images highlighting how the synthetic protein DHR81 supports chemically-induced condensates. (C) Apoptosis is further reduced in the presence of dimerizer (i.e. with induced DHR81 condensates). The FKBP-modifications alone, which enable chemically-induced multi-valency, have a minimal effect on apoptosis. (D) Activated caspase-7 (green) co-localizes with DHR81 condensates but not PARRC CAHS2 or ApoE4 (47-300) condensates (magenta). White indicates colocalization.

Figure 5C), while caspase-3 did not localize to any of the examined condensates

(Supplemental Figure 6B). Even though condensate properties may confound (i.e.

physically occlude) antibody binding during immunofluorescence, the positive DHR81

result suggests sequestration of key apoptotic factors can indeed be a potential

mechanism by which these condensates reduce apoptosis.

Discussion

To broadly evaluate the ability of ExTol proteins to protect orthogonal systems from

environmental stress, we studied over one hundred different ExTol proteins. We used

these as a basis to design approximately 300 variants, including truncations and fusion

proteins. These designs were evaluated on their ability to protect human HT-1080 cells

against CPT-induced apoptosis. The results showed that known ExTol proteins, ApoE

variants, and the synthetic DHR81 protein could modulate apoptosis levels when

transferred to human cells.

Protection against apoptosis is a key feature of many ExTol proteins. As apoptosis is a

common cellular response to stress, the ability to mitigate this response is important in

providing cell-autonomous protection in stressful environments. While the mechanism

resilient organisms use to survive is still not fully understood, it is becoming increasingly

clear that phase separation may play an important role. Some ExTol proteins are

proposed to form expansive higher order structures (e.g. gels or glassy systems) that

reduce the viscosity of the cytoplasm, especially in desiccated states [3]. Other reports

show that these proteins could mediate formation of protective phase-separated

compartments (i.e. condensates) that sequester damage-inducing proteins and/or protect

important targets [11]. Based on our data, it appears the latter may be at play for proteins

like ApoE, ARTSF LEA3m, and DHR81, which exhibit anti-apoptotic effects while also

being condensate prone.

Condensates alone are not sufficient for apoptosis protection. While we observe

condensate formation in many of our top performing hits, we also observe condensate

formation in targets like HYPDU CAHS3 and PARRC CAHS2, which do not reduce

apoptosis very much. Additionally, some proteins like PARRC CAHS1 (101-189) are

truncated in ways predicted to reduce condensate formation[44, 46-47], but they still perform

well in apoptosis protection. These observations suggest yet-to-be-discovered

mechanisms are likely responsible for apoptosis protection in these cases. However, for

those proteins that exhibit both anti-apoptotic effects as well as elevated phase separation

propensities, increasing the amount of condensates via an engineered inducible system

led to stronger apoptosis protection (Supplemental Figure 5). In these cases,

condensates could be a key part of the protective mechanism.

Specifically, we observed that activated caspase-7 partitions into DHR81 condensates,

suggesting caspase sequestration as a potential anti-apoptotic mechanism. Phase-

separated condensates and higher order assemblies are already implicated in mediating

apoptosis: stress granules inhibit apoptosis by suppressing MAPK pathways [48]; the

apoptosome relies on Apaf-1 multimerization, which has been reported to undergo

amyloid fibrillation in specific conditions [49]; and stress-dependent phase separation

modulates proteasome formation, whose function affects apoptosis [50-51]. It is interesting

to note that we only observe clear caspase-7 partitioning into DHR81 condensates,

suggesting the possibility of a specific binding interaction in this case. ApoE and ARTSF

LEA3m condensates may be sequestering other apoptotic factors or using a different

mechanism to reduce apoptosis.

It is also possible that these ExTol proteins achieve their anti-apoptotic effect by

implementing a more general physical mechanism, namely slowing the intracellular

mobility of biological effectors. There are various reports describing how the effective

viscosity within the cytoplasm can be increased in response to external stresses [4-5] or

mediated by protein composition, specifically intrinsically disordered proteins [52]. Notably,

a pervasive cell-spanning gel or network is not necessary to achieve such an effect; it is

possible that the presence of condensates, coupled with their ability to selectively partition

biomolecules (i.e. control what biomolecules can enter and diffuse through them), could

broadly slow biomolecular movement. That is, by forcing different proteins to either be

trapped in them or diffuse around them, condensates can decrease protein mobility and

perturb e.g. signaling pathways. Our results showing a lack of caspase-3 and -7

sequestration by most of our protein condensates could thus support this more general

mechanism of condensates perturbing biology, though they certainly do not disqualify

sequestration of other effectors not examined in this study.

Condensates are becoming increasingly implicated in disease [13], particularly when their

structural metastability and reversibility is dysregulated (e.g. through mutation). Notably,

drugs that block the formation of amyloid fibrils and plaques, which are products of phase

separation dysregulation, are promising yet strongly debated [53] therapies for many

neurological diseases. As such, our observations concerning ApoE proteins, while

unexpected from an extremotolerance perspective, are particularly salient. We found that

expression of a cytoplasmic form of ApoE is protective against apoptosis. In particular,

the truncated repeat region of ApoE is sufficient and shows the strongest anti-apoptotic

activity of all designs we tested. Others have shown similar effects for exogenously added

ApoE-containing liposomes protecting rat neurons against apoptosis [54], though we show

the effect through overexpression of ApoE inside of human cells. When combined with

our condensate results and the clinical relevance of ApoE in Alzheimer's disease [55], these

observations suggest the anti-apoptotic effect observed here warrants further study, to

further elucidate its role in Alzheimer's disease progression.

Altogether, our work expands upon the understanding of the relationship between

condensate formation and cell physiology, demonstrating how condensate accumulation

can be beneficial in reducing apoptosis while also identifying diverse new protein domains

that confer protection. The proteins and concepts presented here demonstrate the utility

of this class of proteins for engineering new properties into cells and ultimately higher

organisms.

Methods

Cell culturing and maintenance

HT-1080 cells (ATCC, CCL-121) were maintained in EMEM medium supplemented with

10% FBS and Penicillin-Streptomycin according to ATCC's protocol. For additional details

see Supplemental Methods.

DNA Preparation

Apoptosis constructs were synthesized by Twist Bioscience and transformed into DH5α

cells for amplification and plasmid purification. Condensate assay constructs were

subcloned from these synthesized constructs into vectors containing the appropriate

chemically- or light-inducible condensate elements. DNA was prepared using the

QIAGEN Plasmid Plus Midi Kit (Qiagen 12943) for transfection grade DNA purification.

See Supplemental Methods for additional details.

Apoptosis assay

200 µL of HT-1080 cells at 7.5×10^4 cells per mL (15000 cells per well) were added to

each well of a microscope grade 96 well plate (Corning 3603). Plates were then

centrifuged at 330 rpm for 3 min (Eppendorf, Centrifuge 5810). Plates were then

incubated at 37°C for 24 hours. Cells were then transfected with appropriate DNA

constructs with the lipofectamine 3000 kit (Thermo Fisher Scientific, L3000015) as

specified by the manufacturer. 24 hours after transfection, the transfection media was

removed and replaced with FluoroBrite DMEM Media (Thermo A1896702) supplemented

with 1x GlutaMAX (Thermo 35-050-061) and 10% FBS (Thermo Fisher Scientific,

10082147). This media also contained 13.2 µM (S)-(+)-Camptothecin (Sigma C9911-

100MG), 0.3 µM SiR-DNA stain (SpiroChrome CY-SC007), and 1.25 µM of the DEVD

peptide for detecting caspase activation (Sartorius 4440). The plates were sealed with

sterile Breathe-Easy film (USA scientific 9123-6100) and incubated at 37°C with 5% CO_2

for 30 minutes. After 30 min incubation, the plates were placed in an ImageXpress-

confocal plate imager (Molecular Devices) at 37°C with 5% CO_2 and saturated humidity

through the presence of a heated water bath in the chamber. Images of all wells were

acquired every 45 minutes and stored on a remote server for later analysis. See

Supplemental Methods for further details on this method as well as the image analysis

pipeline used.

Confocal microscopy

Confocal microscopy was performed at the Nikon Imaging Center at Harvard Medical

School using a Nikon Ti fluorescence microscope attached to a Yokogawa W1 spinning

disk confocal. See Supplemental Methods for further details on the equipment and

settings used.

Light-inducible condensate assay

HEK293-T cells were seeded on 96 well glass bottom plates from Cellvis (P96-1.5H-N).

Cells within each well were transfected with 50 ng of the LLPS1-iLID::EGFP::FTH1

(Addgene plasmid #122149) construct containing the core FTH1 scaffolding domain

attached to iLID for light-inducible interactions. The cells were also transfected with 50 ng

of a construct containing the target protein fused to mCherry and SspB. The transfection

was allowed to continue overnight before the media was swapped to FluoroBrite media

supplemented with 10% FBS and 1x GlutaMAX. Pre-induction images were acquired

using the settings listed in the Supplemental Methods for mCherry and eGFP. The cells

were then exposed to 1 second of the DAPI imaging protocol described in Supplemental

Methods before new images were captured on the mCherry and eGFP settings.

Chemically-inducible condensate assay

HT-1080 cells were seeded on 96 well glass bottom plates from Cellvis (P96-1.5H-N).

Cells within each well were transfected with 50 ng of target construct fused to a mCherry-

2xFKBP domain. The transfection was allowed to continue overnight before the media

was swapped to FluoroBrite media supplemented with 10% FBS, 1x GlutaMAX, and 300

nM SiR-Hoechst. The cells were then imaged using the settings listed in the Supplemental

Methods for mCherry and SiR-Hoechst for the pre-induction images. Then, B/B dimerizer

(AP20187 ligand; TaKaRa 635058) was added to a final concentration of 500 nM in the

well. Imaging continued for 15 minutes with new images every 45 seconds.

Caspase assay with chemically-induced condensates

These caspase assays are performed similarly to the standard caspase assays with a

few key differences. Briefly, HT-1080 cells were seeded as normal on to 96 well plates

and allowed to adhere overnight. The next day, 12.5 ng of nuclear mCherry marker and

37.5 ng of the target construct fused to mCherry-2xFKBP was transfected into the cells

as described above. The next day, 500 nM of the B/B dimerizer (TaKaRa 635058) was

added 2 hours before the start of the assay (or an Ethanol vehicle control was added).

The media was then removed and replaced with the apoptosis assay media described

above with the addition of 500 nM B/B dimerizer or a vehicle control. See Supplemental

Methods for further details.

Abbreviations

ExTol: extremotolerance-associated

IDP: Intrinsically Disordered Protein

LEA: Late Embryonic Abundant (protein)

ApoE: Apolipoprotein E

DHR: Designed helical repeat (protein)

CPT: Camptothecin

ATCC: American Type Culture Collection

XIAP: X-linked inhibitor of apoptosis

CAHS: Cytosolic Abundant Heat Soluble (protein)

FTH1: Ferritin Heavy Chain 1

iLID: improved light inducible dimer

SspB: Stringent Starvation Protein B

eGFP: Enhanced Green Fluorescent Protein

FKBP: FK506 binding protein

PvLEA: Polypedilum vanderplanki Late Embryonic Abundant (Protein)

MAPK: Mitogen-activated protein kinase

Apaf: Apoptotic protease activating factor

EMEM: Eagle's Minimum Essential Medium

FBS: Fetal bovine serum

DMEM: Dulbecco's Modified Eagle Medium

Author Information.

Correspondences should be sent to: Pamela Silver, Harvard Medical School Department

of Systems Biology 200 Longwood Avenue Warren Alpert Building Boston, MA 02115;

and to Roger Chang, Albert Einstein College of Medicine Department of Systems &

Computational Biology 1301 Morris Park Avenue Price Center Bronx, NY 10461.

Author Contributions

M.T.V. and D.T.N. designed and performed experiments, analyzed the data, and wrote the paper. N.N.T. analyzed the data and wrote the paper. M.E.O provided technical support. N.J.R., K.B.B., N.P.B., and S.L., designed constructs. D.B. and J.C.W. managed the teams and provided key insight. D.S.M., R.L.C., and P.A.S., designed constructs, managed teams, and wrote the paper.

Conflict of interest statement

Conflicts of Interest: None.

Supporting information

Supplemental figures and table descriptions: Contains Supplemental Figures 1 - 6 referenced in the main text, as well as full descriptions of the Supplemental Tables. This includes an analysis of the disorder and repetitive distribution of the identified ExTol proteins, extended analysis of the relationship between the synthetic LEA proteins and apoptosis protection, additional data on the ApoE Western blot and immunofluorescence experiments, data regarding our screen for condensate formation across the spectrum of apoptosis related constructs, a broader look at condensate formation as it relates to apoptosis protection and condensate sequestration, and extended data on DHR81.

Supplemental Methods: Contains detailed information about the methodology used in this study.

Supplemental Table S1: Reports sequences of all the proteins and designs discussed in

this work.

Supplemental Table S2: Reports all normalized apoptosis data collected throughout this

work.

Acknowledgements

Research was sponsored by the NSF, DARPA, the Wyss Institute for Biologically inspired

engineering, and the Army Research Office and was accomplished under Cooperative

Agreement Number W911NF-19-2-0017. M.T.V. was sponsored by the National Science

Foundation under Grant No. 2010370. D.T.N., N.N.T., M.E.O., N.J.R., K.P.B., J.C.W,

D.S.M., and P.A.S. were supported by the Harvard Medical School Department of

Systems Biology. J.C.W. was further supported by the Harvard Medical School

Laboratory of Systems Pharmacology. K.P.B. and D.S.M. were further supported by CZI

Neurodegenerative Challenge Ben Barres Early Career Award (CZI2018-191853). N.P.B

was further supported by the HHMI Hanna Gray fellowship (grant GT11817). We would

like to acknowledge Clarence Yapp for help with microscopy as well as the NCI U54-

CA225088 grant and a gift from the Ludwig Institute for Cancer Research to the

Laboratory for Systems Pharmacology. We would like to acknowledge the Nikon Imaging

Center at Harvard Medical Center for providing microscope support. The views and

conclusions contained in this document are those of the authors and should not be

interpreted as representing the official policies, either expressed or implied, of any funding

agency including the NSF, DARPA, the Army Research Office, or the U.S. Government.

The U.S. Government is authorized to reproduce and distribute reprints for Government

purposes notwithstanding any copyright notation herein. Visual table of contents entry

was created with support from BioRender.com.

593
594   **Bibliography & References**
595

1.   Jonsson, K. I.; Rabbow, E.; Schill, R. O.; Harms-Ringdahl, M.; Rettberg, P., Tardigrades
survive exposure to space in low Earth orbit. *Curr Biol* **2008,** *18* (17), R729-R731.
2.   Thevelein, J. M., Regulation of trehalose mobilization in fungi. *Microbiol Rev* **1984,** *48*
(1), 42-59.
3.   Boothby, T. C.; Tapia, H.; Brozena, A. H.; Piszkiewicz, S.; Smith, A. E.; Giovannini, I.;
Rebecchi, L.; Pielak, G. J.; Koshland, D.; Goldstein, B., Tardigrades Use Intrinsically Disordered
Proteins to Survive Desiccation. *Mol Cell* **2017,** *65* (6), 975-984 e5.
4.   Dijksterhuis, J.; Nijsse, J.; Hoekstra, F. A.; Golovina, E. A., High viscosity and anisotropy
characterize the cytoplasm of fungal dormant stress-resistant spores. *Eukaryot Cell* **2007,** *6* (2),
157-70.
5.   Parry, B. R.; Surovtsev, I. V.; Cabeen, M. T.; O'Hern, C. S.; Dufresne, E. R.; Jacobs-Wagner,
C., The bacterial cytoplasm has glass-like properties and is fluidized by metabolic activity. *Cell*
2014, *156* (1-2), 183-94.
6.   Hand, S. C.; Menze, M. A.; Toner, M.; Boswell, L.; Moore, D., LEA proteins during water
stress: not just for plants anymore. *Annu Rev Physiol* **2011,** *73*, 115-34.
7.   Chakrabortee, S.; Meersman, F.; Kaminski Schierle, G. S.; Bertoncini, C. W.; McGee, B.;
Kaminski, C. F.; Tunnacliffe, A., Catalytic and chaperone-like functions in an intrinsically
disordered protein associated with desiccation tolerance. *Proc Natl Acad Sci U S A* **2010,** *107*
(37), 16084-9.
8.   Boothby, T. C., Water content influences the vitrified properties of CAHS proteins. *Mol*
Cell **2021,** *81* (3), 411-413.
9.   Arakawa, K.; Numata, K., Reconsidering the "glass transition" hypothesis of intrinsically
unstructured CAHS proteins in desiccation tolerance of tardigrades. *Mol Cell* **2021,** *81* (3), 409-
410.
10.   Crilly, C. J.; Brom, J. A.; Warmuth, O.; Esterly, H. J.; Pielak, G. J., Protection by
desiccation-tolerance proteins probed at the residue level. *Protein Sci* **2021**.
11.   Belott, C.; Janis, B.; Menze, M. A., Liquid-liquid phase separation promotes animal
desiccation tolerance. *Proc Natl Acad Sci U S A* **2020,** *117* (44), 27676-27684.
12.   Hyman, A. A.; Weber, C. A.; Julicher, F., Liquid-liquid phase separation in biology. *Annu*
Rev Cell Dev Biol **2014,** *30*, 39-58.
13.   Shin, Y.; Brangwynne, C. P., Liquid phase condensation in cell physiology and disease.
Science **2017,** *357* (6357).
14.   Boeynaems, S.; Alberti, S.; Fawzi, N. L.; Mittag, T.; Polymenidou, M.; Rousseau, F.;
Schymkowitz, J.; Shorter, J.; Wolozin, B.; Van Den Bosch, L.; Tompa, P.; Fuxreiter, M., Protein
Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol* **2018,** *28* (6), 420-435.
15.   Choi, J. M.; Holehouse, A. S.; Pappu, R. V., Physical Principles Underlying the Complex
Biology of Intracellular Phase Transitions. *Annu Rev Biophys* **2020,** *49*, 107-133.

16.   Jiang, S.; Fagman, J. B.; Chen, C.; Alberti, S.; Liu, B., Protein phase separation and its role
in tumorigenesis. *Elife* **2020,** *9*.

17.   Boija, A.; Klein, I. A.; Young, R. A., Biomolecular Condensates and Cancer. *Cancer Cell*
2021, *39* (2), 174-192.

18.   Boija, A.; Klein, I. A.; Sabari, B. R.; Dall'Agnese, A.; Coffey, E. L.; Zamudio, A. V.; Li, C. H.;
Shrinivas, K.; Manteiga, J. C.; Hannett, N. M.; Abraham, B. J.; Afeyan, L. K.; Guo, Y. E.; Rimel, J.
K.; Fant, C. B.; Schuijers, J.; Lee, T. I.; Taatjes, D. J.; Young, R. A., Transcription Factors Activate
Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell* **2018,** *175* (7),
1842-1855 e16.

19.   Hubstenberger, A.; Courel, M.; Benard, M.; Souquere, S.; Ernoult-Lange, M.; Chouaib, R.;
Yi, Z.; Morlot, J. B.; Munier, A.; Fradet, M.; Daunesse, M.; Bertrand, E.; Pierron, G.;
Mozziconacci, J.; Kress, M.; Weil, D., P-Body Purification Reveals the Condensation of Repressed
mRNA Regulons. *Mol Cell* **2017,** *68* (1), 144-157 e5.

20.   Tolleter, D.; Jaquinod, M.; Mangavel, C.; Passirani, C.; Saulnier, P.; Manon, S.; Teyssier,
E.; Payet, N.; Avelange-Macherel, M. H.; Macherel, D., Structure and function of a mitochondrial
late embryogenesis abundant protein are revealed by desiccation. *Plant Cell* **2007,** *19* (5), 1580-
9.

21.   Leon, D., III, Occurrence of a Repeating 11-Mer Amino Acid Sequence Motif in Diverse
Organisms. *Protein & Peptide Letters* **2001,** *8* (2), 115-122.

22.   Bussell, R., Jr.; Eliezer, D., A structural and functional role for 11-mer repeats in alpha-
synuclein and other exchangeable lipid binding proteins. *J Mol Biol* **2003,** *329* (4), 763-78.

23.   Li, S.; Chakraborty, N.; Borcar, A.; Menze, M. A.; Toner, M.; Hand, S. C., Late
embryogenesis abundant proteins protect human hepatoma cells during acute desiccation.
Proc Natl Acad Sci U S A **2012,** *109* (51), 20859-64.

24.   Tanaka, S.; Tanaka, J.; Miwa, Y.; Horikawa, D. D.; Katayama, T.; Arakawa, K.; Toyoda, A.;
Kubo, T.; Kunieda, T., Novel mitochondria-targeted heat-soluble proteins identified in the
anhydrobiotic Tardigrade improve osmotic tolerance of human cells. *PLoS One* **2015,** *10* (2),
e0118272.

25.   Mertens, J.; Aliyu, H.; Cowan, D. A., LEA Proteins and the Evolution of the WHy Domain.
Appl Environ Microbiol **2018,** *84* (15).

26.   Makarova, K. S.; Aravind, L.; Wolf, Y. I.; Tatusov, R. L.; Minton, K. W.; Koonin, E. V.; Daly,
M. J., Genome of the extremely radiation-resistant bacterium Deinococcus radiodurans viewed
from the perspective of comparative genomics. *Microbiol Mol Biol Rev* **2001,** *65* (1), 44-79.

27.   Yamazaki, Y.; Zhao, N.; Caulfield, T. R.; Liu, C. C.; Bu, G., Apolipoprotein E and Alzheimer
disease: pathobiology and targeting strategies. *Nat Rev Neurol* **2019,** *15* (9), 501-518.

28.   Raulin, A. C.; Kraft, L.; Al-Hilaly, Y. K.; Xue, W. F.; McGeehan, J. E.; Atack, J. R.; Serpell, L.,
The Molecular Basis for Apolipoprotein E4 as the Major Risk Factor for Late-Onset Alzheimer's
Disease. *J Mol Biol* **2019,** *431* (12), 2248-2265.

29.   Erdos, G.; Dosztanyi, Z., Analyzing Protein Disorder with IUPred2A. *Curr Protoc*
Bioinformatics **2020,** *70* (1), e99.

30.   Biegert, A.; Soding, J., De novo identification of highly diverged protein repeats by
probabilistic consistency. *Bioinformatics* **2008,** *24* (6), 807-14.

31.   Eisenberg, D.; Weiss, R. M.; Terwilliger, T. C., The helical hydrophobic moment: a
measure of the amphiphilicity of a helix. *Nature* **1982,** *299* (5881), 371-4.

32.   Furuki, T.; Niwa, T.; Taguchi, H.; Hatanaka, R.; Kikawada, T.; Sakurai, M., A LEA model
peptide protects the function of a red fluorescent protein in the dry state. *Biochem Biophys Rep*
2019, *17*, 27-31.
33.   Brunette, T. J.; Parmeggiani, F.; Huang, P. S.; Bhabha, G.; Ekiert, D. C.; Tsutakawa, S. E.;
Hura, G. L.; Tainer, J. A.; Baker, D., Exploring the repeat protein universe through computational
protein design. *Nature* **2015,** *528* (7583), 580-4.
34.   Li, F.; Jiang, T.; Li, Q.; Ling, X., Camptothecin (CPT) and its derivatives are known to target
topoisomerase I (Top1) as their mechanism of action: did we miss something in CPT analogue
molecular targets for treating human disease such as cancer? *Am J Cancer Res* **2017,** *7* (12),
2350-2394.
35.   Wall, M. E.; Wani, M. C.; Cook, C. E.; Palmer, K. H.; McPhail, A. T.; Sim, G. A., Plant
Antitumor Agents. I. The Isolation and Structure of Camptothecin, a Novel Alkaloidal Leukemia
and Tumor Inhibitor from Camptotheca acuminata1,2. *Journal of the American Chemical Society*
1966, *88* (16), 3888-3890.
36.   Bortner, C. D.; Cidlowski, J. A., Cellular mechanisms for the repression of apoptosis.
Annu Rev Pharmacol Toxicol **2002,** *42*, 259-81.
37.   Rasheed, S.; Nelson-Rees, W. A.; Toth, E. M.; Arnstein, P.; Gardner, M. B.,
Characterization of a newly derived human sarcoma cell line (HT-1080). *Cancer* **1974,** *33* (4),
1027-33.
38.   Obexer, P.; Ausserlechner, M. J., X-linked inhibitor of apoptosis protein - a critical death
resistance regulator and therapeutic target for personalized cancer therapy. *Front Oncol* **2014,**
4, 197.
39.   Li, W.; Lee, M. K., Antiapoptotic property of human alpha-synuclein in neuronal cell lines
is associated with the inhibition of caspase-3 but not caspase-9 activity. *J Neurochem* **2005,** *93*
(6), 1542-50.
40.   Geng, X.; Guo, J.; Zang, X.; Chang, C.; Shang, H.; Wei, H.; Xu, C., Proteomic analysis of
eleven tissues in the Chinese giant salamander (Andrias davidianus). *Sci Rep* **2019,** *9* (1), 16415.
41.   Bracha, D.; Walls, M. T.; Brangwynne, C. P., Probing and engineering liquid-phase
organelles. *Nat Biotechnol* **2019,** *37* (12), 1435-1445.
42.   Bracha, D.; Walls, M. T.; Wei, M. T.; Zhu, L.; Kurian, M.; Avalos, J. L.; Toettcher, J. E.;
Brangwynne, C. P., Mapping Local and Global Liquid Phase Behavior in Living Cells Using Photo-
Oligomerizable Seeds. *Cell* **2018,** *175* (6), 1467-1480 e13.
43.   Stanton, B. Z.; Chory, E. J.; Crabtree, G. R., Chemically induced proximity in biology and
medicine. *Science* **2018,** *359* (6380).
44.   Yagi-Utsumi, M.; Aoki, K.; Watanabe, H.; Song, C.; Nishimura, S.; Satoh, T.; Yanaka, S.;
Ganser, C.; Tanaka, S.; Schnapka, V.; Goh, E. W.; Furutani, Y.; Murata, K.; Uchihashi, T.; Arakawa,
K.; Kato, K., Desiccation-induced fibrous condensation of CAHS protein from an anhydrobiotic
tardigrade. *Sci Rep* **2021,** *11* (1), 21328.
45.   Malki, A.; Teulon, J. M.; Camacho-Zarco, A. R.; Chen, S. W.; Adamski, W.; Maurin, D.;
Salvi, N.; Pellequer, J. L.; Blackledge, M., Intrinsically Disordered Tardigrade Proteins Self-
Assemble into Fibrous Gels in Response to Environmental Stress. *Angew Chem Int Ed Engl* **2022,**
61 (1), e202109961.
46.   Hesgrove, C. S.; Nguyen, K. H.; Biswas, S.; Childs, C. A.; Shraddha, K.; Medina, B. X.;
Alvarado, V.; Yu, F.; Sukenik, S.; Malferrari, M.; Francia, F.; Venturoli, G.; Martin, E. W.;

Holehouse, A. S.; Boothby, T. C., Tardigrade CAHS Proteins Act as Molecular Swiss Army Knives
to Mediate Desiccation Tolerance Through Multiple Mechanisms. *bioRxiv* **2021**,
2021.08.16.456555.
47.   Tanaka, A.; Nakano, T.; Watanabe, K.; Masuda, K.; Kamata, S.; Yasui, R.; Kozuka-Hata, H.;
Watanabe, C.; Chinen, T.; Kitagawa, D.; Oyama, M.; Yanagisawa, M.; Kunieda, T., Stress-
dependent dynamic and reversible formation of cytoskeleton-like filaments and gel-transition
by tardigrade tolerance proteins. *bioRxiv* **2021**, 2021.10.02.462891.
48.   Arimoto, K.; Fukuda, H.; Imajoh-Ohmi, S.; Saito, H.; Takekawa, M., Formation of stress
granules inhibits apoptosis by suppressing stress-responsive MAPK pathways. *Nat Cell Biol*
2008, *10* (11), 1324-32.
49.   Rao, P. N.; Reddy, K. S.; Bhuyan, A. K., Amyloid fibrillation of human Apaf-1 CARD.
Biochemistry **2009,** *48* (32), 7656-64.
50.   Yasuda, S.; Tsuchiya, H.; Kaiho, A.; Guo, Q.; Ikeuchi, K.; Endo, A.; Arai, N.; Ohtake, F.;
Murata, S.; Inada, T.; Baumeister, W.; Fernandez-Busnadiego, R.; Tanaka, K.; Saeki, Y., Stress-
and ubiquitylation-dependent phase separation of the proteasome. *Nature* **2020,** *578* (7794),
296-300.
51.   Gupta, I.; Singh, K.; Varshney, N. K.; Khan, S., Delineating Crosstalk Mechanisms of the
Ubiquitin Proteasome System That Regulate Apoptosis. *Front Cell Dev Biol* **2018,** *6*, 11.
52.   Theillet, F. X.; Binolfi, A.; Frembgen-Kesner, T.; Hingorani, K.; Sarkar, M.; Kyne, C.; Li, C.;
Crowley, P. B.; Gierasch, L.; Pielak, G. J.; Elcock, A. H.; Gershenson, A.; Selenko, P.,
Physicochemical properties of cells and their effects on intrinsically disordered proteins (IDPs).
Chem Rev **2014,** *114* (13), 6661-714.
53.   Lalli, G.; Schott, J. M.; Hardy, J.; De Strooper, B., Aducanumab: a new phase in
therapeutic development for Alzheimer's disease? *EMBO Mol Med* **2021,** *13* (8), e14781.
54.   Hayashi, H.; Campenot, R. B.; Vance, D. E.; Vance, J. E., Protection of neurons from
apoptosis by apolipoprotein E-containing lipoproteins does not require lipoprotein uptake and
involves activation of phospholipase Cgamma1 and inhibition of calcineurin. *J Biol Chem* **2009,**
284 (43), 29605-13.
55.   Schellenberg, G. D.; D'Souza, I.; Poorkaj, P., The genetics of Alzheimer's disease. *Curr*
Psychiatry Rep **2000,** *2* (2), 158-64.
751
752
753
754

For Table of Contents Use Only:

Natural and designed proteins inspired by extremotolerant organisms can form condensates and attenuate apoptosis in human cells

Mike T. Veling[1,2,8], Dan T. Nguyen[1,2,8], Nicole N. Thadani[1], Michela E. Oster[1,2], Nathan J. Rollins[1,2], Kelly P. Brock[1], Neville P. Bethel[3,4], Samuel Lim[1,2], David Baker[3,4,5], Jeffrey C. Way[1,2], Debora S. Marks[1,6], Roger L. Chang[1,2,7,*], Pamela A. Silver[1,2,*]