Natural and designed proteins inspired by extremotolerant organisms can form

condensates and attenuate apoptosis in human cells

Mike T. Veling[1,2,7], Dan T. Nguyen[1,2,7], Nicole N. Thadani[1], Michela E. Oster[1,2], Nathan J.

Rollins[1,2], Kelly P. Brock[1], Neville P. Bethel[3,4], David Baker[3,4,5], Jeffrey C. Way[1,2], Debora

S. Marks[1,6], Roger L. Chang[1,2,8], Pamela A. Silver[1,2,8]

Affiliations

1. Department of Systems Biology, Harvard Medical School, Boston, MA, 02115, USA;

2. Wyss Institute for Biologically Inspired Engineering, Boston, MA, 02115, USA;

3. Institute for Protein Design, University of Washington, Seattle, WA, 98195, USA;

4. Department of Biochemistry, University of Washington, Seattle, WA, 98195, USA;

5. Howard Hughes Medical Institute, University of Washington, Seattle, WA, 98195, USA;

6. Broad Institute of Harvard and MIT, Cambridge, MA, 02142, USA;

7. Co-first authors

8. Co-corresponding authors

ABSTRACT

Many organisms can survive extreme conditions and successfully recover to normal life. This extremotolerant behavior has been attributed in part to repetitive, amphipathic, and intrinsically disordered proteins that are upregulated in the protected state. Here, we assemble a library of approximately 300 naturally-occurring and designed extremotolerance-associated proteins to assess their ability to protect human cells from chemically-induced apoptosis. We show that proteins from tardigrades, nematodes, and the Chinese giant salamander are apoptosis protective. Notably, we identify a region of the human ApoE protein with similarity to extremotolerance-associated proteins that also protects against apoptosis. This region mirrors the phase separation behavior seen with such proteins, like the tardigrade protein CAHS2. Moreover, we identify a synthetic protein, DHR81, that shares this combination of elevated phase separation propensity and apoptosis protection. Finally, we demonstrate that driving protective proteins into the condensate state increases apoptosis protection, and highlight the ability for DHR81 condensates to sequester caspase-7. Taken together, this work draws a link between extremotolerance-associated proteins, condensate formation, and human cellular protection.

INTRODUCTION

When exposed to extreme environmental stress, extremotolerance-associated (*ExTol*) organisms enter a state of reduced cellular metabolic activity that promotes survival in harsh conditions. This state has been shown to protect these organisms from high salinity, desiccation, and even the vacuum of outer space (Jonsson et al., 2008). To acclimate to

these conditions, *ExTol* organisms significantly upregulate the production of protective

metabolites, like trehalose (Neuman, 2006), as well as tolerance-conferring proteins

(Boothby et al., 2017), many of which are intrinsically disordered proteins (*IDPs*) with

repetitive sequence and predicted amphipathic helices.

While the mechanism of protection is unknown, one model suggests these proteins

mediate formation of a gel- or glass-like material state within an entire cell to protect it

from damage (Boothby et al., 2017; Hengherr et al., 2008). The model posits that such

proteins promiscuously interact with membranes and proteins, increasing intracellular

viscosity and protecting vulnerable structures (Hand et al., 2011) (Figure 1A Top). Such

interactions would also work to prevent stress-induced protein unfolding and protein

aggregation (Chakrabortee et al., 2010). An alternative model suggests that these

proteins form intracellular phase-separated membraneless compartments with distinct

physical parameters (referred to here as *condensates*) that partition key stress-sensitive

cellular components (Belott et al., 2020) (Figure 1A Bottom). Within these dense protein

structures, essential components are sequestered and protected from damage. The

models are not mutually exclusive; for example, in the case of desiccation tolerance,

continued dehydration could lead phase-separated compartments to fuse, comprise

much (if not all) of the cell, and promote a glassy material state.

The resurgence of interest in biological phase separation, extensively reviewed in

(Boeynaems et al., 2018; Choi et al., 2020; Hyman et al., 2014; Shin and Brangwynne,

2017), has shown that human cells can employ similar strategies to survive stressful

conditions. While the reversible formation of stress granules in healthy cells is the

archetype of stress tolerance, recent studies have revealed phase separation as a key

adaptation that cancer cells use to avoid apoptosis and cell death (Jiang et al., 2020).

Phase-separated condensates have been implicated in sequestering key tumor

repressors (Jiang et al., 2020), enhancing or repressing gene expression in response to

stress (Boija et al., 2018), and concentrating and protecting key metabolites like

nucleotides (Hubstenberger et al., 2017). These processes parallel the mechanisms that

ExTol proteins are thought to use to protect complex biological systems, suggesting there

could be some functional overlap. That is, sequence features and properties within ExTol

proteins could be implicated in mammalian stress tolerance pathways, including

apoptosis resistance.

A key sequence feature of some ExTol proteins is the presence of amphipathic helices

(Tolleter et al., 2007). While disordered in most contexts, ExTol proteins can transiently

form amphipathic helices in certain molecular environments, notably when interacting with

and stabilizing membranes (Hand et al., 2011). The helices often exhibit a hydrophobic

face, which interacts with core lipids within the membrane, flanked by positively charged

residues that might interact with the negatively charged lipid head groups (Figure 1B).

This amphipathic moment often repeats across the length of the protein (Figure 1C). This

feature is a hallmark of the late embryonic abundant (LEA) class of ExTol proteins (Hand

et al., 2011). Human membrane interacting proteins, like alpha-synuclein, also exhibit a

similar repeat motif (Bussell and Eliezer, 2003). Such parallels in sequence features

suggest that orthogonal expression of ExTol proteins may confer their protective

properties to other biological systems, like human cells.

Indeed, previous reports have shown that some of these ExTol proteins provide similar

protective properties when expressed heterologously. Expressing tardigrade-specific

IDPs (TDPs) in bacteria provides protection against desiccation (Boothby et al., 2017).

Anhydrin, an IDP upregulated by *Aphelenchus avenae* during dehydration, expressed in

a human cell line limited PABPN1-A17 aggregation, a hallmark of oculopharyngeal

muscular dystrophy (Chakrabortee et al., 2010). Similar reports demonstrated protection

of human cells provided by expression of LEA proteins originating from the African midge

(Li et al., 2012) and by TDPs from tardigrades (Tanaka et al., 2015). These findings

reinforce the promise of transferring protective phenotypes from their native contexts to

human cells.

Taken together, these data indicate a potential avenue for using this class of proteins to

protect biological systems in a controlled manner. To test this hypothesis, we generated

approximately 300 sequences inspired by ExTol proteins and examined their ability to

protect against apoptotic cell death. Our results show that human proteins containing

sequence hallmarks of ExTol proteins as well as *de-novo* designed condensate forming

proteins can protect human cells from chemically-induced apoptosis. We also

demonstrate that condensate induction using synthetic modifications can strengthen

apoptotic protection. Finally, we show that key components of the apoptotic pathway can

localize to inducible condensates, suggesting enzyme sequestration could be a

mechanism of action.

RESULTS

Assembling a library of stress tolerance proteins

Based on the hallmark sequence features of ExTol proteins, we searched the published

literature to identify proteins that not only contain these features but also exhibit some

evidence for protection of biological structures. The library was generated based on over

one hundred naturally-occurring proteins based on five lines of evidence for stress

tolerance association, including: change in expression during extremotolerance,

knockdown or overexpression changing the tolerant state, *in vitro* protection, orthogonal

expression transferring tolerance, and sequence similarity to other proteins with these

features (Figure 1D and Supplemental table 1).

A PFAM domain analysis of the identified proteins reveals almost half of the identified

targets have two key domains: Late Embryonic Abundant (LEA) and Apolipoprotein E

(ApoE) domains (Figure 1E, Supplemental table 1). LEA proteins are well known to be

upregulated during the last stages of plant seed maturation (Mertens et al., 2018). This

domain is thought to protect the seed embryo during transport to its new destination.

There is also evidence that this domain has been co-opted by bacteria as protection from

desiccation and radiation (Makarova et al., 2001). ApoE proteins interact with lipid

particles to allow for lipid reuptake into cells. Additionally, mutations within human ApoE

have been identified as strong genetic predictors of Alzheimer's disease in human

populations (Yamazaki et al., 2019). Based on their lipid interaction properties and

potential role in neuronal homeostasis, as well as its prevalence in the PFAM domain

analysis, we decided to include human ApoE and its various mutations in our study as an

ExTol protein.

Designing novel extremotolerance-associated proteins

To test hypotheses about sequence features and their relationship to phase separation

and apoptosis, we designed novel proteins based on the set of ExTol proteins. These

new designs take 3 approaches (Figure 2): truncation of proteins to focus on repetitive,

intrinsically disordered, and amphipathic regions; isolation of individual helices with

desired sequence features; and *de novo* design of alpha helix-containing proteins that

self-assemble into higher order structures. Through these designs, we sought to identify

the minimal elements required for protection and phase separation.

First, we identified regions within natural proteins that contain intrinsic disorder (Erdos

and Dosztanyi, 2020), repetitive sequences (Biegert and Soding, 2008), and amphipathic

helices (Gimenez-Andres et al., 2018; Segrest et al., 1992). Figure 2A illustrates the use

of HHRepID on the sequence of full length human ApoE. Based on this analysis, we

identified the repeating domain within ApoE for further testing. Other ExTol proteins were

similarly modified and examined.

We also included short peptides based on the consensus sequence of repetitive regions

that have previously been shown to be protective (Furuki et al., 2019). To test if any of

the sequence features we identified are important for apoptosis protection, we altered

these features within these short peptides. For example, we altered the amphipathic

character of these helices (Figure 2B) to determine if apoptosis protection was changed.

To examine whether generic condensate formation is sufficient to confer apoptosis

protection, we also examined completely synthetic alpha helix-containing proteins that

were designed to form repetitive structure with sequences orthogonal to naturally-

occurring proteins (Brunette et al., 2015). These designed helical repeat (DHR) proteins

interact along the axis of the helix, forming long spiraling fibrils, but may also potentially

self-interact along the face of the protein based on a pattern of positive and negative

charges (Figure 2C). This ability to oligomerize and form multivalent complexes, which

may then interact via weak non-covalent interactions, is an established determinant of

condensate formation (Choi et al., 2020).

Together, our library consists of 104 naturally-occurring proteins, 166 truncations or

isolations of helical regions, and 19 *de novo* designs (Figure 2D). Taken altogether, these

proteins interrogate several hypotheses about how disordered proteins may protect

human cells against apoptosis.

Apoptosis assay identifies extremotolerance-associated proteins in human cells

To model stress in a mammalian cell system, we induced cellular apoptosis by treatment

with the chemotherapeutic camptothecin (CPT), a DNA topoisomerase inhibitor well

known to trigger apoptosis (Li et al., 2017; Wall et al., 1966). Under these conditions,

protection against apoptosis could be achieved through a number of avenues, including

protection against the DNA damage caused by CPT-induced topoisomerase inhibition (Li

et al., 2017), interruption of the signaling cascade, or even protection of the mitochondrial

membrane whose rupture ultimately triggers apoptosis (Bortner and Cidlowski, 2002).

To test our designs for protection against apoptosis, we used a human cell-based

apoptosis assay (Figure 3A). This assay uses HT-1080 fibrosarcoma cells from ATCC

(Rasheed et al., 1974). These cells are seeded in a 96-well plate and allowed to adhere

overnight. The next day, the cells are co-transfected with a plasmid containing the ExTol

candidate proteins of interest as well as a transfection marker. The following day, the

media is swapped to one containing 13.2 μM CPT. After addition of CPT, caspase-3 and

-7 activation is measured by microscopy every 45 minutes for 1 day using a fluorophore

quencher pair attached to either end of a peptide with the sequence "DEVD". As caspase-

3 and -7 are activated, the DEVD peptide is cleaved, releasing the fluorophore from the

quencher. The unquenched fluorophore localizes to the nucleus where it can be

quantified by imaging on a single-cell level. The single-cell resolution allows for the

separation of transfected and non-transfected cells based on the transfection marker.

Only single cells with the transfection marker were subjected to further analysis; Figure

3B shows sample images highlighting the type of data being analyzed.

On every plate, an empty vector negative control is run as well as a positive control

containing the X-linked inhibitor of apoptosis (XIAP). This protein is known to bind to

caspase-3 and -7 to block their activity (Obexer and Ausserlechner, 2014) . Figure 3C

shows example data comparing median DEVD peptide signal intensity over time after

induction with CPT of four wells on the same plate transfected with XIAP and empty

vector. The results show a reduction in DEVD peptide signal intensity when the XIAP-

containing vector is transfected, demonstrating that the assay can identify modulators of

apoptosis.

To convert these time-resolved data to a measure of protection against apoptosis, the

maximum DEVD peptide signal intensity is calculated and normalized to a plate-matched

empty vector control (see methods for further details). The normalized apoptosis induction

intensity is shown in Figure 3D left. The data are slightly skewed towards apoptosis-

protection, which supports our hypothesis that the targeted proteins can protect against

apoptosis.

To highlight the top-performing designs, the top 10% of the targets are shown in Figure

3D right. As expected, the XIAP positive control showed the strongest anti-apoptotic

effect. The two top-performing hits were ApoE designs. Human alpha-synuclein is the 5th

hit, which is expected as it is known to be both intrinsically disordered, and anti-apoptotic

(Li and Lee, 2005). Additionally, several ApoE designs and short LEA helices are top hits.

The data used to generate these figures is included in supplemental table 2.

Known extremotolerance-associated proteins affect apoptosis

Among the top hits in apoptosis reduction are an IGKV protein highly expressed in mucus

of the Chinese giant salamander *Andrias davidianus* (Geng et al., 2019) that is used as

a medical glue in traditional Chinese medicine, an LEA protein from the nematode

Aphelenchus avenae previously shown to have the ability to reduce a form of protein

aggregation associated with oculopharyngeal muscular dystrophy in human cells

(Chakrabortee et al., 2010), and a mitochondrial LEA protein from the brine shrimp

Artemia franciscana shown to protect human cells during desiccation (Li et al., 2012)

(Figure 3D). The prominence of these ExTol proteins suggests the assay can help identify

other proteins that can protect cells from more general stresses beyond CPT-induced

apoptosis.

Short repetitive peptides and *de novo* designs modulate apoptosis

LEA repeat-containing peptides are among the top-performing hits in the screen. While

the original short LEA designs from Furuki et al. 2019 (Furuki et al., 2019) had a minor

effect reducing apoptosis, several of the designs tested here performed better

(Supplemental Figure 3A). As 22 amino acids is small and could lead to peptide stability

issues, longer 50-amino acid designs inspired by the same LEA proteins Furuki et al.

utilized were also developed (Supplemental Figure 3B). Some of these designs were also

able to reduce apoptosis, including the PvLEA4 with extended repeats (notably PvLEA4

repeats extended #2), which fell into the top hits of the entire library (Figure 3D and

Supplemental Figure 3B). These results suggest that optimization of synthetic designs

yields proteins with stronger function than naturally-occurring proteins.

While these proteins were designed to interrogate the relationship between sequence

and apoptosis protection in short peptides, little relationship was found with respect to the

sequence features we analyzed (Supplemental Figure 3C). Several designs were made

that would alter the hydrophobic moments of putative helices these peptides are predicted

to form. To examine if this sequence property relates to the stress-tolerance function of

these proteins, we attempted to correlate this property with assay performance. However,

no significant correlation was found, suggesting (at least in these cases) that degree of

amphipathicity on its own is not the key determinant of stress-tolerance function of these

proteins.

The repeat region of ApoE comprises its minimal anti-apoptotic unit

The top two hits in the assay were ApoE4 (47-300) and ApoE2 (47-300). These truncated

forms of ApoE maintain the repeat domains shown in Figure 4A, while removing the signal

peptide and terminal regions. This suggests that the repeat region of ApoE accounts for

its core anti-apoptotic functionality. Interestingly, the full length ApoE2 and E4 do not

show up as top hits, suggesting that these truncated forms improve on the apoptosis

protection provided by the full-length variants (Figure 4B).

Based on the location of the signal peptide, it is expected that ApoE4 should localize to

the secretory pathway whereas the truncated form may be cytoplasmic. To test this

hypothesis, cells were subjected to Western blot and immunofluorescence analysis using

commercially available antibodies (Abcam ab24139; Figure 4C and Supplemental Figure

4A). The Western blot results show that both the full-length and truncated forms of ApoE4

are expressed at the expected size (Supplemental Figure 4A). The immunofluorescence

shows co-localization between the ApoE4 full length protein and the endoplasmic

reticulum (marked by Thermo's PDI antibody MA3-019, Figure 4C). However, the

truncated form did not show the same co-localization, suggesting as expected that it may

not enter the secretory pathway as the full-length version does.

Strikingly, the truncated version appeared to form more pronounced puncta throughout

the cytoplasm of the cell, suggesting the formation of potentially phase-separated

condensates. In screening through the point mutants of ApoE also tested here, we found

a similar condensate phenotype with the full length ApoE3 D125I mutation (Supplemental

Figure 4B). Together, these data suggest that ApoE may be able to support condensate

formation and confer apoptosis protection in part via the protective compartment

mechanism (Figure 1A bottom).

CAHS2 and ApoE form condensates upon induced multivalency

Established condensate engineering tools (Bracha et al., 2019) were used to test for

latent phase separation propensity within the top performing proteins. We used both light-

and chemically-inducible systems of multivalency. In the light system, a ferritin core

(comprised of self-assembled monomers, FTH1 domain-eGFP-iLID) becomes decorated

with SspB-modified proteins upon exposure to blue light (Bracha et al., 2018) (Figure 4D

top). In the chemical system, multivalency (i.e. multiple arms of the ExTol protein of

interest) is induced using a dimerization agent (Stanton et al., 2018) that brings together

FKBP-modified proteins (Bracha et al., 2019) (Figure 4D bottom). These inducible

methods increase the valency of a system, and thus work to favor phase separation. In

this way, they can help reveal any latent ability to form condensates that is not

immediately observable without modification.

Indeed, condensates were observed in both inducible systems using ExTol proteins.

CAHS2, a protein from the tardigrade *P. richtersi*, exhibited the most dramatic phenotype,

which is consistent with recent work describing the gel forming capabilities of CAHS

proteins (Yagi-Utsumi et al., 2021) (Figure 4E&F). ApoE variants were of particular

interest, based on the puncta observed in the top performing ApoE constructs. All

versions of ApoE tested formed puncta without activation, consistent with our previous

observation using untagged native ApoE (Figure 4C). Upon light- or chemical-activation,

there is a noticeable increase in condensates, exemplifying how phase separation can be

enhanced with increased multivalency (Figure 4E&F).

Induction of condensates reduces apoptosis

Induction of multivalency in ApoE containing constructs reduces apoptosis. We performed

the apoptosis assay using the top hits modified for chemically-induced multivalency,

namely fusion to two FKBP domains. The results show that the presence of the chemical

dimerizer did not affect apoptosis signal in cells transfected with the empty vector; in

contrast, the ApoE constructs exhibit reduced apoptosis with added dimerizer (Figure

4G). That is, artificially increasing ApoE condensates was apoptosis protective. These

data suggest that condensate formation plays a role in the anti-apoptotic effect of these

proteins.

Synthetic DHR81 condensates reduce apoptosis and sequester caspase-7

Induction of DHR81 based condensates protects against apoptosis. DHR81 is formed

from four helical hairpin repeat regions; the core of the protein is hydrophobic to ensure

consistent folding, while the outside of the protein is decorated with checkerboard positive

and negative charges (Figure 5A) (Brunette et al., 2015). This structure ensures the

protein is well folded, while also enabling the potential for higher order structure formation

via electrostatic interactions. Indeed, as with the ApoE variants, DHR81 supported

condensates in the chemically-inducible condensate system (Figure 5B) and exhibited a

stronger anti-apoptotic phenotype when multivalency is induced (Figure 5C and

Supplemental Figure 5A). There was little to no effect in cells containing an empty vector

or expressing the FKBP dimerization domains alone.

To further explore how condensates mediated by ExTol proteins may confer an anti-

apoptotic effect, we examined whether they sequester major components of apoptotic

signaling, namely caspase-3 and -7. The results showed that activated caspase-7

localized to DHR81 condensates but not CAHS2 or ApoE4 condensates (Figure 5D).

Caspase-3, however, does not localize to either DHR81, CAHS2, or ApoE4 condensates

(Supplemental Figure 5B). These results suggest sequestration of key apoptotic factors

could be a mechanism by which these condensates reduce apoptosis.

Discussion

To broadly evaluate the ability of ExTol proteins to protect orthogonal systems from

environmental stress, we studied over one hundred different ExTol proteins. We used

these as a basis to design approximately 300 variants, including truncations and fusion

proteins. These designs were evaluated on their ability to protect human HT-1080 cells

against CPT-induced apoptosis. The results showed that known ExTol proteins, ApoE

variants, and the synthetic DHR81 protein could modulate apoptosis levels when

transferred to human cells.

Protection against apoptosis is a key feature of many ExTol proteins. As apoptosis is a

common cellular response to stress, the ability to mitigate this response is important in

providing cell-autonomous protection in stressful environments. While the mechanism

resilient organisms use to survive is still not fully understood, it is becoming increasingly

clear that phase separation may play an important role.

ExTol proteins are proposed to form expansive higher order structures (e.g. gels or glassy

systems) that reduce the viscosity of the cytoplasm, especially in desiccated states

(Boothby et al., 2017). Other reports show that these proteins could mediate formation of

protective phase-separated compartments (i.e. condensates), sequestering damage-

inducing proteins and/or protecting important targets (Belott et al., 2020). Based on our

condensate data, it appears the latter may be at play for proteins like ApoE or DHR81.

Notably, in these proteins that already exhibit both anti-apoptotic effects as well as

elevated phase separation propensities, increasing the amount of condensates via an

engineered inducible system improved protection against apoptosis. This observation

suggests that condensates could be a key part of the protective mechanism.

Interestingly, we observed that activated caspase-7 partitions into DHR81 condensates,

suggesting caspase sequestration as a potential anti-apoptotic mechanism. Phase-

separated condensates and higher order assemblies are already implicated in mediating

apoptosis: stress granules inhibit apoptosis by suppressing MAPK pathways (Arimoto et

al., 2008); paraspeckles sequester pro-apoptotic tumor suppressors (Jiang et al., 2020);

the apoptosome relies on Apaf-1 multimerization, which has been reported to undergo

amyloid fibrillation in specific conditions (Rao et al., 2009); and stress-dependent phase

separation modulates proteasome formation, whose function affects apoptosis (Gupta et

al., 2018; Yasuda et al., 2020). It is interesting to note that we only observe clear caspase-

7 partitioning into DHR81 condensates, suggesting the possibility of a specific binding

interaction in this case. ApoE condensates may be sequestering other apoptotic factors

or using a different mechanism to reduce apoptosis.

It is also possible that these ExTol proteins achieve their anti-apoptotic effect by

implementing a more general physical mechanism, namely slowing the intracellular

mobility of biological effectors. There are various reports describing how the effective

viscosity within the cytoplasm can be increased in response to external stresses

(Dijksterhuis et al., 2007; Parry et al., 2014) or mediated by protein composition,

specifically intrinsically disordered proteins (Theillet et al., 2014). Notably, a pervasive

cell-spanning gel or network is not necessary to achieve such an effect; it is possible that

the presence of condensates, coupled with their ability to selectively partition

biomolecules (i.e. control what biomolecules can enter and diffuse through them), could

broadly slow biomolecular movement. That is, by forcing different proteins to either be

trapped in them or diffuse around them, condensates can decrease protein mobility and

perturb e.g. signaling pathways. Our results showing a lack of caspase-7 sequestration

by some of our protein condensates could thus support this more general mechanism of

condensates perturbing biology.

Condensates are becoming increasingly implicated in disease (Shin and Brangwynne,

2017), particularly when their structural metastability and reversibility is dysregulated (e.g.

through mutation). Notably, drugs that block the formation of amyloid fibrils and plaques,

which are products of phase separation dysregulation, are promising yet strongly debated

(Lalli et al., 2021) therapies for many neurological diseases. As such, our observations of

ApoE are particularly salient. We found that expression of a cytoplasmic form of ApoE is

protective against apoptosis. In particular, the truncated repeat region of ApoE is sufficient

and shows the strongest anti-apoptotic activity of all designs we tested. Others have

shown similar effects for exogenously added ApoE-containing liposomes protecting rat

neurons against apoptosis (Hayashi et al., 2009), though we show the effect through

overexpression of ApoE inside of human cells. When combined with our condensate

results and the clinical relevance of ApoE in Alzheimer's disease (Schellenberg et al.,

2000), these observations suggest the anti-apoptotic effect observed here warrants

further study, to further elucidate its role in Alzheimer's disease progression.

Altogether, our work expands upon the understanding of the relationship between

condensate formation and cell physiology, demonstrating how condensate accumulation

can be broadly beneficial in reducing apoptosis while also identifying diverse new protein

domains that confer protection.

Author Contributions

M.T.V. and D.T.N. designed and performed experiments, analyzed the data, and wrote

the paper. N.N.T. analyzed the data and wrote the paper. M.E.O provided technical

support. N.J.R., K.B.B., and N.P.B., designed constructs. D.B. and J.C.W. managed the

teams and provided key insight. D.S.M., R.L.C., and P.A.S., designed constructs,

managed teams, and wrote the paper.

Acknowledgements

Research was sponsored by the NSF, DARPA, the Wyss Institute for Biologically inspired

engineering, and the Army Research Office and was accomplished under Cooperative

Agreement Number W911NF-19-2-0017. M.T.V. was sponsored by the National Science

Foundation under Grant No. 2010370. D.T.N., N.N.T., M.E.O., N.J.R., K.P.B., J.C.W,

D.S.M., and P.A.S. were supported by the Harvard Medical School Department of

Systems Biology. J.C.W. was further supported by the Harvard Medical School

Laboratory of Systems Pharmacology. K.P.B. and D.S.M. were further supported by CZI

Neurodegenerative Challenge Ben Barres Early Career Award (CZI2018-191853). N.P.B

was further supported by the HHMI Hanna Gray fellowship (grant GT11817). We would

like to acknowledge Clarence Yapp for help with microscopy as well as the NCI U54-

CA225088 grant and a gift from the Ludwig Institute for Cancer Research to the

Laboratory for Systems Pharmacology. We would like to acknowledge the Nikon Imaging

Center at Harvard Medical Center for providing microscope support. The views and

conclusions contained in this document are those of the authors and should not be

interpreted as representing the official policies, either expressed or implied, of any funding

agency including the NSF, DARPA, the Army Research Office, or the U.S. Government.

The U.S. Government is authorized to reproduce and distribute reprints for Government

purposes notwithstanding any copyright notation herein.

440
441   **Bibliography & References**
442
Arimoto, K., Fukuda, H., Imajoh-Ohmi, S., Saito, H., and Takekawa, M. (2008). Formation of
stress granules inhibits apoptosis by suppressing stress-responsive MAPK pathways. Nat Cell
Biol *10*, 1324-1332.
Belott, C., Janis, B., and Menze, M.A. (2020). Liquid-liquid phase separation promotes animal
desiccation tolerance. Proc Natl Acad Sci U S A *117*, 27676-27684.
Biegert, A., and Soding, J. (2008). De novo identification of highly diverged protein repeats by
probabilistic consistency. Bioinformatics *24*, 807-814.
Boeynaems, S., Alberti, S., Fawzi, N.L., Mittag, T., Polymenidou, M., Rousseau, F., Schymkowitz,
J., Shorter, J., Wolozin, B., Van Den Bosch, L.*, et al.* (2018). Protein Phase Separation: A New
Phase in Cell Biology. Trends Cell Biol *28*, 420-435.
Boija, A., Klein, I.A., Sabari, B.R., Dall'Agnese, A., Coffey, E.L., Zamudio, A.V., Li, C.H., Shrinivas,
K., Manteiga, J.C., Hannett, N.M.*, et al.* (2018). Transcription Factors Activate Genes through
the Phase-Separation Capacity of Their Activation Domains. Cell *175*, 1842-1855 e1816.
Boothby, T.C., Tapia, H., Brozena, A.H., Piszkiewicz, S., Smith, A.E., Giovannini, I., Rebecchi, L.,
Pielak, G.J., Koshland, D., and Goldstein, B. (2017). Tardigrades Use Intrinsically Disordered
Proteins to Survive Desiccation. Mol Cell *65*, 975-984 e975.
Bortner, C.D., and Cidlowski, J.A. (2002). Cellular mechanisms for the repression of apoptosis.
Annu Rev Pharmacol Toxicol *42*, 259-281.
Bracha, D., Walls, M.T., and Brangwynne, C.P. (2019). Probing and engineering liquid-phase
organelles. Nat Biotechnol *37*, 1435-1445.
Bracha, D., Walls, M.T., Wei, M.T., Zhu, L., Kurian, M., Avalos, J.L., Toettcher, J.E., and
Brangwynne, C.P. (2018). Mapping Local and Global Liquid Phase Behavior in Living Cells Using
Photo-Oligomerizable Seeds. Cell *175*, 1467-1480 e1413.
Brunette, T.J., Parmeggiani, F., Huang, P.S., Bhabha, G., Ekiert, D.C., Tsutakawa, S.E., Hura, G.L.,
Tainer, J.A., and Baker, D. (2015). Exploring the repeat protein universe through computational
protein design. Nature *528*, 580-584.
Bussell, R., Jr., and Eliezer, D. (2003). A structural and functional role for 11-mer repeats in
alpha-synuclein and other exchangeable lipid binding proteins. J Mol Biol *329*, 763-778.

Chakrabortee, S., Meersman, F., Kaminski Schierle, G.S., Bertoncini, C.W., McGee, B., Kaminski,
C.F., and Tunnacliffe, A. (2010). Catalytic and chaperone-like functions in an intrinsically
disordered protein associated with desiccation tolerance. Proc Natl Acad Sci U S A *107*, 16084-
16089.
Choi, J.M., Holehouse, A.S., and Pappu, R.V. (2020). Physical Principles Underlying the Complex
Biology of Intracellular Phase Transitions. Annu Rev Biophys *49*, 107-133.
Dijksterhuis, J., Nijsse, J., Hoekstra, F.A., and Golovina, E.A. (2007). High viscosity and anisotropy
characterize the cytoplasm of fungal dormant stress-resistant spores. Eukaryot Cell *6*, 157-170.
Erdos, G., and Dosztanyi, Z. (2020). Analyzing Protein Disorder with IUPred2A. Curr Protoc
Bioinformatics *70*, e99.
Furuki, T., Niwa, T., Taguchi, H., Hatanaka, R., Kikawada, T., and Sakurai, M. (2019). A LEA model
peptide protects the function of a red fluorescent protein in the dry state. Biochem Biophys Rep
17, 27-31.
Geng, X., Guo, J., Zang, X., Chang, C., Shang, H., Wei, H., and Xu, C. (2019). Proteomic analysis of
eleven tissues in the Chinese giant salamander (Andrias davidianus). Sci Rep *9*, 16415.
Gimenez-Andres, M., Copic, A., and Antonny, B. (2018). The Many Faces of Amphipathic
Helices. Biomolecules *8*.
Gupta, I., Singh, K., Varshney, N.K., and Khan, S. (2018). Delineating Crosstalk Mechanisms of
the Ubiquitin Proteasome System That Regulate Apoptosis. Front Cell Dev Biol *6*, 11.
Hand, S.C., Menze, M.A., Toner, M., Boswell, L., and Moore, D. (2011). LEA proteins during
water stress: not just for plants anymore. Annu Rev Physiol *73*, 115-134.
Hayashi, H., Campenot, R.B., Vance, D.E., and Vance, J.E. (2009). Protection of neurons from
apoptosis by apolipoprotein E-containing lipoproteins does not require lipoprotein uptake and
involves activation of phospholipase Cgamma1 and inhibition of calcineurin. J Biol Chem *284*,
29605-29613.
Hengherr, S., Brümmer, F., and Schill, R.O. (2008). Anhydrobiosis in tardigrades and its effects
on longevity traits. Journal of Zoology *275*, 216-220.
Hubstenberger, A., Courel, M., Benard, M., Souquere, S., Ernoult-Lange, M., Chouaib, R., Yi, Z.,
Morlot, J.B., Munier, A., Fradet, M.*, et al.* (2017). P-Body Purification Reveals the Condensation
of Repressed mRNA Regulons. Mol Cell *68*, 144-157 e145.
Hyman, A.A., Weber, C.A., and Julicher, F. (2014). Liquid-liquid phase separation in biology.
Annu Rev Cell Dev Biol *30*, 39-58.
Jiang, S., Fagman, J.B., Chen, C., Alberti, S., and Liu, B. (2020). Protein phase separation and its
role in tumorigenesis. Elife *9*.
Jonsson, K.I., Rabbow, E., Schill, R.O., Harms-Ringdahl, M., and Rettberg, P. (2008). Tardigrades
survive exposure to space in low Earth orbit. Curr Biol *18*, R729-R731.
Lalli, G., Schott, J.M., Hardy, J., and De Strooper, B. (2021). Aducanumab: a new phase in
therapeutic development for Alzheimer's disease? EMBO Mol Med *13*, e14781.
Li, F., Jiang, T., Li, Q., and Ling, X. (2017). Camptothecin (CPT) and its derivatives are known to
target topoisomerase I (Top1) as their mechanism of action: did we miss something in CPT
analogue molecular targets for treating human disease such as cancer? Am J Cancer Res *7*,
2350-2394.

Li, S., Chakraborty, N., Borcar, A., Menze, M.A., Toner, M., and Hand, S.C. (2012). Late
embryogenesis abundant proteins protect human hepatoma cells during acute desiccation.
Proc Natl Acad Sci U S A *109*, 20859-20864.
Li, W., and Lee, M.K. (2005). Antiapoptotic property of human alpha-synuclein in neuronal cell
lines is associated with the inhibition of caspase-3 but not caspase-9 activity. J Neurochem *93*,
1542-1550.
Makarova, K.S., Aravind, L., Wolf, Y.I., Tatusov, R.L., Minton, K.W., Koonin, E.V., and Daly, M.J.
(2001). Genome of the extremely radiation-resistant bacterium Deinococcus radiodurans
viewed from the perspective of comparative genomics. Microbiol Mol Biol Rev *65*, 44-79.
Mertens, J., Aliyu, H., and Cowan, D.A. (2018). LEA Proteins and the Evolution of the WHy
Domain. Appl Environ Microbiol *84*.
Neuman, Y. (2006). Cryptobiosis: a new theoretical perspective. Prog Biophys Mol Biol *92*, 258-
267.
Obexer, P., and Ausserlechner, M.J. (2014). X-linked inhibitor of apoptosis protein - a critical
death resistance regulator and therapeutic target for personalized cancer therapy. Front Oncol
4, 197.
Parry, B.R., Surovtsev, I.V., Cabeen, M.T., O'Hern, C.S., Dufresne, E.R., and Jacobs-Wagner, C.
(2014). The bacterial cytoplasm has glass-like properties and is fluidized by metabolic activity.
Cell *156*, 183-194.
Rao, P.N., Reddy, K.S., and Bhuyan, A.K. (2009). Amyloid fibrillation of human Apaf-1 CARD.
Biochemistry *48*, 7656-7664.
Rasheed, S., Nelson-Rees, W.A., Toth, E.M., Arnstein, P., and Gardner, M.B. (1974).
Characterization of a newly derived human sarcoma cell line (HT-1080). Cancer *33*, 1027-1033.
Schellenberg, G.D., D'Souza, I., and Poorkaj, P. (2000). The genetics of Alzheimer's disease. Curr
Psychiatry Rep *2*, 158-164.
Segrest, J.P., Jones, M.K., De Loof, H., Brouillette, C.G., Venkatachalapathi, Y.V., and
Anantharamaiah, G.M. (1992). The amphipathic helix in the exchangeable apolipoproteins: a
review of secondary structure and function. J Lipid Res *33*, 141-166.
Shin, Y., and Brangwynne, C.P. (2017). Liquid phase condensation in cell physiology and disease.
Science *357*.
Stanton, B.Z., Chory, E.J., and Crabtree, G.R. (2018). Chemically induced proximity in biology
and medicine. Science *359*.
Tanaka, S., Tanaka, J., Miwa, Y., Horikawa, D.D., Katayama, T., Arakawa, K., Toyoda, A., Kubo, T.,
and Kunieda, T. (2015). Novel mitochondria-targeted heat-soluble proteins identified in the
anhydrobiotic Tardigrade improve osmotic tolerance of human cells. PLoS One *10*, e0118272.
Theillet, F.X., Binolfi, A., Frembgen-Kesner, T., Hingorani, K., Sarkar, M., Kyne, C., Li, C., Crowley,
P.B., Gierasch, L., Pielak, G.J.*, et al.* (2014). Physicochemical properties of cells and their effects
on intrinsically disordered proteins (IDPs). Chem Rev *114*, 6661-6714.
Tolleter, D., Jaquinod, M., Mangavel, C., Passirani, C., Saulnier, P., Manon, S., Teyssier, E., Payet,
N., Avelange-Macherel, M.H., and Macherel, D. (2007). Structure and function of a
mitochondrial late embryogenesis abundant protein are revealed by desiccation. Plant Cell *19*,
1580-1589.
Wall, M.E., Wani, M.C., Cook, C.E., Palmer, K.H., McPhail, A.T., and Sim, G.A. (1966). Plant
Antitumor Agents. I. The Isolation and Structure of Camptothecin, a Novel Alkaloidal Leukemia

and Tumor Inhibitor from Camptotheca acuminata1,2. Journal of the American Chemical

Society *88*, 3888-3890.

Yagi-Utsumi, M., Aoki, K., Watanabe, H., Song, C., Nishimura, S., Satoh, T., Yanaka, S., Ganser,

C., Tanaka, S., Schnapka, V.*, et al.* (2021). Desiccation-induced fibrous condensation of CAHS

protein from an anhydrobiotic tardigrade. bioRxiv, 2021.2006.2022.449423.

Yamazaki, Y., Zhao, N., Caulfield, T.R., Liu, C.C., and Bu, G. (2019). Apolipoprotein E and

Alzheimer disease: pathobiology and targeting strategies. Nat Rev Neurol *15*, 501-518.

Yasuda, S., Tsuchiya, H., Kaiho, A., Guo, Q., Ikeuchi, K., Endo, A., Arai, N., Ohtake, F., Murata, S.,

Inada, T.*, et al.* (2020). Stress- and ubiquitylation-dependent phase separation of the

proteasome. Nature *578*, 296-300.

Figure 1: Selection of candidate extremotolerance-associated proteins (a) Illustration of different protective models for cellular stress tolerance. **(b)** Illustration of amphipathic membrane interaction for rotifer LEA4 protein and Alpha-synuclein, thus highlighting a feature indicative of extremotolerance-associated proteins. **(c)** Hydrophobic moment analysis of rotifer LEA4, showing the repetitive nature of the amphipathic signature. Highlighted green segment corresponds to the LEA4 helix in (b). **(d)** Venn diagram of the lines of evidence used to select targets in the final library. Abbreviations: KD, knockdown; OE, overexpression; Δ, change in expression levels. **(e)** Domain analysis of included targets reveals LEA and ApoE domains in almost half of the targets tested. The other half contains proteins with other PFAM domains (19%) or without any affiliated PFAM entry (32%). Of the proteins lacking a known PFAM annotation, many exhibit repetitive sequence (20% of all targets).

a ——— Isolate repetitive units ———

☑ HHRepID repeat

Remove non-repetitive elements

b ——— Modulate amphipathicity ———

LEA 22mer
Mutant #10

LEA 22mer

Minimize
Moment

Maximize
Moment

LEA 22mer
Mutant #5

Positive
Negative
Non-polar
Other

c ——— De novo protein design ———

Alternating
Charges

**Oligomerizing
Interactions**

Electrostatic

Hydrophobic

Multivalency favors
phase separation

Non-polar
Face

d ——— Library composition ———

Natural
(104)

Inspired by
nature
(166)

de novo (19)

15 **Figure 2: Design based on candidate extremotolerance-associated proteins (a)**
16 Truncations were made to isolate regions of repetitive and amphipathic sequence. **(b)**
17 Small helices were selected and altered based on their hydrophobic moment. **(c)**
18 Synthetic alpha helix-containing proteins designed to oligomerize. **(d)** Venn diagram
19 highlighting the library that was tested.

a

Seed Cells — Transfect Cells — Add CPT — Fluorescent microscopy — Image Analysis

Day 0 — Day 1 — Day 2 — ----- — Day 3

b

Cell (SiR-Hoechst) | Transfection (mCherry) | Caspase (DEVD peptide)

50 µm

c

$Log_2\left(\dfrac{Target}{Empty\ Vector}\right)$

d

ApoE derived
Synthetic

XIAP
ApoE4 (47-300)
ApoE2 (47-300)
ANDDA IGKV
Human α-synuclein
ApoE3 (E114I)
APHAV LEA1
PARRC CAHS1 (101-189)
ARTSF LEA3m
Dehydrin VrDHN1a
ApoE3 (80-255)
DHR81
PvLEA4 extended #2
ApoE3 (K164E)
HYPDU CAHS7
CRAPL C4-LEA
ApoE3 (R160W)
HYPDU CAHS2
RAMVA CAHS1 (del 108-197)
LEA 22mer Shuffle #3
ApoE3 (D125I)
PARRC CAHS2 (98-130)
Dehydrin TsDHN-1
PvLEA4 extended #3
PvLEA4 extended #1
RAMVA CAHS2 (94-170)
LEA 22mer Mutant #10
DEIRA DrHD b3 domain 3
DEIRA DR_1172 (48-197)
LEA 22mer Mutant #16
Empty Vector

Protein library

Change in apoptosis (relative to empty vector)

Figure 3: Screen for CPT resistance by extremotolerance-associated proteins (a) Flow diagram of the assay. **(b)** Representative images showing cells during the assay. Cells are identified using SiR-Hoechst marking the nucleus (*blue*). Transfected cells are identified by the presence of the mCherry marker (*green*). Finally, the degree of apoptosis is identified by the presence of the cleaved DEVD peptide (*red*). **(c)** Positive control showing apoptosis reduction upon expression of XIAP. **(d)** Bar graph highlighting the

apoptosis resistance provided by different constructs, including zoomed in area to
highlight the top hits in the assay. Source species of a sequence are abbreviated and
shown in italics; abbreviations are drawn from Uniprot, including *RAMVA* (*Ramazzottius*
varieornatus), *HYPDU* (*Hypsibius dujardini*), *PARRC* (Paramacrobiotus richtersi),
ANDDA (*Andrias davidianus*), *APHAV* (*Aphelenchus avenae*), *ARTSF* (*Artemia*
franciscana), *CRAPL* (*Craterostigma plantagineum*), and *DEIRA* (*Deinococcus*
radiodurans).
38

Fig. 4. ApoE truncations and synthetic condensates further reduce apoptosis.
(a) Gene diagram highlighting the repetitive region of ApoE. **(b)** Boxplots highlighting the apoptosis assay performance of the full length and truncated forms of ApoE2 and E4. **(c)** Immunofluorescence of ApoE4, full length and truncated forms. *White* indicates colocalization, which is more apparent with the full-length protein. The truncated form appears to exhibit larger and more distinguishable puncta. **(d)** Diagram of the induced multi-valency systems used here, including the light-inducible system (*top*) and the chemically-inducible system (*bottom*). **(e)** Condensate formation assay using the light-inducible system, including the CAHS2 positive control showing condensates only upon light induction and several ApoE examples that form condensates both before and after induction. **(f)** Condensate formation assay using the chemically-inducible system, including the CAHS2 positive control showing condensates only upon added dimerizer and ApoE4 (47-300) exhibiting condensates both before and after induction. **(g)**

54   Apoptosis data showing reduction in apoptosis upon condensate induction with dimerizer.
55   Data are normalized to mean apoptosis level of uninduced (i.e. no dimerizer) sample for
56   each construct.

Fig. 5. DHR81 condensate formation and apoptosis reduction. (a) Diagram of
DHR81, illustrating a charge distribution that can support oligomerization. (b) Sample
images highlighting how the synthetic protein DHR81 supports chemically-induced
condensates. (c) Apoptosis is further reduced in the presence of dimerizer (i.e. with
induced DHR81 condensates). The FKBP-modifications alone, which enable chemically-
induced multi-valency, have a minimal effect on apoptosis. (d) Activated caspase-7
(*green*) co-localizes with DHR81 condensates but not CAHS2 or ApoE4 (47-300)
condensates (*magenta*). *White* indicates colocalization.

Figure 3 Supplement (a) Graph and sequences of various LEA repeats tested, with full sequences and performance. **(b)** Graph and sequences of extended LEA repeats tested.

71 **(c)** Correlation analysis for relationship between hydrophobic moment with apoptosis
72 assay performance.

Figure 4 Supplement (a) Western blot data of some ApoE constructs showing expression at the expected molecular weight (b) Sample images showing condensates forming in the full length ApoE3 D125I mutation.

79 **Figure 5 Supplement** (a) DHR81 inducible condensates reduce apoptosis. (b) Sample
80 images showing no co-localization between activated caspase 3 and condensates.