Natural and designed proteins inspired by extremotolerant organisms can form

condensates and attenuate apoptosis in human cells

Mike T. Veling[1,2,7], Dan T. Nguyen[1,2,7], Nicole N. Thadani[1], Michela E. Oster[1,2], Nathan J.

Rollins[1,2], Kelly P. Brock[1], Neville P. Bethel[3,4], David Baker[3,4,5], Jeffrey C. Way[1,2], Debora

S. Marks[1,6], Roger L. Chang[1,2,8], Pamela A. Silver[1,2,8]

Affiliations

1. Department of Systems Biology, Harvard Medical School, Boston, MA, 02115, USA;

2. Wyss Institute for Biologically Inspired Engineering, Boston, MA, 02115, USA;

3. Institute for Protein Design, University of Washington, Seattle, WA, 98195, USA;

4. Department of Biochemistry, University of Washington, Seattle, WA, 98195, USA;

5. Howard Hughes Medical Institute, University of Washington, Seattle, WA, 98195, USA;

6. Broad Institute of Harvard and MIT, Cambridge, MA, 02142, USA;

7. Co-first authors

8. Co-corresponding authors

Key words

Stress tolerance, Phase separation, Intrinsically disordered proteins, Apoptosis

ABSTRACT

Many organisms can survive extreme conditions and successfully recover to normal life. This extremotolerant behavior has been attributed in part to repetitive, amphipathic, and intrinsically disordered proteins that are upregulated in the protected state. Here, we assemble a library of approximately 300 naturally-occurring and designed extremotolerance-associated proteins to assess their ability to protect human cells from chemically-induced apoptosis. We show that proteins from tardigrades, nematodes, and the Chinese giant salamander are apoptosis protective. Notably, we identify a region of the human ApoE protein with similarity to extremotolerance-associated proteins that also protects against apoptosis. This region mirrors the phase separation behavior seen with such proteins, like the tardigrade protein CAHS2. Moreover, we identify a synthetic protein, DHR81, that shares this combination of elevated phase separation propensity and apoptosis protection. Finally, we demonstrate that driving protective proteins into the condensate state increases apoptosis protection, and highlight the ability for DHR81 condensates to sequester caspase-7. Taken together, this work draws a link between extremotolerance-associated proteins, condensate formation, and designing human cellular protection.

INTRODUCTION

When exposed to extreme environmental stress, extremotolerance-associated (*ExTol*) organisms enter a state of reduced cellular metabolic activity that promotes survival in harsh conditions. This state has been shown to protect these organisms from high salinity, desiccation, and even the vacuum of outer space [1]. To acclimate to these conditions,

ExTol organisms significantly upregulate the production of protective metabolites, like

trehalose [2], as well as tolerance-conferring proteins [3], many of which are intrinsically

disordered proteins (*IDPs*) with repetitive sequence and predicted amphipathic helices.

While the mechanism of protection is unknown, one model suggests these proteins

mediate formation of a gel- or glass-like material state within an entire cell to protect it

from damage [3]. The model posits that such proteins promiscuously interact with

membranes and proteins, increasing intracellular viscosity and protecting vulnerable

structures [4-6] (Figure 1A Top). Such interactions would also work to prevent stress-

induced protein unfolding and protein aggregation [7]. An alternative model suggests that

these proteins form intracellular phase-separated membraneless compartments with

distinct physical parameters (referred to here as *condensates*) that partition key stress-

sensitive cellular components [8] (Figure 1A Bottom). Within these dense protein

structures, essential components are sequestered and protected from damage. The

models are not mutually exclusive; for example, in the case of desiccation tolerance,

continued dehydration could lead phase-separated compartments to fuse, comprise

much (if not all) of the cell, and promote a glassy material state.

Figure 1: Selection of candidate extremotolerance-associated proteins (a) Illustration of different protective models for cellular stress tolerance. **(b)** Illustration of amphipathic membrane interaction for rotifer LEA4 protein and Alpha-synuclein, thus highlighting a feature indicative of extremotolerance-associated proteins. **(c)** Hydrophobic moment analysis of rotifer LEA4, showing the repetitive nature of the amphipathic signature. Highlighted green segment corresponds to the LEA4 helix in (b). **(d)** Venn diagram of the lines of evidence used to select targets in the final library. Abbreviations: KD, knockdown; OE, overexpression; Δ, change in expression levels. **(e)** Domain analysis of included targets reveals LEA and ApoE domains in almost half of the targets tested. The other half contains proteins with other PFAM domains (19%) or without any affiliated PFAM entry (32%). Of the proteins lacking a known PFAM annotation, many exhibit repetitive sequence (20% of all targets).

The resurgence of interest in biological phase separation, extensively reviewed in [9-12],

has shown that human cells can employ similar strategies to survive stressful conditions.

While the reversible formation of stress granules in healthy cells is the archetype of stress

tolerance, recent studies have revealed phase separation as a key adaptation that cancer

cells use to avoid apoptosis and cell death [13-14]. Phase-separated condensates have been

implicated in sequestering key tumor repressors[14], enhancing or repressing gene

expression in response to stress [15], and regulating mRNA or protein function through

sequestration [16]. These processes parallel the mechanisms that ExTol proteins are

thought to use to protect complex biological systems, suggesting there could be some

functional overlap. That is, sequence features and properties within ExTol proteins could

be implicated in mammalian stress tolerance pathways, including apoptosis resistance.

A key sequence feature of some ExTol proteins is the presence of amphipathic helices

[17]. While disordered in most contexts, ExTol proteins can transiently form amphipathic

helices in certain molecular environments, notably when interacting with and stabilizing

membranes [6]. The helices often exhibit a hydrophobic face, which interacts with core

lipids within the membrane, flanked by positively charged residues that might interact with

the negatively charged lipid head groups (Figure 1B). This amphipathic moment often

repeats across the length of the protein (Figure 1C). This feature is a hallmark of the late

embryonic abundant (LEA) class of ExTol proteins [18]. Human membrane interacting

proteins, like alpha-synuclein, also exhibit a similar repeat motif [19]. Such parallels in

sequence features suggest that orthogonal expression of ExTol proteins may confer their

protective properties to other biological systems, like human cells.

Indeed, previous reports have shown that some of these ExTol proteins provide similar

protective properties when expressed heterologously. Expressing tardigrade-specific

IDPs (TDPs) in bacteria provides protection against desiccation [3]. Anhydrin, an IDP

upregulated by *Aphelenchus avenae* during dehydration, expressed in a human cell line

limited PABPN1-A17 aggregation, a hallmark of oculopharyngeal muscular dystrophy [7].

Similar reports demonstrated protection of human cells provided by expression of LEA

proteins originating from the African midge [20] and by TDPs from tardigrades [21]. These

findings reinforce the promise of transferring protective phenotypes from their native

contexts to human cells.

Taken together, these data indicate a potential avenue for using this class of proteins to

protect biological systems in a controlled manner. To test this hypothesis, we generated

approximately 300 sequences inspired by ExTol proteins and examined their ability to

protect against apoptotic cell death. Our results show that human proteins containing

sequence hallmarks of ExTol proteins as well as *de novo* designed condensate forming

proteins can protect human cells from chemically-induced apoptosis. We also

demonstrate that condensate induction using synthetic modifications can strengthen

apoptotic protection. Finally, we show that key components of the apoptotic pathway can

localize to inducible condensates, suggesting enzyme sequestration could be a

mechanism of action.

RESULTS

Assembling a library of stress tolerance proteins

Based on the hallmark sequence features of ExTol proteins, we searched the published

literature to identify proteins that not only contain these features but also exhibit some

evidence for protection of biological structures. The library was generated based on over

one hundred naturally-occurring proteins based on five lines of evidence for stress

tolerance association, including: change in expression during extremotolerance,

knockdown or overexpression changing the tolerant state, *in vitro* protection, orthogonal

expression transferring tolerance, and sequence similarity to other proteins with these

features (Figure 1D and Supplemental table 1).

A PFAM domain analysis of the identified proteins reveals almost half of the identified

targets have two key domains: Late Embryonic Abundant (LEA) and Apolipoprotein E

(ApoE) domains (Figure 1E, Supplemental table 1). LEA proteins are well known to be

upregulated during the last stages of plant seed maturation [22]. This domain is thought to

protect the seed embryo during transport to its new destination. There is also evidence

that this domain has been co-opted by bacteria as protection from desiccation and

radiation [23]. ApoE proteins interact with lipid particles to allow for lipid reuptake into cells.

Additionally, mutations within human ApoE have been identified as strong genetic

predictors of Alzheimer's disease in human populations [24]. Based on their lipid interaction

properties and potential role in neuronal homeostasis, as well as its prevalence in the

a — Isolate repetitive units —

☒ HHRepID repeat

Remove non-repetitive elements

b — Modulate amphipathicity —

LEA 22mer
Mutant #10

LEA 22mer

Minimize
Moment

LEA 22mer
Mutant #5

Maximize
Moment

Positive
Negative
Non-polar
Other

c — De novo protein design —

Alternating
Charges

**Oligomerizing
Interactions**

Electrostatic

Multivalency favors
phase separation

Hydrophobic

Non-polar
Face

d — Library composition —

Natural
(104)

Inspired by
nature
(166)

de novo (19)

Figure 2: Design based on candidate extremotolerance-associated proteins. (a) Truncations were made to isolate regions of repetitive and amphipathic sequence. **(b)** Small helices were selected and altered based on their hydrophobic moment. **(c)** Synthetic alpha helix-containing proteins designed to oligomerize. **(d)** Venn diagram highlighting the library that was tested.

PFAM domain analysis, we decided to include human ApoE and its various mutations as ExTol proteins in our study.

Designing novel extremotolerance-associated proteins

To test hypotheses about sequence features and their relationship to phase separation and apoptosis, we designed novel proteins based on the set of ExTol proteins. These new designs take 3 approaches (Figure 2): truncation of proteins to focus on repetitive, intrinsically disordered, and amphipathic regions; isolation of individual helices with desired sequence features; and *de novo* design of alpha helix-containing proteins that self-assemble into higher order structures. Through these designs, we sought to identify the minimal elements required for protection and phase separation.

First, we identified regions within natural proteins that contain intrinsic disorder [25],

repetitive sequences [26], and amphipathic helices[27]. Figure 2A illustrates the use of

HHRepID on the sequence of full length human ApoE. Based on this analysis, we

identified the repeating domain within ApoE for further testing. Other ExTol proteins were

similarly modified and examined.

We also included short peptides based on the consensus sequence of repetitive regions

that have previously been shown to be protective [28]. To test if any of the sequence

features we identified are important for apoptosis protection, we altered these features

within these short peptides. For example, we altered the amphipathic character of these

helices (Figure 2B) to determine if apoptosis protection was changed.

To examine whether generic condensate formation is sufficient to confer apoptosis

protection, we also examined completely synthetic alpha helix-containing proteins that

were designed to form repetitive structure with sequences orthogonal to naturally-

occurring proteins [29]. These designed helical repeat (DHR) proteins interact along the

axis of the helix, forming long spiraling fibrils, but may also potentially self-interact along

the face of the protein based on a pattern of positive and negative charges (Figure 2C).

This ability to oligomerize and form multivalent complexes, which may then interact via

weak non-covalent interactions, is an established determinant of condensate formation

[12].

Together, our library consists of 104 naturally-occurring proteins, 166 truncations or

isolations of helical regions, and 19 *de novo* designs (Figure 2D). Taken altogether, these

proteins interrogate several hypotheses about how disordered proteins may protect

human cells against apoptosis.

Apoptosis assay identifies extremotolerance-associated proteins in human cells

To model stress in a mammalian cell system, we induced cellular apoptosis by treatment

with the chemotherapeutic camptothecin (CPT), a DNA topoisomerase inhibitor well

known to trigger apoptosis [30-31]. Under these conditions, protection against apoptosis

could be achieved through a number of avenues, including protection against the DNA

damage caused by CPT-induced topoisomerase inhibition [30], interruption of the signaling

cascade, or even protection of the mitochondrial membrane whose rupture ultimately

triggers apoptosis [32].

To test our designs for protection against apoptosis, we used a human cell-based

apoptosis assay (Figure 3A). This assay uses HT-1080 fibrosarcoma cells from ATCC [33].

These cells are seeded in a 96-well plate and allowed to adhere overnight. The next day,

the cells are co-transfected with a plasmid containing the ExTol candidate proteins of

interest as well as a transfection marker. The following day, the media is swapped to one

containing 13.2 µM CPT. After addition of CPT, caspase-3 and -7 activation is measured

by microscopy every 45 minutes for 1 day using a fluorophore quencher pair attached to

either end of a peptide with the sequence "DEVD". As caspase-3 and -7 are activated,

the DEVD peptide is cleaved, releasing the fluorophore from the quencher. The

unquenched fluorophore localizes to the nucleus where it can be quantified by imaging

on a single-cell level. The single-cell resolution allows for the separation of transfected

and non-transfected cells based on the transfection marker. Only single cells with the

transfection marker were subjected to further analysis; Figure 3B shows sample images

highlighting the type of data being analyzed.

On every plate, an empty vector negative control is run as well as a positive control

containing the X-linked inhibitor of apoptosis (XIAP). This protein is known to bind to

caspase-3 and -7 to block their activity [34] . Figure 3C shows example data comparing

median DEVD peptide signal intensity over time after induction with CPT of four wells on

the same plate transfected with XIAP and empty vector. The results show a reduction in

DEVD peptide signal intensity when the XIAP-containing vector is transfected,

demonstrating that the assay can identify modulators of apoptosis.

To convert these time-resolved data to a measure of protection against apoptosis, the

maximum DEVD peptide signal intensity is calculated and normalized to a plate-matched

empty vector control (see methods for further details). The normalized apoptosis induction

intensity is shown in Figure 3D left. The data are slightly skewed towards apoptosis-

protection, which supports our hypothesis that the targeted proteins can protect against

apoptosis.

Figure 3: Screen for CPT resistance by extremotolerance-associated proteins (a) Flow diagram of the assay. **(b)** Representative images showing cells during the assay. Cells are identified using SiR-Hoechst marking the nucleus (*blue*). Transfected cells are identified by the presence of the mCherry marker (*green*). Finally, the degree of apoptosis is identified by the presence of the cleaved DEVD peptide (*red*). **(c)** Positive control showing apoptosis reduction upon expression of XIAP. **(d)** Bar graph highlighting the apoptosis resistance provided by different constructs, including zoomed in area to highlight the top hits in the assay. Source species of a sequence are abbreviated and shown in italics; abbreviations are drawn from Uniprot, including *RAMVA* (*Ramazzottius varieornatus*), *HYPDU* (*Hypsibius dujardini*), *PARRC* (Paramacrobiotus richtersi), *ANDDA* (*Andrias davidianus*), *APHAV* (*Aphelenchus avenae*), *ARTSF* (*Artemia franciscana*), *CRAPL* (*Craterostigma plantagineum*), and *DEIRA* (*Deinococcus radiodurans*).

To highlight the top-performing designs, the top 10% of the targets are shown in Figure

3D right. As expected, the XIAP positive control showed the strongest anti-apoptotic

effect. The two top-performing hits were ApoE designs. Human alpha-synuclein is the 5th

hit, which is expected as it is known to be both intrinsically disordered, and anti-apoptotic

[35]. Additionally, several ApoE designs and short LEA helices are top hits. The data used

to generate these figures is included in supplemental table 2.

Known extremotolerance-associated proteins affect apoptosis

Among the top hits in apoptosis reduction are an IGKV protein highly expressed in mucus

of the Chinese giant salamander *Andrias davidianus* [36] that is used as a medical glue in

traditional Chinese medicine, an LEA protein from the nematode *Aphelenchus avenae*

previously shown to have the ability to reduce a form of protein aggregation associated

with oculopharyngeal muscular dystrophy in human cells [7], and a mitochondrial LEA

protein from the brine shrimp *Artemia franciscana* shown to protect human cells during

desiccation [20] (Figure 3D). The prominence of these ExTol proteins suggests the assay

can help identify other proteins that can protect cells from more general stresses beyond

CPT-induced apoptosis.

Short repetitive peptides and *de novo* designs modulate apoptosis

LEA repeat-containing peptides are among the top-performing hits in the screen. While

the original short LEA designs from Furuki et al. 2019 had a minor effect reducing

apoptosis, several of the designs tested here performed better (Supplemental Figure 3A)

[28]. As 22 amino acids is small and could lead to peptide stability issues, longer 50-amino

acid designs inspired by the same LEA proteins Furuki et al. utilized were also developed

(Supplemental Figure 3B). Some of these designs were also able to reduce apoptosis,

including the PvLEA4 with extended repeats (notably PvLEA4 repeats extended #2),

which fell into the top hits of the entire library (Figure 3D and Supplemental Figure 3B).

These results suggest that optimization of synthetic designs yields proteins with stronger

function than naturally-occurring proteins.

While these proteins were designed to interrogate the relationship between sequence

and apoptosis protection in short peptides, little relationship was found with respect to the

sequence features we analyzed (Supplemental Figure 3C). Several designs were made

that would alter the hydrophobic moments of putative helices these peptides are predicted

to form. To examine if this sequence property relates to the stress-tolerance function of

these proteins, we attempted to correlate this property with assay performance. However,

no significant correlation was found, suggesting (at least in these cases) that degree of

amphipathicity on its own is not the key determinant of stress-tolerance function of these

proteins.

The repeat region of ApoE comprises its minimal anti-apoptotic unit

The top two hits in the assay were ApoE4 (47-300) and ApoE2 (47-300). These truncated

forms of ApoE maintain the repeat domains shown in Figure 4A, while removing the signal

peptide and terminal regions. This suggests that the repeat region of ApoE accounts for

its core anti-apoptotic functionality. Interestingly, the full length ApoE2 and E4 do not

show up as top hits, suggesting that these truncated forms improve on the apoptosis

protection provided by the full-length variants (Figure 4B).

Based on the location of the signal peptide, it is expected that ApoE4 should localize to

the secretory pathway whereas the truncated form may be cytoplasmic. To test this

hypothesis, cells were subjected to Western blot and immunofluorescence analysis using

commercially available antibodies (Abcam ab24139; Figure 4C and Supplemental Figure

4A). The Western blot results show that both the full-length and truncated forms of ApoE4

are expressed at the expected size (Supplemental Figure 4A). The immunofluorescence

shows co-localization between the ApoE4 full length protein and the endoplasmic

reticulum (marked by Thermo's PDI antibody MA3-019, Figure 4C). However, the

truncated form did not show the same co-localization, suggesting as expected that it may

not enter the secretory pathway as the full-length version does.

Strikingly, the truncated version appeared to form more pronounced puncta throughout

the cytoplasm of the cell, suggesting the formation of potentially phase-separated

condensates. In screening through the point mutants of ApoE also tested here, we found

a similar condensate phenotype with the full length ApoE3 D125I mutation (Supplemental

Figure 4B). Together, these data suggest that ApoE may be able to support condensate

formation and confer apoptosis protection in part via the protective compartment

mechanism (Figure 1A bottom).

Fig. 4. ApoE truncations and synthetic condensates further reduce apoptosis.
(a) Gene diagram highlighting the repetitive region of ApoE. **(b)** Boxplots highlighting the apoptosis assay performance of the full length and truncated forms of ApoE2 and E4. **(c)** Immunofluorescence of ApoE4, full length and truncated forms. *White* indicates colocalization, which is more apparent with the full-length protein. The truncated form appears to exhibit larger and more distinguishable puncta. **(d)** Diagram of the induced multi-valency systems used here, including the light-inducible system (*top*) and the chemically-inducible system (*bottom*). **(e)** Condensate formation assay using the light-inducible system, including the CAHS2 positive control showing condensates only upon light induction and several ApoE examples that form condensates both before and after induction. **(f)** Condensate formation assay using the chemically-inducible system, including the CAHS2 positive control showing condensates only upon added dimerizer and ApoE4 (47-300) exhibiting condensates both before and after induction. **(g)** Apoptosis data showing reduction in apoptosis upon condensate induction with dimerizer. Data are normalized to mean apoptosis level of uninduced (i.e. no dimerizer) sample for each construct.

CAHS2 and ApoE form condensates upon induced multivalency

Established condensate engineering tools [37] were used to test for latent phase separation

propensity within the top performing proteins. We used both light- and chemically-

inducible systems of multivalency. In the light system, a ferritin core (comprised of self-

assembled monomers, FTH1 domain-eGFP-iLID) becomes decorated with SspB-

modified proteins upon exposure to blue light [38] (Figure 4D top). In the chemical system,

multivalency (i.e. multiple arms of the ExTol protein of interest) is induced using a

dimerization agent [39] that brings together FKBP-modified proteins [37] (Figure 4D bottom).

These inducible methods increase the valency of a system, and thus work to favor phase

separation. In this way, they can help reveal any latent ability to form condensates that is

not immediately observable without modification.

Indeed, condensates were observed in both inducible systems using ExTol proteins.

CAHS2, a protein from the tardigrade *P. richtersi*, exhibited the most dramatic phenotype,

which is consistent with recent work describing the gel forming capabilities of CAHS

proteins [40] (Figure 4E&F). ApoE variants were of particular interest, based on the puncta

observed in the top performing ApoE constructs. All versions of ApoE tested formed

puncta without activation, consistent with our previous observation using untagged native

ApoE (Figure 4C). Upon light- or chemical-activation, there is a noticeable increase in

condensates, exemplifying how phase separation can be enhanced with increased

multivalency (Figure 4E&F).

Induction of condensates reduces apoptosis

Induction of multivalency in ApoE containing constructs reduces apoptosis. We performed

the apoptosis assay using the top hits modified for chemically-induced multivalency,

namely fusion to two FKBP domains. The results show that the presence of the chemical

dimerizer did not affect apoptosis signal in cells transfected with the empty vector; in

contrast, the ApoE constructs exhibit reduced apoptosis with added dimerizer (Figure

4G). That is, artificially increasing ApoE condensates was apoptosis protective. These

data suggest that condensate formation plays a role in the anti-apoptotic effect of these

proteins.

Synthetic DHR81 condensates reduce apoptosis and sequester caspase-7

Induction of DHR81 based condensates protects against apoptosis. DHR81 is formed

from four helical hairpin repeat regions; the core of the protein is hydrophobic to ensure

consistent folding, while the outside of the protein is decorated with checkerboard positive

and negative charges (Figure 5A) [29]. This structure ensures the protein is well folded,

while also enabling the potential for higher order structure formation via electrostatic

interactions. Indeed, as with the ApoE variants, DHR81 supported condensates in the

chemically-inducible condensate system (Figure 5B) and exhibited a stronger anti-

apoptotic phenotype when multivalency is induced (Figure 5C and Supplemental Figure

5A). There was little to no effect in cells containing an empty vector or expressing the

FKBP dimerization domains alone.

Fig. 5. DHR81 condensate formation and apoptosis reduction. (a) Diagram of DHR81, illustrating a charge distribution that can support oligomerization. **(b)** Sample images highlighting how the synthetic protein DHR81 supports chemically-induced condensates. **(c)** Apoptosis is further reduced in the presence of dimerizer (i.e. with induced DHR81 condensates). The FKBP-modifications alone, which enable chemically-induced multi-valency, have a minimal effect on apoptosis. **(d)** Activated caspase-7 (*green*) co-localizes with DHR81 condensates but not CAHS2 or ApoE4 (47-300) condensates (*magenta*). *White* indicates colocalization.

To further explore how condensates mediated by ExTol proteins may confer an anti-apoptotic effect, we examined whether they sequester major components of apoptotic signaling, namely caspase-3 and -7. The results showed that activated caspase-7 localized to DHR81 condensates but not CAHS2 or ApoE4 condensates (Figure 5D). Caspase-3, however, does not localize to either DHR81, CAHS2, or ApoE4 condensates (Supplemental Figure 5B). These results suggest sequestration of key apoptotic factors could be a mechanism by which these condensates reduce apoptosis.

Discussion

To broadly evaluate the ability of ExTol proteins to protect orthogonal systems from environmental stress, we studied over one hundred different ExTol proteins. We used these as a basis to design approximately 300 variants, including truncations and fusion proteins. These designs

were evaluated on their ability to protect human HT-1080 cells against CPT-induced

apoptosis. The results showed that known ExTol proteins, ApoE variants, and the

synthetic DHR81 protein could modulate apoptosis levels when transferred to human

cells.

Protection against apoptosis is a key feature of many ExTol proteins. As apoptosis is a

common cellular response to stress, the ability to mitigate this response is important in

providing cell-autonomous protection in stressful environments. While the mechanism

resilient organisms use to survive is still not fully understood, it is becoming increasingly

clear that phase separation may play an important role.

ExTol proteins are proposed to form expansive higher order structures (e.g. gels or glassy

systems) that reduce the viscosity of the cytoplasm, especially in desiccated states [3].

Other reports show that these proteins could mediate formation of protective phase-

separated compartments (i.e. condensates), sequestering damage-inducing proteins

and/or protecting important targets [8]. Based on our condensate data, it appears the latter

may be at play for proteins like ApoE or DHR81. Notably, in these proteins that already

exhibit both anti-apoptotic effects as well as elevated phase separation propensities,

increasing the amount of condensates via an engineered inducible system improved

protection against apoptosis. This observation suggests that condensates could be a key

part of the protective mechanism.

Interestingly, we observed that activated caspase-7 partitions into DHR81 condensates,

suggesting caspase sequestration as a potential anti-apoptotic mechanism. Phase-

separated condensates and higher order assemblies are already implicated in mediating

apoptosis: stress granules inhibit apoptosis by suppressing MAPK pathways [41]; the

apoptosome relies on Apaf-1 multimerization, which has been reported to undergo

amyloid fibrillation in specific conditions [42]; and stress-dependent phase separation

modulates proteasome formation, whose function affects apoptosis [43-44]. It is interesting

to note that we only observe clear caspase-7 partitioning into DHR81 condensates,

suggesting the possibility of a specific binding interaction in this case. ApoE condensates

may be sequestering other apoptotic factors or using a different mechanism to reduce

apoptosis.

It is also possible that these ExTol proteins achieve their anti-apoptotic effect by

implementing a more general physical mechanism, namely slowing the intracellular

mobility of biological effectors. There are various reports describing how the effective

viscosity within the cytoplasm can be increased in response to external stresses [4-5] or

mediated by protein composition, specifically intrinsically disordered proteins [45]. Notably,

a pervasive cell-spanning gel or network is not necessary to achieve such an effect; it is

possible that the presence of condensates, coupled with their ability to selectively partition

biomolecules (i.e. control what biomolecules can enter and diffuse through them), could

broadly slow biomolecular movement. That is, by forcing different proteins to either be

trapped in them or diffuse around them, condensates can decrease protein mobility and

perturb e.g. signaling pathways. Our results showing a lack of caspase-7 sequestration

by some of our protein condensates could thus support this more general mechanism of

condensates perturbing biology.

Condensates are becoming increasingly implicated in disease [10], particularly when their

structural metastability and reversibility is dysregulated (e.g. through mutation). Notably,

drugs that block the formation of amyloid fibrils and plaques, which are products of phase

separation dysregulation, are promising yet strongly debated [46] therapies for many

neurological diseases. As such, our observations of ApoE are particularly salient. We

found that expression of a cytoplasmic form of ApoE is protective against apoptosis. In

particular, the truncated repeat region of ApoE is sufficient and shows the strongest anti-

apoptotic activity of all designs we tested. Others have shown similar effects for

exogenously added ApoE-containing liposomes protecting rat neurons against apoptosis

[47], though we show the effect through overexpression of ApoE inside of human cells.

When combined with our condensate results and the clinical relevance of ApoE in

Alzheimer's disease [48], these observations suggest the anti-apoptotic effect observed

here warrants further study, to further elucidate its role in Alzheimer's disease

progression.

Altogether, our work expands upon the understanding of the relationship between

condensate formation and cell physiology, demonstrating how condensate accumulation

can be broadly beneficial in reducing apoptosis while also identifying diverse new protein

domains that confer protection. The proteins and concepts presented here demonstrate

the utility of this class of proteins for engineering new properties into cells and ultimately

higher organisms.

Methods

Cell culturing and maintenance

HT-1080 cells (ATCC, CCL-121) were maintained in EMEM medium supplemented with

10% FBS and Penicillin-Streptomycin according to ATCC's protocol. For additional details

see supplemental methods.

DNA Preparation

Apoptosis constructs were synthesized from TWIST and transformed into DH5α cells for

amplification and plasmid purification. Condensate assay constructs were subcloned from

these synthesized constructs into vectors containing the appropriate chemically or light

inducible condensate elements. DNA was prepared using the QIAGEN Plasmid Plus Midi

Kit (Qiagen 12943) for transfection grade DNA purification. See supplemental methods

for additional details.

Apoptosis assay

200 μL of HT-1080 cells at to 7.5×10^4 cells per mL (15000 cells per well) were added to

each well of a microscope grade 96 well plate (Corning 3603). Plates were then

centrifuged at 330 rpm for 3 mins (Eppendorf, Centrifuge 5810). Plates were then

incubated at 37°C for 24 hours. Cells were then transfected with appropriate DNA

constructs with the lipofectamine 3000 kit (Thermo Fisher Scientific, L3000015) as

specified by the manufacturer. 24 hours after transfection, the transfection media was

removed and replaced with FluoroBrite DMEM Media (Thermo A1896702) supplemented

with 1x GlutaMAX (Thermo 35-050-061) and 10% FBS (Thermo Fisher Scientific,

10082147). This media also contained 13.2 µM (S)-(+)-Camptothecin (Sigma C9911-

100MG), 0.3 µM SiR-DNA stain (SpiroChrome CY-SC007), and 1.25 µM of the DEVD

peptide for detecting caspase activation (Sartorius 4440). The plates were sealed with

sterile Breathe-Easy film (USA scientific 9123-6100) and incubated at 37°C with 5% CO_2

for 30 minutes. After 30 min incubation, the plates were placed in an ImageXpress-

confocal plate imager with CO_2, humidity, and temperature control set to 37°C with 5%

CO_2 (Molecular Devices). Images of all wells were taken every 45 minutes and stored on

a remote server for later analysis. See supplemental methods for further details on this

method as well as the image analysis pipeline used.

Confocal microscopy

Confocal microscopy was done at the Nikon Imaging Center at Harvard Medical School

using a Nikon Ti fluorescence microscope attached to a Yokogawa W1 spinning disk

confocal. See supplemental methods for further details on the equipment and settings

used.

Light inducible condensate assay

HEK293-T cells were seeded on 96 well glass bottom plates from Cellvis (P96-1.5H-N).

Cells within each well were transfected with 50 ng of the LLPS1-iLID::EGFP::FTH1

(Addgene plasmid #122149) construct containing the core FTH1 scaffolding domain

attached to iLID for light inducible interactions. The cells were also transfected with 50 ng

of a construct containing the target protein attached to mCherry and SspB. The

transfection was allowed to continue overnight before the media was swapped to

FluoroBrite media supplemented with 10% FBS and 1x GlutaMAX. The cells were then

imaged using the settings listed above for mCherry and eGFP for the pre-induction

images. The cells were then exposed to 1 second of the DAPI imaging protocol described

in supplemental methods before new images were captured on the mCherry and eGFP

settings.

Chemically inducible condensate assay

HT-1080 cells were seeded on 96 well glass bottom plates from Cellvis (P96-1.5H-N).

Cells within each well were transfected with 50 ng of target construct fused to a mCherry-

2xFKBP domain. The transfection was allowed to continue overnight before the media

was swapped to FluoroBrite media supplemented with 10% FBS, 1x GlutaMAX, and 300

nM SiR-Hoechst. The cells were then imaged using the settings listed in the supplemental

methods for mCherry and SiR-Hoechst for the pre-induction images. Then, B/B dimerizer

(AP20187 ligand; TaKaRa 635058) was added to a final concentration of 500 nM in the

well. Imaging continued for 15 minutes with new images every 45 seconds.

Caspase assay with chemically induced condensates

These caspase assays are performed similarly to the standard caspase assays with a

few key differences. Briefly, HT-1080 cells were seeded as normal on to 96 well plates

and allowed to adhere overnight. The next day, 12.5 ng of nuclear mCherry marker and

37.5 ng of the target construct fused to mCherry-2xFKBP was transfected into the cells

as described above. The next day, 500 nM of the B/B dimerizer (TaKaRa 635058) was

added 2 hours before the start of the assay (or an Ethanol vehicle control was added).

The media was then removed and replaced with the apoptosis assay media described

above with the addition of 500 nM B/B dimerizer or a vehicle control. The rest of the assay

proceeded as normal.

Abbreviations

ExTol: extremotolerance-associated

IDP: Intrinsically Disordered Protein

LEA: Late Embryonic Abundant (protein)

ApoE: Apolipoprotein E

DHR: Designed helical repeat (protein)

CPT: Camptothecin

ATCC: American Type Culture Collection

XIAP: X-linked inhibitor of apoptosis

CAHS: Cytosolic Abundant Heat Soluble (protein)

FTH1: Ferritin Heavy Chain 1

iLID: light inducible dimer

SspB: Stringent Starvation Protein B

eGFP: Enhanced Green Fluorescent Protein

FKBP: FK506 binding protein

PvLEA: *Polypedilum vanderplanki* Late Embryonic Abundant (Protein)

MAPK: Mitogen-activated protein kinase

Apaf: Apoptotic protease activating factor

EMEM: Eagle's Minimum Essential Medium

FBS: Fetal bovine serum

DMEM: Dulbecco's Modified Eagle Medium

Author Information.

Correspondences should be sent to Pamela Silver, Harvard Medical School Department

of Systems Biology 200 Longwood Avenue Warren Alpert Building Boston, MA 02115.

Author Contributions

M.T.V. and D.T.N. designed and performed experiments, analyzed the data, and wrote

the paper. N.N.T. analyzed the data and wrote the paper. M.E.O provided technical

support. N.J.R., K.B.B., and N.P.B., designed constructs. D.B. and J.C.W. managed the

teams and provided key insight. D.S.M., R.L.C., and P.A.S., designed constructs,

managed teams, and wrote the paper.

Acknowledgements

Research was sponsored by the NSF, DARPA, the Wyss Institute for Biologically inspired

engineering, and the Army Research Office and was accomplished under Cooperative

Agreement Number W911NF-19-2-0017. M.T.V. was sponsored by the National Science

Foundation under Grant No. 2010370. D.T.N., N.N.T., M.E.O., N.J.R., K.P.B., J.C.W,

D.S.M., and P.A.S. were supported by the Harvard Medical School Department of

Systems Biology. J.C.W. was further supported by the Harvard Medical School

Laboratory of Systems Pharmacology. K.P.B. and D.S.M. were further supported by CZI

Neurodegenerative Challenge Ben Barres Early Career Award (CZI2018-191853). N.P.B

was further supported by the HHMI Hanna Gray fellowship (grant GT11817). We would

like to acknowledge Clarence Yapp for help with microscopy as well as the NCI U54-

CA225088 grant and a gift from the Ludwig Institute for Cancer Research to the

Laboratory for Systems Pharmacology. We would like to acknowledge the Nikon Imaging

Center at Harvard Medical Center for providing microscope support. The views and

conclusions contained in this document are those of the authors and should not be

interpreted as representing the official policies, either expressed or implied, of any funding

agency including the NSF, DARPA, the Army Research Office, or the U.S. Government.

The U.S. Government is authorized to reproduce and distribute reprints for Government

purposes notwithstanding any copyright notation herein.

549
550 **Bibliography & References**
551

1.   Jonsson, K. I.; Rabbow, E.; Schill, R. O.; Harms-Ringdahl, M.; Rettberg, P., Tardigrades
survive exposure to space in low Earth orbit. *Curr Biol* **2008,** *18* (17), R729-R731.
2.   Thevelein, J. M., Regulation of trehalose mobilization in fungi. *Microbiol Rev* **1984,** *48*
(1), 42-59.
3.   Boothby, T. C.; Tapia, H.; Brozena, A. H.; Piszkiewicz, S.; Smith, A. E.; Giovannini, I.;
Rebecchi, L.; Pielak, G. J.; Koshland, D.; Goldstein, B., Tardigrades Use Intrinsically Disordered
Proteins to Survive Desiccation. *Mol Cell* **2017,** *65* (6), 975-984 e5.
4.   Dijksterhuis, J.; Nijsse, J.; Hoekstra, F. A.; Golovina, E. A., High viscosity and anisotropy
characterize the cytoplasm of fungal dormant stress-resistant spores. *Eukaryot Cell* **2007,** *6* (2),
157-70.
5.   Parry, B. R.; Surovtsev, I. V.; Cabeen, M. T.; O'Hern, C. S.; Dufresne, E. R.; Jacobs-Wagner,
C., The bacterial cytoplasm has glass-like properties and is fluidized by metabolic activity. *Cell*
2014, *156* (1-2), 183-94.
6.   Hand, S. C.; Menze, M. A.; Toner, M.; Boswell, L.; Moore, D., LEA proteins during water
stress: not just for plants anymore. *Annu Rev Physiol* **2011,** *73*, 115-34.
7.   Chakrabortee, S.; Meersman, F.; Kaminski Schierle, G. S.; Bertoncini, C. W.; McGee, B.;
Kaminski, C. F.; Tunnacliffe, A., Catalytic and chaperone-like functions in an intrinsically
disordered protein associated with desiccation tolerance. *Proc Natl Acad Sci U S A* **2010,** *107*
(37), 16084-9.
8.   Belott, C.; Janis, B.; Menze, M. A., Liquid-liquid phase separation promotes animal
desiccation tolerance. *Proc Natl Acad Sci U S A* **2020,** *117* (44), 27676-27684.
9.   Hyman, A. A.; Weber, C. A.; Julicher, F., Liquid-liquid phase separation in biology. *Annu*
Rev Cell Dev Biol **2014,** *30*, 39-58.
10.   Shin, Y.; Brangwynne, C. P., Liquid phase condensation in cell physiology and disease.
Science **2017,** *357* (6357).
11.   Boeynaems, S.; Alberti, S.; Fawzi, N. L.; Mittag, T.; Polymenidou, M.; Rousseau, F.;
Schymkowitz, J.; Shorter, J.; Wolozin, B.; Van Den Bosch, L.; Tompa, P.; Fuxreiter, M., Protein
Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol* **2018,** *28* (6), 420-435.

12.   Choi, J. M.; Holehouse, A. S.; Pappu, R. V., Physical Principles Underlying the Complex
Biology of Intracellular Phase Transitions. *Annu Rev Biophys* **2020,** *49*, 107-133.
13.   Jiang, S.; Fagman, J. B.; Chen, C.; Alberti, S.; Liu, B., Protein phase separation and its role
in tumorigenesis. *Elife* **2020,** *9*.
14.   Boija, A.; Klein, I. A.; Young, R. A., Biomolecular Condensates and Cancer. *Cancer Cell*
2021, *39* (2), 174-192.
15.   Boija, A.; Klein, I. A.; Sabari, B. R.; Dall'Agnese, A.; Coffey, E. L.; Zamudio, A. V.; Li, C. H.;
Shrinivas, K.; Manteiga, J. C.; Hannett, N. M.; Abraham, B. J.; Afeyan, L. K.; Guo, Y. E.; Rimel, J.
K.; Fant, C. B.; Schuijers, J.; Lee, T. I.; Taatjes, D. J.; Young, R. A., Transcription Factors Activate
Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell* **2018,** *175* (7),
1842-1855 e16.
16.   Hubstenberger, A.; Courel, M.; Benard, M.; Souquere, S.; Ernoult-Lange, M.; Chouaib, R.;
Yi, Z.; Morlot, J. B.; Munier, A.; Fradet, M.; Daunesse, M.; Bertrand, E.; Pierron, G.;
Mozziconacci, J.; Kress, M.; Weil, D., P-Body Purification Reveals the Condensation of Repressed
mRNA Regulons. *Mol Cell* **2017,** *68* (1), 144-157 e5.
17.   Tolleter, D.; Jaquinod, M.; Mangavel, C.; Passirani, C.; Saulnier, P.; Manon, S.; Teyssier,
E.; Payet, N.; Avelange-Macherel, M. H.; Macherel, D., Structure and function of a mitochondrial
late embryogenesis abundant protein are revealed by desiccation. *Plant Cell* **2007,** *19* (5), 1580-
9.
18.   Leon, D., III, Occurrence of a Repeating 11-Mer Amino Acid Sequence Motif in Diverse
Organisms. *Protein & Peptide Letters* **2001,** *8* (2), 115-122.
19.   Bussell, R., Jr.; Eliezer, D., A structural and functional role for 11-mer repeats in alpha-
synuclein and other exchangeable lipid binding proteins. *J Mol Biol* **2003,** *329* (4), 763-78.
20.   Li, S.; Chakraborty, N.; Borcar, A.; Menze, M. A.; Toner, M.; Hand, S. C., Late
embryogenesis abundant proteins protect human hepatoma cells during acute desiccation.
Proc Natl Acad Sci U S A **2012,** *109* (51), 20859-64.
21.   Tanaka, S.; Tanaka, J.; Miwa, Y.; Horikawa, D. D.; Katayama, T.; Arakawa, K.; Toyoda, A.;
Kubo, T.; Kunieda, T., Novel mitochondria-targeted heat-soluble proteins identified in the
anhydrobiotic Tardigrade improve osmotic tolerance of human cells. *PLoS One* **2015,** *10* (2),
e0118272.
22.   Mertens, J.; Aliyu, H.; Cowan, D. A., LEA Proteins and the Evolution of the WHy Domain.
Appl Environ Microbiol **2018,** *84* (15).
23.   Makarova, K. S.; Aravind, L.; Wolf, Y. I.; Tatusov, R. L.; Minton, K. W.; Koonin, E. V.; Daly,
M. J., Genome of the extremely radiation-resistant bacterium Deinococcus radiodurans viewed
from the perspective of comparative genomics. *Microbiol Mol Biol Rev* **2001,** *65* (1), 44-79.
24.   Yamazaki, Y.; Zhao, N.; Caulfield, T. R.; Liu, C. C.; Bu, G., Apolipoprotein E and Alzheimer
disease: pathobiology and targeting strategies. *Nat Rev Neurol* **2019,** *15* (9), 501-518.
25.   Erdos, G.; Dosztanyi, Z., Analyzing Protein Disorder with IUPred2A. *Curr Protoc*
Bioinformatics **2020,** *70* (1), e99.
26.   Biegert, A.; Soding, J., De novo identification of highly diverged protein repeats by
probabilistic consistency. *Bioinformatics* **2008,** *24* (6), 807-14.
27.   Eisenberg, D.; Weiss, R. M.; Terwilliger, T. C., The helical hydrophobic moment: a
measure of the amphiphilicity of a helix. *Nature* **1982,** *299* (5881), 371-4.

28.   Furuki, T.; Niwa, T.; Taguchi, H.; Hatanaka, R.; Kikawada, T.; Sakurai, M., A LEA model
peptide protects the function of a red fluorescent protein in the dry state. *Biochem Biophys Rep*
2019, *17*, 27-31.
29.   Brunette, T. J.; Parmeggiani, F.; Huang, P. S.; Bhabha, G.; Ekiert, D. C.; Tsutakawa, S. E.;
Hura, G. L.; Tainer, J. A.; Baker, D., Exploring the repeat protein universe through computational
protein design. *Nature* **2015,** *528* (7583), 580-4.
30.   Li, F.; Jiang, T.; Li, Q.; Ling, X., Camptothecin (CPT) and its derivatives are known to target
topoisomerase I (Top1) as their mechanism of action: did we miss something in CPT analogue
molecular targets for treating human disease such as cancer? *Am J Cancer Res* **2017,** *7* (12),
2350-2394.
31.   Wall, M. E.; Wani, M. C.; Cook, C. E.; Palmer, K. H.; McPhail, A. T.; Sim, G. A., Plant
Antitumor Agents. I. The Isolation and Structure of Camptothecin, a Novel Alkaloidal Leukemia
and Tumor Inhibitor from Camptotheca acuminata1,2. *Journal of the American Chemical Society*
1966, *88* (16), 3888-3890.
32.   Bortner, C. D.; Cidlowski, J. A., Cellular mechanisms for the repression of apoptosis.
Annu Rev Pharmacol Toxicol **2002,** *42*, 259-81.
33.   Rasheed, S.; Nelson-Rees, W. A.; Toth, E. M.; Arnstein, P.; Gardner, M. B.,
Characterization of a newly derived human sarcoma cell line (HT-1080). *Cancer* **1974,** *33* (4),
1027-33.
34.   Obexer, P.; Ausserlechner, M. J., X-linked inhibitor of apoptosis protein - a critical death
resistance regulator and therapeutic target for personalized cancer therapy. *Front Oncol* **2014,**
4, 197.
35.   Li, W.; Lee, M. K., Antiapoptotic property of human alpha-synuclein in neuronal cell lines
is associated with the inhibition of caspase-3 but not caspase-9 activity. *J Neurochem* **2005,** *93*
(6), 1542-50.
36.   Geng, X.; Guo, J.; Zang, X.; Chang, C.; Shang, H.; Wei, H.; Xu, C., Proteomic analysis of
eleven tissues in the Chinese giant salamander (Andrias davidianus). *Sci Rep* **2019,** *9* (1), 16415.
37.   Bracha, D.; Walls, M. T.; Brangwynne, C. P., Probing and engineering liquid-phase
organelles. *Nat Biotechnol* **2019,** *37* (12), 1435-1445.
38.   Bracha, D.; Walls, M. T.; Wei, M. T.; Zhu, L.; Kurian, M.; Avalos, J. L.; Toettcher, J. E.;
Brangwynne, C. P., Mapping Local and Global Liquid Phase Behavior in Living Cells Using Photo-
Oligomerizable Seeds. *Cell* **2018,** *175* (6), 1467-1480 e13.
39.   Stanton, B. Z.; Chory, E. J.; Crabtree, G. R., Chemically induced proximity in biology and
medicine. *Science* **2018,** *359* (6380).
40.   Yagi-Utsumi, M.; Aoki, K.; Watanabe, H.; Song, C.; Nishimura, S.; Satoh, T.; Yanaka, S.;
Ganser, C.; Tanaka, S.; Schnapka, V.; Goh, E. W.; Furutani, Y.; Murata, K.; Uchihashi, T.; Arakawa,
K.; Kato, K., Desiccation-induced fibrous condensation of CAHS protein from an anhydrobiotic
tardigrade. *bioRxiv* **2021**, 2021.06.22.449423.
41.   Arimoto, K.; Fukuda, H.; Imajoh-Ohmi, S.; Saito, H.; Takekawa, M., Formation of stress
granules inhibits apoptosis by suppressing stress-responsive MAPK pathways. *Nat Cell Biol*
2008, *10* (11), 1324-32.
42.   Rao, P. N.; Reddy, K. S.; Bhuyan, A. K., Amyloid fibrillation of human Apaf-1 CARD.
Biochemistry **2009,** *48* (32), 7656-64.

43. Yasuda, S.; Tsuchiya, H.; Kaiho, A.; Guo, Q.; Ikeuchi, K.; Endo, A.; Arai, N.; Ohtake, F.;
Murata, S.; Inada, T.; Baumeister, W.; Fernandez-Busnadiego, R.; Tanaka, K.; Saeki, Y., Stress-
and ubiquitylation-dependent phase separation of the proteasome. *Nature* **2020,** *578* (7794),
296-300.
44. Gupta, I.; Singh, K.; Varshney, N. K.; Khan, S., Delineating Crosstalk Mechanisms of the
Ubiquitin Proteasome System That Regulate Apoptosis. *Front Cell Dev Biol* **2018,** *6*, 11.
45. Theillet, F. X.; Binolfi, A.; Frembgen-Kesner, T.; Hingorani, K.; Sarkar, M.; Kyne, C.; Li, C.;
Crowley, P. B.; Gierasch, L.; Pielak, G. J.; Elcock, A. H.; Gershenson, A.; Selenko, P.,
Physicochemical properties of cells and their effects on intrinsically disordered proteins (IDPs).
Chem Rev **2014,** *114* (13), 6661-714.
46. Lalli, G.; Schott, J. M.; Hardy, J.; De Strooper, B., Aducanumab: a new phase in
therapeutic development for Alzheimer's disease? *EMBO Mol Med* **2021,** *13* (8), e14781.
47. Hayashi, H.; Campenot, R. B.; Vance, D. E.; Vance, J. E., Protection of neurons from
apoptosis by apolipoprotein E-containing lipoproteins does not require lipoprotein uptake and
involves activation of phospholipase Cgamma1 and inhibition of calcineurin. *J Biol Chem* **2009,**
284 (43), 29605-13.
48. Schellenberg, G. D.; D'Souza, I.; Poorkaj, P., The genetics of Alzheimer's disease. *Curr*
Psychiatry Rep **2000,** *2* (2), 158-64.
684