Partitioning of ribonucleoprotein complexes from the cellular actin cortex

C. Isaac Angert[1], Siddarth Reddy Karuka[1], Louis M. Mansky[2,3,4], Joachim D. Mueller[1,2,5]

School of Physics and Astronomy[1], University of Minnesota, Minneapolis, MN 55455
Institute of Molecular Virology[2], University of Minnesota, Minneapolis, MN 55455
Division of Basic Sciences, School of Dentistry[3], University of Minnesota, Minneapolis, MN 55455
Masonic Cancer Center[4], University of Minnesota, Minneapolis, MN 55455
Department of Biomedical Engineering[5], University of Minnesota, Minneapolis, MN 55455

Abstract

The cell cortex plays a crucial role in cell mechanics, signaling, and development. However, little is known about the influence of the cortical meshwork on the spatial distribution of cytoplasmic biomolecules. Here, we describe a new fluorescence microscopy method to infer the intracellular distribution of labeled biomolecules with sub-resolution accuracy. Unexpectedly, we find that RNA-binding proteins are partially excluded from the cytoplasmic volume adjacent to the plasma membrane that corresponds to the actin cortex. Complementary diffusion measurements of RNA-protein complexes suggest that a rudimentary model based on excluded volume interactions can explain this partitioning effect. Our results suggest the actin cortex meshwork may play a role in regulating the biomolecular content of the volume immediately adjacent to the plasma membrane.

Teaser

A novel microscopy technique reveals partial exclusion of RNA-protein complexes from the actin cortex due to their large hydrodynamic size.

Introduction

Both the actin cortex and RNA are fundamental building blocks of eukaryotic cells. The cortex, acting in concert with other cytoskeletal elements, is involved in nearly all aspects of cell mechanics, such as cell structure, membrane rigidity, migration and division (1). Due to its location adjacent to the plasma membrane, the cortex plays important roles in signal transduction and mechanical response pathways (1, 2). In contrast, RNA molecules serve as genomic messengers and constitute major components of the cellular translational machinery. Specific localization of mRNA to sub-cellular regions has been widely observed (3). RNA localization to sub-cellular compartments such as the endoplasmic reticulum (ER),

processing (P)-bodies or distal cytoplasm in neurons is critical to aspects of development (4) and is an

important regulator of cellular function (5). Due to their highly structured and extended nature, mRNAs have

hydrodynamic radii that are dramatically larger than those of the globular proteins they encode (6). Based

on a recent study (7), a typical mRNA of $1 - 2$ kilobase length (8, 9) is expected to have a hydrodynamic

radius in the range of ~10 nm. The radii of ribosomes are of comparable size $(10 - 15$ nm) (10, 11).

Previous biophysical studies have established that the cell cytoplasm is a complex medium (12, 13) that

modulates the diffusion and localization of large (>10 nm) particles in a size-dependent manner (14–18).

These observations have prompted the question of whether the large size of the cellular translational

machinery might broadly influence its subcellular localization (19). Here, the novel observation that large

riboneucleoprotein (RNP) complexes are partially excluded, or partitioned, from the interior of the actin cortex

is reported. These observations are facilitated by recent developments in two-photon dual-color (DC) z-scan

fluorescence microscopy (20), which is adapted in the present work to infer protein localizations inside living

cells with sub-diffraction resolution. DC z-scans are acquired by moving the two-photon excitation volume

vertically through the cytoplasm of adherent cells. This movement of the excitation volume yields fluorescent

intensity traces that encode the vertical intracellular distribution of a fluorescently labeled protein species as

well as that of a distinctly colored reference fluorescent protein. Although different reference species are

possible, our typical choice is a soluble, purely cytoplasmic protein, which serves as a "paint" that marks the

cytoplasmic extent. Differential vertical localization between the fluorescently labeled protein and the

reference protein is encoded as subtle differences in the shape of the DC z-scan traces. The magnitude of

the differential vertical localization is quantitatively recovered from the data by fitting the z-scan traces (20).

Results

Initial proof-of-principle experiments were conducted with U2OS cells to demonstrate the utility of DC z-scan

to quantify differences in intracellular protein distributions. For detailed descriptions of the z-scan model

geometries used in the following, see the Materials and Methods section. First, cells were co-transfected

with mCherry-RXR (a protein that localizes in the nucleus) and EGFP (which diffuses freely throughout the

cell). DC z-scans through the cell nucleus were fit to the S^G-S^{CH} (Eq. 17) model to identify the slab length

associated with the mCherry fluorescence, which corresponds to the nuclear thickness, and the slab length

of the EGFP signal, which corresponds to the slightly longer thickness of the entire cell (Fig. 1A). A

representative z-scan with fit shows that the mCherry-RXR and EGFP z-scans are well described by this

model, with flat residuals and $\chi_v^2 = 1.2$ (Fig. 1B). The fit identified the intensity amplitudes of mCherry-RXR

and EGFP, the locations of the bottom and top cell edges ($a^G = 2.16$ µm and $b^G = 5.80$ µm), as well as the

locations of the bottom and top nuclear edges ($a^{CH} = 2.51$ µm and $b^{CH} = 5.03$ µm), respectively. The

distances between the edges of the EGFP and mCherry slab were defined at the top and bottom of the cell

by $\Delta^T = b^{CH} - b^G = $ -770 nm and $\Delta^B = a^G - a^{CH} = $ -350 nm, respectively. Since Δ measures the displacement

between the borders of the EGFP and mCherry layers, its value can be positive or negative in general.

Distances are reported by taking the absolute value $\left|\Delta\right|$, which in this instance determines the thickness of

the cytoplasmic layers surrounding the top and bottom of the nucleus (Fig. 1A). Repeated DC z-scan

experiments on multiple cells yielded multiple estimates of Δ^T and Δ^B which are summarized by a histogram

(Fig. 1C) with averages of $<\Delta^T> = $ -340 ± 50 nm and $<\Delta^B> = $ -260 ± 30 nm (SEM, $n = 19$).

As a control, DC z-scans were performed through the cytoplasm of U2OS cells co-expressing EGFP

and mCherry fluorescent proteins (Fig. 1D). Since both proteins are soluble species within the cell cytoplasm,

DC z-scans are expected to recover $\Delta = 0$, on average. As shown in Fig. 1E, a representative DC z-scan

fits to a S^G-S^{CH} (Eq. 17) model with flat residuals and $\chi_v^2 = 1.1$. Repeated DC z-scans in multiple cells

recovered $<\Delta^B> = $ -2.3 ± 1.2 nm and $<\Delta^T> = 1.0 ± 3.1$ nm (SEM, $n = 59$). The histogram of the combined

displacements is shown in Fig. 1F. For the EGFP and mCherry co-expression, the weighted average of the

combined data yielded $<\Delta> = $ -0.5 ± 1.7 nm (SEM, $n = 118$), which is consistent with $\Delta = 0$ nm within 0.3 SD.

Note that since the diameter of globular proteins is typically several nanometer, differences of less than

10 nm were considered insignificant in this study and treated as consistent with zero.

Next, DC z-scans were performed in U2OS cells co-expressing EGFP-HRas and mCherry (Fig. 1G).

EGFP-HRas was previously reported to be a peripheral membrane protein that binds to the plasma

membrane as a monomer ([21]). Previous z-scan studies established that the axial distribution of EGFP-HRas

in the cytoplasm and at the PM membranes was successfully modeled by the δSδ layer geometry ([21, 22]).

Thus, DC z-scans from cells co-expressing mCherry and EGFP-HRas were modeled by the $(S^{CH}\text{-}\delta S\delta)^G$

geometry (Eq. 17). A fit of a representative z-scan attained good quality of fit with $\chi_v^2 = 1.2$ and flat residuals.

Repeated z-scans in the cytoplasm of multiple cells co-expressing EGFP-HRas and mCherry (Fig. 1I)

determined $<\Delta> = $ -2.6 ± 5.4 nm (SEM, $n = 30$), which was within 0.5 SD of the expected value of $\Delta = 0$. This

result indicated that the location of the cytoplasm slab edge, as determined by fitting the soluble mCherry

species to the S model, coincided with the location of the PM, as determined by fitting the EGFP-HRas

species to the δSδ model.

Next, the axial cytoplasmic distribution of APOBEC 3G (A3G) labeled with EGFP and co-expressed

with mCherry (Fig. 1J) was examined. A representative z-scan acquired in the cytoplasm of a HeLa cell

showed a single broad peak for both mCherry and A3G-EGFP (Fig. 1K). Since the trace shape was

suggestive of a slab geometry, both species were fit to slab models (S^{CH}-S^G, Eq. 17). This fit provided a

good description of the DC z-scan intensity profiles with $\chi_v^2 = 1.3$ and flat residuals (Fig. 1K). Interestingly,

this fit identified non-zero values for the displacements at the top and bottom PM ($\Delta^B = 184$ nm and $\Delta^T = 167$

nm). Repeated z-scan acquisitions at the same location in this cell determined $\overline{\Delta^B} = 180 \pm 6$ nm and $\overline{\Delta^T} =$

162 ± 12 nm (SEM, $n = 12$). This surprising observation suggests that the thickness of the soluble A3G-

EGFP layer is less than the thickness of the mCherry layer, unlike the results obtained for EGFP and

mCherry (Fig. 1D), which had equal lengths ($\Delta = 0$) for both soluble proteins. Repeated z-scan acquisitions

in multiple HeLa cells (Fig. 1L) found $<\Delta>_{HeLa} = 155 \pm 10$ nm (SEM, $n = 64$), indicating that A3G does not

occupy the entire cytoplasmic volume. To further validate this observation, DC z-scan measurements were

collected using a different cell line (U2OS), which resulted in $<\Delta>_{U2OS} = 110 \pm 9$ nm (SEM, $n = 32$).

Intriguingly, this result implies that cytoplasmic A3G-EGFP was excluded from regions adjacent to

the PM with apparent thickness of 100 - 200 nm. Given the magnitude of this value and its differential value

in U2OS and HeLa cell lines, further experiments were conducted to test whether A3G-EGFP might be

partially excluded from the dense meshwork of the actin cortex. To do this, HeLa cells were treated with

ethanol (see Materials and Methods), which disrupts actin assemblies and leads to the formation of large

plasma membrane blebs (23) (Fig. 2A) that lack an intact actin cortex (24–26). A representative z-scan

performed near the center of these blebs showed good quality of fit to a S^{CH}-S^G model (Eq. 17 and Fig. 2C).

Repeated z-scans in blebs from HeLa cells co-expressing A3G-EGFP and mCherry determined $<\Delta> = 31 \pm$

10 nm (SEM, $n = 14$), which differs from the results obtained in untreated HeLa cells by 8.8 SD. The

distribution of Δ in blebs and in untreated cells are markedly different (Fig. 2D). The strong reduction in Δ

found in blebs supports the conclusion that A3G-EGFP concentration is depleted from the vicinity of the PM

by the presence of the actin cortex. To reflect these results Δ is also referred to as the effective depletion

length.

Given that dependence of cortical architecture on the cell type has been observed (27–29), the

differences in effective depletion length observed between HeLa and U2OS cells might reflect natural

differences in cortex thickness between these cell types. To investigate this, direct measurements of the

cortex thickness were performed using distinct fluorescent colors to mark the actin cortex and the plasma

membrane. A previous study used this approach to estimate the actin cortex thickness from the displacement

between intensity peaks corresponding to the actin and the PM marker in confocal images (*30*). In the

present study two-photon DC z-scan was employed instead of confocal imaging to determine the cortex

thickness from the intensity profiles of EGFP-HRas (which binds to the PM) and Lifeact-mApple (which binds

to actin and marks the cortex) (*31*) by fitting to on a quantitative model of the entire intensity traces (Fig. 3A).

The axial concentration profile of both labels is depicted in Fig. 3B, which shows an enrichment of EGFP-

HRas and Lifeact-mApple at the PM and within the cortex, respectively. A small amount of both fluorescent

proteins is also present in the cytoplasm. Based on this profile, EGFP-HRas is again modeled by the δSδ

geometry. The distance separating the two HRas δ layers represents the cell thickness L_{cell}.

The concentration profile of Lifeact-mApple implies a wide cytoplasmic slab of length L_{cyto}

separating two thin slabs representing the cortex at the bottom and the top membrane with thickness L_C^B

and L_C^T (Fig. 3B), giving rise to an SSS^{CH} (triple S-layer) geometry. Choosing a uniform slab with a sharp

transition between the cortical and cytoplasmic spaces to model the cortex has been previously used in (*30*)

where it is estimated that even a substantial non-uniformity of the cortical density leads to only a small

change in the estimated cortex thickness. Because the expected slab length L_C of the cortex is smaller than

500 nm, its length cannot be directly identified in single-color (SC) z-scan experiments. Such thin slabs are

experimentally indistinguishable from a δ layer (*32*). However, the centroid position of the thin cortical layer

is expected to coincide with the midpoint of the slab as depicted in Fig. 3B. Given this, modeling the Lifeact-

mApple species as a δSδ^{CH} geometry can identify the cortex centroid from the fitted location of each δ layer.

Combined with the EGFP-HRas species, this DC z-scan fit (δSδ^{CH}-δSδ^G) identifies the locations of the δ

layers corresponding to the cortex centroid and the PM as well as their displacement *Δ*. The distance |*Δ*|

separating both δ layers is half of the actin cortex thickness, |*Δ*| = L_C / 2 (Fig. 3B).

The relation |*Δ*| = L_C / 2 was verified by computer modeling of the DC z-scan intensity profile of the

SSS^{CH}-δSδ^G geometry (Fig. 3B), while systematically varying the cortical thickness $L_{C,in}$, as well as the cell

thickness and fluorescence intensities of HRas and Lifeact (see Materials and Methods). These simulated

scans were fit to a δSδ^{CH}- δSδ^G model to determine the locations of each δ layer. The difference *Δ* between

the fitted cortical δ layer and the PM δ layer was computed for each fit and the averaged *Δ* recovered from

fits to simulated DC z-scans were plotted as a function of the cortical thickness $L_{C,in}$ (Fig. S1). The data

confirm that the relation $|\Delta| = L_C / 2$ is accurate to within 10 nm for cortex thicknesses up to 200 nm. This

result demonstrates that this approach is experimentally accurate and produces negligible error.

Repeated DC z-scans were then collected in cells co-expressing EGFP-HRas and Lifeact-mApple

proteins in order to determine cortical thickness. A representative DC z-scan and fit to the $\delta S \delta^{CH}$- $\delta S \delta^{G}$

model (Fig. 3C) for both proteins is shown in Fig. 3D. Data taken from multiple cells resulted in absolute

values $|<\Delta>|_{U2OS} = 63 \pm 8$ nm (SEM, $n = 7$) and $|<\Delta>|_{HeLa} = 100 \pm 13$ nm (SEM, $n = 15$). These values

correspond to uniform cortex thicknesses of 126 ± 16 nm for U2OS cells and 200 ± 24 nm for HeLa cells

(Fig. 3E).

Since both the depletion length Δ for A3G-EGFP and the cortex thickness differed between the two

cell lines, $<\Delta>$ in HeLa and U2OS cells were plotted for comparison (Fig. 3F). DC z-scans were performed

in HeLa and U2OS cells expressing additional proteins, HIV-1 Gag^{G2A}-EGFP, and its matrix (MA) domain,

MA^{G2A}-EGFP (*33*) with mCherry again as a reference marker of the cytoplasmic volume for each instance.

A cell-line dependent $<\Delta>$ was observed for both proteins (Fig. S2), and their $<\Delta>$ values were added to Fig.

3F. Interestingly, all 3 data points cluster near a line of slope 0.66, indicating that, although the magnitude

of the depletion effect varies for each protein, it is systematically scaled by ~66% between these two cell

lines. This same ratio corresponds well to the cortex thickness ratio estimated by DC z-scan between HeLa

and U2OS cells (0.66 ± 0.11, shaded area in Fig. 3F), suggesting that the cell-type dependent differences

in $<\Delta>$ are due to differences in their cortical actin thicknesses. This result further implicates the actin cortex

as a central driver of the observed depletion effect near the PM.

All 3 proteins in Fig. 3F are known to be RNA-interacting proteins: HIV-1 Gag^{G2A} binds specifically

and non-specifically to RNA sequences through basic residues and two zinc fingers in its nucleocapsid (NC)

domain (*34–36*), while the MA domain of Gag separately interacts with RNA via a stretch of basic residues

in its highly basic region (HBR) (*37–39*) and has preferential tRNA binding affinity (*40, 41*). A3G interacts

with RNA primarily through its N-terminal domain zinc finger (*42, 43*). A3G-RNA binding is required for

packaging of A3G into retroviral particles (*43–46*) and is critical to its antiretroviral activity (*43, 44*). To test

whether the depletion effect is specific to RNA-binding proteins, measurements with two EGFP-labeled

proteins that lack a known RNA binding function were conducted. In particular, DC z-scans of HeLa cells co-

expressing mCherry with the GABA-A receptor associated protein fused to EGFP (GABARAP-EGFP), and

mCherry with a tandem hexamer of the Venus fluorescent protein, Venus$_6$ were collected. The average $<\Delta>$

value of both proteins was found to be < 10 nm (Fig. 4A) and therefore treated as consistent with zero. The

previously presented depletion lengths in HeLa cells from Fig. 3D were added to Fig. 4A to highlight the

contrast with the RNA-binding proteins. Since RNA binding appeared to be important for observing depletion,

DC z-scans were performed with the mutant A3G[C97A]-EGFP, which has impaired RNA binding due to

mutation of the N-terminal zinc finger (*44*). These measurements revealed a large reduction in $<\Delta>$

compared to that of wildtype A3G-EGFP (Fig. 4A). Lastly, the $<\Delta>$ value was measured in mCherry

expressing HeLa cells that were stained with the small molecule RNA binding dye, SYTO RNASelect. These

measurements revealed that the labeled cellular RNA had a significant degree of depletion adjacent to the

PM (Fig. 4A).

Based on these results, it was hypothesized that the exclusion of large RNA species from the dense

cortical meshwork may be responsible for the observed PM adjacent depletion of RNA binding proteins (Fig.

4B). The average pore diameter of the actin cortex has been reported to range between 50 to 200 nm (*28*),

depending on cell type. These values are only slightly larger than or overlap with the size of large RNAs (*6*,

7). Thus, a partial partitioning of RNAs from the interior of the cortex could be driven by excluded volume

interactions. In a simple model of the cortical partitioning (Fig. 4C), the ratio of the concentration c_C in the

cortical layer to the concentration c_{cyto} in the cytoplasm defines the partition coefficient $R_C = c_C / c_{cyto}$. This

partitioning of the protein concentration between the cytoplasmic and cortical spaces introduces a SSS[G]

concentration profile for the EGFP-labeled RNA-binding protein (Fig. 4C), which consists of two thin cortical

slabs of length L_C^B and L_C^T separated by a thick cytoplasmic slab of length L_{cyto} with a higher protein

concentration. As in the case of the Lifeact measurements (Fig. 3), the cortical layer is too narrow to be

directly resolved by z-scan analysis. However, since the concentration within the cortex layer is reduced

(Fig. 4C), the z-scan intensity profile can be effectively approximated by a single slab layer with a width that

slightly exceeds the cytoplasmic thickness L_{cyto} (Fig. 4D). Consequently, the apparent depletion lengths Δ^B

and Δ^T will be less than the corresponding cortex thickness L_C^B and L_C^T in a manner that depends on R_C.

To quantitatively analyze how Δ recovered from DC z-scans is related to R_C and L_C, simulation of

DC z-scan traces for an SSS[G]-S[CH] geometry was performed (see Materials and Methods). These simulated

traces were fit to the S[CH]-S[G] DC z-scan model (Fig. 4D). The cortex length L_C and partition ratio R_C (Fig. 4C)

were varied for each simulation and the averaged Δ recovered from the fits was plotted (Fig. 4E, diamonds).

The results of these simulations indicate a nearly linear relationship between Δ and R_C for a given L_C. This

linear relationship can also be derived from preserving the integrated area Λ under the z-scan intensity

profile between the SSS^G model and the fitted S^G model. Since Λ and the axially integrated concentration A

are directly proportional (Eq. 11), the concentration areas of both models (Fig. 4C and D) must match. In

particular, the cortical concentration area $A_C^i = L_C^i \cdot c_C$ (Fig. 4C) has to be equal to the added extra area

$A_x^i = L_C^i \cdot R_C c_{cyto}$ (Fig. 4D) at both the top (i = T) and bottom (i = B) of the cell, which predicts an apparent

depletion length Δ (Fig. 4D) of

$$\Delta = \left(1 - R_C\right)L_C \,, \qquad\qquad 1$$

where the superscript has been omitted. This model (Fig. 4E, solid lines) is in excellent agreement with the

results from the simulations. Eq. 1 was used to estimate the partition coefficient of proteins from their

measured depletion lengths (Fig. 3D) using the experimentally determined cortex thicknesses L_C. R_C = 0.76

± 0.02, 0.35 ± 0.04, and 0.24 ± 0.05 for MA^{G2A}-EGFP, Gag^{G2A}-EGFP, and A3G-EGFP, respectively were

determined from HeLa cells. Similar values (0.76 ± 0.02, 0.41 ± 0.02, and 0.17 ± 0.07) were obtained from

the measurements in U2OS cells.

Because these estimates of the partition coefficient R_C rely on the measured displacement Δ, which

is a new method, a second approach was developed to estimate R_C independently of Δ by using DC z-scans

in very thin cytoplasmic regions. The partial exclusion of fluorescently labeled protein from the cortex reduces

the fluorescence area Λ^G as compared to the area Λ_{ref}^G corresponding to the hypothetical case of no

exclusion. The ratio of these two areas is denoted as the axially averaged partition coefficient,

$$\bar{R}_z = \frac{\Lambda^G}{\Lambda_{ref}^G} = \frac{A^G}{A_{ref}^G} = \frac{L_{cyto}c_{cyto} + 2L_C R_C c_{cyto}}{L_{cell}c_{cyto}} \,, \qquad\qquad 2$$

230 where the proportionality between Λ and A (Eq. 11) was used to express the ratio in terms of the integrated

231 concentration areas. Since the cell thickness is defined by (Fig. 3B),

$$L_{cell} = L_{cyto} + L_C^T + L_C^B = L_{cyto} + 2L_C \,, \qquad\qquad 3$$

233 where $L_C = (L_C^T + L_C^B)/2$ defines the average cortex thickness, the axially averaged partition coefficient

234 simplifies to

$$\bar{R}_z = 1 + (R_C - 1)\frac{2L_C}{L_{cell}} \qquad\qquad 4$$

This equation predicts $\overline{R}_z \rightarrow 1$ for very thick sections ($L_{cyto} \gg L_C$). As the cell thickness L_{cell} decreases,

the value of $\overline{R}_z$ drops. It reaches a limiting value of $\overline{R}_z = R_C$ for very thin cell section composed entirely of

cortex $L_{cell} = 2L_C$. Thus, DC z-scans in thin regions near the cell perimeter combined with DC z-scans in

thick regions of the same cells allow direct estimation of the partition coefficient.

While this relatively simple model sets a minimum cell thickness of $L_{cell} = 2L_C$, where L_C denotes

the average cortical thickness in a given cell type, very thin cellular regions were found where the realized

cell thickness is less than this value, $L_{cell} < 2L_C$. This implies that in these regions of the cell the local cortex

thickness is less than L_C. Assuming that these thin sections are composed entirely of cortex, Eq. 4 is

generalized to,

$$\overline{R}_z = \begin{cases} R_C & \text{if} \quad L_{cell} < 2L_C \\ 1 + (R_C - 1)\dfrac{2L_C}{L_{cell}} & \text{if} \quad L_{cell} > 2L_C \end{cases} .$$

245   5

As shown in Fig. 5A & B, cytoplasmic DC z-scans in thick and thin regions of a cell co-expressing

EGFP and mCherry indicate that the relative concentrations of the EGFP and mCherry species are identical

regardless of cytoplasm thickness. In contrast, cytoplasmic DC z-scans in thick and thin regions of a cell co-

expressing HIV-1 Gag^{G2A}-EGFP and mCherry show that the relative intensity of Gag^{G2A}-EGFP is sharply

reduced in thin regions (Fig. 5C & D), demonstrating that the depletion effect is reflected in the decrease of

the integrated fluorescence area Λ^G as a function of thickness. To quantify the effect and utilize Eq. 2, an

estimate was needed for the reference area $\Lambda_{ref}^G = L_{cell} g_{cyto}^G$, where $g_{cyto}^G = c_{cyto}^G \eta$ (Eqs. 11 and 2). The cell

thickness L_{cell} was determined from the mCherry signal in the red detection channel as described in the

Material and Methods, while the amplitude g_{cyto}^G was estimated from a scan through a thick cell section (L_{cell}

> 1.5 µm) where g_{cyto}^G can be reliably extracted from fitting the DC z-scan traces. Repeated z-scans were

performed within the same cell covering thick and thin regions (Fig. 5E) to graph $\overline{R}_z = \Lambda^G / \Lambda_{ref}^G$ versus L_{cell}.

As expected, $\overline{R}_z$ is approximately one for a HeLa cell coexpressing mCherry and EGFP (Fig. 5F), which

implies the absence of partitioning of EGFP from the cortex according to Eq. 5 ($R_C = 1$). In contrast, DC z-

scan estimates of $\overline{R}_z$ for Gag^{G2A}-EGFP for a HeLa cell (magenta diamonds) and a U2OS cell (blue

diamonds) exhibit a decrease at thin cell sections (Fig. 5F). The HeLa cell data in Fig. 5F closely resemble

the simple model (Eq. 5), and cluster near 0.2 for $L_{cell} < L_C$, suggesting a cortical partition coefficient near

0.2 in this cell line. The data from U2OS cells appears to deviate from (Eq. 5) in thin cell sections, which

may reflect the presence of distinct peripheral F-actin structures in this cell line as discussed later.

The cortical partitioning effect documented here supports a model where protein depletion within

the cortex is driven by protein association into ribonucleoprotein complexes that are much larger than the

typical protein. In such a case, the large <Δ> values should be correlated with substantially slowed

cytoplasmic diffusion. FCS measurements were performed in HeLa cells in order to estimate the effective

diffusion coefficient D_{eff} of labeled proteins in the cytoplasm (see Materials and Methods). In Fig. 6A <Δ>

values from Fig. 4 were plotted versus the median D_{eff} that was obtained from FCS measurements. Two

other members of the APOBEC3 protein family, A3A-EGFP and A3C-EGFP, which have differing

antiretroviral activities (*47–49*), were also measured and added to Fig. 6A. These data show that large <Δ>

values are associated with remarkably low cytoplasmic mobility. This relationship is summarized by the

cartoons in Fig. 6 B&C. In order to model the data in Fig. 6A, the cortical partitioning was described as arising

from excluded volume interactions of particles that are allowed to enter spherical voids (*50–52*),

275
$$R_C = (1 - r_p / r_v)^3 \, , \qquad\qquad\qquad 6$$

276   where r_p is the particle radius, and r_v is the void radius. A hindered cytoplasmic diffusion coefficient D_{eff}

277   was modeled as arising from a size-dependent viscosity imposed by macromolecular crowding within the

278   cytoplasm (*53, 54*) according to

279
$$\frac{D_{eff}}{D_0} = \exp\left[-\left(\frac{r_{eff}}{\xi} \right)^a \right] , \qquad\qquad\qquad 7$$

280   where D_0 is a reference diffusion coefficient, r_{eff} is an effective hydrodynamic radius of the diffusing

281   particles, a is a scaling exponent of order 1, and ξ is a length interpreted as the characteristic distance

282   between points of entanglement within the crowded solution. Along with equation 1, setting $r_{eff} = r_p$ in the

283   above yields a relation between D_{eff} and Δ,

284
$$\Delta = L_c \left[1 - \left(1 - \frac{\xi}{r_v} \ln \left(\frac{D_0}{D_{eff}} \right)^{1/a} \right)^3 \right].$$
8

This curve is plotted as the magenta line in Fig. 6A using values $\underline{L_C}$ = 170 nm, D_0 equal to D_{eff}

measured for EGFP (27.5 $\mu m^2 s^{-1}$), r_v = 60 nm, ξ = 4 nm and a = 0.7. The values for ξ and a are within the

range estimated by (*53*) for HeLa cells. r_v is similar to published estimates of the cortical mesh size (*28*).

Cytoplasmic crowding in cell associated PM blebs may differ from crowding within undisturbed

cytoplasm due at least in part to the lack of a cytoskeletal mesh in the bleb interior. Thus, the mobility of

A3G-EGFP in HeLa cell blebs was investigated. FCS measurements (Fig. 6A, open circle) showed an order

of magnitude increase in D_{eff} compared to A3G in the cytoplasm of untreated HeLa cells. This result

supports our choice to model cytoplasmic diffusion of RNP complexes as arising from a size-dependent

viscosity (Eq. 717). The value from blebs is plotted as an open circle in order to emphasize that the model

parameters (especially r_v and ξ) that describe the rest of the data most likely do not describe data from blebs.

It was recently reported that the cytoplasmic diffusion coefficient of fluorescent proteins depended

on their net charge, with positively charged proteins diffusing slowly. It was suggested that this slow diffusion

was due to transient interactions with large, negatively charged cell components such as RNA and the

cytoskeleton (*55*). An examination was performed to determine whether the D_{eff} values in Fig. 6A might be

explained by the net charge of each protein-FP fusion. As shown in Fig. S3, the mobility of the proteins in

Fig. 6A was only weakly correlated with net charge. Thus, the mobility of these proteins in living cells is likely

to be better understood as arising predominantly from their RNA binding function.

Discussion

In this study, DC z-scan microscopy measurements of the fluorescence intensity profiles of green and red

labeled proteins provided the foundation for identifying the cortical partitioning effect and allowed its

quantitative description through the effective depletion length Δ. Implementation of the DC z-scan

measurements using two-photon microscopy eliminates systematic biases, such as chromatic aberration,

from the collected data. DC z-scan permitted precise quantification of the relative localizations of

fluorescently labeled proteins. These measurements indicated an apparent restriction in the cytoplasmic

volume of RNA binding proteins (e.g., A3G and Gag[G2A]). The partitioning effect was markedly reduced in

ethanol induced cell blebs, which largely lack a cortex (*24–26*) and was observed to be cell-type dependent

in a manner that correlated with DC z-scan estimates of cortex thickness in adherent interphase HeLa and

U2OS cells. The depletion effect is interpreted as being a partitioning, or partial exclusion, of these proteins

from the volume of the actin cortex. Furthermore, Δ values measured for proteins with and without RNA-

binding function suggest that the cortical partitioning is driven by association into RNP complexes and is not

strongly associated with protein molecular mass or net charge. This conclusion is supported by

measurements of the diffusion coefficient, which shows that a marked decrease in mobility is associated

with proteins that have the largest values of Δ.

In vitro and in vivo (*56*) observations of actin assemblies suggest that gel-like properties are nearly

ubiquitous, including for the actin cortex specifically (*57*). The cortical partitioning effect may be analogous

to solute partitioning from the interior of gels, a well-known chemical phenomenon (*58*, *59*). Furthermore,

the subcellular heterogeneity driven by the dense cortical environment is reminiscent of those molecular

crowding effects which have been reported to drive association or phase separations in cells (*60*, *61*). This

suggests that the crowded environment within gels may drive biophysical phenomena similar to, but distinct

from those associated with molecular crowding in the cytoplasm. The observation of cortical partitioning

described in this report builds on earlier work that found strong exclusion of labeled colloidal particles with

radii exceeding ~10 nm from distal regions of mammalian cell cytoplasm, and regions near the centrosome

(*15*, *62*). Subsequent studies documented that the distal excluding regions were associated with a dense

cytoskeletal mesh and were thinner than non-excluding regions (*62*). These and other observations (*50*)

have suggested that this partitioning was driven by the cytoskeletal meshwork.

The cortical partitioning of RNPs described in the present study also leads to strong exclusion from

distal regions of the cytoplasm (Fig. 5). However, unlike previous studies, which linked partitioning to the

actin cytoskeleton in general, our observations identified the actin cortex as the causative agent of

partitioning. Differentiation between the cortical space and the bulk cytoplasm was not previously achieved

and probably reflects experimental limitations in these earlier studies, which are overcome by the use of DC

z-scan method employed here. Thus, the cortical partitioning of RNPs and the distal exclusion of

microinjected particles documented previously (*15*, *63*) likely are manifestations of the same physical

phenomenon, namely the size-dependent partitioning of particles from the actin cortex.

The measurement of the actin cortex thickness, L_C, in HeLa (~200 nm) and U2OS (~130 nm) cells

in our study represents a novel application of the DC z-scan technique. This application was inspired by a

conceptually similar approach based on confocal microscopy that was used to estimate a cortex thickness

of ~190 nm in mitotic HeLa cells (*30*) and of 300 – 400 nm in interphase HeLa cells with imposed spherical

shapes (*29*). Interestingly, DC z-scan identified a cortical thickness ≤200 nm for adherent interphase HeLa

and U2OS. This is consistent with literature reports which indicate that cortical architecture is strongly

associated with cell morphology (*27, 29*). In particular, cell adhesion and cell spreading on the glass coverslip

can result in a thinning of the actin cortex (*27, 64*).

DC z-scan measurement of the axially averaged partition coefficient $\overline{R}_z$ (Fig. 5E) provided further

evidence that partitioning is associated with the cellular cortex as evidenced by the decreasing $\overline{R}_z$ values

in increasingly thinner cell sections. The value of $\overline{R}_z$ in very thin cell sections provides a direct estimate of

the cortical partition coefficient R_C, assuming that these thin sections are composed entirely of the actin

cortical mesh. This R_C value can in turn provide an additional estimate of cortex thickness that does not rely

on DC z-scans of cortical Lifeact. Utilizing Eq. 1, the R_C ~0.2 measured for HIV-1 Gag^{G2A}-EGFP in thin

sections of HeLa cells would suggest L_C ~170 nm given <Δ> = 135 ± 8 nm observed for HIV-1 Gag^{G2A}-EGFP

in thick cytoplasmic DC z-scans in this same cell line. This value of L_C is within 1.3 SD of the value estimated

from DC z-scans of Lifeact. The agreement between these two different DC z-scan approaches provides

further validation of the DC z-scan technique and demonstrates its utility for interrogating biophysical

properties of the bulk actin cortex.

Interestingly, DC z-scan estimates of $\overline{R}_z$ in U2OS cells exhibited more scatter than observed in

HeLa cells, particularly in thin cell sections. This may reflect cell-type differences in F-actin assemblies, in

particular large structures such as lamellar actin sheets which may be present near the cell periphery. In

electron micrographs of F-actin, these lamellae appear distinct from cortical actin fibers (*27, 65*). Their

extensive presence may invalidate the use of Eq. 5 near the periphery of U2OS cells. Future work is needed

to investigate whether distinct types of actin structures modulate the cortical partitioning and whether this

effect is cell-type and -function dependent.

The strong correlation between the ability of a protein to bind RNA and measured values of <Δ>

(Fig. 4A) suggests that many RNA-protein complexes are too large to efficiently enter the dense actin cortical

meshwork. The cortical partitioning of RNA-binding proteins measured by DC z-scan is thus interpreted as

arising from an underlying partitioning associated with the large RNA species with which these proteins

associate. In this framework, DC z-scans of RNA interacting proteins indirectly probe the spatial distribution

of cellular RNAs. More directly, when cellular RNA was labeled with an RNA-binding fluorescent dye, DC z-

scan measurements of $<\Delta>$ revealed that the RNA itself is partitioned from the cortex with R_C in the range

0.35 – 0.2 (given L_C 170 – 200 nm). This R_C value is interpreted as representing the average R_C of a broad

spectrum of differently sized cellular RNAs (i.e., mRNAs, tRNAs, etc) and RNP complexes (e.g., ribosomes)

with different degrees of cortical penetration. Similarly, the $<\Delta>$ values for fluorescently labeled proteins

likely represent the average R_C of a heterogenous mix of differently sized RNA complexes to which each

protein preferentially binds. Alternatively, it is possible that different $<\Delta>$ values represent differing levels of

RNA-bound and -unbound protein, where the unbound state is identified as the non-partitioned fraction.

However, the unbound protein population is expected to be negligible in many cases. This follows from the

high concentration of protein-RNA binding sites in cell cytoplasm (*66*, *67*), which is estimated to greatly

exceed both the protein-RNA affinities (likely tens of nanomolar for Gag and A3G binding to single strand

RNAs (*42*, *68*, *69*)) and the protein concentrations (~1 µM) in DC z-scan measurements. The potential for

protein-RNA binding equilibria or selective protein-RNA binding to influence the cortical partitioning effect

may be explored with further studies utilizing DC z-scan.

FCS measurements were acquired in order to characterize protein mobility in the cytoplasm. Given

that these data were acquired with the PSF focused near the middle of thick (>1.5 µm) cytoplasmic sections,

the experimental estimates mainly report the mobility within the cytoplasm and not the mobility within the

actin cortex. The diffusion time of RNA binding proteins revealed that large $<\Delta>$ values were associated with

very low mobility (Fig. 6A), as expected for proteins that are present in large complexes. Slow diffusion of

large sized objects in the cytoplasm has been previously reported (*14*, *17*, *70*, *71*), including mRNAs (*72*).

Previous studies have suggested that this low cytoplasmic mobility may reflect a hindrance created by

cytoskeletal elements (*15*, *16*, *18*, *19*, *62*, *73*, *74*) or by cytoplasmic crowding created due to ribosomes,

organelles and proteins (*17*, *70*, *75–77*). Although the precise determinants for cytoplasmic mobility have

not been fully characterized, existing literature provides extensive evidence that the cytoplasm cannot be

described as a simple fluid (*12*, *53*, *75*, *78*), and it is generally agreed that particles in the range of 10 –

100nm show a size-dependent drop in cytoplasmic mobility much greater than that predicted by the Stokes-

Einstein relation (*14*, *17*, *18*, *70*). In this study, a marked increase in A3G-EGFP mobility in cell-associated

blebs (Fig. 6A) was observed. This suggests a role for the cytoskeleton in the mobility of large

macromolecular complexes.

To capture the size dependence of RNP diffusion, an empirical hydrodynamic scaling model (Eq. 7

) was used. This and similar models have been previously employed to describe diffusion in polymer

solutions and in living cells (*53*, *54*). This model led to good agreement with our experimental data (Fig. 6A)

when combined with a relatively simple description of the size dependence of cortical partitioning (*50–52*)

(Eq. 6). This result demonstrates the feasibility of a quantitative description of the actin cortical partitioning

effect. The choice to model the partition coefficient as arising from the excluded volume due to spherical

cavity (Eq. 6) was driven primarily by the simplicity of this approach. The excluded volume due to a mesh of

randomly oriented fibers, which may provide a more realistic model of cortical actin, is expected to give

similar results to Eq. 6 (*50*) at the expense of an additional free parameter.

This study presents data suggesting that large RNP complexes are disproportionately partitioned

from the space immediately adjacent to the PM due to the presence of the dense cortical actin meshwork.

In this phenomenon the physical properties of the cortical space appear to drive subcellular segregation of

macromolecules in the absence of a clear boundary such as a membrane. Other instances of biomolecule

compartmentalization in the absence of membranes have been recognized, such as that observed between

P-bodies and the cytoplasm (*79*), and between nucleoli and the nucleoplasm (*80*). In such cases, the distinct

physical properties of each subcellular compartment regulate their molecular content (*81*). Partitioning of

large complexes from the cortical space is consistent with observations demonstrating that a highly

orchestrated coupling between exocytic pathways and cortical actin polymerization is needed to facilitate

passage of vesicles through the cortex when the actin fiber density is transiently lowered (*82*, *83*). Moreover,

the cortical partitioning effect might hinder the replication of enveloped viruses, which target large

macromolecular complexes to the PM as part of their assembly processes. In conclusion, new developments

in DC z-scan microscopy have enabled the observation and characterization of an unexpected cortical

partitioning effect that applies to large RNP complexes. A simple quantitative model was introduced to

provide a rudimentary foundation for understanding the origins of this effect. The DC z-scan toolkit should

be of interest to probe the spatial relation of biomolecules with other cellular structures as well as continue

to provide additional insights into the function of the actin cortex.

Materials and Methods

Cell Culture and Plasmids

HeLa and U2OS (Numbers CCL-2 and HTB-96, American Type Culture Collection, Manassas, VA) cells

were cultured in Dulbecco's modified Eagle's medium supplemented with 10% FBS (HyClone Laboratories,

Logan, UT) at 37°C and 5% CO_2. Cells were plated at ~30% confluency into 8-well chambered slides ~24 h

prior to measurement and transfected with vectors as indicated in the text ~16 h prior to measurement using

GenJet transfection reagent (SignaGen Laboratories, Fredrick, MD) according to the manufacturer's

instructions. The cell growth medium was replaced with phosphate buffered saline immediately prior to

measurement. All measurements were performed at room temperature.

HIV-1 Gag (*84*) was ligated into pEGFP-N1 (Clontech, Takara Bio, Mountain View, CA) to generate the HIV-

1 Gag-EGFP vector. HIV-1 Gag^{G2A}-EGFP was made using the QuikChange XL Site-Directed Mutagenesis

Kit (Stratagene, La Jolla, CA). Venus$_6$ was a gift from Steven Vogel (*85*) (Addgene plasmid # 27813;

http://n2t.net/addgene:27813; RRID: Addgene_27813). mApple-Lifeact-7 (denoted Lifeact-mApple here)

was a gift from Michael Davidson (Addgene plasmid # 54747; http://n2t.net/addgene:54747; RRID:

Addgene_54747). All other vectors have been previously described: A3G-EGFP, A3G^{C97A}-EGFP, A3C-

EGFP, A3A-EGFP (*44*); HIV-1 MA^{G2A}-EGFP (*20*); mCherry (*86*); EGFP-HRas (*21*); mCherry-RXR (*87*);

GABARAP-EGFP (*88*).

Instrumentation

DC z-scans and fluorescence fluctuation spectroscopy (FFS) data were acquired on a Zeiss Axiovert 200

microscope modified for 2-photon excitation (*20*). Excitation light was provided by a femtosecond pulsed

Ti:S laser (MaiTai or Tsunami models, Spectra Physics, Santa Clara, CA) operated at 1000 nm. This light

was focused to a diffraction limited point by a water immersion objective (C-Apochromat, Zeiss, Oberkochen,

Germany) with numerical aperture 1.2 and correction collar set according to the glass thickness of the

sample holder. Emission light was collected by the same objective and separated from the excitation light

by a dichroic filter (740DCSPXR, Chroma Technology, Bellows Falls, VT), then split into red and green

detection channels by a low-pass dichroic (FF580-FDi01, Semrock, Rochester, NY). A 515/50nm band pass

filter placed before the green channel detector eliminated Fresnel reflected mCherry emission light from the

green channel. Emission light in each channel was detected by single photon counting hybrid

photomultipliers (HPM-100-40, Becker and Hickl, Berlin, Germany), recorded by hardware (FastFLIM, ISS

Inc., Champaign, IL or DPC-230, Becker and Hickl, Berlin, Germany), and saved for later analysis using

routines written in IDL 8.6 (Harris Geospatial Solutions, Broomfield, CO).

Axial z-scans utilized a piezo stage (PZ2000, ASI, Eugene OR) driven by a voltage waveform from

an arbitrary waveform generator (model 33250A, Agilent Technologies, Santa Clara, CA) to move the sample

vertically across the two-photon point spread function (PSF). All z-scans were acquired by driving the piezo

stage with a triangle waveform with a 10 s period and peak-to-peak displacement corresponding to 24 μm.

Each period contained two z-scans: the first z-scan passed upward through the cell and the second passed

downward through the cell. Each z-scan covered a 24 μm range.

464   XY scanning of the PSF for acquisition of two-photon images was provided by a galvanometer scan

465   head (Yanus IV, FEI Deutschland GmbH, Planegg, Germany) aligned in the two-photon excitation path. XY

466   positioning was driven by a DAC card (3-axis, ISS Inc., Champaign, IL) controlled by SimFCS software

467   (version 3; G-SOFT Inc., Champaign, IL). All z-scans were acquired within 24 μm of the XY scan field center.

Z-scan modeling

470   The single-channel (SC) z-scan fluorescence intensity profile $F(z)$ is determined by the convolution of the

471   axial concentration profile $c(z)$ of the fluorescently-tagged protein and the radially integrated point spread

472   function (RIPSF) of the two-photon instrument as modeled in (*22*, *32*),

473
$$F(z) = \lambda (c * \mathrm{RIPSF})(z), \qquad\qquad\qquad 9$$

474   where λ is the brightness of the fluorescence tag. Evaluation of Eq. 9 for different concentration profiles

475    $c(z)$ gives rise to an alternative formulation with a fluorescence amplitude f and a function $\mathrm{v}(z)$

476   describing the shape of the concentration profile (*32*),

477
$$F(z) = f\,\mathrm{v}(z). \qquad\qquad\qquad 10$$

478   The axially integrated concentration $A = \int c(z)\,dz$ and the axially integrated fluorescence

479    $\Lambda = \int F(z)\,dz$ are related according to Eq. 9 by

480
$$\Lambda = \eta A \qquad\qquad\qquad 11$$

481   with the constant $\eta = \lambda \int \mathrm{RIPSF}(z)\,dz$.

Z-scan and FFS acquisition in live cells

Cells were selected by verifying the desired expression levels for the mCherry and EGFP labeled constructs

using 1-photon epi illumination provided by a mercury lamp and using an EGFP/mCherry filter set. Points

within cells for z-scan and FFS data collection were chosen where the fluorescence appeared locally (~3 μm

radius) uniform as visualized in epi illumination and where a cell lacked nearby internal structures when

visualized by using transmitted light. For data acquisition, the microscope was switched to two-photon mode

and the excitation volume was focused to the middle of the cell. In most instances, 60 s of z-scan data

acquisition was followed by an optional 60 s of FFS data acquisition at up to 4 unique locations within each

cell. Z-scan acquisition times at a given location were lengthened to a maximum of 10 min in dim samples

or where increased statistical precision was desired.

Effective diffusion coefficient

The effective diffusion coefficient was calculated as ([89]) $D_{\mathit{eff}} = \omega_0^2/(8\tau_D)$, where ω_0^2 is the $1/e^2$ radial

beam waist of the 2-photon excitation light focus and τ_D was estimated by fitting the log-averaged

autocorrelation of photon counts $G(\tau)$ to $G_{2D}(\tau) = G_0/(1+(\tau/\tau_D))^{-1} + G_{\mathit{offset}}$. FCS data quality control was

performed by redacting data where G_{offset} showed significantly negative values or where τ_D indicated by

the fit was >50% higher than the time at which the correlation amplitude reached half its initial value.

Modeling of DC z-scan intensity profiles

The DC fluorescence intensity $\mathbf{F^G}(z)$ of a z-scan intensity profile of EGFP (G)-labeled protein through a

series of horizontal layers, each containing a uniform concentration of labels, is described by ([20–22])

504
$$\mathbf{F^G}(z) = \sum_i \mathbf{f}_i^{\mathbf{G}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right) = \boldsymbol{\alpha}^{\mathbf{G}} \sum_i f_i^{\mathrm{G}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right), \qquad\qquad 12$$

where f_i^{G} and $\mathrm{v}\left(z, a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right)$ are the fluorescence amplitude and shape function associated with the *i*-th

layer. The shape function $\mathrm{v}\left(z, a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right)$ characterizes a uniform slab or S layer starting at $z = a_i^{\mathrm{G}}$ and

ending at $z = b_i^{\mathrm{G}}$, while $\mathrm{v}\left(z, a_i^{\mathrm{G}}, a_i^{\mathrm{G}}\right)$ defines an infinitesimally thin or δ layer located at $z = a_i^{\mathrm{G}}$ ([20–22]).

Vector notation is used to describe properties of the green (g) and red (r) channel, i.e.

$\mathbf{F^G}(z) = \left(F_g^{\mathrm{G}}(z), F_r^{\mathrm{G}}(z)\right)$ and $\mathbf{f}_i^{\mathbf{G}} = \left(f_{i,g}^{\mathrm{G}}, f_{i,r}^{\mathrm{G}}\right)$. The broad emission spectrum of EGFP is split at the

dichroic mirror, resulting in identical shapes of the z-scan profile in both channels. Their intensity ratio is

specified by the crosstalk vector $\boldsymbol{\alpha}^{\mathbf{G}}$, which is defined by $\mathbf{f}_i^{\mathbf{G}} = \boldsymbol{\alpha}^{\mathbf{G}} f_i^{\mathrm{G}}$, where $f_i^{\mathrm{G}} = f_{i,g}^{\mathrm{G}} + f_{i,r}^{\mathrm{G}}$ is the sum

amplitude of both channels. Similarly, the DC z-scan intensity profiles of the red fluorescent protein mCherry

(CH) are given by

514
$$\mathbf{F}^{\mathbf{CH}}(z) = \sum_i \mathbf{f}_i^{\mathbf{CH}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{CH}}, b_i^{\mathrm{CH}}\right) = \boldsymbol{\alpha}^{\mathbf{CH}} \sum_i f_i^{\mathrm{CH}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{CH}}, b_i^{\mathrm{CH}}\right),$$
13

515 and the presence of both proteins is described by the sum of Eqs. 12 and 13,

516
$$\mathbf{F}(z) = \mathbf{F}^{\mathbf{G}}(z) + \mathbf{F}^{\mathbf{CH}}(z).$$
14

517 Only a few scan geometries are of interest for modeling of the data in this study. The first uses a single slab

518 layer to describe the distribution of EGFP (X = G) and mCherry (X = CH) labeled species,

519
$$\mathbf{F}_{\mathbf{S}}^{\mathbf{X}}(z) = \boldsymbol{\alpha}^{\mathbf{X}} f^{\mathrm{X}} \, \mathrm{v}\left(z; a^{\mathrm{X}}, b^{\mathrm{X}}\right),$$
15

520 as illustrated in Fig. 1A. The second geometry consists of a delta layer followed by a slab layer and another

521 delta layer, which is referred to as (δSδ) layer model (20–22). This model was introduced to describe

522 peripheral membrane proteins with the δ layers representing the top and bottom plasma membrane, while

523 the S layer describes the cytoplasmic space (21). The DC z-scan (δSδ) model is given by

524
$$\mathbf{F}_{\boldsymbol{\delta S \delta}}^{\mathbf{X}}(z) = \boldsymbol{\alpha}^{\mathbf{X}} \left(f_{\mathrm{B}}^{\mathrm{X}} \cdot \mathrm{v}\left(z; a^{\mathrm{X}}, a^{\mathrm{X}}\right) + f_{\mathrm{Cy}}^{\mathrm{X}} \cdot \mathrm{v}\left(z; a^{\mathrm{X}}, b^{\mathrm{X}}\right) + f_{\mathrm{T}}^{\mathrm{X}} \cdot \mathrm{v}\left(z; b^{\mathrm{X}}, b^{\mathrm{X}}\right) \right),$$
16

where a^{X} and b^{X} mark the axial height of the bottom (B) and top (T) membrane (see Fig. 1G).

The fluorescence amplitudes of the B, T and cytoplasmic (Cy) layer are given by $f_{\mathrm{B}}^{\mathrm{X}}$, $f_{\mathrm{T}}^{\mathrm{X}}$, and

$f_{\mathrm{Cy}}^{\mathrm{X}}$.

The DC z-scan of cells co-expressing EGFP and mCherry labeled proteins is described by a superposition

of their individual profiles (Eq. 14). The notation X^{G}-Y^{CH} was used to describe a model where the z-scan

intensity of the EGFP-labeled protein is described by the X profile, while the z-scan intensity of the mCherry-

labeled protein is given by the Y profile,

532
$$\mathbf{F}_{\mathbf{X\text{-}Y}}(z) = \mathbf{F}_{\mathbf{X}}^{\mathbf{G}}(z) + \mathbf{F}_{\mathbf{Y}}^{\mathbf{CH}}(z),$$
17

533 where X and Y denote either the S or δSδ model (Eqs. 15 and 16). For example, the S^{G}-S^{CH} model is based

534 on independent slab layers for the EGFP and mCherry-labeled proteins, while the $δSδ^{G}$-S^{CH} model uses the

535 (δSδ) model for the EGFP label and a slab layer for mCherry.

Determining cell thickness from the z-scan intensity profile

For a cell section that is thick with respect to the RIPSF width, a direct fit of the intensity profile to a slab model (Eq. 15) determines the cell thickness by $L_{cell} = b^{CH} - a^{CH}$. For cell sections that are thin with respect to the RIPSF, cell thickness cannot be reliably determined from fitting to a slab model (*32*). However, because the fluorescent protein mCherry is distributed uniformly throughout the cell interior, its z-scan intensity provides a measure of local cell thickness L_{cell}. In this manner the cell thickness in thin sections can be determined from the integrated fluorescence Λ^{CH} of the intensity profile. Λ^{CH} is proportional to the integrated concentration area A^{CH} (Eq. 11), which is $L_{cell}c^{CH}$ for the slab geometry, where the mCherry concentration c^{CH} is assumed constant throughout the cell. Thus, the ratio of the integrated intensity profiles from a thick and a thin section measured in the same cell equals their thickness ratio, $\Lambda_{thin}^{CH}/\Lambda_{thick}^{CH} = L_{cell,thin}/L_{cell,thick}$, which was used to determine the thickness of thin cell sections. Although previous work (*32*) defined the cutoff between thick and thin cell lections as L_{cell} = 0.5 µm, a more conservative value of 1.5 µm was used in this study to improve the precision of length measurements .

Production of PM blebs

Bulges along the plasma membrane (i.e., blebs) were induced along the PM of cultured HeLa cells by adding 10-15% ethanol (*23*) to the cell growth medium and incubating for ~15 min at 37°C and 5% CO_2. The ethanol-treated cells were subsequently analyzed by transmitted light microscopy to assess the number and size of blebs with the goal of producing blebs with ~5µm diameter in sufficient quantity for measurement. If insufficient blebbing was observed, cells were returned to the incubator for ~10 min before re-checking. Once sufficient blebbing was obtained, the growth medium with ethanol was exchanged by PBS containing ethanol immediately before measurement.

SYTO RNASelect cell stain

Live cells were stained with SYTO RNASelect (Thermo Fisher Scientific, Waltham, MA) by diluting the stock dye 1000x (5 µM final concentration) into the cell culture medium followed by incubation for 1 h at 37 °C and 5% CO_2. For measurement, the growth medium with SYTO RNASelect was replaced by PBS containing the same concentration of the dye.

Z-scan redaction and statistical analysis

Despite the selection of scan locations as described earlier, some of the measured z-scan intensity profiles were not adequately described by our fit model. This was attributed to scans that inadvertently passed through complex intracellular structures. Since these structures are not captured by the model framework, the spatial autocorrelation of the normalized residual from the fit was calculated to identify patterns that are indicative of the presence of extraneous cellular features not described by the fit model. Scans with a correlation amplitude exceeding a threshold value were removed. Furthermore, z-scan measurements of the cortex thickness were selected to have a cell thickness $L_{cell} \geq 2.5$ µm and a membrane fraction (22) $m \geq 0.5$ in order to ensure the absence of biases in the fitted values as supported by our study using simulated DC z-scans (Fig. S1). Quality control for the z-scan fits of cortex thickness was provided by averaging the normalized residuals within ±1 µm of the fitted PM locations. Scans where these averaged residuals exceeded 2.5 in either the red or green channel were redacted.

A typical z-scan acquisition at a single location contained 12 repeated scans, which were fit individually, yielding 12 estimates of each parameter of interest. This set of parameters served to calculate the average effective depletion length $\overline{\Delta}_i$ and its standard deviation σ_i at the i-th z-scan location. Multiple measurement locations were summarized using the weighted average $\langle \Delta \rangle$ over all locations with $1/\sigma_i^2$ as weight. The uncertainty in the weighted average was computed using the dispersion corrected standard error of the mean (SEM),

$$SEM^2 = \sigma_{we}^2 \left(\frac{1}{n-1} \sum_i \frac{(\overline{\Delta}_i - \langle \Delta \rangle)^2}{\sigma_i^2} \right)$$

where σ_{we}^2 is the variance of the weighted average. Since it was observed that $\langle \Delta \rangle$ at the top (T) and bottom (B) PM interfaces are in agreement ($\langle \Delta^T \rangle \approx \langle \Delta^B \rangle$), data from both interfaces were grouped together when calculating $\langle \Delta \rangle$. This is indicated by dropping the T and B superscript.

Simulations of z-scan traces

Simulated DC z-scan traces were obtained following a previously described procedure (*20*) to evaluate

models and estimate the statistical uncertainty in DC z-scan fit results. Simulated DC z-scan traces were fit

using the same algorithms that were applied to experimental data. All data derived from simulations

represent the average result from fits of *n* = 1000 z-scan traces.

References

1.   P. Chugh, E. K. Paluch, The actin cortex at a glance. *J Cell Sci*. **131** (2018),

doi:10.1242/jcs.186254.

2.   A. Kusumi, T. K. Fujiwara, R. Chadda, M. Xie, T. A. Tsunoyama, Z. Kalay, R. S. Kasai,

K. G. N. Suzuki, Dynamic organizing principles of the plasma membrane that regulate signal

transduction: commemorating the fortieth anniversary of Singer and Nicolson's fluid-mosaic

model. *Annu Rev Cell Dev Biol*. **28**, 215–250 (2012).

3.   M. D. Blower, Molecular insights into intracellular RNA localization. *Int Rev Cell Mol

Biol*. **302**, 1–39 (2013).

4.   C. Medioni, K. Mowry, F. Besse, Principles and roles of mRNA localization in animal

development. *Development*. **139**, 3263–3276 (2012).

5.   E. Lécuyer, H. Yoshida, N. Parthasarathy, C. Alm, T. Babak, T. Cerovina, T. R. Hughes,

P. Tomancak, H. M. Krause, Global analysis of mRNA localization reveals a prominent role in

organizing cellular architecture and function. *Cell*. **131**, 174–187 (2007).

6.   A. Gopal, Z. H. Zhou, C. M. Knobler, W. M. Gelbart, Visualizing large RNA molecules in

solution. *RNA*. **18**, 284–299 (2012).

7.   A. Borodavka, S. W. Singaram, P. G. Stockley, W. M. Gelbart, A. Ben-Shaul, R. Tuma,

Sizes of Long RNA Molecules Are Determined by the Branching Patterns of Their Secondary

Structures. *Biophys J*. **111**, 2077–2085 (2016).

8.   S. S. Sommer, J. E. Cohen, The size distributions of proteins, mRNA, and nuclear RNA. *J

Mol Evol*. **15**, 37–57 (1980).

9.   I. Legnini, J. Alles, N. Karaiskos, S. Ayoub, N. Rajewsky, FLAM-seq: full-length mRNA

sequencing reveals principles of poly(A) tail length control. *Nat Methods*. **16**, 879–886 (2019).

10.   M. F. Tam, J. A. Dodd, W. E. Hill, Physical characteristics of 16 S rRNA under

reconstitution conditions. *J Biol Chem*. **256**, 6430–6434 (1981).

11.   A. Verschoor, J. Frank, Three-dimensional structure of the mammalian cytoplasmic

ribosome. *J Mol Biol*. **214**, 737–749 (1990).

12.   G. T. Charras, T. J. Mitchison, L. Mahadevan, Animal cell hydraulics. *J Cell Sci*. **122**,

3233–3241 (2009).

13.   E. Moeendarbary, L. Valon, M. Fritzsche, A. R. Harris, D. A. Moulding, A. J. Thrasher, E.

Stride, L. Mahadevan, G. T. Charras, The cytoplasm of living cells behaves as a poroelastic

material. *Nat Mater*. **12**, 253–261 (2013).

14.   K. Luby-Phelps, P. E. Castle, D. L. Taylor, F. Lanni, Hindered diffusion of inert tracer

particles in the cytoplasm of mouse 3T3 cells. *Proc Natl Acad Sci U S A*. **84**, 4910–4913 (1987).

15.   K. Luby-Phelps, D. L. Taylor, Subcellular compartmentalization by local differentiation of

cytoplasmic structure. *Cell Motil Cytoskeleton*. **10**, 28–37 (1988).

16.   L. Hou, F. Lanni, K. Luby-Phelps, Tracer diffusion in F-actin and Ficoll mixtures. Toward

a model for cytoplasm. *Biophys J*. **58**, 31–43 (1990).

17.   G. L. Lukacs, P. Haggie, O. Seksek, D. Lechardeur, N. Freedman, A. S. Verkman, Size-

dependent DNA mobility in cytoplasm and nucleus. *J Biol Chem*. **275**, 1625–1629 (2000).

18.   E. Dauty, A. S. Verkman, Actin cytoskeleton as the principal determinant of size-

dependent DNA mobility in cytoplasm: a new barrier for non-viral gene delivery. *J Biol Chem*.

280, 7823–7828 (2005).

19.   K. Luby-Phelps, Effect of cytoarchitecture on the transport and localization of protein

synthetic machinery. *Journal of Cellular Biochemistry*. **52**, 140–147 (1993).

20.   I. Angert, S. R. Karuka, J. Hennen, Y. Chen, J. P. Albanesi, L. M. Mansky, J. D. Mueller,

Sensitive Detection of Protein Binding to the Plasma Membrane with Dual-Color Z-Scan

Fluorescence. *Biophys J*. **118**, 281–293 (2020).

21.   E. M. Smith, P. J. Macdonald, Y. Chen, J. D. Mueller, Quantifying protein-protein

interactions of peripheral membrane proteins by fluorescence brightness analysis. *Biophys J*. **107**,

66–75 (2014).

22.   E. M. Smith, J. Hennen, Y. Chen, J. D. Mueller, Z-scan fluorescence profile

deconvolution of cytosolic and membrane-associated protein populations. *Anal Biochem*. **480**,

11–20 (2015).

23.   S. Saha, I.-H. Lee, A. Polley, J. T. Groves, M. Rao, S. Mayor, Diffusion of GPI-anchored

proteins is influenced by the activity of dynamic cortical actin. *Mol Biol Cell*. **26**, 4033–4045

(2015).

24.   R. E. Scott, R. G. Perkins, M. A. Zschunke, B. J. Hoerl, P. B. Maercklein, Plasma

membrane vesiculation in 3T3 and SV3T3 cells. I. Morphological and biochemical

characterization. *J Cell Sci*. **35**, 229–243 (1979).

25.   T. Baumgart, A. T. Hammond, P. Sengupta, S. T. Hess, D. A. Holowka, B. A. Baird, W.

W. Webb, Large-scale fluid/fluid phase separation of proteins and lipids in giant plasma

membrane vesicles. *Proc Natl Acad Sci U S A*. **104**, 3165–3170 (2007).

26.   F. Schneider, D. Waithe, M. P. Clausen, S. Galiani, T. Koller, G. Ozhan, C. Eggeling, E.

Sezgin, Diffusion of lipids and GPI-anchored proteins in actin-free plasma membrane vesicles

measured by STED-FCS. *Mol Biol Cell*. **28**, 1507–1518 (2017).

27.   T. M. Svitkina, Actin Cell Cortex: Structure and Molecular Organization. *Trends Cell

Biol*. **30**, 556–565 (2020).

28.   N. Morone, T. Fujiwara, K. Murase, R. S. Kasai, H. Ike, S. Yuasa, J. Usukura, A. Kusumi,

Three-dimensional reconstruction of the membrane skeleton at the plasma membrane interface by

electron tomography. *J Cell Biol*. **174**, 851–862 (2006).

29.   P. Chugh, A. G. Clark, M. B. Smith, D. A. D. Cassani, K. Dierkes, A. Ragab, P. P. Roux,

G. Charras, G. Salbreux, E. K. Paluch, Actin cortex architecture regulates cell surface tension.

Nat Cell Biol. **19**, 689–697 (2017).

30.   A. G. Clark, K. Dierkes, E. K. Paluch, Monitoring actin cortex thickness in live cells.

Biophys J. **105**, 570–580 (2013).

31.   J. Riedl, A. H. Crevenna, K. Kessenbrock, J. H. Yu, D. Neukirchen, M. Bista, F. Bradke,

D. Jenne, T. A. Holak, Z. Werb, M. Sixt, R. Wedlich-Soldner, Lifeact: a versatile marker to

visualize F-actin. *Nat Methods*. **5**, 605–607 (2008).

32.   P. J. Macdonald, Y. Chen, X. Wang, Y. Chen, J. D. Mueller, Brightness analysis by Z-

scan fluorescence fluctuation spectroscopy for the study of protein interactions within living cells.

Biophys J. **99**, 979–988 (2010).

33.   P. R. Tedbury, E. O. Freed, HIV-1 gag: an emerging target for antiretroviral therapy. *Curr*

Top Microbiol Immunol. **389**, 171–201 (2015).

34.   J. G. Levin, J. Guo, I. Rouzina, K. Musier-Forsyth, Nucleic acid chaperone activity of

HIV-1 nucleocapsid protein: critical role in reverse transcription and molecular mechanism. *Prog*

Nucleic Acid Res Mol Biol. **80**, 217–286 (2005).

35.   R. N. De Guzman, Z. R. Wu, C. C. Stalling, L. Pappalardo, P. N. Borer, M. F. Summers,

Structure of the HIV-1 nucleocapsid protein bound to the SL3 psi-RNA recognition element.

Science. **279**, 384–388 (1998).

36.   A. Rein, Nucleic acid chaperone activity of retroviral Gag proteins. *RNA Biol*. **7**, 700–705

(2010).

37.   V. Chukkapalli, S. J. Oh, A. Ono, Opposing mechanisms involving RNA and lipids

regulate HIV-1 Gag membrane binding through the highly basic region of the matrix domain.

Proc Natl Acad Sci U S A. **107**, 1600–1605 (2010).

38.   P. Purohit, S. Dupont, M. Stevenson, M. R. Green, Sequence-specific interaction between

HIV-1 matrix protein and viral genomic RNA revealed by in vitro genetic selection. *RNA*. **7**, 576–

584 (2001).

39.   D. Thornhill, B. Olety, A. Ono, Relationships between MA-RNA Binding in Cells and

Suppression of HIV-1 Gag Mislocalization to Intracellular Membranes. *J Virol*. **93** (2019),

doi:10.1128/JVI.00756-19.

40.   C. R. Gaines, E. Tkacik, A. Rivera-Oven, P. Somani, A. Achimovich, T. Alabi, A. Zhu, N.

Getachew, A. L. Yang, M. McDonough, T. Hawkins, Z. Spadaro, M. F. Summers, HIV-1 Matrix

Protein Interactions with tRNA: Implications for Membrane Targeting. *J Mol Biol*. **430**, 2113–

2127 (2018).

41.   S. B. Kutluay, T. Zang, D. Blanco-Melo, C. Powell, D. Jannain, M. Errando, P. D.

Bieniasz, Global changes in the RNA binding specificity of HIV-1 gag regulate virion genesis.

Cell. **159**, 1096–1109 (2014).

42.   Y. Iwatani, H. Takeuchi, K. Strebel, J. G. Levin, Biochemical activities of highly purified,

catalytically active human APOBEC3G: correlation with antiviral effect. *J Virol*. **80**, 5992–6002

(2006).

43.   Y. N. Friew, V. Boyko, W.-S. Hu, V. K. Pathak, Intracellular interactions between

APOBEC3G, RNA, and HIV-1 Gag: APOBEC3G multimerization is dependent on its association

with RNA. *Retrovirology*. **6**, 56 (2009).

44.   J. Li, Y. Chen, M. Li, M. A. Carpenter, R. M. McDougle, E. M. Luengas, P. J. Macdonald,

R. S. Harris, J. D. Mueller, APOBEC3 multimerization correlates with HIV-1 packaging and

restriction activity in living cells. *J Mol Biol*. **426**, 1296–1307 (2014).

45. E. S. Svarovskaia, H. Xu, J. L. Mbisa, R. Barr, R. J. Gorelick, A. Ono, E. O. Freed, W.-S.

Hu, V. K. Pathak, Human apolipoprotein B mRNA-editing enzyme-catalytic polypeptide-like 3G

(APOBEC3G) is incorporated into HIV-1 virions through interactions with viral and nonviral

RNAs. *J Biol Chem*. **279**, 35822–35828 (2004).

46. A. Burnett, P. Spearman, APOBEC3G multimers are recruited to the plasma membrane

for packaging into human immunodeficiency virus type 1 virus-like particles in an RNA-

dependent process requiring the NC basic linker. *J Virol*. **81**, 5000–5013 (2007).

47. B. R. Cullen, Role and mechanism of action of the APOBEC3 family of antiretroviral

resistance factors. *J Virol*. **80**, 1067–1076 (2006).

48. J. S. Albin, R. S. Harris, Interactions of host APOBEC3 restriction factors with HIV-1 in

vivo: implications for therapeutics. *Expert Rev Mol Med*. **12**, e4 (2010).

49. J. F. Hultquist, J. A. Lengyel, E. W. Refsland, R. S. LaRue, L. Lackey, W. L. Brown, R. S.

Harris, Human and rhesus APOBEC3D, APOBEC3F, APOBEC3G, and APOBEC3H

demonstrate a conserved capacity to restrict Vif-deficient HIV-1. *J Virol*. **85**, 11220–11234

(2011).

50. L. W. Janson, K. Ragsdale, K. Luby-Phelps, Mechanism and size cutoff for steric

exclusion from actin-rich cytoplasmic domains. *Biophys J*. **71**, 1228–1234 (1996).

51. A. P. Minton, Confinement as a determinant of macromolecular structure and reactivity.

Biophys J. **63**, 1090–1100 (1992).

52. J. C. Giddings, E. Kucera, C. P. Russell, M. N. Myers, Statistical theory for the

equilibrium distribution of rigid molecules in inert porous networks. Exclusion chromatography.

J. Phys. Chem. **72**, 4397–4408 (1968).

53. T. Kalwarczyk, N. Ziebacz, A. Bielejewska, E. Zaboklicka, K. Koynov, J. Szymański, A.

Wilk, A. Patkowski, J. Gapiński, H.-J. Butt, R. Hołyst, Comparative analysis of viscosity of

complex liquids and cytoplasm of mammalian cells at the nanoscale. *Nano Lett*. **11**, 2157–2163

(2011).

54. R. Holyst, A. Bielejewska, J. Szymański, A. Wilk, A. Patkowski, J. Gapiński, A.

Zywociński, T. Kalwarczyk, E. Kalwarczyk, M. Tabaka, N. Ziebacz, S. A. Wieczorek, Scaling

form of viscosity at all length-scales in poly(ethylene glycol) solutions studied by fluorescence

correlation spectroscopy and capillary electrophoresis. *Phys Chem Chem Phys*. **11**, 9025–9032

(2009).

55. L. Xiang, K. Chen, R. Yan, W. Li, K. Xu, Single-molecule displacement mapping unveils

nanoscale heterogeneities in intracellular diffusivity. *Nat Methods*. **17**, 524–530 (2020).

56. T. P. Stossel, P. A. Janmey, K. S. Zaner, in *Cytomechanics: The Mechanical Basis of Cell*

Form and Structure, J. Bereiter-Hahn, O. R. Anderson, W.-E. Reif, Eds. (Springer, Berlin,

Heidelberg, 1987; https://doi.org/10.1007/978-3-642-72863-1_9), pp. 131–153.

57. J.-F. Joanny, J. Prost, Active gels as a description of the actin-myosin cytoskeleton. *HFSP*

J. **3**, 94–104 (2009).

58. T. C. Laurent, J. Killander, A theory of gel filtration and its exeperimental verification.

Journal of Chromatography A. **14**, 317–330 (1964).

59. K. C. Duong-Ly, S. B. Gabelli, Gel filtration chromatography (size exclusion

chromatography) of proteins. *Methods Enzymol*. **541**, 105–114 (2014).

60. T. Kaur, I. Alshareedah, W. Wang, J. Ngo, M. M. Moosa, P. R. Banerjee, Molecular

Crowding Tunes Material States of Ribonucleoprotein Condensates. *Biomolecules*. **9** (2019),

doi:10.3390/biom9020071.

61. A. A. M. André, E. Spruijt, Liquid-Liquid Phase Separation in Crowded Environments. *Int*

J Mol Sci. **21** (2020), doi:10.3390/ijms21165908.

62. D. W. Provance, A. McDowall, M. Marko, K. Luby-Phelps, Cytoarchitecture of size-

excluding compartments in living cells. *J Cell Sci*. **106 (Pt 2)**, 565–577 (1993).

63.   K. Luby-Phelps, D. L. Taylor, F. Lanni, Probing the structure of cytoplasm. *J Cell Biol*. **102**, 2015–2022 (1986).

64.   R. Kumar, S. Saha, B. Sinha, Cell spread area and traction forces determine myosin-II-based cortex thickness regulation. *Biochim Biophys Acta Mol Cell Res*. **1866**, 118516 (2019).

65.   T. M. Svitkina, A. A. Shevelev, A. D. Bershadsky, V. I. Gelfand, Cytoskeleton of mouse embryo fibroblasts. Electron microscopy of platinum replicas. *Eur J Cell Biol*. **34**, 64–74 (1984).

66.   F. Feijó Delgado, N. Cermak, V. C. Hecht, S. Son, Y. Li, S. M. Knudsen, S. Olcum, J. M. Higgins, J. Chen, W. H. Grover, S. R. Manalis, Intracellular water exchange for measuring the dry mass, water mass and changes in chemical composition of living cells. *PLoS One*. **8**, e67590 (2013).

67.   M. A. Urbaneja, B. P. Kane, D. G. Johnson, R. J. Gorelick, L. E. Henderson, J. R. Casas-Finet, Binding properties of the human immunodeficiency virus type 1 nucleocapsid protein p7 to a model RNA: elucidation of the structural determinants for function. *J Mol Biol*. **287**, 59–75 (1999).

68.   W. Ouyang, S. Okaine, M. P. McPike, Y. Lin, P. N. Borer, Probing the RNA binding surface of the HIV-1 nucleocapsid protein by site-directed mutagenesis. *Biochemistry*. **52**, 3358–3368 (2013).

69.   M. Comas-Garcia, S. A. Datta, L. Baker, R. Varma, P. R. Gudla, A. Rein, Dissection of specific binding of HIV-1 Gag to the "packaging signal" in viral RNA. *Elife*. **6** (2017), doi:10.7554/eLife.27055.

70.   M. Delarue, G. P. Brittingham, S. Pfeffer, I. V. Surovtsev, S. Pinglay, K. J. Kennedy, M. Schaffer, J. I. Gutierrez, D. Sang, G. Poterewicz, J. K. Chung, J. M. Plitzko, J. T. Groves, C. Jacobs-Wagner, B. D. Engel, L. J. Holt, mTORC1 Controls Phase Separation and the Biophysical Properties of the Cytoplasm by Tuning Crowding. *Cell*. **174**, 338-349.e20 (2018).

71.   K. Kwapiszewska, K. Szczepański, T. Kalwarczyk, B. Michalska, P. Patalas-Krawczyk, J.

Szymański, T. Andryszewski, M. Iwan, J. Duszyński, R. Hołyst, Nanoscale Viscosity of

Cytoplasm Is Conserved in Human Cell Lines. *J Phys Chem Lett*. **11**, 6914–6920 (2020).

72.   J. Chen, D. Grunwald, L. Sardo, A. Galli, S. Plisov, O. A. Nikolaitchik, D. Chen, S.

Lockett, D. R. Larson, V. K. Pathak, W.-S. Hu, Cytoplasmic HIV-1 RNA is mainly transported

by diffusion in the presence or absence of Gag protein. *Proc Natl Acad Sci U S A*. **111**, E5205-

5213 (2014).

73.   J. J. Blum, G. Lawler, M. Reed, I. Shin, Effect of cytoskeletal geometry on intracellular

diffusion. *Biophys J*. **56**, 995–1005 (1989).

74.   K. Luby-Phelps, F. Lanni, D. L. Taylor, The submicroscopic properties of cytoplasm as a

determinant of cellular function. *Annu Rev Biophys Biophys Chem*. **17**, 369–396 (1988).

75.   D. S. Banks, C. Fradin, Anomalous diffusion of proteins due to molecular crowding.

Biophys J. **89**, 2960–2971 (2005).

76.   A. Miermont, F. Waharte, S. Hu, M. N. McClean, S. Bottani, S. Léon, P. Hersen, Severe

osmotic compression triggers a slowdown of intracellular signaling, which can be explained by

molecular crowding. *Proc Natl Acad Sci U S A*. **110**, 5725–5730 (2013).

77.   M. Weiss, M. Elsner, F. Kartberg, T. Nilsson, Anomalous subdiffusion is a measure for

cytoplasmic crowding in living cells. *Biophys J*. **87**, 3518–3524 (2004).

78.   C. I. Lacayo, J. A. Theriot, Listeria monocytogenes actin-based motility varies depending

on subcellular location: a kinematic probe for cytoarchitecture. *Mol Biol Cell*. **15**, 2164–2175

(2004).

79.   J.-Y. Youn, B. J. A. Dyakov, J. Zhang, J. D. R. Knight, R. M. Vernon, J. D. Forman-Kay,

A.-C. Gingras, Properties of Stress Granule and P-Body Proteomes. *Mol Cell*. **76**, 286–294

(2019).

80.   A. A. Hyman, C. A. Weber, F. Jülicher, Liquid-liquid phase separation in biology. *Annu*

Rev Cell Dev Biol. **30**, 39–58 (2014).

81.   L.-P. Bergeron-Sandoval, N. Safaee, S. W. Michnick, Mechanisms and Consequences of

Macromolecular Phase Separation. *Cell*. **165**, 1067–1079 (2016).

82.   R. Wollman, T. Meyer, Coordinated oscillations in cortical actin and Ca2+ correlate with

cycles of vesicle secretion. *Nat Cell Biol*. **14**, 1261–1269 (2012).

83.   P. Li, A. T. Bademosi, J. Luo, F. A. Meunier, Actin Remodeling in Regulated Exocytosis:

Toward a Mesoscopic View. *Trends Cell Biol*. **28**, 685–697 (2018).

84.   A. Derdowski, L. Ding, P. Spearman, A novel fluorescence resonance energy transfer

assay demonstrates that the human immunodeficiency virus type 1 Pr55Gag I domain mediates

Gag-Gag interactions. *J Virol*. **78**, 1230–1242 (2004).

85.   C. Thaler, S. V. Koushik, H. L. Puhl, P. S. Blank, S. S. Vogel, Structural rearrangement of

CaMKIIalpha catalytic domains encodes activation. *Proc Natl Acad Sci U S A*. **106**, 6369–6374

(2009).

86.   B. Wu, Y. Chen, J. D. Müller, Fluorescence fluctuation spectroscopy of mCherry in living

cells. *Biophys J*. **96**, 2391–2404 (2009).

87.   B. Wu, Y. Chen, J. D. Müller, Heterospecies partition analysis reveals binding curve and

stoichiometry of protein interactions in living cells. *Proc Natl Acad Sci U S A*. **107**, 4117–4122

(2010).

88.   Y. Chen, H.-Q. Sun, J. P. Eichorst, J. P. Albanesi, H. Yin, J. D. Mueller, Comobility of

GABARAP and Phosphatidylinositol 4-Kinase 2A on Cytoplasmic Vesicles. *Biochemistry*. **57**,

3556–3559 (2018).

89.   J. D. Müller, Y. Chen, E. Gratton, Fluorescence correlation spectroscopy. *Methods*

Enzymol. **361**, 69–92 (2003).

Acknowledgments

Non-author contributions: The authors thank Yan Chen for help with logistics and experiment planning.

Author contributions: CIA, JDM, and SRK developed the dual color z-scan method. CIA and SRK performed experiments and acquired the data. LMM helped interpret the data and guide the project. CIA and JDM wrote analysis code, performed data analysis, and drafted the manuscript. All authors reviewed and revised the manuscript.

Funding: This work was supported by NIH grants GM064589, GM124279 and GM098550. C. I. A. was supported by NIH T32 AI083196

Competing interests: The authors declare that they have no financial or other competing interests.

Data and materials availability: The raw data and code necessary to perform the analyses will be made available through the Dryad data repository.

Fig. 1: DC z-scans in living cells. The z-scan trajectory through the cell is depicted in the cartoons along

with crosshatched regions indicating the subcellular localization of the EGFP- and mCherry-labeled proteins

(A, D, G, J, left panel) together with the concentration profile indicated by z-scan fitting for the EGFP (green

lines) and mCherry (magenta lines) protein species (A, D, G, J, right panel). B, E, H, K) Fits (solid lines) of

representative z-scan intensity traces of the green (green diamonds) and red (magenta diamonds) detection

channel are shown along with their normalized residuals sr. The cell edge locations a^G, b^G, a^{CH}, b^{CH} extracted

from the fits determine the vertical distance Δ separating the EGFP- and mCherry-labeled slab edges in

each z-scan. C, F, I, L) Histograms of $\overline{\Delta}$ from repeated z-scan experiments in many cells. Experiments with

different EGFP- and mCherry-labeled protein species are indicated in rows: (A, B, C) EGFP and mCherry-

RXR, (D, E, F) EGFP and mCherry, (G, H, I) EGFP-HRas and mCherry, and (J, K, L) A3G-EGFP and

mCherry.

Fig. 2: Z-scans in blebs produced from HeLa cells co-expressing A3G-EGFP and mCherry. A) A

graphic depicting the production of large PM blebs from adherent cells via ethanol treatment is shown. In

this example, the cytoskeleton is denoted by brown colors. B) A DC 2-photon laser scanning z-stack image

of a HeLa cell with two large PM blebs is shown. The x-y image was created by the summation of the z-

stack over the z-dimension. The y-z image was produced by the sum of the x-dimension over two volume-

sections passing roughly through the center of each bleb. While significant red autofluorescence is produced

by the ethanol treatment near the cell center, it is typically absent in the blebs. Z-scan locations were selected

to pass near the center of each bleb, while avoiding regions possessing autofluorescence or identifiable

structures. C) Fit of a representative z-scan passing through a HeLa cell bleb (χ_v^2= 1.2). D) Histogram of $\overline{\Delta}$

values in the cytoplasm of HeLa cells not treated with ethanol (magenta bars) and in blebs produced by

ethanol treatment (gray bars).

Fig. 3: DC z-scan measurement of cortex thickness. A) Lifeact-mApple demarcates the actin cortex, and EGFP-HRas demarcates the PM location. These fluorescent proteins lead to the SSS^{CH}-$\delta S\delta^G$ concentration profile depicted in (B). C) DC z-scans were fit to a $\delta S\delta^{CH}$-$\delta S\delta^G$ model. The location of the mApple δ layers corresponds to the centroid of the cortical slabs, so that the effective depletion length is half the thickness of the cortex slab. D) A fit of a DC z-scan through a HeLa cell co-expressing EGFP-HRas and Lifeact-mApple. E) Average cortex thickness values estimated by DC z-scan in HeLa and U2OS cells. F) $\langle\Delta\rangle$ is plotted in U2OS cells versus the same vales in HeLa cells for MA^{G2A}-EGFP (brown), Gag^{G2A}-EGFP (green) and A3G-EGFP (magenta). The ratio of cortex thicknesses in the cell lines corresponds to the slope of the data as is indicated by the shaded area and dashed lines, which correspond to the slope-uncertainty derived from the data in panel (E).

Fig. 4 Effective depletion length. A) $\langle\Delta\rangle$ in HeLa cells for the indicated proteins and an RNA dye. B) Model

of cortical partitioning leading to a SSS^G distribution of fluorescence. C) Axial concentration profile

corresponding to the model. The concentration drops from c_{cyto} in the cytoplasm to c_C in the cortex. D)

Approximation of the SSS^G by a single S^G concentration profile. Fitting of the experimental z-scan data to

this profile was used to identify apparent depletion lengths Δ^B and Δ^T. The mCherry concentration profile

has been omitted in panels C and D. E) $\langle\Delta\rangle$ vs partition coefficient R_C extracted from simulated DC z-scans

for three different cortex lengths ($L_C^B = L_C^T$ of 200, 100, and 50 nm). The solid lines represent a theoretical

model based on Eq. 1.

Fig. 5: Axially averaged partition coefficient $\overline{R}_z$ vs cell thickness. A-D) Representative DC z-scan traces in a thin and thick section of a HeLa cell co-expressing EGFP and mCherry (A, B) and representative DC z-scan traces in thin and thick sections of a HeLa cell co-expressing Gag^{G2A}-EGFP and mCherry. E) Schematic illustration of DC z-scans taken at multiple cytoplasmic locations varying from thick to thin sections to determine $\overline{R}_z$ as a function of thickness L_{cell}. F) $\overline{R}_z$ vs L_{cell}. Black diamonds correspond to control measurements in HeLa cells co-expressing EGFP and mCherry. Magenta and blue diamonds correspond to data from HeLa and U2OS cells co-expressing HIV-1 Gag^{G2A}-EGFP and mCherry. Magenta and blue lines correspond to Eq. 5 with $L_C = 200\,\text{nm}$ and $L_C = 130\,\text{nm}$, respectively, and $R_C = 0.2$.

Fig. 6: Protein mobility and effective depletion length. A) Depletion length <Δ> versus effective diffusion

coefficient D_{eff} in HeLa cells (diamonds). Eq. 8 (magenta curve) is plotted with parameter values as

described in the text. A3G-EGFP measured in HeLa cell blebs (green open circle) displayed increased

mobility along with decreased effective depletion length. B) Conceptual model of the effects quantified in

panel A. The partitioning effect (center left) arises due to hindered entry of large RNP complexes into the

cortical mesh, while hindered mobility of large RNP complexes due to macromolecular crowding occurs in

the cytoplasm (right). Binding of labeled proteins to partitioned RNP complexes yields the concentration

gradient (far left) observed by DC z-scan measurements. C) The cytoplasmic mobility and partition

coefficient are connected by the average RNP complex radius, as indicated in the graphical depiction and

suggested by Eqs. 7 and 8.