Partitioning of ribonucleoprotein complexes from the cellular actin cortex

Isaac Angert[1,2,3], Siddarth Reddy Karuka[1], Louis M. Mansky[2,3,4], Joachim D. Mueller[1,2,5,†]

School of Physics and Astronomy[1], University of Minnesota, Minneapolis, MN 55455
Institute of Molecular Virology[2], University of Minnesota, Minneapolis, MN 55455
Division of Basic Sciences, School of Dentistry[3], University of Minnesota, Minneapolis, MN 55455
Masonic Cancer Center[4], University of Minnesota, Minneapolis, MN 55455
Department of Biomedical Engineering[5], University of Minnesota, Minneapolis, MN 55455

†Corresponding author. Email: jochen@umn.edu

Abstract

The cell cortex plays a crucial role in cell mechanics, signaling, and development. However, little is known about the influence of the cortical meshwork on the spatial distribution of cytoplasmic biomolecules. Here, we describe a new fluorescence microscopy method to infer the intracellular distribution of labeled biomolecules with sub-resolution accuracy. Unexpectedly, we find that RNA-binding proteins are partially excluded from the cytoplasmic volume adjacent to the plasma membrane that corresponds to the actin cortex. Complementary diffusion measurements of RNA-protein complexes suggest that a rudimentary model based on excluded volume interactions can explain this partitioning effect. Our results suggest the actin cortex meshwork may play a role in regulating the biomolecular content of the volume immediately adjacent to the plasma membrane.

Teaser

A novel microscopy method reveals exclusion of RNA-protein complexes from the actin cortex due to their large hydrodynamic size.

Introduction

Both the actin cortex and RNA are fundamental building blocks of eukaryotic cells. The cortex, acting in concert with other cytoskeletal elements, is involved in nearly all aspects of cell mechanics, such as cell structure, membrane rigidity, migration and division (*1*). Due to its location adjacent to the plasma membrane, the cortex plays important roles in signal transduction and mechanical response pathways (*1*,

2). In contrast, RNA molecules serve as genomic messengers and constitute major components of the cellular translational machinery. Specific localization of mRNA to sub-cellular regions has been widely observed (3). RNA localization to sub-cellular compartments such as the endoplasmic reticulum (ER), processing (P)-bodies or distal cytoplasm in neurons is critical to aspects of development (4) and is an important regulator of cellular function (5). Due to their highly structured and extended nature, mRNAs have hydrodynamic radii that are dramatically larger than those of the globular proteins they encode (6). Based on a recent study (7), a typical mRNA of 1 – 2 kilobase length (8, 9) is expected to have a hydrodynamic radius in the range of ~10 nm. The radii of ribosomes are of comparable size (10 – 15 nm) (10, 11).

Previous biophysical studies have established that the cell cytoplasm is a complex medium (12, 13) that modulates the diffusion and localization of large (>10 nm) particles in a size-dependent manner (14–18). These observations have prompted the question of whether the large size of the cellular translational machinery might broadly influence its subcellular localization (19). Here, the novel observation that large riboneucleoprotein (RNP) complexes are partially excluded, or partitioned, from the interior of the actin cortex is reported. These observations are facilitated by recent developments in two-photon dual-color (DC) z-scan fluorescence microscopy (20), which is adapted from its previous application quantifying binding of peripheral membrane proteins to now infer protein localizations inside living cells with sub-diffraction resolution of lengths. DC z-scans are acquired by moving the two-photon excitation volume vertically through the cytoplasm of adherent cells. This movement of the excitation volume yields fluorescent intensity traces that encode the vertical intracellular distribution of a fluorescently labeled protein species as well as that of a distinctly colored reference fluorescent protein. Although different reference species are possible, our typical choice is a soluble, purely cytoplasmic protein, which serves as a "paint" that marks the cytoplasmic extent. Differential vertical localization between the fluorescently labeled protein and the reference protein is encoded as subtle differences in the shape of the DC z-scan traces. The magnitude of the differential vertical localization is quantitatively recovered from the data by fitting the z-scan traces. In a typical sample, this procedure attains a resolution of 20 – 50 nm in after collection of 60 s of z-scan data.

Results

DC z-scan measures sub-diffraction distances in living cells

Initial proof-of-principle experiments were conducted with U2OS cells to demonstrate the utility of DC z-scan to quantify sub-diffraction differences in intracellular protein localizations. For detailed descriptions of the z-

scan model geometries used in the following, see the Materials and Methods section. First, cells were co-

transfected with mCherry-RXR (a protein that localizes in the nucleus) and EGFP (which diffuses freely

throughout the cell). DC z-scans through the cell nucleus were fit to the S^G-S^{CH} (Eq. 13) model to identify

the slab length associated with the mCherry fluorescence, which corresponds to the nuclear thickness, and

the slab length of the EGFP signal, which corresponds to the slightly longer thickness of the entire cell (Fig.

1A). A single representative z-scan, acquired in 5 s and subsequently fitted, shows that the mCherry-RXR

and EGFP z-scans are well described by this model, with flat residuals and χ_v^2 = 1.2 (Fig. 1B). The fit

identified the intensity amplitudes of mCherry-RXR and EGFP, the locations of the bottom and top cell edges

(a^G = 2.16 µm and b^G = 5.80 µm), as well as the locations of the bottom and top nuclear edges (a^{CH} = 2.51

µm and b^{CH} = 5.03 µm), respectively. The distances between the edges of the EGFP and mCherry slab were

defined at the top and bottom of the cell by $\Delta^T = b^{CH} - b^G$ = -770 nm and $\Delta^B = a^G - a^{CH}$ = -350 nm, respectively.

Since Δ measures the displacement between the borders of the EGFP and mCherry layers, its value can be

positive or negative in general. Distances are reported by taking the absolute value, which in this instance

determines the thickness of the cytoplasmic layers surrounding the top and bottom of the nucleus (Fig. 1A).

To improve the resolution and identify statistical uncertainties, 60 s of repeated z-scans were then acquired

at the same location within this cell and subsequently fit to obtain an average of Δ^B = -410 ± 16 nm and Δ^T

= -800 ± 15 nm (SEM from 12 scans). Multiple 60 s DC z-scan measurements were then acquired in multiple

cells. Each of these experiments yielded estimates of Δ^B and Δ^T at each location within a cell. These values

are summarized by a histogram (Fig. 1C) and the resulting distribution is characterized by the averages (see

Materials and Methods) $<\Delta^T>$ = -340 ± 50 nm and $<\Delta^B>$ = -260 ± 30 nm (SEM, n = 19, where n is the number

of scan locations averaged).

Importantly, the DC z-scan experiments reported in the present work were performed using two-

photon microscopy, which allows EGFP and mCherry to be co-excited by using the same laser focus. This

results in red and green channel point spread functions (PSFs) that are identical at the nanometer scale

(Fig. S1). This lack of chromatic shift can be seen in control DC z-scans that were performed through the

cytoplasm of U2OS cells co-expressing EGFP and mCherry fluorescent proteins (Fig. 1D). Since both

proteins are soluble species within the cell cytoplasm, DC z-scans are expected to recover Δ = 0, on average.

As shown in Fig. 1E, a representative DC z-scan fits to a S^G-S^{CH} (Eq. 13) model with flat residuals and χ_v^2 =

1.1. Repeated DC z-scans in multiple cells recovered mean values of $<\Delta^B>$ = -2.3 ± 1.2 nm and $<\Delta^T>$ = 1.0

± 3.1 nm (SEM, n = 59). In the absence of a statistically significant difference between the results from the

bottom and top of the cell, the data were combined, and the superscript dropped. The histogram of the

combined displacements (Fig. 1F) determined $<\Delta> = -0.5 \pm 1.7$ nm (SEM, $n = 118$).

Next, DC z-scans were performed in U2OS cells co-expressing EGFP-HRas and mCherry (Fig. 1G).

EGFP-HRas was previously reported to be a peripheral membrane protein that binds to the plasma

membrane as a monomer (*21*). Previous z-scan studies established that the axial distribution of EGFP-HRas

in the cytoplasm and at the PM was successfully modeled by the δSδ layer geometry (*21*, *22*), where the

two δ layers account for PM bound protein and the S layer accounts for protein in the cytoplasm. Thus, DC

z-scans from cells co-expressing mCherry and EGFP-HRas were modeled by the $(S^{CH}\text{-}\delta S\delta^G)$ geometry (Eq.

13). A fit of a representative z-scan attained good quality of fit with $\chi_v^2 = 1.2$ and flat residuals. Repeated z-

scans in the cytoplasm of multiple cells co-expressing EGFP-HRas and mCherry (Fig. 1I) determined a

mean displacement of $<\Delta> = -2.6 \pm 5.4$ nm (SEM, $n = 30$), which is consistent with the expected value of Δ

$= 0$. This result indicated that the location of the cytoplasm slab edge, as determined by fitting the soluble

mCherry species to the S model, coincided with the location of the PM, as determined by fitting the EGFP-

HRas species to the δSδ model.

APOBEC 3G is depleted from a thin volume adjacent to the PM

Next, the axial cytoplasmic distribution of APOBEC 3G (A3G) labeled with EGFP and co-expressed

with mCherry (Fig. 1J) was examined. A3G is part of the innate immune system and acts as a cytidine

deaminase that restricts replication of Vif-negative HIV-1. A3G binds strongly to both cellular and viral RNAs,

the latter of which is essential to its retroviral restriction activity (*23*, *24*). A representative z-scan acquired in

the cytoplasm of a HeLa cell showed a single broad peak for both mCherry and A3G-EGFP (Fig. 1K). Since

the trace shape was suggestive of a slab geometry, both species were fit to slab models $(S^{CH}\text{-}S^G$, Eq. 13).

This fit provided a good description of the DC z-scan intensity profiles with $\chi_v^2 = 1.3$ and flat residuals (Fig.

1K). Interestingly, this fit identified non-zero values for the displacements at the top and bottom PM ($\Delta^B =$

184 nm and $\Delta^T = 167$ nm). Repeated z-scan acquisitions at the same location in this cell determined average

values $\Delta^B = 180 \pm 6$ nm and $\Delta^T = 162 \pm 12$ nm (SEM from 12 scans). This surprising observation suggests

that the thickness of the soluble A3G-EGFP layer is less than the thickness of the mCherry layer. Repeated

z-scan acquisitions in multiple HeLa cells (Fig. 1L) found an average of $<\Delta>_{HeLa} = 155 \pm 10$ nm (SEM, $n =$

64), indicating that A3G does not occupy the entire cytoplasmic volume. To reflect these results Δ is also

referred to as the effective depletion length. To further validate this observation, DC z-scan measurements

were collected using a different cell line (U2OS), which resulted in an average of $<\Delta>_{U2OS} = 110 \pm 9$ nm

(SEM, $n = 32$). Rescaled histograms of Fig. 1 are shown in Fig. S2 to facilitate visual comparison of the

different samples. Because the uncertainty in the measured depletion length may arise from intracellular or

intercellular variability, the variation of the depletion length at distinct locations within the same cell and

between multiple cells was determined (Table S1), which indicates that the majority of the variability is

accounted for by intracellular heterogeneity. Additionally, the data in Table S1 demonstrate that DC z-scan

attains a resolution of 20- 50 nm in typical samples after acquisition of 60 s of z-scan data.

PM adjacent depletion is reduced by disrupting F-actin

Intriguingly, this result implies that cytoplasmic A3G-EGFP was excluded from regions adjacent to

the PM with apparent thickness of 100 - 200 nm. Given the magnitude of this value, further experiments

were conducted to test whether A3G-EGFP might be partially excluded from the dense meshwork of the

actin cortex. To do this, HeLa cells were treated with ethanol, which disrupts actin assemblies and leads to

the formation of large plasma membrane blebs (*25*) (Fig. 2A) that lack an intact actin cortex (*26*, *27*). A

representative z-scan performed near the center of these blebs showed good quality of fit to a S^{CH}-S^{G} model

(Eq. 13 and Fig. 2C). Repeated z-scans in blebs from HeLa cells co-expressing A3G-EGFP and mCherry

(Fig. 2D) determined an average of $<\Delta> = 31 \pm 10$ nm (SEM, $n = 14$), which differs from the results obtained

in untreated HeLa cells by 8.8 SD.

Because the ethanol treatment may induce pleiotropic effects, it was verified that blebs induced in

HeLa cells expressing A3G-EGFP by cytochalasin D treatment showed a similar reduction in the depletion

effect (Fig. S3). To provide additional support for the role of the cortex in PM adjacent depletion, DC z-scans

were performed in HeLa cells treated with latrunculin A (LA), a toxin which prevents actin polymerization. In

contrast to experiments with cytochalasin D, LA was used at a concentration low enough to maintain gross

cell morphology (i.e., below the bleb-forming concentration). DC z-scans performed in the cytoplasm of LA

treated cells indicated a reduction in $<\Delta>$ compared to a solvent control (Fig. S3). Taken together, the strong

reduction in Δ found in blebs and the decrease observed in the presence of latrunculin A support the

conclusion that PM adjacent depletion is driven by the presence of the actin cortex.

The depletion length depends on the cortical thickness

Given that dependence of cortical architecture on the cell type has been observed (*28–30)*, the

differences in effective depletion length observed between HeLa and U2OS cells might reflect natural

differences in cortex thickness between these cell types. To investigate this, direct measurements of the

cortex thickness were performed using distinct fluorescent colors to mark the actin cortex and the plasma

membrane. A previous study used this approach to estimate the actin cortex thickness from the displacement

between intensity peaks corresponding to the actin and the PM marker in confocal images (*31)*. In the

present study, two-photon DC z-scan was employed instead of confocal imaging to determine the cortex

thickness from the intensity profiles of EGFP-HRas (which binds to the PM) and Lifeact-mApple (which binds

to actin and marks the cortex) (*32)* by fitting to on a quantitative model of the entire intensity traces (Fig. 3A).

The axial concentration profile of both labels is depicted in Fig. 3B, which shows an enrichment of EGFP-

HRas and Lifeact-mApple at the PM and within the cortex, respectively. A small amount of both fluorescent

proteins is also present in the cytoplasm. Based on this profile, EGFP-HRas is again modeled by the δSδ

geometry. The distance separating the two HRas δ layers represents the cell thickness L_{cell}.

The concentration profile of Lifeact-mApple implies a wide cytoplasmic slab of length L_{cyto}

separating two thin slabs representing the cortex at the bottom and the top membrane with thickness L_C^B

and L_C^T (Fig. 3B), giving rise to an SSS^{CH} (triple S-layer) geometry. A uniform slab with a sharp transition

between the cortical and cytoplasmic spaces model of the cortex has been previously used (*31)*. Because

the expected slab length L_C of the cortex is smaller than 500 nm, its length is experimentally indistinguishable

from a δ layer (*33)* by single-color (SC) z-scans. However, the centroid position of the cortical layer is

expected to coincide with the midpoint of the slab as depicted in Fig. 3B. Given this, modeling the Lifeact-

mApple species as a δSδ^{CH} geometry can identify the cortex centroid from the fitted location of the δ layers.

Combined with the EGFP-HRas species, this DC z-scan fit (δSδ^{CH}-δSδ^G) identifies the locations of the δ

layers corresponding to the cortex centroid and the PM as well as their displacement *Δ*. The distance |*Δ*|

separating both δ layers is half of the actin cortex thickness, |*Δ*| = L_C / 2 (Fig. 3B). This relation was verified

by fitting simulated DC z-scan intensity profiles from the SSS^{CH}-δSδ^G geometry to the δSδ^{CH}-δSδ^G model

(Fig. S4). Repeated DC z-scans were then collected in cells co-expressing EGFP-HRas and Lifeact-mApple

proteins in order to determine cortical thickness. A representative DC z-scan and fit to the δSδ^{CH}- δSδ^G

model (Fig. 3C) for both proteins is shown in Fig. 3D. Data taken from multiple cells resulted in cortex

thickness estimates of 126 ± 16 nm (SEM, n = 7) for U2OS cells and 200 ± 24 nm (SEM, n = 15). for HeLa

cells (Fig. 3E).

Since both the depletion length Δ for A3G-EGFP and the cortex thickness differed between the two

cell lines, <Δ> in HeLa and U2OS cells were plotted for comparison (Fig. 3F). Additional DC z-scans were

performed in cells expressing HIV-1 Gag^{G2A}-EGFP and its matrix (MA) domain, MA^{G2A}-EGFP, with mCherry

again as a reference marker of the cytoplasmic volume for each instance. These <Δ> values were added to

Fig. 3F (for histograms, see Fig. S5). Interestingly, all 3 data points cluster near a line of slope 0.66, indicating

that, although the magnitude of the depletion effect varies for each protein, it is systematically scaled by

~66% between these two cell lines. This same ratio corresponds well to the cortex thickness ratio estimated

by DC z-scan between HeLa and U2OS cells (0.66 ± 0.11, shaded area in Fig. 3F), suggesting that the cell-

type dependent differences in <Δ> are due to differences in their cortical actin thicknesses.

PM adjacent depletion of proteins is driven by RNA-binding

All 3 proteins in Fig. 3F are known to be RNA-interacting proteins: HIV-1 Gag^{G2A} binds specifically

and non-specifically to RNA sequences through basic residues and two zinc fingers in its nucleocapsid (NC)

domain (*34–36*), while the MA domain of Gag separately interacts with RNA via a stretch of basic residues

in its highly basic region (HBR) (*37*) and has preferential tRNA binding affinity (*38*). A3G interacts with RNA

primarily through its N-terminal domain zinc finger (*23, 24*). To test whether the depletion effect is specific

to RNA-binding proteins, DC z-scans were conducted with the GABA-A receptor associated protein fused

to EGFP (GABARAP-EGFP) and a tandem hexamer of the Venus fluorescent protein, Venus$_6$. Both proteins

lack a known RNA binding function. The average <Δ> value of both proteins was found to be < 10 nm

(Fig. 4A) and therefore treated as consistent with zero, since globular proteins typically have an intrinsic

diameter of a few nanometers.

For comparison, the previously presented depletion lengths from RNA binding proteins (from Fig.

3D) were added to Fig. 4A. Since RNA binding appeared to be important for observing depletion, DC z-

scans were performed with the mutant A3G^{C97A}-EGFP, which has impaired RNA binding due to mutation of

the N-terminal zinc finger (*39*). These measurements revealed a large reduction in <Δ> compared to that of

wildtype A3G-EGFP (Fig. 4A). Lastly, the <Δ> value was measured in mCherry expressing HeLa cells that

were stained with the small molecule RNA binding dye, SYTO RNASelect. These measurements revealed

that the labeled cellular RNA had a robust depletion length (Fig. 4A).

Based on these results, it was hypothesized that a partial exclusion, or partitioning, of large RNA species from the dense cortical meshwork may be responsible for the observed PM adjacent depletion of RNA binding proteins (Fig. 4B). To determine whether engagement of RNAs at ribosomes might contribute to this effect, DC z-scans were performed in U2OS cells expressing HIV-1 Gag[G2A]-EGFP and mCherry that were treated with puromycin to induce premature chain termination and dislodge ribosomes from mRNAs. Puromycin treatment resulted in $<\Delta>$ values that were unchanged compared to untreated cells (Fig. S3), suggesting that engagement of mRNAs at ribosomes is not required for cortical partitioning of HIV-1 Gag.

DC z-scan modeling of cortical partitioning

In a simple model of the cortical partitioning (Fig. 4C), the ratio of the concentration c_C in the cortical layer to the concentration c_{cyto} in the cytoplasm defines the partition coefficient $R_C = c_C / c_{\mathrm{cyto}}$. This partitioning of the protein concentration between the cytoplasmic and cortical spaces introduces a SSS[G] concentration profile for the EGFP-labeled RNA-binding protein (Fig. 4C), which consists of two thin cortical slabs of length L_C^B and L_C^T separated by a thick cytoplasmic slab of length L_{cyto} with a higher protein concentration. As before (Fig. 3), the cortical layer is too narrow to be directly resolved by SC z-scan analysis. However, since the concentration within the cortex layer is reduced (Fig. 4C), the z-scan intensity profile can be effectively approximated by a single slab layer with a width that slightly exceeds the cytoplasmic thickness L_{cyto} (Fig. 4D). Consequently, the apparent depletion lengths Δ^B and Δ^T will be less than the corresponding cortex thickness L_C^B and L_C^T in a manner that depends on R_C. In z-scan experiments, this dependence is well approximated by

$$\Delta = (1 - R_C) L_C, \qquad\qquad 1$$

where the superscript has been omitted. Eq. 1 is derived by preserving the integrated fluorescent areas (i.e., $A_C = A_x$) between the partitioning model (Fig. 4C) and the fit model (Fig. 4D). See Supplemental Text 1 for details. This model (Fig. 4E, solid lines) is in excellent agreement with the results from fitting simulated DC z-scan traces of an SSS[G]-S[CH] geometry to the S[CH]-S[G] DC z-scan model (Fig. S4). Eq. 1 was used to estimate the partition coefficient of proteins from their measured depletion lengths (Fig. 3D) using the experimentally determined cortex thicknesses L_C. R_C = 0.76 ± 0.03, 0.33 ± 0.09, and 0.22 ± 0.1 for MA[G2A]-EGFP, Gag[G2A]-

237 EGFP, and A3G-EGFP, respectively, were determined from HeLa cells. Similar values (0.75 ± 0.04, 0.39 ±

238 0.08, and 0.13 ± 0.13) were obtained from the measurements in U2OS cells.

Direct observation of the partition coefficient in the periphery of HeLa cells

Because these estimates of the partition coefficient R_C rely on measurements of Δ and L_C, which is

a new method, a second approach was developed to directly estimate R_C in thin cytoplasmic regions. To

illustrate this approach, a ratio image of the red and green detection channels was calculated from a two-

photon z-stack acquired in cells co-expressing HIV-1 Gag[G2A]-EGFP and mCherry (Fig. S6). The ratio image

reveals that Gag[G2A]-EGFP is more excluded from thin peripheral regions than thick regions of the cytoplasm

when compared with the mCherry reference. A similar effect was previously documented using large

fluorescent molecules that were microinjected into the cytoplasm (*15, 40*). Based on the partitioning model

presented in Fig. 4C, it is anticipated that this relative exclusion seen in peripheral regions is caused by close

opposition of the top and bottom cortical layers in thin cytoplasmic sections. As a control, Gag[G2A]-EGFP was

replaced with EGFP alone. The resulting ratio image showed an approximately constant intensity ratio

throughout the cytoplasm (Fig. S6), as expected given the lack of cortical partitioning for EGFP.

The same relative exclusion of HIV-1 Gag[G2A]-EGFP compared to mCherry is detected by DC z-

scans in thick and thin regions of a HeLa cell (Fig. 5C & D). In contrast, DC z-scans in thick and thin regions

of a cell co-expressing EGFP and mCherry indicated that the relative concentrations of the EGFP and

mCherry species are identical regardless of cytoplasm thickness (Fig. 5A & B). Next, a sequence of z-scans

was taken within a single cell along a line that extended from thick to thin cytoplasmic regions. The axially

averaged partition coefficient $\overline{R}_z$, which describes the cumulative partitioning inclusive of both cortical and

unpartitioned cytoplasmic layers was constructed from the data (see Supplemental Text 2). Assuming that

the cortex thickness L_C and the partition coefficient R_C are constant throughout the cell as illustrated in Fig.

5E, an analytical expression for $\overline{R}_z$ was derived (Eq. S4; see Supplemental Text 2). This theoretical $\overline{R}_z$

approaches 1 in thick cell sections where the cortical layers are small compared to the cytoplasmic layer,

while it approaches R_C in thin cell sections where the unpartitioned layer is small compared to the cortex.

While realistic cortices are unlikely to have homogeneous F-actin structures extending from the center to the

periphery of cells, (i.e., to have constant values of R_C and L_C in the cell periphery) this model (Fig. 5E)

remains useful as baseline against which to assess experimental data.

As shown in Fig. 5F, the theoretical $\bar{R}_z$ model agrees with the DC z-scan estimates of $\bar{R}_z$ from

HeLa cells. The HeLa cell data in Fig. 5F cluster near 0.23 ± 0.02 (SEM, *n* = 6) in thin cell sections. This

value is consistent with the R_C value determined above. As a control, the same experiment was carried out

in HeLa cells co-expressing EGFP and mCherry. These values remain near 1 independent of thickness (Fig.

5F), as expected due to the lack of cortical partitioning for EGFP. Interestingly, although DC z-scan estimates

of $\bar{R}_z$ for Gag[G2A]-EGFP in U2OS cells are close to 1 for thick cell sections, they appear to deviate from the

model (Eq. S4) in thin cell sections (Fig. 5F). This deviation may reflect the presence of distinct peripheral

F-actin structures in this cell line (*28*), which imply changes in the R_C and L_C.

Cortical partitioning and cytoplasmic diffusion are quantitatively related

The cortical partitioning effect documented here supports a model where protein depletion within

the cortex is driven by protein association into ribonucleoprotein complexes that are much larger than the

typical protein. In such a case, the large <*Δ*> values should be correlated with substantially slowed

cytoplasmic diffusion. FCS measurements were performed in HeLa cells in order to estimate the effective

diffusion coefficient D_{eff} of labeled proteins in the cytoplasm (see Materials and Methods and representative

FCS curves in Fig. S7). Because FCS data were acquired with the two-photon PSF focused near the middle

of thick (>1.5 μm) cytoplasmic sections, the estimates of D_{eff} correspond to the mobility within the cytoplasm

and not the mobility within the actin cortex. In Fig. 6A <*Δ*> values from Fig. 4 were plotted versus the median

D_{eff} that was obtained from FCS measurements. Two other members of the APOBEC3 protein family, A3A-

EGFP and A3C-EGFP, which have differing antiretroviral activities (*41–43*), were also measured and added

to Fig. 6A. These data show that large <*Δ*> values are associated with remarkably low cytoplasmic mobility.

This relationship is summarized by the cartoons in Fig. 6 B&C.

The average pore diameter of the actin cortex has been reported to range between 50 to 200 nm

(*29*), depending on cell type. These values are only slightly larger than or overlap with the size of large RNAs

(*6*, *7*), so that a partial partitioning of RNAs from the interior of the cortex could be driven by excluded volume

interactions. Consequently, the cortical partitioning was described as arising from excluded volume

interactions of particles that are allowed to enter spherical voids (*44–46*),

$$R_C = (1 - r_p/r_v)^3 , \qquad\qquad 2$$

where r_p is the particle radius, and r_v is the void radius. A hindered cytoplasmic diffusion coefficient D_{eff}

was modeled as arising from a size-dependent viscosity imposed by macromolecular crowding within the

cytoplasm (*47*, *48*) according to

297
$$\frac{D_{eff}}{D_0} = \exp\left[-\left(\frac{r_{eff}}{\xi}\right)^a\right],$$
3

298   where D_0 is a reference diffusion coefficient, r_{eff} is an effective hydrodynamic radius of the diffusing

299   particles, a is a scaling exponent of order 1, and ξ is a length interpreted as the characteristic distance

300   between points of entanglement within the crowded solution. Using Eqs. 1, 2, & 3, and setting $r_{eff} = r_p$ yields

301   a relation between D_{eff} and Δ,

302
$$\Delta = L_c\left[1-\left(1-\frac{\xi}{r_v}\ln\left(\frac{D_0}{D_{eff}}\right)^{1/a}\right)^3\right].$$
4

This curve is plotted as the magenta line in Fig. 6A using values L_C = 170 nm, D_0 equal to D_{eff}

measured for EGFP (27.5 μm²s⁻¹), r_v = 60 nm, ξ = 4 nm and a = 0.7. The values for ξ and a are within the

range estimated by (*47*) for HeLa cells. r_v is similar to published estimates of the cortical mesh size (*29*). The

hydrodynamic radius r_{eff} of the diffusing protein complexes was calculated (Eq. 3), yielding values in the

range of 10 to 40 nm for the slowly diffusing proteins in Fig. 6A (Table S2).

The above model postulates that the cortical partitioning effect arises solely from large

hydrodynamic size and is independent of any other interactions between RNPs and the cellular milieu. To

check this assumption, DC z-scans and FCS measurements were performed in U2OS cells expressing

mCherry that were loaded with 500 kDa green-fluorescent dextrans via electroporation. The measurement

of 500 kDa dextrans yielded a relatively large <*Δ*> value and slow diffusion (red star in Fig. 6A), which is

consistent with our model. A separate FCS measurement of the 500 kDa dextrans in water estimated a

hydrodynamic radius of ~44 nm, which is comparable to the r_{eff} values obtained from Eq. 3 for slowly diffusing

protein complexes in Fig. 6A (Table S2).

Cytoplasmic crowding in cell associated PM blebs may differ from crowding within undisturbed

cytoplasm due at least in part to the lack of a cytoskeletal mesh in the bleb interior. Thus, the mobility of

A3G-EGFP in HeLa cell blebs was investigated. FCS measurements (Fig. 6A, open circle) showed an order

of magnitude increase in D_{eff} compared to A3G in the cytoplasm of untreated HeLa cells. The value from

blebs is plotted as an open circle in order to emphasize that the model parameters (especially r_v and ξ) that

describe the rest of the data most likely do not apply to the measurements in blebs.

It was recently reported that the cytoplasmic diffusion coefficient of fluorescent proteins depended

on their net charge, with positively charged proteins diffusing slowly. It was suggested that this slow diffusion

was due to transient interactions with large, negatively charged cell components such as RNA and the

cytoskeleton (*49*). An examination was performed to determine whether the D_{eff} values in Fig. 6A might be

explained by the net charge of each protein-FP fusion. As shown in Fig. S8, the mobility of the proteins in

Fig. 6A was only weakly correlated with net charge. Thus, the mobility of these proteins in living cells is likely

to be better understood as arising predominantly from their hydrodynamic size as a consequence of their

RNA binding function.

Discussion

In this study, extension of the DC z-scan microscopy technique to measurements of protein localization at

sub-diffraction length scales provided the foundation for identifying the cortical partitioning effect and allowed

its quantitative description through the effective depletion length *Δ*. Implementation of the DC z-scan

measurements using two-photon microscopy eliminates systematic biases, such as chromatic aberration,

from the collected data. This absence of biases together with accurate models describing the DC z-scan

intensity trace identifies subtle changes in the distribution of fluorescently labeled proteins along the scan

path. DC z-scan measurements indicated an apparent restriction in the cytoplasmic volume occupied by

RNA binding proteins (e.g., A3G and Gag^{G2A}). This depletion effect was markedly reduced in cell blebs and

cells treated with latrunculin A, and was observed to be cell-type dependent in a manner that correlated with

DC z-scan estimates of cortex thickness in adherent interphase HeLa and U2OS cells. Thus, the depletion

effect is interpreted as being a partitioning, or partial exclusion, of these proteins from the volume of the actin

cortex. Furthermore, *Δ* values measured for proteins with and without RNA-binding function suggest that the

cortical partitioning is driven by association into RNP complexes with large hydrodynamic size and is not

associated with protein molecular mass. The independence of *Δ* from protein mass implies that these RNP

complexes have hydrodynamic radii far larger than those attained by globular proteins. This conclusion is

supported by measurements of the diffusion coefficient, which shows that a marked decrease in mobility is

associated with proteins that have the largest values of Δ (Fig. 6).

In vitro and in vivo (*50*) observations of actin assemblies suggest that gel-like properties are nearly

ubiquitous, including for the actin cortex specifically (*51*). The cortical partitioning effect may be analogous

to solute partitioning from the interior of gels, a well-known chemical phenomenon (*52*, *53*). Furthermore,

the subcellular heterogeneity driven by the dense cortical environment is reminiscent of those molecular

crowding effects which have been reported to drive association or phase separations in cells (*54*, *55*). This

suggests that the crowded environment within gels may drive biophysical phenomena similar to, but distinct

from those associated with molecular crowding in the cytoplasm. The observation of cortical partitioning

described in this report builds on earlier work that found strong exclusion of labeled colloidal particles with

radii exceeding ~10 nm from distal regions of mammalian cell cytoplasm, and regions near the centrosome

(*15*, *40*). Subsequent studies documented that the distal excluding regions were associated with a dense

cytoskeletal mesh and were thinner than non-excluding regions (*56*). These and other observations (*44*)

have suggested that this partitioning was driven by the cytoskeletal meshwork.

The cortical partitioning of RNPs described in the present study also leads to strong exclusion from

distal regions of the cytoplasm (Fig. 5 & S6). However, unlike previous studies, which linked partitioning to

the actin cytoskeleton in general (*44*), our observations identified the actin cortex as the causative agent of

partitioning. Differentiation between the cortical space and the bulk cytoplasm was not previously achieved

and probably reflects experimental limitations in these earlier studies, which are overcome by the use of DC

z-scan method employed here. Thus, the cortical partitioning of RNPs observed in this work and the distal

exclusion of microinjected particles documented previously (*15*, *40*) likely are manifestations of the same

physical phenomenon, namely the size-dependent partitioning of particles from the actin cortex.

The measurement of the actin cortex thickness, L_C, in HeLa (~200 nm) and U2OS (~130 nm) cells

in our study represents a novel application of the DC z-scan technique. This application was inspired by a

conceptually similar approach based on confocal microscopy that was used to estimate a cortex thickness

of ~190 nm in mitotic HeLa cells (*31*) and of 300 – 400 nm in interphase HeLa cells with imposed spherical

shapes (*30*). Interestingly, DC z-scan identified a cortical thickness ≤200 nm for adherent interphase HeLa

and U2OS. This is consistent with literature reports which indicate that cortical architecture is strongly

associated with cell morphology (*28*, *30*). In particular, cell adhesion and cell spreading on the glass coverslip

can result in a thinning of the actin cortex (*28*, *57*). Thus, differing cortical thicknesses reported here and in

literature most likely represent inherent differences associated with the disparate cell morphologies studied.

Measurements of the axially averaged partition coefficient $\bar{R}_z$ provided further evidence linking

partitioning with the cellular cortex by observing $\bar{R}_z$ as a function of cell thickness (Fig. 5F). Modeling of $\bar{R}_z$

by assuming uniform cortex properties recovered parameters consistent with DC z-scan analysis in the case

of HeLa cells. Interestingly, estimates of $\bar{R}_z$ in U2OS cells exhibited more scatter than observed in HeLa

cells, particularly in thin cell sections. This may reflect cell-type differences in F-actin assemblies. In EM

micrographs, HeLa cells show an unusually uniform cortex, even near the cell periphery, while large

structures such as peripheral lamellar actin sheets are more pronounced in other cell types (*28*, *58*). Because

F-actin structures in lamellae appear distinct from cortical actin fibers, these structures are anticipated to

partition large complexes in a manner distinct from the uniform cortex model of $\bar{R}_z$.

The strong correlation between the ability of a protein to bind RNA and measured values of <*Δ*>

(Fig. 4A) suggests that many RNA-protein complexes are too large to efficiently enter the dense actin cortical

meshwork. The cortical partitioning of RNA-binding proteins measured by DC z-scan is thus interpreted as

arising from an underlying partitioning associated with the large RNA species with which these proteins

associate. In this framework, DC z-scans of RNA interacting proteins indirectly probe the spatial distribution

of cellular RNAs. More directly, when cellular RNA was labeled with an RNA-binding fluorescent dye, DC z-

scan measurements of <*Δ*> revealed that the RNA itself is partitioned from the cortex with R_C in the range

0.35 – 0.2 (given L_C 170 – 200 nm). This R_C value is interpreted as representing the average R_C of a broad

spectrum of differently sized cellular RNAs (i.e., mRNAs, tRNAs, etc) and RNP complexes (e.g., ribosomes)

with different degrees of cortical penetration. Similarly, the <*Δ*> values for fluorescently labeled proteins

likely represent the average R_C of a heterogenous mix of differently sized RNA complexes to which each

protein preferentially binds. Because polysomes can form extremely large structures, the potential for mRNA

engagement at ribosomes to drive cortical partitioning was explored by treating cells with puromycin, which

causes dissociation of mRNA-ribosome complexes (*59*). This experiment, performed with Gag[G2A]-EGFP,

showed no detectable change in <*Δ*> (Fig. S3A), demonstrating that the hydrodynamic size of the probed

mRNAs is sufficient for establishing cortical depletion.

FCS measurements were acquired in order to characterize protein mobility in the cytoplasm. The

diffusion time of RNA binding proteins revealed that large <*Δ*> values were associated with very low mobility

(Fig. 6A), as expected for proteins that are present in large complexes. Slow diffusion of large sized objects

in the cytoplasm has been previously reported (*14*, *17*, *60*, *61*), including mRNAs (*62*). Previous studies have

suggested that this low cytoplasmic mobility may reflect a hindrance created by cytoskeletal elements (*15,*

16, 18, 19, 56, 63, 64) or by cytoplasmic crowding created due to ribosomes, organelles and proteins (*17,*

60, 65–67). Although the precise determinants for cytoplasmic mobility have not been fully characterized,

existing literature provides extensive evidence that the cytoplasm cannot be described as a simple fluid (*12,*

47, 65, 68), and it is generally agreed that particles in the range of 10 – 100 nm show a size-dependent drop

in cytoplasmic mobility much greater than that predicted by the Stokes-Einstein relation (*14, 17, 18, 60*). In

this study, a marked increase in A3G-EGFP mobility in cell-associated blebs (Fig. 6A) was observed. This

suggests a role for the cytoskeleton in the mobility of large macromolecular complexes.

To capture the size dependence of RNP diffusion, an empirical hydrodynamic scaling model (Eq. 3

) was used. This and similar models have been previously employed to describe diffusion in polymer

solutions and in living cells (*47, 48*). This model led to good agreement with our experimental data (Fig. 6A)

when combined with a relatively simple description of the size dependence of cortical partitioning (*44–46*)

(Eq. 2). Although large RNP complexes are the focus of the current work, the physical model should apply

to any large, freely diffusing complex. This generality of the cortical partitioning phenomena is supported by

data from 500 kDa dextran polymers, which are not native to the cellular milieu but nonetheless show a

robust partitioning effect in line with their hydrodynamic size (Fig. 6A). This demonstrates that cortical

partitioning may be understood as arising from the simple physical interactions captured by Eqs. 1 – 3 without

appeal to complex biochemical interactions that occur in the living cell.

In the cortical partitioning phenomenon, the physical properties of the cortical space appear to drive

subcellular segregation of macromolecules in the absence of a clear boundary such as a membrane. Other

instances of biomolecule compartmentalization in the absence of membranes have been recognized, such

as that observed between P-bodies and the cytoplasm (*69*), and between nucleoli and the nucleoplasm (*70*).

In such cases, the distinct physical properties of each subcellular compartment regulate their molecular

content (*71*). Partitioning of large complexes from the cortical space is consistent with observations

demonstrating that a highly orchestrated coupling between exocytic pathways and cortical actin

polymerization is needed to facilitate passage of vesicles through the cortex when the actin fiber density is

transiently lowered (*72, 73*). Moreover, the cortical partitioning effect might hinder the replication of

enveloped viruses which target large macromolecular complexes to the PM as part of their assembly

processes. In the particular case of HIV-1, a large RNA genome must transit the cortical space in order be

packaged into the nascent viral particles that assemble at the PM. Recruitment of the viral RNA to assembly

sites is driven by the Gag protein, which binds to viral and cellular RNAs (*74*). Because both Gag and the

viral RNA are subject to cortical partitioning, the effect may present a barrier to HIV-1 replication that must

be overcome during the assembly process.

In conclusion, the extension of DC z-scan microscopy to measure sub-diffraction length scales

enabled the observation and characterization of an unexpected cortical partitioning effect that applies to

large RNP complexes. A simple quantitative model was introduced to provide a rudimentary foundation for

understanding the origins of this effect. The DC z-scan toolkit should be of interest to probe the spatial

relation of biomolecules with other cellular structures as well as continue to provide additional insights into

the function of the actin cortex.

447 Materials and Methods

448 Cell Culture and Plasmids

HeLa and U2OS (Numbers CCL-2 and HTB-96, American Type Culture Collection, Manassas, VA) cells

were cultured in Dulbecco's modified Eagle's medium supplemented with 10% FBS (HyClone Laboratories,

Logan, UT) at $37^{\circ}C$ and 5% CO_2. Cells were plated at ~30% confluency into 8-well chambered slides ~24 h

prior to measurement and transfected with vectors as indicated in the text ~16 h prior to measurement using

GenJet transfection reagent (SignaGen Laboratories, Fredrick, MD) according to the manufacturer's

instructions. The cell growth medium was replaced with phosphate buffered saline immediately prior to

measurement. All measurements were performed at room temperature.

HIV-1 Gag (75) was ligated into pEGFP-N1 (Clontech, Takara Bio, Mountain View, CA) to generate the HIV-

1 Gag-EGFP vector. HIV-1 Gag^{G2A} -EGFP was made using the QuikChange XL Site-Directed Mutagenesis

Kit (Stratagene, La Jolla, CA). Venus$_6$ was a gift from Steven Vogel (76) (Addgene plasmid # 27813;

http://n2t.net/addgene:27813; RRID: Addgene_27813). mApple-Lifeact-7 (denoted Lifeact-mApple here)

was a gift from Michael Davidson (Addgene plasmid # 54747; http://n2t.net/addgene:54747; RRID:

Addgene_54747). All other vectors have been previously described: A3G-EGFP, A3G^{C97A}-EGFP, A3C-

EGFP, A3A-EGFP (39); HIV-1 MA^{G2A}-EGFP (20); mCherry (77); EGFP-HRas (21); mCherry-RXR (78);

GABARAP-EGFP (79).

465 Instrumentation

466 DC z-scans and fluorescence fluctuation spectroscopy (FFS) data were acquired on a Zeiss Axiovert 200

467 microscope modified for 2-photon excitation (20). Excitation light was provided by a femtosecond pulsed

Ti:S laser (MaiTai or Tsunami models, Spectra Physics, Santa Clara, CA) operated at 1000 nm. This light was focused to a diffraction limited point by a water immersion objective (C-Apochromat, Zeiss, Oberkochen, Germany) with numerical aperture 1.2 and correction collar set according to the glass thickness of the sample holder. Emission light was collected by the same objective and separated from the excitation light by a dichroic filter (740DCSPXR, Chroma Technology, Bellows Falls, VT), then split into red and green detection channels by a low-pass dichroic (FF580-FDi01, Semrock, Rochester, NY). A 515/50nm band pass filter placed before the green channel detector eliminated Fresnel reflected mCherry emission light from the green channel. Emission light in each channel was detected by single photon counting hybrid photomultipliers (HPM-100-40, Becker and Hickl, Berlin, Germany), recorded by hardware (FastFLIM, ISS Inc., Champaign, IL or DPC-230, Becker and Hickl, Berlin, Germany), and saved for later analysis using routines written in IDL 8.6 (Harris Geospatial Solutions, Broomfield, CO).

Axial z-scans utilized a piezo stage (PZ2000, ASI, Eugene OR) driven by a voltage waveform from an arbitrary waveform generator (model 33250A, Agilent Technologies, Santa Clara, CA) to move the sample vertically across the two-photon PSF. All z-scans were acquired by driving the piezo stage with a triangle waveform with a 10 s period and peak-to-peak displacement corresponding to 24 μm. Each period contained two z-scans: the first z-scan passed upward through the cell and the second passed downward through the cell. Each z-scan covered a 24 μm range.

XY scanning of the PSF for acquisition of two-photon images was provided by a galvanometer scan head (Yanus IV, FEI Deutschland GmbH, Planegg, Germany) aligned in the two-photon excitation path. XY positioning was driven by a DAC card (3-axis, ISS Inc., Champaign, IL) controlled by SimFCS software (version 3; G-SOFT Inc., Champaign, IL). All z-scans were acquired within 24 μm of the XY scan field center.

Z-scan modeling

The single-channel (SC) z-scan fluorescence intensity profile $F(z)$ is determined by the convolution of the axial concentration profile $c(z)$ of the fluorescently-tagged protein and the radially integrated point spread function (RIPSF) of the two-photon instrument,

$$F(z) = \lambda (c * \text{RIPSF})(z), \qquad\qquad 5$$

where λ is the brightness of the fluorescence tag. The RIPSF was described using the modified Gaussian-Lorentzian model (*22, 33*). This model is specified by a Rayleigh length z_R and exponent y, which are

497   calibrated from z-scans through a thin cell section or dye layer (*33*). Calibration values $z_R \sim 0.65$ μm and

498   exponent $y \sim 1.25$ are characteristic of the two-photon instrument used in this study. Evaluation of Eq. 5 for

499   different concentration profiles $c(z)$ gives rise to an alternative formulation with a fluorescence amplitude

500   f and a function $\mathrm{v}(z)$ describing the shape of the concentration profile (*33*),

501
$$F(z) = f\,\mathrm{v}(z).$$
6

502   The axially integrated concentration $A = \int c(z)\,dz$ and the axially integrated fluorescence

503    $\Lambda = \int F(z)\,dz$ are related according to Eq. 5 by

504
$$\Lambda = \eta A$$
7

505   with the constant $\eta = \lambda \int \mathrm{RIPSF}(z)\,dz$.

Z-scan and FFS acquisition in live cells

Cells were selected by verifying the desired expression levels for the mCherry and EGFP labeled constructs

using 1-photon epi illumination provided by a mercury lamp and using an EGFP/mCherry filter set. Points

within cells for z-scan and FFS data collection were chosen where the fluorescence appeared locally (~3 μm

radius) uniform as visualized in epi illumination and where a cell lacked nearby internal structures when

visualized by using transmitted light. All measurements of cortical partitioning were performed in the

cytoplasm. For data acquisition, the microscope was switched to two-photon mode and the excitation volume

was focused to the middle of the cell. At each location chosen, a z-scan acquisition consisted of multiple

repeated vertical scans (see Instrumentation section). In most instances, 60 s of z-scan data acquisition

(i.e., 12 repeated scan passes) was followed by an optional 60 s of FFS data acquisition at up to 4 unique

locations within each cell. Repeated z-scan acquisitions at a given location were measured for a maximum

of 10 min in dim samples or where increased statistical precision was desired.

Effective diffusion coefficient

521   The effective diffusion coefficient was calculated as (*80*) $D_{eff} = \omega_0^2/(8\tau_D)$, where ω_0^2 is the $1/e^2$ radial

522   beam waist of the 2-photon excitation light focus and τ_D was estimated by fitting the log-averaged

523   autocorrelation of photon counts $G(\tau)$ to $G_{2D}(\tau) = G_0/(1+(\tau/\tau_D))^{-1} + G_{offset}$. FCS data quality control was

524   performed by redacting data where G_{offset} showed significantly negative values or where τ_D indicated by

525   the fit was >50% higher than the time at which the correlation amplitude reached half its initial value.

Modeling of DC z-scan intensity profiles

528   The DC fluorescence intensity $\mathbf{F^G}(z)$ of a z-scan intensity profile of EGFP (G)-labeled protein through a

529   series of horizontal layers, each containing a uniform concentration of labels, is described by (*20–22*)

$$\mathbf{F^G}(z) = \sum_i \mathbf{f}_i^{\mathbf{G}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right) = \boldsymbol{\alpha}^{\mathbf{G}} \sum_i f_i^{\mathrm{G}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right) , \qquad 8$$

where f_i^{G} and $\mathrm{v}\left(z, a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right)$ are the fluorescence amplitude and shape function associated with the *i*-th

layer. The shape function $\mathrm{v}\left(z, a_i^{\mathrm{G}}, b_i^{\mathrm{G}}\right)$ characterizes a uniform slab or S layer starting at $z = a_i^{\mathrm{G}}$ and

ending at $z = b_i^{\mathrm{G}}$, while $\mathrm{v}\left(z, a_i^{\mathrm{G}}, a_i^{\mathrm{G}}\right)$ defines an infinitesimally thin or δ layer located at $z = a_i^{\mathrm{G}}$ (*20–22*).

Vector notation is used to describe properties of the green (g) and red (r) channel, i.e.

$\mathbf{F^G}(z) = \left(F_g^{\mathrm{G}}(z), F_r^{\mathrm{G}}(z)\right)$ and $\mathbf{f}_i^{\mathbf{G}} = \left(f_{i,g}^{\mathrm{G}}, f_{i,r}^{\mathrm{G}}\right)$. The broad emission spectrum of EGFP is split at the

dichroic mirror, resulting in identical shapes of the z-scan profile in both channels. Their intensity ratio is

specified by the crosstalk vector $\boldsymbol{\alpha}^{\mathbf{G}}$, which is defined by $\mathbf{f}_i^{\mathbf{G}} = \boldsymbol{\alpha}^{\mathbf{G}} f_i^{\mathrm{G}}$, where $f_i^{\mathrm{G}} = f_{i,g}^{\mathrm{G}} + f_{i,r}^{\mathrm{G}}$ is the sum

amplitude of both channels. Similarly, the DC z-scan intensity profiles of the red fluorescent protein mCherry

(CH) are given by

$$\mathbf{F^{CH}}(z) = \sum_i \mathbf{f}_i^{\mathbf{CH}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{CH}}, b_i^{\mathrm{CH}}\right) = \boldsymbol{\alpha}^{\mathbf{CH}} \sum_i f_i^{\mathrm{CH}} \cdot \mathrm{v}\left(z; a_i^{\mathrm{CH}}, b_i^{\mathrm{CH}}\right), \qquad 9$$

541   and the presence of both proteins is described by the sum of Eqs. 8 and 9,

$$\mathbf{F}(z) = \mathbf{F^G}(z) + \mathbf{F^{CH}}(z). \qquad 10$$

543   Only a few scan geometries are of interest for modeling of the data in this study. The first uses a single slab

544   layer to describe the distribution of EGFP (X = G) and mCherry (X = CH) labeled species,

$$\mathbf{F_S^X}(z) = \boldsymbol{\alpha}^{\mathbf{X}} f^{\mathrm{X}} \mathrm{v}\left(z; a^{\mathrm{X}}, b^{\mathrm{X}}\right) , \qquad 11$$

546 as illustrated in Fig. 1A. The second geometry consists of a delta layer followed by a slab layer and another

547 delta layer, which is referred to as (δSδ) layer model (20–22). This model was introduced to describe

548 peripheral membrane proteins with the δ layers representing the top and bottom plasma membrane, while

549 the S layer describes the cytoplasmic space (21). The DC z-scan (δSδ) model is given by

550
$$\mathbf{F}_{\delta S \delta}^{\mathbf{X}}(z) = \boldsymbol{\alpha}^{\mathbf{X}}\left(f_{\mathrm{B}}^{\mathrm{X}} \cdot \mathrm{v}\left(z; a^{\mathrm{X}}, a^{\mathrm{X}}\right) + f_{\mathrm{Cy}}^{\mathrm{X}} \cdot \mathrm{v}\left(z; a^{\mathrm{X}}, b^{\mathrm{X}}\right) + f_{\mathrm{T}}^{\mathrm{X}} \cdot \mathrm{v}\left(z; b^{\mathrm{X}}, b^{\mathrm{X}}\right)\right), \qquad 12$$

where a^{X} and b^{X} mark the axial height of the bottom (B) and top (T) membrane (see Fig. 1G). The

fluorescence amplitudes of the B, T and cytoplasmic (Cy) layer are given by $f_{\mathrm{B}}^{\mathrm{X}}$, $f_{\mathrm{T}}^{\mathrm{X}}$, and $f_{\mathrm{Cy}}^{\mathrm{X}}$.

The DC z-scan of cells co-expressing EGFP and mCherry labeled proteins is described by a superposition

of their individual profiles (Eq. 10). The notation X^{G}-Y^{CH} was used to describe a model where the z-scan

intensity of the EGFP-labeled protein is described by the X profile, while the z-scan intensity of the mCherry-

labeled protein is given by the Y profile,

557
$$\mathbf{F}_{\mathbf{X}\text{-}\mathbf{Y}}(z) = \mathbf{F}_{\mathbf{X}}^{\mathbf{G}}(z) + \mathbf{F}_{\mathbf{Y}}^{\mathbf{CH}}(z), \qquad 13$$

558 where X and Y denote either the S or δSδ model (Eqs. 11 and 12). For example, the S^{G}-S^{CH} model is based

559 on independent slab layers for the EGFP and mCherry-labeled proteins, while the $\delta S \delta^{G}$-S^{CH} model uses the

560 (δSδ) model for the EGFP label and a slab layer for mCherry.

Production of PM blebs

Bulges along the plasma membrane (i.e., blebs) were induced along the PM of cultured HeLa cells by adding

10-15% ethanol (25) or 5 µM cytochalasin D (Cayman Chemical, Ann Arbor, MI) to the cell growth medium

and incubating for ~15 min at 37°C and 5% CO_2. The drug-treated cells were subsequently analyzed by

transmitted light microscopy to assess the number and size of blebs with the goal of producing blebs with

~5µm diameter in sufficient quantity for measurement. If insufficient blebbing was observed, cells were

returned to the incubator for ~10 min before re-checking. Once sufficient blebbing was obtained, the growth

medium with drug was exchanged by PBS containing the same concentration of drug immediately before

measurement. For the conditions used to generate ethanol induced blebs, it was observed that a near normal

HeLa cell morphology was obtained after ethanol containing media of blebbing cells was exchanged for

clean media and cells were incubated for ~1 h.

Although rapid bleb retraction is often observed in live cell cultures, the blebs produced by ethanol

and cytochalasin D treatment were stable during z-scan acquisitions. Bleb stability is likely due to the

impairment of the F-actin assemblies that are required for bleb retraction. Over the course of the experiment

(several hours), the population of ethanol induced blebs were observed to coalesce into one or several large

blebs per cell. Such large blebs were not measured since they tended to contain large internal structures

which confound z-scan analysis.

Puromycin treatment of U2OS cells

Puromycin (Sigma-Aldrich, St. Louis, MO) in water was added to the cell growth medium of U2OS cells to a

final concentration of 100 µg/mL. The cells were cultured with the drug for approximately 1 h before the start

of measurements. For z-scans, the cell growth medium was exchanged for PBS containing the same

concentration of puromycin.

Latrunculin A (LA) treatment of HeLa cells

LA (Cayman Chemical, Ann Arbor, MI) in DMSO was added to the cell growth medium of HeLa cells to a

concentration of 0.15 µg/mL. The cells were cultured with the drug for approximately 1 h before the start of

measurements. The experimental LA concentration corresponded to a DMSO concentration of 0.1 %. For

z-scans, the cell growth medium was exchanged for PBS containing the same concentration of LA.

Loading of labeled dextran in U2OS cells

U2OS cells previously transfected with mCherry were trypsinized from a T-25 flask near confluency (~ 3×10^6

cells total) and pelleted via low-speed centrifugation. Cells were resuspended in 30 µL PBS. Dextran

fluorescein (500 kDa MW, Life Technologies Corp. Eugene, OR) was diluted in PBS to a concentration of

10 mg/mL. A volume of 30 µL of this dextran solution was added to the cells. Next, cells were electroporated

using the MammoZapper electroporation system (Tritech Research, Inc. Los Angeles, CA) according to the

manufacturer's recommended protocol. To remove excess dextran, 1 mL growth medium was added to the

cells, which was followed by pelleting at low-speed centrifugation. This wash step was repeated a total of

three times. Cells were then plated into 8-well chambered slides and incubated for 24 h. The cell growth

medium was exchanged with PBS immediately prior to measurement.

Imaging of TetraSpeck beads

0.1 µm TetraSpeck beads (Thermo Fisher Scientific, Waltham, MA) were diluted in water and immobilized on imaging slides by addition of PBS buffer followed by quick washing with clean buffer. For measurement, an isolated bead was selected and a z-stack of 200 XY images was acquired with z-steps of 50 nm. Each XY image (256x256 pixels) was taken with a pixel resolution of 26.6 nm.

SYTO RNASelect cell stain

Live cells were stained with SYTO RNASelect (Thermo Fisher Scientific, Waltham, MA) by diluting the stock dye 1000x (5 µM final concentration) into the cell culture medium followed by incubation for 1 h at 37 ^{o}C and 5% CO_2. For measurement, the growth medium with SYTO RNASelect was replaced by PBS containing the same concentration of the dye.

Z-scan redaction and statistical analysis

Despite the selection of scan locations as described earlier, some of the measured z-scan intensity profiles were not adequately described by our fit model. This was attributed to scans that inadvertently passed through complex intracellular structures. Since these structures are not captured by the model framework, the spatial autocorrelation of the normalized residual from the fit was calculated to identify patterns that are indicative of the presence of extraneous cellular features not described by the fit model. Scans with a correlation amplitude exceeding a threshold value were removed. Furthermore, z-scan measurements of the cortex thickness were selected to have a cell thickness $L_{cell} \geq 2.5$ µm and a membrane fraction (*22*) $m \geq$ 0.5 in order to ensure the absence of biases in the fitted values as supported by our study using simulated DC z-scans (Fig. S4). Quality control for the z-scan fits of cortex thickness was provided by averaging the magnitude of the normalized residuals within ±1 µm of the fitted PM locations. Scans where these averaged residuals exceeded 2.5 in either the red or green channel were redacted.

A typical 60 s z-scan acquisition at a single location contained 12 repeated scans, which were fit individually, yielding 12 estimates of each parameter of interest. This set of parameters served to calculate the average effective depletion length Δ_i and its standard deviation σ_i at the *i*-th z-scan location. An average over multiple measurement locations, denoted by angle brackets, was estimated by the weighted

631 average $\langle\Delta\rangle$ over all locations with $1/\sigma_i^2$ as weight. The uncertainty in the weighted average was computed

632 using the dispersion corrected standard error of the mean (SEM),

633
$$SEM^2 = \sigma_{we}^2 \left(\frac{1}{n-1} \sum_i \frac{(\Delta_i - \langle\Delta\rangle)^2}{\sigma_i^2} \right)$$
14

where σ_{we}^2 is the variance of the weighted average. Since it was observed that $\langle\Delta\rangle$ at the top (T) and bottom

(B) PM interfaces are in agreement ($\langle\Delta^T\rangle \approx \langle\Delta^B\rangle$), data from both interfaces were grouped together when

calculating $\langle\Delta\rangle$. This is indicated by dropping the *T* and *B* superscript. Note that when datasets are grouped

in this manner, estimates of Δ^T and Δ^B are considered as independent, which is reflected by a doubling of

the *n* value when calculating SEM.

Simulations of z-scan traces

Simulated DC z-scan traces were obtained following a previously described procedure (*20*) to evaluate

models and estimate the statistical uncertainty in DC z-scan fit results. Simulated DC z-scan traces were fit

using the same algorithms that were applied to experimental data. All data derived from simulations

represent the average result from fits of *n* = 1000 z-scan traces.

References

1.   P. Chugh, E. K. Paluch, The actin cortex at a glance. *J Cell Sci*. **131** (2018), doi:10.1242/jcs.186254.

2.   A. Kusumi, T. K. Fujiwara, R. Chadda, M. Xie, T. A. Tsunoyama, Z. Kalay, R. S. Kasai, K. G. N. Suzuki,
Dynamic organizing principles of the plasma membrane that regulate signal transduction: commemorating the
fortieth anniversary of Singer and Nicolson's fluid-mosaic model. *Annu Rev Cell Dev Biol*. **28**, 215–250
(2012).

3.   M. D. Blower, Molecular insights into intracellular RNA localization. *Int Rev Cell Mol Biol*. **302**, 1–39 (2013).

4.   C. Medioni, K. Mowry, F. Besse, Principles and roles of mRNA localization in animal development.
Development. **139**, 3263–3276 (2012).

5.   E. Lécuyer, H. Yoshida, N. Parthasarathy, C. Alm, T. Babak, T. Cerovina, T. R. Hughes, P. Tomancak, H. M.
Krause, Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and
function. *Cell*. **131**, 174–187 (2007).

6.   A. Gopal, Z. H. Zhou, C. M. Knobler, W. M. Gelbart, Visualizing large RNA molecules in solution. *RNA*. **18**,
284–299 (2012).

7.   A. Borodavka, S. W. Singaram, P. G. Stockley, W. M. Gelbart, A. Ben-Shaul, R. Tuma, Sizes of Long RNA
Molecules Are Determined by the Branching Patterns of Their Secondary Structures. *Biophys J*. **111**, 2077–
2085 (2016).

8.   S. S. Sommer, J. E. Cohen, The size distributions of proteins, mRNA, and nuclear RNA. *J Mol Evol*. **15**, 37–57
(1980).

9.   I. Legnini, J. Alles, N. Karaiskos, S. Ayoub, N. Rajewsky, FLAM-seq: full-length mRNA sequencing reveals
principles of poly(A) tail length control. *Nat Methods*. **16**, 879–886 (2019).

10.   M. F. Tam, J. A. Dodd, W. E. Hill, Physical characteristics of 16 S rRNA under reconstitution conditions. *J
Biol Chem*. **256**, 6430–6434 (1981).

11.   A. Verschoor, J. Frank, Three-dimensional structure of the mammalian cytoplasmic ribosome. *J Mol Biol*. **214**,
737–749 (1990).

12.   G. T. Charras, T. J. Mitchison, L. Mahadevan, Animal cell hydraulics. *J Cell Sci*. **122**, 3233–3241 (2009).

13.   E. Moeendarbary, L. Valon, M. Fritzsche, A. R. Harris, D. A. Moulding, A. J. Thrasher, E. Stride, L.
Mahadevan, G. T. Charras, The cytoplasm of living cells behaves as a poroelastic material. *Nat Mater*. **12**,
253–261 (2013).

14.   K. Luby-Phelps, P. E. Castle, D. L. Taylor, F. Lanni, Hindered diffusion of inert tracer particles in the
cytoplasm of mouse 3T3 cells. *Proc Natl Acad Sci U S A*. **84**, 4910–4913 (1987).

15.   K. Luby-Phelps, D. L. Taylor, Subcellular compartmentalization by local differentiation of cytoplasmic
structure. *Cell Motil Cytoskeleton*. **10**, 28–37 (1988).

16.   L. Hou, F. Lanni, K. Luby-Phelps, Tracer diffusion in F-actin and Ficoll mixtures. Toward a model for
cytoplasm. *Biophys J*. **58**, 31–43 (1990).

17.   G. L. Lukacs, P. Haggie, O. Seksek, D. Lechardeur, N. Freedman, A. S. Verkman, Size-dependent DNA
mobility in cytoplasm and nucleus. *J Biol Chem*. **275**, 1625–1629 (2000).

18.   E. Dauty, A. S. Verkman, Actin cytoskeleton as the principal determinant of size-dependent DNA mobility in
cytoplasm: a new barrier for non-viral gene delivery. *J Biol Chem*. **280**, 7823–7828 (2005).

19.   K. Luby-Phelps, Effect of cytoarchitecture on the transport and localization of protein synthetic machinery.
Journal of Cellular Biochemistry. **52**, 140–147 (1993).

20.   I. Angert, S. R. Karuka, J. Hennen, Y. Chen, J. P. Albanesi, L. M. Mansky, J. D. Mueller, Sensitive Detection
of Protein Binding to the Plasma Membrane with Dual-Color Z-Scan Fluorescence. *Biophys J*. **118**, 281–293
(2020).

21.   E. M. Smith, P. J. Macdonald, Y. Chen, J. D. Mueller, Quantifying protein-protein interactions of peripheral
membrane proteins by fluorescence brightness analysis. *Biophys J*. **107**, 66–75 (2014).

22.   E. M. Smith, J. Hennen, Y. Chen, J. D. Mueller, Z-scan fluorescence profile deconvolution of cytosolic and
membrane-associated protein populations. *Anal Biochem*. **480**, 11–20 (2015).

23.   Y. N. Friew, V. Boyko, W.-S. Hu, V. K. Pathak, Intracellular interactions between APOBEC3G, RNA, and
HIV-1 Gag: APOBEC3G multimerization is dependent on its association with RNA. *Retrovirology*. **6**, 56
(2009).

24.   Y. Iwatani, H. Takeuchi, K. Strebel, J. G. Levin, Biochemical activities of highly purified, catalytically active
human APOBEC3G: correlation with antiviral effect. *J Virol*. **80**, 5992–6002 (2006).

25.   S. Saha, I.-H. Lee, A. Polley, J. T. Groves, M. Rao, S. Mayor, Diffusion of GPI-anchored proteins is influenced
by the activity of dynamic cortical actin. *Mol Biol Cell*. **26**, 4033–4045 (2015).

701 26. T. Baumgart, A. T. Hammond, P. Sengupta, S. T. Hess, D. A. Holowka, B. A. Baird, W. W. Webb, Large-scale
702   fluid/fluid phase separation of proteins and lipids in giant plasma membrane vesicles. *Proc Natl Acad Sci U S*
703   *A*. **104**, 3165–3170 (2007).

704 27. F. Schneider, D. Waithe, M. P. Clausen, S. Galiani, T. Koller, G. Ozhan, C. Eggeling, E. Sezgin, Diffusion of
705   lipids and GPI-anchored proteins in actin-free plasma membrane vesicles measured by STED-FCS. *Mol Biol*
706   *Cell*. **28**, 1507–1518 (2017).

707 28. T. M. Svitkina, Actin Cell Cortex: Structure and Molecular Organization. *Trends Cell Biol*. **30**, 556–565
708   (2020).

709 29. N. Morone, T. Fujiwara, K. Murase, R. S. Kasai, H. Ike, S. Yuasa, J. Usukura, A. Kusumi, Three-dimensional
710   reconstruction of the membrane skeleton at the plasma membrane interface by electron tomography. *J Cell*
711   *Biol*. **174**, 851–862 (2006).

712 30. P. Chugh, A. G. Clark, M. B. Smith, D. A. D. Cassani, K. Dierkes, A. Ragab, P. P. Roux, G. Charras, G.
713   Salbreux, E. K. Paluch, Actin cortex architecture regulates cell surface tension. *Nat Cell Biol*. **19**, 689–697
714   (2017).

715 31. A. G. Clark, K. Dierkes, E. K. Paluch, Monitoring actin cortex thickness in live cells. *Biophys J*. **105**, 570–580
716   (2013).

717 32. J. Riedl, A. H. Crevenna, K. Kessenbrock, J. H. Yu, D. Neukirchen, M. Bista, F. Bradke, D. Jenne, T. A.
718   Holak, Z. Werb, M. Sixt, R. Wedlich-Soldner, Lifeact: a versatile marker to visualize F-actin. *Nat Methods*. **5**,
719   605–607 (2008).

720 33. P. J. Macdonald, Y. Chen, X. Wang, Y. Chen, J. D. Mueller, Brightness analysis by Z-scan fluorescence
721   fluctuation spectroscopy for the study of protein interactions within living cells. *Biophys J*. **99**, 979–988
722   (2010).

723 34. J. G. Levin, J. Guo, I. Rouzina, K. Musier-Forsyth, Nucleic acid chaperone activity of HIV-1 nucleocapsid
724   protein: critical role in reverse transcription and molecular mechanism. *Prog Nucleic Acid Res Mol Biol*. **80**,
725   217–286 (2005).

726 35. R. N. De Guzman, Z. R. Wu, C. C. Stalling, L. Pappalardo, P. N. Borer, M. F. Summers, Structure of the HIV-
727   1 nucleocapsid protein bound to the SL3 psi-RNA recognition element. *Science*. **279**, 384–388 (1998).

728 36. A. Rein, Nucleic acid chaperone activity of retroviral Gag proteins. *RNA Biol*. **7**, 700–705 (2010).

729 37. V. Chukkapalli, S. J. Oh, A. Ono, Opposing mechanisms involving RNA and lipids regulate HIV-1 Gag
730   membrane binding through the highly basic region of the matrix domain. *Proc Natl Acad Sci U S A*. **107**, 1600–
731   1605 (2010).

732 38. S. B. Kutluay, T. Zang, D. Blanco-Melo, C. Powell, D. Jannain, M. Errando, P. D. Bieniasz, Global changes in
733   the RNA binding specificity of HIV-1 gag regulate virion genesis. *Cell*. **159**, 1096–1109 (2014).

734 39. J. Li, Y. Chen, M. Li, M. A. Carpenter, R. M. McDougle, E. M. Luengas, P. J. Macdonald, R. S. Harris, J. D.
735   Mueller, APOBEC3 multimerization correlates with HIV-1 packaging and restriction activity in living cells. *J*
736   *Mol Biol*. **426**, 1296–1307 (2014).

737 40. K. Luby-Phelps, D. L. Taylor, F. Lanni, Probing the structure of cytoplasm. *J Cell Biol*. **102**, 2015–2022
738   (1986).

739 41. B. R. Cullen, Role and mechanism of action of the APOBEC3 family of antiretroviral resistance factors. *J*
740   *Virol*. **80**, 1067–1076 (2006).

741 42. J. S. Albin, R. S. Harris, Interactions of host APOBEC3 restriction factors with HIV-1 in vivo: implications for
742   therapeutics. *Expert Rev Mol Med*. **12**, e4 (2010).

43.   J. F. Hultquist, J. A. Lengyel, E. W. Refsland, R. S. LaRue, L. Lackey, W. L. Brown, R. S. Harris, Human and
rhesus APOBEC3D, APOBEC3F, APOBEC3G, and APOBEC3H demonstrate a conserved capacity to restrict
Vif-deficient HIV-1. *J Virol*. **85**, 11220–11234 (2011).

44.   L. W. Janson, K. Ragsdale, K. Luby-Phelps, Mechanism and size cutoff for steric exclusion from actin-rich
cytoplasmic domains. *Biophys J*. **71**, 1228–1234 (1996).

45.   A. P. Minton, Confinement as a determinant of macromolecular structure and reactivity. *Biophys J*. **63**, 1090–
1100 (1992).

46.   J. C. Giddings, E. Kucera, C. P. Russell, M. N. Myers, Statistical theory for the equilibrium distribution of rigid
molecules in inert porous networks. Exclusion chromatography. *J. Phys. Chem.* **72**, 4397–4408 (1968).

47.   T. Kalwarczyk, N. Ziebacz, A. Bielejewska, E. Zaboklicka, K. Koynov, J. Szymański, A. Wilk, A. Patkowski,
J. Gapiński, H.-J. Butt, R. Hołyst, Comparative analysis of viscosity of complex liquids and cytoplasm of
mammalian cells at the nanoscale. *Nano Lett*. **11**, 2157–2163 (2011).

48.   R. Holyst, A. Bielejewska, J. Szymański, A. Wilk, A. Patkowski, J. Gapiński, A. Zywociński, T. Kalwarczyk,
E. Kalwarczyk, M. Tabaka, N. Ziebacz, S. A. Wieczorek, Scaling form of viscosity at all length-scales in
poly(ethylene glycol) solutions studied by fluorescence correlation spectroscopy and capillary electrophoresis.
Phys Chem Chem Phys. **11**, 9025–9032 (2009).

49.   L. Xiang, K. Chen, R. Yan, W. Li, K. Xu, Single-molecule displacement mapping unveils nanoscale
heterogeneities in intracellular diffusivity. *Nat Methods*. **17**, 524–530 (2020).

50.   T. P. Stossel, P. A. Janmey, K. S. Zaner, in *Cytomechanics: The Mechanical Basis of Cell Form and Structure*,
J. Bereiter-Hahn, O. R. Anderson, W.-E. Reif, Eds. (Springer, Berlin, Heidelberg, 1987;
https://doi.org/10.1007/978-3-642-72863-1_9), pp. 131–153.

51.   J.-F. Joanny, J. Prost, Active gels as a description of the actin-myosin cytoskeleton. *HFSP J*. **3**, 94–104 (2009).

52.   T. C. Laurent, J. Killander, A theory of gel filtration and its exeperimental verification. *Journal of
Chromatography A*. **14**, 317–330 (1964).

53.   K. C. Duong-Ly, S. B. Gabelli, Gel filtration chromatography (size exclusion chromatography) of proteins.
Methods Enzymol. **541**, 105–114 (2014).

54.   T. Kaur, I. Alshareedah, W. Wang, J. Ngo, M. M. Moosa, P. R. Banerjee, Molecular Crowding Tunes Material
States of Ribonucleoprotein Condensates. *Biomolecules*. **9** (2019), doi:10.3390/biom9020071.

55.   A. A. M. André, E. Spruijt, Liquid-Liquid Phase Separation in Crowded Environments. *Int J Mol Sci*. **21**
(2020), doi:10.3390/ijms21165908.

56.   D. W. Provance, A. McDowall, M. Marko, K. Luby-Phelps, Cytoarchitecture of size-excluding compartments
in living cells. *J Cell Sci*. **106 (Pt 2)**, 565–577 (1993).

57.   R. Kumar, S. Saha, B. Sinha, Cell spread area and traction forces determine myosin-II-based cortex thickness
regulation. *Biochim Biophys Acta Mol Cell Res*. **1866**, 118516 (2019).

58.   T. M. Svitkina, A. A. Shevelev, A. D. Bershadsky, V. I. Gelfand, Cytoskeleton of mouse embryo fibroblasts.
Electron microscopy of platinum replicas. *Eur J Cell Biol*. **34**, 64–74 (1984).

59.   G. Blobel, D. Sabatini, Dissociation of mammalian polyribosomes into subunits by puromycin. *Proc Natl Acad
Sci U S A*. **68**, 390–394 (1971).

60.   M. Delarue, G. P. Brittingham, S. Pfeffer, I. V. Surovtsev, S. Pinglay, K. J. Kennedy, M. Schaffer, J. I.
Gutierrez, D. Sang, G. Poterewicz, J. K. Chung, J. M. Plitzko, J. T. Groves, C. Jacobs-Wagner, B. D. Engel, L.
J. Holt, mTORC1 Controls Phase Separation and the Biophysical Properties of the Cytoplasm by Tuning
Crowding. *Cell*. **174**, 338-349.e20 (2018).

61.   K. Kwapiszewska, K. Szczepański, T. Kalwarczyk, B. Michalska, P. Patalas-Krawczyk, J. Szymański, T.
Andryszewski, M. Iwan, J. Duszyński, R. Hołyst, Nanoscale Viscosity of Cytoplasm Is Conserved in Human
Cell Lines. *J Phys Chem Lett*. **11**, 6914–6920 (2020).

62.   J. Chen, D. Grunwald, L. Sardo, A. Galli, S. Plisov, O. A. Nikolaitchik, D. Chen, S. Lockett, D. R. Larson, V.
K. Pathak, W.-S. Hu, Cytoplasmic HIV-1 RNA is mainly transported by diffusion in the presence or absence of
Gag protein. *Proc Natl Acad Sci U S A*. **111**, E5205-5213 (2014).

63.   J. J. Blum, G. Lawler, M. Reed, I. Shin, Effect of cytoskeletal geometry on intracellular diffusion. *Biophys J*.
56, 995–1005 (1989).

64.   K. Luby-Phelps, F. Lanni, D. L. Taylor, The submicroscopic properties of cytoplasm as a determinant of
cellular function. *Annu Rev Biophys Biophys Chem*. **17**, 369–396 (1988).

65.   D. S. Banks, C. Fradin, Anomalous diffusion of proteins due to molecular crowding. *Biophys J*. **89**, 2960–2971
(2005).

66.   A. Miermont, F. Waharte, S. Hu, M. N. McClean, S. Bottani, S. Léon, P. Hersen, Severe osmotic compression
triggers a slowdown of intracellular signaling, which can be explained by molecular crowding. *Proc Natl Acad
Sci U S A*. **110**, 5725–5730 (2013).

67.   M. Weiss, M. Elsner, F. Kartberg, T. Nilsson, Anomalous subdiffusion is a measure for cytoplasmic crowding
in living cells. *Biophys J*. **87**, 3518–3524 (2004).

68.   C. I. Lacayo, J. A. Theriot, Listeria monocytogenes actin-based motility varies depending on subcellular
location: a kinematic probe for cytoarchitecture. *Mol Biol Cell*. **15**, 2164–2175 (2004).

69.   J.-Y. Youn, B. J. A. Dyakov, J. Zhang, J. D. R. Knight, R. M. Vernon, J. D. Forman-Kay, A.-C. Gingras,
Properties of Stress Granule and P-Body Proteomes. *Mol Cell*. **76**, 286–294 (2019).

70.   A. A. Hyman, C. A. Weber, F. Jülicher, Liquid-liquid phase separation in biology. *Annu Rev Cell Dev Biol*. **30**,
39–58 (2014).

71.   L.-P. Bergeron-Sandoval, N. Safaee, S. W. Michnick, Mechanisms and Consequences of Macromolecular
Phase Separation. *Cell*. **165**, 1067–1079 (2016).

72.   R. Wollman, T. Meyer, Coordinated oscillations in cortical actin and Ca2+ correlate with cycles of vesicle
secretion. *Nat Cell Biol*. **14**, 1261–1269 (2012).

73.   P. Li, A. T. Bademosi, J. Luo, F. A. Meunier, Actin Remodeling in Regulated Exocytosis: Toward a
Mesoscopic View. *Trends Cell Biol*. **28**, 685–697 (2018).

74.   E. D. Olson, K. Musier-Forsyth, Retroviral Gag protein-RNA interactions: Implications for specific genomic
RNA packaging and virion assembly. *Semin Cell Dev Biol*. **86**, 129–139 (2019).

75.   A. Derdowski, L. Ding, P. Spearman, A novel fluorescence resonance energy transfer assay demonstrates that
the human immunodeficiency virus type 1 Pr55Gag I domain mediates Gag-Gag interactions. *J Virol*. **78**,
1230–1242 (2004).

76.   C. Thaler, S. V. Koushik, H. L. Puhl, P. S. Blank, S. S. Vogel, Structural rearrangement of CaMKIIalpha
catalytic domains encodes activation. *Proc Natl Acad Sci U S A*. **106**, 6369–6374 (2009).

77.   B. Wu, Y. Chen, J. D. Müller, Fluorescence fluctuation spectroscopy of mCherry in living cells. *Biophys J*. **96**,
2391–2404 (2009).

78.   B. Wu, Y. Chen, J. D. Müller, Heterospecies partition analysis reveals binding curve and stoichiometry of
protein interactions in living cells. *Proc Natl Acad Sci U S A*. **107**, 4117–4122 (2010).

79.   Y. Chen, H.-Q. Sun, J. P. Eichorst, J. P. Albanesi, H. Yin, J. D. Mueller, Comobility of GABARAP and
Phosphatidylinositol 4-Kinase 2A on Cytoplasmic Vesicles. *Biochemistry*. **57**, 3556–3559 (2018).

80.   J. D. Müller, Y. Chen, E. Gratton, Fluorescence correlation spectroscopy. *Methods Enzymol*. **361**, 69–92
828   (2003).

829   81.   T. Kalwarczyk, K. Kwapiszewska, K. Szczepanski, K. Sozanski, J. Szymanski, B. Michalska, P. Patalas-
830   Krawczyk, J. Duszynski, R. Holyst, Apparent Anomalous Diffusion in the Cytoplasm of Human Cells: The
831   Effect of Probes' Polydispersity. *J Phys Chem B*. **121**, 9831–9837 (2017).

Acknowledgments

Non-author contributions: The authors thank Yan Chen for help with logistics and experiment planning.

Author contributions: CIA, JDM, and SRK developed the dual color z-scan method. CIA and SRK performed experiments and acquired the data. LMM helped interpret the data and guide the project. CIA and JDM wrote analysis code, performed data analysis, and drafted the manuscript. All authors reviewed and revised the manuscript.

Funding: This work was supported by NIH grants GM064589, GM124279 and GM098550. C. I. A. was supported by NIH T32 AI083196 and NIH T90 DE 022732

Competing interests: The authors declare that they have no financial or other competing interests.

Data and materials availability: Data and z-scan analysis code are available at Dryad: https://doi.org/10.1016/j.bpj.2019.12.002

Fig. 1: DC z-scans in living cells. The z-scan trajectory through the cell is depicted in the cartoons along with crosshatched regions indicating the subcellular localization of the EGFP- and mCherry-labeled proteins (A, D, G, J, left panel) together with the concentration profile indicated by z-scan fitting for the EGFP (green lines) and mCherry (magenta lines) protein species (A, D, G, J, right panel). B, E, H, K) Fits (solid lines) of representative z-scan intensity traces of the green (green diamonds) and red (magenta diamonds) detection channel are shown along with their normalized residuals sr. The cell edge locations a^G, b^G, a^{CH}, b^{CH} extracted from the fits determine the vertical distance Δ separating the EGFP- and mCherry-labeled slab edges in each z-scan. C, F, I, L) Histograms of the average Δ values estimated from each scan location in many cells. Experiments with different EGFP- and mCherry-labeled protein species are indicated in rows: (A, B, C) EGFP and mCherry-RXR, (D, E, F) EGFP and mCherry, (G, H, I) EGFP-HRas and mCherry, and (J, K, L) A3G-EGFP and mCherry.

Fig. 2: Z-scans in blebs produced from HeLa cells co-expressing A3G-EGFP and mCherry. A) A

graphic depicting the production of large PM blebs from adherent cells via ethanol treatment is shown. In

this example, the cytoskeleton is denoted by brown colors. B) A DC 2-photon laser scanning z-stack image

of a HeLa cell with two large PM blebs is shown. The x-y image was created by the summation of the z-

stack over the z-dimension. The y-z image was produced by the sum of the x-dimension over two volume-

sections passing roughly through the center of each bleb. While significant red autofluorescence is produced

by the ethanol treatment near the cell center, it is typically absent in the blebs. Z-scan locations were selected

to pass near the center of each bleb, while avoiding regions possessing autofluorescence or identifiable

structures. C) Fit of a representative z-scan passing through a HeLa cell bleb (χ_v^2= 1.2). D) Histogram of the

average Δ values estimated from each scan location in the cytoplasm of many HeLa cells not treated with

ethanol (magenta bars) and in blebs produced by ethanol treatment (gray bars).

Fig. 3: DC z-scan measurement of cortex thickness. A) Lifeact-mApple demarcates the actin cortex, and EGFP-HRas demarcates the PM location. These fluorescent proteins lead to the SSS^{CH}-δSδ^G concentration profile depicted in (B). C) DC z-scans were fit to a δSδ^{CH}-δSδ^G model. The location of the mApple δ layers corresponds to the centroid of the cortical slabs, so that the effective depletion length is half the thickness of the cortex slab. D) A fit of a DC z-scan through a HeLa cell co-expressing EGFP-HRas and Lifeact-mApple. E) Average cortex thickness values estimated by DC z-scan in HeLa and U2OS cells. F) <$Δ$> is plotted in U2OS cells versus the same vales in HeLa cells for MA^{G2A}-EGFP (brown), Gag^{G2A}-EGFP (green) and A3G-EGFP (magenta). The ratio of cortex thicknesses in the cell lines corresponds to the slope of the data as is indicated by the shaded area and dashed lines, which correspond to the slope-uncertainty derived from the data in panel (E).

Fig. 4 Effective depletion length. A) Average $\langle\Delta\rangle$ values in HeLa cells for the indicated proteins and an

RNA dye. B) Model of cortical partitioning leading to a SSS[G] distribution of fluorescence. C) Axial

concentration profile corresponding to the model. The concentration drops from c_{cyto} in the cytoplasm to c_C

in the cortex. D) Approximation of the SSS[G] by a single S[G] concentration profile. Fitting of the experimental

z-scan data to this profile was used to identify apparent depletion lengths Δ^B and Δ^T. The mCherry

concentration profile has been omitted in panels C and D. E) $\langle\Delta\rangle$ vs partition coefficient R_C extracted from

simulated DC z-scans for three different cortex lengths ($L_C^B = L_C^T$ of 200, 100, and 50 nm). The solid lines

represent a theoretical model based on Eq. 1.

Fig. 5: Axially averaged partition coefficient $\overline{R}_z$ vs cell thickness. A-D) Representative DC z-scan

traces in a thin and thick section of a HeLa cell co-expressing EGFP and mCherry (A, B) and representative

DC z-scan traces in thin and thick sections of a HeLa cell co-expressing Gag^{G2A}-EGFP and mCherry. E)

Schematic illustration of DC z-scans taken at multiple cytoplasmic locations varying from thick to thin

sections to determine $\overline{R}_z$ as a function of thickness L_{cell}. F) $\overline{R}_z$ vs L_{cell}. Black diamonds correspond to

control measurements in HeLa cells co-expressing EGFP and mCherry. Magenta and blue diamonds

correspond to data from HeLa and U2OS cells co-expressing HIV-1 Gag^{G2A}-EGFP and mCherry. Magenta

and blue lines correspond to Eq. S4 with $L_C = 200\,\mathrm{nm}$ and $L_C = 130\,\mathrm{nm}$, respectively, and $R_C = 0.2$.

Fig. 6: Protein mobility and effective depletion length. A) Depletion length *<Δ>* versus effective diffusion

coefficient D_{eff} in HeLa cells (diamonds). Eq. 4 (magenta curve) is plotted with parameter values as

described in the text. A3G-EGFP measured in HeLa cell blebs (green open circle) displayed increased

mobility along with decreased effective depletion length. B) Conceptual model of the effects quantified in

panel A. The partitioning effect (center left) arises due to hindered entry of large RNP complexes into the

cortical mesh, while hindered mobility of large RNP complexes due to macromolecular crowding occurs in

the cytoplasm (right). Binding of labeled proteins to partitioned RNP complexes yields the concentration

gradient (far left) observed by DC z-scan measurements. C) The cytoplasmic mobility and partition

coefficient are connected by the average RNP complex radius, as indicated in the graphical depiction and

suggested by Eqs. 3 and 4.