Stimulating music supports attention in listeners with attentional difficulties

Kevin JP Woods, Gonçalo Sampaio, Tedra James, Emily Przysinda, Adam Hewett, Andrea E Spencer, Benjamin Morillon, Psyche Loui

ABSTRACT

Background music is widely used to sustain attention, but little is known about what musical properties aid attention. This may be due to inter-individual variability in neural responses to music. We test the hypothesis that music can sustain attention by affecting oscillations via acoustic amplitude modulation, differentially for those with varying levels of attentional difficulty. We first show that heavily-modulated music improves sustained attention for participants with more ADHD symptoms. FMRI showed this music elicited greater activity in attentional networks in this group only, and EEG showed greater stimulus-brain coupling for this group in response to the heavily-modulated music. Finally, we parametrically manipulated the depth and rate of amplitude modulations inserted in otherwise-identical music, and found that beta-range modulations helped more than other frequency ranges for participants with more ADHD symptoms. Results suggest the possibility of an oscillation-based neural mechanism for targeted music to support improved cognitive performance.

Music often has practical uses beyond aesthetic appeal[1], and from mothers' lullabies to laborers' work-songs, the music we make to fill these roles reflects its function[2,3]. One possible use of music is to aid cognitive performance[4–6]. This has become increasingly important with the shift to knowledge-work[7–10], along with widespread adoption of technologies like streaming and personal audio. To date, many different kinds of music have been used to aid focus in the workplace[11]. The diversity in music used for focus may reflect individual differences in cognitive styles: for example, personality differences are associated with the ability to sustain attention[12,13,14]. Preference and familiarity also contribute to effects of music on cognition[15–17].

Another factor deserving special consideration is an individual's ability to focus. Prior work has shown that auditory stimulation can aid performance in individuals with ADHD[18–21]. This has been explained by optimal stimulation theory, which poses that some individuals, specifically those with ADHD, require more stimulation than others to function best[22–25]. However, all these cases compare stimulation (music or noise) to silence, and no studies of this kind to date have used experimental conditions with different types of music.

If people who have difficulty focusing have distinct needs for focus-music, there is a chance to provide a targeted solution for those who could use it most. We were thus interested to see if music with different levels of arousal would affect people differently depending on their attentional capacity. If so, people with attentional deficits, such as symptoms of ADHD, may need specifically-designed focus-music. We hypothesize that arousal in music can affect performance differently in people with different levels of attentional difficulties[19,21,26]. Specifically, we hypothesize that more stimulating (i.e. more arousing) music should benefit people who experience more attentional difficulties as quantified by the ADHD Self-Report Scale (ASRS); i.e.,

high-ASRS individuals should improve in sustained-attention performance over time with more
stimulating music.
We first compared performance (Experiment 1) on the Sustained Attention to Response
Task (SART[27,28]) under three types of background acoustic conditions: AM+Music (i.e. music with
fast amplitude modulations), Control-Music (with slow amplitude modulations), and Pink Noise.
The AM+Music had fast modulations added that do not usually occur in music, and acoustic
analyses (Figure 1) showed that despite similar frequency content, the tracks differed in the
modulation domain due to this added modulation. We then used the same stimuli and task in
experiments with fMRI (Experiment 2) and EEG (Experiment 3). Finally, additional behavioral
experiments (Experiments 4A and 4B) tested the effects of modulation on sustained attention in
an acoustically controlled manner.
56
57

58
59
Fig. 1 | Acoustic analyses of auditory stimuli used in Experiments 1-3. Analysis of a 30-
second excerpt from each stimulus type used in Experiments 1-3. Pressure over time (top row)
first undergoes frequency decomposition via cochlear filtering. The energy in each cochlear
channel varies over time (depicted on the cochleagram, 2nd row). These envelope fluctuations
are then frequency-decomposed to produce a modulation spectrum representation (3rd row).
The broadband modulation spectrum (bottom row) is the sum of modulation spectra across the
cochlear channels. This broadband modulation shows peak in the AM+Music condition only,
where rapid modulation was added to the music.
68

Results

Experiment 1: Sustained Attention to Response Task (SART)

Participants were recruited and tested online via Amazon's Mechanical Turk web service. The ASRS was obtained in all participants, and analyses were performed after a median split on ASRS score into high-ASRS and low-ASRS groups. In Experiment 1A, 62 participants (31 high-ASRS and 31 low-ASRS) rated the AM+Music, Control-Music, and Pink Noise for valence and arousal. In Experiment 1B, another 87 participants (39 high-ASRS and 48 low-ASRS) completed the SART under the three acoustic conditions.

a   **Valence and arousal ratings for stimuli in Experiments 1-3**

b   **Experiment 1: Performance over time in sustained attention task**

Fig. 2 | Stimulus valence and arousal ratings and performance over time in the Sustained Attention to Response Task (SART). a, Valence and Arousal ratings for the music used in Experiments 1-3, with listeners split by their level of self-reported attentional difficulty (ASRS score, median split). N=62 overall, N=31 in each group. **b,** Performance on the SART in a separate group of participants by music condition and by ASRS score. Each participant completed

84 3 blocks (the music conditions) presented in random order. N=87 overall, N=48 and N=39 in the
85 Low- and High-ASRS groups respectively. Error bars depict +/- 1 standard error of the mean
86 (within-subject).
87
88
Valence and arousal ratings for all three sound stimuli were entered into a multivariate
two-factor mixed ANOVA, with the within-subjects factor of Music (3 levels: AM+Music, Control-
Music, and Pink Noise) and the between-subjects factor of ASRS (high-ASRS vs. low-ASRS
groups). A main effect of Music was observed on both valence and arousal (valence:
$F(2,120)=105, p<.001$; arousal: $F(2,120)=45, p<.001$). A main effect of ASRS was significant for
arousal ($F(1,60)=6.0, p=.017$) but not for valence ($F(1,60)=0.166$, n.s.). Participants rated
AM+Music as positive in valence and high in arousal, Control-Music as positive in valence and
low in arousal, and Pink Noise as low in valence but high in arousal. High-ASRS participants
additionally rated the two music conditions (AM+ and Control-Music) as more stimulating (higher
in arousal) than their low-ASRS counterparts.
Since the SART is a test of sustained attention over time, performance on the SART over
time was analyzed with a 3-factor mixed ANOVA incorporating time as a within-subjects factor (2
levels: early vs. late trials), Music as a within-subjects factor (3 levels: AM+Music, Control-Music,
and Pink Noise), and ASRS group as a between-subjects factor (2 levels: high- vs. low-ASRS).
Overall performance (d-prime) in the two groups was not different (High-ASRS: mean = 3.50,
stdv=0.71; Low-ASRS mean = 3.48, stdv=0.75). A three-way interaction was significant
($F(2,170)=5.02$, $p=0.008$): the music affected high-ASRS and low-ASRS participants differently
over time, with high-ASRS participants showing an improvement in d' over time during AM+Music
but not during Control-Music. In contrast, low-ASRS participants showed an improvement over
time during Control-Music but not AM+Music, despite no main effect of ASRS group
($F(1,85)=0.01$, $p=0.92$) and no main effect of music ($F(2,170)=0.02$, $p=0.98$). Since these two
stimuli were rated as different in arousal but not in valence by both groups, differences in arousal
may be related to these different patterns of SART performance over time.
The direction of this interaction--with fast modulations benefiting high-ASRS listeners in
particular--aligns with our hypothesis that arousal in music can affect sustained attention
differently in people because the optimal level of stimulation is greater for those with attentional
deficits. To better understand the neural bases of this effect and how this might differ across the
ASRS groups, we ran the same task and background conditions in neuroimaging experiments,
with fMRI (Experiment 2) and EEG (Experiment 3), looking not only at the brain's response to the
different types of music, but also task-related activity.

119 Experiment 2: SART fMRI During Background Music

In an fMRI study, 34 participants completed the SART under the same three background music
conditions used in Experiment 1: AM+Music, Control-Music, and Pink Noise.
A within-subjects ANOVA comparing overall brain activity during the three conditions
showed significantly higher activation during the AM+Music condition than in the other two
conditions ($p<.05$ FDR-corrected) in multiple regions including the bilateral superior temporal
lobes, frontal lobes, parietal lobes, and mesial and lateral occipital cortices, encompassing the
default mode, executive function, and salience networks (Figure 3A). No other contrast showed
positive suprathreshold clusters (all $p>.05$ FDR-corrected).
128

a   Effect of AM+Music

Voxel-level p<.05 FDR-corrected
Cluster-level p<.05 FDR-corrected

AM+Music > (Control-music + Pink Noise)/2

b   ASRS x Music Interaction

p<.05 FDR-corrected

AM+Music > (Control-music + Pink Noise)/2
x high-ASRS > low-ASRS individuals

Fig. 3 | fMRI results comparing AM+Music, Control-Music, and Pink noise during the Sustained Attention to Response Task (SART). a, Contrast between AM+Music and the average of Control-Music and Pink Noise while participants performed a sustained attention (SART) task in Experiment 2. Higher levels of activity for AM+Music are widespread across many regions. **b,** Interaction between group (high- minus low-ASRS) and Music (AM+Music minus other conditions) was significant in the midcingulate (sometimes referred to as dorsal anterior cingulate; implicated in reward-mediated responding), anterior insula and ventral parietal areas (classic regions of the salience network), and middle frontal regions and anterior temporal poles. All results are significant at the p < .05 FDR cluster-corrected level.

Individual differences in self-reported attention difficulties.

When we divided the participants by ASRS (median split, as in Experiment 1) and compared their main effects of AM+Music (AM+Music > (Control-Music + Pink Noise) contrast), both participant groups showed activity in occipital and medial parietal regions. The interaction between music and group showed higher activity in the high-ASRS group than in the low-ASRS group during AM+Music compared to the other listening conditions. This interaction was significant in the insula and the anterior cingulate cortex, regions in the salience network[29], with high-ASRS group showing larger differences than low-ASRS group in these regions (Figure 3B). Additional areas that survived cluster-wise FDR correction included the bilateral middle frontal gyri and frontal operculum, medial prefrontal cortex, bilateral temporal lobes, and lateral occipital cortex. The

152 latter regions are part of the default network and the ventral attention network[30], and their
153 additional involvement is consistent with the role of the salience network in facilitating attention
154 resources and accessing the motor system upon the detection of salient events.
155

Hits > False Alarms

Voxel-level p<.05 FDR-corrected
Cluster-level p<.05 FDR-corrected

156
Fig. 4 | fMRI results comparing Hits and False Alarm trials from the Sustained Attention to
Response Task (SART) during AM+Music, Control-Music, and Pink Noise. Brain activity
during Hits contrasted against False Alarms on the SART showing increased activity during
correct trials despite similar motor output, centering around the sensorimotor network and the
salience network during AM+Music compared to Control-Music and Pink Noise. Warm colors
show greater activity for Hits; cool colors show greater activity for False Alarms.
163

Higher activity in motor network linked to successful behavior during AM+Music

To relate behavior to brain activity during the different music tracks, we fit separate parametric
models in SPM12[31] for hit trials and false alarm trials for each auditory condition. Brain activity
during successful responses, quantified as a contrast between activity during hits and activity
during false alarms, showed significantly higher activity during the hits in all auditory conditions
(**Figure 4**). Importantly, the Hits vs. FA contrast at the $p < .05$ FDR-corrected level showed more
significant clusters during AM+Music than during any other condition (Control-Music, Pink Noise).
These clusters centered around the sensorimotor network (supplementary motor area (SMA),

precentral gyrus (PCG), the salience network (anterior cingulate cortex, anterior insula), and the
visual association network (lateral and mesial occipital regions). These differences were observed
despite overall similar hit and FA rates across the auditory conditions (one-way ANOVAs all Fs <
1). While the interaction between auditory conditions and groups was not significant at the whole-
brain level, both high-ASRS and low-ASRS groups showed more activity for AM+Music than for
the other auditory conditions during Hit trials compared to FA trials (Supplementary Materials).
Together, these results show that AM+Music is linked to higher levels of brain activity in multiple
networks, especially during successful behavioral performance of a sustained attention task.
Furthermore, differences in brain activity during AM+Music listening (under Hits>FAs) were
observed between people who report high and low levels of attentional difficulties, with more
activity observed in (which regions) in the high-ASRS group. These individual differences in brain
activity could underlie group differences in performance from Experiment 1.
Since the musical stimuli were highly rhythmic, we expected that they might affect rhythmic
activity in the brain, and that group differences in such activity could provide further insight into
the mechanisms by which music affects sustained attention. To capture rhythmic neural activity
and relate it to stimulus rhythms with high temporal precision, we turned to an EEG study with the
same stimulus and task conditions as the fMRI study reported above.

190   Experiment 3: SART EEG During Background Music

191   Forty participants had their EEG recorded while they performed the SART task under the same
192   three background music conditions used in Experiments 1 and 2: AM+Music (containing fast
193   modulation rates; more arousing), Control-Music (containing slow modulation rates; less
194   arousing), and Pink Noise. Hit and FA rates were overall similar across the auditory conditions
195   (one-way ANOVAs all Fs < 1).
196

197   Stimulus-brain coupling EEG shows phase locking at peak frequencies of
198   amplitude modulation

We assessed coupling between the EEG and the acoustic signal by computing the stimulus-brain
phase-locking value (PLV) for every frequency (in 1-Hz bins). During the AM+Music, stimulus-
brain PLV showed prominent peaks at 8, 12, 14, 16, 24, and 32 Hz (Figure 5A). Since the
AM+Music was at 120bpm (i.e., quarter-notes at 2Hz), 8, 16, 24, and 32 Hz are harmonics of the
note rate, previously observed to entrain cortical activity[32], while 12, 14, and 16 Hz reflect the
amplitude-modulated frequencies.
PLV at 8 Hz during AM+Music was strongest at frontal recording sites. In contrast, PLV
during Control-Music was much lower and less sharply tuned, reflecting less focused neural
tracking of acoustic rhythms when listening to Control-Music. Looking across the whole brain
(Figure 5B), PLV at 8Hz was stronger in the AM+Music condition, even over the same frontal
recording sites. We computed an effect size (Cohen's d) measure for every 4-Hz bin (thus
capturing 8, 12, 16, 20, 24, 28, 32 Hz) across the full frequency spectrum, comparing AM+Music
against Control-Music, averaging across all frontal, central, and parietal electrodes. This resulted
in a Cohen's d of 3.74, confirming a highly statistically significant difference between AM+Music
and Control-Music at these frequencies of interest. In contrast, the same analysis for 1-Hz bins
showed a Cohen's d of 0.227, suggesting higher stimulus-brain coupling at multiples of 4-Hz,
consistent with the note rate and with the amplitude-modulation rates in the music.

Fig. 5 | EEG reveals stimulus-brain coupling. a, For Experiment 3, we conducted hertz-by-hertz Morlet wavelet analysis of each acoustic stimulus (amplitude-modulated and control) and evaluated the phase-locking value (stimulus-brain coupling) between acoustics of the stimulus and the EEG. Red trace = Mean±SE Phase Locking Values (PLVs) of frontocentral electrodes for AM+Music. Black trace = Mean±SE PLVs of the same electrodes for the Control-Music. Results show peaks of phase-locking activity at the note rates and its harmonics (8, 16, 24, 32 Hz) as well as the amplitude-modulated frequencies (12, 14, 16 Hz). **b,** Topographic distributions of the peaks of phase-locked activity at 8, 16, and 32 Hz for amplitude-modulated and control stimuli.

Phase-locking across ASRS groups

We compared stimulus-brain PLV between high-ASRS and low-ASRS groups. The high-ASRS group showed higher PLV in high frequencies, especially during the AM+Music condition (Figure 6A). This pattern of higher PLV in high-ASRS participants was mainly observed during the second half of the AM+Music sessions (Figure 6c-d), suggesting that the effects of AM+Music emerge over time, consistent with the behavioral results from Experiment 1. This higher PLV over time for high-ASRS participants was observed at 12, 16, and 24Hz, and not observed below ~12Hz, even though the 8Hz band in particular was strongly affected by modulated music in the overall PLV (Figure 6A), and lower modulation frequencies were still acoustically strong in the AM+Music (Figure 1A). Instead, the selective increase in phase-locking at higher modulation frequencies may hint at the role of particular oscillatory regimes in the behavioral effects of this music. Increments of 4 Hz Cohen's d comparing AM+Music against Control-Music was 3.7391 for low-ASRS and 2.5161 for high-ASRS, but for increments of 1 Hz: Cohen's d was 0.4587 for low-ASRS and 0.0296 for high-ASRS individuals. Comparing late vs. early in the listening session, Cohen's d was -0.3960 for low-ASRS individuals, and 0.5343 for high-ASRS individuals when PLV was tested at increments of 4 Hz, but when PLV was tested at 1-Hz increments, Cohen's d was -0.1945 for low-ASRS and 0.0566 for high-ASRS individuals. The consistently higher effect size when PLV was tested at 4-Hz increments supports the finding that PLV was highest at integer multiples of 4 Hz. And while low-ASRS individuals were more effective at tracking the AM+Music

246   than were high-ASRS individuals, the high-ASRS individuals showed more of a change over time,
247   resulting in higher PLV late in the listening session.
248

249
250
Fig. 6 | Stimulus-brain coupling for high-ASRS and low-ASRS groups. Top row: PLV during
AM+Music (red) and Control-Music (black). **a,** low-ASRS participants; **b,** high-ASRS participants.
High-ASRS participants show higher PLV in higher frequencies, especially while listening to
AM+Music. The inset topos plots show the scalp distribution of PLV at 16 Hz during AM+Music.
Bottom row: PLV during AM+Music listening in the first half (black) and second half (red) of the
recording session. **c,** low-ASRS participants; **d,** high-ASRS participants. Black: first half of the
session; Red: second half of the session. Higher PLV for high-ASRS participants emerge over
time, and is higher late during the session. This is consistent with the behavioral improvement
seen in the second half of the AM+Music session for high-ASRS participants.

Experiment 4: Parametric manipulation of modulation rate and depth

The stimuli in Experiments 1-3 were taken from commercially available focus music that one might encounter if searching for music to work to. They were chosen to have a dramatic difference in modulation characteristics, as was apparent in acoustic analyses (Figure 1). However, they also differed in low-level acoustic properties (e.g., overall spectral balance) as well as musical features (e.g., tonality, instrumentation). To control for these differences and to isolate the effect of amplitude modulation, we developed new stimuli in which otherwise identical music was manipulated to impose modulation of varying rates and depths (Figure 7).

Parametrically testing the effects of amplitude modulations: Rate and Depth

Acoustic amplitude modulations are known to drive neural oscillations, i.e., to induce a selective amplification of neural activity at this frequency. This effect occurs along the auditory pathway but also in cortical networks such as the attentional network [33–36], and may thereby impact cognitive processes. We chose to test rates of 8, 16, and 32Hz for two reasons: First, these fall within ranges of distinct neural oscillatory regimes that are known to have different functions in the brain. Alpha (8-12Hz), beta (14-25Hz) and gamma (25-100Hz) rhythms are three such ranges, and our experimental conditions using 8, 16, and 32Hz modulation thus fall into each of these oscillatory regimes. Second, these rates were chosen to correspond to note values. As we used music that was composed at 120 beats per minute (2 Hz), amplitude modulation rates at 8, 16, and 32 Hz correspond to 16^{th}, 32^{nd}, and 64^{th} notes respectively.

Fig. 7 | Controlling modulation in music stimuli. Modulation in sound can be summarized by the modulation spectrum: The vertical axis on these 2D plots depicts audio frequency (0-8kHz). The horizontal axis shows the amplitude modulation spectrum (amplitude fluctuations from 0-100Hz). **a,** Applying modulation to music results in a modulation spectrum with a peak corresponding to the rate of the added modulation. Upper: The pressure wave is multiplied by a modulator (in this case 16Hz) to produce a modulated signal. Lower: This pressure wave is bandpass filtered, and fluctuation rates in each channel are shown in the modulation spectrum. **b,** Validation of stimulus manipulations used in Experiment 4. The stimulus space is illustrated by

these panels, each of which shows the difference in modulation spectrum (normalized power)
between an unmodulated track and the experimental conditions derived from this track. Rate of
added modulation increases moving rightward, while depth increases moving upward. The
absence of differences elsewhere on the modulation spectrum shows that the experimental
conditions were altered in a controlled way, with modulation properties different from the original
only as specified (stimulus validation).
299
300
Beta-band cortical activity is implicated in the maintenance of sensorimotor or cognitive
states[37] and top-down processing in general[38] including attentional control[39,40]. Moreover, in
spatial attention tasks, beta-band increase is observed in the hemisphere that represents the
attended stimulus, resulting in enhanced processing of the attended stimulus[41]. In contrast to the
other bands, stimulating neural activity in the beta band with the 16Hz modulation condition thus
seemed most likely to confer a performance benefit on sustained attention, and based on results
from Experiments 1-2 we would expect to see a greater effect in our high-ASRS participants. We
thus hypothesized that for higher-ASRS individuals, the 16Hz rate would produce effects on
performance superior to a no-modulation control condition, while the other rates would not. We
also hypothesized that greater modulation depth would have a greater effect on higher-ASRS
listeners.
Depth of modulation refers to how heavily the sound is modulated, rather than the rate of
modulation. A maximal depth of modulation would mean that sound energy is reduced to zero at
the troughs of the applied modulating waveform, while a very low depth would mean barely-
perceptible modulation. While a greater modulation depth is expected to impact neural oscillations
more strongly, beyond a point the underlying music suffers aesthetically as the sound becomes
distracting due to increased auditory salience from the sudden changes in loudness over time[42].
175 participants (81 high-ASRS and 94 low-ASRS) completed two SART experiments
online (4A and 4B) each with four conditions, testing modulation depth and modulation rate
separately. In both cases two groups of participants each heard four conditions, presented for five
minutes each in counterbalanced order. The first group of participants heard no-modulation, 8Hz
medium-depth, 16Hz medium-depth, and 32Hz medium-depth modulation. The second group
heard no-modulation, 16Hz low-depth, 16Hz medium-depth, and 16Hz high-depth modulation.
Thus, rate and depth (4 conditions each) were tested on separate groups of participants, but each
participant heard all four possible rates or all four possible depths for five minutes each. This
limited the duration of each condition, but maximized statistical power in comparing across all the
conditions in one dimension of the modulation spectrum, while controlling for intrinsic between-
subject differences in performance that are unrelated to our conditions of interest.

Figure 8. Stimulus valence and arousal ratings and performance over time in the Sustained
Attention to Response Task (SART) for stimuli parametrically manipulated in modulation
rate and depth. a, Valence and Arousal ratings for the music used in Experiment 4, with listeners
split by their level of self-reported attentional difficulty (ASRS score, median split). N=62 overall,
N=31 in each group (same participants as Experiment 1). **b,** Performance on the SART for varying
rates of added modulation. N=82 overall, N=46 and N=36 in the Low and High ASRS groups
respectively. **c,** Performance on the SART for varying depths of added modulation. N=93 overall,
N=48 and N=45 in the Low and High ASRS groups respectively. In Experiments 4A & 4B, each
participant completed 4 blocks (the music conditions) presented in random order; the blocks are
overlaid in the figures.

Valence and arousal ratings for all six sound stimuli were entered into a multivariate two-factor
mixed ANOVA (as in Experiment 1), with the within-subjects factor of Music (6 levels: No
Modulation, 16Hz Modulation at Low, Medium, and High Depth, 8Hz and 32Hz at Medium Depth)
and the between-subjects factor of ASRS (high-ASRS vs. low-ASRS groups). A main effect of
Music was observed on both valence and arousal (valence: $F_{(5,300)}=32.3$, $p<.001$; arousal:
$F_{(5,300)}=7.1$, $p<.001$). A main effect of ASRS was significant for arousal ($F_{(1,60)}=6.0, p=.017$)
but not for valence ($F_{(1,60)}=0.166$, n.s.).
Overall performance (d-prime) for the two groups in the rate experiment: Low-ASRS,
mean=3.68, stdv=0.72; High-ASRS, mean=3.36, stdv=0.69; and in the depth experiment: Low-
ASRS, mean=3.56, stdv=0.52; High-ASRS, mean=3.52, stdv=0.84.

The modulation depth experiment (Figure 8B) yielded overall d-prime values of Low-ASRS
mean=3.56 stdv=0.52; High-ASRS mean=3.52 stdv=0.84. The depth experiment revealed a trend
toward an interaction between ASRS group and SART performance over time (3-factor mixed
ANOVA; Between-Subjects factor of ASRS group: $F(1,270)=3.70$, $p=0.070$), but no effects of
music condition. The modulation rate experiment (Figure 8C) showed a significant main effect of
ASRS group (3-factor mixed ANOVA; between-subjects factor of ASRS group: $F(1,80) = 4.14$,
$p=0.045$), but also found interactions between ASRS and Time ($F(1,270) = 4.43$, $p=0.038$), as
well as ASRS and Music ($F(3,240) = 2.89$, $p=0.036$). This effect of Music by ASRS appears to be
driven by high-ASRS participants' better performance when hearing 16Hz-modulated music; this
finding is consistent with Experiment 1 and suggests rate-specific effects of modulation in music,
possibly subserved by oscillatory mechanisms in the brain.

365 Discussion

Performance on a variety of everyday tasks requires sustained attention. Cognitive failures,
specifically failures in sustained attention, are linked to mind wandering, which is associated with
decreased productivity and happiness[43]. Music is widely used to help with sustained attention,
with many use cases in everyday life ranging from café music to noise generators for work
environments. Here, we show that effects of music on sustained attention depend on one's level
of attentional difficulty. In high-ASRS participants, who experience more attentional difficulties, a
specific type of amplitude-modulated music (AM+Music) was more effective at engaging the
salience, executive control, sensorimotor, and visual networks, and at coupling with rhythmic brain
activity at multiple frequencies. While previous studies have suggested that differences in
perceived valence and/or arousal of music may explain differences in sustained attention, our
results show that valence and arousal do not explain effects of modulation across groups. Rather,
the effect of music on sustained attention appears to result from specific interactions between
functional differences in brain activity and amplitude modulation patterns in the musical sounds.
Individuals who self-report attentional difficulties are more sensitive to the effects of AM+Music in
behavioral and neural measures. Individual differences in sensitivity to background music may be
attributable to "optimal stimulation level"[44,45], i.e., the effect of external stimulation on cognitive
performance, which differs across individuals, e.g. between introverts and extroverts[12,46,47] and
between groups with and without attentional deficits[25,26,48]. These stimulus-brain interactions could
partly explain—in addition to preference and familiarity[15–17]—why people use such different types
of music to help them focus, and may suggest routes to more effective personalized focus music
in the future.
Behavioral experiments (Experiment 1) showed that high-ASRS participants performed
better on a sustained attention task when listening to music that contains rapid amplitude
modulations, compared to music with slower amplitude modulation or pink noise
(broadband/noisy modulation). While the AM+Music also elicited higher arousal ratings than the
Control-Music especially in the high-ASRS individuals, in a follow-up experiment (Experiment 4)
high-ASRS participants did best in a condition with relatively low arousal; together the
experiments suggest that the parameters of amplitude modulation causally affected behavior
independent of arousal ratings. This suggests that music can be tailor-made with added acoustic
modulations in order to aid performance in attentionally-demanding tasks, and that this can be
done on an individualized basis with ASRS as an important factor.
FMRI and EEG experiments (Experiments 2 & 3) identify some mechanisms behind these
effects. The fMRI experiment (Experiment 2) showed higher activity overall in response to
AM+Music. For the high-ASRS group in particular, AM+Music elicited higher activity in the anterior
cingulate cortex and the insula, core nodes of the salience network. Directly contrasting Hits
against False Alarms showed a greater extent of correct response-related activity during

AM+Music than other conditions, particularly in motor regions. This could indicate that behavioral advantage for AM+Music relates to activity in the motor system, i.e. that the AM+Music could be more effective at priming the motor system to respond correctly when receiving the task-relevant sensory signals from the sensory cortex.

Our EEG experiment (Experiment 3) showed that phase-locking activity was strongest during AM+Music and particularly stronger in the high-ASRS group, in whom the phase-locking activity also increased over time. These results suggested that amplitude modulation could underlie the difference in performance between ASRS groups observed in Experiment 1. To isolate the effects of amplitude modulation, a final behavioral experiment was conducted using music that differed only in the rate or depth of added modulation (Experiment 4). Here we found that 16Hz and high-depth amplitude modulation resulted in overall best performance in the high-ASRS group compared to the other conditions, resulting in performance for high-ASRS participants that was similar to the low-ASRS group.

Effects of music depend on one's level of attentional difficulty

A consistent result across our experiments was that effects of the music varied depending on an individual's level of attentional deficit, as measured by self-report on the ASRS. While previous studies have observed that individuals with attentional deficits are differently affected by music versus silence[5,14,49] or noise[12,20,21,50], here we extend the findings to show that differences between pieces of music are sufficient to affect sustained attention. In our EEG experiment (Experiment 3) we found that the difference between AM+Music and Control-Music was greater for the high-ASRS participants (Figure 6AB) especially in the beta range (12-30Hz); furthermore the change over time, i.e., the difference from the start of the block to the end of the block, was observable in the high-ASRS participants only (Figure 6CD). These results mirror our behavioral results from Experiment 1, where the low-ASRS group saw little difference in performance between music conditions and little change over time, whereas the high-ASRS group saw significant change over time across the music conditions (Figure 2B; significant interaction). Thus, in both brain and behavior, differences between Fast- and Control-Music were more apparent in high-ASRS individuals, and grew over the duration of the music; AM+Music both improved performance over time and stimulated neural activity in the beta range over time. This aligns with the idea that greater relative levels of beta-band activity are associated with improved performance on sustained attention tasks for individuals with ADHD[51–53].

Arousal does not explain effects of modulation across groups

While Experiment 1 suggested that the interaction between music for sustained attention and attentional deficit might depend on arousal, detailed examination of the effects of Fast- and Control-Music on phase-locking activity in the brain over time across ASRS groups (Figure 6CD) suggested that differences in acoustic modulation could underlie performance differences between high-arousal and low-arousal stimuli. In Experiment 4 the stimuli were made to vary only in modulation rate and depth, andhe condition with lowest arousal (16Hz medium depth condition) produced best performance in the high-ASRS group. Taken together the experiments suggest that acoustic modulation, rather than arousal per se, is the main driver behind the effects of music on sustained attention. Likewise, valence also cannot explain the differences in task performance. Although the increase in modulation depth from medium to high was sufficient to reverse the valence and arousal ratings of those stimuli, these ratings did not differ substantially between ASRS groups in the conditions that most affected sustained attention.

Background music has widespread effects on cognitive control networks implicated in ADHD brain activity

fMRI results showed higher activity during AM+Music than during other acoustic conditions, with significant effects in widely distributed regions encompassing the salience, executive function, and default mode networks, especially in the high-ASRS group. When contrasting activity between hits and false alarms to isolate successful task-related activity, AM+Music elicited highest activity in the sensorimotor network, the salience network, and the visual association network. The widespread increases in activity during AM+Music confirm that background music affects sustained attention by influencing multiple interconnected networks that are normally coupled to subserve performance on a variety of cognitive tasks. The default mode and executive function network are typically anticorrelated in activity[54], with the salience network being a consistent regulator of both other networks[29]. The interplay between salience, executive function, and default mode networks is critical in tasks requiring cognitive control, and this relationship is altered in ADHD by aberrant connectivity between these networks[55]. Here, when listening to AM+Music participants showed increased activity in the three networks. The interaction between group and music shows strong effects in the mid-cingulate cortex[56] and the right anterior insula, key components of the salience network, especially in high-ASRS individuals during AM+Music. The simultaneously observed effects in the salience network and the late visual areas / ventral attention network may suggest that the AM+Music affects performance by motivating attention to the visual task. It may also suggest increased functional connectivity among the networks as neural activity becomes coupled to the music. Increased coupling may also explain results from the EEG experiment, which found increased phase-locking to the stimulus at specific frequencies that were targeted by the note rate and the added amplitude modulation patterns of the AM+Music. Within the phase-locking patterns across different frequencies during AM+Music, scalp topography showed highest phase locking around frontocentral channels at low frequencies (8Hz) but more widespread activity across the scalp at higher frequencies (32Hz). These topographical differences in phase-locked activity across different frequencies may be explained by cross-frequency coupling mechanisms which are known to underlie communication across different regions in the brain[57].

Oscillatory mechanisms mediate effects of music on sustained attention

Our subsequent behavioral results also point to the possibility that the music's differential impact on oscillatory activity could underlie its effect on sustained attention. In Experiment 4 our stimuli were parameterized by modulation rate and showed a distinct 16Hz benefit for high-ASRS listeners, which could point to this effect being mediated by oscillatory processes in the brain. Individuals with ADHD have atypical oscillatory activity[36], including a higher theta/beta ratio[58]. The observed rate-specific effect confined to the high-ASRS group in Experiment 4 could indicate that the AM+Music impacts the atypical brain activity in ADHD. Taken together our results suggest that there could be a process by which music at high modulation rates drives brain activity to benefit ADHD-like individuals in particular. Importantly, such modulation rates are not normally found in music, but were added into the AM+Music.

Brain stimulation methods with controlled frequencies have been used to enhance cognitive performance in recent years[59–62]. Sound is another means by which to stimulate the brain[63–65]; however, the frequency of sound stimulation has often resulted in stimuli that are perceptually unpleasant (e.g., click trains, binaural beats). The observation that music strongly affects neural oscillations[32,65,66] motivates an approach whereby targeted acoustic modulation added to music might be used for neuromodulation, with particular goals outside the aesthetic

and/or social uses of music (i.e., 'functional' music rather than 'art' music). One barrier to this has
been individual variability, as different neurotypes and cognitive functions may require a range of
targets for their mechanisms. Here we targeted subpopulations (by ASRS/ADHD) with
hypotheses based on theoretical considerations from systems neuroscience, and found effects
on brain and behavior that have implications for the use of music to enhance cognition in everyday
life.

499
Fast and slow modulations in music can differently affect sustained attention. Results are not
explained by valence and arousal, and may instead be explained by differences in phase-locking
in specific frequencies as well as higher activity in widespread regions across the brain during
task performance. Individuals who self-report ADHD symptoms are most sensitive to these
modulations. This suggests that amplitude modulations in music could be used specifically to
mitigate the negative effects of cognitive failures in sustained attention that come from attentional
difficulties.
507

Methods
508

Experiment 1: Behavioral study
509

510   Experiment 1 presented three music background conditions to each participant (i.e., a within-
511   subjects design), in randomized order. Each of the three conditions lasted 6 minutes and 54
512   seconds, the time required to complete 360 trials of the SART; the experiment ran 1080 trials in
513   total (approximately 21 minutes).

514
Participants
We used Amazon's Mechanical Turk platform to recruit and enroll participants. 114 participants
were recruited for Experiment 1 (67 male, 40 female, 7 other/chose not to respond; mean age =
34). 102 participants were recruited to obtain valence and arousal ratings for the stimuli used in
Experiments 1 and 4 (obtained in the same participants). Participants were asked to wear
headphones, and not to change their volume or turn off audio during the experiment. They were
told that the background music was unrelated to the task, but that they should nonetheless ensure
they could hear the background music, because it was needed to control the acoustic environment
across participants. To ensure compliance we employed a headphone screening task [67] as well
as audio checks after each block, and a test of volume level at the end of the experiment.
Participants who failed any of these were removed from data analysis. The final dataset for
Experiment 1 comprised 87 of the initial 114 participants (76% passed screenings); the final
dataset for the valence and arousal ratings comprised 62 of the initial 102 participants (60.7%
passed screenings);

529
Stimuli
For Experiments 1 to 3, we chose as background auditory stimuli two commercially-available
tracks of music that were predicted to span different arousal levels while being similar in valence
levels, in addition to a Pink noise control stimulus. Background auditory stimuli were selected from
commercially available options used to help people focus while working. The tracks we used were:
(1) 'Techno March' by Brain.fm (high-arousal track) and (2) 'Tracking Aeroplanes' by The Echelon
Effect (low-arousal track). Our listener ratings (Figure 1A) confirmed that the tracks differed in
arousal but not in valence, suggesting that their effects on performance were attributable to

arousal rather than valence. (3) Pink noise was also chosen as a control stimulus; it is often used
for focused work while also being used in behavioral experiments because it has a spectrum that
falls off with increasing frequency, similar to many auditory environments (unlike white noise
which is spectrally flat). We limited our auditory stimuli to these three tracks in Experiments 1 in
order to obtain experimentally well-validated auditory stimuli for neuroimaging and
electrophysiology experiments in Experiments 2 and 3, both of which necessitated time-locked
analyses on a smaller number of participants.
The tracks were musically dissimilar (to drive differences in valence and arousal) but in
terms of low level acoustic features they differed most strikingly in the modulation domain--the
less arousing music was at a slow tempo and contained slow modulations, while the more
arousing music was at a fast tempo and contains fast modulations. The difference in the tracks'
modulation characteristics can be visualized with the modulation spectrum (Figure 1). The slow
music contained a broad region of modulation energy from 0-8Hz, while the fast music contained
a clear peak around 14-16Hz. This suggests tempo was not the sole cause of the difference in
modulation characteristics, as tempo would simply shift the modulation spectrum. The difference
in shape of the modulation spectrum instead reflects differences in the music, particularly note
values (event durations). That is, the fast music was at a faster tempo, but also had faster note
values. In particular, the modulation peak in the fast music was due to a consistent 32nd-note-
rate amplitude modulation that had been deliberately applied as a feature of this kind of focus
music. In contrast, the slow music contained long notes at several durations (e.g., whole, half)
resulting in the absence of a single modulation peak. To complement the music tracks, the Pink
Noise contained modulation energy broadly distributed in the higher modulation ranges
(roughness). The modulation-domain differences were large and may partly underlie the music's
effects, but the tracks varied in other ways (tonality, instrumentation, etc.) which likely contributed
to arousal. Due to the differences in frequency content of the three stimuli, rms-normalization was
not appropriate (resulting in a large difference in perceived loudness across the stimuli); instead
the stimuli were loudness-normalized by ear which produced rms values (average of left and right
channels) of 0.059, 0.061, and 0.014 for AM+, Control-Music, and Pink Noise respectively.
566

Procedure

Users enrolled via Amazon's Mechanical Turk platform, and provided informed consent as
approved by IRB # 120180271 of New England IRB. To enroll, users must be over 18 and have
normal hearing by self-report. If they chose to participate in our experiment, they were directed to
a cover page with consent documentation and a simple description of the task, followed by a page
with payment information. If they still chose to participate, they initiated a volume calibration task
in which they heard a train of 1kHz tones at alternating levels, 10dB apart. They were told to set
the volume on their computer so that only every other tone was audible, and told not to change
their volume after this calibration step. To ensure compliance a short task after the main
experiment required the participant to count the audible tones in a decrementing series of 1kHz
tones (-5dB per step); those who counted fewer tones than expected were excluded from
analysis. Following the initial volume calibration participants were directed to a headphone
screening task[67] composed of six 3AFC questions (<1 min). If they passed the headphone
screening they were directed to the main task instructions and could begin when ready.
In Experiment 1 our participants completed 1080 trials of a sustained attention to response
task (SART)[27,28,68]. In this task, a single digit ranging from 0 to 9 appeared on the screen for each
trial. Each digit was presented for 250 ms followed by a 900 ms mask, resulting in a 1150 ms
inter-trial interval. Participants' task was to respond to any digit except for 0. Instructions for the
task were as follows: "Numbers will appear on the screen. If you see 1-9, hit any key; if you see
0 do not hit any key." Participants were paid at a rate of $0.01 per correct response and -$0.10

per commission error (misses were given no pay; $0.00), resulting in an average of ~$12/hr for
the overall task. Participants are not penalized in any way for leaving, but any who did were not
granted the performance bonus (this was communicated at the outset), and were excluded from
analysis as incomplete data. Participants were told the experiment would run for about 20
minutes, and they should try to complete the entire experiment.
Valence and arousal ratings for these stimuli were obtained in a separate group of 62
participants under the same screening procedures. In this rating task, a webpage displayed
several audio player bars, each corresponding to a music track. Under each player were two
sliders (100-pixel resolution) with the ends of the sliders labeled 'Positive' - 'Negative' and
'Calming' - 'Stimulating'. Participants were given as much time as they liked to listen to the tracks
and decide on the placement of the sliders, and were permitted to move between tracks to better
judge them relative to one another. The stimuli for Experiment 1 and Experiment 4 were rated for
valence and arousal together (by the same participants).
600
Data Analysis
Raw data from Amazon's Mechanical Turk were exported to Matlab for analysis. The dependent
variable was accuracy (d-prime), and independent variables were music condition and time (first
half versus second half of each block). To test the effects of individual differences in attention
difficulties, as quantified by the ASRS[69], we did a median split on all subjects' ASRS scores,
yielding a high-ASRS group (i.e. those with more ADHD-like symptoms) and a low-ASRS group
(those with less ADHD-like symptoms).
608

Experiment 2: fMRI study

610 **Participants**
611 34 Wesleyan undergraduates (16 males, 18 females; mean age = 20.4, SD = 1.94) participated
612 for course credit.
613
Stimuli and Procedure
The same three background music conditions from Experiment 1 (AM+Music, Control-Music, and
Pink noise) were used in the fMRI study. During task fMRI, participants completed the SART while
listening to AM+Music, Control-Music, and Pink noise in counterbalanced order. All experiment
conditions were the same as Experiment 1 except inter-trial interval was 1425 ms, which was set
to be equivalent to 3 TRs (as TR = 475 ms). Before each block the volume of the auditory stimulus
was adjusted to a comfortable level (by communicating with the experimenter).
MRI Acquisition. High-resolution T1 and functional images were acquired in a 3T Siemens Skyra
MRI scanner at the Olin Neuropsychiatry Research Center at the Institute of Living.
The anatomical images were acquired using a T1-weighted, 3D, magnetization-prepared, rapid-
acquisition, gradient echo (MPRAGE) volume acquisition with a voxel resolution of 0.8 x 0.8 x 0.8
mm^3 (TR = 2.4 s, TE = 2.09 ms, flip angle = 8º, FOV = 256 mm). Task functional MRI was acquired
as 1268 contiguous fast-TR echo planar imaging (EPI) functional volumes (TR = 475 ms; TE =
30 ms; flip angle = 90, 48 slices; FOV = 240 mm; acquisition voxel size = 3 x 3 x 3 mm^3), resulting
in a sequence that lasted approximately 10 minutes. Each background music condition was
administered in a 10-minute sequence, resulting in approximately 30 minutes of scan time for
each participant.
631
Data analysis
MRI Preprocessing. Task and structural MRI preprocessing was carried out using the Statistical
Parametric Mapping 12 (SPM12) software (http://www.fil.ion.ucl.ac.uk/spm/) with the CONN

Toolbox (http://www.nitrc.org/projects/conn)[71]. In order, this consisted of functional realignment
and unwarp, functional centering, functional slice time correction, functional outlier detection using
the artifact detection tool (http://www.nitrc.org/projects/artifact_detect), functional direct
segmentation and normalization to MNI template, structural centering, structural segmentation
and normalization to MNI template, and functional smoothing to an 8mm gaussian kernel [72].
Denoising steps for functional connectivity analysis included correction for confounding effects of
white matter and cerebrospinal fluid[73], and bandpass filtering to 0.008-0.09 Hz.
Univariate Task-fMRI Analysis. Task fMRI analyses were done in SPM12 [31]. Task fMRI analyses
included: 1) Within-subject ANOVA comparing the three sessions (AM+, Control-Music, and Pink
noise). 2) Hits, misses, false alarms, and correct rejections were separately modeled. 3) Seed-
based functional connectivity was assessed using Conn and compared between the AM+,
Control-Music, and Pink noise conditions.

Seed-Based Connectivity Analyses. Since we were interested in whole-brain connectivity patterns
of known whole-brain networks, we used a functional network atlas that related 14 cortical
networks to known cognitive functions[74]. Of these 14 networks, 7 aligned with our a priori
hypotheses, including three DMN associated regions (vDMN, dDMN, Precuneus network), two
ECN associated networks (LECN, RECN), and two salience associated networks (posterior
salience network, anterior salience network). For the other 7 networks, hereafter referred to as
post-hoc networks, we performed correction for multiple comparisons by dividing all p-value
height thresholds by a factor of 7. Furthermore, noting a tendency towards false positives within
resting state fMRI analyses [75], we used a conservative p-value height threshold of 0.001 for group
comparisons and familywise error (p-FWE) correction for global effects of group and behavioral
measures, so as to minimize the possibility of Type I Errors.
Mean time-course of each network was used as a single region-of-interest (ROI), and
networks that resulted in clusters demonstrating a significant main effect of group were identified
at the height threshold p < 0.05 FWE-corrected level. For each cluster that showed significant
between-group differences, group-level beta values were extracted in order to determine the
driving factors of these group differences. These networks were then used to extract group
connectivity profiles at the height threshold p < 0.05, p-FWE corrected level.

665 Experiment 3: EEG study

666 Participants
667 40 Wesleyan undergraduates (10 males; 30 females; mean age = 19, SD = 0.75) participated for
668 course credit.

670 Stimuli and Procedure
Participants completed 680 trials of SART, a visual GO/NOGO task with inter-trial interval of 1150
ms, same as Experiment 1. This was done under four auditory conditions, presented in
counterbalanced order: AM+, Silence (within-subjects), Control-Music, and Pink noise (between-
subjects).
EEG was recorded with a 64-channel BrainVision actiCHamp system with PyCorder software in
a sound attenuated and electrically shielded chamber.

678 Data analysis
Behavioral data: RT coefficient of variation (SD/M) was calculated for every block of 10 trials.
EEG data were filtered using .5 Hz high pass filter and 60 Hz notch filter for electrical noise. Data
were re-referenced to channels TP9 and TP10 and corrected for ocular artifacts using ICA. Matlab
with EEGLAB toolbox [76] were used for analyses.

Preprocessing: First, bad channels were rejected and then interpolated using EEGLAB's rejchan
and eeg_interp functions. Stimulus-brain coupling was assessed by first applying Morlet wavelet
filtering at every single Hz from 1 to 50 Hz. Then, the Hilbert transform was done on each Hertz
to get phase angle of stimulus and of EEG data, and the coupling between stimulus and EEG was
assessed as the phase-locking value where plv(ch) = abs(sum(exp(1i*(phase_filtered_EEG(ch,:)
- phase_filtered_music)))). This was applied in parallel to EEG data and to sound stimuli. The
Pink Noise stimulus was generated in real time using Max/MSP (i.e., was not a stored file) and
so the stimulus-brain coupling analysis was not applied to the Pink Noise data.

691 Experiment 4: Behavioral study

692 Participants
693 Recruitment and experimental procedure (including screening) was identical to Experiment 1.
694 Experiment 4 involved 221 participants (120 males; 99 females; 2 other/chose not to respond;
695 mean age = 36, SD = 11.10). The final dataset for Experiment 4 comprised 175 of the initial 221
696 participants (79% passed screenings).
697

698 Stimuli
The stimuli (background music) were based on two different musical tracks; each had variants
created that added amplitude modulation at three rates (8, 16, 32 Hz) and depths (low, medium,
high). Modulation depth differences were quantified after processing to account for interactions
between the music and modulator. We used the difference between original and processed tracks'
modulation spectra (in each cochlear channel; Fig 7) as a metric of applied modulation depth, and
set the modulator such that our three depth conditions stepped up evenly in terms of this metric
going from low to high depth. The transitions between conditions were implemented with smooth
crossfades preserving track location (rather than a break and starting the music from the
beginning).
Modulation patterns were aligned to the metrical grid of the music. This scheme meant
that the relationship of the underlying music to the added modulation was as consistent as
possible over very different rates. This is desirable in controlling for differences between
conditions that arise from intrinsic properties of the underlying (pre-modulation) acoustic signal.
For example, a musical event (e.g. a drum hit) transiently amplified by a modulation peak in the
8Hz condition would also be so in the higher-rate conditions. This is only the case because the
different modulation rates are aligned to the music and are integer multiples of each other.
As can be seen in Figure 7, the applied modulation differences exist predominantly in the
low-mid range of the frequency spectrum, with little modulation difference at high frequencies.
This was by design, due to aesthetic considerations given the spectrotemporal density of the
underlying music: Modulation applied in a broadband manner tended to interact with sparse
events in the high frequency regions, which was occasionally annoying (salient). We confined
added modulation to lower frequencies by applying it to only the frequency range 200Hz-1kHz.
Acoustic analysis before and after processing showed that the manipulated stimuli differed
from the originals only in the modulation domain, and not in the audio frequency spectrum. Our
conditions were therefore identical in terms of musical content and spectral balance ('EQ'),
eliminating important confounding factors and ensuring any behavioral differences can be
attributed to applied modulation alone.   Due to the similarity across the stimuli they were
simply peak-normalized to 0.5, giving rms levels (average across left and right channels) in all
cases between 0.080 and 0.092, where the faster and higher-depth modulation conditions had
lower rms levels and the unmodulated tracks had the highest rms levels.
729

730 Procedure

Experimental procedure in Experiment 4 was similar to Experiment 1 with the addition of a fourth
block of trials. The total number of trials in Experiment 4 was the same as in Experiment 1 (1080
trials) and so each block in Experiment 4 was 270 trials (~5 minutes).

Data Analysis
For Experiment 4, a mixed-effects ANOVA was run on the dependent variable of accuracy, with
the within-subjects factors of rate (4 levels: no-modulation, 8 Hz, 16 Hz, 32 Hz), and the between-
subjects factors of ASRS (high vs. low) and extroversion (high vs. low).

Supplementary Materials

Listener volume settings. In Experiments 1 and 4, online participants were asked to first
calibrate their volume with an alternating sequence of low-level tones separated by 10dB; they
were told to set their computer volume so that only half the beeps could be heard. This
approximate calibration was intended to somewhat narrow the distribution of volume settings
participants used. In a post-test for volume, participants counted the audible tones in a series
decrementing by 5dB per tone.

748
749

750 **Fig. S1 | Online volume settings. a,** Relative volume distribution for participants in Experiment
751 1 (N=87); measured as the distribution of tones counted in a volume-setting post-test where 15
752 tones decremented in volume by 5db per tone. **b,** ASRS scores across participants in the high-
753 and low-volume groups.

754
755

756 Volume level might be expected to influence levels of arousal, or the effect of music in general [78].
757 To see if volume levels interact with the factors addressed in our main results, we split our
758 participants into two groups based on their volume settings (closest median split) and noted that
759 the ASRS scores did not differ between the high-volume and low-volume listeners (Fig S1B).

760
761
762

763
764 **Fig. S2 | Stimulus-brain coupling across EEG electrode channels**. For Experiment 3, PLV
765 from EEG data are separately plotted here for left (green), midline (blue), and right (yellow)
766 channels for frontal lateral sites (solid lines), frontal sites (dotted lines), and parietal sites (dashed
767 lines). Results show largely similar stimulus-brain coupling across multiple recording sites.
768

References

1. Sloboda, J. A., O'Neill, S. A. & Ivaldi, A. Functions of Music in Everyday Life: An Exploratory Study Using the Experience Sampling Method. *Music Sci.* **5**, 9–32 (2001).

2. North, A. C., Hargreaves, D. J. & Hargreaves, J. J. Uses of music in everyday life. *Music Percept.* **22**, 41–77 (2004).

3. Krueger, J. W. Doing things with music. *Phenomenol. Cognitive Sci.* **10**, 1–22 (2011).

4. Rickard, N. S., Toukhsati, S. R. & Field, S. E. The effect of music on cognitive performance: insight from neurobiological and animal studies. *Behav. Cogn. Neurosci. Rev.* **4**, 235–261 (2005).

5. Angel, L. A., Polzella, D. J. & Elvers, G. C. Background music and cognitive performance. *Percept. Mot. Skills* **110**, 1059–1064 (2010).

6. Schellenberg, E. G. & Weiss, M. W. Music and cognitive abilities. *The psychology of music., 3rd ed.* **3**, 499–550 (2013).

7. Bredemeier, H. C. & Drucker, P. F. Landmarks of Tomorrow. *American Sociological Review* vol. 24 917 (1959).

8. Davenport, T. H., Thomas, R. J. & Cantrell, S. The mysterious art and science of knowledge-worker performance. *MIT Sloan management review* **44**, 23 (2002).

9. Powell, W. W. & Snellman, K. The Knowledge Economy. *Annu. Rev. Sociol.* **30**, 199–220 (2004).

10. Drucker, P. F. Knowledge-worker productivity: the biggest challenge. *IEEE Eng. Manage. Rev.* **34**, 29–29 (2006).

11. Haake, A. B. Individual music listening in workplace settings: An exploratory survey of offices in the UK. *Music Sci.* **15**, 107–129 (2011).

12. Furnham, A. & Strbac, L. Music is as distracting as noise: the differential distraction of

background music and noise on the cognitive test performance of introverts and extraverts.

Ergonomics **45**, 203–217 (2002).

13. Furnham, A. & Bradley, A. Music while you work: The differential distraction of background

music on the cognitive test performance of introverts and extraverts. *Applied Cognitive*

Psychology: The Official Journal of the Society for Applied Research in Memory and

Cognition **11**, 445–455 (1997).

14. Reynolds, J., McClelland, A. & Furnham, A. An investigation of cognitive test performance

across conditions of silence, background noise and music as a function of neuroticism.

Anxiety Stress Coping **27**, 410–421 (2014).

15. Feng, S. & Bidelman, G. M. Music familiarity modulates mind wandering during lexical

processing. in *CogSci* (2015).

16. Perham, N. & Currie, H. Does listening to preferred music improve reading comprehension

performance? (2015).

17. Nantais, K. M. & Glenn Schellenberg, E. The Mozart Effect: An Artifact of Preference.

Psychological Science vol. 10 370–373 (1999).

18. Zentall, S. S. & Shaw, J. H. Effects of classroom noise on performance and activity of

second-grade hyperactive and control children. *J. Educ. Psychol.* **72**, 830–840 (1980).

19. Abikoff, H., Courtney, M. E., Szeibel, P. J. & Koplewicz, H. S. The Effects of Auditory

Stimulation on the Arithmetic Performance of Children with ADHD and Nondisabled

Children. *Journal of Learning Disabilities* vol. 29 238–246 (1996).

20. Söderlund, G., Sikström, S. & Smart, A. Listen to the noise: noise is beneficial for cognitive

performance in ADHD. *Journal of Child Psychology and Psychiatry* vol. 48 840–847 (2007).

21. Baijot, S. *et al.* Neuropsychological and neurophysiological benefits from white noise in

children with and without ADHD. *Behavioral and Brain Functions* vol. 12 (2016).

22. Zentall, S. Optimal stimulation as theoretical basis of hyperactivity. *American Journal of*

Orthopsychiatry vol. 45 549–563 (1975).

23. Zentall, S. S. & Meyer, M. J. Self-regulation of stimulation for ADD-H children during

reading and vigilance task performance. *J. Abnorm. Child Psychol.* **15**, 519–536 (1987).

24. Geissler, J., Romanos, M., Hegerl, U. & Hensch, T. Hyperactivity and sensation seeking as

autoregulatory attempts to stabilize brain arousal in ADHD and mania? *ADHD Attention*

Deficit and Hyperactivity Disorders vol. 6 159–173 (2014).

25. Strauß, M. *et al.* Brain arousal regulation in adults with attention-deficit/hyperactivity

disorder (ADHD). *Psychiatry Research* vol. 261 102–108 (2018).

26. Zentall, S. S. & Zentall, T. R. Optimal stimulation: A model of disordered activity and

performance in normal and deviant children. *Psychological Bulletin* vol. 94 446–471 (1983).

27. Robertson, I. H., Manly, T., Andrade, J., Baddeley, B. T. & Yiend, J. `Oops!': Performance

correlates of everyday attentional failures in traumatic brain injured and normal subjects.

Neuropsychologia **35**, 747–758 (1997).

28. Smilek, D., Carriere, J. S. A. & Allan Cheyne, J. Failures of sustained attention in life, lab,

and brain: Ecological validity of the SART. *Neuropsychologia* vol. 48 2564–2570 (2010).

29. Menon, V. & Uddin, L. Q. Saliency, switching, attention and control: a network model of

insula function. *Brain Struct. Funct.* **214**, 655–667 (2010).

30. Corbetta, M. & Shulman, G. L. Control of goal-directed and stimulus-driven attention in the

brain. *Nat. Rev. Neurosci.* **3**, 201–215 (2002).

31. Penny, W. D., Friston, K. J., Ashburner, J. T., Kiebel, S. J. & Nichols, T. E. *Statistical*

Parametric Mapping: The Analysis of Functional Brain Images. (Elsevier, 2011).

32. Doelling, K. B. & Poeppel, D. Cortical entrainment to music and its modulation by expertise.

Proc. Natl. Acad. Sci. U. S. A. **112**, E6233–42 (2015).

33. Lakatos, P., Karmos, G., Mehta, A. D., Ulbert, I. & Schroeder, C. E. Entrainment of

neuronal oscillations as a mechanism of attentional selection. *Science* **320**, 110–113

(2008).

34. Schroeder, C. E. & Lakatos, P. Low-frequency neuronal oscillations as instruments of

sensory selection. *Trends in Neurosciences* vol. 32 9–18 (2009).

35. Besle, J. *et al.* Tuning of the human neocortex to the temporal dynamics of attended

events. *J. Neurosci.* **31**, 3176–3185 (2011).

36. Calderone, D. J., Lakatos, P., Butler, P. D. & Xavier Castellanos, F. Entrainment of neural

oscillations as a modifiable substrate of attention. *Trends in Cognitive Sciences* vol. 18

300–309 (2014).

37. Engel, A. K. & Fries, P. Beta-band oscillations—signalling the status quo? *Current Opinion*

in Neurobiology vol. 20 156–165 (2010).

38. Bressler, S. L. & Richter, C. G. Interareal oscillatory synchronization in top-down

neocortical processing. *Current Opinion in Neurobiology* vol. 31 62–66 (2015).

39. Gross, J. *et al.* Modulation of long-range neural synchrony reflects temporal limitations of

visual attention in humans. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 13050–13055 (2004).

40. Lee, J. H., Whittington, M. A. & Kopell, N. J. Top-down beta rhythms support selective

attention via interlaminar interaction: a model. *PLoS Comput. Biol.* **9**, e1003164 (2013).

41. Richter, C. G., Coppola, R. & Bressler, S. L. Top-down beta oscillatory signaling conveys

behavioral context in early visual cortex. *Scientific Reports* vol. 8 (2018).

42. Kayser, C., Petkov, C. I., Lippert, M. & Logothetis, N. K. Mechanisms for Allocating Auditory

Attention: An Auditory Saliency Map. *Current Biology* vol. 15 1943–1947 (2005).

43. Killingsworth, M. A. & Gilbert, D. T. A wandering mind is an unhappy mind. *Science* **330**,

932 (2010).

44. Raju, P. S. Optimum Stimulation Level: Its Relationship to Personality, Demographics, and

Exploratory Behavior. *Journal of Consumer Research* vol. 7 272 (1980).

45. Fischer, T., Langner, R., Birbaumer, N. & Brocke, B. Arousal and Attention: Self-chosen

Stimulation Optimizes Cortical Excitability and Minimizes Compensatory Effort. *Journal of*

Cognitive Neuroscience vol. 20 1443–1453 (2008).

46. Geen, R. G. Preferred stimulation levels in introverts and extroverts: Effects on arousal and

performance. *Journal of Personality and Social Psychology* vol. 46 1303–1312 (1984).

47. Standing, L., Lynn, D. & Moxness, K. Effects of noise upon introverts and extroverts.

Bulletin of the Psychonomic Society vol. 28 138–140 (1990).

48. Antrop, I., Roeyers, H., Van Oost, P. & Buysse, A. Stimulation Seeking and Hyperactivity in

Children with ADHD. *Journal of Child Psychology and Psychiatry* vol. 41 225–231 (2000).

49. Kämpfe, J., Sedlmeier, P. & Renkewitz, F. The impact of background music on adult

listeners: A meta-analysis. *Psychology of Music* **39**, 424–448 (2011).

50. Helps, S. K., Bamford, S., Sonuga-Barke, E. J. S. & Söderlund, G. B. W. Different Effects of

Adding White Noise on Cognitive Performance of Sub-, Normal and Super-Attentive School

Children. *PLoS ONE* vol. 9 e112768 (2014).

51. Bresnahan, S. M. & Barry, R. J. Specificity of quantitative EEG analysis in adults with

attention deficit hyperactivity disorder. *Psychiatry Res.* **112**, 133–144 (2002).

52. Loo, S. K. *et al.* Cortical activity patterns in ADHD during arousal, activation and sustained

attention. *Neuropsychologia* **47**, 2114–2119 (2009).

53. Ogrim, G., Kropotov, J. & Hestad, K. The quantitative EEG theta/beta ratio in attention

deficit/hyperactivity disorder and normal controls: Sensitivity, specificity, and behavioral

correlates. *Psychiatry Research* vol. 198 482–488 (2012).

54. Fox, M. D. *et al.* The human brain is intrinsically organized into dynamic, anticorrelated

functional networks. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 9673–9678 (2005).

55. Cai, W., Griffiths, K., Korgaonkar, M. S., Williams, L. M. & Menon, V. Inhibition-related

modulation of salience and frontoparietal networks predicts cognitive control ability and

inattention symptoms in children with ADHD. *Mol. Psychiatry* (2019) doi:10.1038/s41380-

019-0564-4.

56. van Heukelum, S. *et al.* Where is Cingulate Cortex? A Cross-Species View. *Trends*

Neurosci. **43**, 285–299 (2020).

57. Fiveash, A. *et al.* A stimulus-brain coupling analysis of regular and irregular rhythms in

adults with dyslexia and controls. *Brain Cogn.* **140**, 105531 (2020).

58. Woltering, S., Jung, J., Liu, Z. & Tannock, R. Resting state EEG oscillatory power

differences in ADHD college students and their peers. *Behav. Brain Funct.* **8**, 60 (2012).

59. Marshall, L., Helgadóttir, H., Mölle, M. & Born, J. Boosting slow oscillations during sleep

potentiates memory. *Nature* vol. 444 610–613 (2006).

60. Nelson, J. T., Andy McKinley, R., Golob, E. J., Warm, J. S. & Parasuraman, R. Enhancing

vigilance in operators with prefrontal cortex transcranial direct current stimulation (tDCS).

NeuroImage vol. 85 909–917 (2014).

61. Albouy, P., Weiss, A., Baillet, S. & Zatorre, R. J. Selective Entrainment of Theta

Oscillations in the Dorsal Stream Causally Enhances Auditory Working Memory

Performance. *Neuron* vol. 94 193–206.e5 (2017).

62. Reinhart, R. M. G. & Nguyen, J. A. Working memory revived in older adults by

synchronizing rhythmic brain circuits. *Nature Neuroscience* vol. 22 820–827 (2019).

63. Ngo, H.-V. V., Martinetz, T., Born, J. & Mölle, M. Auditory closed-loop stimulation of the

sleep slow oscillation enhances memory. *Neuron* **78**, 545–553 (2013).

64. Lerousseau, J. P., Trébuchon, A., Morillon, B. & Schön, D. Frequency selectivity of

persistent cortical oscillatory responses to auditory rhythmic stimulation. *bioRxiv* 834226

(2021) doi:10.1101/834226.

65. Wollman, I., Arias, P., Aucouturier, J.-J. & Morillon, B. Neural entrainment to music is

sensitive to melodic spectral complexity. *J. Neurophysiol.* **123**, 1063–1071 (2020).

66. Trost, W. *et al.* Getting the beat: entrainment of brain activity by musical rhythm and

pleasantness. *Neuroimage* **103**, 55–64 (2014).

67. Woods, K. J. P., Siegel, M. H., Traer, J. & Mcdermott, J. Headphone screening to facilitate

web-based auditory experiments. doi:10.31234/osf.io/aez6f.

68. Helton, W. S. Impulsive responding and the sustained attention to response task. *Journal of*

Clinical and Experimental Neuropsychology vol. 31 39–47 (2009).

69. Kessler, R. C. *et al.* The World Health Organization adult ADHD self-report scale (ASRS): a

short screening scale for use in the general population. *Psychological Medicine* vol. 35

245–256 (2005).

70. Gosling, S. D., Rentfrow, P. J. & Swann, W. B. A very brief measure of the Big-Five

personality domains. *J. Res. Pers.* **37**, 504–528 (2003).

71. Whitfield-Gabrieli, S. & Nieto-Castanon, A. Conn: A Functional Connectivity Toolbox for

Correlated and Anticorrelated Brain Networks. *Brain Connectivity* vol. 2 125–141 (2012).

72. Friston, K. J. *et al.* Spatial registration and normalization of images. *Hum. Brain Mapp.* **3**,

165–189 (1995).

73. Behzadi, Y., Restom, K., Liau, J. & Liu, T. T. A component based noise correction method

(CompCor) for BOLD and perfusion based fMRI. *Neuroimage* **37**, 90–101 (2007).

74. Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V. & Greicius, M. D. Decoding subject-

driven cognitive states with whole-brain connectivity patterns. *Cereb. Cortex* **22**, 158–165

(2012).

75. Eklund, A., Nichols, T. E. & Knutsson, H. Cluster failure: Why fMRI inferences for spatial

extent have inflated false-positive rates. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 7900–7905

(2016).

76. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-trial EEG

dynamics including independent component analysis. *J. Neurosci. Methods* **134**, 9–21

(2004).

77. Nozaradan, S., Peretz, I. & Mouraux, A. Selective neuronal entrainment to the beat and

meter embedded in a musical rhythm. *J. Neurosci.* **32**, 17572–17581 (2012).

78. Wolfe, D. E. Effects of Music Loudness on Task Performance and Self-Report of College-

Aged Students. *Journal of Research in Music Education* **31**, 191–201 (1983).