Title: Maintenance of community function through compensation breaks down over time in a

desert rodent community

Authors: Renata M. Diaz[1] and S. K. Morgan Ernest[2]

1. School of Natural Resources and Environment, University of Florida, Gainesville, FL.

Corresponding author. renata.diaz@weecology.org

2. Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL.

Original submission: This submission analyzes long-term data on rodent community abundance

and energy use from the Portal Project. Sections of this timeseries have been analyzed in

numerous other publications, but this is the first to analyze data from 2007-2020 on

compensation on experimental and control plots.

No prior publication: This submission is posted as a preprint on bioRxiv at

https://www.biorxiv.org/content/10.1101/2021.10.01.462799v1.

Animal welfare: Rodent censuses were conducted with IACUC approval, most recently under

protocol 201808839_01 at the University of Florida.

Open research: All data and code to reproduce these analyses are archived on Zenodo at

https://doi.org/10.5281/zenodo.5544362 and https://doi.org/10.5281/zenodo.5539881.

Analytic methods: All analyses were conducted in R version 4.0.3.

Abstract

Understanding the ecological processes that maintain community function in systems

experiencing species loss, and how these processes change over time, is key to understanding the

relationship between community structure and function and predicting how communities may

respond to perturbations in the Anthropocene. Using a 30-year experiment on desert rodents, we

show that the impact of species loss on community-level energy use has changed repeatedly and

dramatically over time, due to 1) the addition of new species to the community, and 2) a

reduction in functional redundancy among the same set of species. Although strong

compensation, initially driven by the dispersal of functionally redundant species to the local

community, occurred in this system from 1997-2010, since 2010, compensation has broken

down due to decreasing functional overlap within the same set of species. Simultaneously, long-

term changes in sitewide community composition due to niche complementarity have decoupled

the dynamics of compensation from the overall impact of species loss on community-level

energy use. Shifting, context-dependent compensatory dynamics, such as those demonstrated

here, highlight the importance of explicitly long-term, metacommunity, and eco-evolutionary

perspectives on the link between species-level fluctuations and community function in a

changing world.

Key words: community function, compensation, zero-sum dynamic, environmental fluctuations,

functional redundancy

Introduction

Determining the extent to which community-level properties are affected by species loss,

and how and why this changes over time, is key for understanding how communities are

structured and how community function may respond to future perturbations (Gonzalez and

Loreau 2009). When species are lost from a community, their contributions to community

function (e.g. total productivity or resource use) are also directly lost. Community function may

be maintained, however, if in the new community context, species that remain perform similar

functions to the species that were lost, and compensate for the decline in function directly caused

by species loss - i.e., functional redundancy (Walker 1992, 1995; Ernest and Brown 2001;

Rosenfeld 2002; Gonzalez and Loreau 2009). When compensation via functional redundancy

occurs among consumers with a common resource base, it is consistent with a zero-sum

competitive dynamic, in which resources not used by one species are readily absorbed by

competitors, and any increases in the abundance of one species must come at the direct expense

of others (Van Valen 1973; Ernest et al. 2008).

Because the response of system-level function to species loss is partially determined by

degree of functional redundancy in a community, processes that cause functional redundancy to

change over time can have important consequences for the long-term maintenance of ecosystem

function. Colonization events may buffer community function against species loss, if a

community gains species that perform similar functions to the species that were lost (Ernest and

Brown 2001; Leibold et al. 2017). The ability of colonization to supply functionally redundant

species depends on the species (and traits) present in the broader metacommunity, and on the rate

of dispersal supplying appropriate species to local communities (Leibold et al. 2017).

Even without the addition of new species and traits, however, functional redundancy

within a consistent set of coexisting species may fluctuate over time. While, in theory, functional

redundancy may occur via the special case of complete niche neutrality (e.g. Hubbell 2001), it

may also occur in niche-structured systems that contain species that share some traits but differ

along other niche axes (Thibault et al. 2010). In these systems, if similar, but non-identical,

species respond to environmental change in similar ways, functional overlap can be maintained

or even strengthened. However, if niche differences cause species to respond differently to

changing conditions, the degree of functional overlap between those species may decline,

resulting in a breakdown in compensation (Loreau 2004; Fetzer et al. 2015). Over time, as

metacommunity dynamics and changing environmental conditions modulate functional

redundancy within a community, the extent to which community function is robust to species

loss - and the strength of zero-sum competition - may also be dynamic and context-dependent.

Despite logical conceptual support, and evidence from experimental microcosms (Fetzer

et al. 2015), there is little empirical documentation of how, and through which mechanisms,

temporal changes in functional redundancy modulate the effect of species loss on ecosystem

function in natural assemblages. Although relatively plentiful, observational data cannot

unambiguously detect compensation through functional redundancy, and even short-term

experiments may not be sufficient to capture temporal variation in compensation (Ernest and

Brown 2001; Houlahan et al. 2007). In contrast, long-term manipulative experiments are

uniquely suited to address this question. In long-term experiments in which key species are

removed from a community over an extended period of time, the impact of species loss on

community function can be directly quantified by comparing community function between

complete and manipulated assemblages. As metacommunity dynamics and environmental

conditions shift over time, long-term monitoring can reveal how these processes contribute to

changes in functional redundancy and ecosystem function across different time periods. Due to

the financial and logistical resources required to maintain and monitor whole-community

manipulations over long timescales, these experiments are rare in natural systems representative

of realistic evolutionary, geographic, and environmental constraints (Hughes et al. 2017).

Here, we use a 30-year experiment on desert rodents to investigate how shifts in

functional redundancy alter the effect of species loss on community function over time. In this

study, kangaroo rats (*Dipodomys* spp.), the largest and competitively dominant species in the

rodent community, have been removed from a subset of experimental plots to explore how the

loss of key species affects community function, measured as community-level metabolic flux

("total energy use", or *Etot*) or total biomass (Ernest et al. 2019). For systems of consumers with

a shared resource base, such as this community of granivorous rodents, *Etot* reflects the total

amount of resources being processed by an assemblage, and total biomass directly reflects

standing biomass. Both are important metrics of community function (Lawton 1994; Ernest and

Brown 2001). Long-term monitoring of this experiment has documented repeated shifts in the

habitat and species composition of this system, resulting in distinct time periods characterized by

different habitat conditions and configurations of the rodent community (Christensen et al.

2018). Abrupt reorganization events in community composition occurred in 1997 and in 2010,

associated with the establishment and subsequent decline of the pocket mouse *Chaetodipus*

baileyi. *C. baileyi* is similar in size, and presumably other traits, to kangaroo rats, and its

establishment in 1996-97 drove a pronounced increase in compensation due to functional

redundancy between *C. baileyi* and kangaroo rats (Ernest and Brown 2001; Thibault et al. 2010).

Over the course of this experiment, shifting environmental conditions have caused the habitat at

the study site to transition from desert grassland to scrub, driving a shift in baseline rodent

community composition away from kangaroo rats and favoring other, smaller, granivores

(Brown et al. 1997; Ernest et al. 2008). By making comparisons across these time periods, we

explored how shifts in community composition and functional overlap among the same species

have contributed to long-term changes in the effect of species loss on community function.

Methods

The Portal Project

The Portal Project consists of a set of 24 fenced experimental plots located approximately

7 miles east of Portal, AZ, USA, on unceded land of the Chiricahua Apache. Beginning in 1977,

kangaroo rats (*Dipodomys spectabilis, D. merriami,* and *D. ordii*) have been experimentally

excluded from a subset of these plots (exclosures), while all other rodents are allowed access

through small holes cut in the plot fencing. Control plots, with larger holes, are accessible to all

rodents, including kangaroo rats. Rodents on all plots are censused via monthly bouts of live-

trapping.Each individual captured is identified to species and weighed. For additional details on

the site and methodology of the Portal Project, see Ernest et al. (2019).

Data

We used data for control and exclosure plots from February 1988 until January 2020. The

experimental treatments for some plots have changed over time, and we used the subset of plots

that have had the same treatments for the longest period of time (Ernest et al. 2019). Four control

plots, and five exclosure plots, met these criteria. In order to achieve a balanced sample, we

randomly selected four exclosure plots for analysis. We divided the timeseries into three time

periods defined by major transitions in the rodent community surrounding the establishment and

decline of *C. baileyi* (Ernest and Brown 2001; Christensen et al. 2018). The first time period

(February 1988-June 1997) precedes *C. baileyi*'s establishment at site. We defined *C. baileyi*'s

establishment date as the first census period in which *C. baileyi* was captured on all exclosure

plots (following Bledsoe and Ernest, 2019). During the second time period (July 1997-January

2010), *C. baileyi* was abundant on both exclosure and control plots. This time period ended with

a reorganization event in which *C. balieyi* became scarce sitewide. We used January 2010, the

midpoint of the 95% credible interval for the date of this reorganization event as estimated in

Christensen et al. (2018), as the end date for this time period. The last time period spans from

Feburary 2010-January 2020. For each individual rodent captured, we estimated the individual-

level metabolic rate using the scaling relationship between individual body mass and metabolic

rate $b = 5.69 * (m^{0.75})$, where m is body mass in grams and b is metabolic rate (for details, see

White et al. 2004). We calculated treatment and species-level energy use as the sum of the

appropriate individuals' metabolic rates, and total biomass as the sum of individuals' body mass

measurements.

Statistical analysis of rodent community energy use and biomass

Here, we describe analyses for energy use. For biomass, we repeated these analyses

substituting biomass values for energy use throughout. For all variables, we combined data for

all plots within a treatment in each monthly census period and calculated treatment-level means.

This is necessary to calculate compensation, and we treated other variables in the same way to

maintain consistency. A provisional plot-level analysis yielded qualitatively equivalent results

(Appendix S1). To measure the overall impact of kangaroo rat removal on $Etot$, we calculated a

"total energy ratio" as the ratio of treatment-level $Etot$ for kangaroo-rat exclosure plots relative to

unmanipulated control plots, i.e. $Etot_E/Etot_C$ where $Etot_E$ and $Etot_C$ are total energy use on

exclosures and controls, respectively (Thibault et al 2010; Bledsoe and Ernest 2019). This ratio is

distinct from compensation, which we defined as the proportion of the energy made available by

kangaroo rat removal taken up via compensatory increases in energy use by small granivores (all

granivores other than kangaroo rats; *Baiomys taylori, C. baileyi, Chaetodipus hispidus,*

Chaetodipus intermedius, Chaetodipus penicillatus, Perognathus flavus, Peromyscus eremicus,

Peromyscus leucopus, Peromyscus maniculatus, Reithrodontomys fulvescens, Reithrodontomys

megalotis, and *Reithrodontomys montanus*). We calculated this as $(SG_E - SG_C)/KR_C$, where SG_E

and SG_C are the amount of energy used by small granivores (SG) on exclosure and control plots,

respectively, and KR_C is the amount of energy used by kangaroo rats (KR) on control plots

(Ernest and Brown 2001). To compare these variables across time periods, we used generalized

least squares models (GLS; the R package *nlme*; Pinheiro et al. 2020) of the form (SG_E –

$SG_C)/KR_C \sim$ *time period*, for compensation, and $Etot_E/Etot_C \sim$ *time period*, for the total energy

ratio. We included a continuous-time autoregressive temporal autocorrelation term to account for

temporal autocorrelation between values from monthly census periods within each multi-year

time period (for details of model selection, see Appendix S2). To evaluate change in baseline

community composition over time, we calculated the proportion of treatment-level energy use

accounted for by kangaroo rats on control plots in each census period ($KR_C/Etot_C$). Proportional

energy use is bounded 0-1 and is therefore not appropriate for GLS, so we compared values

across time periods using a binomial generalized linear model (GLM) of the form $KR_C/Etot_C \sim$

time period. Finally, we calculated the proportional energy use accounted for by *C. baileyi* (CB)

on exclosure and control plots in each census period ($CB_E/Etot_E$ and $CB_C/Etot_C$, respectively). *C.*

baileyi was not present at the site prior to 1996, and we restricted the analysis of *C. baileyi*

proportional energy use to the second two time periods. We compared *C. baileyi* proportional

energy use over time and across treatments using a binomial GLM of the form $CB_E/Etot_E \sim$ *time*

period + treatment. For all models, we calculated estimated means and 95% confidence or

credible intervals for time-period (and, for *C. baileyi*, treatment) level values, and contrasts

between time periods (and, for *C. baileyi*, treatments), using the R package *emmeans* (Lenth

2021). Analyses were conducted in R 4.0.3 (R Core Team 2020). Data and code are archived at

https://doi.org/10.5281/zenodo.5544362 and https://doi.org/10.5281/zenodo.5539881.

Results

180   The impact of kangaroo rat removal on community function has changed repeatedly

over time, through a combination of abrupt shifts in compensation associated with *C. baileyi*, and

long-term changes in baseline community composition sitewide (Figure 1). These dynamics are

qualitatively identical whether function is measured as total energy use (Figure 1; Appendix S2)

or total biomass (Appendix S3). The first shift coincided with *C. baileyi*'s establishment in the

community beginning in 1996-97 (Figure 1D). *C. baileyi* rapidly became dominant on exclosure

plots and dramatically increased compensation (Figure 1B). From 1997-2010, small granivores

compensated for an average of 58% of kangaroo rat energy use on control plots (95% interval

48-67%), an increase from an average of 18% from 1988-1997 (95% interval 8-29%; contrast p

< 0.001; for complete results of all models, see Appendix S2) from 1997-2010. With *C. baileyi*'s

addition to the community, the total energy ratio (on exclosures relative to controls; Figure 1A)

increased from 30% (20-40%) to 71% (62-79%, contrast $p < 0.014$). In the second shift,

beginning around 2010, *C. baileyi*'s abundance sitewide dropped precipitously (Figure 1D). *C.*

baileyi's proportional energy use dropped from an average of 72% (65-80%) to 26% (18-35%,

contrast $p < 0.001$) on exclosure plots, and from 11% (6-16%) to essentially 0 on control plots

(contrast $p < 0.001$). Other species of small granivore did not make compensatory gains to offset

the decline in *C. baileyi* (Figure 1B). As a result, compensation declined from an average of 58%

(48-67%) to 28% (17-38%, contrast $p = 0.002$), a level not significantly different from the 18%

(8-29%, contrast $p = .44$) observed prior to *C. baileyi*'s establishment at the site. Somewhat

paradoxically, while the total energy ratio also dropped following *C. baileyi*'s decline, from an

average of 71% (62-79%) from 1997-2010 to 50% (40-60%, contrast $p = 0.0056$) from 2010-

2020, it remained higher than its average of 30% (20-40%, contrast $p = 0.0144$) from 1988-1997

(Figure 1A). Over the course of the experiment, community composition shifted sitewide. In

later years, kangaroo rats accounted for a lower proportion of baseline *Etot* than they did at the

beginning of the study (Figure 1C). From 1988-1997, kangaroo rats accounted for 92% (87-97%)

of *Etot* on controls; after 1997, this dropped to an average of approximately 70% (1988-1997

compared to later time periods, both $p = .0004$; 1997-2010 and 2020-2020 not significantly

different, $p = .976$). Because the proportion of *Etot* directly lost to kangaroo rat removal was

smaller from 2010-2020 than from 1988-1997, the total energy ratio was higher from 2010-2020

than it was from 1988-1997 - even though there was not a detectable difference between the two

time periods in the proportion of lost energy being offset through compensation.

Discussion

The dynamics of rodent community energy use at Portal illustrate that the role of

functional redundancy in buffering community function against species loss fluctuates over time,

due to changes in both species composition and in the degree of functional overlap among the

same species. The 1997 increase in compensation, driven by *C. baileyi*'s establishment at the

site, was a clear and compelling instance of colonization from the regional species pool

overcoming limitations on functional redundancy (Ernest and Brown 2001; Leibold et al 2017).

Although the small granivore species originally present in the community did not possess the

traits necessary to compensate for kangaroo rats, *C. baileyi* supplied those traits and

substantially, but incompletely, restored community function. In contrast, following the

community reorganization event in 2010, *C. baileyi* remained present in the community, but

ceased to operate as a partial functional replacement for kangaroo rats. This is consistent with

fluctuating conditions modulating functional redundancy between similar, but non-identical,

competitors. Kangaroo rats and *C. baileyi* are relatively similar in size and are demonstrably

capable of using similar resources. However, *C. baileyi* prefers different, shrubbier microhabitats

than kangaroo rats, and the two groups have been observed to replace each other in adjacent

habitats (Ernest and Brown 2001). We suggest that this study site, which has historically been

dominated by kangaroo rats, constitutes marginal habitat for *C. baileyi*, and that, while

conditions from 1997-2010 aligned sufficiently with *C. baileyi*'s requirements to create

appreciable functional redundancy between kangaroo rats and *C. baileyi*, conditions since have

caused this redundancy to break down. *C. baileyi*'s decline occurred immediately following a

period of low plant productivity and low rodent abundance community-wide, and in the decade

following, the site experienced two long and severe droughts (Appendix S4; Christensen et al.

2018). These extreme conditions may themselves have limited *C. baileyi's* fitness at the site, or

the community-wide low abundance event may have temporarily overcome incumbency effects

and triggered a community shift tracking longer-term habitat trends (Thibault and Brown 2008;

Christensen et al. 2018). Regardless of the proximate cause of *C. baileyi*'s decline, the fact that

C. baileyi remains in the community, but no longer compensates for kangaroo rats, illustrates

that changing conditions can have profound effects on community function by modulating the

degree of functional redundancy within a consistent set of species.

While changes in compensation have contributed to changes in community function in

this system, changes in compensation alone do not fully account for the long-term changes in the

overall impact of kangaroo rat removal on *Etot*. Since 2010, although the ratio of *Etot* on

exclosure plots relative to control plots declined coinciding with the breakdown in compensation

associated with *C. baileyi*, it remained higher than the levels observed prior to 1997 (Figure 1A).

This difference between the first and last time periods cannot be explained by an increase in

compensation, as compensation from 2010-2020 was not greater than pre-1997 levels (Figure

1B). Rather, the increase in *Etot* on exclosure plots relative to control plots was the result of a

long-term decrease in the contribution of kangaroo rats to *Etot* sitewide. Because kangaroo rats

accounted for a smaller proportion of *Etot* on control plots from 2010-2020 than they did prior to

1997, their removal had a smaller impact on community function – even though there was not an

increase in the degree to which small granivores compensated for their absence. In fact, the

comparable levels of compensation achieved in the decades preceding and following *C. baileyi*'s

dominance at the site suggest a relatively stable, and limited, degree of functional overlap

between kangaroo rats and the original small granivores (i.e., excluding *C. bailyei*). Niche

complementarity, combined with changing habitat conditions, may partially explain this

phenomenon. It is well-documented that, while kangaroo rats readily forage in open

microhabitats where predation risk can be relatively high, smaller granivores preferentially

forage in sheltered microhabitats as an antipredator tactic (Kelt 2011). Over the course of this

experiment, the habitat at this study site has transitioned from an arid grassland to a shrubland

(Brown et al. 1997). As sheltered microhabitats became more widespread, small granivores may

have gained access to a larger proportion of resources and increased their share of *Etot* sitewide.

However, kangaroo rats may have continued to use resources in open areas, which would have

remained inaccessible to smaller granivores even on exclosure plots. The long-term reduction in

the impact of kangaroo rat removal on community function, driven by niche complementarity

and consistent niche partitioning, contrasts with the temporary compensatory dynamic driven by

functional redundancy with *C. baileyi*. Although changes in the overall effect of species loss are

sometimes treated interchangeably with compensation (e.g. Ernest and Brown 2001 compared to

Thibault et al. 2010), it is important to recognize that multiple distinct pathways modulate the

long-term impacts of species loss on community function. Particularly in strongly niche-

structured systems, complementarity effects and fluctuations in functional redundancy may occur

simultaneously, with complex and counterintuitive impacts on community function.

Overall, the decadal-scale changes in energy use among the Portal rodents underscore the

importance of long-term metacommunity dynamics to the maintenance of community function

following species loss (see Leibold et al. 2017). Although a single colonization event may allow

for temporary compensation via functional redundancy, as conditions shift, species that once

compensated may no longer perform that function (see also Isbell et al. 2011). Particularly if

limiting similarity prevents similar competitors from specializing on precisely the same habitats

(Rosenfeld 2002), temporary, context-dependent compensation may be common. To maintain

compensation over time, multiple colonization events, supplying species that are functionally

redundant under different conditions, may be required. Depending on dispersal rates, and the

diversity and composition of regional species pools, this may be unlikely or even impossible. At

Portal, dispersal limitation introduced a 20-year delay in the compensatory response driven by *C.*

baileyi. Theoretically, a new species capable of compensating for kangaroo rats, and better-

suited to conditions at the site since 2010, could restore compensation under present conditions –

but it is unclear whether this species exists or if it can disperse to this site. As ecosystems

globally undergo reductions in habitat connectivity and regional beta diversity, and enter novel

climatic spaces, maintenance of community function via functional redundancy may grow

increasingly rare and fragile (Dornelas et al. 2014; Williams and Jackson 2007).

Finally, the long-term variability in functional redundancy documented here adds

important nuance to our understanding of how zero-sum dynamics operate in natural

assemblages. Theories invoking zero-sum dynamics, and tests for compensatory dynamics in

empirical data, often treat a zero-sum dynamic as a strong and temporally consistent constraint

(Hubbell 2001; Houlahan et al. 2007). In this framing, any resources made available via species

loss should immediately be taken up by other species. This is not consistent with the dynamics

that occur at Portal, which has seen extended periods of time when resources are available on

exclosure plots but are not used. Rather, these results are more consistent with a zero-sum

constraint operating at metacommunity or evolutionary scales (Van Valen 1973; Terry and Rowe

2015; Leibold et al. 2017). Over short timescales, or within a closed local assemblage, niche

differences may weaken zero-sum effects, especially under fluctuating conditions. However,

over larger temporal and spatial scales, dispersal or evolution may supply new species equipped

to use available resources - via either functional redundancy, or niche complementarity allowing

them to exploit novel niches. A long-term, metacommunity, and even macroevolutionary

approach may be necessary to fully understand how zero-sum constraints, functional

redundancy, and niche complementarity contribute to the maintenance of community-level

function in the face of species extinctions and changing conditions over time.

Acknowledgements

The Portal Project has been supported by numerous grants from the U.S. National Science

Foundation, most recently NSF DEB-1929730. RMD was supported in part by the NSF Graduate

Research Fellowship under Grant No. DGE-1315138 and DGE-1842473.

Literature cited

Bledsoe, Ellen K., and S. K. Morgan Ernest. 2019. "Temporal Changes in Species Composition

Affect a Ubiquitous Species' Use of Habitat Patches." *Ecology* 100 (11): e02869.

https://doi.org/10.1002/ecy.2869.

Brown, J. H., T. J. Valone, and C. G. Curtin. 1997. "Reorganization of an Arid Ecosystem in

Response to Recent Climate Change." *Proceedings of the National Academy of Sciences* 94

(18): 9729–33. https://doi.org/10.1073/pnas.94.18.9729.

Christensen, Erica M., David J. Harris, and S. K. Morgan Ernest. 2018. "Long-Term Community

Change through Multiple Rapid Transitions in a Desert Rodent Community." *Ecology* 99

(7): 1523–29. https://doi.org/10.1002/ecy.2373.

Dornelas, M., N. J. Gotelli, B. McGill, H. Shimadzu, F. Moyes, C. Sievers, and A. E. Magurran.

2014. "Assemblage Time Series Reveal Biodiversity Change but Not Systematic Loss."

Science 344 (6181): 296–99. https://doi.org/10.1126/science.1248484.

Ernest, S. K. Morgan, and James H. Brown. 2001. "Delayed Compensation for Missing

Keystone Species by Colonization." *Science* 292 (5514): 101–4.

https://doi.org/10.1126/science.292.5514.101.

Ernest, S. K. Morgan, Glenda M. Yenni, Ginger Allington, Ellen K. Bledsoe, Erica M.

Christensen, Renata M. Diaz, Keith Geluso, et al. 2020. "The Portal Project: A Long-Term

Study of a Chihuahuan Desert Ecosystem." *BioRxiv*, January, 332783.

https://doi.org/10.1101/332783.

Ernest, S. K. Morgan, James H. Brown, Katherine M. Thibault, Ethan P. White, and Jacob R.

Goheen. 2008. "Zero Sum, the Niche, and Metacommunities: Long-Term Dynamics of

Community Assembly." *The American Naturalist* 172 (6): E257–69.

https://doi.org/10.1086/592402.

Fetzer, Ingo, Karin Johst, Robert Schäwe, Thomas Banitz, Hauke Harms, and Antonis

Chatzinotas. 2015. "The Extent of Functional Redundancy Changes as Species' Roles Shift

in Different Environments." *Proceedings of the National Academy of Sciences* 112 (48):

14888–93. https://doi.org/10.1073/pnas.1505587112.

Gonzalez, Andrew, and Michel Loreau. 2009. "The Causes and Consequences of Compensatory

Dynamics in Ecological Communities." *Annual Review of Ecology, Evolution, and*

Systematics 40 (1): 393–414. https://doi.org/10.1146/annurev.ecolsys.39.110707.173349.

Houlahan, J. E., D. J. Currie, K. Cottenie, G. S. Cumming, S. K. M. Ernest, C. S. Findlay, S. D.

Fuhlendorf, et al. 2007. "Compensatory Dynamics Are Rare in Natural Ecological

Communities." *Proceedings of the National Academy of Sciences. 104(9): 3273-3277*.

https://www.srs.fs.usda.gov/pubs/29712.

Hubbell, Stephen P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography (MPB-*

32). Princeton University Press. https://www.jstor.org/stable/j.ctt7rj8w.

Hughes, Brent B., Rodrigo Beas-Luna, Allison K. Barner, Kimberly Brewitt, Daniel R.

Brumbaugh, Elizabeth B. Cerny-Chipman, Sarah L. Close, et al. 2017. "Long-Term Studies

Contribute Disproportionately to Ecology and Policy." *BioScience* 67 (3): 271–81.

https://doi.org/10.1093/biosci/biw185.

Isbell, Forest, Vincent Calcagno, Andy Hector, John Connolly, W. Stanley Harpole, Peter B.

Reich, Michael Scherer-Lorenzen, et al. 2011. "High Plant Diversity Is Needed to Maintain

Ecosystem Services." *Nature* 477 (7363): 199–202. https://doi.org/10.1038/nature10282.

Kelt, Douglas A. 2011. "Comparative Ecology of Desert Small Mammals: A Selective Review

of the Past 30 Years." *Journal of Mammalogy* 92 (6): 1158–78. https://doi.org/10.1644/10-

MAMM-S-238.1.

Kelt, Douglas A., Jaclyn R. Aliperti, Peter L. Meserve, W. Bryan Milstead, M. Andrea Previtali,

and Julio R. Gutierrez. 2015. "Energetic Compensation Is Historically Contingent and Not

Supported for Small Mammals in South American or Asian Deserts." *Ecology* 96 (6):

1702–12. https://doi.org/10.1890/14-1569.1.

Lawton, John H. 1994. "What Do Species Do in Ecosystems?" *Oikos* 71 (3): 367–74.

https://doi.org/10.2307/3545824.

Leibold, Mathew A., Jonathan M. Chase, and S. K. Morgan Ernest. 2017. "Community

Assembly and the Functioning of Ecosystems: How Metacommunity Processes Alter

Ecosystems Attributes." *Ecology* 98 (4): 909–19. https://doi.org/10.1002/ecy.1697.

Lenth, Russell V. 2021. *Emmeans: Estimated Marginal Means, Aka Least-Squares Means*.

https://CRAN.R-project.org/package=emmeans.

Loreau, Michel. 2004. "Does Functional Redundancy Exist?" *Oikos* 104 (3): 606–11.

https://doi.org/10.1111/j.0030-1299.2004.12685.x.

Nagy, K. A., I. A. Girard, and T. K. Brown. 1999. "Energetics of Free-Ranging Mammals,

Reptiles, and Birds." *Annual Review of Nutrition* 19 (1): 247–77.

https://doi.org/10.1146/annurev.nutr.19.1.247.

Pinheiro, Jose, Douglas Bates, Saikat DebRoy, Deepayan Sarkar, and R Core Team. 2020. *Nlme:*

Linear and Nonlinear Mixed Effects Models. https://CRAN.R-project.org/package=nlme.

R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Rosenfeld, Jordan S. 2002. "Functional Redundancy in Ecology and Conservation." *Oikos* 98

(1): 156–62. https://doi.org/10.1034/j.1600-0706.2002.980116.x.

Terry, Rebecca C., and Rebecca J. Rowe. 2015. "Energy Flow and Functional Compensation in

Great Basin Small Mammals under Natural and Anthropogenic Environmental Change."

Proceedings of the National Academy of Sciences 112 (31): 9656–61.

https://doi.org/10.1073/pnas.1424315112.

Thibault, Katherine M., and James H. Brown. 2008. "Impact of an Extreme Climatic Event on

Community Assembly." *Proceedings of the National Academy of Sciences of the United*

States of America 105 (9): 3410–15. https://doi.org/10.1073/pnas.0712282105.

Thibault, Katherine M., S. K. Morgan Ernest, and James H. Brown. 2010. "Redundant or

Complementary? Impact of a Colonizing Species on Community Structure and Function."

Oikos 119 (11): 1719–26.

Van Valen, Leigh. 1973. "A New Evolutionary Law." *Evolutionary Theory* 1 (1): 1–30.

Walker, Brian. 1995. "Conserving Biological Diversity through Ecosystem Resilience."

Conservation Biology 9 (4): 747–52. https://doi.org/10.1046/j.1523-1739.1995.09040747.x.

Walker, Brian H. 1992. "Biodiversity and Ecological Redundancy." *Conservation Biology* 6 (1):

18–23. https://doi.org/10.1046/j.1523-1739.1992.610018.x.

White, Ethan P., S. K. Morgan Ernest, and Katherine M. Thibault. 2004. "Trade-offs in

Community Properties through Time in a Desert Rodent Community." *The American*

Naturalist 164 (5): 670–76. https://doi.org/10.1086/424766.

Williams, John W., and Stephen T. Jackson. 2007. "Novel Climates, No-Analog Communities,

and Ecological Surprises." *Frontiers in Ecology and the Environment* 5 (9): 475–82.

https://doi.org/10.1890/070037.

Figure legends

Figure 1. Dynamics of energy use and rodent community composition over time. Lines represent

the ratio of total energy use on exclosure plots to control plots (a), 6-month moving averages of

energetic compensation (b), and the share of community energy use accounted for by kangaroo

rats on control plots (c), and by *C. baileyi* (d), on control (gold) and exclosure (blue) plots.

Dotted vertical lines mark the boundaries between time periods used for statistical analysis.

Horizontal lines are time-period estimates from generalized least squares (a, b) and generalized

linear (c, d) models, and the semitransparent envelopes mark the 95% confidence or credible

intervals.

411
412

a. Total energy use ratio

b. Energetic compensation

c. Kangaroo rat proportional energy use

d. *C. baileyi* proportional energy use

Control
Exclosure

Date