Potential of breadfruit cultivation to contribute to climate-resilient low latitude food systems

Lucy Yang[1], Nyree Zerega[2,3], Anastasia Montgomery[4], and Daniel E. Horton[4,5]

[1]Program in Environmental Sciences, Northwestern University

[2]Program in Plant Biology and Conservation, Northwestern University

[3]Chicago Botanic Garden

[4]Department of Earth & Planetary Sciences, Northwestern University

[5]Institute for Sustainability and Energy at Northwestern University

Keywords

food security, climate change adaptation, bioclimactic niches, neglected and underutilized species (NUS),

CMIP6, species distribution modeling, breadfruit

Abstract

The number of people in food crisis around the world is increasing, exacerbated by the challenges of

COVID-19 and a rapidly changing climate. Major crop yields are projected to decrease in low-latitude

regions due to anthropogenic climate change, making tropical and sub-tropical food systems particularly

vulnerable to climate shocks. Increased cultivation of breadfruit (*Artocarpus altilis*), often categorized as

a neglected and underutilized species (NUS), has been suggested as an agricultural adaptation pathway

for food insecure tropical and subtropical regions, due to its potential to enhance climate resilience and

overall sustainability of low-latitude agricultural systems. To better understand breadfruit's cultivation

suitability and geographic range in current and future climates, we employ a diverse set of observations

and models to delineate the current climatically viable breadfruit range and assess the climatically viable

breadfruit range in the future (2061-2080) under stabilization and high emission scenarios. We find that

the area of suitable breadfruit range within the tropics and subtropics is projected to decrease ~4.4% in the

stabilization scenario and ~4.5% in the high emission scenario. In Southeast Asia and the Pacific Islands,

yield quality and consistency show minimal decreases under the high emission scenario, with increases in

total suitable area under both scenarios. In contrast, in Latin America and the Caribbean, the current range

of breadfruit suitability is projected to contract ~10.1-11.5% (stabilization-high emission). Present and

future model suitability outputs suggest that opportunities to successfully expand breadfruit cultivation

over the next several decades exist in sub-Saharan Africa, where food insecurity is coincidentally high.

However, in all regions, high emission scenario conditions reduce the overall consistency and quality of

breadfruit yields compared to the stabilization scenario. Our results have the potential to inform global

food security adaptation planning and highlight breadfruit as an ideal NUS to incorporate in food security

adaptation strategies in a changing climate.

1 Introduction

The number of people in the world lacking regular access to nutritious and sufficient food has been

on the rise since 2015, well before the COVID-19 pandemic exacerbated existing global food security

challenges and increased the prevalence of undernourishment (PoU) by +1.5% to ~9.9% in 2020 (1,2). Food

insecurity has increased in all global regions since 2014, except in North America and Europe, and the

world is considerably off track to achieve the UN's global goal of Zero Hunger by 2030 (3). Climate change

is a major driver of the observed rise in global hunger (4), with both gradual climatic change and punctuated

shocks working to multiply threats and disrupt the food system at various levels, most notably food

production and availability, but also in terms of access, quality, utilization, and stability (5). Climate

adaptation solutions are needed to lessen risk in the global agricultural production system, particularly in

regions with high populations of undernourished peoples (6).

In the face of changing climate conditions and increases in the frequency and intensity of extremes,

scaling up the production of NUS is an attractive strategy to build global food system resilience, particularly

in the tropics and sub-tropics. Studies suggest that the agricultural adaptations made between 1981-2010

were insufficient to offset the negative impacts of climate change on common staple yields and production

losses (e.g., wheat, rice, soybean, and maize), particularly at lower latitudes (7). While the geography of

these major crops may expand to higher-latitudes under various climate change scenarios over the 21st

century (8), production of major crops in low latitudes is projected to be consistently and significantly

negatively impacted (9–11), leading to a projected decrease in food availability and dietary diversity in

tropical and subtropical regions (12). Furthermore, many of the world's current food security challenges lie

within tropical latitudes, where anthropogenic warming is also projected to impact agricultural production

most substantially (6,12). These regions are particularly sensitive to increases in temperature due to low

historical variability, with the time of emergence (TOE) of unprecedented temperature and precipitation

regimes arriving earlier in terrestrial tropical biomes, compared to higher latitude regions, and potentially

crossing a signal-to-noise threshold of 2 as early as the 2020s (13–15). Tropical agricultural production has

already been adversely affected by extreme weather events and increased climatic variability, and yields

will likely worsen with increased temperature stress and rainfall variability (16). Given these realized and

projected impacts on staple crops, scaling up production of NUS has been suggested as an important

adaptation measure in food-insecure regions. Expansion of NUS agriculture has the potential to enhance

the climate resilience of food systems while simultaneously improving other negative environmental and

social characteristics of current global food systems, including improving global crop and varietal diversity,

increasing water efficiency, reducing excessive use of agrochemicals, increasing more locally produced

food, and implementing practices that protect and conserve traditional knowledge and biodiversity (6).

A traditional crop originating in Oceania, where it is an important and widely used staple, breadfruit

is a tropical NUS in other regions where it has a promising role in alleviating present and future food

insecurity and improving the overall sustainability and resiliency of tropical and sub-tropical food systems

(17,18). Breadfruit offers practical advantages in low-latitude regions as a nutritious "superfood" grown

from a long-lived tree that does not need annual replanting, requires lower energy input to cultivate

(especially when cultivated in sustainable agroforestry systems), sequesters carbon as a long-lived tree, and

can be used as a staple food in a variety of dishes and cultural cuisines (19). Breadfruit compares favorably,

in some cases surpassing, major staples (including wheat, rice, and corn) in both yield and nutritional

content (20). Breadfruit also shows great potential to reduce economic barriers to food access when

implemented as part of economically sustainable development strategies in food-insecure regions of the

world (21). As such, breadfruit has been presented as a potential solution to the world hunger crisis and

labeled a "superfood of the future" by popular media, although its potential implementation as a practical

adaptation to food insecurity under a changing climate has not been thoroughly explored. That said, its

potential for more intensive cultivation over its broad current range and expansion to new regions shows

promise as a climate adaptation strategy for sustainable food systems. Because the most commonly grown

breadfruit varieties are seedless and are clonally propagated via root cuttings, grafting, air layering, or

micropropagation methods, they have little to no likelihood of becoming invasive when introduced to new

regions. In fact, over the last more than ~200 years seedless varieties have been introduced to many tropical

regions with no recorded invasive behavior.

Breadfruit is a domesticated crop with a rich global history, connecting various cultures and regions.

Its wild, seeded ancestor is believed to have originated in New Guinea, and the seedless breadfruit was

domesticated in the islands of Oceania, where hundreds of cultivars exist (22). Breadfruit played a major

role as a key staple food in the human colonization of these islands. By about 3,000 years ago humans had

reached Samoa and few-seeded and seedless varieties began to appear. When Hawaii was one of the last

Pacific islands to be colonized by humans (between 1400 and 1700 years ago), seedless breadfruit was

brought by Polynesian voyagers to islands, where it is known as "ulu" (23). Breadfruit continues to play a

large part of cultural and spiritual life throughout Oceania. Europeans encountered breadfruit in Oceania in

the 1500s (24). By the 18th century it had been introduced to Central and South America, the Caribbean,

tropical Asia, and other tropical regions, and it remains especially important in the Caribbean today (25).

Given the success of breadfruit cultivation in various tropical and subtropical regions around the world,

farmers in sub-Saharan Africa are increasingly looking to breadfruit as a potential crop for alleviating local

food insecurity concerns (26). Despite its large promise, breadfruit remains more of a subsistence crop than

commercial commodity in most regions (e.g., parts of West Africa, Southeast Asia, Latin America, and

India) with major production areas limited to the Pacific and Caribbean Islands (Ragone, 2018). However,

interest in commercial uses is growing (27).

The potential expansion of breadfruit cultivation as an adaptation strategy is constrained by present

and future climatic uncertainties. A newly planted breadfruit tree takes several years before it begins fruiting

and may live and produce fruit for over 50 years—if grown within a suitable environment. While global

increases of temperature in some regions may intuitively seem to allow for the expansion of breadfruit into

areas of higher latitudes, uncertainties around extreme weather, increased variability, and rising

temperatures within the current cultivated extent may concurrently limit its future suitability in its current

range. However, breadfruit yields are still expected to be more resilient than important global staples like

rice to certain climate change stressors. For example, after a breadfruit tree is established, it may withstand

droughts of 3-4 months (28). The extent of breadfruit's resilience must be evaluated amidst an uncertain

climactic future.

To examine the adaptation potential of breadfruit cultivation, we evaluate current and projected

breadfruit niche suitability and assess the potential for range expansion and/or contraction using species

distribution modeling (SDM) and bioclimactic data. Given our knowledge of suitable climate conditions

for breadfruit cultivation, and our understanding that many of these underlying factors are undergoing

anthropogenically-driven changes, we use a weighted ensemble of six species distribution models (SDMs)

created using 7 selected bioclimactic variables (from BIOCLIM) to assess changes in future (2061-2080)

breadfruit suitability range using downscaled simulation data from 8 different General Circulation Models

(GCMs) from phase 6 of the Coupled Model Intercomparison Project (CMIP6) (29). SDMs that rely on

bioclimactic variables over the full extent of a species range facilitate a realization of the fundamental niche

rather than a realized niche. Consideration of seasonality and variability is useful in determining the

fundamental niche of plant species, particularly when an SDM is applied to future climatic conditions. This

approach allows us to assess the potential range of breadfruit in our current climate where it has not yet

been introduced, as well as extrapolate breadfruit cultivation suitability in future climates. This ensemble

SDM approach in conjunction with ensemble GCM projections allows us to evaluate the inherent inter-

model differences/uncertainties in both steps of the process. Gaining an understanding of the SDM and

climatic uncertainties that constrain breadfruit's potential global range can facilitate policy design intended

to mobilize new adaptation strategies for more resilient low-latitude food systems.

2 Materials and Methods

2.1 Species Distribution Modeling (SDM)

To determine current and future breadfruit niche suitability, we used observations and

bioclimactic data to constrain 6 unique SDMs: Generalized Linear Model (GLM), Generalized Additive

Model (GAM), Multiple Adaptive Regression Splines (MARS), Random Forest (RF), Generalized

Boosted Model (GBM), and Maximum Entropy (MAXENT) using the *Biomodhub/Biomod2* software

version 3.4.13 (2018/2021). A weighted-mean ensemble of SDMs was used to determine current

breadfruit niche suitability. The ensemble approach to species distribution modeling has gained

considerable popularity and is useful for examining which bioclimatic variables determine the

fundamental niche of breadfruit, allowing us to project this model to future scenarios and evaluate the

changed distribution of global breadfruit suitability (31).

To constrain the current distribution of breadfruit and provide training data for our SDMs, we use

Global Biodiversity Information Facility (GBIF) historical breadfruit (*Artocarpus altilis*) occurrence point

data (32–35). GBIF presence points are prone to sampling bias because it allows citizen science

contributions to the database. Citizen scientists are prone to misidentify *A. altilis*, particularly by

confusing the plant with its close relative, seeded breadnut (*A. camansi*). To limit this bias, we

downloaded only those data points that were identified as part of herbarium specimens ("Preserved

specimen") or that included picture documentation. We also filtered GBIF presence points using a

number of screens. First, presence data that appeared under alternative names (taxonomic synonyms),

e.g., *A. communis, A. incisa,* and *A. incisus*, were included. However, data of two close relatives (i.e., *A.*

camansi and *A. mariannensis*) were not included (36). Next, we ran the GBIF points using default settings

of the CoordinateCleaner, an R-package tailored for cleaning occurrence records from biological

collection databases (37). This package flags geographical coordinates that are overly imprecise or have

common problems (e.g., points assigned to country centroids due to automated geo-referencing, plants

belonging to botanical gardens or museums, etc.). We also added datapoints from a yield study in Jamaica

(Trees that Feed Foundation, 2020, unpublished data). Finally, we discarded duplicate points within

common grid cells to limit spatial bias. Ultimately, over half of the breadfruit presence points downloaded

from GBIF were removed. The remaining 431 presence points are used for the SDM ensemble training

and prediction.

To determine the climatic variables that best predict the occurrence of current breadfruit

distribution, we used the dataset of bioclimatic variables (BIOCLIM) for 1970-2000 from WorldClim

Version 2.1, a database of high-resolution global climate data (39). To prevent overfitting of the final

SDM, we examined the full BIOCLIM dataset of 19 variables and selected a subset of variables based on

their predictive performance in individual and paired runs. We selected variables largely following the

methodology detailed in Moat et al., 2017 using GLM and GAM, two of the six SDMs used for the final

ensemble. Boxplots, preliminary map projections, and evaluative metrics including sensitivity, specificity,

True Skills Statistics (TSS), relative operating characteristic curve (ROC), and Cohen's Kappa statistic

(KAPPA) were used to determine which combination of variables were best for model fit, i.e., which

combination of bioclimatic variables best predicted the observed range of breadfruit. After evaluating

variable performance, we tested resulting variable combinations for collinearity and multi-collinearity

using a threshold of R^2=0.8 and reduced the selection to 7 variables (Table 1). Annual mean temperature

(BIO7) and annual precipitation (BIO12) were among the highest performing variables in the preliminary

model runs using GLM and GAM. While environmental characteristics such as solar radiation and soil

quality also impact breadfruit growth, they are not used to constrain breadfruit suitability in this study.

The optimal combination of variables to characterize niche suitability differs based on the spatial

domain. A preliminary version of the model covered the full extent of global BIOCLIM data at 10

arcminute resolution (~100 km). However, because over 97% of our preliminary weighted model results

were constrained within the latitudes of 45° N to 45° S, the final analysis presented in this study was

performed within these lesser latitudinal bounds. Despite this change, our analysis covers almost the

entire extent of the current breadfruit range, which is ideal for accurately capturing the fundamental niche

of the species (41). A lesser geographical extent also increases the reliability of pseudoabsence data. We

use a presence-only SDM that relies on pseudoabsence (background) points to model suitability because

no reliable source of true absence data was available. The TSS scores of the three sets of pseudoabsence

points we generated vary by 0.05. Although we latitudinally limited the geographic extent of the SDM in

favor of generating more accurate pseudoabsence points, it is still difficult to produce useful

pseudoabsence data at this large extent, as evidenced by the concentration of pseudoabsence points in

grid-cells of less-suitable categories (Supplementary Figure 1).

We ran the 7 selected BIOCLIM variables and 431 filtered data points using the Biomod2 R

package version 3.4.14 using six independent SDM methods (GLM, GAM, MARS, RF, GBM, and

MAXENT.Phillips). We followed established protocols for model evaluation, execution, replication, and

generation of three sets of pseudo-absence data (42). Presence and pseudoabsence data were split

randomly with a ratio of 70:30 to build training and testing data separately over ten runs. The same

metrics of sensitivity, specificity, TSS, ROC, and KAPPA were used to evaluate model outputs.

Resultant maps and statistics from model outputs were used to evaluate the performance of the

SDMs. Output statistics were reviewed to ensure a lack of difference between model runs, which could

indicate problematic outcomes, such as overfitting. After evaluation, the final SDM configuration was

used to produce our suitability predictions from ten runs and three replicates. The TSS threshold of 0.7

was used to remove poorly performing runs from all five models before producing a mean ensemble

prediction weighted by TSS score. Our weighted ensemble of SDMs demonstrates robust present-range

predictive power, characterizing the distribution of breadfruit presences indicated by GBIF data while

also predicting broader distribution within Africa and other previously unexamined regions with high

sensitivity (0.96) and reasonable accuracy (specificity = 0.83). The included SDMs in this study meet

established criteria (TSS = 0.79, KAPPA = 0.81, ROC = 0.93).

2.2 General Circulation Model (GCM) Data

Projected climate data was then used to determine niche expansion/contraction under future

climate scenarios SSPs 2 and 5 (43,44). We modeled the future projections over the time period of 2061-

2080 using the Biomod2 package Ensemble Forecasting. The projection was performed with the same

seven selected BIOCLIM variables from the GCM data under two shared socioeconomic pathway

scenarios: SSP2-4.5 (stabilization GHG emission scenario) and SSP5-8.5 (high GHG emission scenario)

forcing scenarios from 8 different GCMs. The GCMs from CMIP6 were used based on availability in

WorldClim 2.1 and include BCC-CSM2-MR, CNRM-CM6-1, CNRM-ESM2-1, CanESM5, IPSL-CM6A-

LR, MIROC-ES2L, MIROC6, and MRI-ESM2-0.

2.3 Analysis and Mapping

The SDM outputs from the current (Figure 1) and the two future scenarios were processed and

categorized into three: Good, Fair, and Unsuitable. Niche classes were chosen based on natural inflection

points and breaks between pseudoabsence and presence data (see Supplementary Figure 1). The division

between "Fair" and "Good" categories was placed at 50% to approximately reflect the inflection point of

the data. The Good niche classifications assesses breadfruit suitability based on yield quality and

consistency, marking locations that are conducive to the establishment of new breadfruit trees.

Meanwhile, the Fair range is more liberal and represents climactic conditions that established trees will

likely withstand, ergo persistence is assumed likely.

3 Results

3.1 SDM Prediction of Current Breadfruit Suitability Range

Our weighted ensemble baseline SDM predicts niche suitability in all GBIF-identified locations of

current breadfruit cultivation, as well as substantial swathes of the global tropics and subtropics where

successful cultivation of breadfruit has not yet been confirmed. Predicted suitable areas for cultivation

include large sections of Southeast Asia, Central America, northern South America, the Caribbean, and

equatorial Africa (Figure 1a). Latitudinal limits of modeled suitability extend as far north as northeastern

India in Asia and southern Florida in North America and stretch as far south as Chile in South America and

New Zealand in the Pacific. These regions have previously been identified by the National Tropical Botanic

Garden (NTBG) as suitable for breadfruit cultivation based on current climatic conditions (45). Suitable

niches that extend into the higher latitudes of the tropics are largely concentrated along coastlines, in

agreement with available GBIF occurrence data (Figure 1a; black dots). Notable areas without widespread

occurrence data that are identified as having robust potential for breadfruit cultivation include Cuba,

equatorial Africa, the southeastern coastline of Africa, and the Malay peninsula. In Africa, the lack of

occurrence data in GBIF may be explained by breadfruit's recent arrival to the continent, but recent

successful introductions haven been made in Uganda, Kenya, Tanzania, and Ghana. While no GBIF records

are recorded in Cuba, it is known to grow there (26,46,47).

Our SDM breadfruit suitability predictions employ seven weighted bioclimatic variables to

characterize the climatic conditions within a grid cell. The environmental variables were selected based on

evaluation scores on the Generalized Linear Models and Generalized Additive Models while controlling

for multicollinearity and overfitting (see Methods). Contribution weights of the seven bioclimatic variables

are quantified using True Skill Statistic (TSS) scores. We find that baseline breadfruit suitability is largely

driven by three temperature-related bioclimatic variables (TSS: 60.9). The annual temperature range,

defined as the difference between the maximum temperature of the warmest month and the minimum

temperature of the coldest month is the predominant prediction contributor (Figure 4c; TSS: 40.1).

Secondary contributors include the mean diurnal temperature range (Figure 4a; TSS: 9.2) and the mean

temperature of the coldest month (Figure 4g; TSS: 11.5). The four remaining contributing variables – annual

precipitation, and precipitation during the driest, warmest, and coldest quarters – are lesser contributors

relative to the temperature variables (Figure d-g; TSS: 6.1-7.0).

To refine our suitability findings and identify the niches most suitable for cultivation, we employ

statistical thresholding to subdivide the SDM suitability scores into Unsuitable, Fair, and Good categories

(Supplementary Figure 1). The unsuitable niche classification marks areas where breadfruit cultivation is

barely possible to impossible, whereas fair classification marks areas where breadfruit cultivation is

possible but yield and quality may be negatively affected by adverse seasonal weather conditions (e.g., low

rainfall or high temperatures). Good niche classification indicates where breadfruit cultivation has the

potential for consistent, high quality yields that are less likely to be affected by adverse weather conditions.

For the lower threshold, between unsuitable and fair, we set the cut-off at the inflection point where an

increase in model suitability values and the cumulative percentage of all (including pseudoabsence and

occurrence) points plateaus, which resulted in the loss of four ground control points. The division between

Fair and Good categories indicates a second inflection point where model suitability values begin to

substantially increase.

Individual SDMs are remarkably consistent in their suitable geographic range predictions (Figure

1b-g) but show notable differences in categorizing the level of breadfruit suitability within these ranges.

Across the 6 SDMs, there is only a 3.3% difference in the total "suitable" land area, although the

classification of suitable land into either Fair or Good is less consistent, with ~33.1-64.0% of the total

suitable area identified as Fair depending on SDM and ~36.0-66.9% classified as Good. When considering

the total modeled area of suitability, both MAXENT (35.9% ,10.0M km^2 Good; 64.1%, 17.7M km^2 Fair)

and GAM (44.0%, 12.5M km^2 Good; 56.0%, 15.8M km^2 Fair) predict lesser "Good" areas in proportion to

"Fair", while RF (66.9%, 8.4M km^2 Good; 33.1%, 17.0M km^2 Fair) and GBM (63.9%, 16.2M km^2 Good;

36.1%, 9.1M km^2 Fair) predict greater "Good" suitability (Figure 1). Because MAXENT is a presence-only

SDM, it is known to underestimate the probability of occurrence within areas of observed presence, while

overestimating in areas without known presence (48). Several MAXENT runs (6 out of 30) were removed

from the final weighted SDM ensemble due to low TSS scores, likely due to a lack of convergence.

Marginal areas of Fair suitability are consistently identified by all 6 SDMs in the same locations (e.g., New

Zealand, southwestern coast of Africa, Chile). Southeast Asia, Oceania, and the Pacific Islands are also

consistently valued with the highest proportion of Good to Fair suitability. The 6 SDMs demonstrate

consistent suitability coverage and patterns in regional Good to Fair suitability, although specific values

differ.

Using the weighted-ensemble projection, we focus our narrative on comparative breadfruit

suitability in three main regions: Asia and the Pacific Islands, the Caribbean and Latin America, and Africa.

Compared to the averaged global proportion of area with Good (51.9%, ~14M km^2) to Fair (48.1%, ~12M

km^2) suitability (Figure 3), breadfruit is particularly suited to climactic conditions in Southeast Asia.

Breadfruit is suitable for cultivation within 80.7% of the area (~3.6M km^2) in this region with a high

proportion of Good (77.4%, ~2.8M km^2) to Fair (22.6%, ~800k km^2) quality, although Pacific Islands such

as Hawaii also show similarly high levels of Good (87.7%, ~30k km^2) to Fair (12.3%, ~4k km^2) area . In

Latin America and the Caribbean, island countries show high coverage of suitable area (99.2%, ~2.3M km^2)

while 51.7% (~11.7M km^2) of the overall land area in Latin America and the Caribbean is projected to be

suitable. 41.3% of these suitable areas are classified as Good and 58.7% as Fair. In Africa, 20.8% of the

continent is projected to be suitable for growing breadfruit, with 45.4% of this land showing Good and 54.6%

showing Fair suitability. This baseline SDM prediction gives a comprehensive picture of the range and

quality of suitable breadfruit-growing areas in various low-latitude regions of the world.

Change in Breadfruit Range due to Climate Change

To assess the effects of climate change on breadfruit suitability, we project our SDM to future (2061-1080) climate conditions using an ensemble of opportunity of 8 GCMs from CMIP6. While breadfruit is projected to largely maintain its present range, the resilience and adaptation potential is projected to decrease, with more substantial decreases in Good area projected from the high-emissions scenario (44.4%) than the stabilization scenario (32.1%) (Figure 2). We find a 4.7% decrease in weighted ensemble SDM suitable breadfruit range (~1.2M km^2) in the stabilization scenario and a 4.5% decrease (~1.2M km^2) under high emission scenario conditions (Figure 3). However, the total area of "winners" and "losers" in terms of overall breadfruit suitability changes largely balance on a global scale.

Change in suitable breadfruit range – expansion, contraction, and persistence— from the baseline projection varies considerably across regions (Figure 2). The suitable area primarily persists and even increases ~0.7% (~23k km^2) under the stabilization scenario and 0.8% (~30k km^2) under the high emission scenario in Southeast Asia, although this increase in area is accompanied by a decrease in quality (2.6% and 8.7%) and reduction in total Good area. In Asia and the Pacific Islands more broadly, total suitable breadfruit area increases ~2.8% (~144k km^2) and ~4.1% (~214k km^2), showing areas of potential expansion in the upper and lower latitudinal limits of this region (Figure 2C, D). The majority of suitable breadfruit-growing area is projected to persist in equatorial Africa, although with lower resilience potential to future climate conditions than in Southeast Asia (1.5-3.4% reduction in total suitable area and 58.2-70.0% reduction in Good area depending on emissions scenario). In both the stabilization and high-emission scenarios, the baseline suitability range within equatorial Africa expands northward while the southern bounds of breadfruit suitability are projected to contract. However, Latin America and the Caribbean are projected to experience the largest magnitude range reduction on both the stabilization (10.1%) and high emission pathways (11.5%). These range reductions appear to be concentrated in specific areas, particularly in the Yucatan peninsula and southwestern Brazil. While Caribbean islands experience minimal contraction of the overall suitable breadfruit-growing area (1.4-1.6% reduction) under either emission pathway, the quality of these conditions may decrease significantly (50.6-72.5% reduction in Good area). Despite regional variation, a large portion of global suitable breadfruit range is projected to persist in both scenarios

(particularly in Africa and Asia and the Pacific Islands), suggesting that (1), established trees will likely

withstand future climatic conditions in most of the tropics and subtropics and (2) there is room for expanded

cultivation of breadfruit across a broad section of the global tropics and subtropics in Africa. However, a

greater decrease in quality but not necessarily a greater decrease in total area of suitability is projected under

the high emissions scenario in all regions.

Projected expansion of breadfruit suitability is driven by increases in seasonal temperature and

precipitation. Increased precipitation during the coldest quarter (BIO19, Fig. 4t, u) as well as general

increases in annual precipitation (BIO12, Fig. 4k,l) drive potential expansion of suitable breadfruit area at

the perimeter of the baseline suitable range. Meanwhile, decreases in precipitation of the warmest quarter

(BIO18) drive contraction of suitable breadfruit range in Central and South America (Fig. 4q, r). Mean

Diurnal Range shows little change in Southeast Asia but increases in areas of expansion and decreases in

areas of contraction and reduced breadfruit suitability (Fig. 4b, c).

To characterize the structural uncertainty of our breadfruit projections, we use our CMIP6 ensemble

of opportunity and define projection confidence according to ensemble member agreement (i.e., 4-5

members agree – Strong, 6+ members agree = Very Strong). Projections of future suitability show Very

Strong GCM agreement over the majority of the persistent range of breadfruit suitability under the

stabilization scenario (92.8%) and high-emission scenario (90.5%), with a larger area of weak GCM

agreement under the high-emission scenario. Weak SDM agreement (<4 members agree) only characters

5.1% of the persisting suitable range under the high-emission scenario and 3.6% of the suitable range under

the stabilization scenario. Strong and Weak agreement is concentrated at the outer fringes of breadfruit

distribution and most significantly in central South America in the Mato Grosso Plateau and eastern Africa

in the Ethiopian highlands (Figure 3). This finding suggests higher levels of uncertainty in areas of higher

elevation, although breadfruit suitability is projected to persist in the Andes in South America. Areas of

potential expansion also overlap with the areas of greatest uncertainty. In future scenarios, South Asia and

South America exhibit the greatest variability in area among GCMs (Figure 4). However, the majority of

future global breadfruit cultivation in both scenarios demonstrates strong (4-5) to Very Strong (6+)

agreement among the 8 GCMs in both scenarios.

Discussion

Our SDM prediction captures known locations of breadfruit while also suggesting potential

expansion for the crop into food-insecure regions, most notably within Sub-Saharan Africa. While previous

studies have attempted to constrain global and regional distributions of breadfruit suitability (18,45), direct

comparisons are challenging due to differences in scoring systems. However, the geographic range of

predictions from other studies are in good agreement with our ensemble SDM predictions. For example,

Lucas and Ragone (2012) used Geographic Information Systems (GIS) to produce a map of global

breadfruit suitability using WorldClim v1.4 data and ecologically determined ranges for mean temperature

and rainfall (Ragone, 2006). While this study did not assess the impacts of climate change, their results on

the current distribution of suitable cultivation range is in good agreement with our SDM-based findings.

Mausio et al. (2020) further used fuzzy-set models to extrapolate breadfruit suitability from environmental

parameters derived in Hawaii. Their work offers a useful comparison of an alternative approach to modeling

breadfruit suitability by incorporating FAO EcoCrop parameters and climate data from CMIP5, but it does

not address the impact of seasonal temperature and precipitation data on potential breadfruit distribution

under future scenarios.

Our SDM indicates that the differences between the high-emissions scenario and stabilization

scenario in terms of total breadfruit suitability area is minimal, however the high-emission scenario is

predicted to lead to more substantial reductions in suitability scores in most regions. These findings are

particularly important for tropical and subtropical regions (low to mid latitude regions) where impacts on

the production of conventional staple crops such as maize, wheat, soybean and rice are projected to be more

negative than higher latitudes (49). Expanded cultivation of breadfruit in such regions may contribute to

global food and economic security, as a locally produced staple food that can lead to greater independence
from food imports.

Breadfruit shows the most promising resilience to future climactic conditions within its native
extent in Asia, largely persisting in Southeast Asia and even showing potential areas of expansion within
India and China. Culturally, breadfruit is also a significant traditional crop in many Pacific Islands including
Hawaii, where it has been noted for its promising role as part of local biocultural restoration efforts
(Langston and Lincoln, 2018). Breadfruit can play an important role in local low-latitude food systems,
alleviating the yield losses of major crops such as rice, while promoting sustainable agroforestry systems.
However, the persistence of suitable conditions for growing breadfruit are most at risk in specific regions
within Latin America, namely a section of the Yucatan peninsula and southwestern Brazil in the Mosso
Plateau. These areas experience decreases in annual precipitation, with significant decreases in precipitation
during the warmest quarter of the year. These areas also coincide with weak agreement among the 8 GCMs,
with less than half agreeing on the magnitude of precipitation reduction.

Our SDM predictions also signal opportunities to expand cultivation in Central Africa, building on
nascent cultivation of breadfruit which currently exists in parts of West and East Africa (26). Given the
present state of food insecurity in Africa, the widespread cultivation of breadfruit offers a path towards
increased food security and independence in the present and more distant future for several Sub-Saharan
countries. As a promising NUS, breadfruit could fill immediate and future needs to increase food security
in many tropical regions, especially as crops such as maize and rice become less suited to warming
conditions (11,50). Breadfruit demonstrates resilience to uncertain climactic conditions in many tropical
food-insecure regions. This adaptation potential is further enhanced if breadfruit trees are well-established
and grown within sustainable agroforestry systems. Agroforestry systems can serve as both adaptation and
mitigation, contributing to food productivity while enhancing restoration by combining traditional growing
methods used by Pacific Islanders for centuries with modern horticultural techniques (6,51).

Our study uses species distribution modeling to examine potential areas for breadfruit cultivation
over the next several decades under two emission pathways. Our model prediction is limited to bioclimactic

401 factors (seasonal temperature and precipitation variables) and does not include the potential influence of

402 soil type, solar radiation, or elevated CO_2 concentrations on breadfruit suitability. We are confident that the

403 45° N to 45° S swath covers the majority (~97%) of present and future suitable breadfruit range under

404 these scenarios, capturing the fundamental niche and identifying new areas where breadfruit can be

405 introduced or expanded globally. While additional presence observations may improve the performance of

406 our model, most studies agree that sample size effects usually become less critical above 50 presence points

407 (Guisan et al., 2017). Ideally, models of habitat suitability should also be based on measurements of

408 population fitness at each geographic location. However, this would prevent the use of reliable data

409 available in natural history collections. Although data points from GBIF were cleaned and filtered, absence

410 data and rigorous groundtruthing would further improve the performance of the individual models that

411 made up the final SDM ensemble. When projecting to future conditions, we were limited to using

412 realizations from relatively few (8) CMIP6 GCMs from WorldClim v.2.1 which likely limits the range of

413 internal variability and structural uncertainty considered in our projections.

414   Significant portions of sub-Saharan Africa show room for expansion of breadfruit cultivation over

415 the next several decades, especially as rapid population growth is projected to intensify food insecurity and

416 undernourishment across the continent (52). Closing the yield gap via sustainable intensification of

417 agriculture which utilizes NUS such as breadfruit may be critical in preventing future catastrophic

418 circumstances. Prioritizing a rich diversity of NUS over staple crops such as wheat, maize, and rice will

419 lead to healthier diets and improved nutrition and contribute to sustainable development (53). Similar

420 studies have examined the potential distribution of NUS under climate change, including 4 neglected and

421 underutilized fruit species (NUFS) in Sri Lanka and a yam species in Benin (54,55), but few have examined

422 the viability of expanding cultivation of NUS across different continents at a global scale. As a long-lived

423 crop that can be grown and used locally, breadfruit offers an opportunity for various low-latitude

424 communities around the world to improve the resilience of their local food systems. By building

425 agroforestry systems around breadfruit, low-latitude regions can avoid the pitfalls of strictly monoculture

426 systems while building sustainable food systems with increased resilience to climate change.

4.4 Conclusion

429   Our model of breadfruit niche suitability has the potential to provide projections useful for both short-term decision making and long-term adaptation planning. Because breadfruit cultivation has not yet expanded into its full suitable range, this crop has the potential to alleviate present and future food security burdens, particularly in food-insecure countries in equatorial Africa. Although breadfruit growers may need to contend with a certain level of reduced quality of yields in the future, suitable climatic conditions for breadfruit cultivation exists across a broad range of the tropics and subtropics and persist for the next several decades regardless of emissions pathway, indicating that it is a reliable and promising NUS to implement as a climate adaptation strategy. Depending on scale of production, breadfruit itself may even play a role in contributing to climate change mitigation due to its significant carbon sequestration benefits.

Acknowledgments

Research presented in this paper was supported by Summer Undergraduate Research and Conference Travel Grants from Northwestern University's Office of the Provost to LY. The authors declare that there is no conflict of interest.

References

1. Diffenbaugh NS, Field CB, Appel EA, Azevedo IL, Baldocchi DD, Burke M, et al. The COVID-19 lockdowns: a window into the Earth System. Nat Rev Earth Environ. 2020 Sep;1(9):470–81.

2. FAO, IFAD, UNICEF, WFP, and WHO. The State of Food Security and Nutrition in the World 2021. Rome: FAO; 2021.

3. Global Network Against Food Crises, FSIN. Global Report on Food Crises [Internet]. 2020 [cited 2021 Mar 29]. Available from: https://www.wfp.org/publications/2020-global-report-food-crises

4. FAO, IFAD, UNICEF, WFP, WHO. The State of Food Security and Nutrition in the World: Building climate resilience for food security and nutrition. Rome, Italy: FAO; 2018. 181 p.

5. Vermeulen SJ, Campbell BM, Ingram JSI. Climate Change and Food Systems. Annual Review of Environment and Resources. 2012;37(1):195–222.

6. Mbow C, Rosenzweig C, Barioni LG. Food Security. In: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems [Internet]. 2019 [cited 2021 Mar 29]. Available from: https://www.ipcc.ch/srccl/chapter/chapter-5/

7. Iizumi T, Shiogama H, Imada Y, Hanasaki N, Takikawa H, Nishimori M. Crop production losses associated with anthropogenic climate change for 1981–2010 compared with preindustrial levels. International Journal of Climatology. 2018;38(14):5405–17.

8. Porter JR, Xie L, Challinor AJ, Cochrane K, Howden SM, Iqbal MM, et al. Food security and food production systems. In: Climate Change 2014: Impacts, Adaptation, and Vulnerability Part A:

Global and Sectoral Aspects Contribution of Working Group II to the Fifth Assessment Report of

the Intergovernmental Panel on Climate Change. 2014. p. 488–9.

9.   Brown ME, Funk CC. Food Security Under Climate Change. Science. 2008 Feb 1;319(5863):580–

1.

10.   Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL. Prioritizing Climate

Change Adaptation Needs for Food Security in 2030. Science. 2008 Feb 1;319(5863):607–10.

11.   Zhao C, Liu B, Piao S, Wang X, Lobell DB, Huang Y, et al. Temperature increase reduces global

yields of major crops in four independent estimates. PNAS. 2017 Aug 29;114(35):9326–31.

12.   WFP. Climate impacts on food security and nutrition: a review of existing knowledge [Internet].

2014 [cited 2021 Mar 29]. Available from: https://www.eldis.org/document/A73884

13.   Diffenbaugh NS, Scherer M. Observational and model evidence of global emergence of permanent,

unprecedented heat in the 20th and 21st centuries. Climatic Change. 2011 Aug 1;107(3):615–24.

14.   Tan X, Gan TY, Horton DE. Projected timing of perceivable changes in climate extremes for

terrestrial and marine ecosystems. Global Change Biology. 2018;24(10):4696–708.

15.   Hawkins E, Sutton R. Time of emergence of climate signals: TIME OF EMERGENCE OF

CLIMATE SIGNALS. Geophys Res Lett. 2012 Jan;39(1):n/a-n/a.

16.   Ericksen PJ, Thornton PK, Notenbaert AMO, Cramer L, Jones PG, Herrero MT. Mapping hotspots

of climate change and food insecurity in the global tropics [Internet]. CGIAR Research Program on

Climate Change, Agriculture and Food Security; 2011 Jun [cited 2021 Mar 29]. Available from:

https://cgspace.cgiar.org/handle/10568/3826

17.   Turi CE, Liu Y, Ragone D, Murch SJ. Breadfruit (Artocarpus altilis and hybrids): A traditional crop

with the potential to prevent hunger and mitigate diabetes in Oceania. Trends in Food Science &

Technology. 2015 Oct 1;45(2):264–72.

18.   Mausio K, Miura T, Lincoln NK. Cultivation potential projections of breadfruit (Artocarpus altilis)

under climate change scenarios using an empirically validated suitability model calibrated in

Hawai'i. PLOS ONE. 2020 May 22;15(5):e0228552.

19.   Orwa C, Mutua A, Kindt R, Simons A, Jamnadass R H. Agroforestree database: a tree reference and

selection guide version 4.0. Nairobi, Kenya: World Agroforestry Centre (ICRAF); 2010. 1 p.

20.   Jones AMP, Ragone D, Aiona K, Lane WA, Murch SJ. Nutritional and morphological diversity of

breadfruit (Artocarpus, Moraceae): Identification of elite cultivars for food security. Journal of Food

Composition and Analysis. 2011 Dec 1;24(8):1091–102.

21.   Omobuwajo TO. BREADFRUIT AS A KEY COMPONENT OF SUSTAINABLE LIVELIHOODS

IN NIGERIA: PROSPECTS, OPPORTUNITIES AND CHALLENGES. Acta Hortic. 2007

Nov;(757):121–4.

22.   Zerega N, Wiesner-Hanks T, Ragone D, Irish B, Scheffler B, Simpson S, et al. Diversity in the

breadfruit complex (Artocarpus, Moraceae): genetic characterization of critical germplasm. Tree

Genetics & Genomes. 2015 Jan 20;11(1):4.

23.   Kirch PV. On the Road of the Winds: An Archaeological History of the Pacific Islands before

European Contact, Revised and Expanded Edition [Internet]. 2nd ed. University of California Press;

2017 [cited 2021 Sep 23]. Available from: https://www.jstor.org/stable/10.1525/j.ctv1xxsng

24.   Markham C. The voyages of Pedro Fernandez de Quiros, 1595. Scottish Geographical Magazine.

1904 Dec 1;20(12):649–52.

25. Ragone D. Breadfruit.(Parkinson) Fosberg. Promoting the conservation and use of underutilized and

neglected crops. 10. 1997;

26. Trees That Feed Foundation. Africa [Internet]. [cited 2021 Sep 7]. Available from:

https://www.treesthatfeed.org/countries/africa/

27. Peters A. Why Patagonia wants you to eat breadfruit [Internet]. Fast Company. 2021 [cited 2021

Sep 7]. Available from: https://www.fastcompany.com/90620093/youve-probably-never-eaten-

breadfruit-heres-why-patagonia-wants-to-change-that

28. Taylor M, McGregor A, Dawson B. Vulnerability of Pacific Island agriculture and forestry to

climate change [Internet]. 2016 [cited 2021 Mar 29]. Available from: http://www.fao.org/family-

farming/detail/en/c/432913/

29. Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ, et al. Overview of the Coupled

Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization.

Geoscientific Model Development. 2016 May 26;9(5):1937–58.

30. biomodhub/biomod2 [Internet]. biomod2 package, tools and scripts; 2021 [cited 2021 Mar 29].

Available from: https://github.com/biomodhub/biomod2

31. Hao T, Elith J, Guillera-Arroita G, Lahoz-Monfort JJ. A review of evidence about use and

performance of species distribution modelling ensembles like BIOMOD. Diversity and

Distributions. 2019;25(5):839–52.

32. GBIF Occurrence Download [Internet]. GBIF.org; [cited 2020 Nov 11]. Available from:

https://doi.org/10.15468/dl.mas6f8

33. GBIF Occurrence Download [Internet]. GBIF.org; [cited 2019 Jul 15]. Available from:

https://doi.org/10.15468/dl.fcoapd

34. GBIF Occurrence Download [Internet]. GBIF.org; [cited 2019 Jul 15]. Available from:

https://doi.org/10.15468/dl.jae0fb

35. GBIF Occurrence Download [Internet]. GBIF.org; [cited 2020 Nov 11]. Available from:

https://doi.org/10.15468/dl.x5ubgu

36. Zerega N, Ragone D, Motley TJ. 10. Breadfruit Origins, Diversity, and Human-Facilitated

Distribution. In: Motley T, editor. Darwin's Harvest [Internet]. Columbia University Press; 2006

[cited 2021 Aug 1]. p. 213–38. Available from:

https://www.degruyter.com/document/doi/10.7312/motl13316-011/html

37. Zizka A, Silvestro D, Andermann T, Azevedo J, Ritter CD, Edler D, et al. CoordinateCleaner:

Automated Cleaning of Occurrence Records from Biological Collections [Internet]. 2020 [cited

2021 Mar 29]. Available from: https://CRAN.R-project.org/package=CoordinateCleaner

38. Trees that Feed Foundation. Breadfruit Yield Study in Jamaica, 2018-20 [Internet]. 2020 [cited

2021 Sep 28]. Available from: https://treesthatfeed.org/breadfruit-yield-study/

39. Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global land

areas. Int J Climatol. 2017 Oct;37(12):4302–15.

40. Moat J, Williams J, Baena S, Wilkinson T, Gole TW, Challa ZK, et al. Resilience potential of the

Ethiopian coffee sector under climate change. Nature Plants. 2017 Jun 19;3(7):1–14.

41. Guisan A, Thuiller W, Zimmermann NE. Habitat Suitability and Distribution Models: With

Applications in R [Internet]. Cambridge: Cambridge University Press; 2017 [cited 2021 Sep 28].

(Ecology, Biodiversity and Conservation). Available from:

https://www.cambridge.org/core/books/habitat-suitability-and-distribution-

models/A17F74A3418DBF9ADA191A04C35187F9

42.   Habitat suitability and distribution models applications r | Ecology and conservation | Cambridge

551   University Press [Internet]. [cited 2021 Mar 29]. Available from:

552   https://www.cambridge.org/us/academic/subjects/life-sciences/ecology-and-conservation/habitat-

553   suitability-and-distribution-models-applications-r?format=PB&isbn=9780521758369

554   43.   Fricko O, Havlik P, Rogelj J, Klimont Z, Gusti M, Johnson N, et al. The marker quantification of

555   the Shared Socioeconomic Pathway 2: A middle-of-the-road scenario for the 21st century. Global

556   Environmental Change. 2017 Jan 1;42:251–67.

557   44.   Kriegler E, Bauer N, Popp A, Humpenöder F, Leimbach M, Strefler J, et al. Fossil-fueled

558   development (SSP5): An energy and resource intensive scenario for the 21st century. Global

559   Environmental Change. 2017 Jan 1;42:297–315.

560   45.   Matthew P. Lucas, Ragone D. Will Breadfruit Solve the World Hunger Crisis? ArcNews [Internet].

561   2012 [cited 2021 Jun 7]; Available from:

562   https://www.esri.com/news/arcnews/summer12articles/will-breadfruit-solve-the-world-hunger-

563   crisis.html

564   46.   Berg CC, Corner EJH, Jarrett FM. Moraceae genera other than Ficus. Flora Malesiana - Series 1,

565   Spermatophyta. 2006 Jan 1;17(1):1–146.

566   47.   Miguel E, Castiñeiras L, Knüpffer H, Hammer K. A checklist of the cultivated plants of Cuba. Die

567   Kulturpflanze. 1989 Jun 1;37(2):211–357.

568   48.   Fitzpatrick MC, Gotelli NJ, Ellison AM. MaxEnt versus MaxLike: empirical comparisons with ant

569   species distributions. Ecosphere. 2013;4(5):art55.

570   49.   Levis S, Badger A, Drewniak B, Nevison C, Ren X. CLMcrop yields and water requirements:

571   avoided impacts by choosing RCP 4.5 over 8.5. Climatic Change. 2018 Feb 1;146(3):501–15.

50.   Schlenker W, Roberts MJ. Nonlinear temperature effects indicate severe damages to U.S. crop

yields under climate change. PNAS. 2009 Sep 15;106(37):15594–8.

51.   Elevitch C, Ragone D. Breadfruit Agroforestry Guide: Planning and implementation of regenerative

organic methods. 2018.

52.   Hall C, Dawson TP, Macdiarmid JI, Matthews RB, Smith P. The impact of population growth and

climate change on food security in Africa: looking ahead to 2050. International Journal of

Agricultural Sustainability. 2017 Mar 4;15(2):124–35.

53.   Hunter D, Borelli T, Beltrame DMO, Oliveira CNS, Coradin L, Wasike VW, et al. The potential of

neglected and underutilized species for improving diets and nutrition. Planta. 2019 Sep

1;250(3):709–29.

54.   Gbemavo DSJC, Assani Z, Idohou R, Laly J, Gbaguidi A, Dansi A. Modelling current agro-

ecological zones for the cultivation of Dioscorea dumetorum (Kunth) Pax, a neglected and

underutilized yam species in Benin (West Africa). Model Earth Syst Environ. 2021 Sep

1;7(3):2127–35.

55.   Ratnayake SS, Kumar L, Kariyawasam CS. Neglected and Underutilized Fruit Species in Sri Lanka:

Prioritisation and Understanding the Potential Distribution under Climate Change. Agronomy. 2020

Jan;10(1):34.

    Figures and Tables

    Figure 1. Range of current (1970-2000) breadfruit suitability in the global tropics and subtropics (45° N to 45° S)

    as predicted from bioclimactic variables from 10 arcminute resolution WorldClim v2.1 data and filtered GBIF

    occurrence points. Climatic suitability of breadfruit cultivation is indicated by green shading ranging from 0 (fair) to

    1000 (good). (*A*) The ensemble mean-weighted SDM and 6 different individual SDMs including (*B*) MAXENT, (*C*)

    GLM, (*D*) GAM, (*E*) MARS, (*F*) RF, and (*G*) GBM. Black dots in (*A*) are observed occurrence points from GBIF.

    Pie charts in upper left of each plot indicate fraction of Fair (score=250-750) to Good (score=750+) area.

Figure 2. Future (2061-2080) breadfruit suitability range and change under stabilization and high-emission

scenarios. (a) Future suitability (Fair: score=250-750; Good: score=750+) of weighted SDM ensemble under

stabilization scenario (SSP2-4.5) and (b) high-emission scenario (SSP8-8.5) using averages from a CMIP6 ensemble

of opportunity from 8 GCMs. Change in suitable range (score>250) from baseline suitability score under (c)

stabilization scenario and (d) high-emission scenario.

Figure 3. Change in suitable breadfruit area (value>250) between the ensemble baseline SDM (B) and the stabilization (S) and high-emission (H) scenarios within various regions in the global tropics and subtropics, including Asia and the Pacific Islands, Africa, and Latin America, with Weak (1-3 members), Strong (4-5 members) and Very Strong (6+ members) CMIP6 GCM average agreement.

Figure 4. Difference Plots of Top BIOCLIM Contributors to SDM. Columns show the baseline (1970-2000)

bioclimatic variables (left) and the absolute difference of the specific bioclimatic variables from 2061-2080 under

SSP2-4.5 (middle) and SSP5-8.5 (right) scenarios of the CMIP6 ensemble of opportunity downscaled to 10

arcminute resolution from WorldClim v2.1.

Supplementary Figure 1. Threshold metrics for breadfruit suitability used for SDM classification. Categories

are demarcated in relation to trends between cumulative percent of total number of points (presence and

pseudoabsence) and model suitability value (0-1000). The Fair (250-750) and Good (750-1000) suitability

categories are marked by greater prevalence of presence points, with 4 occurrence points categorized as unsuitable.

Supplementary Table 1. Relative importance (TSS) of 7 selected BIOCLIM variables within and across 6

individual SDMs in baseline (1970-2000) scenario.

	Bioclimactic Variable	MAXENT	GLM	GAM	MARS	RF	GBM	Average
BIO2	Mean Diurnal Range	0.201	0.046	0.123	0.081	0.063	0.042	0.093
BIO7	Temperature Annual Range (BIO5-BIO6)	0.312	0.543	0.403	0.381	0.312	0.454	0.401
BIO11	Mean Temperature of Coldest Quarter	0.194	0.052	0.191	0.124	0.098	0.033	0.115
BIO12	Annual Precipitation	0.232	0.115	0.216	0.130	0.057	0.029	0.130
BIO17	Precipitation of Driest Quarter	0.121	0.027	0.106	0.055	0.031	0.025	0.061
BIO18	Precipitation of Warmest Quarter	0.150	0.063	0.107	0.026	0.038	0.016	0.067
BIO19	Precipitation of Coldest Quarter	0.185	0.012	0.104	0.069	0.034	0.016	0.070

628 **Supplementary Table 2.** SDM Variability of Breadfruit Suitability across 6 individual and ensemble SDMs.

629 Suitability is categorized over 98280727 km^2 of total land area constrained within the latitudes of 45° N to 45° S.

Suitability	SDM	Area (km^2)
Fair	MAXENT	17743876
	GLM	14117175
	GAM	15835734
	MARS	12581785
	RF	8411102
	GBM	9125204
	Ensemble	12725593
Good	MAXENT	9953293
	GLM	14455265
	GAM	12461001
	MARS	13168259
	RF	16982621
	GBM	16180823
	Ensemble	13721639
Unsuitable	MAXENT	70583558
	GLM	69708287
	GAM	69983992
	MARS	72530682
	RF	72887004
	GBM	72974700
	Ensemble	71833495

 Supplementary Table 3. GCM Variability in Breadfruit Suitability Under Stabilization and High Emissions

 Scenarios of individual CMIP6 GCMs. All available downscaled GCMs at 10 arcminute resolution were accessed

 from WorldClim v2.1. Breadfruit suitability is evaluated within land area constrained within the global tropics and

 subtropics.

Suitability	GCM	Area - SSP2-4.5	Area - SSP5-8.5
Fair	BCC-CSM2-MR	16127181	17010437
	CanESM5	16386867	17383498
	CNRM-CM6-1	15938225	17329833
	CNRM-ESM2-1	16025173	17377793
	IPSL-CM6A-LR	15660300	17401963
	MIROC-ES2L	14538802	15165247
	MIROC6	14200092	15245630
	MRI-ESM2-0	13900888	15981020
Good	BCC-CSM2-MR	9364025	8533701
	CanESM5	8101903	7002352
	CNRM-CM6-1	9571089	8025798
	CNRM-ESM2-1	9442967	7896402
	IPSL-CM6A-LR	10249666	8673592
	MIROC-ES2L	10186645	9507514
	MIROC6	10498834	8805485
	MRI-ESM2-0	10430547	8993401
Unsuitable	BCC-CSM2-MR	72784947	72732016
	CanESM5	73778272	73881192
	CNRM-CM6-1	72765857	72919540
	CNRM-ESM2-1	72807031	73000976

	IPSL-CM6A-LR	72358041	72192452
	MIROC-ES2L	73541595	73594281
	MIROC6	73576245	74224056
	MRI-ESM2-0	73944719	73301732

636 **Supplementary Figure 2. Suitable Breadfruit Range** (a) baseline (1970-2000), (b) Future Stabilization (2061-

637 2080), and (c) Future High Emissions (2061-2080) scenarios with country outlines using average values from a

638 weighted SDM ensemble projected using a CMIP6 GCM ensemble of opportunity under SSPs 2 and 5.