Potential of breadfruit cultivation to contribute to climate-resilient low latitude food systems

Lucy Yang[1], Nyree Zerega[2,3], and Daniel E. Horton[4]

[1]Program in Environmental Sciences, Northwestern University

[2]Program in Plant Biology and Conservation, Northwestern University

[3]Chicago Botanic Garden

[4]Department of Earth & Planetary Sciences and Institute for Sustainability and Energy at

Northwestern University

Keywords

food security, climate change adaptation, bioclimactic niches, neglected and underutilized species (NUS),

CMIP6, species distribution modeling, breadfruit

Abstract

The number of people in food crisis around the world is increasing, exacerbated by the challenges of

COVID-19 and a rapidly changing climate. Major crop yields are projected to decrease in low-latitude

regions due to anthropogenic climate change, making tropical and sub-tropical food systems particularly

vulnerable to climate shocks. Increased cultivation of breadfruit (*Artocarpus altilis*), often categorized as

a neglected and underutilized species (NUS), has been suggested as an agricultural adaptation pathway

for food insecure tropical and subtropical regions, due to its potential to enhance climate resilience and

overall sustainability of low-latitude agricultural systems. To better understand breadfruit's cultivation

suitability and geographic range in current and future climates, we employ a diverse set of observations

and models to delineate the current climatically viable breadfruit range, and assess the climatically viable

breadfruit range in the future (2061-2080) under stabilization and high emission scenarios. We find that

the area of suitable breadfruit range within the tropics and subtropics is projected to decrease ~4.4% in the

stabilization scenario and ~4.5% in the high emission scenario. In Southeast Asia and the Pacific Islands,

yield quality and consistency show minimal decreases under the high emission scenario, with increases in

total suitable area under both scenarios. In contrast, in Latin America and the Caribbean, the current range

of breadfruit suitability is projected to contract ~10.1-11.5% (stabilization-high emission). Present and

future model suitability outputs suggest that opportunities to successfully expand breadfruit cultivation

over the next several decades exist in sub-Saharan Africa, where food insecurity is coincidentally high.

However, in all regions, high emission scenario conditions reduce the overall consistency and quality of

breadfruit yields compared to the stabilization scenario. Our results have the potential to inform global

food security adaptation planning and highlight breadfruit as an ideal NUS to incorporate in food security

adaptation strategies in a changing climate.

1 Introduction

The number of people in the world lacking regular access to nutritious and sufficient food has been

on the rise since 2015, well before the COVID-19 pandemic exacerbated existing global food security

challenges and increased the prevalence of undernourishment (PoU) +1.5% to ~9.9% in 2020 (Diffenbaugh

et al., 2020; FAO, IFAD, UNICEF, WFP, and WHO, 2021). Food insecurity has increased in all global

regions since 2014, except in North America and Europe, and the world is considerably off track to achieve

the UN's global goal of Zero Hunger by 2030 (Global Network Against Food Crises & FSIN, 2020).

Climate change is a major driver of the observed rise in global hunger (FAO et al., 2018), with both gradual

climatic change and punctuated shocks working to multiply threats and disrupt the food system at various

levels, most notably food production and availability, but also in terms of access, quality, utilization, and

stability (Vermeulen et al., 2012). Climate adaptation solutions are needed to lessen risk in the global

agricultural production system, particularly in regions with high populations of undernourished peoples

(Mbow et al., 2019).

In the face of changing climate conditions and increases in the frequency and intensity of extremes,

scaling up the production of NUS is an attractive strategy to build global food system resilience, particularly

in the tropics and sub-tropics. Studies suggest that the agricultural adaptations made between 1981-2010

were insufficient to offset the negative impacts of climate change on common staple yields and production

losses (e.g., wheat, rice, soybean, and maize), particularly at lower latitudes (Iizumi et al., 2018). Climate

change over the 21st century is projected to have further significant negative impacts on the production of

common staples at lower latitudes (Brown & Funk, 2008; Lobell et al., 2008; Zhao et al., 2017). While the

geography of these major crops may expand to higher-latitudes under various scenarios (Diffenbaugh et al.,

2012), low latitude crop production is projected to be consistently and negatively impacted (Porter et al.,

2014). Reduced local agricultural production of major crops is projected to decrease food availability and

dietary diversity in tropical and subtropical regions (WFP, 2014). Furthermore, many of the world's current

food security challenges lie within tropical latitudes, where anthropogenic warming is also projected to

impact agricultural production most substantially (Mbow et al., 2019, p. 5; WFP, 2014). These regions are

particularly sensitive to increases in temperature due to low historical variability, with the time of

emergence (TOE) of unprecedented temperature and precipitation regimes arriving earlier in terrestrial

tropical biomes, compared to higher latitude regions, and potentially crossing an emergence signal-to-noise

threshold of 2 as early as the 2020s (Diffenbaugh & Scherer, 2011; Hawkins & Sutton, 2012; Tan et al.,

2018). Tropical agricultural production has already been adversely affected by extreme weather events and

increased climatic variability, and yields will likely worsen with increased temperature stress and rainfall

variability (Ericksen et al., 2011). Given these realized and projected impacts on staple crops, scaling up

production of NUS has been suggested as an important adaptation measure in food-insecure regions.

Expansion of NUS agriculture has the potential to enhance the climate resilience of food systems while

simultaneously improving other negative environmental and social characteristics of current global food

systems, including improving global crop and varietal diversity, increasing water efficiency, reducing

excessive use of agrochemicals, increasing more locally produced food, and implementing practices that

protect and conserve traditional knowledge and biodiversity (Mbow et al., 2019).

A traditional crop originating in Oceania, where it is an important and widely used staple, breadfruit
is a tropical NUS in other regions where it has a promising role in alleviating present and future food
insecurity and improving the overall sustainability and resiliency of tropical and sub-tropical food systems
(Mausio et al., 2020; Turi et al., 2015). Breadfruit offers practical advantages in low-latitude regions as a
nutritious "superfood" grown from a long-lived tree that does not need annual replanting, requires lower
energy input to cultivate (especially when cultivated in sustainable agroforestry systems), sequesters carbon
as a long-lived tree, and can be used as a staple food in a variety of dishes and cultural cuisines (Orwa C et
al., 2010). Breadfruit compares favorably, in some cases surpassing, major staples (including wheat, rice,
and corn) in both yield and nutritional content (Jones et al., 2011). Breadfruit also shows great potential to
reduce economic barriers to food access when implemented as part of economically sustainable
development strategies in food-insecure regions of the world (Omobuwajo, 2007). As such, breadfruit has
been presented as a potential solution to the world hunger crisis and labeled a "superfood of the future" by
popular media, although its potential implementation as a practical adaptation to food insecurity under a
changing climate has not been thoroughly explored. That said, its potential for more intensive cultivation
over its broad current range and expansion to new regions shows promise as a climate adaptation strategy
for sustainable food systems. Because the most commonly grown breadfruit varieties are seedless and are
clonally propagated via root cuttings, grafting, air layering, or micropropagation methods, they have little
to no likelihood of becoming invasive when introduced to new regions. In fact, over the last more than ~200
years seedless varieties have been introduced to many tropical regions with no recorded invasive behavior.

Breadfruit is a domesticated crop with a rich global history, connecting various cultures and regions.
Its wild, seeded ancestor is believed to have originated in New Guinea, and the seedless breadfruit was
domesticated in the islands of Oceania, where hundreds of cultivars exist (Zerega et al., 2015). Breadfruit
played a major role as a key staple food in the human colonization of these islands. By about 3,000 years
ago humans had reached Samoa and few-seeded and seedless varieties began to appear. When Hawaii was
one of the last Pacific islands to be colonized by humans (between 1400 and 1700 years ago), seedless

breadfruit was brought by Polynesian voyagers to islands, where it is known as "ulu" (Kirch, 2017).

Breadfruit continues to play a large part of cultural and spiritual life throughout Oceania. Europeans

encountered breadfruit in Oceania in the 1500s (Markham, 1904). By the 18th century it had been introduced

to Central and South America, the Caribbean, tropical Asia, and other tropical regions, and it remains

especially important in the Caribbean today (Ragone, 1997). Given the success of breadfruit cultivation in

various tropical and subtropical regions around the world, farmers in sub-Saharan Africa are increasingly

looking to breadfruit as a potential crop for alleviating local food insecurity concerns (Trees That Feed

Foundation, n.d.). Despite its large promise, breadfruit remains more of a subsistence crop than commercial

commodity in most regions (e.g., parts of West Africa, Southeast Asia, Latin America, and India) with

major production areas limited to the Pacific and Caribbean Islands (Ragone, 2018). However, interest in

commercial uses is growing (Peters, 2021).

The potential expansion of breadfruit cultivation as an adaptation strategy is constrained by present

and future climatic uncertainties. A newly planted breadfruit tree takes several years before it begins fruiting

and may live and produce fruit for over 50 years—if grown within a suitable environment. While global

increases of temperature in some regions may intuitively seem to allow for the expansion of breadfruit into

areas of higher latitudes, uncertainties around extreme weather, increased variability, and rising

temperatures within the current cultivated extent may concurrently limit its future suitability in its current

range. However, breadfruit yields are still expected to be more resilient than important global staples like

rice to certain climate change stressors. For example, after a breadfruit tree is established, it may withstand

droughts of 3-4 months (Taylor et al., 2016). The extent of breadfruit's resilience must be evaluated amidst

an uncertain climactic future.

To examine the adaptation potential of breadfruit cultivation, we evaluate current and projected

breadfruit niche suitability and assess the potential for range expansion and/or contraction using species

distribution modeling (SDM) and bioclimactic data. Given our knowledge of suitable climate conditions

for breadfruit cultivation, and our understanding that many of these underlying factors are undergoing

anthropogenically-driven changes, we use a weighted ensemble of six species distribution models (SDMs)

created using 7 selected bioclimactic variables (from BIOCLIM) to assess changes in future (2061-2080)

breadfruit suitability range using downscaled simulation data from 8 different General Circulation Models

(GCMs) from phase 6 of the Coupled Model Intercomparison Project (CMIP6) (Eyring et al., 2016). SDMs

that rely on bioclimactic variables over the full extent of a species range facilitate a realization of the

fundamental niche rather than a realized niche. Consideration of seasonality and variability is useful in

determining the fundamental niche of plant species, particularly when an SDM is applied to future climatic

conditions. This approach allows us to assess the potential range of breadfruit in our current climate where

it has not yet been introduced, as well as extrapolate breadfruit cultivation suitability in future climates.

This ensemble SDM approach in conjunction with ensemble GCM projections allows us to evaluate the

inherent inter-model differences/uncertainties in both steps of the process. Gaining an understanding of the

SDM and climatic uncertainties that constrain breadfruit's potential global range can facilitate policy design

intended to mobilize new adaptation strategies for more resilient low-latitude food systems.

2 Materials and Methods

2.1 Species Distribution Modeling (SDM)

To determine current and future breadfruit niche suitability, we used observations and

bioclimactic data to constrain 6 unique SDMs: Generalized Linear Model (GLM), Generalized Additive

Model (GAM), Multiple Adaptive Regression Splines (MARS), Random Forest (RF), Generalized

Boosted Model (GBM), and Maximum Entropy (MAXENT) using the *Biomodhub/Biomod2* software

version 3.4.13 (2018/2021). A weighted-mean ensemble of SDMs was used to determine current

breadfruit niche suitability. The ensemble approach to species distribution modeling has gained

considerable popularity and is useful for examining which bioclimatic variables determine the

fundamental niche of breadfruit, allowing us to project this model to future scenarios and evaluate the

changed distribution of global breadfruit suitability (Hao et al., 2019).

152   To constrain the current distribution of breadfruit and provide training data for our SDMs, we use

153 Global Biodiversity Information Facility (GBIF) historical breadfruit (*Artocarpus altilis*) occurrence point

154 data. GBIF presence points are prone to sampling bias because it allows citizen science contributions to

155 the database. Citizen scientists are prone to misidentify *A. altilis*, particularly by confusing the plant with

156 its close relative, seeded breadnut (*A. camansi*). To limit this bias, we downloaded only those data points

157 that were identified as part of herbarium specimens ("Preserved specimen") or that included picture

158 documentation. We also filtered GBIF presence points using a number of screens. First, presence data that

159 appeared under alternative names (taxonomic synonyms), e.g., *A. communis, A. incisa,* and *A. incisus*,

160 were included. However, data of two close relatives (i.e., *A. camansi* and *A. mariannensis*) were not

161 included (Zerega et al., 2006). Next, we ran the GBIF points using default settings of the

162 CoordinateCleaner, an R-package tailored for cleaning occurrence records from biological collection

163 databases (Zizka et al., 2020). This package flags geographical coordinates that are overly imprecise or

164 have common problems (e.g., points assigned to country centroids due to automated geo-referencing,

165 plants belonging to botanical gardens or museums, etc.). We also added datapoints from a yield study in

166 Jamaica (Trees that Feed Foundation, 2020, unpublished data). Finally, we discarded duplicate points

167 within common grid cells to limit spatial bias. Ultimately, over half of the breadfruit presence points

168 downloaded from GBIF were removed. The remaining 431 presence points are used for the SDM

169 ensemble training and prediction.

170   To determine the climatic variables that best predict the occurrence of current breadfruit

171 distribution, we used the dataset of bioclimatic variables (BIOCLIM) for 1970-2000 from WorldClim

172 Version 2.1, a database of high-resolution global climate data (Fick & Hijmans, 2017). To prevent

173 overfitting of the final SDM, we examined the full BIOCLIM dataset of 19 variables and selected a subset

174 of variables based on their predictive performance in individual and paired runs. We selected variables

175 largely following the methodology detailed in Moat et al., 2017 using GLM and GAM, two of the six

176 SDMs used for the final ensemble. Boxplots, preliminary map projections, and evaluative metrics

including sensitivity, specificity, True Skills Statistics (TSS), relative operating characteristic curve

(ROC), and Cohen's Kappa statistic (KAPPA) were used to determine which combination of variables

were best for model fit, i.e., which combination of bioclimatic variables best predicted the observed range

of breadfruit. After evaluating variable performance, we tested resulting variable combinations for

collinearity and multi-collinearity using a threshold of $R^2=0.8$ and reduced the selection to 7 variables

(Table 1). Annual mean temperature (BIO7) and annual precipitation (BIO12) were among the highest

performing variables in the preliminary model runs using GLM and GAM. While environmental

characteristics such as solar radiation and soil quality also impact breadfruit growth, they are not used to

constrain breadfruit suitability in this study.

The optimal combination of variables to characterize niche suitability differs based on the spatial

domain. A preliminary version of the model covered the full extent of global BIOCLIM data at 10

arcminute resolution (~100 km). However, because over 97% of our preliminary weighted model results

were constrained within the latitudes of 45° N to 45° S, the final analysis presented in this study was

performed within these lesser latitudinal bounds. Despite this change, our analysis covers almost the

entire extent of the current breadfruit range, which is ideal for accurately capturing the fundamental niche

of the species (Guisan et al., 2017). A lesser geographical extent also increases the reliability of

pseudoabsence data. We use a presence-only SDM that relies on pseudoabsence (background) points to

model suitability because no reliable source of true absence data was available. The TSS scores of the

three sets of pseudoabsence points we generated vary by 0.05. Although we latitudinally limited the

geographic extent of the SDM in favor of generating more accurate pseudoabsence points, it is still

difficult to produce useful pseudoabsence data at this large extent, as evidenced by the concentration of

pseudoabsence points in grid-cells of less-suitable categories (Supplementary Figure 1).

We ran the 7 selected BIOCLIM variables and 431 filtered data points using the Biomod2 R

package version 3.4.14 using six independent SDM methods (GLM, GAM, MARS, RF, GBM, and

MAXENT.Phillips). We followed established protocols for model evaluation, execution, replication, and

generation of three sets of pseudo-absence data (*Habitat Suitability and Distribution Models Applications*

r | Ecology and Conservation | Cambridge University Press, n.d.). Presence and pseudoabsence data were

split randomly with a ratio of 70:30 to build training and testing data separately over ten runs. The same

metrics of sensitivity, specificity, TSS, ROC, and KAPPA were used to evaluate model outputs.

Resultant maps and statistics from model outputs were used to evaluate the performance of the

SDMs. Output statistics were reviewed to ensure a lack of difference between model runs, which could

indicate problematic outcomes, such as overfitting. After evaluation, the final SDM configuration was

used to produce our suitability predictions from ten runs and three replicates. The TSS threshold of 0.7

was used to remove poorly performing runs from all five models before producing a mean ensemble

prediction weighted by TSS score. Our weighted ensemble of SDMs demonstrates robust present-range

predictive power, characterizing the distribution of breadfruit presences indicated by GBIF data while

also predicting broader distribution within Africa and other previously unexamined regions with high

sensitivity (0.96) and reasonable accuracy (specificity = 0.83). The included SDMs in this study meet

established criteria (TSS = 0.79, KAPPA = 0.81, ROC = 0.93).

217   2.2 General Circulation Model (GCM)

Projected climate data was then used to determine niche expansion/contraction under future

climate scenarios SSPs 2 and 5 (Fricko et al., 2017; Kriegler et al., 2017). We modeled the future

projections over the time period of 2061-2080 using the Biomod2 package Ensemble Forecasting. The

projection was performed with the same seven selected BIOCLIM variables from the GCM data under

two shared socioeconomic pathway scenarios: SSP2-4.5 (stabilization GHG emission scenario) and

SSP5-8.5 (high GHG emission scenario) forcing scenarios from 8 different GCMs. The GCMs from

CMIP6 were used based on availability in WorldClim 2.1 and include BCC-CSM2-MR, CNRM-CM6-1,

CNRM-ESM2-1, CanESM5, IPSL-CM6A-LR, MIROC-ES2L, MIROC6, and MRI-ESM2-0.

2.3 Analysis and Mapping

The SDM outputs from the current (Figure 1) and the two future scenarios were processed and

categorized into three: Good, Fair, and Unsuitable. Niche classes were chosen based on natural inflection

points and breaks between pseudoabsence and presence data (see Supplementary Figure 1). The division

between "Fair" and "Good" categories was placed at 50% to approximately reflect the inflection point of

the data. The Good niche classifications assesses breadfruit suitability based on yield quality and

consistency, marking locations that are conducive to the establishment of new breadfruit trees.

Meanwhile, the Fair range is more liberal and represents climactic conditions that established trees will

likely withstand, ergo persistence is assumed likely.

3 Results

3.1 SDM Prediction of Current Breadfruit Suitability Range

Our weighted ensemble baseline SDM predicts niche suitability in all GBIF-identified locations of

current breadfruit cultivation, as well as substantial swathes of the global tropics and subtropics where

successful cultivation of breadfruit has not yet been confirmed. Predicted suitable areas for cultivation

include large sections of Southeast Asia, Central America, northern South America, the Caribbean, and

equatorial Africa (Figure 1a). Latitudinal limits of modeled suitability extend as far north as northeastern

India in Asia and southern Florida in North America and stretch as far south as Chile in South America and

New Zealand in the Pacific. These regions have previously been identified by the National Tropical Botanic

Garden (NTBG) as suitable for breadfruit cultivation based on current climatic conditions (Matthew P.

Lucas & Ragone, 2012). Suitable niches that extend into the higher latitudes of the tropics are largely

concentrated along coastlines, in agreement with available GBIF occurrence data (Figure 1a; black dots).

Notable areas without widespread occurrence data that are identified as having robust potential for

breadfruit cultivation include Cuba, equatorial Africa, the southeastern coastline of Africa, and the Malay

peninsula. In Africa, the lack of occurrence data in GBIF may be explained by breadfruit's recent arrival to

the continent, but recent successful introductions haven been made in Uganda, Kenya, Tanzania, and Ghana.

While no GBIF records are recorded in Cuba, it is known to grow there (Berg et al., 2006; Miguel et al.,

1989; Trees That Feed Foundation, n.d.).

Our SDM breadfruit suitability predictions employ seven weighted bioclimatic variables to

characterize the climatic conditions within a grid cell. The environmental variables were selected based on

evaluation scores on the Generalized Linear Models and Generalized Additive Models while controlling

for multicollinearity and overfitting (see Methods). Contribution weights of the seven bioclimatic variables

are quantified using True Skill Statistic (TSS) scores. We find that baseline breadfruit suitability is largely

driven by three temperature-related bioclimatic variables (TSS: 60.9). The annual temperature range,

defined as the difference between the maximum temperature of the warmest month and the minimum

temperature of the coldest month is the predominant prediction contributor (Figure 4c; TSS: 40.1).

Secondary contributors include the mean diurnal temperature range (Figure 4a; TSS: 9.2) and the mean

temperature of the coldest month (Figure 4g; TSS: 11.5). The four remaining contributing variables – annual

precipitation, and precipitation during the driest, warmest, and coldest quarters – are lesser contributors

relative to the temperature variables (Figure d-g; TSS: 6.1-7.0).

To refine our suitability findings and identify the niches most suitable for cultivation, we employ

statistical thresholding to subdivide the SDM suitability scores into Unsuitable, Fair, and Good categories

(Supplementary Figure 1). The unsuitable niche classification marks areas where breadfruit cultivation is

barely possible to impossible, whereas fair classification marks areas where breadfruit cultivation is

possible but yield and quality may be negatively affected by adverse seasonal weather conditions (e.g., low

rainfall or high temperatures). Good niche classification indicates where breadfruit cultivation has the

potential for consistent, high quality yields that are less likely to be affected by adverse weather conditions.

For the lower threshold, between unsuitable and fair, we set the cut-off at the inflection point where an

increase in model suitability values and the cumulative percentage of all (including pseudoabsence and

occurrence) points plateaus, which resulted in the loss of four ground control points. The division between

Fair and Good categories indicates a second inflection point where model suitability values begin to

substantially increase.

Individual SDMs are remarkably consistent in their suitable geographic range predictions (Figure

1b-g) but show notable differences in categorizing the level of breadfruit suitability within these ranges.

Across the 6 SDMs, there is only a 3.3% difference in the total "suitable" land area, although the

classification of suitable land into either Fair or Good is less consistent, with ~33.1-64.0% of the total

suitable area identified as Fair depending on SDM and ~36.0-66.9% classified as Good. When considering

the total modeled area of suitability, both MAXENT (35.9% ,10.0M km^2 Good; 64.1%, 17.7M km^2 Fair)

and GAM (44.0%, 12.5M km^2 Good; 56.0%, 15.8M km^2 Fair) predict lesser "Good" areas in proportion to

"Fair", while RF (66.9%, 8.4M km^2 Good; 33.1%, 17.0M km^2 Fair) and GBM (63.9%, 16.2M km^2 Good;

36.1%, 9.1M km^2 Fair) predict greater "Good" suitability (Figure 1). Because MAXENT is a presence-only

SDM, it is known to underestimate the probability of occurrence within areas of observed presence, while

overestimating in areas without known presence (Fitzpatrick et al., 2013). Several MAXENT runs (6 out

of 30) were removed from the final weighted SDM ensemble due to low TSS scores, likely due to a lack of

convergence. Marginal areas of Fair suitability are consistently identified by all 6 SDMs in the same

locations (e.g., New Zealand, southwestern coast of Africa, Chile). Southeast Asia, Oceania, and the Pacific

Islands are also consistently valued with the highest proportion of Good to Fair suitability. The 6 SDMs

demonstrate consistent suitability coverage and patterns in regional Good to Fair suitability, although

specific values differ.

Using the weighted-ensemble projection, we focus our narrative on comparative breadfruit

suitability in three main regions: Asia and the Pacific Islands, the Caribbean and Latin America, and Africa.

Compared to the averaged global proportion of area with Good (51.9%, ~14M km^2) to Fair (48.1%, ~12M

km^2) suitability (Figure 3), breadfruit is particularly suited to climactic conditions in Southeast Asia.

Breadfruit is suitable for cultivation within 80.7% of the area (~3.6M km^2) in this region with a high

proportion of Good (77.4%, ~2.8M km^2) to Fair (22.6%, ~800k km^2) quality, although Pacific Islands such

as Hawaii also show similarly high levels of Good (87.7%, ~30k km^2) to Fair (12.3%, ~4k km^2) area . In

Latin America and the Caribbean, island countries show high coverage of suitable area (99.2%, ~2.3M km^2)

while 51.7% (~11.7M km^2) of the overall land area in Latin America and the Caribbean is projected to be

suitable. 41.3% of these suitable areas are classified as Good and 58.7% as Fair. In Africa, 20.8% of the

continent is projected to be suitable for growing breadfruit, with 45.4% of this land showing Good and 54.6%

showing Fair suitability. This baseline SDM prediction gives a comprehensive picture of the range and

quality of suitable breadfruit-growing areas in various low-latitude regions of the world.

Change in Breadfruit Range due to Climate Change

To assess the effects of climate change on breadfruit suitability, we project our SDM to future

(2061-1080) climate conditions using an ensemble of opportunity of 8 GCMs from CMIP6. While

breadfruit is projected to largely maintain its present range, the resilience and adaptation potential is

projected to decrease, with more substantial decreases in Good area projected from the high-emissions

scenario (44.4%) than the stabilization scenario (32.1%) (Figure 2). We find a 4.7% decrease in weighted

ensemble SDM suitable breadfruit range (~1.2M km^2) in the stabilization scenario and a 4.5% decrease

(~1.2M km^2) under high emission scenario conditions (Figure 3). However, the total area of "winners" and

"losers" in terms of overall breadfruit suitability changes largely balance on a global scale.

Change in suitable breadfruit range – expansion, contraction, and persistence— from the baseline

projection varies considerably across regions (Figure 2). The suitable area primarily persists and even

increases ~0.7% (~23k km^2) under the stabilization scenario and 0.8% (~30k km^2) under the high emission

scenario in Southeast Asia, although this increase in area is accompanied by a decrease in quality (2.6%

and 8.7%) and reduction in total Good area. In Asia and the Pacific Islands more broadly, total suitable

breadfruit area increases ~2.8% (~144k km^2) and ~4.1% (~214k km^2), showing areas of potential expansion

in the upper and lower latitudinal limits of this region (Figure 2C, D). The majority of suitable breadfruit-

growing area is projected to persist in equatorial Africa, although with lower resilience potential to future

climate conditions than in Southeast Asia (1.5-3.4% reduction in total suitable area and 58.2-70.0%

reduction in Good area depending on emissions scenario). In both the stabilization and high-emission

scenarios, the baseline suitability range within equatorial Africa expands northward while the southern

bounds of breadfruit suitability are projected to contract. However, Latin America and the Caribbean are

projected to experience the largest magnitude range reduction on both the stabilization (10.1%) and high

emission pathways (11.5%). These range reductions appear to be concentrated in specific areas, particularly

in the Yucatan peninsula and southwestern Brazil. While Caribbean islands experience minimal contraction

of the overall suitable breadfruit-growing area (1.4-1.6% reduction) under either emission pathway, the

quality of these conditions may decrease significantly (50.6-72.5% reduction in Good area). Despite

regional variation, a large portion of global suitable breadfruit range is projected to persist in both scenarios

(particularly in Africa and Asia and the Pacific Islands), suggesting that (1), established trees will likely

withstand future climatic conditions in most of the tropics and subtropics and (2) there is room for expanded

cultivation of breadfruit across a broad section of the global tropics and subtropics in Africa. However, a

greater decrease in quality but not necessarily a greater decrease in total area of suitability is projected under

the high emissions scenario in all regions.

Projected expansion of breadfruit suitability is driven by increases in seasonal temperature and

precipitation. Increased precipitation during the coldest quarter (BIO19, Fig. 4t, u) as well as general

increases in annual precipitation (BIO12, Fig. 4k,l) drive potential expansion of suitable breadfruit area at

the perimeter of the baseline suitable range. Meanwhile, decreases in precipitation of the warmest quarter

(BIO18) drive contraction of suitable breadfruit range in Central and South America (Fig. 4q, r). Mean

Diurnal Range shows little change in Southeast Asia but increases in areas of expansion and decreases in

areas of contraction and reduced breadfruit suitability (Fig. 4b, c).

To characterize the structural uncertainty of our breadfruit projections, we use our CMIP6 ensemble

of opportunity and define projection confidence according to ensemble member agreement (i.e., 4-5

members agree – Strong, 6+ members agree = Very Strong). Projections of future suitability show Very

Strong GCM agreement over the majority of the persistent range of breadfruit suitability under the

stabilization scenario (92.8%) and high-emission scenario (90.5%), with a larger area of weak GCM

agreement under the high-emission scenario. Weak SDM agreement (<4 members agree) only characters

5.1% of the persisting suitable range under the high-emission scenario and 3.6% of the suitable range under

the stabilization scenario. Strong and Weak agreement is concentrated at the outer fringes of breadfruit

distribution and most significantly in central South America in the Mato Grosso Plateau and eastern Africa

in the Ethiopian highlands (Figure 3). This finding suggests higher levels of uncertainty in areas of higher

elevation, although breadfruit suitability is projected to persist in the Andes in South America. Areas of

potential expansion also overlap with the areas of greatest uncertainty. In future scenarios, South Asia and

South America exhibit the greatest variability in area among GCMs (Figure 4). However, the majority of

future global breadfruit cultivation in both scenarios demonstrates strong (4-5) to Very Strong (6+)

agreement among the 8 GCMs in both scenarios.

Discussion

Our SDM prediction captures known locations of breadfruit while also suggesting potential

expansion for the crop into food-insecure regions, most notably within Sub-Saharan Africa. While previous

studies have attempted to constrain global and regional distributions of breadfruit suitability (Matthew P.

Lucas & Ragone, 2012; Mausio et al., 2020), direct comparisons are challenging due to differences in

scoring systems. However, the geographic range of predictions from other studies are in good agreement

with our ensemble SDM predictions. For example, Lucas and Ragone (2012) used Geographic Information

Systems (GIS) to produce a map of global breadfruit suitability using WorldClim v1.4 data and ecologically

determined ranges for mean temperature and rainfall (Ragone, 2006). While this study did not assess the

impacts of climate change, their results on the current distribution of suitable cultivation range is in good

agreement with our SDM-based findings. Mausio et al. (2020) further used fuzzy-set models to extrapolate

breadfruit suitability from environmental parameters derived in Hawaii. Their work offers a useful

comparison of an alternative approach to modeling breadfruit suitability by incorporating FAO EcoCrop

parameters and climate data from CMIP5, but it does not address the impact of seasonal temperature and

precipitation data on potential breadfruit distribution under future scenarios.

Our SDM indicates that the differences between the high-emissions scenario and stabilization

scenario in terms of total breadfruit suitability area is minimal, however the high-emission scenario is

predicted to lead to more substantial reductions in suitability scores in most regions. These findings are

particularly important for tropical and subtropical regions (low to mid latitude regions) where impacts on

the production of conventional staple crops such as maize, wheat, soybean and rice are projected to be more

negative than higher latitudes (Levis et al., 2018). Expanded cultivation of breadfruit in such regions may

contribute to global food and economic security, as a locally produced staple food that can lead to greater

independence from food imports.

Breadfruit shows the most promising resilience to future climactic conditions within its native

extent in Asia, largely persisting in Southeast Asia and even showing potential areas of expansion within

India and China. Culturally, breadfruit is also a significant traditional crop in many Pacific Islands including

Hawaii, where it has been noted for its promising role as part of local biocultural restoration efforts

(Langston and Lincoln, 2018). Breadfruit can play an important role in local low-latitude food systems,

alleviating the yield losses of major crops such as rice, while promoting sustainable agroforestry systems.

However, the persistence of suitable conditions for growing breadfruit are most at risk in specific regions

within Latin America, namely a section of the Yucatan peninsula and southwestern Brazil in the Mosso

Plateau. These areas experience decreases in annual precipitation, with significant decreases in precipitation

during the warmest quarter of the year. These areas also coincide with weak agreement among the 8 GCMs,

with less than half agreeing on the magnitude of precipitation reduction.

Our SDM predictions also signal opportunities to expand cultivation in Central Africa, building on

nascent cultivation of breadfruit which currently exists in parts of West and East Africa (Trees That Feed

Foundation, n.d.). Given the present state of food insecurity in Africa, the widespread cultivation of

breadfruit offers a path towards increased food security and independence in the present and more distant

future for several Sub-Saharan countries. As a promising NUS, breadfruit could fill immediate and future

needs to increase food security in many tropical regions, especially as crops such as maize and rice become

less suited to warming conditions (Schlenker & Roberts, 2009; Zhao et al., 2017). Breadfruit demonstrates

resilience to uncertain climactic conditions in many tropical food-insecure regions. This adaptation

potential is further enhanced if breadfruit trees are well-established and grown within sustainable

agroforestry systems. Agroforestry systems can serve as both adaptation and mitigation, contributing to

food productivity while enhancing restoration by combining traditional growing methods used by Pacific

Islanders for centuries with modern horticultural techniques (Elevitch & Ragone, 2018; Mbow et al., 2019).

Our study uses species distribution modeling to examine potential areas for breadfruit cultivation

over the next several decades under two emission pathways. Our model prediction is limited to bioclimactic

factors (seasonal temperature and precipitation variables) and does not include the potential influence of

soil type, solar radiation, or elevated CO_2 concentrations on breadfruit suitability. We are confident that the

45° N to 45° S swath covers the majority (~97%) of present and future suitable breadfruit range under

these scenarios, capturing the fundamental niche and identifying new areas where breadfruit can be

introduced or expanded globally. While additional presence observations may improve the performance of

our model, most studies agree that sample size effects usually become less critical above 50 presence points

(Guisan et al., 2017). Ideally, models of habitat suitability should also be based on measurements of

population fitness at each geographic location. However, this would prevent the use of reliable data

available in natural history collections. Although data points from GBIF were cleaned and filtered, absence

data and rigorous groundtruthing would further improve the performance of the individual models that

made up the final SDM ensemble. When projecting to future conditions, we were limited to using

realizations from relatively few (8) CMIP6 GCMs from WorldClim v.2.1 which likely limits the range of

internal variability and structural uncertainty considered in our projections.

Significant portions of sub-Saharan Africa show room for expansion of breadfruit cultivation over

the next several decades, especially as rapid population growth is projected to intensify food insecurity and

undernourishment across the continent (Hall et al., 2017). Closing the yield gap via sustainable

intensification of agriculture which utilizes NUS such as breadfruit may be critical in preventing future

catastrophic circumstances. Prioritizing a rich diversity of NUS over staple crops such as wheat, maize, and

rice will lead to healthier diets and improved nutrition and contribute to sustainable development (Hunter

et al., 2019). Similar studies have examined the potential distribution of NUS under climate change,

including 4 neglected and underutilized fruit species (NUFS) in Sri Lanka and a yam species in Benin

(Gbemavo et al., 2021; Ratnayake et al., 2020), but few have examined the viability of expanding

cultivation of NUS across different continents at a global scale. As a long-lived crop that can be grown and

used locally, breadfruit offers an opportunity for various low-latitude communities around the world to

improve the resilience of their local food systems. By building agroforestry systems around breadfruit, low-

latitude regions can avoid the pitfalls of strictly monoculture systems while building sustainable food

systems with increased resilience to climate change.

4.4 Conclusion

Our model of breadfruit niche suitability has the potential to provide projections useful for both

short-term decision making and long-term adaptation planning. Because breadfruit cultivation has not yet

expanded into its full suitable range, this crop has the potential to alleviate present and future food security

burdens, particularly in food-insecure countries in equatorial Africa. Although breadfruit growers may need

to contend with a certain level of reduced quality of yields in the future, suitable climatic conditions for

breadfruit cultivation exists across a broad range of the tropics and subtropics and persist for the next several

decades regardless of emissions pathway, indicating that it is a reliable and promising NUS to implement

as a climate adaptation strategy. Depending on scale of production, breadfruit itself may even play a role in

contributing to climate change mitigation due to its significant carbon sequestration benefits.

Acknowledgments

Research presented in this paper was supported by a Conference Travel Grant and summer Undergraduate

Research Grant from Northwestern University's Office of the Provost to LY. The authors declare that

there is no conflict of interest.

References

Berg, C. C., Corner, E. J. H., & Jarrett, F. M. (2006). Moraceae genera other than Ficus. *Flora Malesiana*

- Series 1, Spermatophyta, *17*(1), 1–146.

Biomodhub/biomod2. (2021). [R]. biomod2 package, tools and scripts.

https://github.com/biomodhub/biomod2 (Original work published 2018)

Brown, M. E., & Funk, C. C. (2008). Food Security Under Climate Change. *Science*, *319*(5863), 580–

581. https://doi.org/10.1126/science.1154102

Diffenbaugh, N. S., Field, C. B., Appel, E. A., Azevedo, I. L., Baldocchi, D. D., Burke, M., Burney, J. A.,

Ciais, P., Davis, S. J., Fiore, A. M., Fletcher, S. M., Hertel, T. W., Horton, D. E., Hsiang, S. M.,

Jackson, R. B., Jin, X., Levi, M., Lobell, D. B., McKinley, G. A., … Wong-Parodi, G. (2020).

The COVID-19 lockdowns: A window into the Earth System. *Nature Reviews Earth &*

Environment, *1*(9), 470–481. https://doi.org/10.1038/s43017-020-0079-1

Diffenbaugh, N. S., Hertel, T. W., Scherer, M., & Verma, M. (2012). Response of corn markets to climate

volatility under alternative energy futures. *Nature Climate Change*, *2*(7), 514–518.

https://doi.org/10.1038/nclimate1491

Diffenbaugh, N. S., & Scherer, M. (2011). Observational and model evidence of global emergence of

permanent, unprecedented heat in the 20th and 21st centuries. *Climatic Change*, *107*(3), 615–624.

https://doi.org/10.1007/s10584-011-0112-y

Elevitch, C., & Ragone, D. (2018). *Breadfruit Agroforestry Guide: Planning and implementation of*

regenerative organic methods.

Ericksen, P. J., Thornton, P. K., Notenbaert, A. M. O., Cramer, L., Jones, P. G., & Herrero, M. T. (2011).

Mapping hotspots of climate change and food insecurity in the global tropics [Report]. CGIAR

Research Program on Climate Change, Agriculture and Food Security.

https://cgspace.cgiar.org/handle/10568/3826

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016).

Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design

and organization. *Geoscientific Model Development*, *9*(5), 1937–1958.

https://doi.org/10.5194/gmd-9-1937-2016

FAO, IFAD, UNICEF, WFP, and WHO. (2021). *The State of Food Security and Nutrition in the World*

2021. FAO.

FAO, IFAD, UNICEF, WFP, & WHO. (2018). *The State of Food Security and Nutrition in the World:*

Building climate resilience for food security and nutrition. FAO.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for

global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

https://doi.org/10.1002/joc.5086

Fitzpatrick, M. C., Gotelli, N. J., & Ellison, A. M. (2013). MaxEnt versus MaxLike: Empirical

comparisons with ant species distributions. *Ecosphere*, *4*(5), art55. https://doi.org/10.1890/ES13-

00066.1

Fricko, O., Havlik, P., Rogelj, J., Klimont, Z., Gusti, M., Johnson, N., Kolp, P., Strubegger, M., Valin, H.,

Amann, M., Ermolieva, T., Forsell, N., Herrero, M., Heyes, C., Kindermann, G., Krey, V.,

McCollum, D. L., Obersteiner, M., Pachauri, S., … Riahi, K. (2017). The marker quantification

of the Shared Socioeconomic Pathway 2: A middle-of-the-road scenario for the 21st century.

Global Environmental Change, *42*, 251–267. https://doi.org/10.1016/j.gloenvcha.2016.06.004

Gbemavo, D. S. J. C., Assani, Z., Idohou, R., Laly, J., Gbaguidi, A., & Dansi, A. (2021). Modelling

current agro-ecological zones for the cultivation of Dioscorea dumetorum (Kunth) Pax, a

neglected and underutilized yam species in Benin (West Africa). *Modeling Earth Systems and*

Environment, *7*(3), 2127–2135. https://doi.org/10.1007/s40808-020-00973-w

Global Network Against Food Crises, & FSIN. (2020). *Global Report on Food Crises*.

https://www.wfp.org/publications/2020-global-report-food-crises

Guisan, A., Thuiller, W., & Zimmermann, N. E. (2017). *Habitat Suitability and Distribution Models:*

With Applications in R. Cambridge University Press. https://doi.org/10.1017/9781139028271

Habitat suitability and distribution models applications r | Ecology and conservation | Cambridge

University Press. (n.d.). Retrieved March 29, 2021, from

https://www.cambridge.org/us/academic/subjects/life-sciences/ecology-and-conservation/habitat-

suitability-and-distribution-models-applications-r?format=PB&isbn=9780521758369

Hall, C., Dawson, T. P., Macdiarmid, J. I., Matthews, R. B., & Smith, P. (2017). The impact of population

growth and climate change on food security in Africa: Looking ahead to 2050. *International*

Journal of Agricultural Sustainability, *15*(2), 124–135.

https://doi.org/10.1080/14735903.2017.1293929

Hao, T., Elith, J., Guillera-Arroita, G., & Lahoz-Monfort, J. J. (2019). A review of evidence about use and

performance of species distribution modelling ensembles like BIOMOD. *Diversity and*

Distributions, *25*(5), 839–852. https://doi.org/10.1111/ddi.12892

Hawkins, E., & Sutton, R. (2012). Time of emergence of climate signals: TIME OF EMERGENCE OF

CLIMATE SIGNALS. *Geophysical Research Letters*, *39*(1), n/a-n/a.

https://doi.org/10.1029/2011GL050087

Hunter, D., Borelli, T., Beltrame, D. M. O., Oliveira, C. N. S., Coradin, L., Wasike, V. W., Wasilwa, L.,

Mwai, J., Manjella, A., Samarasinghe, G. W. L., Madhujith, T., Nadeeshani, H. V. H., Tan, A.,

Ay, S. T., Güzelsoy, N., Lauridsen, N., Gee, E., & Tartanac, F. (2019). The potential of neglected

and underutilized species for improving diets and nutrition. *Planta*, *250*(3), 709–729.

https://doi.org/10.1007/s00425-019-03169-4

Iizumi, T., Shiogama, H., Imada, Y., Hanasaki, N., Takikawa, H., & Nishimori, M. (2018). Crop

production losses associated with anthropogenic climate change for 1981–2010 compared with

preindustrial levels. *International Journal of Climatology*, *38*(14), 5405–5417.

https://doi.org/10.1002/joc.5818

Jones, A. M. P., Ragone, D., Aiona, K., Lane, W. A., & Murch, S. J. (2011). Nutritional and

531   morphological diversity of breadfruit (Artocarpus, Moraceae): Identification of elite cultivars for

532   food security. *Journal of Food Composition and Analysis*, *24*(8), 1091–1102.

533   https://doi.org/10.1016/j.jfca.2011.04.002

534   Kirch, P. V. (2017). *On the Road of the Winds: An Archaeological History of the Pacific Islands before*

535   *European Contact, Revised and Expanded Edition* (2nd ed.). University of California Press.

536   https://www.jstor.org/stable/10.1525/j.ctv1xxsng

537   Kriegler, E., Bauer, N., Popp, A., Humpenöder, F., Leimbach, M., Strefler, J., Baumstark, L., Bodirsky,

538   B. L., Hilaire, J., Klein, D., Mouratiadou, I., Weindl, I., Bertram, C., Dietrich, J.-P., Luderer, G.,

539   Pehl, M., Pietzcker, R., Piontek, F., Lotze-Campen, H., … Edenhofer, O. (2017). Fossil-fueled

540   development (SSP5): An energy and resource intensive scenario for the 21st century. *Global*

541   *Environmental Change*, *42*, 297–315. https://doi.org/10.1016/j.gloenvcha.2016.05.015

542   Levis, S., Badger, A., Drewniak, B., Nevison, C., & Ren, X. (2018). CLMcrop yields and water

543   requirements: Avoided impacts by choosing RCP 4.5 over 8.5. *Climatic Change*, *146*(3), 501–

544   515. https://doi.org/10.1007/s10584-016-1654-9

545   Lobell, D. B., Burke, M. B., Tebaldi, C., Mastrandrea, M. D., Falcon, W. P., & Naylor, R. L. (2008).

546   Prioritizing Climate Change Adaptation Needs for Food Security in 2030. *Science*, *319*(5863),

547   607–610. https://doi.org/10.1126/science.1152339

548   Markham, C. (1904). The voyages of Pedro Fernandez de Quiros, 1595. *Scottish Geographical Magazine*,

549   *20*(12), 649–652. https://doi.org/10.1080/14702540408555042

550   Matthew P. Lucas, & Ragone, D. (2012). Will Breadfruit Solve the World Hunger Crisis? *ArcNews*.

551   https://www.esri.com/news/arcnews/summer12articles/will-breadfruit-solve-the-world-hunger-

552   crisis.html

553   Mausio, K., Miura, T., & Lincoln, N. K. (2020). Cultivation potential projections of breadfruit

554   (Artocarpus altilis) under climate change scenarios using an empirically validated suitability

model calibrated in Hawai'i. *PLOS ONE*, *15*(5), e0228552.

https://doi.org/10.1371/journal.pone.0228552

Mbow, C., Rosenzweig, C., & Barioni, L. G. (2019). Food Security. In *Climate Change and Land: An*

IPCC special report on climate change, desertification, land degradation, sustainable land

management, food security, and greenhouse gas fluxes in terrestrial ecosystems.

https://www.ipcc.ch/srccl/chapter/chapter-5/

Miguel, E., Castiñeiras, L., Knüpffer, H., & Hammer, K. (1989). A checklist of the cultivated plants of

Cuba. *Die Kulturpflanze*, *37*(2), 211–357. https://doi.org/10.1007/BF01984616

Moat, J., Williams, J., Baena, S., Wilkinson, T., Gole, T. W., Challa, Z. K., Demissew, S., & Davis, A. P.

(2017). Resilience potential of the Ethiopian coffee sector under climate change. *Nature Plants*,

3(7), 1–14. https://doi.org/10.1038/nplants.2017.81

Omobuwajo, T. O. (2007). BREADFRUIT AS A KEY COMPONENT OF SUSTAINABLE

LIVELIHOODS IN NIGERIA: PROSPECTS, OPPORTUNITIES AND CHALLENGES. *Acta*

Horticulturae, *757*, 121–124. https://doi.org/10.17660/ActaHortic.2007.757.15

Orwa C, Mutua A, Kindt R, Simons A, & Jamnadass R H. (2010). *Agroforestree database: A tree*

reference and selection guide version 4.0. World Agroforestry Centre (ICRAF).

Peters, A. (2021). *Why Patagonia wants you to eat breadfruit*. Fast Company.

https://www.fastcompany.com/90620093/youve-probably-never-eaten-breadfruit-heres-why-

patagonia-wants-to-change-that

Porter, J. R., Xie, L., Challinor, A. J., Cochrane, K., Howden, S. M., Iqbal, M. M., & Travasso, M. I.

(2014). Food security and food production systems. In *Climate Change 2014: Impacts,*

Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working

Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (Vol.

2014, pp. 488–489).

Ragone, D. (1997). *Breadfruit.(Parkinson) Fosberg. Promoting the conservation and use of underutilized*

and neglected crops. 10.

Ratnayake, S. S., Kumar, L., & Kariyawasam, C. S. (2020). Neglected and Underutilized Fruit Species in

Sri Lanka: Prioritisation and Understanding the Potential Distribution under Climate Change.

Agronomy, *10*(1), 34. https://doi.org/10.3390/agronomy10010034

Schlenker, W., & Roberts, M. J. (2009). Nonlinear temperature effects indicate severe damages to U.S.

crop yields under climate change. *Proceedings of the National Academy of Sciences*, *106*(37),

15594–15598. https://doi.org/10.1073/pnas.0906865106

Tan, X., Gan, T. Y., & Horton, D. E. (2018). Projected timing of perceivable changes in climate extremes

for terrestrial and marine ecosystems. *Global Change Biology*, *24*(10), 4696–4708.

https://doi.org/10.1111/gcb.14329

Taylor, M., McGregor, A., & Dawson, B. (2016). *Vulnerability of Pacific Island agriculture and forestry*

to climate change. http://www.fao.org/family-farming/detail/en/c/432913/

Trees That Feed Foundation. (n.d.). *Africa*. Retrieved September 7, 2021, from

https://www.treesthatfeed.org/countries/africa/

Trees that Feed Foundation. (2020). *Breadfruit Yield Study in Jamaica, 2018-20*.

https://treesthatfeed.org/breadfruit-yield-study/

Turi, C. E., Liu, Y., Ragone, D., & Murch, S. J. (2015). Breadfruit (Artocarpus altilis and hybrids): A

traditional crop with the potential to prevent hunger and mitigate diabetes in Oceania. *Trends in*

Food Science & Technology, *45*(2), 264–272. https://doi.org/10.1016/j.tifs.2015.07.014

Vermeulen, S. J., Campbell, B. M., & Ingram, J. S. I. (2012). Climate Change and Food Systems. *Annual*

Review of Environment and Resources, *37*(1), 195–222. https://doi.org/10.1146/annurev-environ-

020411-130608

WFP. (2014). *Climate impacts on food security and nutrition: A review of existing knowledge*.

https://www.eldis.org/document/A73884

Zerega, N., Ragone, D., & Motley, T. J. (2006). 10. Breadfruit Origins, Diversity, and Human-Facilitated

Distribution. In T. Motley (Ed.), *Darwin's Harvest* (pp. 213–238). Columbia University Press.

https://doi.org/10.7312/motl13316-011

Zerega, N., Wiesner-Hanks, T., Ragone, D., Irish, B., Scheffler, B., Simpson, S., & Zee, F. (2015).

Diversity in the breadfruit complex (Artocarpus, Moraceae): Genetic characterization of critical

germplasm. *Tree Genetics & Genomes*, *11*(1), 4. https://doi.org/10.1007/s11295-014-0824-z

Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., Huang, M., Yao, Y., Bassu, S., Ciais, P.,

Durand, J.-L., Elliott, J., Ewert, F., Janssens, I. A., Li, T., Lin, E., Liu, Q., Martre, P., Müller,

C., … Asseng, S. (2017). Temperature increase reduces global yields of major crops in four

independent estimates. *Proceedings of the National Academy of Sciences*, *114*(35), 9326–9331.

https://doi.org/10.1073/pnas.1701762114

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Ritter, C. D., Edler, D., Farooq, H., Herdean, A.,

Ariza, M., Scharn, R., Svanteson, S., Wengstrom, N., Zizka, V., Antonelli, A., ropensci, I. S.

(Irene reviewed the package for, see <https://github.com/ropensci/onboarding/issues/210>),

ropensci, F. R.-S. (Francisco reviewed the package for, & see

<https://github.com/ropensci/onboarding/issues/210>). (2020). *CoordinateCleaner: Automated*

Cleaning of Occurrence Records from Biological Collections (2.0-18) [Computer software].

https://CRAN.R-project.org/package=CoordinateCleaner

Figures and Tables

Figure 1. Range of current (1970-2000) breadfruit suitability in the global tropics and subtropics (45° N to 45° S) as predicted from bioclimactic variables from 10 arcminute resolution WorldClim v2.1 data and filtered GBIF occurrence points. Climatic suitability of breadfruit cultivation is indicated by green shading ranging from 0 (fair) to 1000 (good). (*A*) The ensemble mean-weighted SDM and 6 different individual SDMs including (*B*) MAXENT, (*C*) GLM, (*D*) GAM, (*E*) MARS, (*F*) RF, and (*G*) GBM. Black dots in (*A*) are observed occurrence points from GBIF. Pie charts in upper left of each plot indicate fraction of Fair (score=250-750) to Good (score=750+) area.

Figure 2. Future (2061-2080) breadfruit suitability range and change under stabilization and high-emission

scenarios. (a) Future suitability (Fair: score=250-750; Good: score=750+) of weighted SDM ensemble under

stabilization scenario (SSP2-4.5) and (b) high-emission scenario (SSP8-8.5) using averages from a CMIP6 ensemble

of opportunity from 8 GCMs. Change in suitable range (score>250) from baseline suitability score under (c)

stabilization scenario and (d) high-emission scenario.

Figure 3. Change in suitable breadfruit area (value>250) between the ensemble baseline SDM (B) and the

stabilization (S) and high-emission (H) scenarios within various regions in the global tropics and subtropics,

including Asia and the Pacific Islands, Africa, and Latin America, with Weak (1-3 members), Strong (4-5 members)

and Very Strong (6+ members) CMIP6 GCM average agreement.

Figure 4. Difference Plots of Top BIOCLIM Contributors to SDM. Columns show the baseline (1970-2000)

bioclimatic variables (left) and the absolute difference of the specific bioclimatic variables from 2061-2080 under

SSP2-4.5 (middle) and SSP5-8.5 (right) scenarios of the CMIP6 ensemble of opportunity downscaled to 10

arcminute resolution from WorldClim v2.1.

Supplementary Figure 1. Threshold metrics for breadfruit suitability used for SDM classification. Categories

are demarcated in relation to trends between cumulative percent of total number of points (presence and

pseudoabsence) and model suitability value (0-1000). The Fair (250-750) and Good (750-1000) suitability

categories are marked by greater prevalence of presence points, with 4 occurrence points categorized as unsuitable.

Supplementary Table 1. Relative importance (TSS) of 7 selected BIOCLIM variables within and across 6

individual SDMs in baseline (1970-2000) scenario.

	Bioclimactic Variable	MAXENT	GLM	GAM	MARS	RF	GBM	Average
BIO2	Mean Diurnal Range	0.201	0.046	0.123	0.081	0.063	0.042	0.093
BIO7	Temperature Annual Range (BIO5-BIO6)	0.312	0.543	0.403	0.381	0.312	0.454	0.401
BIO11	Mean Temperature of Coldest Quarter	0.194	0.052	0.191	0.124	0.098	0.033	0.115
BIO12	Annual Precipitation	0.232	0.115	0.216	0.130	0.057	0.029	0.130
BIO17	Precipitation of Driest Quarter	0.121	0.027	0.106	0.055	0.031	0.025	0.061
BIO18	Precipitation of Warmest Quarter	0.150	0.063	0.107	0.026	0.038	0.016	0.067
BIO19	Precipitation of Coldest Quarter	0.185	0.012	0.104	0.069	0.034	0.016	0.070

661   **Supplementary Table 2.** SDM Variability of Breadfruit Suitability across 6 individual and ensemble SDMs.

662   Suitability is categorized over 98280727 km^2 of total land area constrained within the latitudes of 45° N to 45° S.

Suitability	SDM	Area (km^2)
Fair	MAXENT	17743876
	GLM	14117175
	GAM	15835734
	MARS	12581785
	RF	8411102
	GBM	9125204
	Ensemble	12725593
Good	MAXENT	9953293
	GLM	14455265
	GAM	12461001
	MARS	13168259
	RF	16982621
	GBM	16180823
	Ensemble	13721639
Unsuitable	MAXENT	70583558
	GLM	69708287
	GAM	69983992
	MARS	72530682
	RF	72887004
	GBM	72974700
	Ensemble	71833495

663 **Supplementary Table 3. GCM Variability in Breadfruit Suitability Under Stabilization and High Emissions**

664 **Scenarios of individual CMIP6 GCMs**. All available downscaled GCMs at 10 arcminute resolution were accessed

665 from WorldClim v2.1. Breadfruit suitability is evaluated within land area constrained within the global tropics and

666 subtropics**.**

Suitability	GCM	Area - SSP2-4.5	Area - SSP5-8.5
Fair	BCC-CSM2-MR	16127181	17010437
	CanESM5	16386867	17383498
	CNRM-CM6-1	15938225	17329833
	CNRM-ESM2-1	16025173	17377793
	IPSL-CM6A-LR	15660300	17401963
	MIROC-ES2L	14538802	15165247
	MIROC6	14200092	15245630
	MRI-ESM2-0	13900888	15981020
Good	BCC-CSM2-MR	9364025	8533701
	CanESM5	8101903	7002352
	CNRM-CM6-1	9571089	8025798
	CNRM-ESM2-1	9442967	7896402
	IPSL-CM6A-LR	10249666	8673592
	MIROC-ES2L	10186645	9507514
	MIROC6	10498834	8805485
	MRI-ESM2-0	10430547	8993401
Unsuitable	BCC-CSM2-MR	72784947	72732016
	CanESM5	73778272	73881192
	CNRM-CM6-1	72765857	72919540
	CNRM-ESM2-1	72807031	73000976

	IPSL-CM6A-LR	72358041	72192452
	MIROC-ES2L	73541595	73594281
	MIROC6	73576245	74224056
	MRI-ESM2-0	73944719	73301732

Supplementary Figure 2. Suitable Breadfruit Range (a) baseline (1970-2000), (b) Future Stabilization (2061-2080), and (c) Future High Emissions (2061-2080) scenarios with country outlines using average values from a weighted SDM ensemble projected using a CMIP6 GCM ensemble of opportunity under SSPs 2 and 5.