1    **Oligogenic combinations of rare variants influence specific phenotypes in**

2    **complex disorders**

4    Vijay Kumar Pounraja[1,2] and Santhosh Girirajan[1,2,3]

6        1.  Department of Biochemistry and Molecular Biology, Pennsylvania State University,

7            University Park, PA 16802

8        2.  Bioinformatics and Genomics Graduate Program, The Huck Institute of the Life

9            Sciences, University Park, PA 16802

3.  Department of Anthropology, Pennsylvania State University, University Park, PA 16802

**Correspondence:**

Santhosh Girirajan

205A Life Sciences Building

Pennsylvania State University

University Park, PA 16802

E-mail: sxg47@psu.edu

**Classification**: Biological Sciences

**Keywords**: oligogenic, complex disorders, autism, rare variants

**ABSTRACT**

Genetic studies of complex disorders such as autism and intellectual disability (ID) are often based on enrichment of individual rare variants or their aggregate burden in affected individuals compared to controls. However, these studies overlook the influence of combinations of rare variants that may not be deleterious on their own due to statistical challenges resulting from rarity and combinatorial explosion when enumerating variant combinations, limiting our ability to study oligogenic basis for these disorders. We present a framework that combines the apriori algorithm and statistical inference to identify specific combinations of mutated genes associated with complex phenotypes. Our approach overcomes computational barriers and exhaustively evaluates variant combinations to identify non-additive relationships between simultaneously mutated genes. Using this approach, we analyzed 6,189 individuals with autism and identified 718 combinations significantly associated with ID, and carriers of these combinations showed lower IQ than expected in an independent cohort of 1,878 individuals. These combinations were enriched for nervous system genes such as *NIN* and *NGF*, showed complex inheritance patterns, and were depleted in unaffected siblings. We found that an affected individual can carry many oligogenic combinations, each contributing to the same phenotype or distinct phenotypes at varying effect sizes. We also used this framework to identify combinations associated with multiple comorbid phenotypes, including mutations of *COL28A1* and *MFSD2B* for ID and schizophrenia and *ABCA4*, *DNAH10* and *MC1R* for ID and anxiety/depression. Our framework identifies a key component of missing heritability and provides a novel paradigm to untangle the genetic architecture of complex disorders.

**SIGNIFICANCE**

While rare mutations in single genes or their collective burden partially explain the genetic basis for complex disorders, the role of specific combinations of rare variants is not completely understood. This is because combinations of rare variants are rarer and evaluating all possible combinations would result in a combinatorial explosion, creating difficulties for statistical and computational analysis. We developed a data mining approach that overcomes these limitations to precisely quantify the influence of combinations of two or more mutated genes on a specific clinical feature or multiple co-occurring features. Our framework provides a new paradigm for

53    dissecting the genetic causes of complex disorders and provides an impetus for its utility in

54    clinical diagnosis.

## INTRODUCTION

Recent human population growth has led to a rapid increase in the load of rare variants affecting functionally important regions of the genome[1–3]. Thus, rare variants are collectively more abundant in the population compared to common variants, many of which confer significant risk for neurodevelopmental disorders such as autism and intellectual disability[4]. In fact, recent studies have directly implicated rare damaging mutations that are very recent or *de novo* in more than one hundred genes towards neurodevelopmental disorders[5–7]. The ability to establish robust associations between rare variants of high effect size and complex disease has made this class of variants the primary focus of recent studies. However, a much larger class of rare and variably expressive variants that are individually less deleterious but, in combination, exert large effects towards disease is often overlooked. Variants in this category are often transmitted across generations without adverse effects on their carriers until they encounter other similar variants that, when combined, lead to genetic interactions conferring a higher risk for disease than their individual risks[8,9]. While this phenomenon underpins oligogenic models proposed over the years, studies so far have not focused on detecting combinatorial effects of specific sets of rare variants towards disease phenotypes[10–13].

Identifying the effects of specific combinations of rare variants towards disease etiology has been challenging for many reasons. *First*, combinations of rare variants are rarer, and extremely large cohorts are required to observe even a few recurrent instances of specific variant combinations[14]. Prior studies of oligogenic models for rare variants evaded this problem by aggregating variant information at the sample level and comparing the overall burden of rare variants between groups of individuals (such as cases and controls)[6,7,15,16]. *Second*, the combinatorial explosion resulting from even a small set of rare variants makes it difficult to exhaustively evaluate all combinations. While sophisticated frameworks such as network analysis and machine learning provide powerful tools to model the composite effects of thousands of variables on a complex system and predict emergent behaviors and quantitative outcomes, adapting them to exhaustively search and delineate the effects of specific combinations of variables is daunting[17,18]. Furthermore, incorporating an efficient search tool into these frameworks and extending them to detect higher-order combinatorial effects would be nearly impossible. *Third*, even when all combinations of rare variants could be exhaustively evaluated within a large cohort, there is a lack of methods that are sensitive enough to detect

small differences between comparison groups to establish statistical significance. Therefore, an

alternate approach that is highly flexible, scalable, and sensitive is necessary to address

computational and statistical challenges associated with assessing rare variant combinations.

Here, we present a combinatorial framework called *RareComb* that couples the apriori

algorithm[19] with binomial tests to overcome the limitations of data sparsity and high

dimensionality, and systematically analyzes patterns of rare variants between groups of interest

to identify specific combinations that are significantly associated with phenotypes[20]. We apply

our analysis framework to a discovery cohort of 6,189 children with autism to identify genetic

interactions involving pairs and triplets of mutated genes that are enriched in individuals with

intellectual disability compared to individuals without intellectual disability. We demonstrate

that the carriers of mutations in these specific gene pairs and triplets within an independent

cohort of 1,878 children have significantly lower-than-expected intelligence quotient (IQ) scores.

We also demonstrate the adaptability of our framework by leveraging it to identify mutated gene

pairs and triplets significantly associated with two or more comorbid phenotypes among children

with autism. Finally, we show how this generalizable and modular framework can be easily

extended to identify higher order interactions beyond pairs and triplets of variants. Our stand-

alone framework does not depend on *a priori* knowledge and can detect rare patterns from high-

dimensional genetic data to generate interpretable results, making it readily applicable for

analyzing cohorts of all size ranges to dissect the genetic basis of complex disorders.

## RESULTS

We hypothesized that two or more genes disrupted simultaneously by rare deleterious mutations contribute to a highly penetrant phenotype, as in an oligogenic model, or lead to a more severe phenotype than when each of the same genes are disrupted individually. We developed *RareComb* as a framework that combines data mining and statistical analysis to identify specific combinations (such as pairs, triplets, etc.) of rare variants that show significant associations with one or more phenotypes. *RareComb* analyzes an '*n×p*' sparse Boolean matrix with '*p*' genes in '*n*' individuals in three discrete steps (**Figure 1**). *First*, it applies the apriori algorithm independently in cases and controls to enumerate the frequency of all simultaneously mutated combinations that meet a pre-set minimum frequency threshold (**Supp. Figure 1**). *Second*, for each qualifying combination of variants, the method derives the expected frequency of simultaneously observing mutations in the constituent genes under the assumption of independence. It then independently quantifies the magnitude of deviation of the observed from the expected frequencies using binomial tests in cases and controls, and uses multiple-testing adjusted p-values to identify combinations that are statistically enriched in cases but not in controls. *Finally*, the method calculates effect sizes using Cohen's d and statistical power at 1% and 5% significance thresholds, to enable prioritization of a high confidence set of combinations that contribute to the phenotype in an oligogenic manner.

### *RareComb* identifies oligogenic combinations associated with ID and autism

We sought to identify pairs and triplets of mutated genes that are significantly associated with intellectual disability (ID) phenotypes by analyzing 6,189 affected individuals from the SPARK[21] cohort for discovery and 1,878 affected individuals from the SSC[22] cohort for validation. To facilitate cross-cohort comparison, we identified 10,217 rare variants (MAF≤1%) that were predicted to be deleterious by multiple methods and observed in both cohorts, and aggregated these variants to genes for the analysis (see **Methods**). We first categorized 1,215 probands from the SPARK cohort diagnosed with ID as cases and 4,974 probands without ID as controls (**Figure 2A**). We then applied *RareComb* to cases after constraining it to only evaluate those gene combinations in which simultaneous mutations are observed in at least five probands. We identified 25,602 pairs involving 1,956 mutated genes in cases that were observed at a higher frequency than expected under the assumption of independence. Similarly, analyzing the controls

using only the 1,956 genes mutated in cases, *RareComb* identified 148 pairs of mutated genes

that were significantly enriched in cases but not in controls (**Supp. Table 1)**, with moderate to

high effect sizes (Cohen's d, 0.08-0.15) and adequate statistical power (70%-100% at 5%

significance threshold) (**Supp. Figure 2**). These 148 gene pairs belonged to 142 probands, with

74% (105/142) of them carrying more than one significant pair. These observations suggest that

an individual can carry multiple combinations, each contributing to the same phenotype at

varying effect sizes (**Supp. Figure 3**).

151            We next sought to validate the association of these 148 mutated gene pairs towards

intellectual disability. We hypothesized that if the association of the gene pairs with ID in the

SPARK cohort were truly significant, carriers of mutations in those gene pairs would tend to

have lower than average IQ scores in the independent SSC cohort. We found that 90 of the 148

significant pairs identified in the SPARK cohort were observed in at least one proband in the

SSC cohort. These 90 mutated gene pairs were carried by 91 unique probands, whose average

full-scale IQ scores (average IQ=68.52) were lower than those of all ascertained probands in the

SSC cohort (average IQ=86). To assess the significance of this result, we performed 10,000

random draws of 91 probands from the SSC cohort to generate a simulated distribution of their

average IQ scores. The average IQ of carriers of mutated gene pairs (average IQ=68.52) was

significantly lower than the overall distribution of average IQ derived from simulations (average

IQ ranged from 73 to 92; empirical p=0) (**Figure 2B**). Furthermore, the average IQ of the 91

SSC probands with both mutated genes was significantly lower than the average IQ of 1,252

carriers of mutations in only one of the two genes (68.5 versus 82.8; Kolmogorov-Smirnov p =

$1.302\times10^{-16}$) (**Figure 2C**). When each of the 90 combinations was evaluated individually,

carriers of mutations in both genes for 73% (66/90) of the combinations showed lower IQ than

individuals with mutations in individual genes of the same combination, with 39/90 remaining

significant after multiple testing correction (**Supp. Table 2; Supp. Figure 4**). These results

provide evidence for synergistic effects of deleterious mutations within specific pairs of genes

towards ID phenotypes.

171            We also applied *RareComb* to identify gene triplets associated with intellectual disability

using the two cohorts and repeated the simulations to identify 1,593 significant combinations in

the SPARK cohort. We selected 570 high-confidence triplets (with ≥90% statistical power at 5%

significance threshold; **Supp. Table 3**) and found that 79 probands in the SSC cohort carried at

least one of these deleterious triplets. The average IQ score of individuals carrying significant

gene triplets (average IQ score=73) was significantly lower than a distribution of average IQ

scores from 10,000 draws of 79 SSC probands (average IQ score=82.5; min=72, max=94;

empirical p=0.0011; see **Supp. Figure 5**). This result reiterated that carriers of mutations in the

significant gene combinations have lower IQ than a random group of probands. Our results also

demonstrate the ability of the framework to identify higher order combinations of mutations that

are significantly associated with specific phenotypes in individuals with complex disorders.

**Oligogenic combinations are enriched for specific inheritance patterns**

As individual variants can arise *de novo* or be inherited maternally or paternally, variants in pairs

of genes can have six possible patterns of transmission (**Supp. Figure 6A**). We identified a total

of 926 occurrences of the 148 pairs of mutated genes enriched among SPARK probands with ID

(n=142 probands), of which inheritance could be determined without ambiguity for 887

instances. We found that one variant occurred *de novo* and the other variant was inherited from

the mother in 244/887 instances (27.5%). Similarly, both mutated genes were inherited from the

mother in 226/887 instances (25.4%) or occurred *de novo* in 221/887 instances (24.9%), while

the remaining fraction (~22%) of variant pairs were either inherited from both parents, inherited

from the father, or transmitted *de novo* and paternally. To assess the significance of our

observations, we performed simulations to establish a baseline expectation of proportions for

each category of parental inheritance pattern. We selected 926 pairs of genes in 1000 random

draws of all possible mutated gene pairs among SPARK probands and calculated the fraction of

instances that fell into each of the six transmission categories. The observed proportion was

higher than the simulated proportions for instances when both variants occurred *de novo* (24.9%

versus 17%, empirical p=0) and when one variant was *de novo* and the other was inherited

maternally (27.5% versus 25%, p=0.028) (**Figure 3A**). We repeated this analysis for 7,596

children affected with autism in the SPARK cohort compared to 11,740 unaffected parents and

identified 110 gene pairs significantly associated with autism (**Supp. Table 4**). Similar to the

results obtained for the ID phenotype, we found that both variants of a gene pair were more

likely to occur *de novo* (24% versus 18%, empirical p=0) or one variant occurring *de novo* and

the other inherited maternally (33% versus 26%, p=0) than expected based on simulation studies

(**Supp. Figure 7**). The enrichment of *de novo* or maternally inherited variants for significant

gene pairs aligns with published reports that severely affected children tend to carry multiple *de*
*novo* mutations or inherit pathogenic rare variants from mildly affected or unaffected carrier
mothers[16,23,24].

We then assessed whether the mutated gene pairs associated with ID were also found in
siblings of carrier probands. Restricting our analysis to families with unaffected siblings whose
probands had mutations in ID-enriched gene pairs, we found that both variants were present in
the corresponding sibling for only 53/219 (24.2%) instances of gene pairs, while 102/219
(46.6%) had variants in only one of the two genes and 64/219 (29.2%) instances had no variants
in either of the genes in the siblings (**Supp. Figure 6B)**. Using simulations, we found a
significantly higher proportion of instances with only one of the two variants present in siblings
compared to the expected values (46.6% versus 38.5%, p=0.007). Furthermore, the proportion of
observed instances with neither of the variants present in siblings (29.2% versus 33.1%,
empirical p=0.098) or both variants present in siblings (24.2% versus 28.4%, p=0.079) was
lower than expected (**Figure 3B)**. The observation that only a small fraction of unaffected
siblings carried both mutated gene pairs suggests a strong association of these gene pairs with ID
phenotypes. These results suggest that mutations in pairs of genes significantly associated with a
severe phenotype in probands are more likely to occur individually than simultaneously in
unaffected siblings of the same family.

**Genes forming oligogenic combinations are distinct from canonical autism genes**
We expanded our analysis to include all 16,556 mutated genes in the SPARK cohort, as opposed
to genes with mutations present in both the SPARK and SSC cohorts, and identified 52
significant gene pairs (**Supp. Table 5)** and 230 triplets associated with the ID phenotype (with
≥90% statistical power at 1% significance threshold; **Supp. Table 6)**. Due to the expanded
search space, the mutated gene pairs showed more significant p-values from the binomial tests
when compared to those obtained from the more restricted set of variants overlapping both
SPARK and SSC cohorts (**Supp. Figure 8**). Mutated genes within these combinations included
several genes related to nervous system development, such as *NIN*, *HDC*, *NGF*, and *BRD8*.
Furthermore, 5/52 pairs and 59/230 triplets contained at least one gene associated with autism in
the SFARI database, including *FGFR1*, associated with multiple disorders including Kallmann
syndrome[25] and Pfeiffer syndrome[26]; *RELN*, associated with temporal lobe epilepsy[27]; *SYNE1*,

associated with spinocerebellar ataxia[28,29]; and *PNPLA7*, associated with autism and ID[30]. Thus,

most genes forming combinations are not involved in canonical autism or ID disorders,

suggesting synergistic effects of these genes without prior association to disease.

We also performed gene ontology enrichment analysis for genes within the combinations

and identified seven out of nine significantly enriched GO terms to be exclusively associated

with nervous system-related functions, including synthesis and metabolism of catecholamines,

axon/neuron regeneration, and neuron generation and differentiation (**Supp. Figure 9**)[31].

Furthermore, the differences in the type and specificity of GO terms enriched for significant

pairs versus triplets were apparent, with genes forming pairs involved in nervous system function

and genes forming triplets associated with both nervous system as well as other biological

processes. We next assessed the enrichment and depletion of Human Phenotype Ontology (HPO)

terms for genes forming significant pairs towards ID phenotypes[32]. *First*, we calculated the

fraction of all 4,484 genes within the HPO database associated with each HPO term. *For*

*example*, 30% (1,366/4,484) of all genes in HPO were associated with ID. We compared these

expected values calculated for each HPO term with the corresponding fractions observed within

the 95 genes forming 52 ID-associated pairs using binomial tests. Interestingly, genes associated

with HPO terms related to neurodevelopmental phenotypes, such as ID, global developmental

delay, seizure, and microcephaly, were significantly depleted within the set of 95 genes forming

gene pairs (**Supp. Table 7**). *Next*, we evaluated whether genes within each of the 52 significant

pairs shared one or more common HPO phenotype or disease. Of the 52 pairs, only one pair

(*DNASE1 & MTR*) shared an HPO phenotype ("epilepsy"). This was significantly lower than the

expected value obtained from the distribution of the number of shared HPO phenotypes between

all possible pairs of genes in the HPO database (1/52, 1.9% ID gene pairs compared to 31.5% of

all HPO gene pairs shared one HPO phenotype, p= $2.2 \times 10^{-16}$; one-sided binomial test) (**Supp.**

**Figure 10; Supp. Table 8**). We note that the 4,484 genes within HPO are potentially biased

towards well-studied disorders, making pairs of genes drawn from HPO more likely to share

phenotypes than random pairs of genes from the genome. Overall, GO and HPO analyses show

that genes forming oligogenic combinations are involved in neuronal processes but have not

been previously connected to neurodevelopmental phenotypes, indicating the novelty of the

associations between these genes and ID phenotypes.

**Identifying variant combinations towards specific patterns of comorbid phenotypes**

We adapted our framework to identify significant associations of two or more genotypes with multiple comorbid phenotypes. To identify novel comorbid associations, we eliminated phenotypes that were highly correlated with each other, such as ADHD and reading disorder[33]. We analyzed variant profiles of 6,189 autism probands from the SPARK cohort with records of comorbid features, including 1,215 individuals with ID, 1,825 with anxiety and depression, and 332 with schizophrenia features. We assessed for significant co-occurrences of two or more mutated genes with two or more of the above phenotypes (**Figure 4**). Using one-tailed binomial tests to compare the observed frequency of combinations of genotypes and phenotypes to the expected frequency, we first identified 169 significant associations between pairs of mutated genes and two comorbid phenotypes as well as 82 combinations of three mutated genes and two comorbid phenotypes (**Supp. Tables 9 & 10**). As some of these significant genotype-phenotype combinations can be confounded by high degree of co-occurrence of mutated genes, we next calculated genotype-only p-values using binomial tests for all significant genotype-phenotype associations. For 32/169 combinations of two mutated genes and two comorbid phenotypes and 5/82 combinations of three mutated genes and two comorbid phenotypes, the composite genotype-phenotype p-values were significant while genotype-only p-values were not significant, suggesting stronger associations between these variant combinations and phenotypes. *For example*, even when variants in genes *COL28A1* and *MFSD2B* did not co-occur more frequently than expected under the assumption of independence, these mutated genes co-occurred more frequently than expected among probands with ID and schizophrenia phenotypes. Loss-of-function and rare missense mutations in *COL28A1* have been reported in individuals with autism[34,35], and *MFSD2A, a* paralog of *MFSD2B,* has been directly implicated in an autosomal recessive disorder associated with progressive microcephaly, spasticity and brain imaging abnormalities[36]. Similarly, we found *ARVCF* and *FAT1* to be significantly associated with ID and schizophrenia, with *ARVCF* mapping within the 22q11.2 DiGeorge syndrome region[37], while rare *de novo* mutations in *FAT1* being associated with autism and schizophrenia[6,38]. Finally, we found that the mutations in genes *ABCA4*, *DNAH10* and *MC1R* significantly co-occurred in individuals with ID and anxiety/depression phenotypes. These

results demonstrate the utility of identifying higher-order associations between genotypes and

phenotypes in complex disorders such as autism.

**DISCUSSION**

Current rare variant analysis strategies are geared towards either searching for individual variants

of high effect size whose influence on the phenotype is evident, such as *de-novo* gene-disruptive

mutations, or comparing rare variant burden to explain collective effects on phenotypes[7,39,40].

The wider space between these two extremes of the analysis spectrum that involves

combinations of rare variants has largely remained understudied. Although digenic diseases and

multi-hit models of complex diseases have been used to provide post-hoc explanations for an

observed phenomenon, they are not equipped to serve as a framework to actively search and

identify rare variant combinations that fit oligogenic models for specific phenotypes[9,12,13]. While

machine learning has become the de-facto approach for disease outcome predictions, the lack of

holy-grail predictors and reduced interpretability due to data sparsity makes it less fit to detect

combinatorial effects[17]. In addition, the common practice of evaluating feature importance

metrics of machine learning classifiers falls short of the objective to identify combinations of

features that exert higher effect on the phenotype than evident from their independent effects[17,18].

Furthermore, prior studies to assess combinatorial effects have been inherently biased due to

their need to minimize the search space by restricting the analysis to only a subset of genes

chosen based on *a priori* knowledge[41–43]. Here, we provide a proof-of-concept analytical

framework that remains agnostic to prior evidence and performs exhaustive searches to identify

combinatorial effects among rare variants while retaining high granularity of data and

interpretability of results.

We use our framework to identify gene pairs and triplets significantly associated with

intellectual disability and show that several constituent genes are associated with nervous system

processes. These mutated gene combinations are more likely to be inherited maternally or occur

*de novo*, are depleted in unaffected siblings from the same family, and are less likely to involve

canonical autism or ID genes, suggesting that genes forming significant combinations are less

deleterious on their own but manifest effects only when combined with other similar genes

carrying rare mutations. While previous studies have linked aggregate rare variant burden

towards intellectual disability[44,45], our results fine map the association to specific combinations

of constituent genes contributing to the burden. We propose a novel paradigm for dissecting the

complexity of genetic disorders, where an affected individual carries multiple combinations of

rare variants, and each combination contributes to either the same phenotype or distinct

phenotypes at varying effect sizes (**Figure 5**). A limitation of our method is that it tends to be

biased towards genes that are mutated frequently enough to be observed in a combination. This

limitation can be addressed by fixing specific primary variants of interest irrespective of their

frequency and screening for "second-hit" modifiers that significantly co-occur with the primary

variant, such as the co-occurrence of *RBM8A* variants in distal 1q21.1 deletion carriers

manifesting thrombocytopenia-absent-radius syndrome and *TBX6* variants in 16p11.2 deletion

carriers with scoliosis[46,47].

Our method is fast and scalable, allows for fine-tuning combinatorial searches based on

frequency, statistical power, and multiple testing criteria, and can be adapted to enable

computational approximations to further improve run time and assess higher-order combinations

beyond triplets. While larger sample sizes are generally required for detecting smaller frequency

differences, we note that our framework achieves reliable statistical power even with modest

sample sizes, implying that our framework could be applied to exome sequencing studies of

other neurodevelopmental disorders that have not been explored for combinatorial effects. This

approach can also be used to address a variety of research questions involving rare event

combinations, including searching for protective effects of rare variants where simultaneous

mutations are enriched in controls but not in cases, and finding combinations that exhibit specific

enrichment or depletion patterns in more than two phenotypic groups. In summary, we provide a

conceptual framework and the necessary tools to identify the oligogenic basis for complex

disorders such as autism and intellectual disability, which hitherto was restricted to the analysis

of canonical disorders such as Hirschsprung disease[48] and Bardet-Biedl syndrome[12].

## MATERIALS AND METHODS

We developed *RareComb* to address computational and statistical challenges associated with combinatorial analysis of rare variants. *RareComb* first uses the apriori algorithm to efficiently count the frequencies of co-occurring variant combinations. It then uses one-tailed binomial tests to compare the observed frequency of each variant combination to the expected frequency derived under the assumption of independence among the constituent variants within each combination (**Figure 1**). This method can be applied to identify variant combinations that are significantly enriched in cases but not in controls. In studies involving multiple comorbid phenotypes, this method can also be used to detect associations between specific combinations of variants and one or more (comorbid) phenotypes (see **Supplementary Note**). The general principles of our method, built using the basic axioms of probability theory, can be easily extended to a variety of problems involving rare higher-order combinations (**Supp. Figure 11**).

**Identifying frequencies of rare variant combinations**

*RareComb* utilizes the apriori algorithm to efficiently calculate frequencies of variant combinations from sparse Boolean matrices (of 0s and 1s) (**Supp. Figure 12A**). The apriori algorithm has been successfully applied to analyze consumer behavior, where identifying products frequently purchased together could benefit a company[49,50]. While an algorithm that is used to derive insights from patterns within highly frequent events (i.e. frequent itemset mining) might not seem like a good fit to analyze rare variant combinations, its ability to perform disciplined search based on both built-in and user-specified constraints makes it an ideal counting tool. *For example*, the apriori algorithm avoids enumerating each of the 50 million pairs or 167 billion triplets from just 10,000 variants, and instead prunes the search-space based on user-defined criteria such as minimum frequency threshold and size of combinations (pairs, triplets, etc.) (**Supp. Figure 12B**). *RareComb* applies an additional constraint to the algorithm to limit its search to co-occurring events, which further reduces the search space (see **Supplementary Note**). *For example*, when considering variants A and B, only the frequency of the presence of both variants (A=1 & B=1) is counted, and not absence of either or both variants (A=1 & B=0; A=0 & B=1; or A=0 & B=0).

**Statistical Inference**

*RareComb* utilizes the p-values of one-tailed binomial tests to establish the magnitude of enrichment for each rare variant combination (**Figure 1**). For each combination, *RareComb* formulates null and alternate hypotheses for the binomial test by considering the event of observing all constituent variants together within a group of individuals as success and all other possibilities as failure in a binomial trial:

$$H_0 : \pi = \pi_0$$
$$H_a : \pi > \pi_0$$

where,

$\pi$ = Probability of *observing* all constituent rare variants of a combination together within a cohort, i.e., P(A=1 & B=1)

$\pi_0$ = *Expected* probability derived from the frequency of individual variants of a combination, under the assumption of independence, i.e., P(A=1) * P(B=1).

*RareComb* then compares the null binomial distribution derived using the sample size of the group (n) and the expected probability ($\pi_0$) (i.e., $X \sim Binom(n, p = \pi_0)$) with the observed probability ($\pi$), and calculates the probability of observing rare variants occurring together at least as frequently as they were observed within the cohort (i.e. p-value).

In case-control analyses, this method is applied independently to each group, and the p-values between them are compared. The combinations exhibiting enrichment in both cases and controls, likely due to proximity of variants in linkage disequilibrium, are eliminated, following which the p-values in cases are adjusted for multiple-testing to identify statistically significant combinations that exhibit enrichment in cases but not in controls. Finally, the effect sizes are calculated using Cohen's d and the statistical power is measured using 2-sample 2-proportion tests, as additional metrics to prioritize the final set of significant rare variant combinations. In genotype-comorbid phenotype association analyses, the method is applied just once to the entire cohort, with multiple-testing adjusted p-values serving as a sufficient metric to identify high quality associations between genotypes and two or more co-occurring phenotypes.

**Statistical power and computational performance of the method**

We measured the relationship between sample size and statistical power for both binomial and 2-sample 2-proportion tests used in the framework. It took 1,356 samples for the binomial test to

achieve a statistical power of 80% to establish statistical enrichment between expected and

observed co-occurrence frequencies of 0.1% and 0.5% (**Supp. Figure 13**). This number

increased to 6,469 when the test needed to be more sensitive to compare frequencies of 0.3% and

0.5%. Similarly, it took 7,840 samples for the 2-sample 2-proportion test to achieve 80% power

to establish statistical difference between co-occurrence frequencies of 2% and 0.5% observed in

two groups (**Supp. Figure 14**). The sample size requirement increased to 14,633 to differentiate

frequencies of 1.5% and 0.5% at 80% statistical power. These results align with the known

relationship between sample size and statistical power, and indicate that our method can be

reliably applied to analyze reasonably modest-size cohorts.

We also measured the run times for the case-control analysis to identify significant pairs

and triplets of mutated genes using simulated data of three discrete sizes of samples (5,000,

10,000, and 50,000 individuals) and genes (5,000, 10,000, and 15,000 genes). The apriori

algorithm was run on single-core CPUs with 256 GB memory and was constrained to analyze

combinations observed in at least 0.15% of the samples. Given the memory-intensive nature of

the apriori algorithm implemented in the 'arules' package, 256 GB was chosen to maintain

uniformity[51]. However, smaller input files could be processed successfully using much less

memory. As expected, the runtimes were proportional to the size of the combination (pairs

versus triplets) and the number of input variables (**Supp. Figure 15**). While the increase in run

time with the increase in sample size is apparent for pairs, lower runtimes observed with running

50,000 samples compared to 5,000 samples for triplets can be attributed to stochasticity of the

input data. Overall, the analysis of gene pairs took between one minute and 12 minutes while

triplets took between two minutes and 150 minutes. Since several factors influence the runtime

of the method, a trial-and-error approach to determine an optimal minimum frequency threshold

for co-occurring events can help identify relevant combinations without resulting in insufficient

memory due to combinatorial explosion.

**Samples**

We used whole exome sequencing data from 6,189 affected males from the Simons Foundation

Powering Autism Research (SPARK)[21] and 1,878 affected males from 2,247 simplex families

from the Simons Simplex Collection (SSC)[52] cohort from the Simons Foundation Autism

Research Initiative (SFARI)[53]. We selected only male probands for our analysis to avoid any

confounding effect due to gender or ascertainment bias[54,55]. While diagnosis information for

intellectual disability (ID), anxiety, attention deficit hyperactivity disorders (ADHD),

schizophrenia, language and sleep disorders were encoded as binary variables for the SPARK

samples, full-scale intelligence quotient (IQ) scores were available for the SSC cohort.

**Data preparation and quality control**

Variant Call Format (VCF) files obtained from exome sequencing data were annotated using

ANNOVAR[56] for rsID information and variant frequency using ExAC[57] and gnomAD[58]. To

overcome the limitations of using a single method to predict pathogenicity, the effects of non-

synonymous mutations were annotated using 11 prediction methods: SIFT[59], Polyphen2[60]

(HDIV), Polyphen2 (HVAR), LRT[61], MutationTaster[62], MutationAssessor[63], FATHMM[64],

MetaSVM[65], PROVEAN[66], REVEL[67], and CADD[68]. Briefly, all missense, stop-loss/gain, and

start-loss/gain variants within exonic, 3', and 5' UTR regions with minor allele frequencies ≤1%

identified based on both ExAC and gnomAD databases were selected. Then, variants with allele

depth of ≥15 and allele balance between 25% and 75% for heterozygous variants and > 90% for

homozygous variants were selected as high-quality variants. Deleteriousness of the variants were

measured and reported differently by each prediction method. REVEL provided a score between

0 and 1, with higher scores indicating higher level of deleteriousness, while Polyphen2 and

MutationAssessor classified variants into one of three categories. *For example*, Polyphen2

classified variants as 'Deleterious', 'Possibly damaging', or 'Tolerated', while MutationAssessor

classified variants as 'High', 'Medium', or 'Low'. The other nine methods classified variants as

either 'Deleterious' or 'Tolerated'. Pathogenicity reported by each tool was encoded as a binary

variable, with the categories 'Possibly damaging' and 'Medium' encoded as 0.5. Thus, the

composite pathogenicity score derived from the 10 tools could range between 0 and 10. Missense

variants with a cumulative score of ≥4 and stop-loss/gain predicted as 'deleterious' either based

on CADD score (CADD phred >30) or MutationTaster were considered deleterious for all

analyses. Indels and other smaller structural variants were not considered, as their functional

impact could not be easily assessed.

**Gene Ontology (GO) and Human Phenotype Ontology (HPO) enrichment analyses**

Gene Ontology term enrichment analyses were performed using the 'Gene Ontology API' accessed using the 'post' command of the python package 'requests' (python version 3.7)[31]. All analyses were performed using parameters for *homo sapiens* (organism = '9606') to identify biological processes enrichment (annotDataSet = 'GO:0008150') using binomial tests. HPO enrichment analyses were performed using data from the 'genes_to_phenotype' file obtained from the HPO website[32]. Since enrichment of phenotypes is not automatically evaluated by HPO, we used customized R scripts to derive baseline expectations that could be compared against the actual observations to determine significance using the p-values from binomial tests.

**Statistical analysis**

All statistical analyses were performed using R v3.6.1 (R Foundation for Statistical Computing, Vienna, Austria)[69] and Python (v3.7)[70]. All data-related plots were generated using the R package ggplot2[71].

**Software Availability**

*RareComb* is available as an open-source (https://github.com/girirajanlab/RareComb) R package that can be downloaded from the Comprehensive R Archive Network (CRAN) repository[72]. It can also be installed into development environments via interfaces such as Rstudio[73] using the command *install.packages('RareComb')*. The tool provides several functionalities that allow users to run the types of analyses described in this manuscript. The functionalities are as follows: (1) Identify rare event combinations statistically enriched within a single group; (2) Identify rare event combinations statistically enriched in cases but not in controls; (3) Identify rare event combinations enriched in cases but depleted in controls; (4) Identify statistically enriched rare event combinations that include at least one element from an user-supplied list; and (5) Identify genotypes statistically enriched within individuals manifesting two or more comorbid phenotypes. Each functionality takes a Boolean matrix as input and provides a set of user-adjustable parameters to customize the analysis, and delivers the results in a tabular format as csv files. Detailed instructions on the available functionalities and parameters built into *RareComb* and their usage can be found on the GitHub page or CRAN website. A shiny app illustrating the ideas behind *RareComb* is available online at https://girirajanlab.shinyapps.io/RareComb/ [74].

**DECLARATIONS**

**Ethics approval and consent to participate**

As these data were de-identified, all our samples were exempt from IRB review and conformed

to the Helsinki Declaration. No other approvals were needed for the study.

**Consent for publication**

All authors agree and consent for publication of the manuscript.

**Competing interests**

The authors declare that no competing interests exist in relation to this work.

**Authors' contributions**

VK and SG conceived the project. VK performed the analyses, generated the plots/images, and

wrote and revised the manuscript; SG supervised the research and wrote and revised the

manuscript. All authors read and approved the final draft of the manuscript.

**Acknowledgements**

We thank Naomi Altman, Yifei Huang, Dajiang Liu, Matthew Jensen, and Corrine Smolen for

constructive comments on the manuscript. This work was supported by R01-MH107431, R01-

GM121907, Seed Grants program from the Institute of Computational and Data Sciences at Penn

State, and resources from the Huck Institutes of the Life Sciences (to SG). The funding bodies

had no role in data collection, analysis, and interpretation. The authors are grateful to all the

families who participated in the SSC and SPARK consortia, as well as the principal

investigators, clinical sites, and staff for the consortia. The authors appreciate obtaining access to

genetic and phenotypic data for SPARK and SSC through the Simons Foundation Autism

Research Initiative (SFARI) Base. Approved researchers can obtain the SSC and SPARK

population datasets described in this study by applying at https://base.sfari.org.

**References**

1. Coventry, A. *et al.* Deep resequencing reveals excess rare recent variants consistent with explosive population growth. *Nat. Commun.* **1**, 131 (2010).

2. Keinan, A. & Clark, A. G. Recent explosive human population growth has resulted in an excess of rare genetic variants. *Science.* **336**, 740–743 (2012).

3. Tennessen, J. A. *et al.* Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science.* **337**, 64–69 (2012).

4. McClellan, J. & King, M. C. Genetic heterogeneity in human disease. *Cell* **141**, 210–217 (2010).

5. Wilfert, A. B. *et al.* Recent ultra-rare inherited variants implicate new autism candidate risk genes. *Nat. Genet.* **53**, 1125–1134 (2021).

6. Iossifov, I. *et al.* The contribution of de novo coding mutations to autism spectrum disorder. *Nature* **515**, 216–221 (2014).

7. Sebat, J. *et al.* Strong Association of De Novo Copy Number Mutations with Autism. *Science.* **316**, 445–449 (2007).

8. Badano, J. L. & Katsanis, N. Beyond mendel: An evolving view of human genetic disease transmission. *Nat. Rev. Genet.* **3**, 779–789 (2002).

9. Gifford, C. *et al.* Oligogenic inheritance of a human heart disease involving a genetic modifier. *Science.* **364**, 865–870 (2019).

10. Pizzo, L. *et al.* Rare variants in the genetic background modulate cognitive and developmental phenotypes in individuals carrying disease-associated variants. *Genet. Med.* **21**, 816–825 (2019).

11. Girirajan, S. *et al.* A recurrent 16p12.1 microdeletion supports a two-hit model for severe developmental delay. *Nat. Genet.* **42**, 203–209 (2010).

12. Badano, J. L. *et al.* Dissection of epistasis in oligogenic Bardet-Biedl syndrome. *Nature* **439**, 326–330 (2006).

13. Leblond, C. S. *et al.* Genetic and functional analyses of SHANK2 mutations suggest a multiple hit model of autism spectrum disorders. *PLoS Genet.* **8**, e1002521 (2012).

14. Uricchio, L. H., Zaitlen, N. A., Ye, C. J., Witte, J. S. & Hernandez, R. D. Selection and explosive growth alter genetic architecture and hamper the detection of causal rare variants. *Genome Res.* **26**, 863–873 (2016).

571    15.    Halvorsen, M. *et al.* Increased burden of ultra-rare structural variants localizing to

572            boundaries of topologically associated domains in schizophrenia. *Nat. Commun.* **11**, 1842

573            (2020).

574    16.    Krumm, N. *et al.* Excess of rare, inherited truncating mutations in autism. *Nat. Genet.* **47**,

575            582–588 (2015).

576    17.    Murdoch, W. J., Singh, C., Kumbier, K., Abbasi-Asl, R. & Yu, B. Definitions, methods,

577            and applications in interpretable machine learning. *Proc. Natl. Acad. Sci. U. S. A.* **116**,

578            22071–22080 (2019).

579    18.    Molnar, C., Casalicchio, G. & Bischl, B. Interpretable Machine Learning – A Brief

580            History, State-of-the-Art and Challenges. *Commun. Comput. Inf. Sci.* **1323**, 417–431

581            (2020).

582    19.    Agarwal, R. & Srikant, R. Fast algorithms for mining association rules. *Proc. 20th VLDB*

583            *Conf.*

584    20.    Agrawal, Rakesh; Ramakriahnan, S. Fast Algorithms for Mining Association Rules. *Proc.*

585            *20th VLDB Conf.* 487–499 (1994).

586    21.    The SPARK Consortium. SPARK: A US Cohort of 50,000 Families to Accelerate Autism

587            Research. *Neuron* **97**, 488–493 (2018).

588    22.    Fischbach, G. D. & Lord, C. The simons simplex collection: A resource for identification

589            of autism genetic risk factors. *Neuron* **68**, 192–195 (2010).

590    23.    Girirajan, S. *et al.* Phenotypic Heterogeneity of Genomic Disorders and Rare Copy-

591            Number Variants. *N. Engl. J. Med.* **367**, 1321–1331 (2012).

592    24.    Turner, T. N. *et al.* Genomic Patterns of De Novo Mutation in Simplex Autism. *Cell* **171**,

593            710–722 (2017).

594    25.    Dodé, C. *et al.* Loss-of-function mutations in FGFR1 cause autosomal dominant Kallmann

595            syndrome. *Nat. Genet.* **33**, 463–465 (2003).

596    26.    Schell, U. *et al.* Mutations in FGFR1 and FGFR2 cause familial and sporadic pfeiffer

597            syndrome. *Hum. Mol. Genet.* **4**, 323–328 (1995).

598    27.    Dazzo, E. *et al.* Heterozygous Reelin Mutations Cause Autosomal-Dominant Lateral

599            Temporal Epilepsy. *Am. J. Hum. Genet.* **96**, 992–1000 (2015).

600    28.    Yoshinaga, T. *et al.* A novel frameshift mutation of SYNE1 in a Japanese family with

601            autosomal recessive cerebellar ataxia type 8. *Hum. Genome Var.* **4**, 17052 (2017).

29.    Synofzik, M. *et al.* SYNE1 ataxia is a common recessive ataxia with major non-cerebellar
features: A large multi-centre study. *Brain* **139**, 1378–1393 (2016).

30.    Prasad, A. *et al.* A Discovery resource of rare copy number variations in individuals with
autism spectrum disorder. *G3 Genes, Genomes, Genet.* **2**, 1665–1685 (2012).

31.    Mi, H., Muruganujan, A., Ebert, D., Huang, X. & Thomas, D. PANTHER version 14 :
more genomes , a new PANTHER GO-slim and improvements in enrichment analysis
tools. *Nucleic Acids Res.* **47**, 419–426 (2019).

32.    Köhler, S. *et al.* The human phenotype ontology in 2021. *Nucleic Acids Res.* **49**, D1207–
D1217 (2021).

33.    Gilger, J. W., Pennington, B. F. & DeFries, J. C. A Twin Study of the Etiology of
Comorbidity: Attention-deficit Hyperactivity Disorder and Dyslexia. *J. Am. Acad. Child*
*Adolesc. Psychiatry* **31**, 343–348 (1992).

34.    Krumm, N. *et al.* Transmission disequilibrium of small CNVs in simplex autism. *Am. J.*
*Hum. Genet.* **93**, 595–606 (2013).

35.    Guo, H. *et al.* Genome-wide copy number variation analysis in a Chinese autism spectrum
disorder cohort. *Sci. Rep.* **7**, 44155 (2017).

36.    Guemez-Gamboa, A. *et al.* Inactivating mutations in MFSD2A, required for omega-3 fatty
acid transport in brain, cause a lethal microcephaly syndrome. *Nat. Genet.* **47**, 809–813
(2015).

37.    Sanders, A. R. *et al.* Haplotypic association spanning the 22q11.21 genes COMT and
ARVCF with schizophrenia. *Mol. Psychiatry* **10**, 353–365 (2005).

38.    Kenny, E. M. *et al.* Excess of rare novel loss-of-function variants in synaptic genes in
schizophrenia and autism spectrum disorders. *Mol. Psychiatry* **19**, 872–879 (2014).

39.    Zheng, G. X. Y. *et al.* Haplotyping germline and cancer genomes with high-throughput
linked-read sequencing. *Nat. Biotechnol.* **34**, 303–311 (2016).

40.    Girirajan, S. *et al.* Relative burden of large CNVs on a range of neurodevelopmental
phenotypes. *PLoS Genet.* **7**, e1002334 (2011).

41.    Papadimitriou, S. *et al.* Predicting disease-causing variant combinations. *Proc. Natl. Acad.*
*Sci. U. S. A.* **116**, 11878–11887 (2019).

42.    Kerner, G. *et al.* A genome-wide case-only test for the detection of digenic inheritance in
human exomes. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 19367–19375 (2020).

43.   Schaaf, C. P. *et al.* Oligogenic heterozygosity in individuals with high-functioning autism
spectrum disorders. *Hum. Mol. Genet.* **20**, 3366–3375 (2011).

44.   Singh, T. *et al.* The contribution of rare variants to risk of schizophrenia in individuals
with and without intellectual disability. *Nat. Genet.* **49**, 1167–1173 (2017).

45.   Fitzgerald, T. W. *et al.* Large-scale discovery of novel genetic causes of developmental
disorders. *Nature* **519**, 223–228 (2015).

46.   Albers, C. A. *et al.* Compound inheritance of a low-frequency regulatory SNP and a rare
null mutation in exon-junction complex subunit RBM8A causes TAR syndrome. *Nat.*
*Genet.* **44**, 435–439 (2012).

47.   Yang, N. *et al.* TBX6 compound inheritance leads to congenital vertebral malformations
in humans and mice. *Hum. Mol. Genet.* **28**, 539–547 (2019).

48.   Gabriel, S. B. *et al.* Segregation at three loci explains familial and population risk in
Hirschsprung disease. *Nat. Genet.* **31**, 89–93 (2002).

49.   Brijs, T., Swinnen, G., Vanhoof, K. & Wets, G. Using association rules for product
assortment decisions. *ACM* (1999). doi:10.1145/312129.312241

50.   Glance, N. *et al.* Deriving marketing intelligence from online discussion. *Proc. ACM*
*SIGKDD Int. Conf. Knowl. Discov. Data Min.* 419–428 (2005).
doi:10.1145/1081870.1081919

51.   Hahsler, M., Grun, B. & Hornik, K. arules – A Computational Environment for Mining
Association Rules and Frequent Item Sets. *J. Stat. Softw.* **14**, 1–6 (2005).

52.   Sanders, S. J. *et al.* Insights into Autism Spectrum Disorder Genomic Architecture and
Biology from 71 Risk Loci. *Neuron* **87**, 1215–1233 (2015).

53.   SFARI Base. *https://www.sfari.org/resource/sfari-base/*

54.   Jacquemont, S. *et al.* A higher mutational burden in females supports a 'female protective
model' in neurodevelopmental disorders. *Am. J. Hum. Genet.* **94**, 415–425 (2014).

55.   Polyak, A., Rosenfeld, J. A. & Girirajan, S. An assessment of sex bias in
neurodevelopmental disorders. *Genome Med.* **7**, 94 (2015).

56.   Wang, K., Li, M. & Hakonarson, H. ANNOVAR: Functional annotation of genetic
variants from high-throughput sequencing data. *Nucleic Acids Res.* **38**, e164 (2010).

57.   Exome aggregation Consortium. Analysis of protein-coding genetic variation in 60,706
humans. *Nature* **536**, 285–291 (2016).

58. Genome Aggregation Database Consortium. The mutational constraint spectrum
quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).

59. Ng, P. C. & Henikoff, S. SIFT: Predicting amino acid changes that affect protein function.
*Nucleic Acids Res.* **31**, 3812–4 (2003).

60. Adzhubei, I. A. *et al.* A method and server for predicting damaging missense mutations.
*Nat. Methods* **7**, 248–249 (2010).

61. Chun, S. & Fay, J. C. Identification of deleterious mutations within three human genomes.
*Genome Res.* **19**, 1553–1561 (2009).

62. Schwarz, J. M., Rödelsperger, C., Schuelke, M. & Seelow, D. MutationTaster evaluates
disease-causing potential of sequence alterations. *Nat. Methods* **7**, 575–576 (2010).

63. Reva, B., Antipin, Y. & Sander, C. Predicting the functional impact of protein mutations:
Application to cancer genomics. *Nucleic Acids Res.* **39**, 37–43 (2011).

64. Shihab, H. A. *et al.* Predicting the Functional, Molecular, and Phenotypic Consequences
of Amino Acid Substitutions using Hidden Markov Models. *Hum. Mutat.* **34**, 57–65
(2013).

65. Kim, S., Jhong, J. H., Lee, J. & Koo, J. Y. Meta-analytic support vector machine for
integrating multiple omics data. *BioData Min.* **10**, 2 (2017).

66. Choi, Y. & Chan, A. P. PROVEAN web server: A tool to predict the functional effect of
amino acid substitutions and indels. *Bioinformatics* **31**, 2745–2747 (2015).

67. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of
Rare Missense Variants. *Am. J. Hum. Genet.* **99**, 877–885 (2016).

68. Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J. & Kircher, M. CADD: Predicting
the deleteriousness of variants throughout the human genome. *Nucleic Acids Res.* **47**,
D886–D894 (2019).

69. R Core Team. R: A language and environment for statistical computing. (2019).

70. Van Rossum, Guido; Drake, F. L. Python 3 Reference Manual. (2009).

71. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York,
2016).

72. CRAN Repository. Available at: https://cran.r-
project.org/web/packages/available_packages_by_name.html.

73. RStudio Team. RStudio: Integrated Development for R. (2020).

74.    Chang, W., Cheng, J., Allaire, J., Xie, Y. & McPherson, J. shiny: Web Application

696           Framework for R. (2020).

 **MAIN FIGURES**

**Figure 1: Conceptual overview of combinatorial analyses using *RareComb*.** A Boolean representation of genotype (mutated genes, G1, G2, etc) and disease status for probands (P1, P2, etc) is shown. In *step 1*, the apriori algorithm is applied to the Boolean input matrix to calculate the frequencies of individual (for example, G1) and simultaneous occurrences of events (G1 and G2) that meet the user-specified criteria, including the size of combinations (pairs, triplets, etc.) and minimum frequency threshold of simultaneous occurrences. In *step 2*, independently in case and control groups, for each combination, the binomial test is applied to compare the observed frequency of simultaneous occurrence of events with its corresponding null binomial distribution of the expected frequencies calculated under the assumption of independence. Binomial test for gene pair G3 and G4 is shown as an example.

**Figure 2: Combinations of rare variants contributing to intellectual disability (ID) phenotype. (A)** An outline of the approach used to identify and validate mutated gene pairs and triplets enriched in probands with ID is shown. We tested whether mutated gene pairs identified as significant in one cohort (SPARK) are also associated with severe phenotypes in an independent cohort (SSC). To test this, we obtained the mean IQ score of individuals from the SSC cohort carrying significant combinations identified from the SPARK cohort. Empirical p-values were then calculated based on the deviation of the mean IQ from the distribution of mean IQ scores obtained from 10,000 random draws in the simulation. **(B)** The mean IQ of individuals with mutated gene pairs in the SSC cohort was significantly lower (empirical p-value=0) when compared to the distribution of mean IQ scores obtained from the simulation. **(C)** Histogram shows the distributions of IQ scores of SSC probands who carried mutations in either genes versus both constituent genes of the significant gene pairs. The distributions were significantly different from each other (p-value = $1.302 \times 10^{-6}$, Kolmogorov-Smirnov test).

**Figure 3: Analysis of parental and sibling inheritance patterns of significant gene pairs associated with ID. (A)** Fraction of all instances of significant gene pairs observed within each of the six possible parental inheritance patterns (red) compared against 1,000 simulations is shown (blue). During each simulation, random mutated gene pairs from the SSC cohort were selected, the inheritance status of the mutations was identified, and the fraction of those instances

belonging to one of the six pre-defined categories was calculated. Comparing the observed

fractions with the simulated fractions indicate statistical enrichment for two specific inheritance

patterns based on empirical p-values: both variants being *de novo*, and one variant being *de novo*

and the other transmitted from the mother. **(B)** Histograms show the carrier status of significant

gene pairs in siblings of carrier probands (red) compared against 1,000 simulations (blue).

Among significant pairs, both genes were mutated in only 24.2% of all siblings (compared to

28.4% in simulations), whereas one of the two genes was mutated in 46.6% of all siblings

(compared to 38.5% in simulations). These results show that mutations are more likely to be

observed in just one of the two genes within the gene pairs and are less likely to be observed

simultaneously in siblings of carrier probands.

**Specific combinations of mutated genes significantly associated with comorbid phenotypes**

**Figure 4: Analysis of comorbid phenotypes using *RareComb*.** We analyzed the genotypes of probands with anxiety/depression, ID, or schizophrenia. The heatmap shows combinations of two or three mutated genes that were significantly enriched in individuals with specific patterns of comorbid phenotypes compared to the expected frequency under the assumption of independence.

**Rare variants and their combinations associated with phenotype**

**Figure 5: Rare variant models for complex disorders.** The schematic shows two models for

the genetic etiology of complex disorders. Circles represent rare variants present that are either

*de novo* or inherited from a parent. On the left, individual high-effect *de novo* variants are

strongly associated with a phenotype of interest. On the right, rare variants within an individual

combine in multiple ways and contribute towards distinct phenotypes. The thickness of the

connecting lines denotes effect sizes, and an affected individual can carry multiple oligogenic

combinations of rare variants, each of which contributes to the same or distinct phenotypes. This

extension of the oligogenic model enables further dissection of the genetic architecture of

complex disorders.