Anther cones increase pollen release in buzz-pollinated *Solanum* flowers

Mario Vallejo-Marín[1], *

Carlos Eduardo Pereira Nunes[1]

Avery Leigh Russell[2]

[1]Biological and Environmental Sciences. University of Stirling, Stirling, FK9 4LA, Scotland, United Kingdom.

[2] Department of Biology, Missouri State University, Springfield, Missouri 65897. U.S.A.

*Author for correspondence. Email: mario.vallejo@stir.ac.uk

Running title: Functional significance of anther cones

Author contributions: MVM, CEPN, and ALR conceived the idea. MVM designed and performed the experiments, wrote the software, and analysed the data. CEPN collected and analysed data. MVM, CEPN, and ALR wrote the manuscript and contributed to reviewing and editing it. MVM acquired the funding.

Acknowledgements: We thank L. Nevard and the rest of the Vallejo-Marin Lab for feedback and discussions during the development of this project, and J. Weir for support in the Controlled Environment Facility. We thank S. C. H. Barrett and A. Etcheverry for sharing their seed collection of *S. sisymbriifolium*. This research was supported by a grant from The Leverhulme Trust (RG-2018-235) to MVM.

Data accessibility: The vibration and pollen data will be made publicly available in DataSTORRE (https://datastorre.stir.ac.uk) upon publication.

Abstract

The widespread evolution of tube-like anthers releasing pollen from apical pores is associated with buzz pollination, in which bees vibrate flowers to remove pollen. The mechanical connection among anthers in buzz-pollinated species varies from loosely held conformations, to anthers tightly held together with trichomes or bio-adhesives forming a functionally joined conical structure (anther cone). Joined anther cones in buzz-pollinated species have evolved independently across plant families and via different genetic mechanisms, yet their functional significance remains mostly untested. We used experimental manipulations to compare vibrational and functional (pollen release) consequences of joined anther cones in three buzz-pollinated species of *Solanum* (Solanaceae). We applied bee-like vibrations to focal anthers in flowers with ("joined") and without ("free") experimentally created joined anther cones, and characterised vibrations transmitted to other anthers and the amount of pollen released. We found that joined anther architectures cause non-focal anthers to vibrate at higher amplitudes than free architectures. Moreover, in the two species with naturally loosely held anthers, anther fusion increases pollen release, while in the species with a free but naturally compact architecture it does not. We discuss hypotheses for the adaptive significance of the convergent evolution of joined anther cones.

Keywords: Anther cone, bees, buzz pollination, convergent evolution, pollen release, pollination.

Introduction

Flowers are extremely morphologically diverse, and establishing how this morphological diversity affects function has long been a focus of research (Darwin 1877; Vogel 1996). Buzz-pollinated plants capture the close relationship between floral form and function. In these species, modifications of floral structures result in morphologies that require the visits of bees that produce vibrations to remove pollen grains (Macior 1968; Thorp and Estes 1975; Buchmann et al. 1977). The floral vibrations produced by the bee cause the anthers to shake, transmitting energy to the pollen grains inside the anthers and causing them to be propelled outwards through the apical pores (Buchmann and Hurley 1978). In buzz-pollinated plants, floral structures, usually the anthers, but sometimes the corolla, have evolved a tubular form that retains the pollen grains inside after anthesis (Buchmann 1983; Vallejo-Marín 2019). A taxonomically widespread floral form of buzz-pollinated plants that has evolved convergently across multiple plant families is the *Solanum*-like or "solanoid" flower, named by Fægri (1986) after the canonical flower form of *Solanum* (Solanaceae) (Endress 1996a; De Luca and Vallejo-Marin 2013; Russell et al. 2016). In *Solanum*-like flowers, the anthers are often arranged in the centre of the flower forming a structure that resembles a cone (Faegri 1986). The degree to which the anthers in the solanoid flower are physically connected to one another varies. In one extreme, the enlarged stamens might be held loosely towards the centre of the flower, with each individual stamen capable of relatively independent movement from the other stamens ("free" stamen architecture). Other species may have connivent anthers that are closely pressed together yet non-joined. In the other extreme, anthers can be physically attached to each other (postgenitally connate; Endress 1996a), forming a single conical structure (Glover et al. 2004). This type of joined anther cone ("joined" architecture) has evolved multiple independent times in different plant groups including species in the families Anthericaceae, Luzuriageaceae, Pittosporaceae, Tecophilaceae and Solanaceae (Endress 1996a; Glover et al. 2004). Despite the repeated evolution of conical anther architecture across different species, to date, no studies have tried to empirically evaluate its functional significance.

During buzz pollination, an individual bee might only vibrate one or few anthers. However, the vibrations generated by the bee's thorax and applied to this focal anther(s), propagate through the flower (Brito et al. 2020; Nevard et al. 2021), and can cause pollen release even in distal anthers that are not in direct contact with the bee's body (Arroyo-Correa et al. 2019). These oscillations in the focal anther(s) can cause other stamens to vibrate via two transmission pathways: **(1) Filament pathway.** Anthers are attached to the corolla or to the base of the flower

via a filament that, in buzz pollinated plants is usually short and relatively stiff. The oscillation of the focal anther can thus cause shaking in the whole flower via the filament attachment, which in turn causes vibrations in other, non-focal anthers. **(2) Anther-anther pathway.** In species with floral architectures in which stamens are held closely together, for example forming a cone (joined anthers) (Glover et al. 2004), vibrations can be transmitted by direct anther-anther contact from the vibrating focal anther to adjacent anthers even when these distal anthers are not touching the bee's body. Recent work across buzz-pollinated flowers with different morphologies in three different plant families (Solanaceae, Primulaceae and Gentianaceae), suggests that stamen architecture, defined as the stamen's relative sizes, degree of fusion and their spatial and functional connections (Endress 1996b), affects the transmission of vibrations (Nevard et al. 2021). Therefore, variation in stamen architecture could be associated with different types of vibrations experienced by distal, non-focal anthers during buzz pollination with potential consequences for pollen release and pollen placement on the pollinator's body (Glover et al. 2004; Nevard et al. 2021).

   Here we use an experimental approach to compare the vibrational and functional (pollen release) consequences of joined anther cones in buzz-pollinated species in the genus *Solanum* (Solanaceae). Specifically, we address the following two questions: **(1)** How do the vibrations experienced in stamens differ between floral configurations with free vs. joined anthers? We hypothesise that when vibrations are applied to a focal anther (proximate anther), the vibration amplitude experienced by distal anthers (those anthers not directly being vibrated) is higher in floral configurations with joined anthers than in floral configurations with free anthers. Our hypothesis assumes that species with loose anthers mainly transmit vibrations to distal anthers via the filament pathway, while anther fusion enables vibration transmission via both the filament and anther-anther pathways. **(2)** How does anther fusion into a cone affects pollen release upon vibrations? We hypothesise that the higher vibration amplitude of anthers in joined architectures result in higher pollen release compared to free anther configurations. Our hypothesis is based on the fact that higher vibration amplitudes (e.g., higher velocity or acceleration amplitude) have been shown to be theoretically (Buchmann and Hurley 1978; Hansen et al. 2021) and empirically associated with higher rates of pollen release (De Luca et al. 2013; Rosi-Denadai et al. 2020; Kemp and Vallejo-Marin 2021). If joined anther architectures are associated with higher vibration magnitudes across more anthers (both focal and distal anthers), then we would expect pollen release to be proportionally higher as well.

Material and Methods

Study system

Solanum is the largest buzz-pollinated genus of flowering plants with approximately 1,400 species (Knapp 2002; Särkinen et al. 2013). Within the genus *Solanum* L. and its close relatives, joined anther cones have evolved independently at least four times: in tomatoes (*S. lycopersicum* L., sect. *Lycopersicon*) and its wild relatives, in *S. dulcamara* L. (sect. *Dulcamara*), in *S. luridifuscescens* Bitter (sect. *Cyphomandropsis*) and related taxa (Glover et al. 2004; Falcão et al. 2016), and in some species of *Lycianthes* such as *L. synanthera* and *L. anomala* (Dean et al. 2020). Strikingly, the joined anther cone in clades of *Solanum* is formed via different attachment mechanisms. In *S. lycopersicum* and *S. luridifuscescens*, the anther cone is formed via interlocking epidermal cells, joining adjacent anthers, while in *S. dulcamara* smooth anthers are held together by adhesive secretions (Glover et al. 2004; Falcão et al. 2016). In some cases, the pores of individual anthers in species with joined cones can secondarily evolve increasingly longitudinally dehiscent slits as in *S. lycopersicum* and related taxa. In these species, the slits open to the interior of the joined cone which functions as a single poricidal unit (Endress 1996a).

We studied three *Solanum* species from the subgenus Leptostemonum that differ in stamen architecture, specifically, the extent to which the anthers are loosely or closely held together: *Solanum sisymbriifolium* Lam. and *S. elaeagnifolium* Cav. have free, relatively loose, stamen architectures, while *S. pyracanthos* Lam. has stamens that, although not joined, are held closely together forming a cone-like structure. In *S. sisymbriifolium* and *S. elaeagnifolium,* the petals become reflexed soon after anthesis, and the anthers, which are loosely connivent at anthesis, become increasingly spread out (Knapp 2014a; Vorontsova and Knapp 2014). In *S. pyracanthos*, the petals are slightly reflexed, and the stamens are free but held closely together forming a conical structure that persists throughout anthesis. All three species are nectarless, andromonoecious (producing both hermaphroditic and staminate flowers in the same individual) (D'Arcy 1992; Knapp 2014a; Vorontsova and Knapp 2014), and present flowers to pollinators more or less horizontally, with the anthers' long axis pointing parallel to the ground (Figure 1). We used seeds of two accessions of *S. sisymbriifolium*, either from seeds collected in the field and outcrossed in the glasshouse or sourced from a commercial provider (Chiltern Seeds, Wallingford, UK) (Table 1)*.* For *S. elaeagnifolium* we used two accessions of this species collected in arid regions in Mexico where they formed locally abundant populations along roadsides and train tracks (Table

1). A summary of the distribution, floral characteristics, and floral visitors of these species as well as the accessions used in this study is presented in Table 1.

Plant growth

Seeds were germinated following a 24h treatment with 1000ppm aqueous solution of gibberellic acid (GA3, Asklepios-seeds, Bad Liebenzell, Germany) at room temperature. Approximately 10 days after germination, seedlings were transplanted to 1.5L plastic pots with an 80:20 compost mix of John Innes No. 2 and medium grade perlite (LBS Horticulture, Colne, UK), and placed in a Snijders Microclimate cabinet in the Controlled Environment Facilities at the University of Stirling with the following growth conditions: 16h-light/8h-dark cycles, at 27ºC/25ºC, with constant 50% RH. A few additional plants of *S. pyracanthos* and *S. elaeagnifolium* were transplanted to 10L pots and placed in the glasshouse with 16h of supplemental light using compact fluorescent lamps and provided with heating to keep temperature above 16C at night. Plants were fertilised as needed with Tomorite Concentrated Tomato Food (Levington, Surrey, UK).

Signal generation

To mimic a vibration produced by bees during buzz pollination, we synthesised a pure tone at 300Hz of two seconds in duration with a 50ms fade-in and fade-out feature using *seewave* (Sueur et al. 2008) and saved it as a Mono WAV file at a sampling rate of 44.1kHz. The frequency and duration of the synthetic buzz is within the range observed in buzz pollinating bees (De Luca and Vallejo-Marin 2013; De Luca et al. 2019). The 50ms fade-in and fade-out features in the synthesised signal were introduced to avoid amplitude spikes in the playback equipment caused by a rapid voltage change.

Vibration system

We built a vibration system to generate and apply vibrations to experimental flowers (Figure 2). A permanent magnetic shaker (LDV210, Brüel & Kjær, Nærum, Denmark) attached to a linear power amplifier (LDS-LPA100, Brüel & Kjær) that received the signal played back in the computer was used to generate the vibration. An M4, 10cm stainless steel screw ("wand") (ACCU, Huddersfield, UK) was attached to the magnetic shaker. A miniature IPC force sensor (209C11, PCB Piezotronics) was placed at the end of the wand. The system was attached to the flower using an entomological pin "00" of 0.30mm in diameter (Austerlitz, Entomoravia, Czech Republic) cut 10 mm from the tip and superglued to a head screw in the force sensor (Figure 2).

Vibration playback

The synthesised signal was played back in Audacity ver. 3.0 (Audacity 2021) using a laptop

computer (HP Elitebook) with the volume set to 70% and output to the linear amplifier. We

adjusted the gain in the linear power amplifier by hand to generate approximately 80mm/s peak

velocity (~56mm/s, Root Mean Square (RMS) velocity) measured in a small piece of retroreflective

tape at the base of the pin. The tape was kept in place after calibration. The value of peak velocity

was chosen to be within the range of values observed in buzz-pollinating bumblebees (De Luca et

al. 2013), and used previously in pollen release experiments (Kemp and Vallejo-Marin 2021).

Calibration measurements were taken and recorded each day at the start and end of the

experimental run.

Experimental treatment

Flowers were collected in the morning from plants growing in the cabinets or, more rarely, the

glasshouses and immediately transferred to a temperature- and humidity-controlled laboratory

(19C, 40% RH) for data acquisition. The age of the flower was recorded as days since anthesis (day

1 = flowered opened that day). Flowers were removed from the inflorescence by cutting at the

base of the pedicel and placing them in wet floral foam (Oasis Ideal Floral Foam Maxlife, Smithers-

Oasis UK Ltd, Washington, UK) on a plastic container. Flowers were measured within 1-3 hours of

being removed from plant.

Once in the lab, each flower was randomly assigned to one of two treatments: (1) **Free**

configuration: This represents the natural arrangement of the flowers. A small amount of PVA

glue was applied as a sham treatment to the outside of the anther. (2) **Joined configuration:** The

anthers were glued together using a small amount of PVA applied along the lateral edge of each

anther without blocking the pores. The treatment was applied with the help of a dissecting scope

(6.7x magnification). Every flower in the experiment experienced both treatments assigned in a

random order. When the free configuration treatment was applied to a flower that had been

previously glued for the joined configuration treatment, the anthers were freed using fine-tip

forceps and carefully separating the anthers from each other while leaving the glue on the anther

(in some cases a small amount of glue fell off in the process of freeing the anthers).

To increase the reflectivity of the anther surface for Doppler vibrometry, we placed a small

square of retroreflective tape (approximately 1-4mm^2) onto a single anther in the adaxial side of

the flower (anther 3 or 4 in Figure 3) depending on which one presented a surface that was

perpendicular to the laser beam and parallel to the axis of the vibrations produced by the shaker

(Figure 2). The tape was placed as close to the tip of the anther as possible, without blocking the

pores. The tape was attached to anthers 3 or 4 (at the top of the flower). The shaker pin was

inserted at the base of anther 1 (the lowest most anther, see Figure 3). Sometimes, we used the

dissecting microscope to draw a small dot made with a black marker to help placing the pin in the

exact desired location. The pin made a microscopic wound in the anther. The pin was carefully

pushed into the anther without going all the way through.

During the experiment, the flower was held by the pedicel using a stainless-steel micro-V

clamp (VK250/M; Thorlabs Ltd., Ely, UK) on a 100mm post (TR100/M), held in a 40mm post holder

(PH40/M), perpendicular to the ground. The post holder was screwed in an adjustable-angle

mounting plate (AP180/M; Thorlabs) placed on a linear stage (M-UMR8.25; Newport Spectra-

Physics Ltd, Didcot, UK) with a standard resolution micrometer (BM17.25; Newport) (see Jankauski

2020). The system was attached to a vertical translation stage (VAP10/M; Thorlabs) fixed to a

250x300x12.7mm aluminium breadboard (MB2530/M). To reduce external vibration noise, the

breadboard was placed on four sorbothane isolators (25.4mm x 27mm in diameter; AV4/M) and

rested on an antivibration table.

Vibration recording

To measure the velocity of the vibrations applied and measured, we used a Doppler laser

vibrometer (PDV-100, Polytec Ltd, Coventry, UK) set to 500mm/s maximum velocity and a Low

Pass Filter at 22kHz. The force applied by the shaker was simultaneously measured using the

miniature force sensor. The signals of both the laser vibrometer and the force sensor were

simultaneously acquired using a two-channel NI9250 Sound and Vibration module (NI Corporation

(UK) Ltd, Newbury, UK) and a USB powered data acquisition module (cDAQ-9171, NI). The

acquisition was done using custom-written software in LabView NXG 5.1 (NI). Samples were

acquired at a rate of 10,240 samples per second. Data was saved in TDMS format and

subsequently converted to text files using a custom program in LabView.

Signal processing

For each recorded vibration, we estimated the dominant frequency (Hz) and the RMS amplitude

for both the velocity (mm/s) measured in the distal anther, and the force (mN) measured in the

proximate anther. The text file containing the velocity and force measurements for each vibration

was processed in R ver. 4.0.5 (R Core Development Team 2021) using the package *seewave* (Sueur

et al. 2008). In brief, we first removed the offset of the signal and used the *timer* function

(threshold = 3, dmin = 1, window size = 30, no overlap) on the force channel to identify the

segment of the recording to be analysed (approximately 2 seconds or 20,480 samples per

channel). For each channel, we then applied a high pass filter at 20Hz (Hanning window, window

length = 520 samples) to remove background noise. The filtered signal was then used to obtain a power spectrum using the function *spec* (PSD = TRUE) with a Hanning window of 1,024 samples and a frequency range of 0-2,000Hz. The RMS amplitude was calculated on the same filtered signal.

Pollen collection

The pollen ejected through floral vibrations was collected on a plate made of black polyethylene measuring 13 cm x 5cm x 4mm (4083829; RS Components Ltd, Corby, UK) (see Ito et al. 2020). The plate had a hole drilled at 2cm from one of the short edges at the midline to allow the shaker wand to go through. The surface of the plate was painted with black acrylic paint (Black 3.0, Culture Hustle, Dorset, UK) (2 layers) to increase the contrast of pollen grains against the background. The polyethylene slide was positioned immediately below the flower with the aid of a micromanipulator (M330 with M3 tilting base and 2.5kg weight; World Precision Instruments, Hitchin, UK). After the vibration was applied, the ejected, light-coloured, pollen fell on the slide and provided a good contrast against the black background. The pollen was collected from the slide using a 2mm^3 cube of fuchsine-glycerol jelly kept at room temperature. The jelly with the collected pollen was then placed in a 1.5mL microcentrifuge tube and stored at 4°C until pollen counting.

Pollen counting

To estimate the number of pollen grains released after experimental buzz, we used a particle counter (Multisizer 4e Coulter Counter, Indianapolis, USA). Each pollen sample (contained in a 2mm^3 piece of fuchsin jelly in a vial) was suspended in 1 ml distilled water by heating it at 80°C for at least 30 min then shaking the vial vigorously with the help of an electric shaker until the vial content looked homogeneous. Prior to counting with the particle counter, the vial contents were added to 19 ml 0.9% NaCl solution, for a total volume of approximately 20 mL. For each sample, the amount of pollen was counted in four 1 ml subsamples. Only particles within the size range of 15-30μm were included in downstream analysis. Although the size of viable pollen grains within the studied species is much less variable and averaged around 24-25μm (Supplementary Figure 1), using a broader particle size range allowed us to also include inviable pollen grains, which were considerable smaller. In total, the particle analysis counted and measured 2,291,835 particles in the 15-30μm range (Supplementary Figure 1). We ran blank samples, containing only 0.9% NaCl solution at the beginning of every daily session and regularly between samples to ensure the equipment calibration accuracy. Finally, we summed the pollen counts for the four subsamples and multiplied by five to obtain each pollen release estimate.

Statistical analysis

To analyse the effect of anther treatment (free vs. joined) on vibration RMS amplitude we used

generalized mixed effects models using the package *lme4* (Bates et al. 2014) in *R* ver. *4.0.5* (R Core

Development Team 2021). We fitted separate models for RMS velocity of the distal anther and for

RMS force of the proximate anther. For each response variable, we started by fitting a full model

with anther fusion treatment (free vs. joined), plant species, the interaction between anther

treatment and plant species, sequential buzz order (first or second), and flower age (in days) as

fixed effects. Plant accession, individual, and flower identity were used as random effects to yield

the following model:

$$A_{RMS} \sim g + s + g \times s + n_{buzz} + a + (1|i_{accession}) + (1|i_{plant}) + (1|i_{flower}),$$

where A_{RMS} is vibration RMS amplitude (either velocity or force), g is anther fusion treatment, s is

plant species, n_{buzz} is sequential buzz order, a is flower age, and $i_{accession}$, i_{plant}, and i_{flower} are indices

corresponding to accession, individual and flower identity, respectively. For distal anther velocity,

the full model yielded a singular fit and thus we reduced model complexity by removing the plant

accession random effect ($i_{accession}$). This simplified model yielded qualitatively identical results to

the full model (results not shown) while avoiding the fit singularity. For proximate anther velocity,

the full model also yielded a singular fit, and thus we sequentially removed the random effects of

plant accession and plant identity ($i_{accession}$, i_{plant}). The simpler model also yielded qualitatively

identical results (coefficients and statistical significance) to the full model (results not shown). No

overdispersion of the residuals of the final models was detected with the statistical package

DHARMa (Hartig 2021).

To test for the effect of anther configuration on pollen release, we also fitted a generalized

mixed effects model using the number of pollen grains removed per buzz as the response variable.

The response variable was square root transformed to improve the distribution of the residuals.

The fitted model was:

$$\sqrt{r} \sim g + s + g \times s + n_{buzz} + a + (1|i_{accession}) + (1|i_{plant}) + (1|i_{flower}),$$

where r is number of pollen grains removed, and other variables are as previously defined. No

singular fit was detected in this case and the full model was used in the final analysis. No

overdispersion of the residuals was detected with the statistical package *DHARMa* (Hartig 2021).

Statistical significance of fixed effects (*P*-values) for all final models were obtained using Type III

analysis of variance and Satterthwaite's estimation of degrees of freedom implemented in the

package *lmerTest* (Kuznetsova et al. 2017). Given that we detected species *x* treatment

interactions (see Results), we calculated the estimated marginal means with treatment nested

within species using the fitted mixed-effects model in the *R* package *emmeans* (Lenth 2021).

Statistical significance of the linear contrast (free – joined) was obtained using a Kenward-Roger

approximation of degrees of freedom.

Results

We found a statistically significant effect of anther fusion treatment, which depended on plant

species, on both the velocity of the distal anther and on the force measured in the proximate

anther (treatment *x* species interaction; Table 2, Figure 4). For distal anther velocity, we found

that anther fusion increased the RMS velocity achieved by distal anthers compared to the free

anther treatment (Figure 4). The effect of treatment on RMS velocity was significant for all within

species comparisons (estimated marginal means contrasts, $P < 0.0001$) but the marginal means

contrast was 73-83% higher for *S. elaeagnifolium* (110.0 ± 8.93 mm/s) and *S. sisymbriifolium*

(116.4 ± 6.88mm/s) than for *S. pyracanthos* (63.3 ± 8.58 mm/s) (Figure 4). For both *S.*

elaeagnifolium and *S. sisymbriifolium*, the RMS velocity at the tip of the distal anther in the free

treatment was lower than the input RMS velocity measured in the shaker, while the anther

velocity in the joined treatment was higher than the input velocity for all species (Figure 5). For

the force measured in the proximate anther, we also found a species-dependent effect of

treatment (Table 2). All species experienced a higher force in the proximate anther in the joined

treatment (estimated marginal means contrasts, $P < 0.005$), although the magnitude of this

difference varied across species (marginal means difference = 2.43 ± 0.827 mN, 2.54 ± 0.637 mN,

6.93 ± 0.795 mN, for *S. elaeagnifolium, S. sisymbriifolium* and *S. pyracanthos*, respectively) (Figure

4). In *S. elaeagnifolium* and *S. pyracanthos*, the force measured in the proximate anther was

generally higher than the input force measured in the shaker before loading the flower, but in *S.*

sisymbriifolium RMS force in the proximate anther varied more from lower to higher compared to

the one measured before loading (Figure 5). Neither the order in which the buzzes were applied

and measured, nor flower age, had a significant effect on distal anther velocity or measured force

in proximate anthers (Table 2).

Floral vibrations removed on average 9,531 ± 679 pollen grains per buzz (mean ± se; range

55-59,665, n = 240). We found a significant effect of anther treatment on pollen release that

depended on plant species (species *x* treatment interaction, Table 3). For *S. elaeagnifolium* and *S.*

sisymbriifolium, more pollen (1,500 – 2,300 more pollen grains; square-root transformed marginal

means estimates = 48.2 ± 9.38 pollen$^{1/2}$ and 38.3 ± 7.18 pollen$^{1/2}$, respectively; $P < 0.0001$) was

released in the joined than in the free anther treatment (Figure 6). Conversely, the average

number of pollen grains released in *S. pyracanthos* was not statistically different between free and

joined treatments (estimated marginal means contrast 12.3 ± 8.91 pollen$^{1/2}$, *P* = 0.169; Figure 6).

The order in which the buzzes were applied and measured did not affect pollen release, but older

flowers released significantly more pollen grains (Table 3).

Discussion

In this study, we set out to address how different anther architectures, specifically the presence or

absence of joined anther cones in buzz-pollinated flowers, affect patterns of vibration

transmission and pollen release. Our results show that when applying vibrations to a focal

(proximate) anther, the vibration velocity experienced in other (distal) anthers is significantly

higher in flowers with joined anther cones, compared to those with free anthers. This result is

consistent with our hypothesis that joined anther cones enable more effective transmission of

applied vibrations from proximate to distal anthers via both the filament and the anther-anther

pathways. Importantly, this difference in vibration transmission across anthers translates to

functional differences in pollen release. In two species that naturally have loosely held, and

sometimes sprawling anther architectures (*S. elaeagnifolium* and *S. sisymbriifolium*), experimental

anther fusion results in more pollen released per buzz. In another species that naturally has non-

joined, but tightly held anthers forming a cone (*S. pyracanthos*), experimental anther fusion did

not increase pollen release. Anther fusion thus increases pollen release more strongly in species

with loosely held, sprawling anthers, than in those in which anthers form a non-joined but tightly

held cone. Our study suggests that anther architecture and the evolution of joined anther cones

might serve to influence the rate of pollen released across the flower following vibrations applied

to individual anthers.

Functional consequences of joined anther cones

The evolution of joined anther cones is likely to incur in different costs and benefits for flower

function depending on its interaction with floral visitors. Given that regardless of anther

architecture, both focal and non-focal anthers release pollen upon vibrations (e.g., Nevard et al.

2021, and the present study), joined anther cones might reduce pollen wastage by directing more

pollen onto the bee's body (Glover et al. 2004). We call this the **reduced pollen wastage**

hypothesis. Reducing the amount of pollen grains that misses the body of floral visitors should

increase pollination success. At the same time, this anther architecture may involve a trade-off if

bees can also benefit from joined anther cones by capturing a greater proportion of pollen that is

released upon vibrating. For example, the stereotypical C-shape posture that bees take during the

production of floral vibrations (De Luca and Vallejo-Marin 2013), should favour the receipt of

pollen grains in the ventral region, where bees are generally capable of grooming and harvesting

pollen grains (Vallejo-Marin et al. 2009; Huang et al. 2015; Koch et al. 2017; Tong et al. 2019).

Reduced pollen wastage could also be achieved in flowers with loosely held anthers if the visiting

bee can gather together all stamens using its legs and mandibles (M. Mayberry, D. McCart, J.

Burrow, T.L. Ashman, and A. Russell, *unpublished data*). Bees that are relatively large compared to

the flower they visit should be capable of such manipulation, although quantitative evidence of

this behaviour remains scarce. At the same time, the evolution of loosely held anthers might be

favored when buzzing bees can remove pollen from only one or a few anthers but only

infrequently contact the stigma, such as when the bees are relatively small compared to the

flower (Li et al. 2015; Solis-Montero and Vallejo-Marin 2017; Telles et al. 2020; Mesquita-Neto et

al. 2021). In this context, loosely held anthers that reduce vibrations being transmitted to non-

focal anthers (as shown in our study) would simultaneously reducing pollen wastage and pollen

theft. We call this the **reduced costs of pollen thieves hypothesis**. The hypotheses of reduced

pollen wastage and reduced costs of pollen thieves highlight how the potential benefits of fused

anther cones depend on ecological context; namely the presence and abundance of floral visitors

differing in size and behaviour. For example, in ecological communities dominated by large bee

pollinators capable of embracing all loosely held anthers and intercepting most of the ejected

pollen with their bodies (e.g. carpenter bees), the benefits of reducing pollen wastage to the plant

(when pollen grains miss the visitor's body) may be relatively minor. Conversely, the conditions for

the reduced cost of pollen thieves hypothesis may prevail in communities where smaller bees

acting as pollen thieves are dominant (Mesquita-Neto et al. 2018).

Additionally, the evolution of joined anther cones may increase the precision of pollen

placement on specific parts of the pollinator's body, which could increase the likelihood of stigma

contact (**unimodal pollen deposition hypothesis**). During buzz-pollination, anther cones are

expected to interact with the bee's ventral side depositing pollen in a single region (unimodal),

making the location of pollen placement and pick-up relatively predictable. At the same time,

unimodal pollen deposition could involve a trade-off, if the pollen were thus more readily

groomed into the bee's pollen baskets (Russell et al. 2021). Accordingly, the evolution of loosely

held anthers, could be favoured if pollen were thereby more frequently deposited int hard-to-

groom "safe sites" on the bee's body. The presence of loosely held anthers may also facilitate the

evolution of anther specialization within a flower, if different sets of loose anthers consistently

deposit pollen on different parts of the bee's body. This preliminary division of labour may

facilitate the evolution of anther morphology that improves the effectiveness of the division of labour, such as heteranthery, which commonly occurs in buzz pollinated plants (Vallejo-Marin et al. 2010; Barrett 2021). Division of labour in heterantherous species is achieved by two or more types of morphologically distinct stamens that release and deposit pollen on distinct parts of the pollinator's body, one of which is more effectively groomed by floral visitors (feeding anthers) and another that is more likely to contact the stigma of other conspecific flowers (pollinating anthers) (Luo et al. 2008; Vallejo-Marin et al. 2009; Papaj et al. 2017; Saab et al. 2021). In heterantherous flowers, pollen release and deposition are thus expected to have a bimodal (or multimodal if more than two stamen types) distribution on the pollinator's body, and such distribution is incompatible with joined anther cones.

Finally, a non-mutually exclusive hypothesis, for which our results provide direct empirical support, is that joined anther cones increase the transmission of vibrations across the stamens and, in some case, result in higher rates of pollen released per buzz (**increased pollen release hypothesis**). Our experiments across three *Solanum* species with slightly different anther architectures, show that in all cases, joined anther cones significantly increase the vibration amplitude (RMS velocity) transmitted to non-focal anthers showing an immediate consequence of joined anther cones. However, our experiments also show that the extent to which experimental fusion of anther cones influence pollen release is contingent on species-specific characteristics. Pollen release was increased in the two species that naturally have loosely arranged anther architectures (*S. elaeagnifolium* and *S. sisymbriifolium*), and in which the experimental treatment notably increases the extent to which anthers contact each other (the anther-anther vibration transmission pathway). An increase in pollen release through anther fusion is not observed in the Malagasy species *S. pyracanthos*, in which anthers are closely held together since the beginning of anthesis and throughout the flower's life. Although vibration amplitude in joined anthers of *S. pyracanthos* increases, we found no effect on pollen release. This might be because, despite being non-joined by either trichomes or bio-adhesives, the closely held androecium of *S. pyracanthos* is sufficient to transmit strong enough vibrations across all anthers to maximise pollen release rate. Pollen release rate in this case might be limited not by vibration amplitude, but by other factors such as the size of the apical pore from which pollen can come out during buzzing, and the amount of freely available pollen inside the anther locules, which in part is determined by flower age (Harder and Barclay 1994; Kemp and Vallejo-Marin 2021).

The four hypotheses mentioned above are not an exhaustive list or mutually exclusive, and they could act in concert, for example by simultaneously reducing pollen wastage and increasing

pollen release rates. Other consequences of free anthers could include lengthening the amount of

time that a visitor spends in a flower, due to having to separately manipulate anthers, which in

some cases might increase pollen deposition on the stigma, including self-pollen deposition.

However, a common feature of these hypotheses is their dependence on specific ecological

factors, particularly the morphological and behavioural characteristics of floral visitors. Evaluating

the relative value of these hypotheses in explaining the evolution of joined anther cones,

therefore requires explicitly considering the interaction of buzz pollinated plants and their wild

bee pollinators.

References

Armbruster, W. S., J. Lee, M. E. Edwards, and B. G. Baldwin. 2013. Floral paedomorphy leads to secondary specialization in pollination of Madagascar *Dalechampia* (Euphorbiaceae). Evolution 67:1196-1203.

Arroyo-Correa, B., C. E. Beattie, and M. Vallejo-Marin. 2019. Bee and floral traits affect the characteristics of the vibrations experienced by flowers during buzz-pollination. Journal of Experimental Biology 222:jeb198176.

Audacity. 2021. Audacity (ver. 3.0.2) https://www.audacityteam.org.

Barrett, S. C. 2021. Heteranthery. Current Biology 31:R774-R776.

Bates, D., M. Maechler, and B. Bolker. 2014. lme4: Linear mixed-effects models using Eigen and S4_. R package version 1.1-7. http://CRAN.R-project.org/package=lme4.

Brito, V. L. G., C. E. P. Nunes, C. R. Resende, F. Montealegre-Zapata, and M. Vallejo-Marín. 2020. Biomechanical properties of a buzz-pollinated flower. Royal Society Open Science 7:201010.

Buchmann, S. L. 1983. Buzz pollination in angiosperms. Pp. 73-113 *in* C. E. Jones, and R. J. Little, eds. Handbook of Experimental Pollination Biology. Scientific and Academic Editions, NY.

Buchmann, S. L. and J. H. Cane. 1989. Bees assess pollen returns while sonicating *Solanum* flowers. Oecologia 81:289-294.

Buchmann, S. L. and J. P. Hurley. 1978. Biophysical model for buzz pollination in Angiosperms. Journal of Theoretical Biology 72:639-657.

Buchmann, S. L., C. E. Jones, and L. J. Colin. 1977. Vibratile pollination of *Solanum douglasii* and *Solanum xanti* (Solanaceae) in Southern California USA. Wasmann Journal of Biology 35:1-25.

D'Arcy, W. G. 1992. Solanaceae of Madagascar: form and geography. Annals of the Missouri Botanical Garden:29-45.

Darwin, C. 1877. The different forms of flowers on plants of the same species. J. Murray, London.

De Luca, P. A., S. Buchmann, C. Galen, A. C. Mason, and M. Vallejo-Marin. 2019. Does body size predict the buzz-pollination frequencies used by bees? Ecology and Evolution 9:4875-4887.

De Luca, P. A., L. F. Bussiere, D. Souto-Vilaros, D. Goulson, A. C. Mason, and M. Vallejo-Marín. 2013. Variability in bumblebee pollination buzzes affects the quantity of pollen released from flowers. Oecologia 172:805-816.

De Luca, P. A. and M. Vallejo-Marin. 2013. What's the 'buzz' about? The ecology and evolutionary significance of buzz-pollination. Current Opinion in Plant Biology 16:429-435.

Dean, E., J. Poore, M. A. Anguiano-Constante, M. H. Nee, H. Kang, T. Starbuck, A. Rodrígues, and M. Conner. 2020. The genus *Lycianthes* (Solanaceae, Capsiceae) in Mexico and Guatemala. PhytoKeys 168:1.

Endress, P. K. 1996a. Diversity and evolutionary trends in angiosperm anthers. Pp. 92-110 *in* W. G. D'Arcy, and R. C. Keating, eds. The Anther: Form, Function and Phylogeny. Cambridge University Press, Cambridge.

Endress, P. K. 1996b. Homoplasy in Angiosperm flowers. Pp. 303-325 *in* S. J. Sanderson, and L. Hufford, eds. Homoplasy - The Recurrence of Similarity in Evolution. San Diego Academic Press, San Diego.

Faegri, K. 1986. The solanoid flower. Pp. 51-59. Transactions of the Botanical Society of Edinburgh. Taylor & Francis.

Falcão, B. F., C. Schlindwein, and J. R. Stehmann. 2016. Pollen release mechanisms and androecium structure in *Solanum* (Solanaceae): Does anther morphology predict pollination strategy? Flora 224:211-217.

Glover, B. J., S. Bunnewell, and C. Martin. 2004. Convergent evolution within the genus *Solanum*: the specialised anther cone develops through alternative pathways. Gene 331:1-7.

Hansen, M., G. C. Lanes, V. L. Brito, and E. D. Leonel. 2021. Investigation of pollen release by poricidal anthers using mathematical billiards. Physical Review E 104:034409.

Harder, L. D. and M. R. Barclay. 1994. The functional significance of poricidal anthers and buzz pollination: controlled pollen removal from *Dodecatheon*. Functional Ecology 8:509-517.

Hartig, F. 2021. DHARMa: Residual diagnostics for hierarchical (multi-level / mixed) regression models.

Huang, Z.-H., H.-L. Liu, and S.-Q. Huang. 2015. Interspecific pollen transfer between two coflowering species was minimized by bumblebee fidelity and differential pollen placement on the bumblebee body. J Plant Ecol 8:109-115.

Ito, S., H. Rajabi, and S. N. Gorb. 2020. A ballistic pollen dispersal strategy hidden in stylar oscillation. bioRxiv 217:109 - 114.

Jankauski, M. A. 2020. Measuring the frequency response of the honeybee thorax. Bioinspiration & Biomimetics 15:046002.

Kemp, J. E. and M. Vallejo-Marin. 2021. Pollen dispensing schedules in buzz-pollinated plants: Experimental comparison of species with contrasting floral morphologies. American Journal of Botany 108:1-13.

Knapp, S. 2002. Floral diversity and evolution in the Solanaceae. Pp. 267-297 *in* Q. C. B. Cronk, R. M. Bateman, and J. A. Hawkins, eds. Developmental Genetics and Plant Evolution. Taylor & Francis, London.

Knapp, S. 2014a. Solanaceae Source: *Solanum elaeagnifolium* Cav. http://solanaceaesource.org/taxonomy/term/106820/descriptions.

Knapp, S. 2014b. Solanaceae Source: *Solanum pyracanthos* Lam. http://solanaceaesource.org/content/solanum-pyracanthos

Koch, L., K. Lunau, and P. Wester. 2017. To be on the safe site - Ungroomed spots on the bee's body and their importance for pollination. PLOS One 12.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest package: Tests in linear mixed effect models. Journal of Statistical Software 82:1-26.

Lenth, R. V. 2021. emmeans: Estimated marginal means, aka least-squares means.

Li, J.-K., Y.-P. Song, H. Xu, Y.-W. Zhang, J.-Y. Zhu, and L.-L. Tang. 2015. High ratio of illegitimate visitation by small bees severely weakens the potential function of heteranthery. Journal of Plant Ecology 8:213-223.

Luo, Z., D. Zhang, and S. S. Renner. 2008. Why two kinds of stamens in buzz-pollinated flowers?
Experimental support for Darwin's division-of-labour hypothesis. Functional Ecology
22:794-800.
Macior, L. W. 1968. Pollination adaptation in *Pedicularis groenlandica*. American Journal of Botany
55:927-932.
Mesquita-Neto, J. N., N. Blüthgen, and C. Schlindwein. 2018. Flowers with poricidal anthers and
their complex interaction networks—Disentangling legitimate pollinators and illegitimate
visitors. Functional Ecology 32:2321-2332.
Mesquita-Neto, J. N., A. L. C. Vieira, and C. Schlindwein. 2021. Minimum size threshold of visiting
bees of a buzz-pollinated plant species: consequences for pollination efficiency. American
Journal of Botany.
Nevard, L., A. L. Russell, K. Foord, and M. Vallejo-Marin. 2021. Transmission of bee-like vibrations
in buzz-pollinated plants with different stamen architectures. Scientific Reports 11:13541.
Papaj, D. R., S. L. Buchmann, and A. L. Russell. 2017. Division of labor of anthers in heterantherous
plants: flexibility of bee pollen collection behavior may serve to keep plants honest.
Arthropod-Plant Interactions 11:307-315.
Petanidou, T., M. V. Price, J. L. Bronstein, A. Kantsa, T. Tscheulin, R. Kariyat, N. Krigas, M. C.
Mescher, C. M. De Moraes, and N. M. Waser. 2018. Pollination and reproduction of an
invasive plant inside and outside its ancestral range. Acta Oecologica 89:11-20.
Plebani, M., O. Imanizabayo, D. M. Hansen, and W. S. Armbruster. 2015. Pollination ecology and
circadian patterns of inflorescence opening of the Madagascan climber *Dalechampia aff.*
bernieri (Euphorbiaceae). Journal of Tropical Ecology 31:99-101.
Ramos, A. F., S. G. Bauermann, L. A. Lopes, and A. C. P. Evaldt. 2016. Produção polínica de *Solanum*
sisymbriifolium Lam. (Solanaceae) e a coleta de pólen pelas abelhas visitantes florais.
Educaçao Ambiental em Açao:1-7.
Robyns, W. 1931. L'Organisation florale des solanacées zygomorphes, par le Dr. Walter Robyns. M.
Lamertin.
Rosi-Denadai, C. A., P. C. S. Araujo, L. A. O. Campos, L. Cosme, Jr., and R. N. C. Guedes. 2020. Buzz-
pollination in Neotropical bees: genus-dependent frequencies and lack of optimal
frequency for pollen release. Insect Sci 27:133-142.
Russell, A. L., A. M. Fetters, E. I. James, and T.-L. Ashman. 2021. Pollinator effectiveness is affected
by intraindividual behavioral variation. Oecologia:1-12.
Russell, A. L., R. E. Golden, A. S. Leonard, and D. R. Papaj. 2016. Bees learn preferences for plant
species that offer only pollen as a reward. Behavioral Ecology 27:731-740.
Saab, G. d. S., V. d. F. Mansano, A. Nogueira, I. C. Maia, P. J. Bergamo, and J. V. Paulino. 2021. A
sophisticated case of division of labour in the trimorphic stamens of the *Cassia fistula*
(Leguminosae) flower. AoB Plants 13:plab054.
Särkinen, T., L. Bohs, R. G. Olmstead, and S. Knapp. 2013. A phylogenetic framework for
evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. Bmc
Evolutionary Biology 13.
Solis-Montero, L. and M. Vallejo-Marin. 2017. Does the morphological fit between flowers and
pollinators affect pollen deposition? An experimental test in a buzz-pollinated species with
anther dimorphism. Ecology and Evolution 7:2706-2715.
Sueur, J., T. Aubin, and C. Simonis. 2008. Seewave, a free modular tool for sound analysis and
synthesis. Bioacoustics 18:213-226.
Team, R. D. C. 2021. R. A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL http://www.R-project.org.

Telles, F. J., C. L. Klunk, F. R. d. Maia, V. L. G. de Brito, and I. G. Varassin. 2020. Towards a new
understanding of the division of labour in heterantherous flowers: the case of *Pterolepis*
glomerata (Melastomataceae). Biological Journal of the Linnean Society 131:1-11.
Thorp, R. W. and J. R. Estes. 1975. Intrafloral behavior of bees on flowers of *Cassia fasciculata*.
Journal of the Kansas Entomological Scoiety 48:175-184.
Tong, Z.-Y., X.-P. Wang, L.-Y. Wu, and S.-Q. Huang. 2019. Nectar supplementation changes
pollinator behaviour and pollination mode in *Pedicularis dichotoma*: implications for
evolutionary transitions. Annals of Botany 123:373-380.
Tscheulin, T., T. Petanidou, S. G. Potts, and J. Settele. 2009. The impact of *Solanum elaeagnifolium*,
an invasive plant in the Mediterranean, on the flower visitation and seed set of the native
co-flowering species *Glaucium flavum*. Plant Ecology 205:77-85.
Vallejo-Marín, M. 2019. Buzz pollination: studying bee vibrations on flowers. New Phytologist
224:1068-1074.
Vallejo-Marin, M., E. M. Da Silva, R. D. Sargent, and S. C. H. Barrett. 2010. Trait correlates and
functional significance of heteranthery in flowering plants. New Phytologist 188:418-425.
Vallejo-Marin, M., J. S. Manson, J. D. Thomson, and S. C. H. Barrett. 2009. Division of labour within
flowers: heteranthery, a floral strategy to reconcile contrasting pollen fates. Journal of
Evolutionary Biology 22:828-839.
Vogel, S. 1996. Christian Konrad Sprengel's theory of the flower: the cradle of floral ecology. Pp.
44-62 *in* D. G. Lloyd, and S. C. H. Barrett, eds. Floral Biology. Chapman & Hall, NY.
Vorontsova, M. S. and S. Knapp. 2014. Solanaceae Source: *Solanum sisymbriifolium.*
http://solanaceaesource.org/content/solanum-sisymbriifolium.
593

595 **Table 1.** Species distribution, flower and anther characteristics, floral visitors and accessions of the three *Solanum* species used in the present study.

596 Accessions 1179F and 1180M were sourced commercially (see Methods) and their exact origin is unknown.

Species	Native range	Corolla diameter (anther length x width) mm	Anther size	Floral visitors	Accession	Population	Lat.	Lon.	References
S. elaeagnifolium	Americas	20-35 (6-10 x 1.5-2)		*Amegilla, Bombus, Centris, Exomalopsis, Melissodes, Nomia, Protandrena, Pseudapis, Ptiloglossa, Xylocopa*					(Knapp 2014a) (Buchmann and Cane 1989; Tscheulin et al. 2009; Petanidou et al. 2018)
					10-QSJ-12	Querétaro, México	20.37	99.99	
					11s36	Puebla, México	18.46	97.40	
S. pyracanthos	Madagascar	23-30 (6-7 x 1.5-2)		Unknown, but likely to include Malagasy *Nomia* and *Xylocopa*					(Armbruster et al. 2013; Knapp 2014b; Plebani et al. 2015)
					1179F	Unknown	--	--	
S. sisymbriifolium	South America	20-35 (9-10 x		*Parapsaenythia sp., Augochlora spp.,*					(Vorontsova and Knapp 2014; Ramos

1.5-2)	*Augochloropsis spp., Dialictus sp., Xylocopa sp.*				et al. 2016)
		F1-sys-3	Salta, Argentina	-24.71 -65.41	
		1180M	Unknown	-- --	

598 **Table 2.** Statistical analysis of the effect of anther fusion treatment (free vs. joined), buzz order (first or second), and flower age on the RMS

599 amplitude velocity measured in the distal anther, and RMS amplitude force experienced by the proximate anther in three *Solanum* species. Model

600 estimates were obtained from a linear mixed-effects model with individual and/or flower identity as random effects (see Methods). Statistical

601 significance (*P*-values) of the fixed effects was calculated with a Type III analysis of variance. Significant effects (*P*-value < 0.05) are shown in bold. The

602 reference level used for coefficient estimation is *S. pyracanthos,* free stamen configuration.

	Velocity (mm/s; distal anther)				Force (mN; proximate anther)		
Variable	**Model coefficient**	**Estimate**	**SE**	***P*-value**	**Estimate**	**SE**	***P*-value**
	Intercept	63.456	12.122		117.397	1.522	
Treatment				**<0.001**			**<0.001**
	Joined	63.273	8.582		6.925	0.795	
Species				0.073			**<0.001**
	S. elaeagnifolium	-48.609	11.553		-1.014	1.371	
	S. sisymbriifolium	-40.225	10.297		-7.234	1.263	
Species × Treatment				**<0.001**			**<0.001**
	S. elaeagnifolium × joined	46.774	12.406		-4.490	1.149	
	S. sisymbriifolium × joined	53.146	10.969		-4.384	1.016	
Buzz order	Order	2.862	4.630	0.537	-0.188	0.431	0.663

Flower age	Days		-3.739	5.077	0.463	0.581	0.823	0.481

Table 3. Statistical analysis of the effect of anther fusion treatment (free vs. joined), buzz order (first or second), and flower age on number of pollen grains released after single buzzes in flowers from three *Solanum* species. Model estimates were obtained from a linear mixed-effects model with accession, individual and flower identity as random effects using a square-root transformation of pollen grains removed. Statistical significance (*P*-values) of the fixed effects was calculated with a Type III analysis of variance. Significant effects (*P*-value < 0.05) are shown in bold. The reference level used for coefficient estimation is *S. pyracanthos,* free stamen configuration.

Variable	Model coefficient	Estimate	SE	*P*-value
	Intercept	76.152	20.960	
Treatment				**<0.001**
	Joined	-12.332	8.915	
Species				0.609
	S. elaeagnifolium	-35.432	23.895	
	S. sisymbriifolium	-47.562	22.725	
Species × Treatment				**<0.001**
	S. elaeagnifolium × joined	60.545	12.966	
	S. sisymbriifolium × joined	50.645	11.422	
Buzz order	Order	-4.269	4.842	0.379
Flower age	Days	18.777	6.165	**0.003**

Figure 1. Flower profiles of the three studied species of *Solanum* before (left-hand side panels) and after (right-hand side panels) applying the experimental treatment joining the anthers in a joined cone. The left hand-side panels show the flowers in their natural form and orientation. Notice the natural variation in the level of contact between the individual anthers across the three species, with *S. sisymbriifolium* and *S. elaeagnifolium* having relatively free anthers, while the anthers of *S. pyracanthos* are naturally arranged in a connivent cone. A, B: *S. sisymbriifolium;* C, D: *S. elaeagnifolium*; E, F: *S. pyracanthos*. Scale bar = 1cm.

Figure 2. Diagram of the experimental set up (for details see Methods). Diagram not to scale.

Figure 3. (**A**) Close up of the experimental set up showing the attachment of the shaker system via an entomological pin to the proximate anther (anther 1 in B) and the laser beam of the Doppler vibrometer targeting one of the distal anthers (either anther 3 or 4 in B). **(B)** Floral diagram of *Solanum* showing the labelled anthers (1-5). Floral diagram modified from (Robyns 1931; Knapp 2002).

Figure 4. Vibration properties measured in anthers of three species of *Solanum* with experimentally joined anther cones ("Joined"; golden symbols) and without ("Free"; grey symbols). The "Joined" anther cones are created by gluing anthers together with water-based polyvinyl acetate glue (PVA). The bottom row shows the vibration force (root mean square, RMS force in mN) measured in the anther to which the mechanical vibrations were applied (proximate anther). The top row shows the vibration velocity (RMS velocity in mm/s) in a different anther in the opposite side of the cone (distal anther). As a reference, the experiment-wide median RMS values of the vibration velocity (53.85 mm/s) and force (116.05 mN) of the mechanical shaker measured both before the flower was attached to the system, and after it was removed, are shown with a dashed line. Dark circles represent observed data, with random noise added in the x-axis to facilitate visualisation. The white symbols inside the violin plots indicate the mean and 95% confidence interval of the mean calculated using 1,000 bootstrap replicates. Asterisks indicate statistical significance of within-species comparisons between treatments (marginal means contrasts): *** = $P < 0.001$; ** = $P < 0.005$.

Figure 5. Relationship between RMS force measured in the anther to which mechanical vibrations are applied (proximate anther, x-axis) and RMS velocity of an anther located in the opposite side of the anther cone (distal anther, y-axis). RMS values are shown relative to the average daily values of RMS force and velocity measured during calibration of the mechanical shaker system. Symbol shape and colour indicate the experimental treatment: "Joined" is shown in golden circles, and "Free" in grey squares. Within each category, each symbol indicates a different flower. All flower received both treatments in random order.

Figure 6. Effect of experimental manipulation of the anther cone on the number of pollen grains removed during a single buzz in three species of buzz-pollinated *Solanum* species. Dark circles represent observed data, with random noise added in the x-axis to facilitate visualisation. The white symbols inside the violin plots indicate the mean and 95% confidence interval of the mean calculated using 1,000 bootstrap replicates. Comparison of the effect of treatment (marginal means contrast): NS = $P > 0.05$, *** = $P < 0.001$.

Supplementary Material

Supplementary Figure 1. Frequency distribution of the size of measured particles (pollen grains) in the electric particle counter as described in the Methods. All samples for each species are presented in the same histogram. Top = *Solanum elaeagnifolium.* Centre = *S. pyracanthos.* Bottom = *S. sisymbriifolium.*

A — Distal anther, Proximate anther, Shaker pin

B

Supplementary Figure 1. Frequency distribution of the size of measured particles (pollen grains) in the electric particle counter as described in the Methods. All samples for each species are presented in the same histogram. Top = *Solanum elaeagnifolium.* Centre = *S. pyracanthos.* Bottom = *S. sisymbriifolium*