1 **Sperm of more colorful males are better adapted to ovarian fluids in lake char**
2 **(Salmonidae)**
3

4 David Nusbaumer, Laura Garaud, Christian de Guttry, Laurie Ançay & Claus Wedekind[*]
5
6 Department of Ecology & Evolution, University of Lausanne, Lausanne, Switzerland
7 [*]correspondence: Claus Wedekind, claus.wedekind@unil.ch, orcid.org/0000-0001-6143-4716

8
Author contributions
DN and CW designed the experiment. DN and LG organized and conducted the field work
(with the help of CdG) and determined sperm characteristics. DN performed the two
experiments, determined male colors, and genotyped the offspring for parental assignments.
CdG did the ddRADseq library preparation and genomic analysis. LA stained the blood
smears and classified the blood cells. DN & CW did the statistics and wrote the manuscript
that was revised by all authors.
16
Abstract

Fish often spawn eggs with ovarian fluids that have been hypothesized to support sperm of
some males over others (cryptic female choice). Alternatively, sperm reactions to ovarian
fluids could reveal male strategies linked to their likely roles during spawning. Sperm of
males who would usually be close to females during spawning are then expected to be better
adapted to the presence of ovarian fluids than to water only, while the reverse would be
expected for males that typically spawn at larger distance to the females. We tested these
predictions with gametes and ovarian fluids from wild-caught lake char (*Salvelinus umbla*).
We found that sperm of more colorful males showed increased sperm velocity in diluted
ovarian fluids while sperm of paler males were fastest in water only. We then let equal
numbers of sperm compete for fertilizations in the presence or absence of ovarian fluids and
used microsatellite markers to assign in total 1,464 embryos (from 70 experimental trials) to
their fathers. Overall, sperm of more colorful males reached higher fertilization success than
sperm of pale males. This difference was enhanced by the presence of ovarian fluids and best
explained by the increased sperm velocity. Sperm competitiveness was not enhanced with
decreasing genetic distance to a given female, although parallel stress tests on embryos had
revealed that females would profit more from mating with least related males rather than most
colored ones. We conclude that sperm of more colorful males are best adapted to ovarian
fluids, and that the observed reaction norms reveal male strategies rather than cryptic female
choice.

37
38 **Key words:** sperm competition, loaded raffle, sperm velocity, male reproductive strategy,
39 cryptic female choice, sperm reaction norms
40

Introduction

Ejaculate economics predicts that males invest strategically into sperm according to the expected microecology they will experience after ejaculation (Parker & Pizzari 2010). This microecology is strongly defined by competitors and the level of sperm competition. Males are predicted to increase sperm number and/or sperm velocity with increasing risk of sperm competition (Parker & Pizzari 2010). This prediction is meanwhile well supported in many taxa (Magris 2021). Another aspect of the microecology that sperm have to deal with is the biochemical environment that females create around their eggs with their reproductive fluids. The functional significance of such female reproductive fluids is not well understood yet (Gasparini *et al.* 2020). In externally spawning fishes, the fluids with which females expel their eggs are called "ovarian fluids" and consist of various inorganic and organic compounds (Lahnsteiner *et al.* 1995).

Salmonids are externally spawning fish whose ovarian fluids vary in amount and composition among species and even within populations (Lahnsteiner *et al.* 1995; Zadmajid *et al.* 2019). Viscosity of undiluted ovarian fluids is about 2 to 3 times higher than that of water (Zadmajid *et al.* 2019), but ovarian fluids will quickly be diluted when expelled with eggs into water. During spawning, freshly spawned eggs and ovarian fluids are swirled around by male and female movements so that some eggs may no more be surrounded by significant amounts of ovarian fluids at the time of fertilizations while others still are. A likely predictor of how much sperm will be exposed to ovarian fluids is the position of the male during spawning. Arguably the closer a male is with its urogenital opening to a female's vent during egg release, the more likely its sperm will be exposed to ovarian fluids.

Ovarian fluids have repeatedly been found to support sperm velocity, motility, and longevity (Zadmajid *et al.* 2019), i.e. key targets of sexual selection (Evans *et al.* 2013). They have also been discussed as allowing for certain forms of cryptic female choice (Firman *et al.* 2017). Their release with eggs can, for example, specifically promote paternity of dominant males over parasitic spawners ("sneaker males") (Alonzo *et al.* 2016; Egeland *et al.* 2016; Makiguchi *et al.* 2016; Lehnert *et al.* 2017). The support of sperm function can be even more specific: Ovarian fluids of guppies (*Poecilia reticulata*) were found to slow down sperm of full-sibs as compared to unrelated males, potentially to reduce inbreeding (Gasparini & Pilastro 2011). The presence of ovarian fluids during spawning has also been found to reduce inbreeding and hence offspring mortality by differentially supporting some sperm over others in Chinook salmon (*Oncorhynchus tshawytscha*) (Rosengrave *et al.* 2016; Gessner *et al.* 2017; Lehnert *et al.* 2017), while ovarian fluids of lake trout (*Salvelinus namaycush*) were found to promote sperm velocity of more related males (Butts *et al.* 2012), potentially to reduce outbreeding depression in this salmonid. Indeed, Yeates et al. (2013) found that ovarian fluids specifically support sperm of the same *Salmo* species over sperm of another *Salmo* species, i.e. they promote reproductive isolation by a species-specific chemoattraction. All these examples have mainly been discussed in the light of adaptive female reproductive strategies. However, many of the observed effects of ovarian fluids on sperm function could reveal variation in male reproductive strategies, as predicted by the theory of ejaculate economics and experimentally demonstrated in other vertebrates (Jeannerat *et al.*).

Salmonid males can play various roles during spawning (Esteve 2005), and many of these roles have been found to affect milt and sperm characteristics (Magris 2021). In migratory populations, for example, there is usually a fraction of males that avoid the costs of migration but then remain small and subdominant to their migratory rivals at spawning. They therefore reproduce mainly surreptitiously and invest disproportionally into sperm number and sperm velocity in order to increase the competitiveness of their sperm (Young *et al.* 2013). Evidence for strategic male investments has also been found within non-migratory populations. Male dominance at the spawning place is usually size dependent (Esteve 2005),

and older males and larger male whitefish (*Coregonus zugensis*) have indeed been found to invest less into gonad weight and to have slower sperm than young and small males who typically spawn further away from the female's vent (Rudolfsen *et al.* 2008). However, the links between male size and sperm traits can vary among populations (Perroud *et al.* 2021) and be plastic in response to the perceived social environment (Bartlett *et al.* 2017). When Rudolfsen et al. (2006), for example, controlled for size effects by confining pairs of size-matched male Arctic char (*Salvelinus alpinus*) in an enclosure and let them develop a dominance hierarchy, sperm characteristics changed quickly: as predicted from ejaculate theory, dominance led to reduced sperm velocity. Egeland et al. (2016) then used an analogous setup to show that sperm of dominant males are better designed to swim in ovarian fluids. One possible explanation for this increased performance is that ovarian fluids somehow recognize sperm of dominant males and promote them specifically (cryptic female choice). Alternatively, dominant males can expect to be close to the females during gamete release and may therefore produce sperm that do well when exposed to ovarian fluids. If so, any other male trait that is likely to affect the distance between male and female during spawning would be predicted to affect sperm reaction to ovarian fluids. Here we use wild-caught lake char (*Salvelinus umbla*) from Lake Geneva (Switzerland) to test whether male sexual ornamentation is such a trait.

The lake char is a non-migratory salmonid endemic to Alpine lakes. It spawns in winter in lek-like mating system where male competition is usually intense. This species is closely related to the Arctic char (Kottelat & Freyhof 2007) that has often been used to study effects of ovarian fluid on sperm motility (e.g.Turner & Montgomerie 2002; Urbach *et al.* 2005; Egeland *et al.* 2016). Both species develop spawning colorations. Arctic char often develop strong red colorations in both sexes (Janhunen *et al.* 2011). Male lake char of our study population are mostly yellow during spawning season while females hardly develop any spawning colorations.

In other fish families, sexual ornaments are often positively linked to aspects of sperm performance (Kortet *et al.* 2004; Mehlis *et al.* 2013; Fukuda & Karino 2014; Kekäläinen *et al.* 2014), probably because both types of traits can react to induced stress (Kekäläinen *et al.* 2014), suggesting that high quality sperm in well ornamented males is an indicator of health and vigor rather than of alternative mating strategies. The pattern in salmonids seems less clear. Janhunen et al. (2009) found in a captive population of Arctic char that the intensity of male coloration was positively correlated with sperm velocity (when tested in water) while dominant males are expected to have reduced sperm velocity (Rudolfsen *et al.* 2006). Combined, these findings suggest that male dominance and coloration are not necessarily linked in char. Nevertheless, if females prefer colorful males, we predict that not only dominant but also colorful males are usually closer to females during spawning than other types of males. If so, sperm of colorful males are predicted to do better when exposed to ovarian fluids than sperm of less colorful males.

Here we test whether (i) male coloration of lake char predicts sperm performance as it does in other fish, (ii) sperm of colorful males react better to presence of ovarian fluids that sperm of less colorful males, as predicted if colorful males are preferred and/or more dominant and therefore able to spawn closer to females, (iii) colorful males reach disproportional high fertilization success in sperm competition, (iv) the presence of ovarian fluids change the outcome of sperm competition because of its effects in sperm performance, and (v) ovarian fluids of different females differ in their effects on sperm. We also test whether (vi) coloration is an indicator of male inbreeding coefficient, and if not, whether (vii) male inbreeding coefficient or (viii) their genetic relatedness to females affect sperm characteristics and the outcome of sperm competition in water or in diluted ovarian fluids.

Methods

Sampling and handling of fish, gametes, and ovarian fluids

Wild char were caught in December 2017 and 2018 from three spawning sites in Lake Geneva (Ripaille, 46.3957N-6.4718E; Locum, 46.4049N-6.7588E; Meillerie, 46.4113N-6.7216E) by local fishermen using gill nets (48 mm mesh size, set over night) and transported to a research facility (INRA, Thonon-les-Bains, France) were they were kept in a 1,000 L circular tank with free-flowing lake water before further processing. In 2017, 10 males and 4 females were haphazardly chosen for the experiment. Males were processed on the first day after catching and females on the second day. Each fish was anaesthetized in 10 L water containing 3 mL of Eugenol (clove oil in ethanol at a 1:9 ratio), photographed in a custom-made photobox under standardized conditions (17 mm, f/5.0, 1/200 s, ISO 400, WB 4000 K, JPG 24 Mpx), and striped to collect gametes. Tissue samples were taken from the anal fin and stored in 70% ethanol at -20°C for further analysis. In 2018, further 24 males were sampled and handled using the same procedures.

Milt was directly striped into 50 mL conical tubes (Falcon, BD Biosciences, Allschwil, Switzerland), carefully avoiding contamination by urine, feces, or water. Between 2 and 5 mL of milt were obtained per male. These samples were stored at 4°C (< 2 hours) until further use. When all males had been stripped, 1.5 mL of each milt sample was diluted 1:9 in Storfish (IMV Technologies, France; an inactivating medium) in 15 mL tubes (Falcon, BD Biosciences, Allschwil, Switzerland). These tubes were then placed on ice and are referred to as 'diluted milt'.

Eggs and ovarian fluids were stripped into individual plastic containers. About 20 mL of ovarian fluid each was separated from the eggs with a syringe and stored in 50 mL tubes at 4°C. Eggs were kept (<1h) in the dark at 4°C in about 20 mL Ovafish (IMV Technologies, France) to prevent drying.

Color measurement

Custom-made macros in Fiji were used to analyze male skin colorations. First, the white balance of each image was standardized with the help of the white and black values of the color scale. Skin gray value was then determined as the overall body gray value in the RGB color space (range from 0 = black, to 255 = white; using mean of all pixels in each of the RGB channels). Fish redness and yellowness were measured from 3 squares (about 1% of the total body area each) in the pectoral region, the ventral region and the anal region, as described by Parolini et al. (2018) (Fig. S1). Redness and yellowness were measured as the *a** and *b** components of the CIE-*L*a*b** color space, respectively (the a* axis for greenness-redness and the b* axis for blueness-yellowness both range from -100 to 100). All measures of yellowness were significantly correlated (Pearson's $r > 0.92$, p always < 0.001). Similarly, all measures of redness were significantly correlated (Pearson's $r > 0.75$, $p < 0.013$). Therefore, the means of the three measurements per fish were used for each color. Gray values were correlated with yellowness (the yellower the lighter) and with redness (the redder the darker; Fig. S1) and were therefore ignored in further analyses.

Sperm characteristics of freshly caught fish

Sperm characteristics were measured on the day of sampling with Qualisperm® (AKYmed, Switzerland). Briefly, 20 µL of the diluted milt was activated in a microtube with 980 µL ice-cooled standardized water (OECD 1992) and vortexed for 5s. Then, 2 µL of this activated solution was pipetted into the well of a 4-well chamber slide (Leja, AKYmed, Switzerland), on a cooling station set at 4.5°C (the expected water temperature at the spawning ground). Sperm concentration, sperm velocity (average path velocity), and percentage of immotile sperm (sperm motility) were recorded under a phase contrast microscope at 20X

magnification 20 s after activation in 2017 and 25 s after activation in 2018. Sperm longevity
was recorded with a stopwatch from the time of activation until no progressive sperm motion
could be observed in the frame of capture. Two replicated measures were taken for each
sperm sample to test for consistency, and means of these measurements were used for
statistics. The concentration of active spermatozoa was calculated for each male by
multiplying the total sperm concentration with the motility (e.g. 2,000 Mio/mL times 80%
motility = 1,600 Mio/mL active spermatozoa).
198
Hematological analyses
Hematological analyses are frequently used to determine stress in fish (Seibel *et al.* 2021).
High leukocyte counts and especially also relative lymphocytosis (percentage of lymphocytes
among leukocytes) have been associated with acute infections, e.g. in eel (*Anguilla anguilla*)
infected with the virulent nematode *Anguillicola crassus* (Haenen *et al.* 2010), or have been
found in response to exposure to pesticides (Oluah *et al.* 2020), and even in anticipation of
predation risk (Meuthen *et al.* 2020). Thrombocytosis (elevated platelet counts) is another
possible indicator of an acute infection (Rose *et al.* 2012; Yan *et al.* 2013).
In order to determine these potential stress indicators, a blood sample (2-3 mL) was
drawn from the caudal vasculature of males with a 3 mL syringe mounted with a 23 G needle
heparinized with heparin 1 IU/µL. The blood was emptied in 2 mL microtubes, centrifuged 10
min at 10,000 g and 4°C. The plasma was then carefully collected, disposed in 1.5 mL
microtubes and frozen in the vapor of liquid nitrogen for transportation and stored at -80°C in
the lab for further analysis. A drop a blood from the syringe was used to make two blood
smears per individual, on microscope frosted slides. Blood smears were dried in the field and
fixed for 3 minutes in absolute ethanol back in the lab. Slides were stained for 20 minutes in
Giemsa stain and rinsed twice in PBS. From each slide, 10 photographs (i.e. 20 per
individual) were taken on an EVOS XL Core inverted light transmission microscope
(Thermofisher, Switzerland) at 40x magnification. Counts of thrombocytes, lymphocytes, and
granulocytes were done manually by a naïve observer (Fig. S2). Counts of total number of
cells were done automatically with a custom-made macro in Fiji (Fig. S2).
220
Experiment 1: Sperm characteristics in response to ovarian fluids
For the 2017 sample, the same protocol was used on the day of experimental fertilizations, i.e.
one day after catching the fish from the wild, to determine each male's sperm characteristics
in each female's ovarian fluid (40 combinations of milt samples and ovarian fluids) and in
water (controls, 4 times replicated each; Table S1). To do so, 10 µL of the diluted milt were
activated either with 490 µL standardized water or with 490 µL ovarian fluid solution. The
concentration of ovarian fluid in this solution was 50% as this would reflect a possible
concentration of ovarian fluid in the water when sperm swim towards the egg (Rosengrave *et*
al. 2016).
230
Experiment 2: Competitive fertilization trials and monitoring of embryos
The 10 males were haphazardly assigned to 5 dyads (pairs of competitors) whose sperm then
competed for fertilization of 24 eggs/female, either activated in water only or in water with
ovarian fluid. Each possible combination of dyad x female was tested in both environments
twice, resulting in 80 sperm competition trials (5 dyads x 4 females x 2 treatments x 2
replicates; Table S2) that involved in total 1,920 eggs. However, due to an accident during
handling, 10 of these 80 experimental cells were lost (all of the same female, prepared to be
exposed to sperm activated in water only, see Table S2), reducing the number of eggs that
could be monitored to N = 1,680.

In preparation of these sperm competition trials, the eggs of each female were first
washed twice with 200 mL Ovafish® (IMV Technologies, France) to remove ovarian fluids.
Then, 20 batches of 24 eggs each were placed in wells of 6-well plates (Falcon, BD
Biosciences, Allschwil, Switzerland). Ten mL of diluted milt of 2 males each was prepared
such that each male was represented with the same concentration of active sperm (25
Mio/mL) in the mix. One mL of each mix was then used to fertilize the 2 batches of 24
eggs/female. This 1 mL of mix was activated in a separated tube with either 4 mL of
standardized water or 4 mL of standardized water with ovarian fluid (ratio ovarian fluid to
water = 1:2) and vortexed for 5s. The solution was poured in a well with eggs. Two minutes
later, i.e. after fertilization could be expected to have happened, standardized water was added
to fill the wells (16.8 mL) and the eggs were left undisturbed in a dark environment for 2
hours to allow hardening.
Each batch of 24 supposedly fertilized eggs was transferred to a 50 mL tube filled with
standardized water and transported on ice to a climate chamber where they were rinsed for
30s under running tap water (4 L/min) in a sterilized tea strainer. The eggs were then
distributed singly to wells of 24-well plates (Falcon, BD Biosciences, Allschwil, Switzerland)
filled with 1.8 mL of autoclaved standardized water (OECD 1992). Embryos were raised at
4.5°C in a 12h:12h light-dark cycle. Fertilization success was assessed at the neurula stage
(14 days post fertilization). No visible embryo at this stage was interpreted as non-fertilized
egg. Embryo mortality, the timing of hatching, hatchling size, and hatchling growth was
monitored from then on in the course of two parallel studies, one that combines these data
with further stress experiments (by a pathogen and a chemical pollutant) on embryos
produced in a regular full-factorial breeding design to test for parental effects on embryo
performance (Garaud *et al.* 2022), and another one on sex-specific stress tolerance
(Nusbaumer *et al.* 2021). Embryos and larvae that died during the observational period were
transferred to individual microtubes and stored in 96% ethanol at -20°C for further analysis.
Larvae were euthanized 14 days after hatching with a spike of 100 µL of Koi Med® Sleep
4.85% (Koi & Bonsai Zimmermann, Germany) in the well before being transferred to
individual microtubes.
269
Genotyping and paternity assignment based on microsatellite markers
In the case of larvae and adults, DNA was extracted from up to 25 mg tissue with an
extraction robot and the DNAeasy Blood & Tissue kit (Qiagen, Hilden, Germany) following
manufacturer's instructions. Unhatched embryos were homogenized and DNA was extracted
using the Qiamp Fast DNA Stool mini kit protocol (Qiagen, Germany) as in Wilkins et al.
(2015). A first multiplex PCR of 3 polymorphic microsatellite markers (Savary *et al.* 2017)
could be used to assign most embryos and larvae to their fathers (Bernatchez & Duchesne
2000). A second multiplex PCR of 3 further microsatellite markers (Savary *et al.* 2017) was
used for to assign the remaining offspring (Table S3). In the first multiplex PCR, the *sdY* sex
marker (Yano *et al.* 2013) was included to sex offspring (Table S3) for a parallel study on
sex-specific stress tolerance (Nusbaumer *et al.* 2021). Multiplex PCR were run on a Biometra
thermocycler in 20 µL reaction volume using HotStarTaq DNA Polymerase (Qiagen,
Germany) reagents (i.e. 2 µL PCR buffer 10X, 4 µL QSolution 5X, 0.4 µL HotStar Taq DNA
Polymerase), 0.4 µL of dNTPs 10 mM and 0.4 µL of each primer). PCR cycles were run as
follow: initial heat activation of 15 min at 95 °C; 35 cycles of 30 s denaturation at 94°C, 90 s
annealing at 57°C and 60 s extension at 72°C; followed by 30 min of final extension at 72°C
and 10 min at 4°C. DNA extracted from unhatched embryo was amplified using the same
protocol, but in 10 µL reaction volume and with 40 cycles. PCR products were diluted 2X and
run on an ABI3100 sequencer (Applied Biosystems, USA). Genotypes were read using
Genemapper v4.0 (Applied Biosystems, USA). Genotyping of 3 dead embryos was not

successful possibly due to poor DNA quality, bringing the final number of genotyped
offspring to N = 1,475.
Adults were genotyped twice (no discrepancies observed). Paternity was assigned
using CERVUS v3.0 (Kalinowski *et al.* 2007) as recommended for categorical assignments in
controlled experiments (Jones *et al.* 2010). Input simulation parameters were set with 6 loci,
10,000 offspring, 2 candidate fathers with a proportion of sampling of 1 and a proportion of
typing of 1, a minimum of 2 typed loci and an allelic dropout/mistyping rate of 0.05 as
recommended by Wang (2004), and an assignment confidence of 95%.
298
Double-digested RAD sequencing and SNPs calling
DNA from all males and females of experiment 2 was used to also determine male multi-
locus heterozygosity (MLH) and the genomic similarity ("kinship") between males and
females of all possible combinations (N = 40). DNA concentration was measured using Qubit
2.0 (Thermo Fisher Scientific, Waltham**,** Massachusetts) while its integrity was verified on a
1% agarose gel. DNA was subsequently standardized to 20 ng/μl. Two sequencing libraries
were produced using all 10 males and 4 females each following the Brelsford et al. (2016)
protocol adapted from Parchman et al. (2012). Briefly, DNA was digested using the enzymes
EcoRI-HF and MspI (New England Biolab, Ipswich, Massachusetts, United States) and a
unique EcoRI barcode was ligated to each individual. After library purification, PCR
amplification were performed and fragments size-selected in between 400-550bp. Libraries
were then single-end genotyped on 2 lanes of Illumina HiSeq 2500 with fragments of 125 bp
length at Lausanne Genomic Technologies Facility (University of Lausanne).
After the quality control on the resulting fastq files done with FASTQC v0.11.7
(Andrews 2010), reads were trimmed to 110bp given the low per-base quality of the
sequences (<20 Phred score). The resulting RAD data were analysed using Stacks 1.48
(Catchen *et al.* 2013) using the default parameters unless otherwise specified. Individuals
sequences were demultiplexed using *process_radtags* (Stacks 1.48) and reads mapped to the
Salvelinus spp reference genome (Christensen *et al.* 2018) using BWA (Li & Durbin 2010).
Pstacks (Stacks 1.48) was done selecting a minimum stack depth of 5 (-*m* 5) and using the
bounded SNP model. The catalogue of loci was created using *Cstacks* (Stacks 1.48) allowing
2 mismatches (-*n* 2) between loci. *Populations* (Stacks 1.48) was used to generate the VCF
file considering only loci present in 100% of the individuals and markers heterozygosity of
0.5 (--*max_obs_het* 0.50). None of the individuals was excluded based on genotyping rate.
Further filtering was done using *vcftools*, v0.1.15 (Danecek *et al.* 2011). To reduce incorrect
heterozygosity call and remove paralogs, loci were filtered for a minimum coverage of 10X (-
-*min-meanDP* 10) and a maximum coverage of 50X *(--max-meanDP* 50; two times the mean
coverage). Only loci under Hardy-Weinberg equilibrium were retained with a P-value
treshold of 0.05 (--*hwe* 0.05). Finally, all the loci that were not shared between all individuals
were discarded (--*max-missing* 1). No filtering was made based on minor allele frequencies. A
total of 4,150 SNPs was retained after filtering with a mean coverage per individual of 29X.
Individual MLH, i.e. the number of heterozygous loci on the genotyped portion of the
genome, was calculated using *vcftools* (--*het*) (Danecek *et al.* 2011). The kinship, i.e. the
expected inbreeding coefficient for the progeny of one family, was estimated using the
beta.dosage function of the package *Hierfstat* (Goudet 2005). This function generates a
genomic matrix based on allele dosage to estimate kinship coefficients considering the
proportion of alleles shared between individuals (Goudet *et al.* 2018). The inbreeding
coefficients for females and males, F_{beta}, were obtained by extracting the diagonal of the
genomic matrix obtained with *beta.dosage*. Fig. S3 shows how MLH (range in males: 0.242
to 0.259) relates to F_{beta} (range in males: 0.029 to 0.092). Further analyses were focused on
F_{beta}.

340
Statistical analysis
Analyses were done in RStudio (RStudioTeam 2015) and the package *lme4* (Bates *et al.*
2015) and in JMP® 15.2.1. Pearson correlation coefficients (r) were used to test for
correlations when visual inspection of the plots suggested that the conditions for parametric
statistics were not significantly violated, else Spearman rank order correlations (r_s) were used.
Generalized linear models (GLM) and linear mixed models (LMM) were run on categorical
or continuous dependent variables, respectively. GLM were based on a binomial error
distribution. Models were reduced in stepwise regression with backward term selection in the
built-in function *drop1*, and model comparison was done with likelihood-ratio test (LRT).
The non-violation of the model assumptions was checked visually using fitted values versus
residuals plot and residuals QQ-plots.

   When testing for effects of ovarian fluids on sperm characteristics and fertilization
success in the sperm competition trials, directed tests were used based on the *a priori*
predictions that if ovarian fluids have an effect on sperm, the are expected to support sperm of
more colorful and/or less inbred and/or less related males (the corresponding p-values are
notated as p_{dir}). Directed tests were used because such tests avoid inflation of the alpha value
while still allowing to discover effects in the non-expected direction if the effect sizes are
strong (Rice & Gaines 1994).

Results
Male phenotypes, leukocyte counts, MLH, and milt characteristics
Males looked silvery with variable intensity of ventral coloration. This coloration was striking
on some individuals while largely absent on others. Colorful males were mostly yellow (Fig.
1), in few cases also slightly reddish (but 19 of 34 males had negative *a** values, and the
mean (±95% CI) *a** value for the males was 0.95±1.78). Redness was positively correlated
with yellowness (r = 0.45, p < 0.005) and therefore ignored in further analyses.

   Yellowness increased with male size (Fig. 1A) but was not significantly correlated
with F_{beta} (Fig. 1B). Yellower males had less concentrated milt (LRT: *p* = 0.03; Table S4A).
When their milt was activated in water only, yellower males had faster sperm (LRT: *p* = 0.05;
Table S4B; Fig. 1C) and fewer immobile sperm than paler males (LRT: *p* = 0.05, Table S4C;
Fig. 1D). Sperm longevity was not significantly correlated to skin coloration (LMM: p = 0.17,
Table S4D; Fig. 1E). Including sampling year in our models explained 0% of the variance in
sperm concentration but 5.1% and 15.8% of the variance in velocity and rate of immobile
sperm, respectively. None of the sperm traits could be predicted by male size (Table S4) nor
F_{beta} except that males with higher F_{beta} had higher counts of immobile sperm Fig. S4).

   Some pale males had high leukocyte counts, but the correlation between leukocyte
counts and skin coloration was not significant (Fig. 1F). However, skin coloration turned out
to be a good predictor of relative lymphocytosis (Fig. 1G) and of thrombocytosis that both
declined with increasing skin yellowness (Fig. 1H). No immune parameter was significantly
correlated to male F_{beta} (Fig. S4).

Experiment 1: Sperm performance in ovarian fluid
The effects of activating sperm with water only or water with ovarian fluid can be analyzed
on two levels, namely ignoring potential effects of male and female identity or including
them. When ignoring male and female identities, activating sperm with or without ovarian
fluids did not seem to affect average sperm velocity (LRT: χ^2 = 1.01, *p* = 0.29, Fig. 2A).
However, activating with ovarian fluids significantly reduced the variance in sperm velocity
(Fig. 2A). Moreover, sperm activated with ovarian fluid were generally more motile (LRT: χ^2

$= 5.22$, $p = 0.022$, Fig. 2B) and showed higher longevity (LRT: $\chi^2 = 124.83$, $p < 0.001$, Fig.
2C).
When including potential effects of male and female identity, they both turned out to
play significant roles. Males significantly varied in all sperm traits that were monitored here
(see main effects of male identity in Table 1). Females significantly varied in how their
ovarian fluids affected average sperm velocity and longevity but not the percentage of
immobile sperm (see main effects of female identity in Table 1). Similar patterns could be
observed with regard to how the sperm of different males responded to the ovarian fluids of
different females (the male $\times$ female interaction terms in Table 1; Fig. S6). Again, average
sperm velocity and longevity but not the percentage of immobile sperm depended on the
combination of male and female identities (Table 1).
Sperm responses to ovarian fluids could partly be predicted by male skin coloration.
With increasing yellowness, sperm velocity increased more in the presence of ovarian fluids
(Fig. 2D). The reduction of immobile sperm that ovarian fluids caused could not be predicted
by male yellowness (Fig. 2E), but the increase in sperm longevity in response to ovarian
fluids was less pronounced in more yellow males than in pale ones (Fig. 2F).
The 40 different combinations of male identity and donor or ovarian fluid allowed to
test whether the genomic distance between males and female (kinship) affected sperm
performance in ovarian fluid. Kinship seemed to have no effects on sperm velocity or
percentage of immobile sperm (Fig. 2G,H). However, kinship between male and female
increased the longevity of sperm when exposed to ovarian fluid (Fig. 2I).
410
Experiment 2: Males success in sperm competition
We genotyped 1,475 offspring of which 1,464 could be assigned to their father (99.3%
success). In the following, three types of focal males are considered: the most yellow, the
least inbred, and the least related to the respective female, because male yellowness did not
correlate with their inbreeding coefficients (Fig. 1B; p = 0.65) nor with kinship (Fig. S4A; p =
0.42), and kinship could not be predicted by male inbreeding coefficients (Fig. S4B; p =
0.47).
Figure 3 shows the competitive success of different types of focal males when tested
in water only or with ovarian fluids (including only the females for which we had both types
of conditions; see Table S5 for the identities of the focal males in the different scenarios).
Overall, the success rates of focal males were 53.6%, 56.2%, and 53.2% if the focal male was
the most yellow, the least inbred, or the least related to the female, respectively (Fig. 3). Dyad
identity always had a strong effect on the focal male's success rate (Table 2). The presence of
ovarian fluid significantly enhanced the focal male's success if it is the yellower of the two
competitors (Table 2A, Fig. 3A) while female identity or the difference in sperm velocity
between the rivals played no significant roles in this scenario (Table 2A). Similar patterns
could be observed if the focal male was the one with the lower F_{beta}, except that the presence
of ovarian fluid did not increase the focal male's fertilization success (Table 2B, Fig. 3B). No
effects of ovarian fluids could be observed if the focal male was the least related to the egg
donor (Table 2C, Fig. 3C). In the latter scenario, the female identity and the difference in
sperm velocity now explained male success (Table 2C).
432
Discussion
The first question of this study was whether sperm traits of wild-caught lake char can be
predicted from male phenotypes (body size and skin coloration) or overall genetic quality
(here determined as F_{beta}) when tested in water only, independently of any potential effects of
ovarian fluids. We found no effects of body size, but skin coloration was a good predictor of
sperm traits: yellower males had faster sperm and lower rates of immobile sperm (i.e. higher

sperm motility) than paler males. The skin coloration of lake char is hence among the
secondary sexual traits that have been found to be positively linked to sperm competitiveness
in different fish families (e.g. Locatello *et al.* 2006; Janhunen *et al.* 2009; Mehlis *et al.* 2013;
Fukuda & Karino 2014). Our findings also suggest that paleness in our sample was linked to
comparatively low health and vigor rather than to a "sneaker" life-history strategy (parasitic
male spawners) that are common in many migratory salmonids, because sneakers are
predicted to invest more into sperm than dominant males (Ball & Parker 2000).
Secondary sexual traits like conspicuous skin colors are predicted to reveal general
health and vigor (Andersson 1994). This has repeatedly been observed in other species
(Milinski & Bakker 1990; Johnson & Fuller 2015). In our study, paler males suffered from
increased relative lymphocytosis and thrombocytosis, i.e. they had increased rates of
lymphocytes and of thrombocytes, two potential indicators of acute infections or other
physiological stress (Haenen *et al.* 2010; Rose *et al.* 2012; Yan *et al.* 2013), than yellower
males. Male yellowness was hence a useful predictor of current male health condition,
potentially revealing, for example, virulence of a tapeworm infection (Johansen *et al.* 2019).
Theory also predicts a link between F_{beta} and general health and vigor (Allendorf &
Luikard 2007), and such links have indeed been found in other studies (Fox & Reed 2011).
We found that the rates of immobile sperm declined with declining male F_{beta}, supporting the
hypothesis that sperm motility can be an indicator of general health and vigor (Kowalski &
Cejko 2019). We did not find any significant correlations between male F_{beta} and immune
parameters, but this may need to be studied again in larger samples that provide more
statistical power.
Despite the similar expectations from theory, F_{beta} and skin colorations turned out to
be not significantly correlated in our sample. In previous studies, such a correlation was
sometimes found (e.g. Zajitschek & Brooks 2010; Herdegen *et al.* 2014) and sometimes not
(e.g. Frommen *et al.* 2008; Marsh *et al.* 2017), including a study on Arctic char (Janhunen *et
al.* 2011). One possibility is that skin coloration is a plastic trait that quickly reacts to acute
stress (Johnson & Fuller 2015) while F_{beta} affects more the overall tolerance to environmental
stress. If so, the link between F_{beta} and skin colorations is expected to change with changing
environmental conditions.
The next major questions of this study were whether and how ovarian fluids affect
sperm traits, and whether such effect differ for different types of males. We found that
ovarian fluids affected all sperm traits that we studied: They reduced the between-male
variance in sperm velocity, reduced the percentage of immobile sperm, and nearly doubled
mean sperm longevity. Sperm velocity was not generally increased by ovarian fluids,
apparently contrary to some previous findings in other fishes (Zadmajid *et al.* 2019).
However, when comparing the effects of ovarian fluids on sperm characteristics of different
types of males, it turned out that the yellower a male, the more does the velocity of its sperm
increase when in contact with ovarian fluids, while sperm of paler males are rather slowed
down in ovarian fluids. Such a correlation between the effects of ovarian fluids and male
coloration suggests that sperm of yellower males are better prepared to ovarian fluids while
sperm of paler males are better prepared to the absence of ovarian fluids. If yellower males
are closer to the female vent during spawning than paler males, these reaction of sperm to the
different environments correspond well to the expected microenvironment that the sperm are
likely to be exposed to.
The effects of ovarian fluids on sperm motility was not correlated to male yellowness,
i.e. all male types profited similarity from the presence of ovarian fluids. Interestingly,
however, sperm longevity was more enhanced by ovarian fluids in paler males than in
yellower males, suggesting that there is a trade-off between increased velocity and increased

longevity. Sperm of yellower males profited from increased velocity while sperm of paler
males profited from increased longevity when in contact with ovarian fluids.
Females varied in how much their ovarian fluids generally increased sperm velocity
and sperm longevity, but not in how much their ovarian fluids reduced sperm motility.
Females also varied in how much their ovarian fluids specifically increased velocity and
longevity of sperm of certain males over others, i.e. we found the corresponding females x
male interactions to be significant (but again not in the case of sperm motility). When we
tested whether some of these interactions can be explained by the genetic distance between
males and females, we found no link between relatedness and sperm velocity or motility, and
a non-expected positive link between relatedness and sperm longevity. Ovarian fluids seemed
to specifically elongate the longevity of sperm of more closely related males. This non-
expected finding needs to be further studied. We can currently not offer an adaptive
explanation for why females should support the longevity of sperm of related males more
than that of sperm of unrelated males. Preferring unrelated males would reduce the average
offspring inbreeding coefficients.
Our next major question was: Whose sperm benefit from exposure or non-exposure to
ovarian fluids when in sperm competition? Here we focused on three types of males: the
yellower ones of the experimental dyads, the less inbred ones, and the ones who were least
related to a given female. We found that the presence of ovarian fluids led to enhanced
fertilization success of the more yellow male but did not specifically affect the sperm of less
inbred or less related males. However, competition trials can only reveal a limited range of
possible situations. In our case, the overall sperm to egg ratios were so high that variation in
sperm longevity may not have affected the outcome of the sperm competition, while late
fertilizations may often be relevant under natural conditions.
In a parallel study, Garaud et al. (2022) used an extended sample of our study
population to test whether females would profit from mating with more yellow, less inbred, or
less related males. They used a full-factorial breeding design (crossing 10 male and 6 females
in all possible combinations) and single rearing of large numbers of offspring to test the
embryos' reaction to the pathogen *Aeromonas salmonicida*, to an ecologically relevant
concentration of a common chemical pollutant (a spike of 1 ng/L ethinylestradiol into the
water), and to no-stress conditions (sham controls). They found that while exposure to
ethinylestradiol showed no effects, exposure to the pathogen reduced embryo development
rate. Contrary to expectations from 'good-genes' hypotheses of sexual selection, this
pathogen-related reduction in embryo development rate increased with increasing male
yellowness, i.e. offspring of yellower males were on average smaller and less tolerant to the
pathogen stress than offspring of paler males. These results confirm previous findings on
Arctic char (Janhunen *et al.* 2011) and suggest that females would not profit from giving
yellower males a selective advantage during sperm competition. However, Garaud et al.
(2022) found strong effects of the relatedness between males and females on offspring
performance. Embryo development rates declined with increasing parental relatedness and
hence increasing mean inbreeding coefficients among the offspring. Similar effects of
relatedness could be observed in the embryos and larvae that resulted from experiment 2 of
the present study and that are also presented in detail in Garaud et al. (2022), while male
yellowness or male F_{beta} were both not significantly linked to the growth of these embryos
(Garaud *et al.* 2022).
In conclusion, male yellowness revealed aspects of current health and vigor and was
generally linked to high sperm velocity and motility. If exposed to diluted ovarian fluids,
sperm velocity of yellower males became even faster and sperm of paler males slowed further
down. In sperm competition experiments, fertilization success of the yellower of the two
competitors was indeed increased when exposed to diluted ovarian fluids. Exposure to

ovarian fluids also generally increased sperm motility and longevity. While motility was similarity increased in sperm differently colored males, longevity was more enhanced in sperm of paler males than in sperm of yellower males, suggesting a trade-off between increased sperm velocity and increased sperm longevity. Male F_{beta} and male genetic relatedness to a given female were poor predictors of sperm performance in water or ovarian fluids. When offspring of the different types of males were tested under various types of stress conditions, offspring of yellower males grew slower and were less stress tolerant than offspring of paler males, while offspring of least related parents showed the best performance in the various stress conditions. Combined, these results suggest that the increased performance of sperm of yellower males in ovarian fluids is an adaptation to the microenvironment that the sperm are likely to be exposed to, rather than cryptic female choice by ovarian fluids.

Ethics

This work complied with the national, cantonal and university regulations where it was carried out. The handling and transport of adults and transport of embryos was approved by the French authorities (INTRA.FR.2017.0109258).

Data accessibility

Data will be deposited on the Dryad depository upon acceptance of the manuscript.

Funding

This project was supported by the Swiss National Science Foundation (31003A_159579 and 31003A_182265).

Acknowledgments

We thank G. Levray from the Pistolet fishery for catching the fish, the APERA association and the *pisciculture des Rives* for access to the fish, L. Adhia Eya, A. Atherton, C. Berney, J. Bideaux, F. Dolivo, L. Espinat, S. Kreuter, E. Longange, L. Marques da Cunha, and N. Sironi for assistance in the field or in the lab, F. Schütz for discussion on statistics, and E. Lasne for discussion and access to the INRA facilities.

References

Allendorf FW, Luikard G (2007) *Conservation and the genetics of populations.* Oxford University Press, Malden, MA, USA.

Alonzo SH, Stiver KA, Marsh-Rollo SE (2016) Ovarian fluid allows directional cryptic female choice despite external fertilization. *Nature Communications* **7**, 12452. 10.1038/ncomms12452

Andersson MB (1994) *Sexual selection.* Princeton University Press, Princeton, New Jersey.

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data, p. http://www.bioinformatics.babraham.ac.uk/projects/fastqc. Babraham Bioinformatics, Cambridge, UK.

Ball MA, Parker GA (2000) Sperm competition games: A comparison of loaded raffle models
and their biological implications. *Journal of Theoretical Biology* **206**, 487-506.

Bartlett MJ, Steeves TE, Gemmell NJ, Rosengrave PC (2017) Sperm competition risk drives
rapid ejaculate adjustments mediated by seminal fluid. *Elife* **6**, e28811.
10.7554/eLife.28811

Bates D, Mächler M, Bolker BM, Walker SC (2015) Fitting linear mixed-effects models
using lme4. *Journal of Statistical Software* **67**, 1-48. 10.18637/jss.v067.i01

Bernatchez L, Duchesne P (2000) Individual-based genotype analysis in studies of parentage
and population assignment: how many loci, how many alleles? *Canadian Journal of*
Fisheries and Aquatic Sciences **57**, 1-12.

Brelsford A, Dufresnes C, Perrin N (2016) High-density sex-specific linkage maps of a
European tree frog (*Hyla arborea*) identify the sex chromosome without information
on offspring sex. *Heredity* **116**, 177-181. 10.1038/hdy.2015.83

Butts IAE, Johnson K, Wilson CC, Pitcher TE (2012) Ovarian fluid enhances sperm velocity
based on relatedness in lake trout, *Salvelinus namaycush*. *Theriogenology* **78**, 2105-
2109. 10.1016/j.theriogenology.2012.06.031

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool
set for population genomics. *Molecular Ecology* **22**, 3124-3140. 10.1111/mec.12354

Christensen KA, Rondeau EB, Minkley DR*, et al.* (2018) The Arctic charr (Salvelinus
alpinus) genome and transcriptome assembly. *PLoS ONE* **13**, e0204076.
10.1371/journal.pone.0204076

Danecek P, Auton A, Abecasis G*, et al.* (2011) The variant call format and VCFtools.
Bioinformatics **27**, 2156-2158. 10.1093/bioinformatics/btr330

Egeland TB, Rudolfsen G, Nordeide JT, Folstad I (2016) Status specific tailoring of sperm
behavior in an external fertilizer. *Frontiers in Ecology and Evolution* **4**, 135-143.
10.3389/fevo.2016.00135

Esteve M (2005) Observations of spawning behaviour in salmoninae: *Salmo*, *Oncorhynchus*

and *Salvelinus*. *Reviews in Fish Biology and Fisheries* **15**, 1-21. 10.1007/s11160-005-

7434-7

Evans JP, Rosengrave P, Gasparini C, Gemmell NJ (2013) Delineating the roles of males and

females in sperm competition. *Proceedings of the Royal Society B-Biological Sciences*

280, 20132047-20132047.

Firman RC, Gasparini C, Manier MK, Pizzari T (2017) Postmating female control: 20 years

of cryptic female choice. *Trends in Ecology & Evolution* **32**, 368-382.

10.1016/j.tree.2017.02.010

Fox CW, Reed DH (2011) Inbreeding depression increases with environmental stress: an

experimental study and meta-analysis. *Evolution* **65**, 246-258. 10.1111/j.1558-

5646.2010.01108.x

Frommen JG, Luz C, Mazzi D, Bakker TCM (2008) Inbreeding depression affects

fertilization success and survival but not breeding coloration in threespine

sticklebacks. *Behaviour* **145**, 425-441. 10.1163/156853908792451458

Fukuda S, Karino K (2014) Male red coloration, female mate preference, and sperm longevity

in the cyprinid fish Puntius titteya. *Environmental Biology of Fishes* **97**, 1197-1205.

10.1007/s10641-013-0207-6

Garaud L, Nusbaumer D, de Guttry C*, et al.* (2022) Genomic compatibility excels possible

good genes effects of sexual selection in lake char. *bioRxiv*.

https://doi.org/10.1101/2022.03.03.482834

Gasparini C, Pilastro A (2011) Cryptic female preference for genetically unrelated males is

mediated by ovarian fluid in the guppy. *Proceedings of the Royal Society B-Biological*

Sciences **278**, 2495-2501. 10.1098/rspb.2010.2369

Gasparini C, Pilastro A, Evans JP (2020) The role of female reproductive fluid in sperm

competition. *Philosophical Transactions of the Royal Society B-Biological Sciences*

375, 20200077. 10.1098/rstb.2020.0077

Gessner C, Johnson SL, Fisher P*, et al.* (2017) Male-female relatedness at specific SNP-
linkage groups influences cryptic female choice in Chinook salmon (*Oncorhynchus*
tshawytscha). *Proceedings of the Royal Society B-Biological Sciences* **284**.
10.1098/rspb.2017.0853

Goudet J (2005) HIERFSTAT, a package for R to compute and test hierarchical F-statistics.
Molecular Ecology Notes **5**, 184-186. 10.1111/j.1471-8286.2004.00828.x

Goudet J, Kay T, Weir BS (2018) How to estimate kinship. *Molecular Ecology* **27**, 4121-
4135. 10.1111/mec.14833

Haenen OLM, Lehmann J, Engelsma MY*, et al.* (2010) The health status of European silver
eels, *Anguilla anguilla*, in the Dutch River Rhine Watershed and Lake IJsselmeer.
Aquaculture **309**, 15-24. 10.1016/j.aquaculture.2010.08.026

Herdegen M, Dudka K, Radwan J (2014) Heterozygosity and orange coloration are associated
in the guppy (*Poecilia reticulata*). *Journal of Evolutionary Biology* **27**, 220-225.
10.1111/jeb.12290

Janhunen M, Rudolfsen G, Kekäläinen J*, et al.* (2009) Spawning coloration and sperm quality
in a large lake population of Arctic charr (Salmonidae: *Salvelinus alpinus* L.).
Biological Journal of the Linnean Society **98**, 794-802. DOI 10.1111/j.1095-
8312.2009.01317.x

Janhunen M, Peuhkuri N, Primmer CR, Kolari I, Piironen J (2011) Does breeding
ornamentation signal genetic quality in Arctic charr, *Salvelinus alpinus*? *Evolutionary*
Biology **38**, 68-78. 10.1007/s11692-010-9100-9

Jeannerat E, Marti E, Berney C*, et al.* (2018) Stallion semen quality depends on major
histocompatibility complex matching to teaser mare. *Molecular Ecology* **27**, 1025–
1035. 10.1111/mec.14490

Johansen IB, Henriksen EH, Shaw JC*, et al.* (2019) Contrasting associations between
breeding coloration and parasitism of male Arctic charr relate to parasite species and
life cycle stage. *Scientific Reports* **9**, 1-9. 10.1038/s41598-019-47083-x

Johnson AM, Fuller RC (2015) The meaning of melanin, carotenoid, and pterin pigments in
the bluefin killifish, *Lucania goodei*. *Behavioral Ecology* **26**, 158-167.
10.1093/beheco/aru164

Jones AG, Small CM, Paczolt KA, Ratterman NL (2010) A practical guide to methods of
parentage analysis. *Molecular Ecology Resources* **10**, 6-30. 10.1111/j.1755-
0998.2009.02778.x

Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program
CERVUS accommodates genotyping error increases success in paternity assignment.
Molecular Ecology **16**, 1099-1106. 10.1111/j.1365-294X.2007.03089.x

Kekäläinen J, Pirhonen J, Taskinen J (2014) Do highly ornamented and less parasitized males
have high quality sperm? - an experimental test for parasite-induced reproductive
trade-offs in European minnow (*Phoxinus phoxinus*). *Ecology and Evolution* **4**, 4237-
4246. 10.1002/ece3.1267

Kortet R, Vainikka A, Rantala MJ, Taskinen J (2004) Sperm quality, secondary sexual
characters and parasitism in roach (*Rutilus rutilus* L.). *Biological Journal of the*
Linnean Society **81**, 111-117.

Kottelat M, Freyhof J (2007) *Handbook of European freshwater fishes.* Kottelat and Freyhof,
Cornol, Switzerland and Berlin, Germany.

Kowalski RK, Cejko BI (2019) Sperm quality in fish: Determinants and affecting factors.
Theriogenology **135**, 94-108. 10.1016/j.theriogenology.2019.06.009

Lahnsteiner F, Weismann T, Patzner RA (1995) Composition of the ovarian fluid in 4
salmonid species: *Oncorhynchus mykiss*, *Salmo trutta* f *lacustris*, *Saivelinus alpinus*
and *Hucho hucho*. *Reproduction Nutrition Development* **35**, 465-474.

Lehnert SJ, Butts IAE, Flannery EW, *et al.* (2017) Effects of ovarian fluid and genetic
differences on sperm performance and fertilization success of alternative reproductive
tactics in Chinook salmon. *Journal of Evolutionary Biology* **30**, 1236-1245.
10.1111/jeb.13088

Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler

689   transform. *Bioinformatics* **26**, 589-595. 10.1093/bioinformatics/btp698

690   Locatello L, Rasotto MB, Evans JP, Pilastro A (2006) Colourful male guppies produce faster

691   and more viable sperm. *Journal of Evolutionary Biology* **19**, 1595-1602.

692   Magris M (2021) Strategic adjustment of ejaculate quality in response to variation of the

693   socio-sexual environment. *Behavioral Ecology and Sociobiology* **75**. 10.1007/s00265-

694   021-03032-1

695   Makiguchi Y, Torao M, Kojima T, Pitcher TE (2016) Reproductive investment patterns and

696   comparison of sperm quality in the presence and absence of ovarian fluid in

697   alternative reproductive tactics of masu salmon, *Oncorhynchus masou*.

698   *Theriogenology* **86**, 2189-2193.e2182. 10.1016/j.theriogenology.2016.07.009

699   Marsh JN, Vega-Trejo R, Jennions MD, Head ML (2017) Why does inbreeding reduce male

700   paternity? Effects on sexually selected traits. *Evolution* **71**, 2728-2737.

701   10.1111/evo.13339

702   Mehlis M, Hilke LK, Bakker TCM (2013) Attractive males have faster sperm in three-spined

703   sticklebacks Gasterosteus aculeatus. *Current Zoology* **59**, 761-768.

704   Meuthen D, Meuthen I, Bakker TCM, Thunken T (2020) Anticipatory plastic response of the

705   cellular immune system in the face of future injury: chronic high perceived predation

706   risk induces lymphocytosis in a cichlid fish. *Oecologia* **194**, 597-607.

707   10.1007/s00442-020-04781-y

708   Milinski M, Bakker TCM (1990) Female sticklebacks use male coloration in mate choice and

709   hence avoid parasitized males. *Nature* **344**, 330-333.

710   Nusbaumer D, Garaud L, Ançay L, Wedekind C (2021) Sex-specific stress tolerance in

711   embryos of lake char (*Salvelinus umbla*). *Frontiers in Ecology and Evolution* **9**,

712   768263. 10.3389/fevo.2021.768263

713   OECD (1992) *OECD Guideline for the testing of chemicals 203 (fish, acute toxicity test)*.

714   OECD Publishing, Paris, France.

Oluah NS, Aguzie IO, Ekechukwu NE*, et al.* (2020) Hematological and immunological
responses in the African catfish *Clarias gairepinus* exposed to sublethal
concentrations of herbicide Ronstar (R). *Ecotoxicology and Environmental Safety* **201**.
10.1016/j.ecoenv.2020.110824

Parchman TL, Gompert Z, Mudge J*, et al.* (2012) Genome-wide association genetics of an
adaptive trait in lodgepole pine. *Molecular Ecology* **21**, 2991-3005. 10.1111/j.1365-
294X.2012.05513.x

Parker GA, Pizzari T (2010) Sperm competition and ejaculate economics. *Biological Reviews*
85, 897-934. 10.1111/j.1469-185X.2010.00140.x

Parolini M, Iacobuzio R, Possenti CD*, et al.* (2018) Carotenoid-based skin coloration signals
antioxidant defenses in the brown trout (*Salmo trutta*). *Hydrobiologia* **815**, 267–280.
10.1007/s10750-018-3571-6

Perroud G, Nusbaumer D, de Guttry C, Wedekind C (2021) Male sexual signaling and
expected effects of hatchery-induced sperm competition vary with water depth at
which whitefish are caught. *Current Zoology* **67**, 337–340. 10.1093/cz/zoab007

Rice WR, Gaines SD (1994) Heads I win, tails you lose - testing directional alternative
hypotheses in ecological and evolutionary research. *Trends in Ecology & Evolution* **9**,
235-237.

Rose SR, Petersen NJ, Gardner TJ, Hamill RJ, Trautner BW (2012) Etiology of
thrombocytosis in a general medicine population: analysis of 801 cases with emphasis
on infectious causes. *Journal of Clinical Medicine Research* **4**, 415-423.
10.4021/jocmr1125w

Rosengrave P, Montgomerie R, Gemmell N (2016) Cryptic female choice enhances
fertilization success and embryo survival in chinook salmon. *Proceedings of the Royal*
Society B-Biological Sciences **283**, 20160001. 10.1098/rspb.2016.0001

RStudioTeam (2015) RStudio: Integrated development environment for R. RStudio, Inc.,
Boston, MA.

Rudolfsen G, Figenschou L, Folstad I, Tveiten H, Figenschou M (2006) Rapid adjustments of
sperm characteristics in relation to social status. *Proceedings of the Royal Society B-*
Biological Sciences **273**, 325-332. 10.1098/rspb.2005.3305

Rudolfsen G, Müller R, Urbach D, Wedekind C (2008) Predicting the mating system from
phenotypic correlations between life-history and sperm quality traits in the Alpine
whitefish *Coregonus zugensis*. *Behavioral Ecology and Sociobiology* **62**, 561-567.
10.1007/s00265-007-0480-5

Savary R, Dufresnes C, Champigneulle A, *et al.* (2017) Stocking activities for the Arctic
charr in Lake Geneva: Genetic effects in space and time. *Ecology and Evolution* **7**,
5201-5211. 10.1002/ece3.3073

Seibel H, Bassmann B, Rebl A (2021) Blood will tell: what hematological analyses can reveal
about fish welfare. *Frontiers in Veterinary Science* **8**, 616955.
10.3389/fvets.2021.616955

Turner E, Montgomerie R (2002) Ovarian fluid enhances sperm movement in Arctic charr.
Journal of Fish Biology **60**, 1570-1579.

Urbach D, Folstad I, Rudolfsen G (2005) Effects of ovarian fluid on sperm velocity in Arctic
charr (*Salvelinus alpinus*). *Behavioral Ecology and Sociobiology* **57**, 438-444.

Wang J (2004) Sibship reconstruction from genetic data with typing errors. *Genetics* **166**,
1963-1979. 10.1534/genetics.166.4.1963

Wilkins LGE, Rogivue A, Fumagalli L, Wedekind C (2015) Declining diversity of egg-
associated bacteria during development of naturally spawned whitefish embryos
(*Coregonus* spp.). *Aquatic Sciences* **77**, 481-497. 10.1007/s00027-015-0392-9

Yan SL, Russell J, Harris NR, *et al.* (2013) Platelet abnormalities during colonic
inflammation. *Inflammatory Bowel Disease* **19**, 1245-1253.
10.1097/MIB.0b013e318281f3df

Yano A, Nicol B, Jouanno E, *et al.* (2013) The sexually dimorphic on the Y-chromosome
gene (sdY) is a conserved male-specific Y-chromosome sequence in many salmonids.
Evolutionary Applications **6**, 486-496. 10.1111/eva.12032

Yeates SE, Diamond SE, Einum S, *et al.* (2013) Cryptic choice of conspecific sperm
controlled by the impact of ovarian fluid on sperm swimming behavior. *Evolution* **67**,
3523-3536. 10.1111/evo.12208

Young B, Conti DV, Dean MD (2013) Sneaker "jack" males outcompete dominant
"hooknose" males under sperm competition in Chinook salmon (*Oncorhynchus*
tshawytscha). *Ecology and Evolution* **3**, 4987-4997. 10.1002/ece3.869

Zadmajid V, Myers JN, Sorensen SR, Ernest Butts IA (2019) Ovarian fluid and its impacts on
spermatozoa performance in fish: A review. *Theriogenology* **132**, 144-152.
10.1016/j.theriogenology.2019.03.021

Zajitschek SRK, Brooks RC (2010) Inbreeding depression in male traits and preference for
outbred males in Poecilia reticulata. *Behavioral Ecology* **21**, 884-891.
10.1093/beheco/arq077
782

Tables

Table 1. Sperm characteristics after activation in water with different types of ovarian fluid (likelihood ratio tests on linear mixed-effects models). Significant *P*-values are highlighted in bold.

Model terms	Effect tested	d.f.	AIC	χ^2	*p*
(A) Sperm velocity					
h + m + f		5	649.0		
m + f	h	4	653.3	6.3	**0.01**
h + f	m	4	660.1	13.2	**<0.001**
h + m	f	4	652.6	5.7	**0.02**
h + m × f	m × f	6	637.8	13.2	**<0.001**
(B) Sperm motility					
m + f		4	587.6		
f	m	3	670.1	84.6	**<0.001**
m	f	3	585.6	0	1
m × f	m × f	5	589.6	0	1
(C) Longevity					
m + f		4	664.7		
f	m	3	687.7	25.0	**<0.001**
m	f	3	694.2	31.5	**<0.001**
m × f	m × f	5	527.6	139.1	**<0.001**

Fixed effect: h, sperm head size; random effects: m, male identity; f, female identity

Table 2. Correlates to fertilization success of the focal male in the sperm competition trials (experiment 2), if the focal male is (A) the more yellow one of the two competitors, (B) the one with the lower inbreeding coefficient F_{beta}, or (C) the least related to the female. Generalized linear model with female (egg donor), dyad (pair of competing males), ovarian fluid (present of absent), and the difference in mean sperm velocity between rivals (focus – opponent, in water or in ovarian fluid, respectively, as determined in experiment 1). Non-significant interaction terms were removed. Final tests were either two-tailed or directed (p_{dir}) based on the *a priori* predictions that ovarian fluids have either no effect or they support sperm of focal males. Final *P*-values are highlighted in bold if significant.

Factor	d.f.	χ^2	$p_{two\text{-}tailed}$	p_{dir}
(A) Focal male: more yellow				
Female	2	3.1	0.21	
Dyad	4	130.7	**<0.001**	
Ovarian fluid	1	4.3	0.039	**0.02**
Difference in sperm velocity	1	0.8	0.37	0.23
(B) Focal male: lower F_{beta}				
Female	2	1.0	0.60	
Dyad	4	119.2	**<0.001**	
Ovarian fluid	1	4.2	0.040	0.10
Difference in sperm velocity	1	0.9	0.35	0.22
(C) Focal male: lower kinship				
Female	2	27.0	**<0.001**	
Dyad	4	57.8	**<0.001**	
Ovarian fluid	1	0.1	0.76	0.48
Difference in sperm velocity	1	8.8	0.003	**0.002**

Figure 1. Yellowness in male lake char and their links to size, inbreeding coefficient, sperm characteristics, and immune parameters. The two fish illustrate the two extremes of observed yellowness in 2017. Yellowness versus (A) body size (r = -0.51, n = 34, p = 0.002), (B) inbreeding coefficients (F_{beta}; r = 0.16, p = 0.65), the sperm characteristics (C) velocity, (D) rate of immobile sperm, and (E) longevity (see Table S3 for statistics on sperm characteristics), and the immune parameters (F) leukocyte counts (% of all blood cells; r = -0.47, p = 0.17), (G) relative lymphocytosis (% of lymphocytes among leukocytes; r = -0.73, p = 0.02), and (H) thrombocytosis (thrombocytes counts per 100 blood cells; r = -0.74, p = 0.01). Regression lines illustrate significant relationships. Symbols and regressions in light grey indicate samples collected in 2017, darker symbols and regression lines those from 2018.

Figure 2. The effects of ovarian fluids on sperm of different types of males. Panels A-C show sperm characteristics (Turkey box plots with quartiles and whiskers) when activated in water only or in diluted ovarian fluids: (A) sperm velocity (μm/sec; comparing means: Welch's $F_{1,12.014} = 0.18$, $p = 0.68$; variances: Levene's $F_{1,18} = 9.9$, $p = 0.006$), (B) rate of immobile sperm (means: $F_{1,17.995} = 0.48$, $p = 0.50$; variances: $F_{1,18} = 0.03$, $p = 0.87$), and (C) sperm longevity (sec; means: $F_{1,14.486} = 207.6$, $p < 0.001$; variances: $F_{1,18} = 0.5$, $p = 0.49$). Panels e-g give the differences between of sperm characteristics in ovarian fluid and in water (8 measurements per male in ovarian fluids compared to the respective mean in water each) relative to male yellowness: (D) differences in sperm velocity (regression on means per male: $r = 0.65$, $n = 10$, $p = 0.04$) , (E) differences in rates of immobile sperm ($r = -0.36$, $p = 0.31$), (F) differences in longevity ($r = 0.65$, $p = 0.04$). Panels G-I give the mean sperm characteristics per combination of male identity and donor of ovarian fluid relative to their kinship: (G) sperm velocity ($r = 0.08$, $n = 40$, $p = 0.64$), (H) rate of immobile sperm ($r = 0.16$, $p = 0.34$), and (I) sperm longevity (Spearman rank order correlation coefficient to account for potential extremes: $r_s = 0.39$, $p_{two-tailed} = 0.012$, $p_{dir} = 0.03$ based on the *a priori* prediction that kinship should be avoided). Regression lines illustrate significant relationships, the dotted line in (D) and (E) mark zero difference.

Supplementary information

Sperm of more colorful males are better adapted to ovarian fluids in lake char (Salmonidae)

David Nusbaumer, Laura Garaud, Christian de Guttry & Claus Wedekind.

Department of Ecology & Evolution, University of Lausanne, Lausanne, Switzerland
[*]correspondence: Claus Wedekind, claus.wedekind@unil.ch, orcid.org/0000-0001-6143-4716

Content

Table S1. Experiment 1: Determining sperm characteristics in various activation media, i.e. diluted ovarian fluid of one of 4 females (F1-F4) or water only. The entries give the number of independent sets of measurements.

Activation medium	Male									
	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10
Ovarian fluid of F1	2	2	2	2	2	2	2	2	2	2
Ovarian fluid of F2	2	2	2	2	2	2	2	2	2	2
Ovarian fluid of F3	2	2	2	2	2	2	2	2	2	2
Ovarian fluid of F4	2	2	2	2	2	2	2	2	2	2
Water only	2	2	2	2	2	2	2	2	2	2

Table S2. Experiment 2: The sperm competition trials, with 2 times 24 eggs per experimental cell. Asterisks indicate experimental cells that were accidentally lost.

Female	Activation	Dyad				
		M1 vs M2	M3 vs M4	M5 vs M6	M7 vs M8	M9 vs M10
F1	ovarian fluid	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24
F1	water	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24
F2	ovarian fluid	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24
F2	water	2 x 24*	2 x 24*	2 x 24*	2 x 24*	2 x 24*
F3	ovarian fluid	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24
F3	water	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24
F4	ovarian fluid	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24
F4	water	2 x 24	2 x 24	2 x 24	2 x 24	2 x 24

Table S3. The 6 microsatellite markers used for paternity assignment, and the sex marker that was added to the first multiplex.

Marker	Multiplex	Sequence forward primer	Sequence reverse primer	Dye	$N_{alleles}$	Range [bp]
Cocl3-N	1	5'-TTCAGGTTTGGTAAGCAAG-3'	5'-AGTGTAATAAATCACCCGAG-3'	atto550	8	226-270
OtsG253b	1	5'-GAGCAGGCCGAGCAGGTGTCT-3'	5'-AATTGGGTCATTAAGGCTCTGTGG-3'	fam	16	88-118
Ssa456	1	5'-CTTCCCAGGAGTCATCATAAATCT-3'	5'-TAAACCCCACTGCTTGTTGAGTGT-3'	hex	4	209-215
Sfo8	2	5'-CAACGAGCACAGAACAGG-3'	5'-CTTCCCCTGGAGAGGAAA-3'	hex	12	248-275
Sfo23	2	5'-GTGTTCTTTTCTCAGCCC-3'	5'-AATGAGCGTTACGAGAGG-3'	atto550	11	136-217
Ssa85	2	5'-AGGTGGGTCCTCCAAGCTAC-3'	5'-ACCCGCTCCTCACTTAATC-3'	fam	11	156-189
Sdy[1]	1	5'-CCCAGCACTGTTTTCTTGTCTCA-3'	5'-CTTAAAACCACTCCACCCTCCAT-3'	atto620	1	226-226

[1]Sex marker, used for a parallel study on sex-specific embryo development (Nusbaumer *et al.* 2021).

Table S4. Characteristics of sperm activated in water only, predicted by male body length, yellowness and year of sampling. Likelihood-ratio tests on linear mixed-effects models were used when year of sampling could be added as random factor (a-c), while linear models were used when measurements were only taken in 2017 (d). The table shows the outcome after removal of least significant parameter and gives the AIC for each model. Bold P-values highlight significant effects ($P \leq 0.05$).

Model terms	Effect tested	d.f.	AIC	χ^2	P
a) Concentration					
y + year			43.7		
year	y	1	46.5	4.8	**0.03**
b) Sperm velocity					
y + year			319.5		
year	y	1	321.3	3.8	**0.05**
c) Immotiles					
y + year			269.2		
year	y	1	271.2	4.0	**0.05**
		d.f.	AIC	SumSq	P
d) Longevity					
l + y			119.4		
y	l	1	118.3	137.4	0.27
l	y	1	117.6	56.9	0.48

Fixed effects: y, yellowness; l, body length. Random effect: year

Table S5. Identity of focal males if it is the more yellow than its competitor, less inbred, or least kin to a given female.

Dyad	More yellow	Less inbred	Least kin to female			
			F1	F2	F3	F4
1	M2	M2	M2	M2	M2	M1
2	M3	M4	M4	M3	M4	M3
3	M5	M6	M5	M6	M6	M6
4	M7	M8	M8	M8	M8	M8
5	M10	M9	M10	M9	M10	M10

Figure S1. Locations of the 3 squares (in green) from which color measurements of the ventral area (yellow and red) were taken. Mean gray value was positively correlated with yellowness, i.e. the yellower the lighter (r = -0.47, p < 0.005) and negatively with redness, i.e. the redder the darker (r = -0.66, p < 0.001).

Figure S2. Example of a blood smear image as taken under the microscope (left) and its output of automated cell count (right). On the left image, black arrow heads point towards lymphocytes, the white arrow towards a granulocyte, and the black and white arrow towards thrombocytes. The rest are erythrocytes.

Figure S3. The link between multi-locus heterozygosity (MLH) and individual inbreeding coefficients (F_{beta}) for the males (blue) and females (red) sampled in 2017 and used in experiments 1 and 2.

Figure S4. Immunological and sperm characteristics of males relative to their inbreeding coefficient (F_{beta}): (A) leukocytes (% of all blood cells; r = 0.17, p = 0.88), (B) relative lymphocytosis (percentage of lymphocytes among leukocytes; r = 0.11, p = 0.61), (C) thrombocytes per 100 blood cells (r = -0.20, p = 0.75), (D) sperm velocity (µm/sec; r = -0.25, p = 0.48), (E) percentage of immotile sperm (r = 0.74, p = 0.01). The regression line is given for the significant relationship. Sperm longevity was not measured in this sample that was taken in 2017.

Figure S5. Relatedness between males and females (kinship) that were used in the sperm competition trials (4 females x 10 males = 40 combinations) versus (A) male skin colorations (yellowness; r = 0.13, n = 40, p = 0.42) and (B) individual male inbreeding coefficients (F_{beta}; r = 0.12, p = 0.47). The dotted lines indicate the non-significant regressions.

Figure S6. Interactions effects between females' ovarian fluids and males' identities on A) sperm velocity and (B) sperm longevity. Symbols depicts means of within subject repeated measures with their 95% confidence intervals. See Table 1 for statistics.

Figure 3. Competitive success of different types of focal males under sperm competition in water only or when exposed to ovarian fluids. The mean fertilization rates of (A) the yellower male of the two competitors each, (B) the less inbred male each, and (C) the male each that is least related to the female whose ovarian fluids the sperm was exposed to. Turkey box plots with quartiles and whiskers ($N_{water} = N_{ovarian\ fluid} = 15$). The asterisk indicates a significant difference linked to the activation medium, n.s. = not significant. See Table 2 for statistics.