1

Stepwise progression of β-selection during T cell development as revealed by histone

deacetylation inhibition

TCR co-receptors are expressed sequentially to fine-tune early αβ T cell development

[1,2]Anchi S Chann, [1,2]Mirren Charnley, [3]Lucas Newton, [2,7]Andrea Newbold, [3,4,5,6]Patrick O Humbert,

[2,7]Ricky W Johnstone, [1,2,7]Sarah M Russell.

Affiliations
[1]Optical Sciences Centre, School of Science, Swinburne University of Technology, Hawthorn, Victoria

3122; Australia.

[2]Peter MacCallum Cancer Centre, Melbourne, Victoria 3000 Australia.

[3] Department of Biochemistry & Chemistry, La Trobe Institute for Molecular Science, La Trobe

University, Melbourne, Victoria 3086, Australia.

[4] Research Centre for Molecular Cancer Prevention, La Trobe University, Melbourne, Victoria 3086,

Australia.

[5] Department of Biochemistry & Pharmacology, University of Melbourne, Melbourne, Victoria 3010,

Australia.

[6] Department of Clinical Pathology, University of Melbourne, Melbourne, Victoria 3010, Australia.

[7]Sir Peter MacCallum Department of Oncology, The University of Melbourne, Victoria 3010, Australia.

*To whom correspondence should be addressed: Sarah Russell: sarah.russell@petermac.org;

613-433-354-996; Optical Sciences Centre, School of Science, Hawthorn, Australia

ORCID ID: 0000-0002-8778-5327 (ASC); ORCID ID: 0000-0003-3845-9290 (MC); ORCID

ID: 0000-0002-1366-6691(POH); 0000-0001-5826-9641 (SMR); 0000-0001-7053-9237

(RWJ); 0000-0002-0481-2431 (AN)

Abstract

During T cell development, the first step in creating a unique T Cell Receptor (TCR) is the genetic

recombination of the TCRβ chain. The quality of the newly recombined TCRβ is assessed at the β-

selection checkpoint. Most cells fail this checkpoint and die, but the coordination of the complex

events that control fate at the β-selection checkpoint is not yet understood. We shed new light on fate

determination during β-selection using a selective inhibitor of histone deacetylase 6, ACY1215.

ACY1215 disrupted the β-selection checkpoint. Characterising the basis for this disruption revealed a

new, pivotal stage in β-selection, bookended by upregulation of the TCR co-receptors, CD28 and CD2

respectively. Within this 'DN3b^{Pre}' stage, CD5 and Lef1 are upregulated to reflect pre-TCR signalling

and their expression correlates with proliferation. During this phase, ACY1215-mediated disruption

of the organisation of the β-selection immunological synapse is associated with a breakdown in

connectivity of expression of pre-TCR, CD5 and Lef1. Subsequent deregulation of pre-TCR-induced

proliferation leads to bypass of the β-selection checkpoint and subsequent failure to progress. We

propose that the progressive expression of CD28, CD5 and Lef1, then CD2 reports and modulates the

pre-TCR signal to orchestrate passage through the β-selection checkpoint. These findings suggest a

refined model of β-selection in which a coordinated increase in expression of pre-TCR, CD28, CD5

and Lef1 provides for escalating TCR signalling strength, and culminates in the expression of CD2 to

enable exit from the β-selection checkpoint.

Introduction

T cells are generated in a multistep, highly orchestrated process guided by stromal cells in the thymus.

T cell development involves massive cell proliferation interspersed with genomic recombination of

the T cell receptor (TCR). If genomic recombination, proliferation and death are not appropriately

orchestrated, the implications for cancer and immunity are profound. Uncontrolled self-renewal or a

sub-optimal TCR repertoire can lead to T cell acute lymphoblastic lymphoma (T-ALL), autoimmunity

or limited response to pathogens or cancers (Ballesteros-Arias et al., 2019; Douek et al., 2000;

Notarangelo et al., 2020). T cell development is most tightly controlled during genomic

recombination to generate the TCR. For the most common αβ T cells, the β chain is generated at the

DN3a phase of T cell development (see **Supp Fig 1A** for a description of T cell development and the

surface markers by which it is assessed), and the appropriateness of this TCRβ sequence is assessed

during the β-selection checkpoint. During β-selection, the strength of the TCRβ chain is tested in

combination with a surrogate 'pre-TCR' before the TCRα chain is created (Chann and Russell, 2019;

Dutta et al., 2021).

β-selection is in part determined by stromal cells in the thymic microenvironment, which provide

Notch and chemokine signals that promote survival and differentiation (Chann and Russell, 2019;

Dutta et al., 2021). Recently, it has become clear that interactions between the pre-TCR and the

MHC-peptide on stromal cells also provide an opportunity to test TCR signalling (Allam et al., 2021;

Dutta et al., 2021; Li et al., 2021) These interactions trigger a sequence of transcriptional changes and

epigenetic modifications that ensure only the cells with an appropriately recombined TCRβ chain

survive, proliferate and differentiate. How pre-TCR signalling integrates fate determination including

proliferation, differentiation and survival during β-selection is still not understood (Chann and

Russell, 2019; Dutta et al., 2021).

Epigenetic therapies such as small-molecule inhibitors of histone deacetylases (HDAC) have shown

great promise in mediating clinical antitumor responses and are emerging as immune modulators

(Hogg et al., 2020). The HDAC6 selective inhibitor ACY1215 (Ricolinostat) can synergize with the

BET inhibitor, JQ1, to recruit T cell mediated responses to lung adenocarcinoma, enhance T cell

responses to *ex vivo* melanoma cultures, impair effector CD8 T cell function during skin

inflammation, and alter Treg function (Adeegbe et al., 2017; Laino et al., 2019; Tsuji et al., 2015; Xu

et al., 2018; Zhang et al., 2021). HDACs 1, 2, and 3 are required at multiple stages of T cell

development, but a role for HDAC6 in T cell development has not been identified (Wang et al., 2020).

Mice lacking HDAC6 showed no overt T cell phenotypes (Shapiro and Shapiro, 2020; Zhang et al.,

2008) and to our knowledge the effect of acute deletion of HDAC6, or pharmacological inhibition of

HDAC6 on T cell development has not been reported (Shapiro and Shapiro, 2020). Here, we show a

striking effect of treatment with the HDAC6-selective inhibitor, ACY1215, on β-selection.

This effect of ACY1215 allowed us to define several stages of β-selection with high resolution. After

the DN3a stage marked by upregulation of the co-receptor, CD28, we observed a transitional stage

marked by low levels of the co-receptor, CD2, that we termed DN3b^{Pre}. DN3b^{Pre} cells are enriched by

ACY1215 treatment. An upregulation of CD2 marks passage through the β-selection checkpoint, to a

phase we termed DN3b^{Post}. The transcription factor, Lef1 is upregulated in DN3b^{Pre} cells, together

with an increasing expression of the reporter of TCR signalling, CD5. These findings implicate a

refined model of β-selection in which, following expression of CD27 and CD28, a coordinated

increase in expression of pre-TCR, CD5 and Lef1 provides for an escalating test of TCR signalling.

Expression of CD2 marks the passing of this test and exit from the β-selection checkpoint.

Results

Treatment with ACY1215 impairs T cell differentiation at the β-selection stage.

To examine the effects of ACY1215 on T cell development, we used two models. Firstly, a well

validated *in vitro* model in which hematopoietic stem cells extracted from mouse fetal liver were co-

cultured with the OP9 mouse bone marrow stromal cell line transfected with the Notch ligand, DL1

('OP9-DL1') (Holmes and Zúñiga-Pflücker, 2009). To assess relevance of our findings to

physiological T cell development in the thymus, we also analysed cells directly taken from young

adult mice.

At Day 8-10 of the co-culture, when cells from mouse fetal liver were predominantly at the DN3 stage

of T cell development **(Supp Fig 1B)**, we applied the HDAC6 inhibitor, ACY1215 at a concentration

previously reported to alter function of mature T cells without affecting lymphocyte viability (Laino

et al., 2019). Compared with DMSO-treated cells, cells treated with ACY1215 for two days (**Figure**

1A) showed reduced proportions of the later stages of differentiation, suggesting HDAC6 is required

for optimal T cell development to the DP stage. This shift in differentiation was accompanied by a

generalised inhibition of expansion in cell numbers at several stages of T cell development (**Supp Fig**

2A). Comparing proportions at different DN stages using CD25 and CD44 expression (Figure 1A)

showed an increase in the proportions of DN3 cells and commensurate decrease in proportions of

DN4, suggesting that the ACY1215-treated cells were blocked at the DN3 stage.

The DN3 stage incorporates the β-selection checkpoint. The passing of the β-selection checkpoint,

conventionally conceptualised as DN3a-to-DN3b progression, is marked by the expression of CD28

(Teague et al., 2010; Williams et al., 2005). Therefore, we assessed whether DN3a (DN3, CD28^{Lo})

cells were inhibited from progressing to DN3b (DN3, CD28^{Hi}) by ACY1215 treatment over one and

two days (**Figure 1B**). Surprisingly, the reverse was observed. Co-staining with CD25 and CD28

indicated a relatively stable distribution between DN3a and DN3b at 1 and 2 days in the control

cultures, but increasingly greater proportions of DN3b cells in the ACY1215-treated cultures. A

similar shift to DN3b and DN4 at the expense of DN3a was observed in thymocytes extracted from an

adult mouse and cultured for 1 day on OP9-DL1, although the relatively low numbers of DN3a in

primary thymocytes *ex vivo* made the shift less obvious (**Figure 1C, Supp Fig 1C**).

We explored the cellular basis of this shift away from DN3a towards DN3b. Cell counts of cultures of

sorted DN3a cells showed that the DN3a cells expanded more than 4-fold over 2 days of control

DMSO-treated culture, but this was dramatically reduced by ACY1215 treatment (Fig 1B**)**. In

contrast, ACY1215 caused a negligible reduction in absolute numbers of DN3b and DN4 cells arising

from DN3a. This stage-specific effect was probably not due to differences in proliferation or death

because CFSE and Ki-67 labelling showed similar inhibition of proliferation of DN3a and DN3b cells

(**Supp Fig 3**) and neither DN3a or DN3b cells were demonstrably susceptible to cell death, although

DP cells died more in the presence of ACY1215 (**Supp Fig 4**). Together, these data suggest that

ACY1215 treatment of DN3a cells led to reduced amplification, but little impact or a compensatory

increase in the number of cells passing through the DN3b stage.

To reconcile the seemingly normal progression through β-selection (as indicated by DN3b and DN4

counts) with the block in DP cells, we first monitored differentiation from sorted DN3a and DN3b

cells that had been treated for four days with ACY1215 or DMSO (**Supp Fig 5**). Progression past the

DN3 stages was inhibited for both DN3a and DN3b starting populations, although less clear for the

DN3a starting population due to limited differentiation by Day 4. To confirm that the effect at β-

selection led to a downstream impact on DP cells, we counted the number of cells at each

differentiation stage that developed from the sorted DN3b population (**Figure 1D**). By day 4, DN3b

cells had expanded more than 50-fold, with a negligible reduction caused by treatment with

ACY1215. DN4 cell numbers were also only slightly reduced by ACY1215 treatment, but DP were

reduced by over two orders of magnitude. These data together suggest that ACY1215 treatment

impacts generally upon cell expansion, but also alters differentiation to DP by acting at the DN3

stage, perhaps by causing precocious differentiation from DN3a to DN3b.

The DN3 cell stage involves fate choice between two T cell lineages ($\alpha\beta$ cells and $\gamma\delta$ cells), each

expressing a different set of T cell receptors (Hayday and Pennington, 2007). To determine which

lineage was altered by ACY1215, we cultured developing T cells for 2 days with and without

HDAC6i and labelled the cells with antibodies specific for these TCRs to assess progression through

the DN and DP stages (**Supp Fig 6**). ACY1215 treatment reduced the proportion of TCRβ–expressing

cells at all stages following β-selection but did not inhibit TCR$\gamma\delta$ development. Together, these data

indicated that the defect in differentiation to DP following ACY1215 treatment was specific to $\alpha\beta$ T

cells, and likely involved the β-selection checkpoint.

One possible explanation for this passage to DN3b without optimal subsequent differentiation could

be that ACY1215-treated cells express some markers suggestive of differentiation, but are not

competent precursors of $\alpha\beta$ T cells. To assess this, we first characterised several phenotypic markers

implicated in progression through the β-selection checkpoint. Co-staining DN3 cells with both CD28

and CD27, both of which can discriminate between DN3a and DN3b cells (Taghon et al., 2006;

Teague et al., 2010) clearly discriminated two subpopulations in DN3 cells pre-treated with DMSO

(**Figure 2A**). A shift from DN3a to DN3b is observed for both fetal liver-derived lymphocytes and *ex*

vivo extracted thymocytes cultured on OP9-DL1 stroma. These data confirm the findings above that,

at least by the canonical markers, CD27 and CD28, ACY1215 treatment promotes differentiation

towards DN3b at the expense of DN3a.

Given that the primary function of this checkpoint is to ensure appropriate TCRβ rearrangement, we

assessed the correlation of each marker with TCRβ expression (**Figure 2B**). As previously shown

(Williams et al., 2005), expression of CD28 was clearly correlated with expression of icTCRβ in cells

derived from fetal liver or from the adult thymus. The CD28[Hi] cells also expressed more of the

nutrient transporters, CD71 and CD98, and more surface TCRβ (**Supp Fig 7A**) indicating that, in

general, they had progressed further through β-selection, but the correlation was not absolute. For

both cell sources, ACY1215 treatment resulted in less icTCRβ expression in the DN3b (whether

measured as CD28[Hi] or CD27[Hi] DN3 cells) (Fig 2B, see also Supp Fig 8C). To validate the icTCRβ

staining, we directly compared it with surface TCRβ (**Supp Fig 7B**). For both cell sources, expression

of surface TCRβ was highly correlated with that of icTCRβ and both were reduced by ACY1215

treatment. Thus, as well as depleting DN3a cells, treatment with ACY1215 increases the proportions

of a cell population that bears many of the phenotypic markers of passage through β-selection, but

these cells appear somewhat defective in expression of TCRβ, and do not progress optimally through

the DP stage. We speculated that further characterisation of these cells might yield valuable

information about requirements at the β-selection checkpoint.

CD2 expression identifies DN3b[Pre] and DN3b[Post] as two functionally distinct stages in β-selection.

A well-established co-stimulator of TCR signalling is CD2 (Skånland et al., 2014). Expression of

CD2 marks TCRβ V(D)J rearrangement and can facilitate pre-TCR signalling for passage through the

β-selection checkpoint (Groettrup et al., 1992; Rodewald et al., 1993; Sasada and Reinherz, 2001). In

the next series of experiments, we find that CD28 and CD2 reliably define 3 sequential stages of β

selection in DN3 cells, which we term DN3a, DN3b[Pre], and DN3b[Pro] to accommodate pre-existing

nomenclature (**Figure 3A**). Co-staining of DMSO-treated DN3 cells with CD28 and CD2 showed that

all DN3a cells (CD28[Lo]) were CD2[Lo], but DN3b (CD28[Hi]) cells had two clearly discernible

populations of CD2[Lo] (DN3b[Pre]) and CD2[Hi] (DN3b[Pro]) (**Figure 3Bi**). ACY1215 treatment enriched for

the DN3b[Pre]. Cell counts indicated that the numbers of DN3b[Pre] cells were not substantially affected

by ACY1215, but DN3a and DN3b[Post] numbers were dramatically reduced (**Supp Fig 8A**). Thus,

ACY1215 treatment led to an enrichment in a DN3b[Pre] population that was low for CD2. The DN3b[Pre]

population **(Fig 3Bii)** was evident in thymocytes from young adult mice, even without *in vitro* culture

(see Fig 4 below), and was again enriched by treatment with ACY1215.

To confirm that DN3b^{Pre} cells were earlier in β-selection than DN3b^{Post} cells, we assessed the

association of CD2 expression with proliferation as measured by CFSE dilution (**Supp Fig 8B**). Two

clear populations were evident, with the CD2^{Hi} (DN3b^{Post}) cells having undergone substantially more

proliferation than the CD2^{Lo} (DN3b^{Pre}) cells. ACY1215 treatment did not disrupt the association

between CD2 and proliferation. Expression of CD27 and surface TCRβ were also increased from

DN3a, DN3b^{Pre}, to DN3b^{Post}, with stepwise correlation which was not altered by ACY1215 despite

lower surface TCRβ on DN3b^{Post} cells (**Supp Fig 8C**). We then assessed icTCRβ and Ki-67 staining,

which should reflect the cell proliferation triggered by successful passage through the β-selection

checkpoint (**Figure 3C**). Indeed, consistent with previous findings that TCRβ V(D)J recombination is

restricted to DN cells positive for CD2 (Rodewald et al., 1993), DN3b^{Pre} cells were intermediate in

expression of icTCRβ and Ki67i.e. between the levels seen on DN3a cells and DN3b^{Post} cells. The

correlated, stepwise increase in icTCRβ and Ki-67 was observed in cultures from both fetal liver and

adult thymus (Figure 3Ci & ii). This pattern was even more striking in thymocytes from adult mice,

where DN3a cells showed no proliferation, DN3b^{Pre} cells were heterogeneous with respect to

proliferation, and proliferation correlated with icTCRβ expression, and DN3b^{Post} cells were all highly

proliferative, with minimal impact from ACY1215 treatment. These data suggest that the dramatic

reduction in TCRβ expression upon ACY1215 treatment observed in Fig 2 was explained more by the

enrichment of DN3b^{Pre} cells, by than a direct effect on TCRβ expression. c-Myc, which mediates the

proliferative response to pre-TCR signalling but is not required for differentiation (Dose et al., 2006),

showed a similar stepwise increase with these differentiation stages (**Supp Fig 8D**). These data are all

compatible with an interim DN3b^{Pre} stage that reflects an early, and pivotal step in β-selection where

rearrangement of the TCRβ chain is translated into proliferation.

To formally assess whether the DN3b^{Pre} stage is a transitional stage before DN3b^{Post}, we sorted DN3a,

DN3b^{Pre} and DN3b^{Post} cells from adult mice and cultured them for further analysis (**Figure 4**).

Analysis of the cells immediately after extraction showed clearly distinct DN3a, DN3b^{Pre} and

DN3b^{Post} cells. After two days, DN3b^{Post} cells had produced substantial numbers of DP cells, with

fewer produced by DN3b^{Pre}, and fewer still by DN3a cells. Also compatible with unidirectional

progression from DN3a to DN3b^{Pre} to DN3b^{Post}, was the proportion of DN4 and DN3 stages from each

sorted population. ACY1215 treatment of thymocytes stabilized the DN3b^{Pre} population and blocked

progression to DP. Similar data was obtained in the fetal liver-derived cultures (**Supp Fig 9**). These

data confirm that DN3b^{Pre} (CD28^{Lo}CD2^{Lo}) is a transitional state between DN3a (CD28^{Hi}CD2^{Lo}) and

DN3b^{Post} (CD28^{Hi}CD2^{Hi}), reflecting partial progression through β-selection.

ACY1215 alters the pre-TCR signalling platform clustered at MTOC

The enrichment of CD28^{Hi}CD2^{Lo} DN3b^{Pre} cells by ACY1215 treatment in both fetal liver-derived

cells and thymocytes from adult mice suggests that a key process in this stage might involve

acetylation. We explored three possible mechanisms by which ACY1215 might disrupt β-selection.

ACY1215 is selective for HDAC6, and can inhibit the catalytic activity of HDAC6 (Eich et al., 2018;

Narita et al., 2019). HDAC6 protein is expressed at every stage of T cell development (**Supp Fig**

10A). The best-known role of HDAC6 is in modification of histones for epigenetic regulation. Indeed,

acetylation of H3K18 was increased upon treatment with ACY1215, suggesting that epigenetic

modification could be affected by the drug (**Supp Fig 10B**). Although best known as a regulator of

histone acetylation, HDAC6 can also acetylate α-tubulin (Narita et al., 2019). In mature T cells, the

regulation of acetylation of α-tubulin by HDAC6 influences TCR-mediated recruitment of the

microtubule-organising centre (MTOC) to the immunological synapse (Serrador et al., 2004). In all T

cell developmental stages, HDAC6 was predominantly in the nucleus, but also detectable in the

cytoplasm and near the MTOC, compatible with effects on MTOC modification (**Supp Fig 10C-E**).

Acetylation of α-tubulin (K40), which can be de-acetylated by HDAC6 (Osseni et al., 2020), was

evident throughout the T cell differentiation stages (**Figure 5A**). Upon a two-hour treatment with

ACY1215, each population showed an increase in acetylated tubulin, suggesting HDAC6 deacetylated

tubulin throughout early T cell development.

We previously found that T cells undergoing β-selection form an immunological synapse, in which

MTOC and other TCR-associated signalling proteins move with the MTOC to the site of antigen

receptor signalling receptor signalling (Allam et al., 2021). We therefore speculated that an alteration

in MTOC organisation might mediate the effects of ACY1215 on β-selection by impacting the

immunological synapse. To address this, we first assessed the association of LAT with the MTOC in

DN3a cells (**Figure 5B**). Acetyl-α-tubulin presented as a single cluster in the site of the MTOC, and

was increased in intensity with ACY1215 treatment, compatible with the flow cytometry results.

LAT was strongly associated with the MTOC in DMSO treated cells, but altered by ACY1215

treatment, with less intense fluorescence and less clustering at the MTOC. Notch is particularly

important during β-selection in mediating assembly of the immunological synapse and transmitting

signals downstream of the pre-TCR signal (Allam et al., 2021; Charnley et al., 2020). We therefore

assessed the co-localisation of Notch1 and the MTOC in DN3a cells (**Figure 5C**). In DMSO treated

cells, Notch was concentrated at the MTOC, but, similar to LAT, was much more diffuse following

ACY1215 treatment. Interestingly, the change in localisation of both Notch and LAT was

accompanied by a loss in their intensity. A reduction in surface Notch expression was observed in

DN3a, DN3b and DN4 cells, suggesting that the reduction in Notch1 transcription that normally

occurs following β-selection (Yashiro-Ohtani et al., 2009) might be accelerated by ACY1215 (**Fig**

5D). Together, these data suggest that a possible mechanistic basis for the effects of ACY1215 on β-

selection is via alterations in recruitment by MTOC of Notch1 and TCR signalling components to the

immunological synapse, associated with enhanced tubulin acetylation at the immunological synapse.

DN3b[Pre] cells have commenced but not completed β-selection, and are controlled by sequential

changes in key regulators of β-selection.

How might such a disruption in the immunological synapse and TCR signalling alter progression

from DN3b[Pre] to DN3b[Post]? A collaboration between pre-TCR and Wnt signalling is well established,

with two core Wnt pathway transcriptional regulators: β-catenin and Lef1/TCF-1 essential to β-

selection (Gounari et al., 2001; López-Rodríguez et al., 2015; Staal and Clevers, 2005; Travis et al.,

1991; Xu et al., 2009; Zhao et al., 2021). TCF-1 and Lef1 can directly modify histone acetylation, and

this function is inhibited by the pharmacological HDAC inhibitors, Tubacin and Vorinostat, (Xing et

al., 2016; Zhao et al., 2021). In support of functional interplay between ACY1215 treatment and TCF-

1 and Lef1, TCF-1 deficiency leads to a similar phenotype as ACY1215 treatment, with disrupted β-

selection and reduced expression of icTCRβ in DN4 cells (Goux et al., 2005), and deletion of both

TCF-1 and Lef1 leads to a complete block at DN4 (Yu et al., 2012). We focused on Lef1 since Lef1 is

specifically expressed at the β-selection stage (Mingueneau et al., 2013). To assess whether Lef1

might mediate the β-selection decisions at the DN3b^{Pre} stage, we explored how its expression changed

with differentiation, and how it was impacted by ACY1215 treatment. Lef1 was minimally expressed

in DN3a cells, uniformly bright in DN3b^{Pro} cells, but heterogeneous in DN3b^{Pre} cells derived from

either fetal liver or adult thymus (**Fig 6Ai & ii**). In the DN3b^{Pre} cells, Lef1-expression was associated

with increased expression of TCRβ, albeit with a wide range in TCRβ expression (**Fig 6B**). ACY1215

treatment did not demonstrably alter icTCRβ expression (Fig 6B) but, prevented the upregulation of

Lef1 expression in DN3b^{Pre} cells, suggesting a role in pre-TCR-mediated induction of Lef1.

The correlation in expression of TCRβ with proliferation and Lef1 expression raised the possibility

that the pre-TCR signal might be directly translated into phenotype during the DN3b^{Pre} stage. To

explore this, we investigated the TCR-associated receptor, CD5. A role for CD5 has not been

demonstrated during β-selection, but during positive selection expression of CD5 reports the strength

of TCR signalling and tunes TCR signalling responsiveness, dampening the response of cells to TCR-

signalling via the NF-κB pathway (Azzam et al., 2001; Azzam et al., 1998; Tarakhovsky et al., 1995;

Voisinne et al., 2018). To explore the dynamics of CD5 expression during β-selection, we stained

fetal liver-derived DN3 cells with CD2, Lef1 and CD5 (**Supp Fig 11**). Gating for DN3b^{Pre} with low

and high Lef1 expression showed strong correlation of CD5 and Lef1expression. Interestingly, Lef1

upregulation was retained in the presence of ACY1215, but the correlation with CD5 expression was

lost. These data suggest a model in which CD5 expression is downstream of Lef1, and suggest the

effects of ACY1215 on progression through β-selection might be mediated by disrupting the

functional connection between Lef1 and CD5. Lef1 and CD5 expression were similarly correlated in

DN3 cells extracted from the adult thymus, particularly in DN3b^{Pre} cells (**Figure 7A**).

To confirm this model, we made use of the fact that progression past DN3 is marked by

downregulation of CD25, and a transitional stage between DN3b and DN4 (sometimes termed DN3c)

can be identified within the DN3 population (Ananias et al., 2008; Teague et al., 2010). We therefore

assessed the relationship between CD2 and CD25 in the DN3b cells. DN3b^{Pre} population was still

high for CD25, but DN3b^{post} included CD25^{Lo} cells. (**Supp Fig 12A**). These data are compatible with

the notion that DN3b^{Post} marks a later stage in differentiation, that includes, but is not exclusive to, the

DN3c population (Ananias et al., 2008). To assess whether the combined expression of CD5 and Lef1

in DN3b cells indicated progression through β-selection, we gated for the two clearly discriminated

sub-populations Lef1^{Lo}CD5^{Lo} and Lef1^{Hi}CD5^{Hi}, and compared their CD25 expression (**Supp Fig**

12B). The Lef1^{Hi}CD5^{Hi} cells were substantially lower for CD25, indicating that they had already

progressed through β-selection. Although as expected there were far fewer cells in the Lef1^{Hi}CD5^{Hi}

population upon ACY1215 treatment, they also exhibited the reduction in CD25, suggesting that

ACY1215-treated cells can precociously differentiate beyond β-selection in the absence of CD5

expression.

To confirm the relationship between pre-TCR and CD5 and to assess the impact of ACY121, we

cultured DN3a or DN3b cells from adult thymus in the presence of ACY1215 or DMSO for 2 days,

gated for DN3b cells, and compared CD5 expression to icTCRβ (**Figure 7B**). Cells from the DN3a

starting population were more likely to be dull for both CD5 and icTCRβ, compared with the more

mature cells from the DN3b starting population. However, in these mature cells, expression of

icTCRβ correlated with expression of CD5, suggesting that, as in other cell types, CD5 expression

might report TCR signalling strength during β-selection. In ACY1215-treated cells, expression of

surface TCRβ was reduced by the DN3b^{Post} stage, and uncoupled from CD5. Similar findings were

obtained using fetal liver cultures and staining for CD5 and icTCRβ (**Supp Fig 13**). The correlation of

TCRβ with CD5, and its uncoupling after ACY1215 treatment, was not observed with CD27 and

CD28 (Figure 2) nor CD2 (Fig 3). Together, a functional link between pre-TCR, Lef1and CD5 is

coordinated at DN3b^{Pre} stage (**Figure 7C**), and is affected by ACY1215.

To assess the functional relationship between TCRβ and CD5, we compared CD5 expression and

proliferation in the presence and absence of ACY1215 (**Figure 8A**). Proliferation was minimal in

most DN3a cells and DN3b^{Pre} cells, but was evident in a portion of DN3b^{Pre}, cells from the DN3b

starting population, and in DN3b^{Post} cells. This proliferation was clearly correlated with CD5

expression in control cells. However, the correlation between CD5 expression and CFSE dilution was

lost in the DN3b^{Pre} cells that had been treated with ACY1215. These data together indicate that CD5

serves as a link between TCRβ expression and proliferation, and that this link is broken by ACY1215

treatment.

Discussion

A critical factor in β-selection is the sequential expression of specific transcriptional regulators, with

differential perdurance and cross-inhibition ensuring checks and balances that prevent inappropriate

survival and differentiation (Chann and Russell, 2019; Zhao et al., 2021). Many such checks and

balances have been identified to play important roles during β-selection, but the stepwise progression

of events that dictate fate have been elusive (Chann and Russell, 2019; Zhao et al., 2021). Our

explorations into effects of an HDAC6 inhibitor have revealed sa new working model (**Figure 8C,**

Supp Fig 14), Once DN3b^{Pre} cells have entered the β-selection checkpoint, they respond to graded

levels of TCRβ by expressing Lef1, then CD5. Normally, proliferation only occurs in cells expressing

CD5. Passing of the β-selection checkpoint is then marked by the expression of CD2. The disruption

in TCRβ-associated expression of CD5 by ACY1215 provides a possible mechanism for the escape of

cells lacking an appropriately rearranged TCRβ chain following ACY1215 treatment, although such a

role remains to be investigated.

The upregulation of CD27 and CD28 is gradual, but the upregulation of CD2 is more abrupt, making

for a convenient demarcation of differentiation stages. Our findings are compatible with previous

observations that CD2 expression marks cells with TCRβ gene rearrangement (Rodewald et al.,

1993), that knockout of CD2 inhibits progression to the DN4 stage but does not influence TCRβ

repertoire (Sasada and Reinherz, 2001), and suggestions of cis-regulation of CD2 and TCRβ

expression involving chromatin structure (Kamoun et al., 1995) Together, these findings suggest that

expression of CD2 (marking entrance to the DN3b^{Post} stage) reflects a completion of TCRβ

recombination and triggers critical fate-determining events that involve progressive upregulation of

CD5 and Lef1. CD2 in mature T cells sets quantitative thresholds for activation (Bachmann et al.,

1999), suggesting the possibility that a similar process could be involved in β-selection.

Ideas and Speculation

A novel finding from this work is that CD5 levels are modified in association with pre-TCR

expression, and that treatment with ACY1215 prevents the upregulation of CD5 (but has a lesser

effect on Lef1 upregulation) during β-selection. At other stages of T cell development and activation,

CD5 reports on the strength of TCR signalling, with increasing expression reflecting a stronger TCR

signal (Azzam et al., 1998; Sood et al., 2021; Tarakhovsky et al., 1995). Our findings that the

correlation of CD5 with TCRβ expression is lost with ACY1215 treatment, but the correlation of CD5

expression with proliferation is retained with ACY1215 treatment, suggest the possibility that CD5

plays a similar role as a reporter of pre-TCR signalling. Interestingly, it is now emerging that

differential CD5 expression also marks the propensity for different fates in thymocytes and CD4$^+$ T

cells, and this reflects transcriptional and epigenetic differences between cells with high and low CD5

expression (Rogers et al., 2021). In other cell types the combined action of responding to, and

dampening, the TCR signal, allows CD5 to tune the TCR signal and so shift the window and modify

the timing of positive selection (Lutes et al., 2021; Matson et al., 2020; Sood et al., 2021; Voisinne et

al., 2018). Whether or not such a tuning role might occur for CD5 during β-selection needs further

exploration, but a possible role for tuning of the TCRβ repertoire during β selection is made relevant

by findings that the pre-TCR can bind with low affinity to MHC-peptide (Li et al., 2021; Mallis et al.,

2015). The pre-TCR also forms an immunological synapse with many similarities to the

immunological synapse formed by the αβTCR, oriented towards MHC (Allam et al., 2021). More

work is needed to determine any functional connection between CD5 and tuning, or epigenetic

modifications during β-selection, but these data point to a role for CD5 during the DN3b^{Pre} stage in

reporting pre-TCR activity to determine subsequent passage through the β-selection checkpoint.

The molecular basis by which ACY1215 mediates the effects on cell fate we observe here needs

more exploration. There are many possible means by which ACY1215 might impact upon

transcriptional and epigenetic alterations during β-selection to mediate the phonotypes we observe

here. At β-selection, transient activation of β-catenin mediates TCF-1 and Lef1 transcriptional

regulation to promote pre-TCR-dependent transition to the DN4 stage (Xu et al., 2009). Ectopic

expression of a constitutively active β-catenin allows the cells to bypass pre-TCR signalling, yielding

DN4 cells lacking expression of TCRβ (Gounari et al., 2001). The similarities between this phenotype

and the apparent bypass of pre-TCR signalling we observe with ACY1215 treatment suggest TCF-

1/Lef might mediate the effects of ACY1215. Indeed, HDAC6 can regulate acetylation of β-catenin to

coordinate its degradation, providing a possible mechanism for the effects of ACY1215 on TCF-1/Lef

activity (Iaconelli et al., 2015). Given the correlation between Lef1 expression and CD5 we observe,

CD5 might be downstream of Lef1 during β-selection, perhaps explaining the ACY1215-mediated

deregulation of CD5 expression. These findings suggest that the induction of Lef1 at the DN3b^{Pre}

stage is coordinated with pre-TCR signalling, in part to promote CD5 expression, and that this activity

can be altered by ACY1215 treatment. Whether the effect of ACY1215 relates to HDAC6, an

alternative HDAC, or the histone deacetylase activity of Lef1/TCF-1 remains to be determined.

Other molecular mediators downstream of ACY1215 effects are also possible. It was recently shown

that the transcriptional repressor, Bcl6, is induced by pre-TCR signalling and required for passage

through the β-selection checkpoint (Solanki et al., 2020). Differentiation of T follicular helper cells

also requires Bcl6, in this case induced by Lef1 and TCF (Choi et al., 2015). Several HDACs have

been shown to complex with and modulate Bcl6 activity to impact on B cell development and

function (Wang et al., 2020). Together with our findings, these studies suggest the possibility that

ACY1215 might mediate at least some of its effects via Bcl6. HDAC6 has also been implicated in the

DNA damage response (Zhang et al., 2019), raising the possibility that ACY1215 effects might relate

to the genetic recombination integral to the creation of a TCRβ chain (Miyazaki et al., 2008). In any

case, the functional outcome of ACY1215 treatment is to disrupt the connections between pre-TCR

signalling and expression of CD5 expression and CD5-associated proliferation. Uncoupling CD5 from

TCRβ and Lef1 expression provides a possible mechanism by which ACY1215 allows the wrong

cells (perhaps cells without suitable TCRβ) to proliferate and differentiate to DN3b^{Post} cells, but these

cells subsequently fail to progress, leading to the block in DP differentiation.

The known role for HDAC6 in linking tubulin acetylation to the immunological synapse of mature T

cells (Serrador et al., 2004), and our finding that acetylation of α-tubulin is increased in DN3 cells,

suggested the possibility that the β-selection immunological synapse might mediate the effects of

ACY125. Indeed, ACY1215 does alter the organisation of the TCR signalling molecules, LAT,

around the MTOC. Similarly, Notch1, which is required to enhance the polarisation of the pre-TCR

(Allam et al., 2021; Charnley et al., 2020), is lost from the MTOC following ACY1215 treatment.

These findings are particularly interesting in light of a recent report that HDAC6 is required for the

formation of aggresomes at the MTOC (Magupalli et al., 2020), and suggest that HDAC6 might play

a more general role in coordinating recruitment to the MTOC. An alternative explanation for the

effect of ACY1215 on β-selection relates to observations that HDAC inhibitors can induce MHC

constituents to promote antigen presentation (Magner et al., 2000). It remains to be determined

whether an influence on stromal-presented peptide-MHC plays any role in disrupting β-selection.

Interestingly, both CD2 and CD5 can regulate activity of the immunological synapse and connectivity

to the MTOC (Brossard et al., 2003; Dustin et al., 1998; Tibaldi et al., 2002) further supporting the

possibility that the immunological synapse provides a hub at which deacetylation and pre-TCR

signalling converge. Together, these findings suggest that the effect of HDAC6 on β-selection that we

see here might occur through disruption of the DN3 immunological synapse.

The molecular basis for the effects of ACY1215 on β-selection might therefore by multifaceted, and

relate to one or more of direct alteration of the epigenome to influence protein expression, alteration

of Lef1 activity, DNA damage response, alteration of pre-TCR signalling via organisation at the

MTOC and the immunological synapse. Irrespective of the molecular mechanism, the finding that

AY1215 disrupts T cell development has potential implications for the development of clinical

applications of small molecule inhibitors of HDAC6, particularly in treatment of young patients,

whose T cell repertoire has not yet been established (Cosenza and Pozzi, 2018; Hogg et al., 2020).

Materials and Methods

***Ex vivo* culture of developing T cells and drug treatments**

Two sources to derive developing T cells were employed in this study: mouse hematopoietic stem

cells (HSCs) and mouse primary thymocytes. HSCs were collected from C57BL/6 mouse E14.5

fetal liver. To induce *ex vivo* T cell development, HSCs were co-cultured with OP9-DL1 stromal

cells (from Juan Carlos Zúñiga-Pflücker, University of Toronto, Toronto, Canada) at a 1:1 ratio in a

6-well plate (2×10^5). The coculture was maintained in αMEM (Sigma-Aldrich, M4526),

supplemented with foetal calf serum (10% v/v), glutamine (1 mM), β-mercaptoethanol (50 μM),

sodium pyruvate (1nM), HEPES (10 mM), penicillin/streptomycin (100 ng/mL), mouse interleukin 7

(1 ng/mL) and mouse FMS-like tyrosine kinase 3 (5 ng/ml). C57BL/6 mice at 4-5 week were

euthanised for thymus collection. The primary thymocytes were released from thymus for sorting,

analysis or further culturing with OP9-DL1 stromal cells under the same culture conditions as the

HSC system described above. For both sources of developing T cells, ACY1215, purchased from

MedChemExpress (NJ, USA), was used at 1 μM as described (Laino et al., 2019). ACY1215 was

dissolved in DMSO to 5 mM, and diluted to 1 μM with culture media immediately prior to application

to cells. The small-molecule inhibitor of Stat5, Pimozide (CAS 2062-78-4 Sigma-Aldrich) was used

at a final concentration of 10 μM. The selective inhibitor of p300 and CBP, C646 (Sigma-Aldrich)

was used at 2.5 μM. OP9-DL1 stromal cells was maintained in αMEM (Sigma-Aldrich, M4526)

supplemented with 20% (v/v) fetal calf serum, glutamine (1 mM) and penicillin/streptomycin (100

ng/mL). When OP9-DL1 stromal cells were confluent after 3-4 days of coculture, primary thymocytes

were dissociated from the stromal cells through forceful pipetting, and were harvested with the fresh

OP9-DL1 stromal cells.

Flow Cytometry and FACS

To purify T cell populations using flow cytometry, primary thymocytes were dissociated from OP9-

DL1 stromal cell, and were resuspended in PBS buffer containing 2% FBS and antibody cocktail for

surface staining. Antibody cocktails were comprised of lineage markers (NK-1.1, CD11b, CD45R,

Ly6G6C, Ter119), and differentiation markers (CD4, CD8, CD25, CD44, CD28 and CD2; details

about fluorochrome, manufacturer and dilution are summarised in the supplementary materials).

Primary thymocytes were incubated with antibody cocktails for 40 minutes before FACS.

Flow cytometry (BD Aria III) was used to analyse and sort the populations of developing T cells.

DN3 cells were identified based on surface expression (Lineage$^-$, CD4$^-$, CD8$^-$, CD25^{Hi}, CD44^{Lo}).

DN3a cells were DN3 cell with CD28^{L0}, by contrast, DN3b cells were DN3 cell with CD28^{Hi}. DP

cells were identified based on surface expression (Lineage$^-$, CD4$^+$, CD8$^+$).

Intracellular staining for flow cytometry

To detect intracellular expression of TCRβ, Ki67, Lef1, c-Myc, Notch1, HDAC6 and H3K18ac, the

Foxp3/Transcription Factor Staining Buffer Kit (eBioscience) was used according to the

manufacturer's instruction. Briefly, the primary thymocytes after surface staining were incubated with

fixative for 15 minutes in dark, room temperature, followed with centrifuge and washing out the

fixative. To detect expression of acetyl-tubulin, the cells were fixed with IC Fixation Buffer

(eBioscience) for 15 minutes in dark, room temperature, followed with centrifuge and washing out the

fixative. The fixed cells were then incubated with antibody for flow cytometry analysis.

Immunofluorescence staining and confocal microscopy

The sorted DN3a or DN3b were grown on glass-bottomed 8-well chambers (ibidi) with OP9-DL1

stromal cells, and were fixed with 4% formaldehyde for 8 minutes at room temperature followed by

0.1% Triton X-100 permeabilization for 5 minutes. The samples were then incubated with primary

antibody overnight at 4°C (details about antibody usage is summarised in Supplementary table 1).

After the primary antibody being washed out by PBS, the cells were labelled with fluorochrome-

conjugated secondary antibodies (as Table S1), DAPI (Thermo Fisher Scientific), and Phalloidin

(Abcam). The samples were examined using a FV3000 confocal microscope (Olympus) and 60X lens

(1.30 NA, UPLSAPO, Olympus). The images were processed with maximum intensity projection

using Image J. The mean intensity of region of interest was examined in Image J.

Data collection and Statistical analyses

482   Python (version 3.8.1) was used to visualise the statistics. Data of flow cytometry were analysed by

483   FlowJo software (version X10.0.7r2. Tree Star, Inc.). The bar plot was shown as mean ± SEM. More

484   than three repeats were included in the statistics and were colour-coded for each individual repeat.

485   Microsoft Excel (version 2016) was used for two sided t-test, by which p value was indicated.

Acknowledgements

This work was done on Wurundjeri land of the Kulin nation, and we pay our respects to the Elders

past and present. We thank Louise Cheng, Callum Dark, Amr Allam and Peter Fraser for technical

assistance and helpful discussions, and Ellis Reinherz, Jonathon Duke-Cohan for helpful comments

on the manuscript. The work was funded by Australian Research Council (FT0990405 to SMR),

National Health and Medical Research Council (APP1099140 to SMR), Swinburne University

Postgraduate Research Award to ASC.

Competing Interest Statement

No author declares any competing interests

Author Contribution Statement

Conceptualization: ASC, SMR
Methodology: ASC, MC, SMR
Experimentation: ASC
Visualization: ASC, SMR
Supervision: POH, RWJ, SMR
Writing—original draft: ASC, SMR
Writing—review & editing: ASC, MC, LN, AN, POH, RWJ, SMR

Ethics

C57bl/6 mouse E14.5 fetal livers and 4-5-week mouse thymi were collected according to

Peter MacCallum Cancer Centre Ethics Committee approval E627 and La Trobe University

Animal Ethics Committee approval AEC-20024.

Figure Legends

Figure 1. ACY1215 inhibits T cell development at the DN3 stage

(A) FL-derived developing T cells were treated with ACY1215 or DMSO for 2 days as indicated, and

the Lin⁻ cells were assessed for CD4 and CD8 expression to determine the proportions of DN, DP,

CD8⁺ and CD4⁺ populations. The DN (Lin⁻, CD4⁻, CD8⁻) cells were assessed for CD44 and CD25

expression to determine the proportions of DN3 and DN4 populations. The frequency of each

population was summarised in a swarm plot, with individual colours representing each experiment.

Sorted, FL-derived DN3a cells **(B)** or thymocytes **(C)** were cultured under ACY1215 or DMSO

treatment for 1 or 2 days as indicated, and the CD44⁻ DN population (representing DN3 and DN4

stages) was assessed for CD25 and CD28 expression. The frequency of each population were

summarised in swarm plots, with individual colours representing each experiment. Cell numbers were

summarised in line plots.

(D) Sorted, FL-derived DN3b cells were cultured for 4 days under ACY1215 or DMSO treatment as

indicated, and cell numbers per population were estimated by multiplying the total cell number by the

frequency and were summarised in the line chart.

Figure 2. ACY1215 inhibits β-selection.

(A) Developing T cells from FL **(i)** or the adult thymus **(ii)** were treated with ACY1215 or DMSO for

1 day or 2 days, and the DN3 cells were assessed for CD27 and CD28 expression to discriminate

DN3a and DN3b. Cell number or frequency were summarised in the swarm plot, with individual

colours representing each experiment.

(B) Sorted, FL-derived DN3a cells **(i)** were cultured under ACY1215 or DMSO treatment for 4 days

as indicated, and DN3 cells were assessed for the expressions of intracellular TCRβ (icTCRβ) versus

CD28 or CD27. Thymocytes **(ii)** were cultured under ACY1215 or DMSO treatment for 1 day as

indicated, and DN3 cells were assessed for the expressions of icTCRβ versus CD28. The frequency of

icTCRβ⁺ cells was summarised in a swarm plot, with individual colours representing each experiment.

Figure 3. CD2 upregulation discriminates DN3b^{Pre} and DN3b^{Post} populations

(A) The strategy for discriminating between three populations (DN3a, DN3b^{Pre}, and DN3b^{Post}) is

indicated in the schematic.

(B) FL-derived developing T cells **(i)** or the sorted thymocyte DN3 **(ii)** were treated with ACY1215

or DMSO for 1 or 2 days, and the DN3 cells were assessed for CD2 and CD28 expression.

(C) FL-derived developing T cells **(i)** or the sorted thymus-derived DN3 **(ii)** were treated with

ACY1215 or DMSO for 1 day, and DN3a, DN3b^{Pre}, and DN3b^{Post} cells were assessed for icTCRβ and

Ki-67 expression. MFI of icTCRβ and Ki-67 were summarised in histogram and swarm plot, with

individual colours representing each experiment. (p value < 0.001 = '***', < 0.01 = '**', <0.05 ='*')

Figure 4. DN3a, DN3b^{Pre} and DN3b^{Post} stages follow a sequential progression.

The DN3a, DN3b^{Pre}, and DN3b^{Post} cells from the adult thymus were sorted as indicated, and were

cultured on OP9-DL1. At two day, the cells were analysed by CD4 versus CD8, the DN cells were

analysed by CD44 versus CD25, and the DN3 cells were analysed by CD2 versus CD28. The

frequencies of each population were summarised in the line plots. (p value < 0.001 = '***', < 0.01 =

'**', <0.05 ='*')

Figure 5. ACY1215 alters pre-TCR signalling clustered at the MTOC.

(A) Sorted FL-derived DN, DP, CD4-SP, CD8-SP, DN1, DN2, DN3a, DN3b were treated with

ACY1215 or DMSO for 2 hours as indicated, and were assessed for acetyl-α-tubulin (K40) as shown

in histograms.

(B, C) Sorted FL-derived DN3a cells were treated with ACY1215 or DMSO for 1 day as indicated,

followed by immunostaining for LAT, acetyl-α-tubulin (K40), Notch1 and α-tubulin. The mean

intensity of acetyl-α-tubulin, LAT and the ratio of Notch1 expressed at MTOC versus at nucleus were

summarized in violin plot overlaid with box plot. Cell number for LAT staining: ACY1215 (n=55)

and DMSO (n=54); Cell number for acetyl-α-tubulin (K40) staining: ACY1215 (n=30) and DMSO

(n=31); Cell number for Notch1 staining: ACY1215 (n=56) and DMSO (n=55).

(D) FL-derived developing T cells were treated with ACY1215 or DMSO for 2 day, and the DN3a,

DN3b and DN4 cells were assessed for Notch1 expression, as summarised in swarm plot, with

individual colours representing each experiment.

Figure 6. Coordinated upregulation of Lef1 and TCRβ in DN3b^{Pre} cells.

(A) FL-derived developing T cells **(i)** or the sorted thymocyte DN3 **(ii)** were treated with ACY1215 or

DMSO for 1 day as indicated, and DN3 cells were assessed for CD2 and CD28 expression. The

DN3a, DN3b^{Pre}, and DN3b^{Post} cells were assessed for Lef1 expression.

(B) DN3a, DN3b^{Pre}, and DN3b^{Post} cells from (A, i) were assessed for Lef1 and icTCRβ expression.

Figure 7. CD5 is upregulated with Lef1 and TCRβ in DN3b^{Post} cells.

(A) Sorted DN3 cells from the adult thymus were treated with ACY1215 or DMSO for 1 day as

indicated, and DN3 cells were assessed for CD5, CD2 and Lef1 expression.

(B) Sorted DN3a and DN3b^{Pre} cells from the adult thymus were treated with ACY1215 or DMSO for

2 day as indicated, and DN3 populations were assessed for CD5 and surface TCRβ expression.

(C) Schematic describes the stepwise expression of TCR co-receptors (CD28, CD5, and CD2) and

Lef1 from DN3a, DN3b^{Pre} to DN3b^{Post} stage.

Figure 8. CD5 upregulation correlates with the passing of β-selection checkpoint

(A) Sorted FL-derived DN3a and DN3b were stained with CFSE after sorting and were cultured with

ACY1215 or DMSO for two days as indicated. At two days, the DN3a, DN3b^{Pre}, and DN3b^{Post} cells

from the sorted DN3a and DN3b were assessed for CD5 expression versus CFSE intensity.

(B) Schematic summarises the working model of molecular interactions at the DN3b^{Pre} stage. Firstly,

CD28 (the defining step in transition from DN3a to DN3b^{Pre}) acts to promote signalling through the

nascent pre-TCR. Second, pre-TCR signals induce expression of CD5 (or *vice versa*) and lead to the

passing of the β-selection checkpoint. By coordinating transcriptional control, functional MTOC and

immunological synapse formation for survival, proliferation and differentiation. Possible molecular

events disrupted by ACY1215 treatment to cause these cellular responses are indicated by the red

double lines.

Figure S1.

(A) A schematic of the stages of T cell development and the cell surface markers used to differentiate

them. Cells were classed as DN (Double Negative, lacking expression of CD4 and CD8), and

progressed from DN1 to DN4 according to their expression of CD44 and CD25 as indicated. β-

selection occurs at the DN3 stage, as cells progress from DN3a to DN3b, indicated by upregulation of

CD28. Cells post β-selection progressed from DN4 to the DP stage (Double Positive, expressing both

CD4 and CD8) before committing to SP (Single Positive for either CD4 or CD8).

(B) Fetal liver cells were collected from E14.5 mice embryos, and cells cultured on OP9-D1L stromal

cells for 8-10 days. The developing T cells were demarcated as the flow indicated.

(C) Thymocytes were collected from 4-to-5-week mice thymus **(i)** and/or were cultured *ex vivo* on

OP9-DL1 stromal cells **(ii)**. The developing T cells were demarcated as the flow indicated.

Figure S2.

Sorted, FL-derived DN2, DN3a, DN3b and DP were cultured for 2 days ACY1215 or DMSO

treatment as indicated. Cell numbers were counted at 3 hours and 2 days, and the proliferation rate (2

days / 3 hours) was summarised in swarm plot, with individual colours representing independent

experiments.

Figure S3.

(A) Sorted, FL-derived DN3a and DN3b were stained with CFSE after sorting and were cultured with

ACY1215 or DMSO for two days as indicated. At one hour post sorting, the intensity of CFSE were

recorded, as indicated by the white histogram. At two days, the DN3 cells from the sorted DN3a and

DN3b were assessed for CD28 expression and CFSE intensity, as shown in the contour plots and the

histograms.

(B) FL-derived developing T cells were treated with ACY1215 or DMSO for 1 day as indicated, and

DN3a and DN3b cells were assessed for FSC and Ki-67 expression.

Figure S4.

(A) Sorted, FL-derived DN3a, DN3b and DP cells were treated with ACY1215 or DMSO for 3 hours

or 6 hours as indicated, followed with propidium iodide (PI) staining to label dead cells. The

frequency of PI^+ population were summarised in swarm plot, with individual colours representing

each experiment.

(B) Sorted, FL-derived DN3a and DN3b cells were treated with ACY1215 or DMSO for 6 hours or 1

day with or without OP9-DL1 coculture as indicated. The cells were assessed for Annexin V

expression and DAPI intensity as shown in the density plot.

Figure S5.

Sorted, FL-derived DN3a **(A)** and sorted, FL-derived DN3b **(B)** were cultured for 4 days under

ACY1215 or DMSO treatment as indicated, and were analysed by CD4 and CD8 **(i)** and CD44 and

CD25 **(ii)**. The frequency of each population was summarised in swarm plot, with individual colours

representing independent experiments.

Figure S6.

(A) FL-derived developing T cells were treated with ACY1215 or DMSO for 2 days as indicated, and

DN3a, DN3b, DN4 and DP cells were assessed for surface TCRβ and surface TCRγδ expression. **(B)**

Sorted, FL-derived DN3b cells were cultured for 2 days under ACY1215 or DMSO treatment as

indicated, and DN3, DN4 and DP cells were assessed for surface TCRβ and surface TCRγδ

expression. **(C)** Sorted FL-derived DN3b cells were cultured for 4 days under ACY1215 or DMSO

treatment as indicated, and DN3, DN4 and DP cells were assessed for intracellular TCRβ (icTCRβ)

and CD28 expression. The frequency and cell number were summarised in swarm plot, with

individual colours representing each experiment.

Figure S7.

(A) CD28 upregulation marks both DN3 cells differentiation and passing β-selection. FL-derived

developing T cells at around day 8-10 of coculture with OP9-DL1 stromal cells were assessed for

CD28, CD71, CD98, CD27 and surface TCRβ expression.

(B) Sorted, FL-derived DN3b **(i)** or thymocytes **(ii)** were treated with ACY1215 or DMSO for 1 day

or 2 days as indicated, and the DN3 cells were assessed for surface TCRβ and icTCRβ expression.

MFI of surface TCRβ from icTCRβ$^+$ cells was summarised in a swarm plot, with individual colours

representing each experiment.

Figure S8

(A) Quantification for Figure 3A. Frequency and cell number of the three populations (DN3a, DN3b^{Pre}, and DN3b^{Post}) was summarised in swarm plots, with individual colours representing each experiment.

(B) Sorted, FL-derived DN3a and DN3b were stained with CFSE after sorting and were cultured with ACY1215 or DMSO for two days as indicated. At two days, the DN3 cells from the sorted DN3a and DN3b were assessed for CD2 expression versus CFSE intensity.

(C) DN3a, DN3b^{Pre}, and DN3b^{Post} thymocytes that treated with ACY1215 for 1 day were assessed for CD27 and surface TCRβ expression.

 (D) FL-derived developing T cells were treated with ACY1215 or DMSO for 2 days, and DN3a, DN3b^{Pre}, and DN3b^{Post} cells were assessed for c-Myc expression.

Figure S9

FL-derived developing T cells were treated with DMSO or ACY1215 for 1.5 day, and DN3a, DN3b^{Pre}, and DN3b^{Post} cells were sorted as indicated **(i)**. At one day, the cells were analysed by CD4 versus CD8 **(ii)** and CD2 versus CD28 **(iii)**. The frequencies of each population were summarised in the line plots. (p value < 0.001 = '***', < 0.01 = '**', <0.05 ='*')

Figure S10.

HDAC6 is expressed in developing T cells. **(A)** To examine HDAC6 protein expression, FL-derived developing T cells treated with DMSO or ACY1215 for two days were fixed by cytoplasmic fixative or nuclear fixative, followed by examination of HDAC6 expression. **(B)** FL-derived developing T cells were treated with ACY1215, the selective inhibitor of p300 and CBP, C646, or DMSO for 1 day as indicated, and the expression of H3K18ac were assessed and shown in the histogram. **(C)** Sorted,

FL-derived DN3a or DN3b cells were assessed for HDAC6 (in white colour) expression by

immunostaining. **(D, E)** Sorted, FL-derived DN3b cells were treated with ACY1215 or DMSO for 1

day, followed by immunostaining for α-tubulin, Arp2 (in white colour), acetyl-α-tubulin (K40) (in

white colour).

Figure S11.

FL-derived developing T cells were treated with ACY1215 or DMSO for 1 day as indicated, and

DN3a, DN3b^{Pre}, and DN3b^{Post} cells were assessed for Lef1 and CD2 expression. DN3b^{Pre} cells were

further characterised by Lef1^{Hi} or Lef1^{Lo}, and were assessed for Lef1 and CD5 expression with DN3a

and DN3b^{Post}.

Figure S12.

(A) Sorted FL-derived DN3b^{Pre} cells were accessed for CD25 expression as schematic indicated.

(B) FL-derived developing T cells were treated with ACY1215 or DMSO for 2 days as indicated, and

DN3a and DN3b were assessed for Lef1 and CD5 expression. The expression of CD25 in the DN3b

subsets were shown in histogram.

Figure S13.

Sorted FL-derived DN3a and DN3b cells were treated with DMSO or ACY1215 for 2 days, and were

assessed for CD5, TCRβ and CD25 expression.

Figure S14.

Schematic shows the sequence of protein expression changes during development from DN3a,

DN3b^{Pre} to DN3b^{Post}.

Adeegbe, D.O., Y. Liu, P.H. Lizotte, Y. Kamihara, A.R. Aref, C. Almonte, R. Dries, Y. Li, S. Liu, X. Wang,
T. Warner-Hatten, J. Castrillon, G.C. Yuan, N. Poudel-Neupane, H. Zhang, J.L. Guerriero, S.
Han, M.M. Awad, D.A. Barbie, J. Ritz, S.S. Jones, P.S. Hammerman, J. Bradner, S.N. Quayle,
and K.K. Wong. 2017. Synergistic Immunostimulatory Effects and Therapeutic Benefit of
Combined Histone Deacetylase and Bromodomain Inhibition in Non-Small Cell Lung Cancer.
Cancer discovery 7:852-867.
Allam, A.H., M. Charnley, K. Pham, and S.M. Russell. 2021. Developing T cells form an immunological
synapse for passage through the beta-selection checkpoint. *J Cell Biol* 220:
Ananias, H.J., I.J. de Jong, R.A. Dierckx, C. van de Wiele, W. Helfrich, and P.H. Elsinga. 2008. Nuclear
imaging of prostate cancer with gastrin-releasing-peptide-receptor targeted
radiopharmaceuticals. *Current pharmaceutical design* 14:3033-3047.
Azzam, H.S., J.B. DeJarnette, K. Huang, R. Emmons, C.-S. Park, C.L. Sommers, D. El-Khoury, E.W.
Shores, and P.E. Love. 2001. Fine Tuning of TCR Signaling by CD5. *The Journal of Immunology*
166:5464-5472.
Azzam, H.S., A. Grinberg, K. Lui, H. Shen, E.W. Shores, and P.E. Love. 1998. CD5 expression is
developmentally regulated by T cell receptor (TCR) signals and TCR avidity. *J Exp Med*
188:2301-2311.
Bachmann, M.F., M. Barner, and M. Kopf. 1999. CD2 sets quantitative thresholds in T cell activation.
J Exp Med 190:1383-1392.
Ballesteros-Arias, L., J.G. Silva, R.A. Paiva, B. Carbonetto, P. Faisca, and V.C. Martins. 2019. T Cell
Acute Lymphoblastic Leukemia as a Consequence of Thymus Autonomy. *J Immunol*
Brossard, C., M. Semichon, A. Trautmann, and G. Bismuth. 2003. CD5 inhibits signaling at the
immunological synapse without impairing its formation. *J Immunol* 170:4623-4629.
Chann, A.S., and S.M. Russell. 2019. An integrated transcriptional switch at the beta-selection
checkpoint determines T cell survival, development and leukaemogenesis. *Biochemical*
Society transactions 47:1077-1089.
Charnley, M., M. Ludford-Menting, K. Pham, and S.M. Russell. 2020. A new role for Notch in the
control of polarity and asymmetric cell division of developing T cells. *J Cell Sci* 133:jcs235358.
Choi, Y.S., J.A. Gullicksrud, S. Xing, Z. Zeng, Q. Shan, F. Li, P.E. Love, W. Peng, H.H. Xue, and S. Crotty.
2015. LEF-1 and TCF-1 orchestrate T differentiation by regulating differentiation circuits
upstream of the transcriptional repressor Bcl6. *Nat Immunol*
Cosenza, M., and S. Pozzi. 2018. The Therapeutic Strategy of HDAC6 Inhibitors in
Lymphoproliferative Disease. *International journal of molecular sciences* 19:
Dose, M., I. Khan, Z. Guo, D. Kovalovsky, A. Krueger, H. von Boehmer, K. Khazaie, and F. Gounari.
2006. c-Myc mediates pre-TCR-induced proliferation but not developmental progression.
Blood 108:2669-2677.
Douek, D.C., R.A. Vescio, M.R. Betts, J.M. Brenchley, B.J. Hill, L. Zhang, J.R. Berenson, R.H. Collins,
and R.A. Koup. 2000. Assessment of thymic output in adults after haematopoietic stem-cell
transplantation and prediction of T-cell reconstitution. *Lancet* 355:1875-1881.
Dustin, M.L., M.W. Olszowy, A.D. Holdorf, J. Li, S. Bromley, N. Desai, P. Widder, F. Rosenberger, P.A.
van der Merwe, P.M. Allen, and A.S. Shaw. 1998. A novel adaptor protein orchestrates
receptor patterning and cytoskeletal polarity in T-cell contacts. *Cell* 94:667-677.

Dutta, A., B. Zhao, and P.E. Love. 2021. New insights into TCR β-selection. *Trends in Immunology*

Eich, C., J. Arlt, C.S. Vink, P. Solaimani Kartalaei, P. Kaimakis, S.A. Mariani, R. van der Linden, W.A. van
Cappellen, and E. Dzierzak. 2018. In vivo single cell analysis reveals Gata2 dynamics in cells
transitioning to hematopoietic fate. *The Journal of Experimental Medicine* 215:233-248.

Gounari, F., I. Aifantis, K. Khazaie, S. Hoeflinger, N. Harada, M.M. Taketo, and H. von Boehmer. 2001.
Somatic activation of beta-catenin bypasses pre-TCR signaling and TCR selection in
thymocyte development. *Nat Immunol* 2:863-869.

Goux, D., J.D. Coudert, D. Maurice, L. Scarpellino, G. Jeannet, S. Piccolo, K. Weston, J. Huelsken, and
W. Held. 2005. Cooperating pre-T-cell receptor and TCF-1-dependent signals ensure
thymocyte survival. *Blood* 106:1726-1733.

Groettrup, M., A. Baron, G. Griffiths, R. Palacios, and H. von Boehmer. 1992. T cell receptor (TCR)
beta chain homodimers on the surface of immature but not mature alpha, gamma, delta
chain deficient T cell lines. *EMBO J* 11:2735-2745.

Hayday, A.C., and D.J. Pennington. 2007. Key factors in the organized chaos of early T cell
development. *Nat Immunol* 8:137-144.

Hogg, S.J., P.A. Beavis, M.A. Dawson, and R.W. Johnstone. 2020. Targeting the epigenetic regulation
of antitumour immunity. *Nature reviews. Drug discovery*

Holmes, R., and J.C. Zúñiga-Pflücker. 2009. The OP9-DL1 system: generation of T-lymphocytes from
embryonic or hematopoietic stem cells in vitro. *Cold Spring Harb Protoc* 2009:pdb.prot5156.

Iaconelli, J., J.H. Huang, S.S. Berkovitch, S. Chattopadhyay, R. Mazitschek, S.L. Schreiber, S.J.
Haggarty, and R. Karmacharya. 2015. HDAC6 inhibitors modulate Lys49 acetylation and
membrane localization of beta-catenin in human iPSC-derived neuronal cells. *ACS Chem Biol*
10:883-890.

Kamoun, M., J.S. Woods, N. Sano, H. Makni, R. Smith, W.B. de Lau, A. van Oers, D. Wotton, M.J.
Owen, Y. Hashimoto, and H.C. Clevers. 1995. Analysis of CD2 and TCR-beta gene expression
in Jurkat cell mutants suggests a cis regulation of gene transcription. *J Immunol* 155:3929-
3937.

Laino, A.S., B.C. Betts, A. Veerapathran, I. Dolgalev, A. Sarnaik, S.N. Quayle, S.S. Jones, J.S. Weber,
and D.M. Woods. 2019. HDAC6 selective inhibition of melanoma patient T-cells augments
anti-tumor characteristics. *Journal for immunotherapy of cancer* 7:33.

Li, X., R. Mizsei, K. Tan, R.J. Mallis, J.S. Duke-Cohan, A. Akitsu, P.W. Tetteh, A. Dubey, W. Hwang, G.
Wagner, M.J. Lang, H. Arthanari, J.H. Wang, and E.L. Reinherz. 2021. Pre-T cell receptors
topologically sample self-ligands during thymocyte beta-selection. *Science* 371:181-185.

López-Rodríguez, C., J. Aramburu, and R. Berga-Bolaños. 2015. Transcription factors and target genes
of pre-TCR signaling. *Cellular and Molecular Life Sciences* 72:2305-2321.

Lutes, L.K., Z. Steier, L.L. McIntyre, S. Pandey, J. Kaminski, A.R. Hoover, S. Ariotti, A. Streets, N. Yosef,
and E.A. Robey. 2021. T cell self-reactivity during thymic development dictates the timing of
positive selection. *eLife* 10:e65435.

Magner, W.J., A.L. Kazim, C. Stewart, M.A. Romano, G. Catalano, C. Grande, N. Keiser, F. Santaniello,
and T.B. Tomasi. 2000. Activation of MHC class I, II, and CD40 gene expression by histone
deacetylase inhibitors. *J Immunol* 165:7017-7024.

Magupalli, V.G., R. Negro, Y. Tian, A.V. Hauenstein, G. Di Caprio, W. Skillern, Q. Deng, P. Orning, H.B.
Alam, Z. Maliga, H. Sharif, J.J. Hu, C.L. Evavold, J.C. Kagan, F.I. Schmidt, K.A. Fitzgerald, T.
Kirchhausen, Y. Li, and H. Wu. 2020. HDAC6 mediates an aggresome-like mechanism for
NLRP3 and pyrin inflammasome activation. *Science* 369:eaas8995.

Mallis, R.J., K. Bai, H. Arthanari, R.E. Hussey, M. Handley, Z. Li, L. Chingozha, J.S. Duke-Cohan, H. Lu,
J.H. Wang, C. Zhu, G. Wagner, and E.L. Reinherz. 2015. Pre-TCR ligand binding impacts
thymocyte development before alphabetaTCR expression. *Proc Natl Acad Sci U S A*
112:8373-8378.

Matson, C.A., S. Choi, F. Livak, B. Zhao, A. Mitra, P.E. Love, and N.J. Singh. 2020. CD5 dynamically calibrates basal NF-kappaB signaling in T cells during thymic development and peripheral activation. *Proc Natl Acad Sci U S A* 117:14342-14353.

Mingueneau, M., T. Kreslavsky, D. Gray, T. Heng, R. Cruse, J. Ericson, S. Bendall, M.H. Spitzer, G.P. Nolan, K. Kobayashi, H. von Boehmer, D. Mathis, C. Benoist, C. Immunological Genome, A.J. Best, J. Knell, A. Goldrath, V. Jojic, D. Koller, T. Shay, A. Regev, N. Cohen, P. Brennan, M. Brenner, F. Kim, T.N. Rao, A. Wagers, T. Heng, J. Ericson, K. Rothamel, A. Ortiz-Lopez, D. Mathis, C. Benoist, N.A. Bezman, J.C. Sun, G. Min-Oo, C.C. Kim, L.L. Lanier, J. Miller, B. Brown, M. Merad, E.L. Gautier, C. Jakubzick, G.J. Randolph, P. Monach, D.A. Blair, M.L. Dustin, S.A. Shinton, R.R. Hardy, D. Laidlaw, J. Collins, R. Gazit, D.J. Rossi, N. Malhotra, K. Sylvia, J. Kang, T. Kreslavsky, A. Fletcher, K. Elpek, A. Bellemare-Pelletier, D. Malhotra, and S. Turley. 2013. The transcriptional landscape of alphabeta T cell differentiation. *Nat Immunol* 14:619-632.

Miyazaki, M., K. Miyazaki, M. Itoi, Y. Katoh, Y. Guo, R. Kanno, Y. Katoh-Fukui, H. Honda, T. Amagai, M. van Lohuizen, H. Kawamoto, and M. Kanno. 2008. Thymocyte proliferation induced by pre-T cell receptor signaling is maintained through polycomb gene product Bmi-1-mediated Cdkn2a repression. *Immunity* 28:231-245.

Narita, T., B.T. Weinert, and C. Choudhary. 2019. Functions and mechanisms of non-histone protein acetylation. *Nature Reviews Molecular Cell Biology* 20:156-174.

Notarangelo, L.D., R. Bacchetta, J.L. Casanova, and H.C. Su. 2020. Human inborn errors of immunity: An expanding universe. *Science Immunology* 5:eabb1662.

Osseni, A., A. Ravel-Chapuis, J.-L. Thomas, V. Gache, L. Schaeffer, and B.J. Jasmin. 2020. HDAC6 regulates microtubule stability and clustering of AChRs at neuromuscular junctions. *Journal of Cell Biology* 219:

Rodewald, H.R., K. Awad, P. Moingeon, L. D'Adamio, D. Rabinowitz, Y. Shinkai, F.W. Alt, and E.L. Reinherz. 1993. Fc gamma RII/III and CD2 expression mark distinct subpopulations of immature CD4-CD8- murine thymocytes: in vivo developmental kinetics and T cell receptor beta chain rearrangement status. *J Exp Med* 177:1079-1092.

Rogers, D., A. Sood, H. Wang, J.J.P. van Beek, T.J. Rademaker, P. Artusa, C. Schneider, C. Shen, D.C. Wong, M.-È. Lebel, S.A. Condotta, M.J. Richer, A.J. Martins, J.S. Tsang, L. Barreiro, P. Francois, D. Langlais, H.J. Melichar, J. Textor, and J.N. Mandl. 2021. Pre-existing chromatin accessibility and gene expression differences among naïve CD4+ T cells influence effector potential. *bioRxiv* 2021.2004.2021.440846.

Sasada, T., and E.L. Reinherz. 2001. A critical role for CD2 in both thymic selection events and mature T cell function. *J Immunol* 166:2394-2403.

Serrador, J.M., J.R. Cabrero, D. Sancho, M. Mittelbrunn, A. Urzainqui, and F. Sanchez-Madrid. 2004. HDAC6 deacetylase activity links the tubulin cytoskeleton with immune synapse organization. *Immunity* 20:417-428.

Shapiro, M.J., and V.S. Shapiro. 2020. Chromatin-Modifying Enzymes in T Cell Development. *Annu Rev Immunol* 38:397-419.

Skånland, Sigrid S., K. Moltu, T. Berge, Einar M. Aandahl, and K. Taskén. 2014. T-cell co-stimulation through the CD2 and CD28 co-receptors induces distinct signalling responses. *Biochemical Journal* 460:399-410.

Solanki, A., D.C. Yánez, C.-I. Lau, J. Rowell, A. Barbarulo, S. Ross, H. Sahni, and T. Crompton. 2020. The transcriptional repressor Bcl6 promotes pre-TCR-induced thymocyte differentiation and attenuates Notch1 activation. *Development* 147:dev192203.

Sood, A., M.E. Lebel, M. Dong, M. Fournier, S.J. Vobecky, E. Haddad, J.S. Delisle, J.N. Mandl, N. Vrisekoop, and H.J. Melichar. 2021. CD5 levels define functionally heterogeneous populations of naive human CD4(+) T cells. *Eur J Immunol* 51:1365-1376.

Staal, F.J., and H.C. Clevers. 2005. WNT signalling and haematopoiesis: a WNT-WNT situation. *Nat Rev Immunol* 5:21-30.

Taghon, T., M.A. Yui, R. Pant, R.A. Diamond, and E.V. Rothenberg. 2006. Developmental and
molecular characterization of emerging beta- and gammadelta-selected pre-T cells in the
adult mouse thymus. *Immunity* 24:53-64.
Tarakhovsky, A., S.B. Kanner, J. Hombach, J.A. Ledbetter, W. Muller, N. Killeen, and K. Rajewsky.
1995. A role for CD5 in TCR-mediated signal transduction and thymocyte selection. *Science*
269:535-537.
Teague, T.K., C. Tan, J.H. Marino, B.K. Davis, A.A. Taylor, R.W. Huey, and C.J. Van De Wiele. 2010.
CD28 expression redefines thymocyte development during the pre-T to DP transition. *Int*
Immunol 22:387-397.
Tibaldi, E.V., R. Salgia, and E.L. Reinherz. 2002. CD2 molecules redistribute to the uropod during T
cell scanning: implications for cellular activation and immune surveillance. *Proc Natl Acad Sci*
U S A 99:7582-7587.
Travis, A., A. Amsterdam, C. Belanger, and R. Grosschedl. 1991. LEF-1, a gene encoding a lymphoid-
specific protein with an HMG domain, regulates T-cell receptor alpha enhancer function
[corrected]. *Genes Dev* 5:880-894.
Tsuji, G., N. Okiyama, V.A. Villarroel, and S.I. Katz. 2015. Histone deacetylase 6 inhibition impairs
effector CD8 T-cell functions during skin inflammation. *J Allergy Clin Immunol* 135:1228-
1239.
Voisinne, G., A. Gonzalez de Peredo, and R. Roncagalli. 2018. CD5, an Undercover Regulator of TCR
Signaling. *Frontiers in Immunology* 9:
Wang, P., Z. Wang, and J. Liu. 2020. Role of HDACs in normal and malignant hematopoiesis.
Molecular cancer 19:5.
Williams, J.A., K.S. Hathcock, D. Klug, Y. Harada, B. Choudhury, J.P. Allison, R. Abe, and R.J. Hodes.
2005. Regulated costimulation in the thymus is critical for T cell development: dysregulated
CD28 costimulation can bypass the pre-TCR checkpoint. *J Immunol* 175:4199-4207.
Xing, S., F. Li, Z. Zeng, Y. Zhao, S. Yu, Q. Shan, Y. Li, F.C. Phillips, P.K. Maina, H.H. Qi, C. Liu, J. Zhu,
R.M. Pope, C.A. Musselman, C. Zeng, W. Peng, and H.H. Xue. 2016. Tcf1 and Lef1
transcription factors establish CD8 T cell identity through intrinsic HDAC activity. *Nat*
Immunol
Xu, K., W.Y. Yang, G.K. Nanayakkara, Y. Shao, F. Yang, W. Hu, E.T. Choi, H. Wang, and X. Yang. 2018.
GATA3, HDAC6, and BCL6 Regulate FOXP3+ Treg Plasticity and Determine Treg Conversion
into Either Novel Antigen-Presenting Cell-Like Treg or Th1-Treg. *Frontiers in immunology*
9:45.
Xu, M., A. Sharma, D.L. Wiest, and J.M. Sen. 2009. Pre-TCR-induced beta-catenin facilitates traversal
through beta-selection. *J Immunol* 182:751-758.
Yashiro-Ohtani, Y., Y. He, T. Ohtani, M.E. Jones, O. Shestova, L. Xu, T.C. Fang, M.Y. Chiang, A.M.
Intlekofer, S.C. Blacklow, Y. Zhuang, and W.S. Pear. 2009. Pre-TCR signaling inactivates
Notch1 transcription by antagonizing E2A. *Genes Dev* 23:1665-1676.
Yu, S., X. Zhou, F.C. Steinke, C. Liu, S.C. Chen, O. Zagorodna, X. Jing, Y. Yokota, D.K. Meyerholz, C.G.
Mullighan, C.M. Knudson, D.M. Zhao, and H.H. Xue. 2012. The TCF-1 and LEF-1 transcription
factors have cooperative and opposing roles in T cell development and malignancy.
Immunity 37:813-826.
Zhang, M., C. Hu, N. Moses, J. Haakenson, S. Xiang, D. Quan, B. Fang, Z. Yang, W. Bai, G. Bepler, G.M.
Li, and X.M. Zhang. 2019. HDAC6 regulates DNA damage response via deacetylating MLH1. *J*
Biol Chem 294:5813-5826.
Zhang, X.-H., M. Qin, H.-P. Wu, M.Y. Khamis, Y.-H. Li, L.-Y. Ma, and H.-M. Liu. 2021. A Review of
Progress in Histone Deacetylase 6 Inhibitors Research: Structural Specificity and Functional
Diversity. *Journal of medicinal chemistry* 64:1362-1391.
Zhang, Y., S. Kwon, T. Yamaguchi, F. Cubizolles, S. Rousseaux, M. Kneissel, C. Cao, N. Li, H.L. Cheng, K.
Chua, D. Lombard, A. Mizeracki, G. Matthias, F.W. Alt, S. Khochbin, and P. Matthias. 2008.

901   Mice lacking histone deacetylase 6 have hyperacetylated tubulin but are viable and develop
902   normally. *Mol Cell Biol* 28:1688-1701.

903   Zhao, X., Q. Shan, and H.-H. Xue. 2021. TCF1 in T cell immunity: a broadened frontier. *Nature*
904   *Reviews Immunology*

Figure 1.

Chann et al. TCR co-receptors are expressed
sequentially to fine-tune early αβ T cell development

Figure 2.

Chann et al. TCR co-receptors are expressed
sequentially to fine-tune early αβ T cell development

Figure 3.

Figure 4.

Chann et al. TCR co-receptors are expressed
sequentially to fine-tune early αβ T cell development

Figure 5.

Figure 6.

Figure 7.

Figure 8.

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

A

B

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

Chann et al. TCR co-receptors are expressed
sequentially to fine-tune early αβ T cell development

Supplementary Figure 7

A

B

(i)

(ii)

Supplementary Figure 8

Chann et al. TCR co-receptors are expressed
sequentially to fine-tune early αβ T cell development

Supplementary Figure 9

Supplementary Figure 10

Supplementary Figure 11

Supplementary Figure 12

Supplementary Figure 13

Supplementary Figure 14