1 **The cerebellum contributes to tonic-clonic seizures by altering neuronal activity in the**
2 **ventral posteromedial nucleus (VPM) of the thalamus**
3

4 Jaclyn Beckinghausen[1,2,5], Joshua Ortiz-Guzman[3,5], Tao Lin[1,4], Benjamin Bachman[4], Yu Liu[6],
5 Detlef H. Heck[6], Benjamin R. Arenkiel[2,3,4,5] and Roy V. Sillitoe[1,2,3,5*]
6
[1]Department of Pathology and Immunology, Baylor College of Medicine
[2]Department of Neuroscience, Baylor College of Medicine
[3]Program in Developmental Biology, Baylor College of Medicine
[4]Departmen of Molecular and Human Genetics, Baylor College of Medicine
[5]Jan and Dan Duncan Neurological Research Institute of Texas Children's Hospital, 1250
Moursund Street, Suite 1325, Houston, Texas 77030, USA
[6]Department of Anatomy and Neurobiology, University of Tennessee Health Science Center,
855 Monroe Ave, Memphis, Tennessee 38163, USA
15
Address correspondence to Dr. Roy V. Sillitoe

Tel: 832-824-8913
Fax: 832-825-1251
Email: sillitoe@bcm.edu

Manuscript contains:
Number of pages: 55
Number of figures: 9 Figures, 8 Supplementary Figures, 7 Videos, 2 Tables
Number of words: Abstract (138)
24
25 Abbreviated title: Optogenetic modulation of cerebellum-VPM input induces seizures
26
27 Keywords: seizures, ventral posteromedial nucleus, cerebellum, electrophysiology, optogenetics
28
29

Summary

Thalamo-cortical networks are central to seizures, yet it's unclear how these circuits initiate the

seizures. Here, we test the hypothesis that a facial region of the thalamus, the VPM, is a source

of convulsive, tonic-clonic seizures. We devised an *in vivo* optogenetic mouse model to elicit

tonic-clonic seizures by driving convergent input to the VPM. With viral tracing, we show dense

cerebellar and cerebral cortical afferent input to the VPM. Lidocaine microinfusions into the

cerebellar nuclei selectively block seizure initiation. We perform single-unit electrophysiology

recordings during awake, convulsive seizures to define the local activity of thalamic neurons

before, during, and after seizure onset. We find highly dynamic activity with biphasic properties,

raising the possibility that heterogenous activity patterns promote seizures. These data reveal the

VPM as a source of tonic-clonic seizures, with cerebellar input providing the predominant

signals.

Introduction

At their worst, seizures can be fatal, and at best, they significantly decrease a patient's quality of life. Regardless of whether a traumatic brain injury or a specific genetic mutation is responsible for starting the seizure, about a third of patients experience persistent seizures despite having pharmacological, dietary, or surgical interventions[1–9]. In over 60% of cases, the catalyst for initial seizure precipitation is unclear and the causes of such "cryptogenic" seizures remain unknown[10]. As a consequence, it is difficult to optimize therapeutic options for seizure patients[3]. Thus, there is a pressing need to uncover how abnormal seizure activity arises and how it impacts behavior.

Of all the types of seizures, convulsive tonic-clonic seizures (TC) are perhaps the most disruptive and generally feared among patients and caregivers. Unfortunately, these are the most common class of seizures in disorders collectively called epilepsy[11]. Two brain regions that have received the majority of focus in these studies are the cerebral cortex and thalamus, as these reciprocally connected regions form a hypersynchronous, self-perpetuating loop in many forms of epilepsy. Numerous studies have demonstrated that the cortico-thalamic circuit is important in seizures; specifically, such studies point to the cerebral cortex as a main site of initiation[12–14]. However, with a large portion of epilepsy patients being refractory to treatment, stemming from attempts to target these well-studied circuits, it may be critical to narrow in on specific seizure foci and to expand the seizure circuit to determine whether other combinations of neural networks contribute to ictogenesis. We therefore set out to delineate specific focal areas within the cortico-thalamic loop and to identify extra-thalamic inputs that contribute to seizure initiation, and subsequently interrogated the neural activity that may underlie convulsive seizures in awake, seizing animals.

Deeper examination of areas outside the cerebral cortex revealed that seizures can arise from a number of subcortical regions, including the amygdala, hippocampus, and basal ganglia[15–18]. Other regions, such as the cerebellum, are thought to play some role in seizures, but the extent of their contribution is controversial[19]. The majority of these regions project to the thalamus, which is not only a well-established area of information integration, but is also understood to serve as a key gateway for mediating seizures that depend upon these various regions[20–22]. Yet, the precise role of individual thalamic nuclei and their unique combination of inputs on the generation of

abnormal brain activity remains unclear. Pharmacological inactivation studies of thalamic

targets, namely the motor or sensory cortices, demonstrate a clear requirement for somatosensory

areas in both initiation and maintenance of spike-wave discharges (SWD), a hallmark EEG

pattern of absence seizures[12,23–27]. Similarly, the pattern of activation based on human MRI and

analysis of multi-electrode recordings in genetic rat models identify SWDs and activity changes

specifically in the peri-oral region of the somatosensory cortex prior to spread in other regions of

the cortico-thalamic loop[12,23,28]. While such findings have been primarily studied in absence

seizures, the possibility of an epileptic origin in peri-oral circuits is particularly intriguing for TC

seizures, as facial clonus is one of the first behavioral readouts observed in this dynamic

phenotype. We therefore hypothesize that specific cortico-thalamic loops involving facial

regions may be particularly important in seizure generation. Furthermore, in patients with

convulsive seizures, recent human studies reported thalamic changes before cortical involvement

based on electrode recordings in putative deep brain stimulation patients[29]. These findings thus

suggest that the thalamus leads the cortex in generalized seizures, whereas frontal seizures

exhibit activity in the opposite direction[29]. With particular interest in cryptogenic, generalized

seizures, we therefore chose to test circuits in thalamic regions that are known to be tightly

interconnected with the facial somatosensory cortex.

The ventral posteromedial thalamic nucleus (VPM) projects directly to layers 4, 5B, and 6A of

the primary facial somatosensory cortex and receives primarily trigeminal input from the head,

face, and oral regions[30–33]. Known to be intimately involved in the production of spindle waves, a

7-15Hz rhythm occurring during sleep and seizures[34], the VPM has been shown to be an

effective locus in aborting post-stroke epileptic seizures[35]. Due to its reciprocal connections with

the reticular thalamic nucleus as well as direct innervation of both excitatory and inhibitory

neurons in the facial somatosensory cortex, the VPM is a strong candidate for initiating

generalized seizure activity. Using optogenetics to mimic cryptogenic seizures, we test whether

activation of afferent inputs specifically to the VPM, but not to other surrounding thalamic

nuclei, drives severe tonic-clonic seizures. We also used viral tracing to identify the specific

neural projections that may contribute to ictogenesis: specifically, those projections that are

monosynaptically connected to the VPM.

The cerebellum is one such source of direct projections to the VPM. Historically, a cerebellar

role in seizures has been controversial. Earlier observations of seizures, especially hemi-facial

spasms (HFS), in patients with cerebellar gangliomas[36,37] inspired the use of the cerebellar cortex

as a deep brain stimulation (DBS) target for epilepsy patients dating back to the 1970's[38,39].

Despite its success in alleviating seizures in early clinical tests, ethical dilemmas in these studies

and variability in success with subsequent trials due to the previously unknown complexity of

the cerebellar circuitry led to shift in attention away from the cerebellum as a seizure locus. Still,

cerebellar pathology is consistently observed in patients with numerous disorders containing a

common denominator: seizures. For example, in Unverrict Lundborg Disease and familial

cortical myoclonic tremor and epilepsy, both of which exhibit seizures, degeneration in

cerebellar granule cells, Purkinje cells, and overall cerebellar atrophy are observed in human

patients[40,41]. Diffusion tensor imaging has shown abnormal white matter connectivity within the

cerebellum in idiopathic epilepsy[42]. With the growing evidence of cerebellar damage in human

patients with seizures[43–46], plus its involvement in the induction[47,48] and resolving of seizures[43,49–

53] when different stimulation paradigms, this previously disregarded seizure-related brain region

may unlock new therapeutic potential. Although recent studies have demonstrated behavioral and

electrophysiological control of seizures using the cerebellum in genetic mouse models of absence

epilepsy[50,53,54], we have a limited understanding about how to (1) elicit a severe, TC seizure

without using drugs, chemicals, current, or gene-specific mutations, or (2) how cerebellar

circuitry and function fit into a broader seizure brain network. With our optogenetics model, we

test the role of the VPM in seizures and address whether the cerebellum may contribute to the

initiation of TC seizures in this mouse model.

Results

Ntsr1^{Cre} transgenic mice drive reporter expression in brain regions involved in seizures

Ntsr1 encodes for a protein that is expressed primarily in the gastrointestinal tract, but is also widely distributed throughout the brain[55]. A previously generated transgenic mouse model[56,57] demonstrated that the gene's regulatory sequences can be used to drive _Cre_ expression across different brain regions including layers 5 and 6 of the cerebral cortex, the thalamus, and the ventral tegmental area[58,59]. Because a key aspect of our study relies on the identities of _Cre_ expressing neurons that may contribute to ictogenesis, we first set out to both confirm and expand upon previous characterizations of this line. To evaluate the _Ntsr1_ transgene expression across the whole brain, we crossed _Ntsr1^{Cre}_ male mice to two different reporter lines: Sun1 (_Ntsr1^{Cre};ROSA$^{lox-stop-lox-Sun1}$_), a small nuclear envelope protein expressed in the nuclear membrane (Fig 1a), and TdTomato (_Ntsr1^{Cre};ROSA$^{lox-stop-lox-TdTomato}$_), which typically has the advantage of brightly labeling cells and showing their architecture in full due to the intensity of its fluorescent emission (Fig 1f). The concomitant use of these lines allowed us to distinguish between areas containing _Cre_ recombined resident neurons (Sun1 positive) versus areas that receive input from _Ntsr1^{Cre}_-positive neurons and therefore exhibit fluorescence specifically from their axonal processes and terminals (TdTomato positive).

Coronal sections from _Ntsr1^{Cre};ROSA$^{lox-stop-lox-Sun1}$_ mouse brains showed Sun1 nuclear membrane labeling in layer 5/6 of the cerebral cortex (Fig 1b, box; Fig 1c). This particular region with _Ntsr1^{Cre}_-positive expression was studied previously and those neurons were identified as excitatory, corticothalamic projection neurons[60]. Another area that was positive for Sun1 reporter expression was the cerebellar nuclei (Fig 1d,e), which was again previously reported to be a region in which _Ntsr1^{Cre}_ drives expression in excitatory, VGLUT2 (vesicular-glutamate transporter 2)-expressing projection neurons[61]. However, we observed that the expression within the cerebellar nuclei was not evenly distributed: the medial cerebellar nuclei (fastigial nucleus) contained very few _Cre_-positive neurons. In contrast, we found that the reporter signal was heavily localized to projection neurons in the interposed (central) and dentate (lateral) nuclei. This is an important consideration when manipulating cerebellar output, as each nucleus has distinct downstream targets[62–64]. Although the cerebral cortex and cerebellum contained the most prominent distribution of cells with reporter expression, we also observed Sun1 labeling in the

VTA[58], caudoputamen and superior colliculus, as well as sparse labeling in hippocampal and

thalamic neurons, and in a small number of cells that were scattered throughout the pontine

nuclei and various brainstem nuclei (Fig S1ai-av).

By crossing the *Ntsr1*[Cre] mouse to a TdTomato reporter line, we were able to mark and clearly

label somata as well as axonal compartments in each of the regions also marked by Sun1. This

allowed us to distinguish between regions in which *Cre*-expressing neurons originate (that is,

where their somas are located) versus regions that these neurons project to. In addition, labeling

of the neurons in their entirety is important for colocalization analyses. To better characterize the

identity of these neurons, we therefore stained 40um sections from our TdTomato reporter

samples with NeuN, a pan-neuronal marker, NFH, an axonal and synaptic marker of neurons,

and calbindin, a calcium binding protein that labels unique populations of neurons in different

regions of the brain. In the cerebral cortex, calbindin is expressed in non-pyramidal cells[65].

Thalamic neurons with calbindin immunoreactivity identify subpopulations within functionally

distinct nuclei[66], whereas in the cerebellum, calbindin is highly expressed in GABAergic

Purkinje cells that project to the cerebellar nuclei neurons. In the *Ntsr1*[Cre] cross with the

TdTomato line, TdTomato positive cells colocalized with NeuN but not calbindin in the cortex

(Fig S2a-f). These data support previous observations showing that *Ntsr1*[Cre] drives

recombination of a reporter allele in excitatory pyramidal cells[67]. We also co-stained with

tyrosine hydroxylase, a dopaminergic cell marker, to confirm the identity of the VTA-localized,

Ntsr1[Cre] marked neurons as dopaminergic (Fig S2g-i). In the cerebellum, we confirmed previous

reports of *Ntsr1*[Cre]-positive neurons as excitatory and NFH-positive cells that receive input from

calbindin-positive, inhibitory Purkinje cells (Fig S3a-o). We also observed a variable number of

Ntsr1[Cre]-positive Purkinje cells (Fig S3p-r) that were distributed throughout the structure without

an obvious pattern of localization. In any given 40um section through cerebellar cortex, we could

identify anywhere from 0 to 20 labeled Purkinje cells scattered throughout the section.

Importantly, the low number of marked Purkinje cells was unlikely to influence our strategy for

modulating neuronal activity with *Ntsr1*[Cre] (see below). We also found that *Ntsr1*[Cre] marks a

subset of molecular layer interneurons in lobules IX and X (Fig S3s-u).

While the TdTomato cross confirmed the presence of *Ntsr1^{Cre}* recombination in cortico-thalamic

(Fig 1g-h, Fig S1 bi-bvi, Fig S2a-f) and cerebellar nuclei projection neurons (Fig 1i-j, Fig S1c),

the most notable difference between our Sun1 and TdTomato reporter mouse lines was the

pattern of staining in thalamic sub-nuclei: while the thalamus was specifically devoid of

fluorescence in the Sun1 mouse line (Fig 1b, Fig S1ai-aiii), the fluorescence present in thalamic

nuclei of the TdTomato mouse was intense (Fig 1g, Fig S1bi-bvi). These data together suggest

that the resident thalamic neurons themselves do not contain *Ntsr1^{Cre}* expression. In fact, high

power magnification images of the thalamus in the TdTomato mice reveals "pockets" where the

fluorescent signal is missing (Fig 2b). We predicted that these ovoid profiles without staining

were likely the somata of resident neurons. To further investigate these "signal void" pockets, we

examined high-power images of calbindin and NeuN immunoreactivity. As hypothesized, the

absence of the TdTomato signal corresponded with both calbindin and NeuN labeling in the cell

body (Fig S4). Therefore, we conclude that the reporter expression observed in the thalamus of

the *Ntsr1^{Cre};ROSA^{lsl-TdTomato}* mice is the result of marking the incoming axons and fibers of

passage that originate in other areas where *Ntsr1^{Cre}* labels the somata. An example of such fibers

providing the robust thalamic reporter signal can be seen in supplemental Figure 1c, which

shows a dense bundle of cerebellar nuclei axons exiting the cerebellum to innervate the various

thalamic nuclei as well as other brain regions.

To gain optogenetic control over these genetically marked neurons, we next crossed the *Ntsr1^{Cre}*

mice to the *ROSA^{lsl-Channelrhodopsin}* (hereafter referred to simply as ChR2) reporter strain. The

offspring of this cross carried light-sensitive neuronal cell bodies in the aforementioned *Cre*-

expressing brain regions as well as in the axons projecting to, or passing through, the thalamic

nuclei. Based on this expression profile, we were able to deliver 465nm light to the thalamic

nuclei in order to optogenetically modulate thalamic neuron activity indirectly by activating the

axons and terminals of their presynaptic, ChR2-expressing afferent inputs (Fig 2a).

High-frequency light pulses to the VPM elicit tonic-clonic seizures in *Ntsr1^{Cre};ChR2* mice

The thalamus is a critical structure in the seizure network[20–22], although the contribution of its

different nuclei to mediating seizures is not fully understood. Using the *Ntsr1^{Cre};ChR2* mice, we

targeted blue light to manipulate the thalamic nuclei, as each of its major divisions were densely

innervated by a convergence of axons from multiple *Cre*-expressing brain regions. We found that
unilateral delivery of 30Hz pulses of light to the ventral posteromedial nucleus (VPM) elicits a
robust, TC seizure phenotype (n=24 mice, supp video 1,2). Bilateral stimulation or co-
stimulation with other *Cre*-expression regions, such as the cerebellar nuclei, did not enhance or
reduce the severity of this phenotype (n=4 mice, supp video 3). Interestingly, stimulating the
surrounding nuclei with the same parameters, including the posterior thalamic nucleus (PO, n=3
mice), the ventroposterior lateral nucleus (VPL, n=3 mice), the reticular thalamic nucleus (RTN,
n=3 mice), and the lateral dorsal nucleus (LD, n=2 mice) did not elicit overt behavioral changes
(supp video 4). We quantified the relative fluorescence in each of these individual thalamic
nuclei (n=4 mice, 5-10 slices per mouse) to determine whether the non-responsive nuclei
received less ChR2-expressing input. While some nuclei, such as the RTN, PO, and CL, had
significantly less fluorescence (Dunnett's multiple comparisons test, Table 1), which may
account for the absence of behavior upon their stimulation, other nuclei like the VPL contained
comparable levels of ChR2 input as the VPM (Fig 3a,b, Dunnett's multiple comparisons test,
Table 1). These data suggest that the initiation of seizures in *Ntsr1^{Cre};ChR2* mice requires the
modulation of unique sets of inputs that converge within the VPM and argues against a global
activation of general input pathways.

Post-hoc anatomical examination of fiber optic targeting further identified a specific locus within
the VPM bordering the PO that reliably elicited seizures (Fig 3c,d, n=7 mice). Targeting of fibers
to other regions within the VPM did not elicit a behavioral change. These data indicate that a
discrete set of VPM inputs are required for seizure initiation. With repeated stimulation, we
determined that all *Ntsr1^{Cre};ChR2* mice with accurate VPM fiber targeting eventually exhibited
stage 6 seizures according to a modified Racine Scale[68] (Table 2), a widely used template for
categorization and ranking of rodent seizure severity. To achieve equally severe seizures, our
paradigm involved epochs of 30-s stimulation that were presented for 1 hour and repeated daily.

Following seizure induction, humans and animal models exhibit a post-ictal state characterized
by lethargy. The mechanisms contributing to the transition from ictal to postictal state remain
elusive: however, candidate neural mechanisms include increased GABA-ergic inhibition,
synaptic effects of neuromodulators such as endocannabinoids or adenosine, neuronal

exhaustion, neurotransmitter depletion, increased extracellular potassium, and decreased extracellular potassium[69,70]. In our model, we could not induce a sequential seizure during this state, which typically lasts about 3-6 minutes from the end of a previously elicited seizure (Fig 3e). These data indicate the possibility of a potential mechanism involving resource depletion as opposed to one that might involve a change in cell inhibition. The support for this distinction is that subsequent light stimulation appears to have no additional effects on seizure-related behavior during this refractory period.

We next compared the phenotype in our optogenetics model with the well-established, chemo-convulsant kainic acid (KA) model[71] (n=4 mice). KA is nondegradable glutamate analog that, when injected intracranially or systemically *in vivo*, induces behavioral seizures and neuronal excitotoxicity[72,73]. A comparison between optogenetic VPM activation and KA injection revealed a number of common behaviors during severe seizure episodes (supp video 5, Fig 4a). Specifically, both models exhibited tensing of the body (stage 1), hunching posture, erect straub tail, facial and forelimb clonus (stages 2-4), and eventual loss of upright postures (stage 5) with convulsions leading to erratic jumping (stage 6, Table 2). The similar seizure-related behaviors observed in both our *Ntsr1^{Cre};ChR2* model and the KA model argues that a core set of thalamic circuits may be central to seizure initiation. The question we next address is how these behaviors relate to activity.

Electrocorticography (EcoG/EEG) activity reflects a widespread presence of seizure activity

Electroencephalography (EEG) is an electrophysiological measure used for diagnosing seizure conditions in humans. It is also an important tool for verifying animal models that mimic these conditions. EEG measures population-wide neuronal activity using a series of electrodes typically placed on the surface of the scalp[74]. In mice, increased signal quality is achieved by placing the electrodes directly above the brain. Due to this difference in electrode position, this technique is referred to as electrocorticography (EcoG); for simplicity, we will hereafter refer to both as EEG.

The presence of seizures as a behavioral phenotype in rodents can have components that overlap
with other disease-like behaviors. For example, paroxysmal dystonia, a condition characterized
by sudden attacks of antagonistic muscle co-contractions, can appear phenotypically similar to
rhythmic tonic-clonic seizures[75,76]. As such, dystonia-like movements and seizure behaviors can
be difficult to distinguish in rodents[77]. To confirm the presence of seizures in the *Ntsr1^{Cre};ChR2*
mice, we recorded EEG signals from the ipsilateral somatosensory cortex, bilateral motor cortex,
and the cerebellum (Fig 4b). We tested whether VPM activation in the *Ntsr1^{Cre};ChR2* mice
results in neuronal hypersynchronization. In accordance with the presentation of seizure
behaviors, EEG activity in *Ntsr1^{Cre};ChR2* mice showed high amplitude, hypersynchronized
activity across the cerebral cortices and cerebellum (Fig 4c, 5a, n=11 mice). We observed similar
EEG responses in the KA model (Fig 4c, n=4 mice). This hallmark seizure activity was
consistent across all optogenetically-induced mice that exhibited behavioral seizures (supp Fig 5,
video 6). Figure 5b shows an example of EEG traces from video-verified seizures in a single
mouse during superimposed periods of stage 2 (head nodding and facial clonus) and stage 6
(severe TC seizure with loss of posture and erratic jumping) seizures. In each scenario, the EEG
signal reflects the seizure severity at that point in time: during stage 2 clonus, the spike
amplitude rose above baseline but was weaker than that during stage 6 convulsions. The
presence of high amplitude and hypersynchronized EEG activity and the resemblance of the
induced behaviors to an established model provided compelling evidence that our optogenetic
paradigm indeed induces TC seizures.
300

Optogenetic activation of the VPM leads to the coherence of cerebral cortical activity
The synchrony exhibited during seizure-derived EEG occurs as a result of summed dipole fields
of contributing neurons in the local electrode region[78]. While it has been argued that this
synchrony is driven by only a sub-population of synchronously co-active neurons,[79] it is
nonetheless a consistent measure of seizure activity. Seizure activity is typically highly rhythmic
covering a broad spectrum of frequencies[80,81]. To evaluate the synchronization of rhythmic
seizure activity between regions in our model, we examined the coherence of EEG oscillations
between the motor cortices, ipsilateral somatosensory cortex, and the cerebellum (Fig 5c, n=4
mice). The left motor cortex and left somatosensory cortices, both having direct neuronal
connectivity with, and physically closest to, the stimulation site (left VPM), were most reflective

of the intense activity at the seizure origin. Interestingly, these regions exhibited the highest

coherence during seizures, particularly in frequencies up to 50Hz (Fig 5c, supp fig 6a).

Conversely, the right motor cortex showed a more modest increase in coherence with the left

hemisphere, likely due to its position contralateral to the seizure origin. Still, even a lower degree

of synchronization supports the generalized spread of the seizure away from its origin, which

ultimately incorporates contralateral brain regions into the pathophysiology that eventually

drives the behavior.

The cerebellum and VPM express highly coherent EEG activity during seizures

The cerebellum is typically used as a location of reference for EEG signal collection, an idea

based on data showing that the cerebellum exhibits minimal EEG changes during seizure

episodes[82–85]. In contrast to these data, we found elevated cerebellar EEG activity[86–89] that was

tightly coupled to the left motor cortex, specifically during seizures occurring between 13 and

50hz (Fig 5c, n=4 mice). The increase in interregional coherence involving the cerebellum was in

accordance with our finding that the cerebellum expresses both high amplitude and

hypersynchronous activity during optogenetic-induced seizures that were triggered by VPM

activation (Fig 4c, 5a, 5c, supp Fig 6c).

A patterned shift in brain activity emerges as the seizures generalize across regions

EEG can be used to understand, predict, and treat seizures by performing spectral analyses of the

signal[90] and testing for an increase in the power of frequencies between 0.5 and 30Hz. Here, we

test whether optogenetic-induced seizures produce a shift in activity by performing wavelet-

based time-frequency analyses of the EEG activity. To do so, we examined changes at the

following frequency bands: delta (0-4Hz), theta (4-8Hz), alpha (8-13Hz), beta (13-30), and

gamma (30-60Hz)[91] (supp Fig 7, n=7 mice). With a fixed power scale across brain regions, we

found that the activation pattern across the brain was inconsistent between mice. We therefore

reran the analysis using an optimized scaling for each individual in which a specific brain

region's own maximal power frequencies were graphed. Scale optimization revealed a patterned

shift in the dominant activity across all sub-domains from the left hemisphere (left

somatosensory and motor cortices, and the central cerebellum) to the right hemisphere (right

motor cortex) (Fig 5d, supp Fig 7).

In human epilepsy, high frequency EEG signals that are above 60Hz are thought to include noise

due to interference from the skin, skull, and muscle[92]. Bypassing the skin and skull with our

electrodes combined with a high sampling rate allowed us to examine these higher frequencies in

mice. We were interested in activity in these domains as high frequency oscillations (HFO's)

recorded during epilepsy surgeries have been suggested to have biological relevance in localizing

the seizure onset zone[93–95]. Thus, we examined whether changes in activity occurred between 60

and 250Hz in our model. Frequency plots with data averaged over multiple mice revealed a

patterned shift in high frequency (HF) power reminiscent of that in the low frequency domains:

from the site of origin (left VPM) to the contralateral side of the brain (Supp fig 7, Fig 5d, n=7

mice).

We next performed power analyses between 0 and 250Hz to gain a better appreciation for

whether neural activity changes on a broader scale (Fig 5e,f, supp Fig 8). Power plot analysis

demonstrated that there is a significant bias towards an increase in power in the lower frequency

band between 0 and 50Hz during the stimulation period (paired t-test of area under the curve,

n=4 mice, average of 4 brain regions each, p<0.0001). In fact, the changes that are represented in

the heat plots at higher frequencies are masked in the power plots due to the dramatic differences

that occur between 0-50Hz. These data suggest that although HFO's may help to localize the

seizure onset, the largest changes in neuronal oscillation power during seizures indeed occur in

the lower frequency ranges.

Optogenetic stimulation of afferent input to the VPM generates a dynamic, heterogeneous

firing pattern in the connected thalamic neurons

We next sought to determine the electrophysiological properties that drive optogenetic-induced

seizures in $Ntsr1^{Cre}$*;ChR2* mice. Specifically, we asked how light stimulation of fibers projecting

to the VPM affect resident thalamic neuronal activity to initiate seizure behavior. To address this

question, we used pulled glass electrodes to record single unit activity in the thalamus of awake,

behaving mice implanted with fibers targeted to the ipsilateral VPM. This technique allowed us

to examine changes in neuronal activity while simultaneously delivering subthreshold levels of

light that are capable of initiating the onset of behavioral seizures but without developing

convulsions. To achieve the necessary stability, mice were secured over a freely rotating wheel

with a headplate and provided with periods of habituation (Fig 6a). With this approach, we

successfully recorded the activity of more than 70 thalamic neurons using an *in vivo* extracellular

technique (n=5 mice).

With the delivery of subthreshold, 2 second light pulses to the ipsilateral VPM, thalamic neurons

expectedly demonstrated a repeatable excitatory response (Fig 6b). However, we found that a 30

second seizure-inducing light simulation did not produce a predictable, single response in

thalamic neurons. Instead, in addition to excitation (Fig 6b, 6f), they exhibited various biphasic

response profiles (Fig 6c-e) or were unaffected by light stimulation (Fig 6g). Most resident

neurons exhibited some degree of firing response – only 17% were unaffected by light

stimulation (Fig 6h). We postulate that these nonresponsive neurons represent a class of

interneurons that do not engage in hypersynchrony[96] or are not innervated by *Cre*-expressing

axons. Interestingly, only 16% of the recorded neurons exhibited purely an increase in frequency.

Most cells engaged in a biphasic response, alternating between periods of rapid firing and silence

(Fig 6c-e). Due to a high concentration of T-type calcium currents, neurons in the VPM are

known to rebound with bursts of excitation following release from hyperpolarization[97–99].

Therefore, during a seizure, thalamic neurons may transition through discrete properties between

states of inhibition and excitation.

Acute optogenetic-induced seizures do not result in cell death, but they do increase dFosB

expression on the side of stimulation

Whether acute or chronic seizures cause neuronal death is still largely debated. In several animal

models, including electroconvulsive and chemically-induced seizure models (KA, pilocarpine),

postmortem analysis of brain tissue provides evidence of programmed cell death[100–105]. However,

kindling models have reported tissue expansion following brief seizure induction rather than

direct cell loss[106–110]. To examine whether the *Ntsr1[Cre];ChR2* model involves cell death

following seizures, we compared the cell density of H&E stained cells in control mice versus

experimental mice that experienced at least one stage-6 seizure. We did not find a significant

difference in the number of H&E stained cells in the control animal thalamus versus the seizure

animals, indicating that with acute stimulation, optogenetic-induced seizures do not lead to a

significant amount of neuronal loss (Fig 7a, unpaired t-test, p=0.4627 n=5 mice per condition, 2-5 tissue sections per animal).

In contrast to cell death, it is well-established that epileptic activity affects long-term neuronal plasticity, which may eventually affect learning and memory[111–115]. Changes in activity are effectively measured by examining the levels of dFosB, a stable immediate-early gene with a long half-life that allows for easy detection[116]. Previous work demonstrates that dFosB is upregulated following seizures, although most studies focus on changes in the hippocampus and cerebral cortex due to their involvement in kindling and temporal lobe epilepsy[117–121]. We therefore tested whether this change in activity is also present in the thalamus, the suspected source of the seizures in our model. We found a significant increase of dFosB in the thalamus ipsilateral to the site of stimulation (Fig 7b-f, Tukey's multiple comparisons test, p<0.0001, n=4 mice, 3-6 tissue sections per animal). The contralateral thalamus exhibited significantly less dFosB expression than that of the ipsilateral thalamus (p<0.0001), although the level was still higher than in stimulated thalamic nuclei of *Cre* negative littermate controls, (p=0.002). These data indicate that optogenetically induced seizures are accompanied by an upregulation of dFosB in the thalamus, but without significant cell loss.

Injection of a *Cre*-dependent virus into the VPM delineates source areas of activation

Based on the hypothesis that the VPM represents a strong seizure locus, we next asked which of the *Cre*-expressing brain regions that project to the VPM are capable of eliciting this robust phenotype, or alternatively whether all regions identified above participate in VPM-mediated seizures. We therefore injected retro-rAAV-Ef1a-mCherry-WPRE into the VPM of *Ntsr1^{Cre}* mice (Fig 8a, n=5). In this experiment, only *Cre*-positive neurons projecting directly to VPM recombine to express mCherry. After allowing up to 5 weeks for viral tracing and recombination, mice were perfused and prepared for immunohistochemistry. Despite the presence of multiple *Cre*-expressing regions observed in the *TdTomato* cross (supp Fig 1), only the cerebral cortex and cerebellar nuclei demonstrated direct projections to the VPM after virus labeling. The most robust axonal bundles filled by the virus injection originated from the somatosensory areas and cerebellar nuclei (Fig 8b-e). We did not find labeling from other areas that are known to project

434 to the VPM, including the internal segment of the globus pallidus, hippocampus, zona incerta,
435 hypothalamus, and amygdala.
436

Drug-induced inhibition of the cerebellum prevents the induction of seizures from the VPM
Based on the virus labeling of the cerebral cortex and cerebellum and that these regions mediate
a strong influence on VPM-induced seizure activity, we asked whether both the cerebral cortex
and cerebellar nuclei are necessary for initiating the phenotype. To parse apart the relative
contribution of each region, we used lidocaine to reversibly block each brain region. Due to the
rapid kinetics of lidocaine and the reversibility of its effects, we performed within-trial
comparisons of seizure induction before administration, immediately after lidocaine delivery,
and after lidocaine washout. We delivered 4% lidocaine through implanted cannulas that
bilaterally targeted the cerebellar nuclei (Fig 9a), somatosensory cortex, and motor cortices (Fig
9b). Based on previous studies, we calculated that the neural inactivation window would extend
approximately 5-30 minutes after initial delivery[122–125]. Furthermore, previous work from our lab
used fluorescent conjugated antibodies to detect the spread of lidocaine and determined that the
drug can travel approximately 1mm from the injection site[126]. By delivering lidocaine to each of
these regions following seizure induction, we evaluated whether ablation of functional
connectivity interrupted the ability to elicit subsequent seizures. We found that pharmacological
inhibition of the cerebellar nuclei (Dunnett's multiple comparisons test, $p<0.0001$), but not
somatosensory ($p=0.9562$) or motor cortices ($p=0.8835$), eliminates seizure induction (Fig 9,
supp video 5, n=4 cerebellar ablation, n=2 per cortex ablation). Following lidocaine washout, the
ability to induce a seizure was rescued. Together, these data support the hypothesis that
cerebellar connectivity to the VPM is essential for seizure initiation in this optogenetic model,
whereas specific regions of the cerebral cortex may be more heavily involved in generalizing and
sustaining the seizure only after it has been initiated elsewhere in the brain.

Discussion

Optogenetics provides temporal and cell-type specific regulation to model seizures

There are now many animal models of seizure including mouse, rat, guinea pig, and non-human primate[127–130]. The broad range of models has been advantageous given the heterogeneity of human seizure conditions[131]. However, although each model has its own limitations, their use is nonetheless vital to dissecting circuit mechanisms and other factors that may contribute to the emergence or worsening of these conditions. Currently, there are at least three classes of seizure research models: genetic, chemoconvulsant, and electroconvulsant. While valuable for heredity conditions, those relying upon genetic mutations often do not reflect the cryptogenic forms of epilepsy, which account for over 60% of human cases[132]. Electro- and chemo-convulsant models begin with a healthy brain at baseline, but can introduce secondary effects that may obstruct the analysis of seizure pathophysiology[133–136]. Furthermore, in all three classes of models, the control of seizure initiation can lack temporal precision. We therefore devised an optogenetics approach to overcome these hurdles. We used cell-type specificity to modulate a genetically defined pool of neurons that evoke seizures in otherwise normal mice and then we tracked the spread of abnormal activity. Using an *Ntsr1*^{Cre} transgene to drive the expression of a light-sensitive channel in major seizure-engaged regions, we were able to test the anatomical, neuronal, and circuit properties of seizure induction. This model provides control over neuronal activity, an ideal feature for disentangling seizure dynamics. In addition, because optogenetic-induced seizures are under precise temporal control, the stimulation paradigm reliably initiates the seizures within a predictable time window.

The use of optogenetics is not new in the seizure field: a number of studies have benefitted from its advantages to abort seizures in established models of genetic or acquired epilepsy[51,137,146–151,138–145], and a few have demonstrated successful induction[138,152–155]. Current models of optogenetic-driven seizures, some of which exhibit kindling[156], provide an important and novel platform for studying the specific role of defined neuronal populations in epileptogenesis without cell loss or glial activation. However, our model differs from previous optogenetic models in a few important ways. First, our mouse model does not include viral injections of channelrhodopsin. Although use of viruses can indeed include specificity of cell type targeted via genetic promotors within the viral construct, this extra surgical step can lead to differential

targeting of neuronal subsets that lacks consistency from mouse-to-mouse. Second, with accurate
fiber optic targeting, our line exhibits severe tonic-clonic seizures upon the first train of light
delivery. Current optogenetic models require delivery of light stimulus over a period of days to
weeks, drastically increasing experimental time[156]. Third, our model does not include or require
the use of seizure-sensitive backgrounds, whether genetic or primed with seizure-inducing drugs.
Some optogenetic stimulation paradigms induce convincing TC-seizures within the first few
stimulation sessions, but require the use of acute seizure-inducing drugs such as 4-AP with or
without the combination of Mg2+ or NMDA[138,153,154,157]. Thus, we greatly simplify and
streamline the procedure for optogenetic induced seizure induction using a genetically tagged
channelrhodopsin mouse line.

Ntsr1^Cre^-ChR2 **mice can be used to mark and manipulate ictogenic brain regions**

Classical brain regions known to play a major role in seizure pathology are the cerebral cortex
and the thalamus. Their roles in seizures are underscored by their reciprocal connectivity with
one another and the emergence of neuronal hypersynchrony that can be observed during diverse
types of seizures. However, there is now compelling evidence that extra-corticothalamic regions,
including the hypothalamus[158,159], hippocampus[152,160,161], olfactory cortex[162–165], amygdala[107,166–
168], striatum[169–171], and the cerebellum[54,172–177] also play critical roles in seizures. In this study,
we used an *Ntsr1^Cre^;ChR2* mouse to test the contributions of the cerebellum and its connections
in initiating seizure activity.

The *Ntsr1^Cre^* mouse line has been previously used to demonstrate cerebral cortical involvement
in absence epilepsy. Specifically, deletion of *Cacna1a*, which encodes for a P/Q-type voltage
gated calcium channel alpha-subunit, in layer VI cortical neurons of *Ntsr1^Cre^* mice generated
spontaneous absence epilepsy[178]. Other studies of absence seizures also attribute layer 5/6
cortical neurons in the facial somatosensory cortex as players in epileptic discharge in a rat
model of epilepsy[23]. These findings are intriguing as a defining feature of rodent TC seizures,
including those elicited in our model, is facial clonus. Our data therefore suggests that absence
and TC seizures may have overlapping networks in their pathology, particularly regions such as
the somatosensory cortices.

It is interesting to note that the optogenetic-induced seizures that we initiated outlast light
delivery, becoming independent of the original stimulation (supp video 4). Combined with the
gradual worsening of the overall seizure phenotype with repeated stimulations, this raised the
possibility that our model was somewhat equivalent to a kindling paradigm[156,179–181]; however,
we never observed spontaneous seizures in any of our experimental mice (n=24*)*. Nonetheless,
we cannot rule out that more subtle seizure-like activity occurred during periods in which the
mice were not monitored. We are further convinced that these mice do not experience kindling
because given an extended period of multiple days without stimulation, the mice return to
baseline seizure severity upon the resumption of stimulation sessions. It is possible, therefore,
that while repeated stimulations over the course of hours or days potentiates seizure-involved
circuitry, presumably due to potential recruitment of nearby seizure loci, a sufficient length of
time between stimulations could allow for recovery of those circuits. This is interesting
considering that a single animal can toggle between periods of seizure and behavioral normalcy.
We validated our seizure phenotype with EEG. This was important as we did not observe
significant changes in the 5-7Hz range (corresponds to 3Hz in humans), a feature used to
diagnose absence epilepsy[182]. We conclude that absence seizures are also an unlikely component
in our model. Rather, the generalized TC seizure activity exhibited behaviorally can be attributed
to a broad increase in low- and high-frequency EEG activity.
539

Co-activation of a discrete set of inputs to the VPM accounts for seizure initiation
Neurons within the thalamic nuclei have distinct genetic profiles, morphologies, firing patterns,
and synaptic properties [183–185]. Despite findings that argue for homogeneous activity during
seizures, calcium imaging and multiarray studies indicate the presence of heterogenous activity
among neuronal populations both at seizure onset as well as throughout the seizure period[96,186–
191]. Furthermore, numerous studies in animal models have established that a proportion of cells
within the epileptic brain remain largely normal during seizure episodes. For example, in a cat
model of focal epilepsy, only 30% of recorded neurons demonstrated an epileptic response[96]. In
humans and monkeys, as many as 51% of neurons recorded by multielectrode arrays were
nonresponsive during seizures[187–191]. Here, we demonstrate that perhaps a higher proportion – as
many as 80% - of VPM neurons may serve an active role. Whether this is true for other regions
has yet to be resolved.

Consistent with the presence of heterogeneous activity, there is now evidence that instead of excitatory neurons being exclusively responsible for initiating seizures, an increase in interneuron activity may first cause local populations of neurons to become quiescent, only to subsequently rebound with sudden bursts of sustained activity[138,192–194]. It has been suggested that parvalbumin-expressing interneurons may be particularly important for this form of ictogenesis[138,192–194]. The VPM is known to target both excitatory cortical pyramidal cells as well as GABAergic parvalbumin interneurons[195–197]. In fact, in an investigative long-range tracing study, the authors determined that the VPM accounts for 72% of subcortical input to the PV neurons[198]. In addition, PV-neurons within the reticular thalamic nucleus were reciprocally connect to the VPM[197,199]. It is possible, therefore, that excitatory-inhibitory neuron interconnectivity between the VPM and surrounding regions partially account for the specific seizure activity observed in our optogenetic induction model.

Optogenetic-induced seizures induce dFosB expression mainly on the side of stimulation

The immediate-early gene *fosB,* a marker of activity and subsequent plasticity, is altered following seizures[200,201]. One study found that *fosB*-null mice exhibit spontaneous seizures that resemble the first stage of forelimb clonus seen in our optogenetic model[202]. dFosB, a splice variant of *fosB,* is used to measure changes in activity due to its long half-life. Again, this variant is specifically upregulated in models of epilepsy[117–121]. In addition, dFosB activity has been recently implicated as a component of the learning and memory deficits that occur in Alzheimer's Disease, and these changes have been attributed to seizure activity in these animal models and human patients[203].

In our model of VPM-induced seizures, we demonstrate that dFosB expression is increased in the thalamus on the side of stimulation (Fig. 7). Upregulation of dFosB also occurred in the contralateral thalamus, though to a lesser degree. Our interpretation is that the largest change in activity occurs at the seizure epicenter and then spreads to the contralateral hemisphere. Overall, this marker of activity is not affected by light delivery: rather, dFosB expression only increases in the thalamus when a seizure-evoking optogenetic train of light successfully induces seizure activity, as demonstrated by ChR2[-/-] control animals that received the same light stimulation

without experiencing seizures. Previous studies of corneal stimulation to induce TC seizures in
rodents observed similar dFosB increases, although in the amygdala and hippocampus[204]. We did
not see significant changes in CA1, CA3, or dentate gyrus, nor in amygdala, perhaps due to the
different method of seizure induction. Lastly, while previous studies have reported differences in
cerebellar activity through decreases in dFosB expression in the cerebellar nuclei[172], we did not
observe such changes. This is likely a result of our experimental paradigm, which involves only
one week of seizure inductions, with likely correspondence between the models if a chronic
paradigm was used.

Cerebellar connectivity is required for tonic-clonic seizure initiation at the VPM

Cerebellar contribution to seizures is one with a long-standing history: as the first deep brain
stimulation target for treatment of epilepsy in the 1970's, the cerebellum was initially a major
site of focus in seizure studies. Evidence for a clinical role of cerebellum in epilepsy is abundant:
patients with tubercular abscesses in cerebellum have been reported to experience seizures.
Likewise, those with gangliomas in the cerebellum noted elimination of seizures upon their
resection[36,205]. In patients with histories of repeated seizure, a correlation between the number of
seizure episodes and cerebellar atrophy through Purkinje cell death is noted through MRI
imaging studies[206,207]. Further evidence of cerebellar involvement lies in epilepsy comorbidities:
in many cases, patients with seizure disorders present with known cerebellar-derived
comorbidities including ataxia, dystonia, and tremor[208,209]. Recently, studies have shown reliable
eradication of the physical symptoms and abnormal EEG with treatment targeted to the
cerebellar nuclei[49,51]. EEG and single unit electrophysiology studies corroborate these data by
demonstrating robust modulation of neuronal activity in the cerebellum during seizure episodes
in various mouse models[210–212].

Our data demonstrates that the cerebellum may in fact also engage in chronic network
hyperactivity during seizures, which supports a growing literature supporting a dynamic role of
the cerebellum in epilepsy[19,43,211,213,214,44,47,50–53,173,205]. Based on our time-frequency analyses, we
tracked the temporal spread of seizure activity from the left motor cortex to the left
somatosensory, cerebellar, and eventually right cortical areas. According to these data, the
cerebellum is activated prior to seizure spread and the greatest peak power across frequency

bands was in the cerebellum (supp Fig 8). It is interesting to note that cFos expression is abundantly activated in granule cells in epilepsy[173]. In cases of cerebellar gangliomas where seizures consequently manifest, resection of the tumor eliminates the seizures[36]. Similarly, clinical studies note the presence of cerebellar gangliomas in cases where the sole behavioral symptom is severe seizures[215]. Ming, et al. (2021) hypothesize that cerebellar disinhibition leads to cortical activation, thereby exacerbating or initiating seizures[216]. Our data support the hypothesis that cerebellar activation can drive seizures.

However, there is also evidence proposing the opposite: that decreasing cerebellar activity worsens seizures. For example, clinical observations since the early 1800's have noted cerebellar atrophy in patients with prolonged seizures; immunohistochemistry experiments report lower dFosB expression in cerebellar nuclei following seizures, indicating a decrease in neuronal activity[172,217]; pharmacological inhibition of the cerebellar nuclei increased abnormal EEG activity in absence mouse models, whereas excitation blocked their occurance[50]. Despite these seemingly contradictory results, both scenarios could co-exist. Depending on the pattern of cerebellar activity, it could either synchronize or disrupt downstream networks to modulate seizures. In genetic models of epilepsy, cerebellar firing is abnormal[172,213,218,219]. During periods of absence seizure episodes, the cerebellum switches from rhythmic to asynchronous firing.[50,219] This rhythm of cerebellar output may force synchrony between regions like the thalamus and cerebral cortex. In these disease models, where baseline cerebellar activity is already disrupted, delivery of stimulation through DBS or optogenetic pulses of light may disrupt synchronized networks and terminate seizures. However, in models with normal baseline cerebellar activity, delivery of a patterned train of light to cerebellar nuclei instead forces the same bursty pattern that is inherent in epilepsy models, thereby initiating synchrony in upstream regions to cause a seizure. In patients with genetic forms of epilepsy, the cerebellum may therefore be an important target for treatment to correct abnormal output. In idiopathic or cryptogenic epilepsies, which make up an overwhelming 65% of cases, abnormalities in the cerebellum may be at fault for the spontaneous emergence of seizures. In both cases, the cerebellum should be considered as a key player in epilepsy pathophysiology.

Conclusion

In these studies, we devised an optogenetics mouse model to demonstrate that 30Hz coactivation
of cerebral cortical and cerebellar input to the VPM induces severe TC seizures. With viral
tracing and reversible inactivation, we show that of these circuit inputs, the cerebellum has a
predominant role in seizure initiation, supporting recent work that links cerebellar connectivity to
epilepsy. Furthermore, using single-unit *in vivo* electrophysiology recordings performed in
seizing mice, we uncover heterogeneous, biphasic activity of thalamic neurons during seizures.
Together, these data suggest that cerebellar input to the VPM is a critical source of abnormal
signals at seizure onset.

Acknowledgements

This work was supported by funds from Baylor College of Medicine (BCM) and Texas
Children's Hospital. RVS received support from The Hamill Foundation and the National
Institutes of Neurological Disorders and Stroke (NINDS) R01NS089664, R01NS100874, and
R01NS119301. RVS is also supported by a grant from the Dystonia Medical Research
Foundation (DMRF). RVS, DHH and YL are supported by National Institute of Mental Health
(NIMH) R01MH112143. Research reported in this publication was supported by the Eunice
Kennedy Shriver National Institute of Child Health & Human Development of the National
Institutes of Health under Award Number P50HD103555 for use of the Cell & Tissue
Pathogenesis Core and the Neuroconnectivity Core. The content is solely the responsibility of the
authors and does not necessarily represent the official views of the National Institutes of Health.
JB received support from F31NS115432.

Contributions

JB, JO-G and RVS designed the experiments. JB, JO-G, TL, and RVS performed the
experiments. JB, BB, YL, DHH, and RVS designed code for electrophysiology analysis. JB, JO-
G, BB, YL, DHH, BRA and RVS analyzed the data. JB and RVS wrote and edited the paper.

670
671 **Competing Interests**

672 All authors declare that no competing interests exist.

673
674

675 **Ethics**

676 All mice were housed in an AALAS-accredited facility on a 14hr light cycle. Husbandry,
677 housing, euthanasia, and experimental guidelines were reviewed and approved by the
678 Institutional Animal Care and Use Committee (IACUC) of Baylor College of Medicine (protocol
679 number: AN-5996).
680

Methods

Animals. To develop transgenic mice with restricted TdTomato or Sun1 expression, we crossed heterozygous *Ntsr1^{Cre}* males (*B6.FVB(Cg)-Tg(Ntsr1-cre)GN220Gsat/Mmucd,* University of California, Davis MMRRC, Mouse Biology Program) to het- or homozygous *ROSA$^{lox-stop-lox-TdTomato}$* or *ROSA$^{lox-stop-lox-Sun1}$* lines (*B6;129-Gt(ROSA)26Sor$^{tm5(CAG-Sun1/sfGFP)Nat}$/J,* Jackson Laboratory, Bar Harbor, Maine). These offspring, whose genotype was confirmed through tail snips or ear punches, were used for immunohistochemistry. To gain light-sensitive control over these neurons, we crossed *Ntsr1^{Cre}* males to *ROSA$^{lox-stop-lox-ChR2-YFP}$* females. Again, their genotypes were confirmed through tail and ear clippings, but also through immunohistochemistry after experimentation. We bred the mice using timed pregnancies, and designated noon on the day a vaginal plug was detected as embryonic day (E) 0.5 and the day of birth as P0. For experimental studies, mice from P60 to P180 were used. Mice of both sexes were studied, and opsin or *Cre*-negative littermates were used as non-expressing seizure controls. The mice were housed on a 14h/10h light/dark cycle. All animal studies were carried out under an approved institutional animal care and use committee animal protocol according to the institutional guidelines at the Baylor College of Medicine.

Surgery. Surgery for awake optogenetics was performed as detailed in Ung and Arenkiel, 2012[8]. Briefly, mice were anesthetized with 4% isoflurane in an induction chamber until unresponsive to the toe-pinch reflex. After transferring to a stereotaxic apparatus, isoflurane was lowered and maintained at 2% for the duration of surgery. Using aseptic technique and a sterile work field, the mouse was cleaned and prepped, the skin above the skull opened, and a small craniotomy of approximately 1mm in diameter made above the VPM (AP -1.6mm, ML +/-1.55mm). A polished optic fiber previously glued into a ceramic ferrule (Thorlabs, Newton, NJ, USA; #CFLC230-10) was then gently lowered to the approximate depth (2.8mm) and secured with Metabond Adhesive Luting Cement. For most surgeries in which EEG was also incorporated, 6 additional craniotomies were made above the super colliculi (reference and ground electrodes, AP -5.21mm, ML +/-1mm), primary sensory cortex (AP +1mm, ML +3.5mm), primary motor cortices (AP -1mm, ML +/-1mm), and cerebellum (AP -6mm, midline). An EEG head mount compatible with a detachable preamplifier (Pinnacle Technology, Inc, Lawrence, KS, USA; #8406) was equipped with six silver ball-tipped wires (A-M Systems, Sequim, WA, USA;

#785500) soldered to individual pins. This head mount was glued to the most rostral portion of

the skull. The wire free ends were inserted into the craniotomies between the skull and dura. All

wires were secured with UV epoxy followed by Metabond Adhesive Luting Cement. Next, the

fiber optic implant as well as an EEG head mount were wrapped in dental cement. After the

surgery, the mouse was placed in a warming box (V500, Peco Services Ltd., Cumbria, UK) to

prevent hypothermia while the anesthesia wears off. Once the mouse was awake and mobile, it

was returned to the home cage. The mouse was allowed to recover for 2-3 days before

optogenetic stimulation and recordings were initiated.

Optogenetics and EEG collection. To induce seizures in this mouse model post-surgery, a fiber

optic cable powered by a 465nm light box (ALA Scientific Instruments Inc, Farmingdale, NY,

USA) was carefully inserted into the unilateral fiber implant surgically targeted to thalamus

(VPM). When EEG was included, a preamplifier (Pinnacle Technology, Inc, Lawrence, KS,

USA; #8406) was connected to the implanted EEG head mount. All EEG and optogenetic

stimulation parameters with subsequent data was recorded through Spike2 software and

delivered using a CED Power1401 data acquisition interface (CED, Cambridge, UK).

Recordings were paired with live video acquisition and were monitored for 5 minutes before

stimulation, the duration of stimulation, and 5 minutes after stimulation. To entrain the circuits to

seizures, 1 hour of 30Hz trains of optogenetic stimulation, 2 minutes light on and two minutes

lights off, was applied daily in the early afternoon for one week. Depending upon the precise

targeting of the fiber optics, severe seizures would typically be triggered upon the first

stimulation of the day beginning on day 2-3 if not on the first day. Seizures were reliably induced

every day after the first successful induction.

Tissue processing and immunohistochemistry. The mice were perfused with 1 M phosphate-

buffered saline (PBS) and 4% paraformaldehyde (PFA). The brain was then carefully dissected

and submerged in 4% PFA for 1-2 days. Following serial sucrose protection (15-30% sucrose in

PBS), samples were submerged in OCT and immediately frozen at -80C. For

immunohistochemistry, free-floating sections cut on the cryostat were incubated in antibodies in

a solution of 10% normal goat or donkey serum and 0.01% Tween-20. For DAB staining,

sections were first bathed in H2O2 to quench endogenous peroxidases and then washed 3 times

in PBS for 5 min each. For all stains, sections were blocked for 2 h with the blocking solution at

room temperature and then incubated in primary antibodies overnight at room temperature.

Sections were then washed with PBS 3 times for 5 min each and then incubated in secondary

antibodies for 2 h. Sections were washed again 3 times in PBS for 5 min each and mounted onto

slides. DAB slides were air-dried overnight and then dehydrated in ethanol before cover-

slipping. Slides with fluorescent signal were immediately cover-slipped using FluoroGel as a

medium. Antibodies used for immunohistochemistry (primary antibodies) were: calbindin (Rb

and mouse (Ms), Swant, 1:10,000); NeuN (Rb, Millipore, 1:500); VGLUT2 (Rb and Gp,

Synaptic Systems; 1:1,000); VGLUT1 (Ms and Gp, Synaptic Systems, 1:1,000); NFH (Ms,

Covance, 1:1,500); TH (Rb, Millipore, 1:500); GFP (Ch, Abcam, 1:1,000); parvalbumin (Rb,

Swant, 1:1,000); calretinin (Rb, Swant, 1:500); Secondary antibodies were: for the DAB

reaction, we used horseradish peroxidase-conjugated goat anti-rabbit and goat anti-mouse

secondary antibodies (diluted 1:200 in PBS; DAKO); staining for fluorescent

immunohistochemistry was performed using donkey anti-mouse, anti-rabbit or anti-guinea pig

secondary antibodies conjugated to Alexa 488, 555 or 647 fluorophores (Invitrogen), all diluted

to 1:1,500. Photomicrographs of tissue sections were captured using Zeiss AxioCam MRm

(fluorescence) and AxioCam MRc5 (DAB-reacted tissue sections) cameras mounted on a Zeiss

Axio Imager.M2 microscope or on a Zeiss Axio Zoom.V16. Images of the tissue sections were

acquired and analyzed using either Zeiss AxioVision software (release 4.8) or Zeiss ZEN

software (2012 edition). After imaging, the raw data were imported into Adobe Photoshop CS5

and corrected for brightness and contrast levels. Schematics were drawn in Adobe Illustrator CS5

and then imported into Adobe Photoshop CS5 to contract the figure.

Basic histology: H&E staining. For H&E staining, tissue sections were mounted on a polarized

slide and dried overnight. Next, the slide was submerged in xylene for 5 minutes. To then

rehydrate the tissue, the sections were placed in serial dilutions of ethanol for 2-5 minutes as

follows: 100% ethanol, 95% ethanol, 70% ethanol, DI water. Following this rehydration step,

slides were placed in hematoxylin staining for approximately 1 minute followed by a water bath

for several rounds of washing. Finally, tissue was placed in an eosin stain for 1 minute, rinsed in

a water bath, and dehydrated in 70%, 95%, and three rounds 100% ethanol followed by xylene.

Slides were then coverslipped with Cytoseal Mounting Media (Fisher Scientific) and left to dry
overnight.

In vivo electrophysiology. For head-fixed, awake recordings, mice were implanted with custom-
made headplates and a craniotomy made above the VPM. A fiber optic was also implanted at a
45-degree angle to target the VPM underneath the headplate. After 72 h of recovery, the mice
were trained for 30 min per day in a head-fixed apparatus for 3 days before recording. Neuronal
spikes were recorded and categorized based on standard stereotaxic coordinates measured from
bregma and firing pattern. We examined a total of 70 units in 5 animals. All recordings were
collected using ~10-20MΩ pulled glass pipettes filled with 0.9% saline. The recorded signals
were digitized at a sampling rate of 5000Hz into Spike2 (CED, England) where single units were
verified with principal components analysis.

Quantifications and Analyses. All data was analyzed and graphed in Prism9.1.0 and p values
were considered significant at values less than 0.05. For differences in relative fluorescence,
thalamic nuclei were segmented in Photoshop and their mean gray values, area, and integrated
densities were calculated. A region of layer 6 cortex was used as reference for each slice. Data
was then transferred to Prism, graphed, and Dunnett's multiple comparisons test performed with
the VPM region used as a control for comparison. All EEG data was collected in Spike2 and
analyzed in MATLAB (code available upon request), except in the case of power analyses which
were exported from Spike2 and plotted in Prism. Statistics were computed using paired t-tests on
area under the curve. Cell count (on H&E stained sections) and dFosB quantifications were
performed by first thresholding, converting to binary images, and automatically counting
positively stained cells in ImageJ before transferring data to Prism for unpaired t-tests. For
lidocaine experiments, presence of seizures was visually scored as 1 (seizure) or 0 (no seizure)
for a train of 30 second, 30-Hz stimulations before lidocaine administration and beginning 5
minutes after administration. For washout experiments, mice were returned to their home cages
and re-evaluated with optogenetic stimulation and visual scoring 1 hour after initial lidocaine
administration. These data were then plotted and analyzed via Dunnett's multiple comparisons
test in Prism.

Data availability. Data generated in the experiments presented in the current study are available

from the corresponding author upon request and also available in the data tables provided.

 Figure Legends

Figure 1: Ntsr1^{Cre} is expressed in corticothalamic layer 6, cerebellar nuclei, and the axons and
fibers of passage within the thalamic nuclei. (a) Schematic of Sun1 expression, which labels the
nuclear envelope of *Cre*-expressing neurons. (b) Expression of Sun1 is seen in layer 6 of the
cerebral cortex (black box, high power image in (c)), but not in thalamic nuclei. (c) High power
image of cortical neurons expressing nuclear Sun1. Scale bar = 20um. (d) Cerebellar nuclei
neurons in the dentate and interposed nuclei express Sun1. (e) High-power image of Sun1 in the
cerebellar nuclei, showing the specificity of fluorescence for the nuclear membrane. Scale bar =
100um. (f) Schematic of Tdtomato expression, which fills the entire cell including dendrites and
axonal processes. (g) Expression of Tdtomato is observed in layer 6 of the cerebral cortex (black
box, high power image in (h)) as well as in the thalamus. However, in context with the lack of
Sun1 expression in (b), this thalamic TdTomato signal can be identified as input from other *Cre*-
expressing areas. (h) High power image demonstrating TdTomato expression in cortical layer 5/6
neurons. Scale bar = 100um. (i) Cerebellar nuclei neurons in the dentate and interposed nuclei
express TdTomato. (j) High power magnification of cerebellar nuclei neurons with Tdtomato
fluorescence in the somas as well as in axons exiting the cerebellum. Scale bar = 100um.

Figure 2: Optogenetic modulation of thalamus instead targets input from extra-thalamic regions.
(a) Because the local thalamic neurons themselves do not express *Cre* in *Ntsr1^{Cre}* animals,
manipulation of their activity is indirect by modulating *Cre*-expressing axons and fibers of
passage from regions outside the thalamus itself. Images of brain regions adapted and modified
from BrainExplorer2, Allen Brain Atlas. (b-e) High power images of *Cre*-induced reporter
expression patterns in the thalamus (scale bar = 50um) (b), cerebellar nuclei (scale bar = 100um)
(c), cortex (scale bar = 100um) (d), and striatum (scale bar = 50um) (e). Note the distinction of
the expression pattern of the thalamus compared to other regions (b), where fibers but not the
resident neurons have expression and these areas are depicted by absence of localized
fluorescence (arrows).

Figure 3: Seizure-inducing stimulations are specific to the VPM. (a) Representative
segmentation of thalamic nuclei (coronal) showing relative fluorescence of various nuclei in
Ntsr1^{Cre};ROSA$^{lox-stop-lox-ChR2-YFP}$ mice. (b) Quantification of fluorescent intensity of various
thalamic nuclei, some of which are segmented in (a), demonstrates that the specificity of VPM as

a seizure locus is not due to biased *Cre* expression, as no behavior is elicited even when VPL is
targeted. A Dunnett's multiple comparison's test determined significance between various
thalamic nuclei compared to the VPM as follows: VPM vs PO ($p<0.0001$), VPM vs LD
($p<0.0001$), VPM vs LP ($p<0.0001$), VPM vs CL ($p<0.0001$), VPM vs VM ($p<0.0001$), VPM vs
VPL (ns), VPM vs RT ($p=0.0156$), VPM vs VAL ($p<0.0001$), VPM vs LGd (ns). (c) Example of
post-hoc targeting of fiber optics to the VPM as demonstrated both by the electrode track and an
area of photobleaching of the ChR2-YFP signal. (d) Targeting of the fiber optics are examined in
mice at the end of experimentation. Each red dot represents the end of a fiber in a single mouse
that successfully exhibited stage 6 seizures. Grey dots depict targeting that did not elicit seizures.
(e) Example of the root mean square of left cortical EEG in a single mouse during multiple
seizure stimulation periods (blue overlays, 30s stimulations). Each subsequent stimulation period
begins 2 minutes following the termination of the previous stimulation. Behavioral seizures are
marked by the yellow rectangles, which coincide with high amplitude EEG activity. Here, we see
that stimulations immediately following a previously elicited seizure (within 2 minutes) do not
induce subsequent seizures. Instead, mice experience a refractory period in which additional
delivery of light does not result in behavioral abnormalities. Scale bars mark two instances in
which the mouse only experiences a seizure 4 minutes and 6 minutes following a previous
seizure, but not 2 minutes after a seizure.

Figure 4: Optogenetic stimulation of the VPM mimics tonic-clonic seizures. (a) Still time-lapse
images of the seizure phenotype in the optogenetic (above) and kainic acid (below) models of
seizure induction demonstrate parallels in behavior beginning with repetitive facial and forelimb
clonus and ending in a loss of posture during convulsions. Arrows point to examples of clonus
and an erect, straub tail, both defining features of rodent seizures. (b) Schematic of electrode
placement for EcoG (EEG) recordings based on stereotaxic coordinates from the skull landmarks
bregma and lambda. Brain overlay adapted from Allen Brain Atlas. (c) EEG signatures during
behavioral seizures (boxed region) are synchronous between cortex (blue) and cerebellum (red).
This pattern is consistent between optogenetic (above) and kainic acid (below)-evoked seizures.

Figure 5: EEG analysis confirms seizure induction. (a) Example of EEG signal from all four recorded brain regions shows high amplitude, synchronous activity during the optogenetic stimulation period (blue box) (b) An example of two superimposed signals (yellow and purple) of averaged EEG (somatosensory cortex, left and right motor cortices, and cerebellum) within a single mouse before, during, and after optogenetic stimulation. Blue shaded region indicates duration of a 30Hz train of light delivered through a fiber optic targeted to the unilateral VPM. Vertical dotted lines demonstrate time at which behavioral seizure is first visualized when both instances are aligned by optogenetic start time. The first plot shows two examples of a stage 6 seizure, whereas the second plot shows two overlaid examples of a stage 2 seizure where only forelimb clonus was observed. The third plot scales the y-axis of the stage 2 seizures to show the change in amplitude from baseline. Black graphed line overlaid on all three plots represents the average root mean square of the EEG signal for both superimposed seizure cases, showing a clear increase from baseline specifically during the behavioral seizure. (c) Example of coherence plots from a single mouse based on the raw EEG signals from four brain regions. The data shows the greatest increase in coherence from before the seizure (gray) to during the seizure (blue) between the left somatosensory cortex/left motor cortex and the left motor cortex/cerebellum. (d) Time frequency heat-plot averaged across mice shows the power of dominant high-power frequencies of EEG shifts from the site of seizure origin (left motor cortex, which is positioned above the VPM) to the contralateral hemisphere (right motor cortex). Time = 0s is aligned to the start of optogenetic stimulation. At 30s, the stimulation is terminated but behavioral and EEG seizure continues independent of the light source and is sustained in cerebellum and left somatosensory cortex. N=7 mice. (e-f) Averaged plot of the power distribution before stimulation (black) versus during stimulation (blue) demonstrate that the most robust changes occur below 100Hz. The same data is shown on a log scale in (f). N=4 mice

Figure 6: Single cell frequency is altered in the VPM during seizure activity. (a) Schematic of the optogenetic setup. We target the VPM (thalamus, green) with a fiber implanted at an angle and create a craniotomy above the cortex to access the VPM with a glass-pulled electrode for recordings. The mouse itself remains awake and is head-fixed on a rotating wheel, which allows for movement during recordings. (b) Short, subthreshold-seizure stimulations (2 second optogenetic deliveries, blue overlays) elicit reliable entraining of VPM neurons characterized by strict excitation. (c-g) Delivery of seizure-inducing trains of 30Hz light (30 second optogenetic

deliveries, blue overlays) elicits heterogeneous responses in thalamic neurons including discreet

biphasic (c), repetitive biphasic (d,e), strict excitation (f), and no change in activity during

seizures. (h) Quantification of neural responses to seizure behavior, n=5 mice, 71 cells

Figure 7: Optogenetic seizure induction does not result in gross cell loss but it does induce a

local increase in dFosB expression. (a) Quantification of total cell count using H&E staining

demonstrated no cell loss in the thalamus of control versus seizure animals (unpaired t-test,

p=0.4627). (b) Quantification of dFosB expression in control animals and seizure animals, both

in the ipsilateral thalamus and contralateral thalamus compared to stimulation, demonstrated a

graded increase in expression from the contralateral to ipsilateral thalamus. This demonstrates

that the most significant changes occurred at the origin of seizure initiation, but also occurred

(though to a lesser degree) as seizures spread to the contralateral hemisphere. (Tukey's multiple

comparisons test: control versus seizure contralateral thalamus, p=0.002; control versus seizure

ipsilateral thalamus p<0.0001; seizure contralateral versus seizure ipsilateral thalamus, p<0.0001.

(c-f) Representative images of DAB visualized immunohistochemical staining of dFosB and

their corresponding binary images within the contralateral (c,e) and (d,f) ipsilateral thalamus.

Figure 8: Cre-dependent viral retrograde tracing from the VPM reveals input primarily from the

cerebral cortex and cerebellar nuclei. (a) Schematic of the experimental setup. *Ntsr1^{Cre}* mice

received unilateral injections with varying volumes of pAAV-Ef1a-flex-RUBY ranging from

138nL to 680nL into the left VPM of the thalamus. Virus taken up by local VPM neurons was

monosynaptically traced in the retrograde direction to the source neurons that send the

projections. In these mice, only the cerebral cortex (b, scale bar = 100um) and cerebellar nuclei

(c, scale bar = 50um, d, e, scale bar = 50um) demonstrated a strong marker signal after the

injections.

Figure 9: Pharmacological inactivation of individual Cre-expression inputs to the thalamus

reveals the requirement of the cerebellum for seizure initiation. (a) Schematic of the

experimental setup for cerebellar ablation includes a surgically implanted fiber optic targeted to

the VPM and bilateral cannulas targeted to the interposed cerebella nuclei for the purpose of

lidocaine delivery. (b) Schematic of the experimental setup for cerebral cortical pharmacological

ablation includes a surgically implanted fiber optic targeted to the VPM and bilateral cannulas

targeted to the somatosensory or motor cortices for lidocaine delivery. (c) Example of a cannula
targeting to the interposed cerebellar nucleus in a 40-um coronal slice. (d) Quantifications of
scored seizures after providing trains of 30Hz blue light pulses to the VPM before lidocaine
delivery, immediately following delivery, and after lidocaine washout. A significant decrease in
ability to elicit seizures occurred following cerebellar interposed nucleus ablation (Dunnett's
multiple comparisons test, p<0.0001), but not somatosensory cortex (p=0.9562) or motor cortical
(p=0.8835) ablation.

936
937

S1: Whole brain expression of the Ntsr1^{Cre} *transgene.* (ai-av) Coronal sections of
Ntsr1^{Cre};ROSA$^{lox-stop-lox-Sun1-GFP}$ mice show nuclear envelope staining primarily in cortical layer
5/6, the caudoputamen, and the cerebellar nuclei. However, there is also noticeable expression in
the substantia nigra/VTA and brainstem nuclei. Although there are some *Ntsr1^{Cre}* positive
neurons in the hippocampus and medial thalamic nuclei, these are sparse, few, and unlikely to
affect optogenetic stimulation paradigms. (bi-bvi) Sagittal sections cut through *Ntsr1^{Cre};ROSA^{lox-}*
$^{stop-lox-TdTomato}$ mouse brains highlight the robust fluorescence in thalamic nuclei stemming
exclusively from incoming axons and fibers of passage, one example of which is seen in panel
(c), a dense bundle of *Ntsr1^{Cre}* positive axons exiting the cerebellum. Segmented atlas images
(adapted from Allen Brain Atlas) are included for reference.

S2: Characterization of Ntsr1^{Cre} positive neurons in the cortex and VTA. (a-c) Co-staining of
NeuN in the cerebral cortex of *Ntsr1^{Cre};ROSA$^{lox-stop-lox-TdTomato}$* mice demonstrate specificity of
TdTomato signal to layer 6 cell bodies. Scale bar = 100um. (d-f) Calbindin staining, which labels
nonpyramidal cells in the cortex, does not colocalize with TdTomato positive cells. Scale bar =
100um. (g-i) In the VTA, TdTomato-positive cells colocalize with tyrosine hydroxylase (TH),
indicating that they are dopaminergic (white arrows). However, not all TH-positive cells are
TdTomato positive (orange arrows), indicating that the *Ntsr1^{Cre}* allele is not expressed in all
VTA neurons. Scale bar = 50um.

S3: Ntsr1^{Cre} expression in the cerebellum is localized to excitatory cerebellar nuclei neurons,
scattered Purkinje cells, and a small population of molecular layer interneurons. (a-f) Calbindin
staining (green) demonstrates the relationship between GABAergic projections from Purkinje
cells to *Ntsr1^{Cre}*-positive cerebellar nuclei neurons (red). Scale bars are 50um (c) and 20um (f).

(j-o) Neurofilament heavy chain (NFH, green) colocalizes with TdTomato neurons in the
interposed and dentate nucleus, identifying these cells as excitatory projection neurons. Scale
bars are 50um (l) and 20 um (o). (p) *Ntsr1^{Cre};TdTomato* coronal section of the cerebellum shows
sparse labeling of individual Purkinje cells (arrows). (q) A single *Ntsr1^{Cre}*-positive Purkinje cell
shows intense expression in the soma and throughout the dendritic tree. Scale bar = 20um. (r)
Ntsr1^{Cre} positive Purkinje cells also express the Purkinje cell marker Plcb4. Scale bar = 100um.
(s-u) RORa staining in lobules IX and X demonstrates that some (blue arrows), but not all
(orange arrows), molecular layer interneurons (basket cells and stellate cells) express the
Ntsr1^{Cre}-transgene. Scale bar = 100um.

S4: Ntsr1^{Cre} *is not expressed in cell bodies of thalamic neurons.* (a-i) Serially magnified images
of calbindin staining in the thalamus, which is differentially expressed in various thalamic nuclei,
demonstrates no colocalization between TdTomato (red) and calbindin-positive (green) thalamic
neurons. Scale bars are 500um (c), 50um (f), and 20um (i). (j-o) Staining with a pan neuronal
marker, NeuN, does not colocalize with TdTomato signal in the thalamus. Instead, *Ntsr1^{Cre}* is
expressed in the space around resident thalamic cell bodies (k, n) as local axons or fibers of
passage originating from other *Cre*-positive brain regions. Scale bars are 100um (l) and 50um
(o).

S5: High amplitude synchrony in EEG activity is consistent across mice during optogenetically-
induced seizures. Here, we show representative examples from 7 mice that all exhibited stage 6
seizures during optogenetic stimulation (blue overlayed box). A hallmark electrophysiological
signature of seizure activity, high amplitude synchrony, is consistently exhibited in the left motor
cortex (blue), right motor cortex (green), left somatosensory cortex (orange), and cerebellum
(red).

S6: Coherence between EEG activity of left motor and left somatosensory cortex and cerebellum
are most consistently increased during seizure activity. Examples of EEG coherence between
brain regions from three representative mice. Here, a consistent pattern of increased coherence is
seen between the before seizure (gray) and during seizure (blue) periods. (a) A large increase in
coherence between left motor and left somatosensory cortex is particularly evident between 0
and 50 Hz. (b) Although coherence does increase between the right motor cortex (contralateral to

the original stimulation) and the rest of the brain (d, e), this change is marginal compared to that
seen in (a) and (c). (c) The left motor cortex and central cerebellum exhibit similar increases in
coherence between 0 and 50Hz as that between left somatosensory and motor cortices.
993
S7: Power spectra with optimized scaling in different brain regions reveals patterned shift in
activity. (a, c) Power-frequency plots from two mice experiencing seizures during the stimulation
period (t = 0-30s). Due to the limitations of a set scale across brain regions, changes in activity
during this period are largely obscured in several regions in Subjects 1 and 2. Here, we compare
a hard/set color scale versus an optimized scale where every region receives its own color scaling
according to its maximum power within the plotted frequency range. Each scaling approach
reveals unique changes in activity. We show specific examples of heat plots from EEG
frequencies in the delta, theta, alpha, beta, gamma, 60-120Hz, and 120-200Hz ranges. White
arrows point to power changes that were previously masked by the set or optimized scaling.
1003
S8: Power of frequencies under 100Hz is significantly increased during seizures. (a-d) Power
plots of EcoG frequencies from 0-250Hz is shown for all four regions measured. In each region,
there is a dramatic increase in the power of all frequencies between 0 and 100Hz. (e-g)
Compilation of all areas onto one plot shows a comparison of changes in each brain region,
demonstrating that this shift is most dramatic in the ipsilateral motor/somatosensory cortices and
cerebellum, as expected due to the proximity to the site of seizure origin (left VPM).
1010
1011 *Video1: Optogenetic delivery of 30Hz light to the VPM elicits tonic-clonic seizures.* Video
1012 representation of a stage 6 seizure showing examples of the progression from facial and forelimb
1013 clonus to loss of posture in a mouse exhibiting a light-induced seizure. All mice, including those
1014 with severe phenotypes, experience a full recovery of general behavior after the seizures resolve.
1015
1016 *Video2: Tonic-clonic seizures that are induced with optogenetics are consistent and robust.*
1017 Examples of multiple mice experiencing phenotypically similar TC-seizures as a result of the
1018 unilateral, VPM-targeted 30Hz light stimulation paradigm.
1019

Video3: Unilateral optogenetic stimulation of the VPM is sufficient to elicit the full seizure

phenotype. Neither bilateral VPM nor co-stimulation of the VPM and cerebellar nuclei, two

heavy *Cre*-expressing regions, amplifies the seizure phenotype. Therefore, to simplify surgeries

and avoid unnecessary stimulation, we use unilateral light delivery to induce the seizures. Here

we show an example of co-stimulation of the VPM and the contralateral interposed cerebellar

nuclei.

1026

1027   *Video4: Optogenetic seizure induction is specific to the thalamic VPM*. Stimulation of

1028   surrounding thalamic nuclei, all *Cre*-expressing, does not elicit any overt behavioral phenotype.

1029   In this example, we show the effects of 30-Hz light delivery to the ventral posterolateral nucleus,

1030   which receives some common input with the VPM and is directly adjacent to the seizure-

1031   inducing region.

1032

1033   *Video5: Behavioral similarities between kainic acid-induced and optogenetic-induced seizures.*

1034   The behavioral phenotype of kainic-acid induced seizures and the optogenetic model are

1035   identical. Both exhibit forelimb and facial clonus, straub tail, and loss of body posture.

1036

1037   *Video6: Abnormal EEG activity is correlated with behavioral seizure.* High amplitude EEG

1038   activity that synchronizes across brain regions becomes evident as the behavioral seizure begins.

1039   For those seizures that outlast the period of light delivery, the EEG becomes increasingly erratic.

1040   When the mice recover, the EEG amplitude returns to baseline values.

1041

1042   *Video7: Pharmacological ablation of the cerebellar nuclei eliminates the ability to induce*

1043   *seizures.* Before administration of lidocaine, the mouse exhibits a severe TC-seizure upon light

1044   delivery to the VPM. After bilateral lidocaine delivery to the interposed cerebellar nuclei, light

1045   stimulation does not evoke any overt behavioral changes. However, after lidocaine washout, the

1046   seizure can once again can be elicited by light stimulation directed into the VPM.

1047

Table 1: Quantification of fluorescence among thalamic nuclei demonstrates regional

differences in channelrhodopsin-expressing input. Despite differences in fluorescent intensity

between different thalamic nuclei such as the PO, CL, and RT, other nuclei such as the VPL do

not demonstrate significant differences in levels of fluorescence when compared to the VPM.

The differences in ChR2 expression alone, therefore, likely do not account for the lack of seizure

behavior upon targeting of the different thalamic nuclei with fiber optics.

1054

1055   *Table 2: Modified Racine Scale is used to categorize and rank seizure behavior.* Using a

1056   modified Racine Scale, we compared the severity of seizure phenotypes by ranking different

1057   categorical behavioral attributes of the mice upon seizure induction with optogenetic stimulation.

1058

1059

1060

1061

References

1. Cockerell, O. C., Sander, J. W. A. S., Hart, Y. M., Shorvon, S. D. & Johnson, A. L. Remission of epilepsy: results from the National General Practice Study of Epilepsy. *Lancet* **346**, 140–144 (1995).

2. Mattson, R. H., Cramer, J. A. & Collins, J. F. Prognosis for total control of complex partial and secondarily generalized tonic clonic seizures. *Neurology* **47**, 68–76 (1996).

3. Kwan, P. & Brodie, M. J. Early Identification of Refractory Epilepsy. *N. Engl. J. Med.* **342**, 314–319 (2000).

4. Elger, C. E. Pharmacoresistance: Modern concept and basic data derived from human brain tissue. in *Epilepsia* **44**, 9–15 (Blackwell Publishing Inc., 2003).

5. Jovel, C. A. E. & Sobrino-Mejía, F. E. Drug resistant epilepsy. Clinical and neurobiological concepts. *Revista de Neurologia* (2015). doi:10.33588/rn.6104.2015181

6. Callaghan, B. C., Anand, K., Hesdorffer, D., Hauser, W. A. & French, J. A. Likelihood of seizure remission in an adult population with refractory epilepsy. *Ann. Neurol.* **62**, 382–389 (2007).

7. Berg, A. T. *et al.* How long does it take for partial epilepsy to become intractable? *Neurology* **60**, 186–190 (2003).

8. Mohanraj, R. & Brodie, M. J. Diagnosing refractory epilepsy: Response to sequential treatment schedules. *Eur. J. Neurol.* (2006). doi:10.1111/j.1468-1331.2006.01215.x

9. Choi, H. *et al.* Seizure remission and relapse in adults with intractable epilepsy: A cohort study. *Epilepsia* (2008). doi:10.1111/j.1528-1167.2008.01601.x

10. FACTS AND FIGURES | American Epilepsy Society. Available at: https://www.aesnet.org/for_patients/facts_figures#Six. (Accessed: 14th January 2021)

11. Kammerman, S. & Wasserman, L. Seizure disorders: Part 1. Classification and diagnosis. *West. J. Med.* **175**, 99–103 (2001).

12. Meeren, H. K. M., Pijn, J. P. M., Van Luijtelaar, E. L. J. M., Coenen, A. M. L. & Da Silva, F. H. L. Cortical focus drives widespread corticothalamic networks during spontaneous absence seizures in rats. *J. Neurosci.* (2002). doi:10.1523/jneurosci.22-04-01480.2002

13. GLOOR, P. Generalized Cortico-Reticular Epilepsies Some Considerations on the

1093   Pathophysiology of Generalized Bilaterally Synchronous Spike and Wave Discharge.

1094   *Epilepsia* (1968). doi:10.1111/j.1528-1157.1968.tb04624.x

1095   14.   Buompadre, M. C. Malformations of cortical development and epilepsy. *Medicina (B.*

1096   *Aires).* (2019). doi:10.31887/dcns.2008.10.1/rjleventer

1097   15.   Jan, M. M. S. & Girvin, J. P. Seizure semiology: Value in identifying seizure origin.

1098   *Canadian Journal of Neurological Sciences* (2008). doi:10.1017/S0317167100007526

1099   16.   Roopun, A. K. *et al.* Detecting seizure origin using basic, multiscale population dynamic

1100   measures: Preliminary findings. *Epilepsy Behav.* (2009). doi:10.1016/j.yebeh.2008.09.008

1101   17.   Dabrowska, N. *et al.* Parallel pathways of seizure generalization. *Brain* (2019).

1102   doi:10.1093/brain/awz170

1103   18.   Norden, A. D. & Blumenfeld, H. The role of subcortical structures in human epilepsy.

1104   *Epilepsy Behav.* **3**, 219–231 (2002).

1105   19.   Marcián, V., Filip, P., Bareš, M. & Brázdil, M. Cerebellar Dysfunction and Ataxia in

1106   Patients with Epilepsy: Coincidence, Consequence, or Cause? *Tremor Other Hyperkinet.*

1107   *Mov. (N. Y).* **6**, 376 (2016).

1108   20.   Blumenfeld, H. The thalamus and seizures. *Archives of Neurology* (2002).

1109   doi:10.1001/archneur.59.1.135

1110   21.   Caciagli, L. *et al.* Thalamus and focal to bilateral seizures. *Neurology* (2020).

1111   doi:10.1212/wnl.0000000000010645

1112   22.   Li, Y. H. *et al.* Involvement of thalamus in initiation of epileptic seizures induced by

1113   pilocarpine in mice. *Neural Plast.* (2014). doi:10.1155/2014/675128

1114   23.   Polack, P. O. *et al.* Deep layer somatosensory cortical neurons initiate spike-and-wave

1115   discharges in a genetic model of absence seizures. *J. Neurosci.* (2007).

1116   doi:10.1523/JNEUROSCI.0753-07.2007

1117   24.   Meeren, H. K. M., Pijn, J. P. M., Van Luijtelaar, E. L. J. M., Coenen, A. M. L. & Lopes da

1118   Silva, F. H. Cortical focus drives widespread corticothalamic networks during

1119   spontaneous absence seizures in rats. *J. Neurosci.* (2002).

1120   25.   Meeren, H., van Luijtelaar, G., Lopes da Silva, F. & Coenen, A. Evolving Concepts on the

1121   Pathophysiology of Absence Seizures. *Arch. Neurol.* (2005).

1122   doi:10.1001/archneur.62.3.371

26.   Polack, P. O., Mahon, S., Chavez, M. & Charpier, S. Inactivation of the somatosensory

cortex prevents paroxysmal oscillations in cortical and related thalamic neurons in a

genetic model of absence epilepsy. *Cereb. Cortex* (2009). doi:10.1093/cercor/bhn237

27. Nersesyan, H., Hyder, F., Rothman, D. L. & Blumenfeld, H. Dynamic fMRI and EEG

recordings during spike-wave seizures and generalized tonic-clonic seizures in WAG/Rij

rats. *J. Cereb. Blood Flow Metab.* (2004). doi:10.1097/01.WCB.0000117688.98763.23

28. Sysoeva, M. V., Lüttjohann, A., van Luijtelaar, G. & Sysoev, I. V. Dynamics of

directional coupling underlying spike-wave discharges. *Neuroscience* (2016).

doi:10.1016/j.neuroscience.2015.11.044

29. Martín-López, D. *et al.* The Role of Thalamus Versus Cortex in Epilepsy: Evidence from

Human Ictal Centromedian Recordings in Patients Assessed for Deep Brain Stimulation.

Int. J. Neural Syst. (2017). doi:10.1142/S0129065717500101

30. Ren, K. Trigeminothalamic Tract Projections. in *Encyclopedia of Pain* (2013).

doi:10.1007/978-3-642-28753-4_4626

31. Henssen, D. J. H. A. *et al.* New insights in trigeminal anatomy: A double orofacial tract

for nociceptive input. *Front. Neuroanat.* (2016). doi:10.3389/fnana.2016.00053

32. Friedberg, M. H., Lee, S. M. & Ebner, F. F. The contribution of the principal and spinal

trigeminal nuclei to the receptive field properties of thalamic VPM neurons in the rat. *J.*

Neurocytol. (2004). doi:10.1023/B:NEUR.0000029649.28599.a5

33. Cerkevich, C. M., Qi, H. X. & Kaas, J. H. Corticocortical projections to representations of

the teeth, tongue, and face in somatosensory area 3b of macaques. *J. Comp. Neurol.*

(2014). doi:10.1002/cne.23426

34. Seneviratne, U., Cook, M. & D'Souza, W. The electroencephalogram of idiopathic

generalized epilepsy. *Epilepsia* (2012). doi:10.1111/j.1528-1167.2011.03344.x

35. Paz, J. T. *et al.* Closed-loop optogenetic control of thalamus as a tool for interrupting

seizures after cortical injury. *Nat. Neurosci.* (2013). doi:10.1038/nn.3269

36. Harvey, A. S. *et al.* Hemifacial seizures and cerebellar ganglioglioma: An epilepsy

syndrome of infancy with seizures of cerebellar origin. *Ann. Neurol.* (1996).

doi:10.1002/ana.410400115

37. Chae, J. H. *et al.* Hemifacial seizure of cerebellar ganglioglioma origin: Seizure control by

tumor resection. *Epilepsia* (2001). doi:10.1046/j.1528-1157.2001.43398.x

38. Cooper, I. S. EFFECT OF CHRONIC STIMULATION OF ANTERIOR CEREBELLUM

ON NEUROLOGICAL DISEASE. *The Lancet* **301**, 206 (1973).

39.   Cooper, I. S. & Upton, A. R. M. Therapeutic implications of modulation of metabolism

and functional activity of cerebral cortex by chronic stimulation of cerebellum and

thalamus. *Biol. Psychiatry* (1985). doi:10.1016/0006-3223(85)90164-7

40.   Lehesjoki, A. E. Molecular background of progressive myoclonus epilepsy. *EMBO*

Journal (2003). doi:10.1093/emboj/cdg338

41.   van den Ende, T., Sharifi, S., van der Salm, S. M. A. & van Rootselaar, A.-F. Familial

Cortical Myoclonic Tremor and Epilepsy, an Enigmatic Disorder: From Phenotypes to

Pathophysiology and Genetics. A Systematic Review. *Tremor Other Hyperkinet. Mov. (N.*

Y). **8**, 503 (2018).

42.   Li, Y. *et al.* Cerebellum abnormalities in idiopathic generalized epilepsy with generalized

tonic-clonic seizures revealed by diffusion tensor imaging. *PLoS One* (2010).

doi:10.1371/journal.pone.0015219

43.   Ming, X., Prasad, N., Thulasi, V., Elkins, K. & Shivamurthy, V. K. N. Possible

contribution of cerebellar disinhibition in epilepsy. *Epilepsy and Behavior* **118**, (2021).

44.   Foit, N. A. *et al.* Lesional cerebellar epilepsy: a review of the evidence. *Journal of*

Neurology (2017). doi:10.1007/s00415-016-8161-9

45.   Rubio, C. *et al.* Increase Signaling of Wnt/β-Catenin Pathway and Presence of Apoptosis

in Cerebellum of Kindled Rats. *CNS Neurol. Disord. - Drug Targets* (2017).

doi:10.2174/1871527316666170117114513

46.   Oyegbile, T. O. *et al.* The nature and extent of cerebellar atrophy in chronic temporal lobe

epilepsy. *Epilepsia* (2011). doi:10.1111/j.1528-1167.2010.02937.x

47.   Miller, J. W., Gray, B. C. & Turner, G. M. Role of the Fastigial Nucleus in Generalized

Seizures as Demonstrated by GABA Agonist Microinjections. *Epilepsia* (1993).

doi:10.1111/j.1528-1157.1993.tb02121.x

48.   Gioanni, Y., Gioanni, H. & Mitrovic, N. Seizures can be triggered by stimulating non-

cortical structures in the quaking mutant mouse. *Epilepsy Res.* (1991). doi:10.1016/0920-

1211(91)90043-F

49.   Kros, L., Eelkman Rooda, O. H. J., De Zeeuw, C. I. & Hoebeek, F. E. Controlling

Cerebellar Output to Treat Refractory Epilepsy. *Trends Neurosci.* **38**, 789–799 (2015).

50.   Kros, L. *et al.* Cerebellar output controls generalized spike-and-wave discharge

occurrence. *Ann. Neurol.* **77**, 1027–1049 (2015).

51.   Krook-Magnuson, E., Szabo, G. G., Armstrong, C., Oijala, M. & Soltesz, I. Cerebellar

1188   Directed Optogenetic Intervention Inhibits Spontaneous Hippocampal Seizures in a

1189   Mouse Model of Temporal Lobe Epilepsy. *eNeuro* **1**, 1–27 (2014).

1190   52.   Streng, M. L. & Krook-Magnuson, E. Excitation, but not inhibition, of the fastigial

1191   nucleus provides powerful control over temporal lobe seizures. *J. Physiol.* (2020).

1192   doi:10.1113/JP278747

1193   53.   Eelkman Rooda, O. H. J. *et al.* Single-pulse stimulation of cerebellar nuclei stops epileptic

1194   thalamic activity. *Brain Stimul.* **14**, 861–872 (2021).

1195   54.   Kros, L., Eelkman Rooda, O. H. J., De Zeeuw, C. I. & Hoebeek, F. E. Controlling

1196   Cerebellar Output to Treat Refractory Epilepsy. *Trends Neurosci.* **38**, 789–799 (2015).

1197   55.   St.-Gelais, F., Jomphe, C. & Trudeau, L. É. The role of neurotensin in central nervous

1198   system pathophysiology: What is the evidence? *Journal of Psychiatry and Neuroscience*

1199   (2006).

1200   56.   Gong, S. *et al.* Targeting Cre recombinase to specific neuron populations with bacterial

1201   artificial chromosome constructs. *Journal of Neuroscience* (2007).

1202   doi:10.1523/JNEUROSCI.2707-07.2007

1203   57.   Gong, S. *et al.* A gene expression atlas of the central nervous system based on bacterial

1204   artificial chromosomes. *Nature* (2003). doi:10.1038/nature02033

1205   58.   Woodworth, H. L. *et al.* Neurotensin Receptor-1 Identifies a Subset of Ventral Tegmental

1206   Dopamine Neurons that Coordinates Energy Balance. *Cell Rep.* (2017).

1207   doi:10.1016/j.celrep.2017.08.001

1208   59.   Sundberg, S. C., Lindström, S. H., Sanchez, G. M. & Granseth, B. Cre-expressing neurons

1209   in visual cortex of Ntsr1-Cre GN220 mice are corticothalamic and are depolarized by

1210   acetylcholine. *J. Comp. Neurol.* **526**, 120–132 (2018).

1211   60.   Sundberg, S. C., Lindström, S. H., Sanchez, G. M. & Granseth, B. Cre-expressing neurons

1212   in visual cortex of Ntsr1-Cre GN220 mice are corticothalamic and are depolarized by

1213   acetylcholine. *J. Comp. Neurol.* (2018). doi:10.1002/cne.24323

1214   61.   Houck, B. D. & Person, A. L. Cerebellar premotor output neurons collateralize to

1215   innervate the cerebellar cortex. *J. Comp. Neurol.* **523**, 2254–2271 (2015).

62.   Sillitoe, R. V., Fu, Y. H. & Watson, C. Cerebellum. in *The Mouse Nervous System* 360–

397 (Academic Press, Elsevier, 2012). doi:10.1016/B978-0-12-369497-3.10011-1

63.   Fujita, H., Kodama, T. & Du Lac, S. Modular output circuits of the fastigial nucleus for

diverse motor and nonmotor functions of the cerebellar vermis. *Elife* (2020).

doi:10.7554/eLife.58613

64.   Miterko, L. N. *et al.* Consensus Paper: Experimental Neurostimulation of the Cerebellum.

Cerebellum (2019). doi:10.1007/s12311-019-01041-5

65.   van brederode, J. F. M., helliesen, M. K. & hendrickson, A. E. Distribution of the calcium-

binding proteins parvalbumin and calbindin-D28k in the sensorimotor cortex of the rat.

Neuroscience (1991). doi:10.1016/0306-4522(91)90258-P

66.   Arai, R., Jacobowitz, D. M. & Deura, S. Distribution of calretinin, calbindin-D28k, and

parvalbumin in the rat thalamus. *Brain Res. Bull.* (1994). doi:10.1016/0361-

9230(94)90086-8

67.   Sundberg, S. C., Lindström, S. H., Sanchez, G. M. & Granseth, B. Cre-expressing neurons

in visual cortex of Ntsr1-Cre GN220 mice are corticothalamic and are depolarized by

acetylcholine. *J. Comp. Neurol.* **526**, 120–132 (2018).

68.   Racine, R. J. Modification of seizure activity by electrical stimulation: II. Motor Seizure.

Electroencephalogr. Clin. Neurophysiol. **32**, 281–294 (1972).

69.   Pottkämper, J. C. M., Hofmeijer, J., Waarde, J. A. van & Putten, M. J. A. M. van. The

postictal state — What do we know? *Epilepsia* **61**, 1045 (2020).

70.   Abood, W. & Bandyopadhyay, S. Postictal Seizure State. *StatPearls* (2021).

71.   Ben-Ari, Y. & Lagowska, J. [Epileptogenic action of intra-amygdaloid injection of kainic

acid]. *C. R. Acad. Sci. Hebd. Seances Acad. Sci. D.* (1978).

72.   Lévesque, M. & Avoli, M. The kainic acid model of temporal lobe epilepsy. *Neuroscience*

and Biobehavioral Reviews (2013). doi:10.1016/j.neubiorev.2013.10.011

73.   Zhu, J., Zheng, X. Y., Zhang, H. L. & Luo, Q. Kainic acid-induced neurodegenerative

model: Potentials and limitations. *Journal of Biomedicine and Biotechnology* (2011).

doi:10.1155/2011/457079

74.   Biasiucci, A., Franceschiello, B. & Murray, M. M. Electroencephalography. *Current*

Biology **29**, (2019).

75.   Zafar, A. & Almajid, A. Paroxysmal dystonia masquerading as focal onset seizure in

patient with multiple sclerosis. *Saudi J. Heal. Sci.* (2016). doi:10.4103/2278-0521.195820

76.   Guerrini, R. Idiopathic epilepsy and paroxysmal dyskinesia. in *Epilepsia* (2001).

1249   doi:10.1046/j.1528-1157.2001.042suppl.3036.x

1250   77.   Löscher, W. *et al.* The sz mutant hamster: A genetic model of epilepsy or of paroxysmal

1251   dystonia? *Mov. Disord.* (1989). doi:10.1002/mds.870040304

1252   78.   Uhlhaas, P. J. & Singer, W. Neural Synchrony in Brain Disorders: Relevance for

1253   Cognitive Dysfunctions and Pathophysiology. *Neuron* (2006).

1254   doi:10.1016/j.neuron.2006.09.020

1255   79.   Farrell, J. S., Nguyen, Q.-A. & Soltesz, I. Review Resolving the Micro-Macro Disconnect

1256   to Address Core Features of Seizure Networks. (2019). doi:10.1016/j.neuron.2019.01.043

1257   80.   Quiroga, R. Q., Garcia, H. & Rabinowicz, A. Frequency evolution during tonic-clonic

1258   seizures. *Electromyogr. Clin. Neurophysiol.* 323- (2002).

1259   81.   RS, F., WR, W., RP, L., S, A. & S, U. High-frequency EEG activity at the start of

1260   seizures. *J. Clin. Neurophysiol.* **9**, 441–448 (1992).

1261   82.   Tieng, Q. M., Anbazhagan, A., Chen, M. & Reutens, D. C. Mouse epileptic seizure

1262   detection with multiple EEG features and simple thresholding technique. *J. Neural Eng.*

1263   (2017). doi:10.1088/1741-2552/aa8069

1264   83.   Cambiaghi, M., Magri, L. & Cursi, M. Importance of EEG in validating the chronic

1265   effects of drugs: Suggestions from animal models of epilepsy treated with rapamycin.

1266   *Seizure* (2015). doi:10.1016/j.seizure.2015.02.015

1267   84.   Bergstrom, R. A. *et al.* Automated identification of multiple seizure-related and interictal

1268   epileptiform event types in the EEG of mice. *Sci. Rep.* (2013). doi:10.1038/srep01483

1269   85.   Arabadzisz, D., Antal, K., Parpan, F., Emri, Z. & Fritschy, J. M. Epileptogenesis and

1270   chronic seizures in a mouse model of temporal lobe epilepsy are associated with distinct

1271   EEG patterns and selective neurochemical alterations in the contralateral hippocampus.

1272   *Exp. Neurol.* (2005). doi:10.1016/j.expneurol.2005.01.029

1273   86.   Andersen, L. M., Jerbi, K. & Dalal, S. S. Can EEG and MEG detect signals from the

1274   human cerebellum? *NeuroImage* (2020). doi:10.1016/j.neuroimage.2020.116817

1275   87.   Samuelsson, J. G., Sundaram, P., Khan, S., Sereno, M. I. & Hämäläinen, M. S.

1276   Detectability of cerebellar activity with magnetoencephalography and

1277   electroencephalography. *Hum. Brain Mapp.* (2020). doi:10.1002/hbm.24951

88.   Paola Calderon, D., Fremont, R., Kraenzlin, F. & Khodakhah, K. The neural substrates of

rapid-onset Dystonia-Parkinsonism. *Nat. Publ. Gr.* **14**, 3–5 (2011).

89.   Pan, M. K. *et al.* Cerebellar oscillations driven by synaptic pruning deficits of cerebellar

climbing fibers contribute to tremor pathophysiology. *Sci. Transl. Med.* (2020).

doi:10.1126/scitranslmed.aay1769

90.   Tsipouras, M. G. Spectral information of EEG signals with respect to epilepsy

classification. *EURASIP J. Adv. Signal Process.* **2019**, (2019).

91.   Crone, N. E., Korzeniewska, A. & Franaszczuk, P. J. Cortical gamma responses:

Searching high and low. *International Journal of Psychophysiology* **79**, (2011).

92.   Gajic, D., Djurovic, Z., Gligorijevic, J., Di Gennaro, S. & Savic-Gajic, I. Detection of

epileptiform activity in EEG signals based on time-frequency and non-linear analysis.

Front. Comput. Neurosci. (2015). doi:10.3389/fncom.2015.00038

93.   Zijlmans, M. *et al.* High-frequency oscillations as a new biomarker in epilepsy. *Ann.*

Neurol. **71**, (2012).

94.   Dümpelmann, M., Jacobs, J. & Schulze-Bonhage, A. Temporal and spatial characteristics

of high frequency oscillations as a new biomarker in epilepsy. *Epilepsia* **56**, (2015).

95.   Liu, S. *et al.* Stereotyped high-frequency oscillations discriminate seizure onset zones and

critical functional cortex in focal epilepsy. *Brain* **141**, (2018).

96.   Matsumoto, H. & Marsan, C. A. Cortical cellular phenomena in experimental epilepsy:

Ictal manifestations. *Exp. Neurol.* (1964). doi:10.1016/0014-4886(64)90026-3

97.   Cheong, E. & Shin, H. S. T-type Ca2+ channels in normal and abnormal brain functions.

Physiological Reviews (2013). doi:10.1152/physrev.00010.2012

98.   Deleuze, C. *et al.* T-type calcium channels consolidate tonic action potential output of

thalamic neurons to neocortex. *J. Neurosci.* (2012). doi:10.1523/JNEUROSCI.1362-

12.2012

99.   Liao, Y. F., Tsai, M. L., Chen, C. C. & Yen, C. T. Involvement of the Cav3.2 T-type

calcium channel in thalamic neuron discharge patterns. *Mol. Pain* (2011).

doi:10.1186/1744-8069-7-43

100. Cavazos, J. E. & Sutula, T. P. Progressive neuronal loss induced by kindling: a possible

mechanism for mossy fiber synaptic reorganization and hippocampal sclerosis. *Brain Res.*

(1990). doi:10.1016/0006-8993(90)91054-K

101. Cavazos, J. E., Das, I. & Sutula, T. P. Neuronal loss induced in limbic pathways by

1310 kindling: Evidence for induction of hippocampal sclerosis by repeated brief seizures. *J.*
1311 *Neurosci.* (1994). doi:10.1523/jneurosci.14-05-03106.1994

1312 102. Sloviter, R. S., Dean, E., Sollas, A. L. & Goodman, J. H. Apoptosis and necrosis induced
1313 in different hippocampal neuron populations by repetitive perforant path stimulation in the
1314 rat. *J. Comp. Neurol.* (1996). doi:10.1002/(SICI)1096-9861(19960311)366:3<516::AID-
1315 CNE10>3.0.CO;2-N

1316 103. Kotloski, R., Lynch, M., Lauersdorf, S. & Sutula, T. Repeated brief seizures induce
1317 progressive hippocampal neuron loss and memory deficits. in *Progress in Brain Research*
1318 (2002). doi:10.1016/S0079-6123(02)35010-6

1319 104. Zarubenko, I. I., Yakovlev, A. A., Stepanichev, M. Y. & Gulyaeva, N. V.
1320 Electroconvulsive shock induces neuron death in the mouse hippocampus: Correlation of
1321 neurodegeneration with convulsive activity. *Neurosci. Behav. Physiol.* (2005).
1322 doi:10.1007/s11055-005-0115-0

1323 105. Pretel, S., Applegate, C. D. & Piekut, D. Apoptotic and necrotic cell death following
1324 kindling induced seizures. *Acta Histochem.* (1997). doi:10.1016/S0065-1281(97)80010-4

1325 106. Bertmam, E. H. & Lothman, E. W. Morphometric effects of intermittent kindled seizures
1326 and limbic status epilepticus in the dentate gyrus of the rat. *Brain Res.* (1993).
1327 doi:10.1016/0006-8993(93)91295-4

1328 107. Brandt, C., Ebert, U. & Löscher, W. Epilepsy induced by extended amygdala-kindling in
1329 rats: Lack of clear association between development of spontaneous seizures and neuronal
1330 damage. *Epilepsy Res.* (2004). doi:10.1016/j.eplepsyres.2004.08.008

1331 108. Khurgel, M. *et al.* Activation of astrocytes during epileptogenesis in the absence of
1332 neuronal degeneration. *Neurobiol. Dis.* (1995). doi:10.1006/nbdi.1995.0003

1333 109. Tuunanen, J. & Pitkänen, A. Do seizures cause neuronal damage in rat amygdala
1334 kindling? *Epilepsy Res.* (2000). doi:10.1016/S0920-1211(99)00123-0

1335 110. Watanabe, Y. *et al.* Null mutation of c-fos impairs structural and functional plasticities in
1336 the kindling model of epilepsy. *Journal of Neuroscience* (1996). doi:10.1523/jneurosci.16-
1337 12-03827.1996

1338 111. Breier, J. I. *et al.* Effects of temporal lobe epilepsy on spatial and figural aspects of
1339 memory for a complex geometric figure. *J. Int. Neuropsychol. Soc.* (1996).
1340 doi:10.1017/s1355617700001703

112. Pedersen, B. & Dam, M. Memory disturbances in epileptic patients. *Acta Neurol. Scand.*
(1986). doi:10.1111/j.1600-0404.1986.tb04859.x

113. Dodrill, C. B. Progressive cognitive decline in adolescents and adults with epilepsy. in
Progress in Brain Research (2002). doi:10.1016/S0079-6123(02)35037-4

114. LaBar, K. S., LeDoux, J. E., Spencer, D. D. & Phelps, E. A. Impaired fear conditioning
following unilateral temporal lobectomy in humans. *J. Neurosci.* (1995).
doi:10.1523/jneurosci.15-10-06846.1995

115. Holley, A. J. & Lugo, J. N. Effects of an acute seizure on associative learning and
memory. *Epilepsy Behav.* (2016). doi:10.1016/j.yebeh.2015.11.001

116. Ulery, P. G., Rudenko, G. & Nestler, E. J. Regulation of ΔFosB stability by
phosphorylation. *J. Neurosci.* (2006). doi:10.1523/JNEUROSCI.4970-05.2006

117. Chen, J. *et al.* Induction of cyclin-dependent kinase 5 in the hippocampus by chronic
electroconvulsive seizures: Role of ΔFosB. *J. Neurosci.* (2000). doi:10.1523/jneurosci.20-
24-08965.2000

118. Morris, T. A., Jafari, N. & Delorenzo, R. J. Chronic ΔFosB expression and increased AP-1
transcription factor binding are associated with the long term plasticity changes in
epilepsy. *Mol. Brain Res.* (2000). doi:10.1016/S0169-328X(00)00112-1

119. Biagini, G. *et al.* Impaired activation of CA3 pyramidal neurons in the epileptic
hippocampus. *NeuroMolecular Med.* (2005). doi:10.1385/NMM:7:4:325

120. Curia, G., Gualtieri, F., Bartolomeo, R., Vezzali, R. & Biagini, G. Resilience to
audiogenic seizures is associated with p-ERK1/2 dephosphorylation in the subiculum of
Fmr1 knockout mice. *Front. Cell. Neurosci.* (2013). doi:10.3389/fncel.2013.00046

121. Giordano, C. *et al.* Progressive seizure aggravation in the repeated 6-Hz corneal
stimulation model is accompanied by marked increase in hippocampal p-ERK1/2
immunoreactivity in neurons. *Front. Cell. Neurosci.* (2016).
doi:10.3389/fncel.2016.00281

122. Martin, J. H. Autoradiographic estimation of the extent of reversible inactivation produced
by microinjection of lidocaine and muscimol in the rat. *Neurosci. Lett.* **127**, 160–164
(1991).

123. Demer, J. L. & Robinson, D. A. Effects of reversible lesions and stimulation of
olivocerebellar system on vestibuloocular reflex plasticity. *J. Neurophysiol.* (1982).

doi:10.1152/jn.1982.47.6.1084

124. Sandktihler, J., Maisch, B. & Zimmermann, M. The use of local anaesthetic

microinjections to identify central pathways: a quantitative evaluation of the time course

and extent of the neuronal block. *Exp. Brain Res.* **68**, 168–178 (1987).

125. Welsh, J. P. Acute inactivation of the inferior olive blocks associative learning. *Eur. J.*

Neurosci. **10**, 3321–3332 (1998).

126. White, J. J. & Sillitoe, R. V. Genetic silencing of olivocerebellar synapses causes

dystonia-like behaviour in mice. *Nat. Commun.* **8**, 14912 (2017).

127. Szabó, C. A. & Salinas, F. S. *Baboon Model of Genetic Generalized Epilepsy. Models of*

Seizures and Epilepsy: Second Edition (Elsevier Inc., 2017). doi:10.1016/B978-0-12-

804066-9.00031-6

128. Pitkänen, A., Galanopoulou, A. S., Moshé, S. L. & Buckmaster, P. S. *What Can We*

Model? Models of Seizures and Epilepsy (Elsevier Inc., 2017). doi:10.1016/B978-0-12-

804066-9.00001-8

129. Hablitz, J. J., McSherry, J. W. & Kellaway, P. Cortical seizures following cerebellar

stimulation in primates. *Electroencephalogr. Clin. Neurophysiol.* (1975).

doi:10.1016/0013-4694(75)90266-7

130. Löscher, W. Animal Models of Seizures and Epilepsy: Past, Present, and Future Role for

the Discovery of Antiseizure Drugs. *Neurochem. Res.* (2017). doi:10.1007/s11064-017-

2222-z

131. Galanopoulou, A. S., Pitkänen, A., Buckmaster, P. S. & Moshé, S. L. *What Do Models*

Model? What Needs to Be Modeled? Models of Seizures and Epilepsy: Second Edition

(Elsevier Inc., 2017). doi:10.1016/B978-0-12-804066-9.00077-8

132. The Epilepsies and Seizures: Hope Through Research | National Institute of Neurological

Disorders and Stroke. Available at: https://www.ninds.nih.gov/Disorders/Patient-

Caregiver-Education/Hope-Through-Research/Epilepsies-and-Seizures-Hope-Through.

(Accessed: 27th May 2020)

133. Löscher, W. Critical review of current animal models of seizures and epilepsy used in the

discovery and development of new antiepileptic drugs. *Seizure* **20**, 359–368 (2011).

134. Sarkisian, M. R. Overview of the Current Animal Models for Human Seizure and

Epileptic Disorders. *Epilepsy Behav.* **2**, 201–216 (2001).

135.   Barker-haliski, M. & White, H. S. *Use of Animal Models for Epilepsy Research and*
Therapy Development. Models of Seizures and Epilepsy (Elsevier Inc., 2014).
doi:10.1016/B978-0-12-804066-9/00007-9

136.   Lopes-aguiar, C., Ruggiero, R. N. & Umeoka, E. H. Animal models of epilepsy⬚: use and
limitations. 1693–1705 (2014). doi:10.2147/NDT.S50371

137.   Wykes, R. C. *et al.* Epilepsy: Optogenetic and potassium channel gene therapy in a rodent
model of focal neocortical epilepsy. *Sci. Transl. Med.* (2012).
doi:10.1126/scitranslmed.3004190

138.   Assaf, F. & Schiller, Y. The antiepileptic and ictogenic effects of optogenetic
neurostimulation of PV-expressing interneurons. *J. Neurophysiol.* (2016).
doi:10.1152/jn.00744.2015

139.   Berglind, F., Andersson, M. & Kokaia, M. Dynamic interaction of local and
transhemispheric networks is necessary for progressive intensification of hippocampal
seizures. *Sci. Rep.* (2018). doi:10.1038/s41598-018-23659-x

140.   Berglind, F. *et al.* Optogenetic inhibition of chemically induced hypersynchronized
bursting in mice. *Neurobiol. Dis.* (2014). doi:10.1016/j.nbd.2014.01.015

141.   Bui, A. D. *et al.* Dentate gyrus mossy cells control spontaneous convulsive seizures and
spatial memory. *Science (80-.).* (2018). doi:10.1126/science.aan4074

142.   Chen, S. *et al.* Near-infrared deep brain stimulation via upconversion nanoparticle–
mediated optogenetics. *Science (80-.).* (2018). doi:10.1126/science.aaq1144

143.   Chiang, C. C., Ladas, T. P., Gonzalez-Reyes, L. E. & Durand, D. M. Seizure suppression
by high frequency Optogenetic stimulation using in vitro and in vivo animal models of
epilepsy. *Brain Stimul.* (2014). doi:10.1016/j.brs.2014.07.034

144.   Krook-Magnuson, E., Armstrong, C., Oijala, M. & Soltesz, I. On-demand optogenetic
control of spontaneous seizures in temporal lobe epilepsy. *Nat. Commun.* (2013).
doi:10.1038/ncomms2376

145.   Ladas, T. P., Chiang, C. C., Gonzalez-Reyes, L. E., Nowak, T. & Durand, D. M. Seizure
reduction through interneuron-mediated entrainment using low frequency optical
stimulation. *Exp. Neurol.* (2015). doi:10.1016/j.expneurol.2015.04.001

146.   Lu, Y. *et al.* Optogenetic dissection of ictal propagation in the hippocampal-entorhinal
cortex structures. *Nat. Commun.* (2016). doi:10.1038/ncomms10962

147.   Wang, Y. *et al.* Depolarized GABAergic Signaling in Subicular Microcircuits Mediates

Generalized Seizure in Temporal Lobe Epilepsy. *Neuron* (2017).

doi:10.1016/j.neuron.2017.06.004

148.   Yekhlef, L., Breschi, G. L., Lagostena, L., Russo, G. & Taverna, S. Selective activation of

parvalbumin- or somatostatin-expressing interneurons triggers epileptic seizurelike

activity in mouse medial entorhinal cortex. *J. Neurophysiol.* (2015).

doi:10.1152/jn.00841.2014

149.   Kros, L. *et al. Cerebellar output controls generalized spike-and-wave discharge*

occurrence. Annals of neurology **77**, (2015).

150.   Soper, C., Wicker, E., Kulick, C. V., N'Gouemo, P. & Forcelli, P. A. Optogenetic

activation of superior colliculus neurons suppresses seizures originating in diverse brain

networks. *Neurobiol. Dis.* (2016). doi:10.1016/j.nbd.2015.12.012

151.   Tønnesen, J., Sørensen, A. T., Deisseroth, K., Lundberg, C. & Kokaia, M. Optogenetic

control of epileptiform activity. *Proc. Natl. Acad. Sci. U. S. A.* (2009).

doi:10.1073/pnas.0901915106

152.   Osawa, S. I. *et al.* Optogenetically Induced Seizure and the Longitudinal Hippocampal

Network Dynamics. *PLoS One* (2013). doi:10.1371/journal.pone.0060928

153.   Sessolo, M. *et al.* Parvalbumin-positive inhibitory interneurons oppose propagation but

favor generation of focal epileptiform activity. *J. Neurosci.* (2015).

doi:10.1523/JNEUROSCI.5117-14.2015

154.   Shiri, Z., Manseau, F., Lévesque, M., Williams, S. & Avoli, M. Interneuron activity leads

to initiation of low-voltage fast-onset seizures. *Ann. Neurol.* (2015).

doi:10.1002/ana.24342

155.   Weitz, A. J. *et al.* Optogenetic fMRI reveals distinct, frequency-dependent networks

recruited by dorsal and intermediate hippocampus stimulations. *Neuroimage* (2015).

doi:10.1016/j.neuroimage.2014.10.039

156.   Cela, E. *et al.* An Optogenetic Kindling Model of Neocortical Epilepsy. *Sci. Rep.* **9**,

(2019).

157.   Chang, M. *et al.* Brief activation of GABAergic interneurons initiates the transition to ictal

events through post-inhibitory rebound excitation. *Neurobiol. Dis.* (2018).

doi:10.1016/j.nbd.2017.10.007

158. Roodakker, K. R. *et al.* Ecstatic and gelastic seizures relate to the hypothalamus. *Epilepsy*
Behav. Reports **14**, 100358 (2020).

159. Maguire, J. & Salpekar, J. A. Stress, Seizures, and Hypothalamic-Pituitary-Adrenal Axis
Targets for the Treatment of Epilepsy. *Epilepsy Behav.* **26**, 352–362 (2013).

160. A, C. Epilepsy and the hippocampus. *Front. Neurol. Neurosci.* **34**, 121–142 (2014).

161. Avoli, M. The Epileptic Hippocampus Revisited: Back to the Future. *Epilepsy Curr.* **7**,
116 (2007).

162. Uva, L. *et al.* A Novel Focal Seizure Pattern Generated in Superficial Layers of the
Olfactory Cortex. *J. Neurosci.* **37**, 3544–3554 (2017).

163. U, E. & W, L. Strong olfactory stimulation reduces seizure susceptibility in amygdala-
kindled rats. *Neurosci. Lett.* **287**, 199–202 (2000).

164. W, L. & U, E. The role of the piriform cortex in kindling. *Prog. Neurobiol.* **50**, 427–481
(1996).

165. Ryu, B. *et al.* Chronic loss of inhibition in piriform cortex following brief, daily
optogenetic stimulation. *Cell Rep.* **35**, 109001 (2021).

166. Aroniadou-Anderjaska, V., Fritsch, B., Qashu, F. & Braga, M. F. M. Pathology and
Pathophysiology of the Amygdala in Epileptogenesis and Epilepsy. *Epilepsy Res.* **78**, 102
(2008).

167. Lv, R.-J. *et al.* Temporal lobe epilepsy with amygdala enlargement: a subtype of temporal
lobe epilepsy. *BMC Neurol. 2014 141* **14**, 1–8 (2014).

168. Paesschen, W. Van, King, M. D., Duncan, J. S. & Connelly, A. The Amygdala and
Temporal Lobe Simple Partial Seizures: A Prospective and Quantitative MRI Study.
Epilepsia **42**, 857–862 (2001).

169. C, D., V, R., B, L., C, M. & A, D. Dopamine in the striatum modulates seizures in a
genetic model of absence epilepsy in the rat. *Neuroscience* **100**, 335–344 (2000).

170. Miyamoto, H. *et al.* Impaired cortico-striatal excitatory transmission triggers epilepsy.
Nat. Commun. 2019 101 **10**, 1–13 (2019).

171. Vuong, J. & Devergnas, A. The Role of the Basal Ganglia in the Control of Seizure. *J.*
Neural Transm. **125**, 531 (2018).

172. Rijkers, K. *et al.* Sustained Reduction of Cerebellar Activity in Experimental Epilepsy.
Biomed Res. Int. (2015). doi:10.1155/2015/718591

173.   Tóth, Z. *et al.* Immunohistochemistry of cerebellar seizures: Mossy fiber afferents play an
1497   important role in seizure spread and initiation in the rat. *Acta Histochem.* (2015).
1498   doi:10.1016/j.acthis.2014.11.002

1499   174.   S, B., J, W., K, V. P., A, M. & FA, B. Cerebellar seizures. *J. Neurosurg. Pediatr.* **12**, 288–
1500   292 (2013).

1501   175.   Bohnen, N. I., O'Brien, T. J., Mullan, B. P. & So, E. L. Cerebellar Changes in Partial
1502   Seizures: Clinical Correlations of Quantitative SPECT and MRI Analysis. *Epilepsia* **39**,
1503   640–650 (1998).

1504   176.   Li, Y. *et al.* Cerebellum abnormalities in idiopathic generalized epilepsy with generalized
1505   tonic-clonic seizures revealed by diffusion tensor imaging. *PLoS One* **5**, (2010).

1506   177.   JOHNSON, H. C., BROWNE, K. M. & MARKHAM, J. W. EXPERIMENTAL
1507   CEREBELLAR SEIZURES. *A.M.A. Arch. Neurol. Psychiatry* **67**, 473–482 (1952).

1508   178.   Bomben, V. C. *et al.* Isolated P/Q calcium channel deletion in layer VI corticothalamic
1509   neurons generates absence epilepsy. *J. Neurosci.* (2016). doi:10.1523/JNEUROSCI.2555-
1510   15.2016

1511   179.   Sutula, T. P. & Kotloski, R. J. *Kindling: A Model and Phenomenon of Epilepsy. Models of*
1512   *Seizures and Epilepsy: Second Edition* (Elsevier Inc., 2017). doi:10.1016/B978-0-12-
1513   804066-9.00055-9

1514   180.   Ergul Erkec, O. Pentylenetetrazol Kindling Epilepsy Model. *J. Turkish Epilepsi Soc.*
1515   (2015). doi:10.5505/epilepsi.2015.08108

1516   181.   Dhir, A. Pentylenetetrazol (PTZ) kindling model of epilepsy. *Curr. Protoc. Neurosci.* **1**,
1517   (2012).

1518   182.   Marescaux, C., Vergnes, M. & Depaulis, A. Genetic absence epilepsy in rats from
1519   Strasbourg - A review. *Journal of Neural Transmission, Supplement* (1992).
1520   doi:10.1007/978-3-7091-9206-1_4

1521   183.   Phillips, J. W. *et al.* A repeated molecular architecture across thalamic pathways. *Nat.*
1522   *Neurosci.* (2019). doi:10.1038/s41593-019-0483-3

1523   184.   Phillips, J. W. *et al.* A single spectrum of neuronal identities across thalamus. *bioRxiv*
1524   (2018). doi:10.1101/241315

1525   185.   Lam, Y. W. & Sherman, S. M. Functional organization of the somatosensory cortical layer
1526   6 feedback to the thalamus. *Cereb. Cortex* (2010). doi:10.1093/cercor/bhp077

1527   186.   Keller, C. J. *et al.* Heterogeneous neuronal firing patterns during interictal epileptiform
1528   discharges in the human cortex. *Brain* (2010). doi:10.1093/brain/awq112

1529   187.   Wyler, A. R., Ojemann, G. A. & Ward, A. A. Neurons in human epileptic cortex:
1530   Correlation between unit and EEG activity. *Ann. Neurol.* (1982).
1531   doi:10.1002/ana.410110311

1532   188.   Schevon, C. A. *et al.* Evidence of an inhibitory restraint of seizure activity in humans. *Nat.*
1533   *Commun.* (2012). doi:10.1038/ncomms2056

1534   189.   Truccolo, W. *et al.* Single-neuron dynamics in human focal epilepsy. *Nat. Neurosci.*
1535   (2011). doi:10.1038/nn.2782

1536   190.   Babb, T. L., Wilson, C. L. & Isokawa-Akesson, M. Firing patterns of human limbic
1537   neurons during stereoencephalography (SEEG) and clinical temporal lobe seizures.
1538   *Electroencephalogr. Clin. Neurophysiol.* (1987). doi:10.1016/0013-4694(87)90093-9

1539   191.   Wyler, A. R. & Ward, A. A. Neuronal firing patterns from epileptogenic foci of monkey
1540   and human. *Adv. Neurol.* (1986).

1541   192.   Miri, M. L., Vinck, M., Pant, R. & Cardin, J. A. Altered hippocampal interneuron activity
1542   precedes ictal onset. *Elife* (2018). doi:10.7554/eLife.40750

1543   193.   Magloire, V., Mercier, M. S., Kullmann, D. M. & Pavlov, I. GABAergic Interneurons in
1544   Seizures: Investigating Causality With Optogenetics. *Neuroscientist* (2019).
1545   doi:10.1177/1073858418805002

1546   194.   Khoshkhoo, S., Vogt, D. & Sohal, V. S. Dynamic, Cell-Type-Specific Roles for
1547   GABAergic Interneurons in a Mouse Model of Optogenetically Inducible Seizures.
1548   *Neuron* (2017). doi:10.1016/j.neuron.2016.11.043

1549   195.   Cruikshank, S. J., Lewis, T. J. & Connors, B. W. Synaptic basis for intense
1550   thalamocortical activation of feedforward inhibitory cells in neocortex. *Nat. Neurosci.*
1551   (2007). doi:10.1038/nn1861

1552   196.   Zucca, S., Pasquale, V., Lagomarsino de Leon Roig, P., Panzeri, S. & Fellin, T. Thalamic
1553   Drive of Cortical Parvalbumin-Positive Interneurons during Down States in Anesthetized
1554   Mice. *Curr. Biol.* (2019). doi:10.1016/j.cub.2019.04.007

1555   197.   Sermet, B. S. *et al.* Pathway-, layer-and cell-type-specific thalamic input to mouse barrel
1556   cortex. *Elife* (2019). doi:10.7554/eLife.52665

1557   198.   Hafner, G. *et al.* Mapping Brain-Wide Afferent Inputs of Parvalbumin-Expressing

GABAergic Neurons in Barrel Cortex Reveals Local and Long-Range Circuit Motifs. *Cell Rep.* (2019). doi:10.1016/j.celrep.2019.08.064

199. Clemente-Perez, A. *et al.* Distinct Thalamic Reticular Cell Types Differentially Modulate Normal and Pathological Cortical Rhythms. *Cell Rep.* (2017). doi:10.1016/j.celrep.2017.05.044

200. Chen, J. *et al.* Regulation of delta FosB and FosB-like proteins by electroconvulsive seizure and cocaine treatments. *Mol. Pharmacol.* **48**, (1995).

201. Madsen, T. M., Bolwig, T. G. & Mikkelsen, J. D. Differential regulation of c-fos and fosB in the rat brain after amygdala kindling. *Cell. Mol. Neurobiol.* (2006). doi:10.1007/s10571-006-9202-1

202. Yutsudo, N. *et al.* FosB-null mice display impaired adult hippocampal neurogenesis and spontaneous epilepsy with depressive behavior. *Neuropsychopharmacology* (2013). doi:10.1038/npp.2012.260

203. Corbett, B. F. *et al.* ΔFosB Regulates Gene Expression and Cognitive Dysfunction in a Mouse Model of Alzheimer's Disease. *Cell Rep.* (2017). doi:10.1016/j.celrep.2017.06.040

204. Giordano, C., Vinet, J., Curia, G. & Biagini, G. Repeated 6-Hz corneal stimulation progressively increases FosB/δFosB levels in the lateral amygdala and induces seizure generalization to the Hippocampus. *PLoS One* (2015). doi:10.1371/journal.pone.0141221

205. Mesiwala, A. H. *et al.* Focal motor seizures with secondary generalization arising in the cerebellum. Case report and review of the literature. *J Neurosurg* (2002). doi:10.3171/jns.2002.97.1.0190

206. Crooks, R., Mitchell, T. & Thom, M. Patterns of cerebellar atrophy in patients with chronic epilepsy: A quantitative neuropathological study. *Epilepsy Res.* (2000). doi:10.1016/S0920-1211(00)00133-9

207. Afifi, A. K. & Van Allen, M. W. Cerebellar atrophy in epilepsy. Pneumographic and histological documentation of a case with psychosis. *J. Neurol. Neurosurg. Psychiatry* (1968). doi:10.1136/jnnp.31.2.169

208. Marcián, V., Filip, P., Bareš, M. & Brázdil, M. Cerebellar Dysfunction and Ataxia in Patients with Epilepsy: Coincidence, Consequence, or Cause? *Tremor Other Hyperkinet. Mov. (N. Y).* **6**, 376 (2016).

209. van den Ende, T., Sharifi, S., van der Salm, S. M. A. & van Rootselaar, A.-F. Familial

Cortical Myoclonic Tremor and Epilepsy, an Enigmatic Disorder: From Phenotypes to

Pathophysiology and Genetics. A Systematic Review. *Tremor Other Hyperkinet Mov (N*

Y) (2018). doi:10.7916/D85155WJ

210. Campbell, D. B. & Hess, E. J. Cerebellar circuitry is activated during convulsive episodes

in the tottering (tg/tg) mutant mouse. *Neuroscience* **85**, 773–783 (1998).

211. Krauss, G. L. & Koubeissi, M. Z. Cerebellar and thalamic stimulation treatment for

epilepsy. *Acta Neurochir. Suppl.* **97**, 347–356 (2007).

212. Kros, L. *et al.* Corrigendum: Synchronicity and Rhythmicity of Purkinje Cell Firing

during Generalized Spike-and-Wave Discharges in a Natural Mouse Model of Absence

Epilepsy. *Front. Cell. Neurosci.* **11**, 1–18 (2017).

213. Kros, L., Eelkman Rooda, O. H. J., De Zeeuw, C. I. & Hoebeek, F. E. Controlling

Cerebellar Output to Treat Refractory Epilepsy. *Trends in Neurosciences* **38**, 789–799

(2015).

214. Harvey, A. S. *et al.* Hemifacial seizures and cerebellar ganglioglioma: An epilepsy

syndrome of infancy with seizures of cerebellar origin. *Ann. Neurol.* (1996).

doi:10.1002/ana.410400115

215. Işik, U., Dinçer, A., Sav, A. & Özek, M. M. Seizures as the sole manifestation of

cerebellar lesions in three children. *J. Pediatr. Neurol.* (2009). doi:10.3233/JPN-2009-

0286

216. Ming, X., Prasad, N., Thulasi, V., Elkins, K. & Shivamurthy, V. K. N. Possible

contribution of cerebellar disinhibition in epilepsy. *Epilepsy and Behavior* (2021).

doi:10.1016/j.yebeh.2021.107944

217. Krisztin-Péva, B., Mihály, A. & Tóth, Z. Differential expression of the c-fos protein and

synaptophysin in zebrin ii positive and zebrin ii negative cerebellar cortical areas in 4-

aminopyridine seizures. *Acta Neurobiol. Exp. (Wars).* (2019). doi:10.21307/ane-2019-022

218. Kros, L. *et al.* Synchronicity and rhythmicity of purkinje cell firing during generalized

spike-and-wave discharges in a natural mouse model of absence epilepsy. *Front. Cell.*

Neurosci. (2017). doi:10.3389/fncel.2017.00346

219. Alva, P. *et al.* A potential role for the cerebellar nuclei in absence seizures. *BMC*

Neurosci. (2013). doi:10.1186/1471-2202-14-s1-p170

1619

Figure 1

Ntsr1^Cre;ROSA^lox-stop-lox-Sun1

Ntsr1^Cre;ROSA^lox-stop-lox-TdTomato

cortex

thalamus

cerebellar nuclei

cortex

thalamus

cerebellar nuclei

Figure 2

Figure 3

a

LD
CL
PO
RT
VPM
VPL
VAL
VM

b

Corrected Fluoresence (a.u.)

n.s.
****
*
****
n.s.
****
****
****
****
****

Thalamic Region: VPM, PO, LD, LP, CL, VM, VPL, RT, VAL, LGd

c

VPM

500 um

d

VPM

VPM

● non-seizure inducing target
● seizure inducing target

e

Optogenetic seizure

1s   3s   5s   10s

Kainic Acid seizure

1s   4s   7s   23s

f

Bregma
Lambda
Ground
Reference

Somatosensory cortex
Motor cortex, left
Motor cortex, right
Ground
Reference
Cerebellum

g

Optogenetic seizure

Kainic Acid seizure

Cortex EEG
Cerebellum EEG
Behavioral seizure

Figure 4

Figure 5

Figure 6

Figure 7

a Cell Count H&E

b Cell Activity dFosB

Control
Seizure

Contralateral thalamus | Ipsilateral thalamus

c dFosB DAB stain

d

e filtered dFosB

f

Figure 8

a

retro-rAAV-Ef1a
-mCherry-WPRE

cortex

inferior
colliculus

striatum

VPM

zona incerta

cerebellum

brainstem nuclei

b

VPM

injection site

c

L4

L5

L6

White matter

d

SIM

Crus1

LIV,V

Crus2

L III

PM

Interposed Nucleus

L VIII

e

Interposed Nucleus

Figure 9

Ntsr1^{Cre};ROSA$^{lox-stop-lox-Sun1}$ *Ntsr1^{Cre};ROSA$^{lox-stop-lox-TdTomato}$*

ai Layer 5/6 cortex

Caudoputamen

aii Layer 5/6 cortex

Thalamus

aiii Layer 5/6 cortex

Hippocampus

aiv Layer 5/6 cortex

Superior Colliculus

Substantia Nigra
VTA

Endopiriform
Nucleus

av Cerebellar
nuclei

Brainstem nuclei

bi

bii

biii

biv

bv

bvi

c

Supplementary Figure 2

Supplementary Figure 4

Increased EEG amplitude is consistent between brain regions across mice

KEY
Left motor cortex EEG
Right motor cortex EEG
Left somatosensory cortex EEG
Midline cerebellum EEG
30s 30Hz light stimulation

Supplementary Figure 6

Example Mouse 1 **Example Mouse 2** **Example Mouse 3**

a — Left Motor Cortex v. Left Somatosensory Cortex

b — Left Motor Cortex v. Right Motor Cortex

c — Left Motor Cortex v. Cerebellum

d — Right Motor Cortex v. Cerebellum

e — RIght Motor Cortex v. Left Somatosensory

Coherence (n.u.)

Frequency (Hz)

BEFORE STIM
DURING SEIZURE

Supplementary Figure 7