Emergence of rare carbapenemases (FRI, GES-5, IMI, SFC, and SFH-1) in Enterobacterales isolated from surface waters in Japan

Ryota Gomi[a,*], Yasufumi Matsumura[b], Michio Tanaka[b], Masaru Ihara[c], Yoshinori Sugie[c], Tomonari Matsuda[c], and Masaki Yamamoto[b]

[a]Department of Environmental Engineering, Graduate School of Engineering, Kyoto University, Katsura, Nishikyo-ku, 615-8540, Kyoto, Japan

[b]Department of Clinical Laboratory Medicine, Kyoto University Graduate School of Medicine, 54 Shogoin-kawahara-cho, Sakyo-ku, 606-8507, Kyoto, Japan

[c]Research Center for Environmental Quality Management, Kyoto University, 1-2 Yumihama, Otsu, 520-0811, Shiga, Japan

[*]Corresponding author

E-mail address: gomi.ryota.34v@kyoto-u.jp; Tel: +81-75-383-3354; Fax: +81-75-383-3358

ABSTRACT

Objectives: Carbapenemase-producing Enterobacterales (CPE) pose serious threats to public health. Compared with clinical CPE, the genetic characteristics of environmental CPE are not well understood. This study aimed to characterize the genetic determinants of carbapenem resistance in CPE isolated from environmental waters in Japan.

Methods: Eighty-five water samples were collected from rivers and a lake in Japan. CPE were identified using selective media, and genome sequencing was performed for the obtained isolates (n = 21).

Results: Various rare/novel carbapenemases were identified: GES-5 in *Raoultella planticola* (n = 1), FRI-8 and FRI-11 in *Enterobacter* spp. (n = 8), IMI-22 and IMI-23 in *Serratia ureilytica* (n = 3), and SFC-1, SFC-2 and SFH-1 in *Serratia fonticola* (n = 9). Genomes of 11 isolates could be closed, allowing the elucidation of the genetic contexts of the carbapenemase genes. The bla_{GES-5} gene was located within a class 1 integron, In2071 (cassette array, bla_{GES-5}-*aacA3-aadA16*), on a 33 kb IncP6 plasmid. The bla_{FRI-8} genes were carried on IncFII(Yp) plasmids ranging in size from 191 kb to 244 kb, and the bla_{FRI-11} genes were carried on 70 kb and 74 kb IncFII(pECLA)/IncR plasmids. The bla_{IMI-22} and bla_{IMI-23} genes were colocated on a 107 kb plasmid. The bla_{SFC} and bla_{SFH-1} genes were found on putative genomic islands inserted at tRNA-Phe genes in chromosomes.

Conclusions: This study revealed the presence of rare/novel carbapenemases among CPE in

aquatic environments, suggesting that the environment may act as a potential reservoir of these

minor carbapenemases.

INTRODUCTION

The worldwide spread of carbapenemase-producing Enterobacterales (CPE) is a major threat

to public health. CPE can be resistant to carbapenems and many other classes of commonly

used drugs, which significantly limits treatment options.[1, 2] The most clinically relevant

carbapenemases detected in CPE are the 'big five' carbapenemases (Ambler class A KPC

enzymes, class B enzymes of IMP, VIM, and NDM, and class D OXA-48-like enzymes), while

Enterobacterales producing other 'minor' carbapenemases have also been sporadically

reported.[3]

Historically, antibiotic resistance has been considered to be a clinical problem, but a growing

number of studies have suggested that the environment represents an important reservoir for

antibiotic resistant bacteria and antibiotic resistance genes.[4-6] Previous studies have reported

the occurrence of CPE in environmental waters, with the aforementioned 'big five'

carbapenemases being the predominant types.[7, 8] Minor carbapenemases, such as GES and IMI,

were also detected among environmental CPE,[9-11] raising the possibility that environmental

waters may act as reservoirs for these minor carbapenemases.

Previous studies have shown that Japanese clinical CPE usually carry bla_{IMP} and sporadically

carry other carbapenemase genes such as bla_{KPC}, bla_{NDM}, and bla_{OXA-48}.[12, 13] However, the genetic

characteristics of CPE in the natural environment, in particular in surface waters, are not well

understood in Japan.[14] Here, we isolated CPE from river and lake water in Japan and

characterized the obtained CPE isolates in terms of resistance determinants, genetic contexts

of the carbapenemase genes, and phylogeny.

MATERIALS AND METHODS

Sample collection and bacterial isolation

In total, 72 water samples were collected from 12 sites in Lake Biwa and 13 water samples

were collected from 8 sites in its surrounding rivers between February and November 2020

(**Figure 1**). Samples were processed using the membrane filter method with chromID CARBA

agar (bioMérieux, Marcy-l'Étoile, France). For each sample, up to three colonies showing

Escherichia coli profiles (pink to burgundy) or KESC (*Klebsiella*, *Enterobacter*, *Serratia*,

Citrobacter) profiles (bluish green to bluish grey) with different colony morphologies were

isolated. The isolates were restreaked on fresh chromID CARBA agar and incubated again.

Streaking on chromID CARBA agar and incubation were repeated until pure colonies were

obtained. The oxidase test was performed using a cytochrome oxidase test strip (Nissui, Tokyo,

Japan), and isolates with negative test results were stored at -85°C in 35% glycerol.

Figure 1. Map of sampling sites. Sampling sites in Lake Biwa are indicated with L1 to L12,

and sampling sites in rivers are indicated with R1 to R8. R8 is in a small river and located near

L11. Red filled circles indicate sampling points where CPE isolates were detected.

Antibiotic susceptibility testing

Antibiotic susceptibility was assessed by microdilution using a Dry Plate Eiken (Eiken,

Tokyo, Japan). The results were interpreted using CLSI criteria.[15] The Carba NP test was

performed to test the carbapenemase activity.[16] *Klebsiella pneumoniae* ATCC BAA-1705 was

used for the positive control, and *K. pneumoniae* ATCC BAA-1706 was used for the negative

control.

Cloning experiments

Novel carbapenemase genes (bla_{FRI-11} in JBIWA008, bla_{SFC-2} in JSRIV001, bla_{IMI-22} and bla_{IMI-}

$_{23}$ in JBIWA004) were amplified using the primers listed in **Table S1**, cloned into the pCR2.1-

TOPO vector using the TOPO TA cloning kit (Thermo Fisher Scientific, Waltham, MA, USA),

and transformed into *E. coli* DH5α (TaKaRa Bio, Inc., Shiga, Japan). Antibiotic susceptibilities

of the recipient and transformants were assessed by microdilution as described above.

Genome sequencing and assembly

DNA was extracted from each isolate using a DNeasy Blood and Tissue kit (Qiagen, Hilden,

Germany) for DNBSEQ short-read sequencing. Libraries were prepared with the MGIEasy FS

PCR-free DNA library prep set (MGI Tech, Shenzhen, China), and paired-end sequencing

(2×150-bp) was performed on DNBSEQ-G400RS. Reads were subsampled using seqtk (v1.3,

https://github.com/lh3/seqtk), trimmed using fastp (v0.20.0),[17] and assembled using Unicycler

(v0.4.8) with the --no_correct option.[18]

Selected isolates (n=12) were also subjected to Oxford Nanopore Technologies (ONT) long-

read sequencing. DNA was isolated as described previously,[19] and libraries were prepared using

the SQK-LSK109 kit (Oxford Nanopore Technologies, Oxford, UK) and the Short Read

Eliminator XS kit (Circulomics, Inc., Baltimore, MD, USA). Sequencing was performed using

ONT GridION X5 with a FLO-MIN106 flow cell. ONT reads were filtered using Filtlong

(v0.2.0, https://github.com/rrwick/Filtlong) to remove reads with low quality or short length.

De novo hybrid assemblies using trimmed DNBSEQ reads and filtered ONT reads were

performed using Unicycler (v0.4.8)[18] with the --no_correct option. Genomes that could not be

completed by Unicycler were subjected to long-read assembly by Flye (v2.8)[20] with the --

plasmids option followed by long-read polishing using Medaka (v1.2.6,

https://github.com/nanoporetech/medaka) and short-read polishing using Pilon (v1.23).[21]

Genomic analysis

Species identification was performed by comparing the obtained genome sequences to the

genome sequences of the type strains. The average nucleotide identity (ANI) analysis was

performed using OrthoANIu,[22] and digital DNA-DNA hybridization (dDDH) analysis was

performed using the GGDC website (formula 2).[23] A ≥96% ANI value or ≥ a 70% DDH value

was used for defining the species.[23, 24] Sequence types (STs) were assigned to *Enterobacter*

isolates using the multilocus sequence typing (MLST) scheme established previously.[25] Whole-

genome SNP-based trees were built using kSNP3.[26] The program Kchooser was used to select

the optimum k-mer length, and SNP loci occurring in at least 80% of the strains were used for

building trees.

Plasmid replicons, relaxase types, mate-pair formation types, and transferability of plasmids

were analysed using PlasmidFinder 2.1 and MOB-typer.[27, 28] Annotation was performed using

ISfinder,[29]   the   RAST   server,[30]   DFAST,[31]   ResFinder   4.1,[32]   Abricate   (v1.0.1,

https://github.com/tseemann/abricate),   INTEGRALL,[33]   and   the   BLAST   algorithm

(https://blast.ncbi.nlm.nih.gov/Blast.cgi).   Genomic   islands   (GIs)   were   detected   using

IslandViewer 4.[34] New allele numbers for carbapenemase genes were assigned by NCBI

through submission of sequence data according to instructions on the following website:

https://www.ncbi.nlm.nih.gov/pathogens/submit-beta-lactamase/.

Accession numbers

The complete genomes and sequence read files were deposited in GenBank and the NCBI

SRA under BioProject PRJNA725976. The BioSample accession number for each isolate is

also available in **Table 1**.

Table 1. Characteristics of 21 CPE isolates recovered from surface waters in the present study.

Isolate[a]	Sampling site	Collection date	Species[b]	Carbapenemase gene(s)	Location of carbapenemase gene(s)[c]	BioSample accession number
JBIWA001	L12	2/5/2020	*Raoultella planticola*	*bla*GES-5	pJBIWA001_5 (IncP6, 32,930 bp)	SAMN18916998
JBIWA002	L12	2/5/2020	*Enterobacter asburiae* (ST1585)	*bla*FRI-8	pJBIWA002_1 (IncFII(Yp), 243,962 bp)	SAMN18916999
JBIWA003	L12	3/4/2020	*Enterobacter quasiroggenkampii* (ST929)	*bla*FRI-8	pJBIWA003_1 (IncFII(Yp), 221,552 bp)	SAMN18917000
JBIWA004	L12	3/4/2020	*Serratia ureilytica*	*bla*IMI-22, *bla*IMI-23	pJBIWA004_1 (IncFII(pRSB107), pSM22, 106,702 bp)	SAMN18917001
JBIWA005	L12	3/4/2020	*Enterobacter genomospecies 8* (ST930)	*bla*FRI-8	pJBIWA005_1 (IncFII(Yp), 191,037 bp)	SAMN18917002

JBIWA006	L9	9/2/2020	*Enterobacter asburiae* (ST1586)	*bla*FRI-11	UD	SAMN19 031978
JBIWA007	L2	9/2/2020	*Enterobacter asburiae* (ST1586)	*bla*FRI-11	pJBIWA007_1 (IncFII(pECLA), IncR, 69,715 bp)	SAMN18 917003
JBIWA008	L1	9/2/2020	*Enterobacter* sp. (ST1587)	*bla*FRI-11	pJBIWA008_3 (IncFII(pECLA), IncR, 73,531 bp)	SAMN18 917004
JBIWA009	L1	9/2/2020	*Enterobacter asburiae* (ST1586)	*bla*FRI-11	UD	SAMN19 031979
JBIWA010	L11	9/2/2020	*Enterobacter* sp. (ST1587)	*bla*FRI-11	UD	SAMN19 031980
JBIWA011	L2	10/5/2020	*Serratia ureilytica*	*bla*IMI-22, *bla*IMI-23	UD	SAMN19 031981
JBIWA012	L1	10/5/2020	*Serratia ureilytica*	*bla*IMI-22, *bla*IMI-23	UD	SAMN19 031982

JSRIV001	R8	3/4/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	chromosome	SAMN18 917005
JSRIV002	R4	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	chromosome	SAMN18 917006
JSRIV003	R2	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	UD	SAMN19 031973
JSRIV004	R2	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	chromosome	SAMN18 917007
JSRIV005	R1	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	UD	SAMN19 031974
JSRIV006	R1	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-1}, bla_{SFH-1}	chromosome	SAMN18 917008
JSRIV007	R3	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	UD	SAMN19 031975
JSRIV008	R5	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	UD	SAMN19 031976
JSRIV009	R5	3/17/2020	*Serratia fonticola* (clade III)	bla_{SFC-2}, bla_{SFH-1}	UD	SAMN19 031977

[a]Genomes of isolates in bold were sequenced by both DNBSEQ and ONT sequencers. Among

the 12 isolates in bold, all the genomes except JBIWA006 were completed. Also see **Table S2**

for the accession numbers of the closed replicons and **Table S3** for the SRA accession numbers

of the incomplete genomes.

[b]STs/clades are shown in parentheses.

[c]UD: undetermined.

RESULTS AND DISCUSSION

Detection of CPE in water samples

Twelve isolates and nine isolates were obtained from lake water samples and river water

samples, respectively (**Table 1**). Lake isolates were identified as *Enterobacter* spp. (n = 8),

Serratia ureilytica (n = 3), and *Raoultella planticola* (n = 1), and all nine river isolates were

identified as *Serratia fonticola* clade III (discussed in detail below) by ANI and dDDH analyses

(**Table 1**).

Among the 21 isolates, 12 were subjected to both DNBSEQ short-read sequencing and ONT

long-read sequencing. Nine genomes were completed by Unicycler, and two of the remaining

three genomes were completed by Flye. The genome of JBIWA006 could not be completed by

either approach. The completed genomes contained one to 12 plasmids (see **Table S2** for

details). The draft genome assemblies of the remaining isolates had 28 to 217 contigs and

189,689 bp to 2,472,998 bp N50 values after discarding contigs shorter than 100 bp (see **Table**

S3 for details).

All the carbapenemase genes detected in this study were found to encode minor

carbapenemases (**Table 1**). Briefly, bla_{GES-5} was detected in a *R. planticola* isolate, bla_{FRI-8} and

bla_{FRI-11} were detected in *Enterobacter* isolates, bla_{IMI-22} and bla_{IMI-23} were detected in *S.*

ureilytica isolates, and bla_{SFC-1}, bla_{SFC-2} and bla_{SFH-1} were detected in *S. fonticola* clade III

isolates. bla_{FRI-11}, bla_{IMI-22}, bla_{IMI-23}, and bla_{SFC-2} are the genes encoding novel carbapenemase

variants, and their protein accession numbers are as follows: FRI-11 (WP_223174987.1), IMI-

22 (WP_223174988.1), IMI-23 (WP_223525081.1), and SFC-2 (WP_223174986.1).

Phenotypic resistance was determined by microdilution. All the isolates were nonsusceptible

to doripenem and meropenem, and all but GES-5-producing *R. planticola* (JBIWA001) were

nonsusceptible to imipenem (**Table S4**). The Carba NP test confirmed carbapenemase activity

in all isolates carrying bla_{SFC} and bla_{SFH-1}, whereas isolates carrying bla_{GES-5}, bla_{FRI-8}, and bla_{IMI}

were negative by the test. Results of the Carba NP test for the bla_{FRI-11}-carrying isolates were

inconsistent (two were positive, one was negative, and two were indeterminate). Previous

studies reported that Enterobacterales harbouring carbapenemases such as bla_{GES-5} and bla_{IMI}

can be falsely negative in the Carba NP test.[35-37] Novel carbapenemase genes were cloned into

the pCR2.1-TOPO vector and transformed into *E. coli* DH5α. An increase in MICs of

carbapenems was detected in *E. coli* DH5α carrying the recombinant plasmids pCR2.1-TOPO-

bla$_{SFC-2}$ and pCR2.1-TOPO-*bla*$_{IMI-23}$ (**Table S5**). However, no detectable increase in MICs of

carbapenems was observed for *E. coli* DH5α carrying the recombinant plasmid pCR2.1-TOPO-

bla$_{FRI-11}$ or pCR2.1-TOPO-*bla*$_{IMI-22}$ under the conditions in the present study, which indicates

that FRI-11 and IMI-22 are enzymes with low carbapenemase activity.

Genomic analysis of *Raoultella planticola* isolate

River isolate JBIWA001 showed 99.15% ANI and 93.90% dDDH values with the type strain

of *R. planticola*. The assembly resulted in a chromosome (5,626,881 bp) and 12 plasmids

ranging from 1,443 bp to 136,413 bp (see **Table S2** for details). The *bla*$_{GES-5}$ gene was on a

32,930 bp plasmid, pJBIWA001_5, with an IncP6 replicon and a MOBP-type relaxase. *bla*$_{GES-}$

$_5$ was found to be part of a novel class 1 integron, In2071. In2071 contains a *bla*$_{GES-5}$-*aacA3*-

aadA16 cassette array, followed by the 3′-conserved segment (3′-CS) (*qacEΔ1*, *sul1*, orf5, orf6),

IS*1326*, *tniBΔ2*, *tniA*, and IRt (inverted repeat at *tni* end). A BLASTN search against the

GenBank database using pJBIWA001_5 as a query sequence identified some closely related

plasmids. pN260-3, which was carried by an *E. roggenkampii* strain detected in a patient with

cholecystitis in Japan in 2019,[38] showed the highest query coverage (97%) and also showed the

high percent identity (>96%) (see **Figure S1A** for the comparison of these two plasmids).

pN260-3 also carried *bla*$_{GES-5}$ but within a different class 1 integron structure (*intI1-bla*$_{GES-5}$-

aac(6′)-31-intI1Δ-qacG2-aadA5-qacEΔ1-sul1) (**Figure S1B**).

Genomic analysis of *Enterobacter* isolates

The *Enterobacter* genus comprises 22 species and 22 additional genomospecies.[39] The lake *Enterobacter* isolates in the present study were identified as *E. asburiae* (n = 4), *E. quasiroggenkampii* (n = 1), and *Enterobacter* genomospecies 8 (n = 1) based on the ANI and dDDH analyses (also see **Figure S2** for the *Enterobacter* phylogenetic tree). The remaining two isolates, JBIWA008 and JBIWA010, showed < 96% ANI values and < 70% dDDH values against all type strains and reference strains (closest to *E. asburiae*, 95.2% ANI and 63.0% dDDH). It should be noted that although we adopted a 96% ANI cutoff value, which is the cutoff value used in a recent study defining *Enterobacter* species,[40] a 95% ANI cutoff value is also used for defining the species and 95-96% ANI values can be an inconclusive zone.[40, 41] However, dDDH values were lower than the 70% cutoff and thus these two isolates were considered to belong to a novel genomospecies. MLST assigned two novel STs (ST1585 and ST1586) to *E. asbriae* isolates, ST929 to an *E. quasiroggenkampii* isolate, ST930 to an *Enterobacter* genomospecies 8 isolate, and a novel ST (ST1587) to two isolates belonging to the novel genomospecies. Interestingly, *Enterobacter* ST929 and ST930 were first reported in our previous study analysing wastewater CPE isolates in the same region.[42] The location where ST929 and ST930 isolates were recovered in the present study is near the outlets of wastewater treatment plants (WWTPs), indicating the origin of these isolates could be those WWTPs.

We could determine the complete sequences of three bla_{FRI-8}-carrying plasmids, namely

pJBIWA002_1 (243,962 bp) from *E. asbriae* isolate JBIWA002, pJBIWA003_1 (221,552 bp)

from *E. quasiroggenkampii* isolate JBIWA003, and pJBIWA005_1 (191,037 bp) from

Enterobacter genomospecies 8 isolate JBIWA005. A BLASTN search against the GenBank

database using bla_{FRI-8} as a query sequence identified one bla_{FRI-8}-carrying plasmid,

Enterobacter sp. 18A13 plasmid pECC18A13-1 (150,509 bp), in which bla_{FRI-8} was first

detected.[43] The BLASTN search also identified the bla_{FLC-1}-carrying plasmid p3442-FLC-1

(93,658 bp) from the *E. cloacae* complex strain E3442.[44] FLC-1 (FRI-like carbapenemase-1)

shares 99.66% amino acid identity with FRI-8 and was suggested that it be reclassified as an

FRI variant.[3] All five plasmids (pJBIWA002_1, pJBIWA003_1, pJBIWA005_1, pECC18A13-

1, p3442-FLC-1) carried an IncFII(Yp)-like replicon and were typed as Y10:A-:B- by plasmid

MLST. All five plasmids also carried a MOBP-type relaxase, though pECC18A13-1 also

carried a MOBF-type relaxase. Although the five plasmids carried some sequences in common,

insertions/deletions contributed to variable region diversification (**Figure 2A**). Analysis of

genetic contexts of bla_{FRI-8} and bla_{FLC-1} revealed the presence of an IS*Peat2*-like insertion

sequence (IS) downstream of bla_{FRI-8}/bla_{FLC-1} and a LysR-type transcriptional regulator

upstream of bla_{FRI-8}/bla_{FLC-1} (**Figure 2B**). A truncated IS*Eae1*-like element disrupted the

IS*Peat2*-like IS in pJBIWA003_1, changing the downstream structure in this plasmid. RAST

identified a coding sequence encoding a hypothetical protein between bla_{FRI-8}/bla_{FLC-1} and a

LysR-type transcriptional regulator. BLASTN analysis showed this coding sequence to be

absent near the other bla_{FRI} genes.

We were able to determine the complete sequences of two bla_{FRI-11}-carrying plasmids, namely

pJBIWA007_1 (69,715 bp) from *E. asburiae* isolate JBIWA007 and pJBIWA008_3 (73,531

bp) from *Enterobacter* sp. isolate JBIWA008. Online BLASTN analysis identified a 108,672

bp plasmid, pJF-587, from *E. asburiae*,[45] which showed >90% coverage to pJBIWA007_1 and

pJBIWA008_3. pJF-587 carried bla_{FRI-2}, which shares 98.5% nucleotide identity with bla_{FRI-11}.

All three plasmids carried IncFII(pECLA) and IncR replicons. Although pJF-587 contained a

MOBF-type relaxase and was predicted to be conjugative by MOB-typer, pJBIWA007_1 and

pJBIWA008_3 did not contain a relaxase and were predicted to be non-mobilizable. Alignment

of the three plasmids revealed conjugation genes present on pJF-587 were deleted in

pJBIWA007_1 and pJBIWA008_3 (**Figure 3A**). The genetic contexts of bla_{FRI-11}/bla_{FRI-2} were

slightly different among the three plasmids. A DNA segment containing *ltrA* was present in

pJBIWA008_3, while this region was absent in pBIWA007_1 (**Figure 3B**). In pJF-587, this

segment was inverted compared to pJBIWA008_3, and an IS*Sm3*-like element was absent.

Although complete plasmid sequences were not available for JBIWA006, JBIWA009, and

JBIWA010, we could determine the genetic contexts of bla_{FRI-11} in these isolates using short-

read assemblies: a pBIWA007_1-type genetic context was detected in JBIWA006 and

JBIWA009, while a pJBIWA008_3-type genetic context was detected in JBIWA010.

Figure 2. (A) Alignment of plasmids pJBIWA003_1, pJBIWA002_1, pJBIWA005_1,

pECC18A13-1, and p3442-FLC-1. Positions of bla_{FRI-8} and bla_{FLC-1} are indicated by blue and

red triangles, respectively. (B) Genetic contexts of bla_{FRI-8} and bla_{FLC-1}. Blue arrows indicate

bla_{FRI-8}, a red arrow indicates bla_{FLC-1}, green boxes indicate insertion sequences, and orange

arrows indicate other coding sequences. hp: hypothetical protein. The alignments were

generated with Easyfig.[46]

Figure 3. (A) Alignment of pJBIWA007_1, pJBIWA008_3, and pJF-587. Positions of $bla_{\text{FRI-11}}$ and $bla_{\text{FRI-2}}$ are indicated by blue and red triangles, respectively. (B) Genetic contexts of $bla_{\text{FRI-11}}$ and $bla_{\text{FRI-2}}$. Blue arrows indicate $bla_{\text{FRI-11}}$, a red arrow indicates $bla_{\text{FRI-2}}$, green boxes indicate insertion sequences, and orange arrows indicate other coding sequences.

Genomic analysis of *Serratia* isolates

Although three *Serratia* isolates recovered from the lake could be classified as *S. ureilytica*, nine *Serratia* isolates from rivers could not be classified based on the ANI and dDDH analyses. The pairwise ANI values and dDDH values among the nine river isolates were all above 96%

and 70%, respectively, indicating the river isolates belong to a single species. The river isolates

seem to be related to *S. fonticola* (**Figure S3**), but the ANI and dDDH values between the river

isolates and the type strain of *S. fonticola* ranged from 95.5% to 95.7% and 65.1% to 65.4%,

respectively, which are lower than the 96% ANI and 70% dDDH cutoff values. These results

suggest the 9 river isolates could belong to a novel species whose closest relative is *S. fonticola*.

It should be noted that genome sequences for type strains were not available for some *Serratia*

species (*Serratia aquatilis*, *Serratia bozhouensis*, *Serratia entomophila*, *Serratia myotis*, and

Serratia profundus), and ANI and dDDH analyses could not be performed for these species.

However, 16S rRNA gene sequences were available for them and all showed < 98.7% identity

with the 16S rRNA gene sequences of the river isolates, which implies they belong to different

species.[41]

In order to gain further insight into the phylogenetic characteristics of river *Serratia* isolates,

we downloaded RefSeq assemblies labelled *S. fonticola* and built a phylogenetic tree (**Figure**

4, also see **Figure S4** for phylogenetic characteristics of these strains in relation to other

Serratia species). Interestingly, three clades were identified in the phylogenetic tree, namely

clade I (containing the genomes of type strains of *S. fonticola*), clade II, and clade III

(containing the genomes of river isolates). Mean within-clade ANI values were > 98% for all

clades, and within-clade pairwise ANI values always exceeded 96%. On the other hand, mean

between-clade ANI values were < 96%, and between-clade pairwise ANI values were always

< 96% (**Table S6**). As there are limitations in the use of the website-based GGDC dDDH

calculation tool to compare large numbers of genomes, we performed dDDH analysis only for

comparisons including the type strain from clade I or the reference strains from clade II and

clade III (**Table S7**). The results show that within-clade pairwise dDDH values were always

above 70% while between-clade pairwise dDDH values were always below 70%. These results

suggest clades I-III could represent different species, although further studies are needed to

confirm this.

River *Serratia* isolates carried two types of carbapenemase genes, bla_{SFH} and bla_{SFC}. bla_{SFH-1}

(a subclass B2 carbapenemase gene) and bla_{SFC-1} (a class A carbapenemase gene) have been

described only once in *S. fonticola* clade II strain UTAD54 (GCF_000476315.1) from

Portugal.[3, 47-49] Although no variants were reported previously for either of these two

carbapenemases, online BLASTN analysis revealed that *S. fonticola* clade III strain AU-AP2C

(GCF_000477615.1)[50] carried a variant of bla_{SFC} (the product shows 99.4% amino acid identity

to SFC-1). All river isolates except JSRIV006 carried bla_{SFH-1} and a variant of bla_{SFC} that is

different from bla_{SFC-1} and bla_{SFC} of AU-AP2C (the product shows 99.4% amino acid identities

to SFC-1 and SFC of AU-AP2C). Novel alleles SFC-2 and SFC-3 were assigned to SFC

enzymes in river isolates and AU-AP2C, respectively. JSRIV006, in contrast, carried bla_{SFH-1}

and bla_{SFC-1}, though a synonymous mutation was detected in bla_{SFH-1} when compared with

bla_{SFH-1} in UTAD54. Although detailed genetic contexts of the bla_{SFH} and bla_{SFC} genes are not

known, these genes were previously shown to be located on the chromosome of UTAD54 by

probe hybridization.[47, 48] The bla_{SFH-1} and bla_{SFC} genes in the river isolates sequenced to closure

in this study (JSRIV001, JSRIV002, JSRIV004, JSRIV006) were also located on chromosomes

and situated close to each other (separated by 76 kbp to 131 kbp). Given the sporadic

phylogenetic distribution (**Figure 4**), the two genes were suspected to be located on GIs.[51]

Comparative analysis of the chromosomes of four river isolates and the chromosome of *S.*

fonticola clade III strain GS2 (GCF_001514455.1) revealed that the two genes were located on

putative GIs inserted at tRNA-Phe genes. (**Figure 5A**). Importantly, IslandViewer 4 predicted

almost the same regions as potential GIs (**Figure S5**). Perfect (5′-

TCGATTCCGAGTCCGGGCACCA-3′, detected in JSRIV004) or imperfect (5′-

TCGATTCCGAGTCC-GGCACCA-3′, detected in JSRIV001, JSRIV002, and JSRIV006)

repeat sequences corresponding to the 3′-end of the tRNA-Phe gene were detected at the right

ends of the GIs. Some previous studies also reported imperfect repeats at the right ends of GIs.[52,]

[53] An integrase gene was detected ~200 bp downstream of the tRNA-Phe gene in each GI

(**Figure 5B**), which indicates that the GIs were integrated at tRNA-Phe genes by site-specific

recombination mediated by these integrases. Genetic contexts of bla_{SFH-1} were highly conserved

and the bla_{SFH-1} gene was surrounded by a ΔIS*Ec18*-like element and an IS*1N*-like element in

all four completed chromosomes (**Figure 5B**). Although the complete genome was not

available for strain UTAD54, analysis of a contig carrying bla_{SFH-1} revealed that the gene was

embedded within the same genetic context and also carried by a putative GI which was inserted

at the tRNA-Phe gene (**Figure 5B**). In draft assemblies of JSRIV003, JSRIV005, JSRIV007,

JSRIV008 and JSRIV009, contigs were broken near bla_{SFH-1} and genetic contexts of bla_{SFH-1}

could not be determined. Genetic surroundings of bla_{SFC} were also highly conserved, though

variations were observed in regions >3,000 bp away from bla_{SFC} (**Figure 5C**). bla_{SFC} was

surrounded by genes such as *smvA* (encoding a methyl viologen resistance protein), *acrR*

(encoding a transcriptional regulator), and genes encoding hypothetical proteins. The genetic

surroundings of bla_{SFC} (<3,000 bp from bla_{SFC}) were also conserved in draft assemblies of

JSRIV003, JSRIV005, JSRIV007, JSRIV008 and JSRIV009. Although bla_{SFH} and bla_{SFC} seem

to be mobilized together within the same GIs, strain AU-AP2C carried only bla_{SFC-3} and lacked

bla_{SFH}. A contig carrying bla_{SFC-3} in the draft genome assembly of AU-AP2C was long enough

for the comparative analysis. In AU-AP2C, a putative GI was inserted at the tRNA-Phe gene,

and bla_{SFC-3} was also carried on this putative GI (**Figure S6A**). However, the region containing

the integrase gene and bla_{SFH-1} was replaced by another segment as shown in **Figure S6B**.

Three *S. ureilytica* isolates from the lake, JBIWA004, JBIWA011 and JBIWA012, carried

bla_{IMI-22} and bla_{IMI-23}. Among the three isolates, the genome of JBIWA004 was sequenced to

closure. The two bla_{IMI} genes in JBIWA004 were located on a 106,702 bp plasmid, named

pJBIWA004_1, with IncFII(pRSB107) and pSM22 replicons. Online BLASTN analysis

revealed the plasmid p2-125 from *Serratia liquefaciens* showed the highest coverage (55%)

and also showed the high percent identity (96%) to pJBIWA004_1, while other plasmids

showed <30% coverage. Alignment of pJBIWA004_1 and p2-125 revealed that the two

plasmids shared conjugation regions, but the region containing the bla_{IMI} genes was only

present in pJBIWA004_1 (**Figure 6A**). Comparison of the genetic contexts of the two bla_{IMI}

genes and that of bla_{IMI-1} in *Enterobacter cloacae* complex strain N11-1168 showed that the

regulator *imiR* gene and a gene encoding a hypothetical protein, which are usually present

upstream and downstream of bla_{IMI} genes, were truncated or deleted in pJBIWA004_1 (**Figure**

6B). bla_{IMI-22} and bla_{IMI-23} were surrounded by remnants of ISs, which might have contributed

to their mobilization. Although only draft genome assemblies were available for JBIWA011

and JBIWA012, the bla_{IMI-22} and bla_{IMI-23} genes were located on a single contig in each assembly.

Moreover, those contigs showed >94% coverage and >99.9% identity to pJBIWA004_1,

indicating that JBIWA011 and JBIWA012 also carried bla_{IMI-22} and bla_{IMI-23} on pJBIWA004_1-

like plasmids.

Figure 4. Phylogenetic tree constructed using RefSeq assemblies labelled *S. fonticola* and

assemblies of nine river *Serratia* isolates. The RefSeq assemblies were downloaded in March

2021. The tree was constructed based on 177,349 SNPs occurring in at least 80% of the

genomes. Four RefSeq assemblies (GCF_006714955.1, GCF_006715025.1,

GCF_007827525.1, GCF_014205655.1) were removed because these assemblies showed <

85% ANI values with the type strain of *S. fonticola*. The mid-point rooted tree was visualized

with ggtree.[54] The black blocks represent the presence of a carbapenemase gene.

Figure 5. (A) Alignment of the putative GIs and the surrounding regions of river *Serratia*

isolates sequenced to closure and the corresponding chromosomal region of *S. fonticola* clade

III strain GS2. Positions of the bla_{SFC-1}, bla_{SFC-2}, bla_{SFH-1}, and tRNA-Phe genes, and (imperfect)

repeats are indicated by blue, light blue, red, brown, and purple triangles, respectively. Double-

headed arrows indicate genomic regions corresponding to GIs. Genetic contexts of (B) bla_{SFH}

and (C) bla_{SFC}. Red arrows indicate bla_{SFH-1}, blue arrows indicate bla_{SFC-1}, a light blue arrow

indicates bla_{SFC-2}, a yellow arrow indicates bla_{SFC-3}, green boxes indicate insertion sequences,

and orange arrows indicate other coding sequences. Dashed lines indicate contig breaks.

Figure 6. (A) Alignment of pJBIWA004_1 and p2-125. Positions of bla_{IMI-22} and bla_{IMI-23} are

indicated by blue and red triangles, respectively. (B) Genetic contexts of bla_{IMI} genes. A blue

arrow indicates bla_{IMI-22}, a red arrow indicates bla_{IMI-23}, a light blue arrow indicates bla_{IMI-1}, green

boxes indicate insertion sequences, and orange arrows indicate other coding sequences.

Conclusions

Here, we identified the prevalence of minor carbapenemases among environmental CPE

isolates in Japan. Of note, bla_{SFH} and bla_{SFC}, which have been described only in one *S. fonticola*

isolate in Portugal, were prevalent among river *S. fonticola* isolates obtained from different

sampling sites. The results of this study indicate aquatic environments as potential reservoirs

of minor carbapenemases and highlight the need to continuously monitor antibiotic resistance

in the environment.

Acknowledgements

Computations were partially performed on the NIG supercomputer at ROIS National Institute

of Genetics.

Funding

This work was supported by the Japan Society for the Promotion of Science KAKENHI (grant

number JP19K20461) and the Environment Research and Technology Development Fund

(JPMEERF20205006) of the Ministry of the Environment Japan.

Transparency declarations

None to declare.

References

1.   Tzouvelekis, L. S.;   Markogiannakis, A.;   Piperaki, E.;   Souli, M.; Daikos, G. L.,
Treating infections caused by carbapenemase-producing Enterobacteriaceae. *Clin Microbiol*
Infect **2014,** *20* (9), 862-72.
2.   Doi, Y.; Paterson, D. L., Carbapenemase-producing *Enterobacteriaceae*. *Semin Respir*
Crit Care Med **2015,** *36* (1), 74-84.
3.   Bonnin, R. A.;   Jousset, A. B.;   Emeraud, C.;   Oueslati, S.;   Dortet, L.; Naas, T.,
Genetic Diversity, Biochemical Properties, and Detection Methods of Minor Carbapenemases
in Enterobacterales. *Front Med (Lausanne)* **2020,** *7*, 616490.

4.   Marti, E.; Variatza, E.; Balcazar, J. L., The role of aquatic ecosystems as reservoirs
of antibiotic resistance. *Trends Microbiol* **2014,** *22* (1), 36-41.

5.   Zainab, S. M.; Junaid, M.; Xu, N.; Malik, R. N., Antibiotics and antibiotic resistant
genes (ARGs) in groundwater: A global review on dissemination, sources, interactions,
environmental and human health risks. *Water Res* **2020,** *187*, 116455.

6.   Berglund, B., Environmental dissemination of antibiotic resistance genes and
correlation to anthropogenic contamination with antibiotics. *Infect Ecol Epidemiol* **2015,** *5*,
28564.

7.   Bleichenbacher, S.; Stevens, M. J. A.; Zurfluh, K.; Perreten, V.; Endimiani,
A.; Stephan, R.; Nuesch-Inderbinen, M., Environmental dissemination of carbapenemase-
producing Enterobacteriaceae in rivers in Switzerland. *Environ Pollut* **2020,** *265* (Pt B),
115081.

8.   Tafoukt, R.; Touati, A.; Leangapichart, T.; Bakour, S.; Rolain, J. M.,
Characterization of OXA-48-like-producing *Enterobacteriaceae* isolated from river water in
Algeria. *Water Res* **2017,** *120*, 185-189.

9.   Suzuki, Y.; Nazareno, P. J.; Nakano, R.; Mondoy, M.; Nakano, A.;
Bugayong, M. P.; Bilar, J.; Perez, M. t.; Medina, E. J.; Saito-Obata, M.; Saito, M.;
Nakashima, K.; Oshitani, H.; Yano, H., Environmental Presence and Genetic Characteristics
of Carbapenemase-Producing *Enterobacteriaceae* from Hospital Sewage and River Water in
the Philippines. *Appl Environ Microbiol* **2020,** *86* (2).

10.   Teixeira, P.; Tacao, M.; Pureza, L.; Goncalves, J.; Silva, A.; Cruz-Schneider,
M. P.; Henriques, I., Occurrence of carbapenemase-producing *Enterobacteriaceae* in a
Portuguese river: *bla*$_{NDM}$, *bla*$_{KPC}$ and *bla*$_{GES}$ among the detected genes. *Environ Pollut* **2020,**
260, 113913.

11.   Mathys, D. A.; Mollenkopf, D. F.; Feicht, S. M.; Adams, R. J.; Albers, A. L.;
Stuever, D. M.; Grooters, S. V.; Ballash, G. A.; Daniels, J. B.; Wittum, T. E.,
Carbapenemase-producing *Enterobacteriaceae* and *Aeromonas* spp. present in wastewater
treatment plant effluent and nearby surface waters in the US. *PLoS One* **2019,** *14* (6), e0218650.

12.   Ohno, Y.; Nakamura, A.; Hashimoto, E.; Matsutani, H.; Abe, N.; Fukuda,
S.; Hisashi, K.; Komatsu, M.; Nakamura, F., Molecular epidemiology of carbapenemase-
producing Enterobacteriaceae in a primary care hospital in Japan, 2010-2013. *J Infect*
Chemother **2017,** *23* (4), 224-229.

13.   Eda, R.; Nakamura, M.; Takayama, Y.; Maehana, S.; Nakano, R.; Yano, H.;
Kitasato, H., Trends and molecular characteristics of carbapenemase-producing
Enterobacteriaceae in Japanese hospital from 2006 to 2015. *J Infect Chemother* **2020,** *26* (7),
667-671.

14.   Cherak, Z.; Loucif, L.; Moussi, A.; Rolain, J. M., Carbapenemase-producing
Gram-negative bacteria in aquatic environments: a review. *J Glob Antimicrob Resist* **2021,** *25*,

287-309.

15.   CLSI, Performance Standards for Antimicrobial Susceptibility Testing. 30th ed. CLSI

supplement M100. Clinical and Laboratory Standards Institute, Wayne, PA. **2020**.

16.   Nordmann, P.;   Poirel, L.; Dortet, L., Rapid detection of carbapenemase-producing

Enterobacteriaceae. *Emerg Infect Dis* **2012,** *18* (9), 1503-7.

17.   Chen, S.;   Zhou, Y.;   Chen, Y.; Gu, J., fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **2018,** *34* (17), i884-i890.

18.   Wick, R. R.;   Judd, L. M.;   Gorrie, C. L.; Holt, K. E., Unicycler: Resolving bacterial

genome assemblies from short and long sequencing reads. *PLoS Comput Biol* **2017,** *13* (6),

e1005595.

19.   Morita, H.;   Kuwahara, T.;   Ohshima, K.;   Sasamoto, H.;   Itoh, K.;   Hattori, M.;

Hayashi, T.; Takami, H., An improved DNA isolation method for metagenomic analysis of the

microbial flora of the human intestine. *Microbes Environ* **2007,** *22* (3), 214-222.

20.   Kolmogorov, M.;   Yuan, J.;   Lin, Y.; Pevzner, P. A., Assembly of long, error-prone

reads using repeat graphs. *Nat Biotechnol* **2019,** *37* (5), 540-546.

21.   Walker, B. J.;   Abeel, T.;   Shea, T.;   Priest, M.;   Abouelliel, A.;   Sakthikumar,

S.;   Cuomo, C. A.;   Zeng, Q.;   Wortman, J.;   Young, S. K.; Earl, A. M., Pilon: an

integrated tool for comprehensive microbial variant detection and genome assembly

improvement. *PLoS One* **2014,** *9* (11), e112963.

22.   Yoon, S. H.;   Ha, S. M.;   Lim, J.;   Kwon, S.; Chun, J., A large-scale evaluation of

algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* **2017,** *110* (10),

1281-1286.

23.   Meier-Kolthoff, J. P.;   Auch, A. F.;   Klenk, H. P.; Goker, M., Genome sequence-

based species delimitation with confidence intervals and improved distance functions. *BMC*

Bioinformatics **2013,** *14*, 60.

24.   Ciufo, S.;   Kannan, S.;   Sharma, S.;   Badretdin, A.;   Clark, K.;   Turner, S.;

Brover, S.;   Schoch, C. L.;   Kimchi, A.; DiCuccio, M., Using average nucleotide identity to

improve taxonomic assignments in prokaryotic genomes at the NCBI. *Int J Syst Evol Micr*

2018, *68* (7), 2386-2392.

25.   Miyoshi-Akiyama, T.;   Hayakawa, K.;   Ohmagari, N.;   Shimojima, M.; Kirikae,

T., Multilocus sequence typing (MLST) for characterization of *Enterobacter cloacae*. *PLoS*

One **2013,** *8* (6), e66358.

26.   Gardner, S. N.;   Slezak, T.; Hall, B. G., kSNP3.0: SNP detection and phylogenetic

analysis of genomes without genome alignment or reference genome. *Bioinformatics* **2015,** *31*

(17), 2877-8.

27.   Carattoli, A.;   Zankari, E.;   Garcia-Fernandez, A.;   Voldby Larsen, M.;   Lund,

O.;   Villa, L.;   Moller Aarestrup, F.; Hasman, H., In silico detection and typing of plasmids

using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother*

2014, *58* (7), 3895-903.

28.   Robertson, J.; Nash, J. H. E., MOB-suite: software tools for clustering, reconstruction and typing of plasmids from draft assemblies. *Microb Genom* **2018,** *4* (8).

29.   Siguier, P.; Perochon, J.; Lestrade, L.; Mahillon, J.; Chandler, M., ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* **2006,** *34* (Database issue), D32-6.

30.   Aziz, R. K.; Bartels, D.; Best, A. A.; DeJongh, M.; Disz, T.; Edwards, R. A.; Formsma, K.; Gerdes, S.; Glass, E. M.; Kubal, M.; Meyer, F.; Olsen, G. J.; Olson, R.; Osterman, A. L.; Overbeek, R. A.; McNeil, L. K.; Paarmann, D.; Paczian, T.; Parrello, B.; Pusch, G. D.; Reich, C.; Stevens, R.; Vassieva, O.; Vonstein, V.; Wilke, A.; Zagnitko, O., The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* **2008,** *9*, 75.

31.   Tanizawa, Y.; Fujisawa, T.; Nakamura, Y., DFAST: a flexible prokaryotic genome annotation pipeline for faster genome publication. *Bioinformatics* **2018,** *34* (6), 1037-1039.

32.   Bortolaia, V.; Kaas, R. S.; Ruppe, E.; Roberts, M. C.; Schwarz, S.; Cattoir, V.; Philippon, A.; Allesoe, R. L.; Rebelo, A. R.; Florensa, A. F.; Fagelhauer, L.; Chakraborty, T.; Neumann, B.; Werner, G.; Bender, J. K.; Stingl, K.; Nguyen, M.; Coppens, J.; Xavier, B. B.; Malhotra-Kumar, S.; Westh, H.; Pinholt, M.; Anjum, M. F.; Duggett, N. A.; Kempf, I.; Nykasenoja, S.; Olkkola, S.; Wieczorek, K.; Amaro, A.; Clemente, L.; Mossong, J.; Losch, S.; Ragimbeau, C.; Lund, O.; Aarestrup, F. M., ResFinder 4.0 for predictions of phenotypes from genotypes. *J Antimicrob Chemother* **2020**.

33.   Moura, A.; Soares, M.; Pereira, C.; Leitao, N.; Henriques, I.; Correia, A., INTEGRALL: a database and search engine for integrons, integrases and gene cassettes. *Bioinformatics* **2009,** *25* (8), 1096-8.

34.   Bertelli, C.; Laird, M. R.; Williams, K. P.; Simon Fraser University Research Computing, G.; Lau, B. Y.; Hoad, G.; Winsor, G. L.; Brinkman, F. S. L., IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Res* **2017,** *45* (W1), W30-W35.

35.   Tijet, N.; Boyd, D.; Patel, S. N.; Mulvey, M. R.; Melano, R. G., Evaluation of the Carba NP test for rapid detection of carbapenemase-producing Enterobacteriaceae and *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* **2013,** *57* (9), 4578-80.

36.   Boyd, D. A.; Mataseje, L. F.; Davidson, R.; Delport, J. A.; Fuller, J.; Hoang, L.; Lefebvre, B.; Levett, P. N.; Roscoe, D. L.; Willey, B. M.; Mulvey, M. R., *Enterobacter cloacae* Complex Isolates Harboring bla_{NMC-A} or bla_{IMI}-Type Class A Carbapenemase Genes on Novel Chromosomal Integrative Elements and Plasmids. *Antimicrob Agents Chemother* **2017,** *61* (5).

37.   Ho, P. L.; Wang, Y.; Wing-Sze Tse, C.; Fung, K. S.; Cheng, V. C.; Lee, R.; To, W. K.; Lai, R. W.; Luk, W. K.; Que, T. L.; Tsang, D. N., Rapid Detection of

Carbapenemase Production in Enterobacteriaceae by Use of a Modified Paper Strip Carba NP Method. *J Clin Microbiol* **2018,** *56* (1).

38. Umeda, K.; Nakamura, H.; Fukuda, A.; Matsumoto, Y.; Motooka, D.; Nakamura, S.; Yasui, Y.; Yoshida, H.; Kawahara, R., Genomic characterization of clinical *Enterobacter roggenkampii* co-harbouring bla_{IMP-1}- and bla_{GES-5}-encoding IncP6 and *mcr-9*-encoding IncHI2 plasmids isolated in Japan. *J Glob Antimicrob Resist* **2020,** *24*, 220-227.

39. Zong, Z.; Feng, Y.; McNally, A., Carbapenem and Colistin Resistance in *Enterobacter*: Determinants and Clones. *Trends Microbiol* **2021**.

40. Wu, W.; Feng, Y.; Zong, Z., Precise Species Identification for *Enterobacter*: a Genome Sequence-Based Study with Reporting of Two Novel Species, *Enterobacter quasiroggenkampii* sp. nov. and *Enterobacter quasimori* sp. nov. *mSystems* **2020,** *5* (4).

41. Chun, J.; Oren, A.; Ventosa, A.; Christensen, H.; Arahal, D. R.; da Costa, M. S.; Rooney, A. P.; Yi, H.; Xu, X. W.; De Meyer, S.; Trujillo, M. E., Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Micr* **2018,** *68* (1), 461-466.

42. Gomi, R.; Matsuda, T.; Yamamoto, M.; Chou, P. H.; Tanaka, M.; Ichiyama, S.; Yoneda, M.; Matsumura, Y., Characteristics of Carbapenemase-Producing Enterobacteriaceae in Wastewater Revealed by Genomic Analysis. *Antimicrob Agents Chemother* **2018,** *62* (5).

43. Adachi, F.; Sekizuka, T.; Yamato, M.; Fukuoka, K.; Yamaguchi, N.; Kuroda, M.; Kawahara, R., Characterization of FRI carbapenemase-producing *Enterobacter* spp. isolated from a hospital and the environment in Osaka, Japan. *J Antimicrob Chemother* **2021**.

44. Brouwer, M. S. M.; Tehrani, K.; Rapallini, M.; Geurts, Y.; Kant, A.; Harders, F.; Mashayekhi, V.; Martin, N. I.; Bossers, A.; Mevius, D. J.; Wit, B.; Veldman, K. T., Novel Carbapenemases FLC-1 and IMI-2 Encoded by an *Enterobacter cloacae* Complex Isolated from Food Products. *Antimicrob Agents Chemother* **2019,** *63* (6).

45. Meunier, D.; Findlay, J.; Doumith, M.; Godoy, D.; Perry, C.; Pike, R.; Gronthoud, F.; Shryane, T.; Poirel, L.; Welfare, W.; Woodford, N.; Hopkins, K. L., FRI-2 carbapenemase-producing *Enterobacter cloacae* complex in the UK. *J Antimicrob Chemother* **2017,** *72* (9), 2478-2482.

46. Sullivan, M. J.; Petty, N. K.; Beatson, S. A., Easyfig: a genome comparison visualizer. *Bioinformatics* **2011,** *27* (7), 1009-10.

47. Henriques, I.; Moura, A.; Alves, A.; Saavedra, M. J.; Correia, A., Molecular characterization of a carbapenem-hydrolyzing class A beta-lactamase, SFC-1, from *Serratia fonticola* UTAD54. *Antimicrob Agents Chemother* **2004,** *48* (6), 2321-4.

48. Saavedra, M. J.; Peixe, L.; Sousa, J. C.; Henriques, I.; Alves, A.; Correia, A., Sfh-I, a subclass B2 metallo-beta-lactamase from a *Serratia fonticola* environmental isolate. *Antimicrob Agents Chemother* **2003,** *47* (7), 2330-3.

49.   Henriques, I.;   Juca Ramos, R. T.;   Barauna, R. A.;   de Sa, P. H.;   Marinho
Almeida, D.;   Carneiro, A. R.;   Barbosa, S.;   Pereira, A.;   Alves, A.;   Saavedra, M. J.;
Egas, C.;   Silva, A.; Correia, A., Draft Genome Sequence of *Serratia fonticola* UTAD54, a
Carbapenem-Resistant Strain Isolated from Drinking Water. *Genome Announc* **2013,** *1* (6).
50.   Devi, U.;   Khatri, I.;   Kumar, N.;   Sharma, D.;   Subramanian, S.; Saini, A. K.,
Draft Genome Sequence of Plant-Growth-Promoting Rhizobacterium *Serratia fonticola* Strain
AU-AP2C, Isolated from the Pea Rhizosphere. *Genome Announc* **2013,** *1* (6).
51.   Langille, M. G. I.;   Hsiao, W. W. L.; Brinkman, F. S. L., Detecting genomic islands
using bioinformatics approaches. *Nature Reviews Microbiology* **2010,** *8* (5), 372-382.
52.   Al-Hasani, K.;   Rajakumar, K.;   Bulach, D.;   Robins-Browne, R.;   Adler, B.;
Sakellaris, H., Genetic organization of the she pathogenicity island in *Shigella flexneri* 2a.
Microb Pathog **2001,** *30* (1), 1-8.
53.   Williams, K. P., Integration sites for genetic elements in prokaryotic tRNA and tmRNA
genes: sublocation preference of integrase subfamilies. *Nucleic Acids Res* **2002,** *30* (4), 866-
75.
54.   Yu, G. C.;   Smith, D. K.;   Zhu, H. C.;   Guan, Y.; Lam, T. T. Y., GGTREE: an R
package for visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol Evol* **2017,** *8* (1), 28-36.

584