Sequential Temporal Anticipation Characterized by Neural Power Modulation and in Recurrent Neural Networks

Authors: Xiangbin Teng [a, 1, *], Ru-Yuan Zhang [b, c, 1]

Affiliation:

[a] Department of Psychology, The Chinese University of Hong Kong, 999077, Hong Kong SAR, China
[b] School of Psychology, Shanghai Jiao Tong University, Shanghai, China. 200030
[c] Shanghai Mental Health Center, School of Medicine, Shanghai Jiao Tong University, Shanghai, China. 200030
[1] These authors contributed equally.

* Corresponding author:

Xiangbin Teng
Department of Psychology
The Chinese University of Hong Kong
Shatin, New Territories
999077 Hong Kong SAR
xiangbin.teng@gmail.com

Keywords: auditory temporal processing, memory-driven prediction, brain oscillations, dynamic gain control, recurrent neural network

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgement: We thank Johannes Messerschmidt, Anna-Lena Possmayer, and Dominik Thiele for their assistance with data collection. We thank Xing Tian and Nai Ding for their comments on a previous version of the manuscript. We thank David Poeppel for support.

Funding: This work was supported by the Max-Planck-Society. R.Y.Z was supported by National Natural Science Foundation of China (32100901) and Natural Science Foundation of Shanghai (21ZR1434700). X.T. was supported by Improvement on Competitiveness in Hiring New Faculties Funding Scheme, the Chinese University of Hong Kong (4937113).

Author contributions:
Conceived and designed the experiments: X.T; Performed the experiments: X.T; Analyzed the data: X.T; Contributed reagents/materials/analysis tools: X.T & R.Y.Z; Wrote the first draft of the paper: X.T; Wrote the paper: X.T & R.Y.Z

Abstract

Relevant prospective moments arise intermittently, while most of time is filled with irrelevant events, or noise, that constantly bombard our sensory systems. Thus, anticipating a few key moments necessitates disregarding what lies between the present and the future – the noise. Here, through examining how the brain and the recurrent neural networks (RNNs) anticipate a sequence of prospective moments without relying on any external timing cues, we provided a reinterpretation of temporal anticipation. We first designed a "premembering" task, where we marked three temporal locations in white noise and asked human listeners to detect a tone at one of these points. Using power modulation analyses, we investigated the memory-guided anticipatory processes in trials involving only flat noise. Our research revealed a unique neural-power modulation pattern for sequential temporal anticipation: neural power within the alpha-beta band range fluctuates over time, accurately identifying the marked locations on a sub-second scale and correlating with tone detection performance. To understand the functional roles of these neural modulations, we utilized RNNs optimized for the behavioral task. The hidden dynamics of the RNNs mirrored the neural modulations, and additional analyses and perturbations on the RNNs indicated that the neural power modulations in the alpha-beta band resulted from selective suppression of irrelevant noise periods and heightened sensitivity to anticipated temporal locations. Our behavioral, neural, and modeling findings collectively indicate that sequential temporal anticipation involves a process of dynamic gain control: anticipating a few significant moments involves actively disregarding irrelevant events that frequently occur.

Introduction

In a common laboratory setup for studying temporal anticipation, a subject - either an animal

or a human - is tasked with predicting a brief sensory target, such as a tone or visual pattern.

These targets usually last for only a few hundred milliseconds or less but are embedded

within a much longer sequence of stimuli, noise, or silence that can last for seconds. This

scenario also happens in our daily lives. For instance, we estimate the moment a traffic light

will turn green to promptly start our car, despite waiting for minutes at a large intersection. In

both the laboratory setting and the daily life, the key prospective moments are akin to small

islands amidst a sea of irrelevant events or noise. Thus, the role of 'temporal anticipation'

likely extends beyond merely predicting a signal; it also involves ignoring the 'noise' that

exists between the present and future. In this study, we explore sequential temporal

anticipation to test how both human participants and recurrent neural networks (RNNs)

predict future events in noisy environments.

An ecological task that the brain solves is to anticipate several time points sequentially

– 'sequential temporal anticipation', such as in communicating speech and music (Nobre &

van Ede, 2018; Rimmele et al., 2018). Take the example of learning to play the piano. While

the musical notes in a melody are played one after the other at specific intervals, piano players

plan a sequence of actions, which necessitates their brain to anticipate a series of future time

points. In this scenario of sequential anticipation, the brain must predict each moment, suppress

what follows, and prepare for the subsequent crucial moment in a timely manner. However,

this dynamic, contiguous process of temporal anticipation is not fully captured by previous

studies, which primarily focus on situations where only the timing of a single target event is

anticipated. Examples of these situations include anticipating the onset of an event, estimating

the duration of an interval, or predicting the end of a lasting event (Coull & Nobre, 1998;

Martens & Johnson, 2005; Niemi & Näätänen, 1981; Nobre et al., 2007; Nobre, 2001; Samaha

et al., 2015). In these studies of 'single' temporal anticipation, the task is considered complete

once the target moment is reached, with no further process required. In this sense, sequential

temporal anticipation offers a more thorough understanding of temporal anticipation compared

to previous studies that only anticipate a single target event. This is the reason we have chosen

to study sequential temporal anticipation.

We aim to investigate the brain's initiation of sequential temporal anticipation, as in the

ecological case of music playing. One experimental model for investigating temporal

anticipation involves presenting a series of sensory events, such as tones or visual images, and

requiring human participants to predict the timing of these events (Haegens & Zion Golumbic,

2018; Jones et al., 1981; Rimmele et al., 2018; Zanto et al., 2006). However, the sensory events

offer external timing cues. When rhythmic sequences are presented, temporal anticipation can

be based on neural oscillatory patterns that are entrained by these sequences that provide

external timing cues (Arnal et al., 2015; Doelling et al., 2019; Doelling & Poeppel, 2015;

Henry et al., 2014; Henry & Obleser, 2012; Herrmann et al., 2016; Morillon et al., 2014;

Morillon et al., 2016; Spaak et al., 2014; Ten Oever et al., 2017; van Bree et al., 2021) (Tal et

al., 2017). Even when the intervals between events in a sequence are varied, the sequence can

still serve as a temporal context for anticipating an incoming event, once the temporal

contingencies of the sequence are learned (A. Breska & L. Y. Deouell, 2017; Breska & Ivry,

2020; Morillon et al., 2016; Paton & Buonomano, 2018; Rimmele et al., 2018; Wilsch et al.,

2015). For example, the timing of an upcoming event can be predicted by using the previous

event as a temporal cue and estimating the duration of a fixed interval. Consequently, it

becomes challenging to isolate the sequential temporal anticipation that is internally initiated

by the brain – a new paradigm is required in this case.

We explored how the brain anticipates a series of prospective moments without the

need for external timing cues. We adapted a straightforward tone-in-noise model, asking

human participants to identify a faint tone at one of three set time points within a segment of

white noise. To successfully detect the faint target tone, participants had to remember the

time points and anticipate them in sequence within the constant white noise. This can be

considered as memory-guided temporal anticipation (Cravo et al., 2017; Grabenhorst et al.,

2019; Mattiesing et al., 2017; Nobre & Stokes, 2019; Nobre & van Ede, 2018), not for the

timing of a single event, but for three potential moments. The white noise segment provides a

global temporal context, limiting the time points of the target tone; the tone time points must

be defined by referencing the onset of the noise or the duration of the noise period. Past

research has indirectly studied temporal anticipation by observing its effects on behavioral

and neural responses to sensory events (Grabenhorst et al., 2019; Tavano et al., 2019) and by

employing an encoding framework (Herbst et al., 2018). However, here, we directly

investigate how the brain internally manages sequential temporal anticipation within flat

white noise.

In addition to testing human participants, to gain insights on the functional aspects of

our neural findings and uncover computational mechanisms underlying the sequential

temporal anticipation, we turned to RNNs (recurrent neural networks) trained under the same

behavioral paradigm in the current study. One advantage of employing neural network

models is that RNNs are flexible to train and can be tested on variations of the current

behavioral task to test different hypotheses (Yang & Wang, 2020). If the trained RNNs can

indeed accomplish the behavioral task used in the current study and show a task performance

comparable to the human participants' performance, we then study RNN hidden state

activities and structures to gain insights on what procedures are conducted to accomplish the

tone detection task and anticipate the temporal locations (Daniel LK Yamins et al., 2014;

Yang et al., 2019).

It is challenging to identify internal anticipating processes without resorting to any

external markers. We developed a data-driven approach, using modulation spectrum analyses

on group-level neural power and phase measurements, to study internal anticipation

processes. This method relies on the premise that if all human participants anticipate the same

temporal structure, their neural dynamics should be similar. We found shared neural

modulation components in the alpha-band range (12 – 30 Hz) that marked anticipated

temporal locations and were correlated with tone detection performance. These could not be

fully explained by expectation violation or false alarms. Next, we analyzed 12 RNNs trained

on task variations to understand the functional role of the neural power modulation. This

comparison revealed a mechanism of dynamic gain control in sequential temporal

anticipation and suggested new hypotheses for future research.

Results

Neural tagging of prospective temporal locations and behavioral results

To aid participants in remembering prospective temporal locations, we presented a 2-second long white noise segment in each trial. A faint tone, calibrated to each participant's tone detection threshold (Fig. S1A), was embedded at one of three time points: 500 ms, 1000 ms, and 1500 ms (Fig. 1A). This consistent spacing was intended to assist in identifying neural correlates of sequential temporal anticipation within the white noise, without the need for external markers (for more details, see *Methods*). Once the noise segment ended, participants indicated if they had heard a tone. In 25 percent of the trials, only white noise segments were used ('no-tone' trials), while in the remaining 75 percent, the target tone was placed at one of the three temporal locations with equal probability ('tone' trials) (Fig. 1B&S1B). The sequence of tone and no-tone trials was randomized, leaving participants unsure of when or if the target tone would occur. As a result, participants had to remember the temporal locations within the noise segments and anticipate each one to effectively detect the faint tone. This task was carried out over three experimental blocks to accumulate an adequate number of 'no-tone' trials, which would help in capturing the neural signature of sequential temporal anticipation in flat noise.

We initially conducted conventional analyses on behavioral measurements and neural responses to the target tone. These analyses not only replicated previous findings but also confirmed that our task effectively manipulated the timing of the target event and modulated the sensory processing of the target tone (Fig. 1C, S1C, S1D&S1E) (Herbst et al., 2018; Ng et al., 2012; Nobre et al., 2007; Rimmele et al., 2011; Tavano et al., 2019). We then shifted our focus to the no-tone trials. The target tone's three temporal locations were deliberately spaced 500 ms apart within the 2-second noise segments. If the brain anticipates the temporal locations on a sub-second scale, we should be able to detect neural signatures in power or phase measurements around 2 Hz (Fig. 1D). Alternatively, the brain may solely encode the elapsed time of the noise segment to continually anticipate the target tone, without considering the exact temporal location. In this scenario, we wouldn't observe a 2-Hz component, but rather a neural signature corresponding to the length of the noise segment or the trial.

Fig. 1. Experimental Paradigm, Target Tone ERPs, and Analysis Procedures. (*A*) Tone Detection Task: Each trial presented participants with a 2-second white noise piece, after which they identified if a tone was heard. The target tone could appear at 500 ms, 1000 ms, or 1500 ms (indicated by colored boxes) or not at all. Two visual feedbacks were given: the first confirmed response accuracy, the second revealed the tone location. (B) Trial Portions and Target Tone Probability: The target tone was present in 75% of trials ('tone trials'), with the remaining trials only featuring white noise ('no-tone trials'). In tone trials, the target tone could appear at any of the three temporal locations, each with a 0.33 probability. (C) ERP Waveforms of the Cz Channel: Larger neural responses were evoked by the target tone at later temporal locations. (D) Hypotheses and Analysis Procedures for Sequential Temporal Anticipation: We hypothesized that participants would anticipate the temporal locations within the noise pieces to improve target tone detection. We expected to observe neural signatures reflecting anticipating processes for the temporal locations on a sub-second scale. To identify relevant neural components, we used a data-driven method, first performing a time-frequency decomposition on recorded EEG signals in no-tone trials to extract neural patterns in phase or power at each EEG frequency. We then conducted another fast Fourier transformation on the decomposed neural data to derive a modulation spectrum. After surrogate tests, we expected to find significant modulation clusters corresponding to the timescale of the temporal locations. We then filtered the EEG data according to each cluster's spectral range for further analysis.

Robust power modulations identified at 2 Hz and in the low frequency range
We sought to identify robust neural modulations for temporal anticipation in power and phase
measurements using a data-driven method, a group-level modulation spectrum analysis (Fig.
1D). We first combined all no-tone trials from the three blocks and, following time-frequency
decomposition, obtained power and inter-trial phase coherence (ITPC). We then applied
another Fourier transform to each frequency of the power and the ITPC to derive their
modulation spectra. Subsequently, we carried out surrogate tests to pinpoint significant
clusters in the modulation spectra. The power in this context was initially derived on each
trial and then averaged over all trials; the ITPC primarily measures phase-locked, or event-
evoked, neural responses (S. Makeig et al., 2004; Mazaheri, 2022). For a more detailed
discussion on these two different measures, please refer to the *Methods*. It's also important to
note that the modulation analysis used here is entirely distinct from the modulation power
spectrum used in acoustic analysis (Elliott & Theunissen, 2009), which involves a 2D FFT
directly applied to the spectrogram of sounds.
The modulation spectrum analyses indeed uncovered a rich profile of neural
modulations in the power modulation spectrum (PMS) (Fig. 1C, 2A, 2B). Although certain
indications can be observed in the spectrograms derived from conventional power analyses
(Fig. 2A, S2A, S2B), the modulation spectrum analysis directly singles out significant power
frequencies and modulation frequencies, eliminating the need for posthoc examination. We
only identified one cluster in the modulation spectrum of ITPC, which signals the onset and
offset responses to the noise segments (Fig. S3). As a result, we did not further investigate the
neural phase domain or the phase-locked neural responses.
In the PMS, four significant modulation clusters were identified, as depicted in the
right plot of Fig. 2B. The power used in the PMS calculation was 1.5 seconds, ranging from
0.25 seconds to 1.75 seconds after the onset of noise. Modulation components below 1.33 Hz,
which did not construct two cycles over 1.5 seconds, likely represented ramping neural
activities. In contrast, modulation components above 1.33 Hz are considered rhythmic
components. Therefore, the modulation frequency of 1.33 Hz acts as a demarcation line,
separating ramping components from rhythmic ones, as seen in the left plot of Fig. 2B,
referred to as the Trend bound. As a result, the two clusters (Cluster 3 and 4) located above
the Trend bound may represent the sequential temporal anticipation of local temporal
locations. The topography of each cluster is displayed in Fig. 2C. Subsequently, we analyzed
the temporal dynamics of these four clusters.

Fig. 2. Exploring Power Modulation Spectrum (PMS) and Power Dynamics. (*A*) Displays a spectrogram of induced power from all no-tone trials. (*B*) Presents raw PMS (left plot) and significant clusters (right plot). The analysis was conducted on the EEG channel of Cz as previous research shows that anticipation-related neural signals can be largely captured by the Cz (see *Methods*). The y-axis signifies the frequency of induced power, or power frequency, while the x-axis represents the modulation frequency. In the right plot, colored bins indicate significant modulation bins with high modulation strength, as determined by a surrogate test. The four dashed boxes highlight the selected power and modulation frequency ranges for the clusters. Modulation frequencies larger than the Trend bound denote rhythmic neural components, while those smaller than Trend bound represent ramping neural components. Modulation bins below the Sampling bound violate the Nyquist theorem and are therefore not meaningful. (*C*) Topographies of significant modulation clusters for visualization, with green dots indicating significant EEG channels. (*D*) Dynamics of significant clusters, where the induced power was filtered according to the clusters selected in (*B*), showing the power dynamics of each cluster. (*E*) Merged clusters. Based on the power-power coupling between each pair of clusters (Fig. S2) and the power frequency overlaps, we grouped Cluster 2, 3, and 4 into one - Cluster 234. Cluster 1 and Cluster 234 collectively capture both the local temporal locations and the global elapsed time of the noise piece. (*F*) Corrected cluster 234, derived by removing the baseline of Cluster 234 from -1 s to 0 s. (*G*) Gaussian fits to each neural peak of Corrected cluster 234, providing peak latencies (mean of Gaussian fits) and peak widths (standard deviation). (*H*) Topography of modulation strength of Corrected cluster 234 from 0.4 s to 1.9 s post noise onset. (*I*) Peak magnitude within one standard deviation (SD) of the Gaussian fits. Each gray dot represents individual participant data. Shaded areas of color and error bars signify ±1 standard error of the mean across participants. * denotes $p < 0.05$.

The power dynamics for each cluster were derived by filtering the power according to

the selected ranges of power frequency and modulation frequency, as shown in Fig. 2B (refer

to *Methods*). Cluster 1's power frequency range falls within the delta band (1 – 4 Hz). The

temporal modulation trajectory of Cluster 1 mirrors the duration of the noise pieces, implying

that the brain is actively monitoring the elapsed time of noise pieces and mentally registering

the length of the noise piece. This aligns with previous research suggesting a correlation

between neural signals in the delta band or slow ramping activities and interval (or event

duration) estimation (A. Breska & L. Y. Deouell, 2017; van Wassenhove & Lecoutre, 2015).

This correspondence serves to validate our method used in this study. However, as this was

not our primary focus, we did not further explore this cluster.

Clusters 2, 3, and 4 do not seem to correspond directly with temporal locations.

However, we hypothesized that these clusters might collectively represent sequential

temporal anticipation. To test this, we measured the power coupling between each pair of

these clusters to understand their interrelation. We not only confirmed the previously found

coupling relationship between the delta (Cluster 1) and the beta (Cluster 2) bands (Arnal et

al., 2015) (Fig. S2C), but also discovered significant coupling relationships post-noise onset

between Clusters 2 and 3, and Clusters 3 and 4. Furthermore, Clusters 2 and 4 shared the

same power frequency range. The coupling relationships and power frequency overlaps

suggested that Clusters 2, 3, and 4 might collectively represent a shared neural process.

Consequently, we grouped them into a single entity, Cluster 234. As shown in Fig. 2E,

Cluster 1 and Cluster 234 concurrently represent the total elapsed time of the noise piece and

the local temporal locations.

The reason Clusters 2, 3, and 4 are inherently related yet individually identified in the

PMS is likely due to the Fourier transform in the PMS calculation, which requires

components to be orthogonal. Other data-driven decomposition methods might be more

effective in this case (Kevric & Subasi, 2017). However, the PMS supports an underlying

psychological hypothesis specific to our study (that, as mentioned in the introduction,

common neural dynamics should be shared among participants performing the same

anticipation task. For more details, refer to *Methods*), and provides clear results. Therefore,

we believe the PMS is suitable for our current purposes, though exploring other

decomposition methods in future research could be beneficial.

We then turned our primary attention to Cluster 234, as it effectively captured the

temporal anticipation of local temporal locations and represented a neural signature that, to

the best of our knowledge, has not been previously identified. We removed the baseline of

Cluster 234, a downward trend that spanned the entire trial (for more details, see *Methods*),

and plotted the dynamics of the corrected Cluster 234 in Fig. 2F. We also depicted the

topography of its modulation strength from 0.4 s to 1.9 s in Fig. 2H (refer to *Methods* for

further details). It is evident that the neural peaks in the corrected Cluster 234 denote the

three temporal locations in the noise pieces (Fig. 2F). After fitting a Gaussian curve to each

peak, we found the peak latencies, relative to the corresponding temporal locations, to be

0.24 s, 0.16 s, and 0.13 s, respectively (Fig. 2G). This suggests a precession of neural

dynamics along the temporal locations. We then calculated the average peak magnitudes for

each temporal location within one standard deviation (SD) of the corresponding Gaussian fit

and conducted a one-way rmANOVA. A significant main effect for Temporal location was

observed ($F(2,52) = 4.30$, $p = .019$, $\eta_p^2 = 0.142$). The peak magnitudes of the first and second

temporal locations were significantly larger than that of the third temporal location (Fig. 2I)

(paired t-test: $t(26) = 3.38$, $p = .038$; $t(26) = 2.41$, $p = .045$). To control the false positive rate

of multiple comparisons, we applied an adjusted False Discovery Rate (FDR) correction

throughout the entire paper (Benjamini & Hochberg, 1995; Yekutieli & Benjamini, 1999).

One might reasonably assume that the corrected Cluster 234 in the no-tone trials

reflects an expectation violation. This is because participants may anticipate a target tone at a

temporal location, and its absence could generate a prediction error (Friston, 2005) or a

mismatch (Heilbron & Chait, 2018; Naatanen et al., 2007), potentially inducing the neural

peaks of the corrected Cluster 234. If this were the case, the magnitudes of these neural peaks

should increase from the first to the third temporal location, as the likelihood of the target

tone's appearance increases with the elapsed time of the noise piece (Janssen & Shadlen,

2005; Nobre et al., 2007) (Fig. S1B). However, our data show that the peak magnitude of the

corrected Cluster 234 actually decreases with the temporal location (Fig. 2I), making the

expectation violation explanation unlikely. This decreasing trend in peak magnitude may

correlate with the estimation of the likelihood of a trial being a tone trial (Grabenhorst et al.,

2021) (Fig. S1B), but our subsequent analyses fitting this likelihood to individual

participants' data did not reveal a significant effect (p > 0.05). Further studies are required to

specifically test this hypothesis.

We analyzed the power dynamics in the conventionally defined frequency bands and

observed indications of power marking the temporal locations in the no-tone trials in both the

beta2 band (21 – 30 Hz) and the alpha band (8 – 12 Hz). This was after we intentionally

divided the beta band (12 to 30 Hz) into the beta1 (13 – 20 Hz) and beta2 bands (21 – 30 Hz)

(Fig. S4A). Our data-driven method here aligns with standard analysis procedures, but it

uniquely extracts key neural features from EEG signals. This is achieved without initially

plotting power dynamics in pre-defined frequency bands for observation and then conducting

post-hoc analyses.

Correlation between behavioral measurements and Corrected cluster 234

In the preceding analyses, we consolidated all no-tone trials from the three experimental

blocks to ensure adequate statistical power for the PMS analysis. This was aimed at

identifying the neural signature associated with sequential temporal anticipation, which we

were uncertain could be detected a prior. After establishing Corrected Cluster 234, we

proceeded to examine each block individually. This allowed us to explore the correlation

between behavioral performance and Corrected Cluster 234, as well as to assess any potential

learning effects across the blocks.

Fig. 3. Analyzing Power Modulation Correlation with Tone Detection Performance and Neural Power Precession. (A) Depicts power dynamics of Corrected cluster 234 in each block, with the colored vertical line representing the group mean of reaction time per block. (B) Illustrates peak magnitudes within one standard deviation (SD) of the Gaussian fits. A significant main effect of Temporal location is observed ($p < 0.05$), but not for Block number ($p > 0.05$). (C) Shows the correlation between d-prime values and neural peak magnitudes. Peak magnitudes of Corrected cluster 234, derived from no-tone trials, positively correlate with tone detection performance. Each dot represents individual participant data. (D) Presents peak latency differences of Corrected cluster 234 between blocks. Filled circles denote peak latencies derived from fitting Gaussian curves to Corrected cluster 234, with different temporal locations coded by the color scheme. Gray square areas indicate the 95% confidence interval of peak latencies from permutation tests across blocks, representing a 'Uniform' bound, a null hypothesis that all peak latencies at a temporal location do not significantly differ across blocks. If a peak latency falls outside the 'Uniform' bound, it significantly differs from other peak latencies. (E) Displays peak latency distribution. Three distributions for three temporal locations were derived from permutation tests. (F) Depicts distribution of shortened latency per temporal location. A set of peak latencies was randomly selected from distributions in (E) and a line was fitted to the selected peak latencies with the temporal location (1, 2, and 3) as the independent variable. This generated a distribution of slopes representing shortened latency per temporal location. 97.5% of the slope values fall below 0, suggesting a significant decrease in peak latency with the temporal location. The shaded areas of color in (A) and the error bars in (B) represent ±1 standard error of the mean across participants. * denotes $p < 0.05$.

Fig. 3A illustrates Corrected Cluster 234 in each block, showing that both the peak

magnitudes and peak latencies varied according to the blocks and temporal locations. By

following the procedure of fitting Gaussian curves to the power peaks (as shown in Fig. 2G),

we derived the peak magnitudes in each block (Fig. 3B & S4A). We then conducted a two-

way rmANOVA, with Temporal location and Block number as the primary factors. The main

effect of Temporal location was significant ($F(2,52) = 5.01$, $p = .010$, $\eta_p^2 = 0.162$), whereas

the main effect of Block number ($F(2,52) = .51$, $p = .604$, $\eta_p^2 = 0.019$) and the interaction

($F(4,104) = .25$, $p = .909$, $\eta_p^2 = 0.010$) were not. The peak magnitude displayed a decrease

with the temporal location, as evidenced by a significant linear trend ($F(1,26) = 9.22$, p

$= .005$, $\eta_p^2 = 0.262$). The peak magnitude around the first temporal location was significantly

larger than the third location (t(25) = 3.04, $p = .016$; adjusted FDR correction was applied).

These findings indicate that the block number did not interact with the neural coding of the

local temporal locations, suggesting that a learning effect was not observed across blocks.

244   Contrary to our initial hypothesis that learning the timing of prospective moments

would be a gradual process, participants quickly grasped the temporal locations of the target

tone within the noise pieces. They were able to anticipate these locations even in the initial

block. This may be attributed to the 10 to 15 trials in the training session we conducted prior

to the formal test, which helped familiarize participants with the task. It's possible that

participants learned the temporal locations during this training phase. Additionally, the visual

feedback provided on the temporal location of the target tone during tone trials likely

facilitated this swift learning process, as it offered clear information about each temporal

location. Future studies could delve deeper into this learning process by eliminating the

visual feedback. However, this efficient learning process proved beneficial in identifying the

neural signatures of sequential temporal anticipation, which was our primary focus.

255   A lingering question is whether the neural power modulation, which marks the local

temporal locations in the no-tone trials, is relevant to the participants' tone detection

performance in the tone trials. By correlating the peak magnitudes of Corrected Cluster 234

with the d-prime values (Fig. S1C&3C), we discovered a significant positive correlation (r =

0.24, $p < 0.001$). This indeed suggests that anticipation of the local temporal locations

enhances tone detection. We further illustrate in Fig. S4B the correlation between d-prime

values and peak magnitudes in each block, and in Fig. S4C the correlation at each temporal

location. One possible explanation for the emergence of Corrected Cluster 234 is that each

neural peak following each temporal location is due to false alarms - participants may

erroneously perceive a tone within the white noise, thereby inducing a neural peak after a

temporal location. If this were the case, the false alarm rate should positively correlate with

the peak magnitudes of Corrected Cluster 234 in the no-tone trials. However, we found that,

for the most part, the false alarm rates negatively correlate with the peak magnitudes (Fig.

S4D). Therefore, Corrected Cluster 234 was not a result of false alarms in the no-tone trials.

Precession of power dynamics suggests active anticipation

We next sought to quantify the observation from Fig. 3A that the peak latencies of Corrected

Cluster 234 varied according to the block number and temporal location. We performed

statistical tests on the peak latencies at the group level using permutation tests (refer to

Methods), as the peak latencies could not be reliably derived using the Gaussian fits

procedure on each participant's data. The permutation tests consisted of two steps. Firstly, we

examined whether the peak latencies changed across blocks. The permutation tests yielded a

95% confidence interval for each temporal location across the three blocks (refer to *Methods*

for details). We discovered that the peak latencies at the first and second temporal locations

of the second block were significantly shortened (Fig. 3D). Secondly, we investigated

whether the peak latencies changed across temporal locations. We combined the peak latency

samples generated from the permutation tests across blocks to form a peak latency

distribution for each temporal location (Fig. 3E). We then fitted a line to three peak latencies

randomly selected from the three distributions. The distribution of the slopes of the fitted

lines indicates that the peak latency significantly decreased with the temporal location (Fig.

3F). This observation is likely due to the sequential anticipation of the temporal locations -

the anticipation of the first temporal location facilitated the anticipation of the second, and so

on. We further examined this observation in the subsequent RNN perturbation analyses.

Task-optimized RNNs replicate the behavioral and neural findings and suggest a process of

dynamic gain control

We have discovered a unique neural signature, labeled as Corrected Cluster 234, which

identifies local temporal locations within the 2-s white noise segments (Fig. 2&3). This

neural signature cannot be explained by either expectation violation (Fig. S1B) or the false

alarm rate (Fig. S4D). The question then arises: what functional role does this neural

signature play in the task at hand? To answer this, we trained continuous-time Recurrent

Neural Networks (CRNNs) to perform the same task. The logic behind this is that the

CRNNs, when trained to perform the same task as human participants, might also perform

certain computations in a manner similar to the human brain (Yamins & DiCarlo, 2016; D. L.

Yamins et al., 2014). If the trained CRNNs demonstrate a behavioral performance similar to

human participants, we will study the internal dynamics of the trained RNNs to understand

how they accomplish the task. Furthermore, the flexibility of RNNs allows us to test various

conditions related to the current study, which can help generate new hypotheses for future

research.

We developed a CRNN structure adhering to Dale's rule (Dale, 1935; H. F. Song et

al., 2016) and created inputs for the CRNNs that encapsulated the essence of the behavioral

task (Fig. 1A&4A). Detailed information about the CRNN parameters and training

procedures can be found in the *SI Appendix, under Methods*. We systematically adjusted the

external noise level of the inputs while maintaining a constant target tone magnitude. This is

because if the timing of temporal locations was expected to assist in detecting the weak target

tone, the sequential temporal anticipation should be deployed more extensively at high

external noise levels (low Signal-to-Noise Ratio - SNR) and less so at lower noise levels

(high SNR). As a result, the hidden activities of the CRNNs trained at high noise levels

should exhibit dynamics with 2 Hz modulation components indicating the temporal locations,

but this should be absent or reduced at lower noise levels. If this is the case, it implies that the

2 Hz neural component depicted in Fig. 2&3 is associated with the brain's use of sequential

temporal anticipation to deal with external sensory noise.

Fig. 4. Behavioral and Neural Findings Explained by Task-Optimized Continuous-Time Recurrent Neural Networks (CRNNs). (A) Training procedures for CRNNs are depicted. The left plot shows the inputs to CRNNs, while the right plot outlines the CRNN structure and training procedures. (B) Task performance of trained CRNNs is presented. Five CRNNs were trained at five different external noise levels and tested at seven noise levels. The solid black line represents the group mean of human behavioral performance (Fig. S1C), with dashed black lines indicating ± 2 standard errors across participants. Filled gray circles mark the testing noise levels chosen for the corresponding CRNNs whose performance matches human performance. (C) Modulation spectra of hidden activities of the CRNNs at the corresponding testing noise level selected in (B) are displayed. Modulation spectra of CRNN hidden activities were derived following the EEG analysis procedures. (D) Hidden activities of two CRNNs derived from 40 no-tone trials are shown. The y-axis specifies whether the units are excitatory (red units) or inhibitory (blue units). (E) Compares CRNN hidden activities with neural data. The neural data are sourced from the group-mean data shown in Fig. 3A. The CRNN analyzed was trained at the noise level of 0.15 SD and tested at the noise level of 0.2 SD. (F) Training CRNN with a uniform probabilistic distribution of tone locations is presented. The upper plot illustrates a trial structure with a uniform distribution of tone locations, where a tone can appear at any time from 0.25 s to 1.75 s post noise onset with equal probability. The lower plot reveals that the hidden activities of the CRNN trained using the 'uniform' trial structure did not exhibit any rhythmic activities. (G) Task performance in the 'uniform' trial structure is shown. At the testing noise level of 0.2 SD, the CRNN trained in (F) and the CRNN trained with three temporal locations at the training noise level of 0.15 SD were tested. (H) Hidden activities of CRNNs with perturbed excitation-inhibition (EI) weights are depicted. The thick colored lines refer to the EI perturbations that caused the peak latencies of CRNN hidden activities to decrease with the temporal location. The violet line represents the unperturbed CRNN selected from (C), with dashed vertical lines denoting peak latencies of this unperturbed CRNN. (I) Task performance of perturbed CRNNs at the testing noise level of 0.2 SD is presented.

We trained five separate CRNNs at five different noise levels, specifically: 0.01, 0.05,

0.1, 0.15, and 0.2 standard deviations (SD) of added Gaussian noise. We then calculated the

d-prime values for each trained CRNN at various testing noise levels (Fig. 4B). For each of

these five CRNNs, we chose a specific testing noise level at which the corresponding CRNN

produced a task performance that matched the average human behavior performance (Fig.

1C&4B). Following this, we analyzed the activities of the hidden units within the selected

CRNNs and calculated the modulation spectra of their hidden activities during the

presentation of 40 no-tone trials. This was done by using the same procedure as the one used

to quantify power modulations of neural signals (Fig. 4C). As expected, the modulation

strength around 2 Hz increased with the external noise level used in training the CRNNs. The

CRNN trained at the noise level of 0.15 SD exhibited the highest modulation strength at 2

Hz, while the CRNN trained at 0.01 SD showed the least modulation strength (Fig. 4C). This

is also reflected in the temporal dynamics of their hidden activities (Fig. 4D). We averaged

the hidden activities of the CRNN trained at 0.15 SD across all hidden units, similar to EEG

recording where neural signals from both excitatory and inhibitory synapses are combined,

and plotted the averaged hidden activities of the CRNN alongside Corrected Cluster 234 (Fig.

4E). The results showed that the dynamics of the CRNN hidden activity closely mirrored the

power dynamics of Corrected Cluster 234.

One potential explanation is that the 2 Hz modulation components observed in both

the brain and the CRNNs performed a noise suppression operation – the noise period outside

of the tone locations was suppressed to facilitate the detection of the target tone. If this is the

case, a target tone might be suppressed along with the noise if it occurs at a time point other

than the three temporal locations used for training the CRNNs. Furthermore, could a CRNN

display such 2 Hz hidden dynamics simply due to high noise levels, and not because of the

specific temporal locations used? To answer these questions, we trained a CRNN with a new

trial structure. In this structure, the training noise level was set at 0.15 SD and the target tone

could appear at any time between 0.25 s and 1.75 s after the noise onset (Fig. 4F, upper

panel) - a 'uniform' trial structure. The newly trained CRNN exhibited flat hidden activities

devoid of any rhythmic patterns (Fig. 4F, lower panel). Subsequently, in the uniform trial

structure at a noise level of 0.2 SD, we tested the newly trained CRNN and the CRNN trained

with the three fixed temporal locations (both trained at a noise level of 0.15 SD). The tone

detection performance of the newly trained CRNN dropped significantly but remained within

a moderate range; in contrast, the CRNN trained with the three fixed temporal locations

demonstrated superior tone detection performance at the temporal locations but much poorer

performance at other time points (Fig. 4G). It should be noted that the calculation of d-prime

values here differed from Fig. 4B as different trial structures were used (refer to *Methods* for

details).

The findings presented in Fig. 4F&G suggest that the 2 Hz neural modulation

observed in Corrected Cluster 234 served to enhance sensory sensitivity to the target tone at

the anticipated temporal locations, while selectively suppressing sensory noise during

irrelevant noise periods. Interestingly, the power frequencies of Corrected Cluster 234 fall

within the range typically defined as the alpha and beta bands (Fig. 2B). Increasing evidence

has shown that neural activities within these alpha-beta bands are closely associated with the

suppression of internal/external noise and the regulation of information flow (Arnal &

Giraud, 2012; Haegens et al., 2012; Jensen & Mazaheri, 2010; Spitzer & Haegens, 2017;

Wostmann et al., 2019; Wöstmann et al., 2017; Zumer et al., 2014).

Our neural findings, in conjunction with the CRNN analyses, indicate that the process

of sequential temporal anticipation was partially executed as a gain control mechanism over

time (Salinas & Sejnowski, 2001; Strauß et al., 2014; van Ede et al., 2018). This is similar to

how the brain selectively filters information for future time points, akin to the way it

processes sensory information in space by focusing on attended locations and disregarding

irrelevant background noise(Hillyard et al., 1998; Reynolds & Chelazzi, 2004) .

Enhanced inhibitory activity explains precession of power dynamics

Previous studies have demonstrated that disrupting the balance between excitation and

inhibition alters the pace of neural oscillatory activities (Brunel & Wang, 2003). In our study,

we hypothesized that if the temporal locations were constant, the expedited neural peak

latencies of Corrected cluster 234 (Fig. 3E&F) could have been stimulated by an increase in

inhibitory or excitatory neural activities during the anticipation process. Following this, we

disrupted the excitation-inhibition balance in the CRNNs to investigate how alterations in this

balance influenced the peak latencies of hidden CRNN activities.

In the CRNN trained at a noise level of 0.15 SD, we manipulated the connectivity

strength of excitatory and inhibitory units in such a way that the original ratio between

excitatory and inhibitory projection weights (E/I ratio) was systematically adjusted (Fig. 4H).

For instance, we reduced the weights of excitatory units by 5% and correspondingly

increased the weights of inhibitory units in a CRNN by 5%. This resulted in weaker

projections from excitatory units to inhibitory units, and stronger projections from inhibitory

units to excitatory units. We then calculated the average hidden activities of the CRNNs with

different E/I ratios (Fig. 4H). When the E/I ratio fell below one (e.g., 95/105) – indicating

stronger inhibitory projections and weaker excitatory ones – the peak latencies decreased

with the temporal location, mirroring the neural finding shown in Fig. 3E&F. The peak

latencies of the original CRNN (E/I ratio: 100/100) remained constant with the temporal

location and were fixed at 0.188 s (the middle time point between the two largest points of

each peak, indicated by the vertical dashed lines in Fig. 4H). We then evaluated the

behavioral performance of the CRNNs with altered E/I ratios and discovered that, when the

E/I ratio of the CRNN was 95/105, the behavioral performance still fell within the range of

human behavioral performance (Fig. 4I).

In conclusion, our examination of E/I ratios in CRNNs suggests that the observed

neural power precession in Corrected cluster 234 may be linked to enhanced neural activities

of inhibitory circuits. The anticipations of earlier temporal locations facilitated the

anticipations of subsequent ones, which is associated with the increased efficiency of

inhibitory circuits. This aligns with previous neurophysiological findings and theories

suggesting that neural signals in the alpha-beta bands correlate with inhibitory activities

(Klimesch et al., 2007; Lakatos et al., 2016; Sherman et al., 2016).

Discussion

To investigate how the brain sequentially anticipates future moments, we developed a 'premembering' task by adapting a basic tone-detection paradigm, applied innovative modulation spectrum analyses to the electrophysiological data, and trained and analyzed 12 RNN models. Our research revealed that a neural modulation component in the alpha-beta band range was utilized to locally anticipate prospective temporal locations on a sub-second scale (Fig. 2). The neural modulation markers for sequential temporal anticipation were found to positively correlate with the performance of tone detection (Fig. 3), and these correlations could not be explained by expectation violation or false alarms. Utilizing task-optimized RNNs (Fig. 4), we delved deeper into the functional role of our neural findings, demonstrating that the neural power modulations underpinned a gain control process over time that reduced irrelevant noise periods and heightened sensory sensitivity at the anticipated temporal locations. Collectively, our behavioral, neural, and neural network modeling findings suggest that temporal anticipation, a psychological term for a general task requirement, was executed in this context as a dynamic gain control process.

A key focus in psychology and neuroscience is understanding how the brain copes with noise and extracts signals (Faisal et al., 2008). The brain is constantly processing sensory inputs, yet relevant and useful information is often sparse in the real world. In the context of our study, meaningful events (i.e., the target tone) potentially occur at only one of a few time points, while sensory noise is a constant presence. Therefore, the functional requirement for temporal anticipation is not only to foresee prospective moments but also to disregard irrelevant noise periods. As shown in our neural and modeling findings, neural operations in the alpha and beta bands enhanced sensory sensitivity to the anticipated temporal locations and suppressed noise outside of these locations. This discovery aligns well with the function of attention in space – in studies of spatial attention, the alpha and beta neural bands have been found to correlate with the suppression of irrelevant objects or backgrounds (Bauer et al., 2012). Our findings suggest a reinterpretation of temporal anticipation, which has often been treated as a unitary construct in various contexts (Nobre & van Ede, 2018) – the precise underlying mechanisms of temporal anticipation can vary according to functional demands.

An alternate interpretation of Corrected Cluster 234 marking the temporal locations of the target tone (as seen in Fig. 2 & 3) could be that this neural signature signifies the probability distribution of an event occurring over time (Grabenhorst et al., 2019; Janssen & Shadlen, 2005) (Fig. 1B). It would be beneficial to further investigate whether Corrected

Cluster 234 aligns with the neural representations of probability distributions observed in

animal studies by altering the distribution shapes (Janssen & Shadlen, 2005). However, it is

not always required to represent a probability distribution of event occurrence, particularly

when the signal-to-noise ratio (SNR) of tone to noise is high, simplifying the tone detection

task. Even when the probability distribution of tone occurrence remained constant, the RNN

modelling results indicated a weakening of the neural dynamics marking the tone locations as

the SNR decreased (i.e., the task became easier) (see Fig. 4C). Therefore, it's plausible that

the need to represent probabilities of event occurrences is influenced by task demands.

Further research could certainly test this hypothesis, which is suggested by our RNN models.

Regarding the brain regions involved in sequential temporal anticipation, research has

shown that, on a supra-second scale, the anticipation of incoming event timings and interval

estimation are primarily processed in the parietal and motor areas (Bolger et al., 2014; Coull

et al., 2016; Janssen & Shadlen, 2005; Merchant & Yarrow, 2016; Morillon & Baillet, 2017;

Wittmann, 2013). In terms of local temporal location anticipation, significant modulation

strength was mostly observed in the EEG channels around Cz (Fig. 2H), leading us to

hypothesize that the corresponding neural modulation component may originate from

subcortical areas. One potential area is the basal ganglia, particularly the striatum, which has

been linked to motor sequence generation and various sub-second scale timing tasks

(McClure et al., 2003; Mello et al., 2015; Motanis & Buonomano, 2015; Nozaradan et al.,

2017). As sequential actions necessitate coordination in a series of moments (Rimmele et al.,

2018; Schubotz, 2007), this functional requirement aligns with the cognitive process required

in our task, which is to anticipate a sequence of prospective moments.

In conclusion, we have identified a neural processing scheme involved in sequential

temporal anticipation. The brain utilizes neural power modulations across several neural

bands, specifically alpha and beta, to anticipate sequential local temporal structures. The task-

optimized artificial neural network models, particularly the Recurrent Neural Networks, have

helped uncover a dynamic gain control process that underpins temporal anticipation. This

process involves not only anticipating a few significant future moments, but also actively

disregarding irrelevant events that frequently occur.

Methods

Data and Code Availability

All study data and codes are available at https://osf.io/8z9h6/. Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Xiangbin Teng (xiangbinteng@cuhk.edu.hk).

Participants.

Thirty participants took part in the experiment (age 18 – 35, right-handed, sixteen females). All participants had normal hearing and no neurological deficits according to their self-report. Written informed consent was obtained from each participant before the experiment; monetary compensation was provided accordingly after the experiment was finished or terminated. Two participants could not undergo the EEG recording because of the metal fillings or decorations above their neck and chose to terminate the experiment. One participant's data was excluded because triggers for stimulus onsets were not recorded during EEG recording. Therefore, the neurophysiological data shown in the main text included twenty-seven participants (age 18 – 35, right-handed, fourteen females). The experimental protocol was approved by the Ethics Council of the Max Planck Society.

Experimental Paradigm and Stimuli

The participants were instructed to detect a 1000 Hz pure tone of 30 ms embedded in a 2-s piece of white noise at one of three temporal locations: 500 ms, 1000 ms, and 1500 ms. In each trial, there was no cue to direct participants' attention to one of the three temporal locations. We hypothesize that the participants, not knowing when the target tone would appear, had to anticipate the three temporal locations of the target tone sequentially. Our current study focused on how the participants memorized the temporal locations and anticipated the temporal locations sequentially.

The experiment was comprised of three blocks and the experimental procedures are illustrated in Fig. 1A. In each trial of each block, the participants were required to keep eyes open and to fix on a white cross in the center of a black screen. The white cross stayed on the screen for a duration randomly chosen between 1 s and 1.5 s. After the white cross disappeared, the 2-s noise piece was presented. The participants pressed one of two buttons to indicate whether they heard a tone after the noise piece was over. Visual feedback after participants' responses was shown on the screen for 0.3 s to inform the participants whether their response was 'correct' or 'wrong'. Then, another feedback was shown on the screen for

0.4 s to indicate whether and when the target tone appeared in the noise piece with one of

four symbols: '_ _ _ _' (no tone was presented), '_ | _ _ _' (a tone appeared at 500 ms after

the noise onset), '_ _ | _ _' (1000 ms), '_ _ _ | _' (1500 ms). After the two visual feedbacks,

the next trial started immediately. The feedbacks were designed to help the participants

quickly learn the tone-detection task and memorize the temporal locations.

In each of three blocks, there were four conditions: a tone appeared at either 500 ms,

1000 ms, or 1500 ms, or no tone was presented. There were 40 trials for each condition in

each block, so there were 160 trials in each of three blocks. 480 trials in total were presented.

The conditions were randomly chosen in each trial and the trial order was randomized across

blocks between participants. After each block, the participants could choose to take a break

of one or two minutes or to start the next block immediately.

The white noise piece in each trial was independently generated using a random

number generation function, 'rand', in MATLAB R2016b (The MathWorks, Natick, MA) at

a sampling rate of 20,000 Hz. As a new white noise piece was generated in each trial, the

participants could not memorize the acoustic details of white noise to help them detect the

target tone. The amplitude of white noise pieces was normalized to 70 dB SPL (sound

pressure level) by referring the noise pieces to a one-minute white noise piece that was

measured beforehand to be 70 dB SPL at the experimental setting. We applied a cosine ramp-

up function in a window of 50 ms at the onset of the noise pieces and a sine ramp-down

function of 50 ms at the offset.

The 1000 Hz target tone of 30 ms duration was generated using the function of 'sin'

in MATLAB with a fixed initial phase of 0 degree. We applied a cosine ramp-up function in

a window of 10 ms at the onset of the tone and a sine ramp-down function of 10 ms at the

offset. As the threshold for detecting a pure tone in white noise varies across individuals, we

set an individualized signal-to-noise ratio (SNR) of the target tone to the white noise for each

participant, so that the difficulty level of the task could be comparable between participants

and be controlled. We measured each participant' tone detection threshold before the formal

experiment (Fig. S1A) and set the SNR of the target tone to be 3 dB above each participant's

threshold.

To measure the tone detection threshold, we employed a two-interval two-alternative

forced choice paradigm and the PSI method (Kontsevich & Tyler, 1999; Prins, 2013) that

were implemented in the Palamedes toolbox 1.5.1 (Prins, 2016). In each trial, participants

were presented with two pieces of white noise of 1 s with an inter-piece interval randomly

sampled between 0.5 s and 1 s. The target tone was randomly inserted into one of the two

noise pieces from 0.2 to 0.8 s after the noise onset. The white noise and the target tone were

generated following the same procedures described above. The participants chose which

noise piece included the target tone by pressing one of two buttons after the second noise

piece was over. Visual feedback was provided to indicate whether the participants made the

correct response in each trial, so that the participants could learn this tone-detection task and

show stable performance in the following sessions. To get robust threshold measurements,

we measured each participant's threshold twice. In the first measurement, the starting point of

the SNR was set to -5 dB and the threshold was measured in 40 trials. In the second

measurement, the starting point of the SNR was the threshold measured from the first

measurement. Every 15 trials, there was a catch trial with a SNR of 5 dB to remind the

participants of the target tone. The threshold measured from the second time was selected as

each participant's threshold. The threshold here represented 75 percentage correct threshold.

The auditory stimuli were delivered through plastic air tubes connected to foam ear

pieces (E-A-R Tone Gold 3A Insert earphones, Aearo Technologies Auditory Systems).

EEG recording

EEG data were recorded using an actiCAP 64-channel, active electrode set (10–20 system,

Brain Vision Recorder, Brain Products, brainproducts.com) with a 0.1 Hz online filter (12

dB/octave roll-off). There were 62 scalp electrodes, and their impedances were kept below 5

kOhm. The first 18 participants of 27 valid participants were recorded at a sampling rate of

500 Hz with one electrode (originally, Oz) placed on the tip of the nose as the reference

channel. The other 9 participants were recorded at a sampling rate of 10000 Hz with one

electrode placed at Oz as the reference channel. The impedance of the reference channel was

kept below ∼10 kΩ. This difference of EEG recording settings was caused by

miscommunication between the experimenter and the research assistants who helped conduct

EEG recording. To make the EEG data compatible between the two recording settings, in the

following data analyses, EEG recordings were off-line referenced to the average of activity at

all scalp electrodes. The recording sampling rate difference should not matter here as we later

applied an off-line low-pass filter of a cut-off frequency of 30 Hz and downsampled all the

data to 100 Hz. Furthermore, the following major analyses included baseline corrections

based on pre-stimulus activities in individual participants' data and focused on modulations

of EEG power (a second-order statistics), the EEG reference differences between participants

should not affect our results.

Behavioral data analysis.

Tone detection performance was analyzed in MATLAB 2016b using the Palamedes toolbox

1.5.1 under the framework of signal detection theory. In each block, we treated the trials with

a tone embedded as the 'target' trials and the trials without a tone as the 'noise' trials. For

each temporal location in each block, we derived a hit rate from the 'target' trials ('yes' when

there was a tone) and a false alarm rate from the 'noise' trials ('yes' where there wasn't a

tone). We then calculated a d-prime value using the hit rate and the false alarm rate assuming

an independent observer model and an unbiased observer. To avoid infinite d-prime values, a

half artificial incorrect trial was added when all 'target' trials were correct; a half artificial

correct trial was added when all 'noise' trials were incorrect (Hautus et al., 2021). Note that

the 'target' trials with the target tone appeared at one of the three temporal locations shared

the same 'noise' trials, so the d-prime values and the criteria derived at each of the three

temporal locations were not independent and the differences between the temporal locations

were dominated by the hit rates.

Raw reaction time data were used in the data analysis. The reason that we did not take

a log transformation of reaction time is that previous literature demonstrated that a log

transformation of reaction time is not necessarily beneficial (Baayen & Milin, 2010).

Nevertheless, we conducted statistical tests on both log-transformed and raw reaction time

data and found that the conclusions of the statistical tests remained the same.

EEG preprocessing

EEG data analysis was conducted in MATLAB 2016b using the Fieldtrip toolbox 20181024

(Oostenveld et al., 2011) and the wavelet toolbox in MATLAB. EEG recordings were off-

line referenced to the average of activity at all scalp electrodes. Raw EEG data were first

filtered through a band-pass filter from 1 Hz to 30 Hz embedded in the Fieldtrip toolbox (a

'fir' two-pass filter with an order of 4). The cutoff frequencies of 1 Hz and 30 Hz were

selected here because previous literature showed that effects of temporal anticipation in the

auditory modality were primarily observed in the delta band (1 – 4 Hz) and the beta band (12

– 30 Hz) (Arnal et al., 2015; Morillon & Baillet, 2017).

Continuous EEG data were first divided into 6-second epochs including a 2-s pre-

stimulus period, a 2-s stimulus period, and a 2-s post-stimulus period for each trial. By

preserving long periods before and after the stimulus onset and offset, we had enough data to

calculate low-frequency EEG power (e.g. at 1 Hz) to conduct the following baseline

corrections and to remove filter artefacts in the beginning and the end of the EEG signals of

each trial. The epoched trials were then visually inspected, and those with artifacts such as

channel jumps and large fluctuations were discarded. An independent component analysis

was applied over all the trials pooling over all the blocks in each session and was used to

correct for artifacts caused by eye blinks and eye movements. After preprocessing, up to 10

trials were removed for each of the four conditions. Baseline in each trial was corrected by

subtracting out the mean of -1 s to 0 s.

We conducted various following analyses on the Cz channel as past research show

that auditory-related and anticipation-related effects can be well captured by the Cz channel.

After we found certain effects at the Cz, for visualization, we conducted the same analyses at

other channels and plotted topographies.

Event-related potential (ERP) to tones

We calculated evoked responses to the target tone at each temporal location. The baseline of

each trial was corrected as described in *EEG preprocessing*. We chose not to conduct

baseline correction again for ERPs to the target tone at different temporal locations, because

the target tone at 500 ms was affected by the noise onset response whereas the tones at 1000

ms and 1500 ms were not. Re-correcting baselines for ERP to the target tone at each temporal

location may introduce artefacts. We focused on the period of ERPs from 0 ms to 500 ms

after the tone onset in the following statistical analyses.

We pooled trials across all the three blocks and calculated ERPs. This pooling

procedure guaranteed that we had enough trials to detect differences of ERPs between three

temporal locations with sufficient statistic power. We conducted a permutation-based cluster

repeated measure ANOVA (rmANOVA) analysis over the three temporal locations (Maris &

Oostenveld, 2007), in which we shuffled the labels on the temporal locations in each

participant's data. By doing this, we constructed a null hypothesis, which was that the ERPs

to the target tone at three temporal locations do not differ from each other. During each time

of shuffling, we constructed a new dataset and conducted a one-way rmANOVA test at each

time point with Temporal location as the main factor. We identified clusters of connected

significant time points (> 2 time points) at a two-sided alpha level of 0.05 and calculated a

cluster-level F value by adding up F values within each cluster across time points. The

biggest cluster F value was selected as the cluster-level F value for each time of shuffling.

We conducted this procedure 1000 times and derived a distribution of cluster-level F values,

from which a threshold of cluster-level F values at a one-sided alpha level of 0.05 was

derived (the 950th cluster F value). We then derived significant clusters from our empirical

data and determined which cluster was significant by comparing their cluster-level F values

with the threshold of cluster-level F values. Then, we conducted paired t-tests, as post-hoc

tests, within the temporal regions of the significant clusters to compare ERPs between each

pair of temporal locations. Adjusted FDR was used for multiple comparison correction.

Time-frequency decomposition of EEG signals

To extract time-frequency information, single-trial data in each EEG channel were

transformed using functions of the Morlet wavelets embedded in the Fieldtrip toolbox, with a

frequency ranging from 1 to 40 Hz in steps of 1 Hz. Note that we did not intend to analyze

the signals above 30 Hz since we applied a low-pass filter at 30 Hz. We simply applied our

previous routine analysis parameters here.

To balance spectral and temporal resolution of the time-frequency transformation,

from 1 to 40 Hz, the window length increased linearly from 2 circles to 20 circles. Phase and

power responses (squared absolute value) were extracted from the wavelet transform output

at each time-frequency point. The temporal resolution was set as 100 Hz and the power

responses from -1 s to 3 s were selected for the following analyses.

In each condition, we first averaged power values over all the trials and then

calculated a baseline by averaging the mean power values from -.75 s to -.25 s. The averaged

power after the noise onset (0 s) was normalized by dividing the baseline and taking

logarithms with base 10, and then was converted into values with unit of decibel by

multiplying 10.

As the power was first calculated on the single-trial level and was then averaged

across trials, the resulting power was sometimes called 'induced power'; in contrast, 'evoked

power' is calculated by first averaging trials in the temporal domain and then conducting the

time-frequency decomposition (Scott Makeig et al., 2004). It has been shown that the evoked

power reflects phase-locked temporal coherence between trials because the averaging

procedure was taken before time-frequency decomposition (Teng & Poeppel, 2020).

However, this labeling of 'evoked' and 'induced' is not entirely correct because the

averaging in time does not necessarily average out non-phase locked activity (Mazaheri,

2022). Thus, we directly analyzed inter-trial phase coherence below to quantify the phase-

locked neural components across trials (Van Diepen & Mazaheri, 2018). As a result, we

simply use 'power' to indicate our calculation of power here and analyzed the phase domain

directly to represent time-locked neural activity that is arguably related to 'evoked power'.

Power modulation spectrum (PMS)

We first quantified how the power was modulated by temporal anticipation for the temporal

locations over the entire trials by calculating power modulation spectrum (PMS). We also

conducted the corresponding analysis on the phase (see *Modulation spectrum analysis on*

inter-trial phase coherence (ITPC)). If the participants could register the three temporal

locations of the target tone in their memory and actively anticipate the tagged temporal

locations every 500 ms, we would find a neural signature around 2 Hz reflecting such

sequential anticipating processes even when only the white noise was presented in each trial.

Note that this method is different from the method of directly applying 2D-FFT on

acoustic spectrograms to derive modulation power spectrum (MPS) (Elliott & Theunissen,

2009). To avoid this confusion, we intentionally named our calculation as Power Modulation

Spectrum (PMS).

We focused on the condition in which only the white noise was presented and pooled

all the no-tone trials over three blocks. We selected the neural data between 0.25 s and 1.75

after the noise onset to avoid effects of onset and offset responses. We did not define any

EEG frequency bands but used a data-driven method to find significant modulation effects on

the power from 1 Hz to 30 Hz.

We first applied a high pass filter at 0.5 Hz on the power at each frequency over entire

trials (-1 s to 3 s) so that we removed neural components over a duration larger than the

length of the white noise pieces (2 s). We next applied a hamming window of 1.5 s (151

points) on the selected time range, 0.25 s to 1.75 s after noise onset, at each frequency of

power and conducted the Fourier transform with a length of 500 points (151 data points

padded with 349 zero points). The spectral resolution of the resulted spectrum of the power

was 0.2 Hz. The PMS derived have two spectral dimensions: one dimension was the

frequency of power ('power frequency') and the other dimension was the 'modulation

frequency' of modulations of power. As the power was normalized to have a unit of dB, the

unit of PMS was also dB since the Fourier transform only includes linear operations.

Here, we laid out two further constraints due to the parameter selections in the above

analyses used to derive the PMS. First, the duration of power selected for the PMS analyses

was 1.5 s long (0.25 s to 1.75 s after noise onset), which meant that the lowest possible

modulation frequency representing more than two cycles of power fluctuations was 1.33 Hz.

At modulation frequencies less than 1.33 Hz, one could not determine whether the

modulation components represented by the frequencies are rhythmic or belong to a duty cycle

of an even lower frequency, or simply indicate a trending signal. Therefore, there is Trend

bound at 1.33 Hz along the axis of modulation frequency - neural components above 1.33 Hz

can be considered to have rhythmic modulation components; neural components below 1.33

Hz likely represent trending responses from 0.25 s to 1.75 s. Secondly, the power at one

power frequency can only fluctuate at a modulation frequency less than one half of the power

frequency according to the Nyquist-Shannon sampling theorem. The power at one power

frequency can be viewed as a signal carrier; the power modulation envelope of this carrier

can only have a modulation frequency as high as half of the carrier frequency. For example,

the modulation frequency derived from the power at the power frequency of 1 Hz can only be

as high as 0.5 Hz in theory. If we examine the modulation frequency of 2 Hz on EEG power,

we have to look at the EEG power at least above 4 Hz of the power frequency. Hence, there

is Sampling bound. We indicated those two constraints, Trend bound and Sampling bound, in

Fig. 2B.

Note that the PMS was not affected by different baseline correction procedures used

in *Time-frequency decomposition of EEG signals*. The reason is that we took a log

transformation of EEG power after the division between the power of the stimulus period and

the baseline power. This is equal to taking a log of the power of the signals and a log of the

baseline power separately and then subtracting the two values linearly. The baseline power

did not change in time and, after the Fourier transform was applied, was separated out from

the EEG power of the signals as a DC component in the modulation spectra (the component

at 0 Hz).

Surrogate test on PMS

To determine the spectral regions of PMS showing significant modulation strength, we

conducted a surrogate test in the spectral domain at the group level. The rationale is that, as

all the participants were doing the same task, neural dynamics reflecting the underlying

cognitive processes should be similar across individual participants. At the group level, we

should be able to identify neural dynamics that are consistent across the participants. Hence,

the null hypothesis here is that no consistent temporal modulation of power at any power

frequency exists across the participants.

To construct the null hypothesis, after we applied the Fourier transform on the EEG

power at each power frequency and obtained the Fourier coefficients (complex numbers), we

multiplied a complex number of a random phase and of a norm of 1 with the Fourier

coefficient at each modulation frequency. By doing this, we disrupted the phase of each

modulation frequency at each power frequency but preserved the modulation strength in each

participant's data – this is equal to disrupting temporal dynamics. We then averaged the

surrogated data over all the participants in the spectral domain and derived a new group-level

PMS. We conducted this procedure 1000 times and constructed a null distribution of PMS at

each power frequency and each modulation frequency, from which we derived a threshold of

a one-sided alpha level of 0.05.

The reason that we chose the surrogate test in the spectral domain but not in the

temporal domain is that we would like to specifically examine the temporal alignment of

modulations on the power across the participants while keeping the modulation strength

unchanged. To conduct a surrogate test in the temporal domain, we would have to change the

time range of individual participants' EEG data by shifting the time range of interest forward

or backward in each surrogate. The shifted time range may include noise onset/offset

responses or data from the fixation period, so both the modulation strength and phase of

power in the shifted time range are changed and are different across participants.

We selected four significant clusters (> 2 spectral bins) that were robustly shown in

each time of the surrogate test with high modulation power (Cluster 1, 2, 3, and 4 in Fig. 2B).

Other significant modulation bins were also observed, but they either had low modulation

strength in the PMS or fell below Sampling bound (likely caused by the ripple artefacts of the

Fourier transform). The power frequency ranges/the modulation frequency ranges for the four

clusters are as following: Cluster 1, 1-4 Hz/0-1.33 Hz; Cluster 2, 22-27 Hz/0-1.33 Hz; Cluster

3, 9-14 Hz/1.8-2.8 Hz; Cluster 4, 24-26 Hz/1.8-2.8 Hz.

The modulation frequency range for each cluster was not selected strictly according to

the significant spectral bins in the PMS of Fig. 2B. The significant modulation frequencies

across different power frequencies in one cluster were not the same, so it was implausible to

filter the power strictly according to the significant frequency bins when the temporal domain

of the power was investigated. Defining a 'square' region of interest in the PMS for each

cluster facilitated filtering procedures in the following analyses of extracting temporal

dynamics; simplistic filtering procedures avoided potential artefacts introduced by filtering

each power frequency using a different modulation frequency range. Therefore, we chose the

spectral bins for the clusters in the following manner: 1) the significant modulation

frequencies of Cluster 1 and 2 were below Trend bound and potentially included trending

responses that lasted longer than the EEG data period (1.5 s) included in the PMS analyses,

so we used 0 Hz as the low cutoff modulation frequency and set the high cutoff frequency as

1.33 Hz; 2) Cluster 3 and 4 were centered around 2 Hz of the modulation frequency and we

selected the same modulation frequency range for both the clusters, so that later we could

compare temporal dynamics of power within the same modulation frequency range.

The PMS analyses were first conducted at Cz to investigate whether our hypothesized

neural phenomena (Fig. 1D) could be observed. We later examined other EEG channels and

created a topography for each cluster to reveal distributions of modulation strength across

EEG channels. We filtered the power according to each defined cluster and conducted the

surrogate test described above on each EEG channel. We derived thresholds at one-sided

alpha level of 0.05 to determine whether the power modulation was significant for each

cluster on each EEG channel, which was indicated by the green dots in Fig. 2C. When we

plotted the topographies in Fig. 2C, we normalized the modulation strength by subtracting the

mean of the null distributions out of the empirical modulation strength so that the distribution

of the modulation strength across EEG channels was highlighted in the topographies.

Temporal dynamics of clusters and power coupling

The above analyses in the spectral domain did not reveal temporal details of the power

fluctuations, so we next investigated temporal dynamics of power within those four clusters.

We first averaged the power within the power frequency range of one ROI/cluster and then

filtered the averaged power using a band-pass filter (for Cluster 3 and 4, 1.8 – 2.8 Hz) or a

low-pass filter (for Cluster 1 and 2, < 1.33 Hz) (two-pass Butterworth filter of an order of 2

implemented in Fieldtrip).

We next checked whether each cluster was independent from or coupled with other

clusters. By doing this, we can validate previous findings on the coupling of neural signals

between the delta and beta bands in prospective timing paradigms and validate our data-

driven procedures here. Considering that the Fourier transform projects signals onto an

orthogonal space of sinusoid waves, a complex neural signal may be broken into several

clusters. For example, Cluster 3 and 4 are probably in a harmonic relationship: Cluster 4 is

likely the harmonic of Cluster 3 as the power frequency of Cluster 4 (24-26 Hz) is

approximately twice of the power frequency of Cluster 3 (9 – 14 Hz). If this is the case,

Cluster 3 and 4 should be coupled.

We calculated phase locking values (PLV) on the power across participants to

measure the coupling between clusters at each time point. We applied the Hilbert

transformation on the power of each cluster and then derived the phase value at each time

point by calculating angle of the Hilbert coefficient. The phase difference between each pair

of clusters for each participant was then used to calculate PLV across all the participants

using the following formula, in which **S** denotes the total number of participants, **s** the

participant number, **c** the cluster, **a** and **b** numbers of two different clusters, **t** the time point,

and ϕ the phase at each time point:

$$PLV_{c_{ab},t} = \left| \sum_{s=1}^{S} e^{(\phi_{s,t,c_a} - \phi_{s,t,c_b})*i} \right| / S$$

If two clusters are consistently coupled across participants, the phase differences

across all the participants at one time point should concentrate around a specific phase value

– a high PLV value (low circular variance across participants). Accordingly, we constructed a

null hypothesis to determine the significant time points of PLV values through a surrogate

test in the spectral domain. In each participant's data, we reset the phase of each cluster at

one time point randomly by multiplying a complex number of a random phase and a norm of

1 with the Hilbert coefficient. By doing this, we disrupted the phase series of each cluster

independently and hence the coupling strength between two clusters. We then calculated new

PLV values of each pair of clusters across all the participants. We conducted this procedure

1000 times and constructed a null distribution of PLV at each time point, from which we

derived a threshold of a one-sided alpha level of 0.05 for each cluster pair.

Merge Cluster 2, 3, and 4 into Cluster 234 and further analyses

From Fig. 2D, we observed that Cluster 1 slowly changed along the entire trials and showed a

global trajectory capturing the length of the noise piece, whereas the dynamics of Cluster 2, 3

and 4 seemingly did not correspond to the temporal structure in each trial. However, a peak

after stimulus onset was observed in Cluster 2 around the first temporal location, Cluster 3

and 4 were coupled and both showed peaks around the second and the third temporal

locations (Fig. 2D&S2C), and Cluster 2 and 4 are within the same EEG power frequency

band (Fig. 2B). Overall, the above observation gave us a clue that Cluster 2, 3 and 4 may

collectively capture the three temporal locations of the target tone. We tried averaging the

EEG power over Cluster 2, 3, and 4 and merged them into Cluster 234. Surprisingly, the

merged EEG power dynamics showed a salient neural signature marking the temporal

locations, even if only the white noise was presented. By plotting Cluster 1 and Cluster 234

together, it can be clearly seen that Cluster 1 in the delta band reflected the global trial

structure (the length of noise pieces) and that Cluster 234 captured the local temporal

locations of the target tone. As we aimed to find neural signatures tracking the temporal

locations and similar findings in the delta band, corresponding to Cluster 1, have been

reported before to encode event duration or fixed intervals (Assaf Breska & Leon Y Deouell,

2017; van Wassenhove & Lecoutre, 2015), we next only focused on Cluster 234.

The dynamics of Cluster 234 contains a downward linear trend and a rhythmic

component (Fig. 2). To single out the rhythmic component, we fitted a line to Cluster 234

from -1 s to 0 s and subtracted out the fitted line. The resulted neural power modulation,

Corrected cluster 234, clearly showed peaks after each temporal location.

Please note that we altered the range used to fit the baseline for Cluster 234, from -1 s

to -0.3 s and from -1 s to -0.2 s, in order to avoid the confounding factor that the power of the

baseline close to 0 s may contain stimulus-evoked signals post the onset of noise at 0 s.

However, the different baseline ranges did not yield significant differences, likely due to the

relatively stable trend of the baseline between -1 s and 0 s, as depicted in Fig. 2E.

Furthermore, there was no 'onset response' after the noise onset for Cluster 234 as illustrated

in Fig. 2E. The power dynamics remained 'stable' until after 0.5 s, when the first peak of

Cluster 234 emerged. In instances where onset responses occur immediately post-stimulus

onset, including neural signals close to 0 s in the baseline could be problematic, but this was

not the case for Cluster 234. Therefore, we assert that using the range from -1 s to 0 s to

derive a baseline for correcting Cluster 234 did not impact subsequent analyses.

We then individually fitted a Gaussian function to Corrected cluster 234 from each

tone onset position to 0.5 s afterwards. From the fitted Gaussian curves, we derived the peak

latency (the position of the mean in the fitted Gaussian curve) and the standard deviation

(SD). We averaged each neural peak of Corrected cluster 234 within one SD centered around

the peak latency (-0.5 SD + tone location to 0.5 SD + tone location) for each participant and

conducted a one-way rmANOVA with Temporal location as the main factor.

To visualize the neural tracking of the temporal locations across all EEG channels, we

calculated the modulation strength of Corrected cluster 234 from 0.4 s to 1.9 s at each

channel (SD over this period of signal) and plotted the topography of the SD. The time range

was chosen based on the first zero-crossing point after the noise onset and the time point after

the third peak showing the lowest power magnitude. The first zero-crossing point indicated

the rising of the EEG power above zero and the second time point the EEG power falling

towards zero. The SD over this time range represented the degree of EEG power fluctuations,

which can be seen as the amplitude summed over all the frequency components of Corrected

cluster 234 within this time range.

Block number on power modulation of Corrected cluster234

We next examined how learning from the first to the third block affected the sequential

temporal anticipation. The participants probably could not precisely anticipate the temporal

locations of the target tone in the first block, but gradually learned the exact temporal

locations in the following blocks. We expected to find that the magnitude of Corrected cluster

234 increases from the first block to the third block.

We extracted the power from the spectral regions defined in Cluster 2, 3, and 4 in the

data from each block and merged the three clusters into Cluster 234 in each block. As

described in the above analyses, the PMS calculation was not affected by differences of

baseline corrections, so the baseline differences between the blocks did not contribute to the

following analyses. To correct Cluster234 in each block, we adopted the fitted baseline from

the above analyses on the trials pooled across all three blocks, so that Corrected cluster 234 in

each block went through the same baseline correction procedure. Similarly, following the

above procedures, we fitted Gaussian curves to peaks after each temporal location in each

block (Fig. 3&S4) to extract peak magnitudes. We conducted a two-way rmANOVA on the

peak magnitudes with Temporal location and Block number as the main factors to examine

the learning effect (Fig. 3B).

Correlation between behavioral performance and Corrected cluster234

We correlated d-prime values with the peak magnitude of Corrected cluster 234 in each block

to examine whether the identified neural signature coding temporal locations in the no-tone

trials correlated with the tone detection performance in the tone trials (Fig. 3&S4). It is

possible that, since the power of Corrected cluster 234 was derived in the no-tone trials, the

power peaks may be caused by false alarms – the participants thought they heard a tone even

though there was no tone, so we then correlated the power peaks with the false alarm rates.

Quantify peak latency change of Corrected cluster 234 across blocks and temporal locations

The peak latencies of Corrected cluster 234 decreased from the first temporal location to the

third and were modulated by the block number (Fig. 2G&3A). We here verified this

observation quantitively. One issue here was that the peak latencies determined by the

Gaussian fits were derived from the group-averaged power; individual participant's data were

noisy so that the Gaussian fits to each participant's neural power were not accurate. This

prevented us from drawing conclusions by conducting parametric statistic tests over

participants (e.g. rmANOVA). To circumvent this issue, we resorted to group-level

permutation tests.

We first used a permutation test to examine one observation from Fig. 3A: the peak

latencies were different between blocks. We constructed a null hypothesis that, at each

temporal location, there is no peak latency difference between blocks. One reason for

constructing this null hypothesis first was that the null distribution constructed at each

temporal location could be viewed as a distribution that averages out the factor of the block

number. Later, by comparing three null distributions of peak latency in the three temporal

locations, we could determine whether the peak latencies of Corrected cluster 234 varied with

the temporal locations.

To construct the null hypothesis of the peak latency at each temporal location across

blocks, we permuted the block label in a way that each participant's data in each block were

randomly assigned with a new block number. We grouped the permuted data into three new

blocks at each temporal location and derived Corrected cluster 234 in each new block. We

fitted a Gaussian curve to each power peak to derive the peak latency by following the above

procedure. We conducted this permutation procedure 1000 times and derived nine null

distributions of peak latencies (3 temporal locations * 3 blocks). As the block labels were

shuffled and were not meaningful any more in the null distributions, we pooled the null

distributions over three blocks and had one null distribution of 3000 samples for each

temporal location. We derived thresholds at a two-sided alpha level of 0.05 (the 76th latency

and the 2925th latency) from each null distribution (Fig. 3D). The thresholds derived can be

viewed as significant boundaries – if a peak latency from one block of the empirical data falls

within the boundaries, the peak latency does not significantly differ from peak latencies of

the other blocks; if a peak latency is outside of those significant boundaries, it is then

significantly different from the other peak latencies. The boundaries, in a sense, represented

an interval that the peak latencies of the three blocks are uniform, statistically speaking, so

we called them as the 'uniform' bound and the permutation test as the 'uniform' test.

The null distributions of peak latency at each temporal location represented

distributions of possible peak latencies without considering the factor of the block number, as

each distribution was derived from random combinations of the participants' data drawn from

the three blocks. Therefore, by randomly selecting a sample from each null distribution and

examining how the peak latencies changed from the null distribution of the first temporal

location to the third, we could further derive a distribution of the peak latency change that

could be used to determine whether the peak latencies significantly varied with the temporal

locations. One simple way to quantify the latency change was to fit a straight line to the peak

latencies with the temporal location as the independent variable of values from 1 to 3 and the

latency as the dependent variable. If the slope is significantly negative, it can be concluded

that the peak latency decreases significantly with the temporal locations. Following this logic,

we fitted a line to peak latencies randomly selected from the three null distributions and

constructed a distribution of slope of 3000 samples. We derived a threshold at a two-sided

alpha level of 0.05 from the slope distribution and compared the thresholds with zero to

determine whether the peak latencies significantly shortened from the first temporal location

to the third (Fig. 3F).

Modulation spectrum analysis on inter-trial phase coherence (ITPC)

Besides the power measurement of neural signals, the anticipation for a future event may

cause neural phases to concentrate around a certain phase value, as suggested in previous

literature (Assaf Breska & Leon Y Deouell, 2017; Samaha & Postle, 2015). If this is the case,

we would expect that the neural phases in the no-tone trials were consistently locked to the

tagged temporal locations and hence the variance of neural phases across trials decreased

before or after the temporal locations. Accordingly, a 2-Hz component should be observed in

the inter-trial phase coherence (ITPC) that measures consistency of neural phases across

trials. Here, we conducted the modulation spectrum analysis on ITPC over the no-tone trials

pooled from the three blocks.

Note that ITPC also reflects routinely-defined 'evoked power', as it has been shown

that the evoked power reflects phase-locked temporal coherence between trials because the

averaging procedure was taken before time-frequency decomposition (Teng & Poeppel,

2020). However, this labeling of 'evoked power' is not entirely correct because the averaging

in time does not necessarily average out non-phase locked activity (Mazaheri, 2022). Here,

we directly analyzed inter-trial phase coherence to quantify the phase-locked neural

components across trials. In this study, we simply use 'power' to indicate our calculation of

power here and analyzed the phase domain directly to represent time-locked neural activity

that is arguably related to 'evoked power'.

The Fourier coefficients were derived in *Time-frequency decomposition of EEG*

signals. We derived the neural phase at each time-frequency bin by calculating angle of the

Fourier coefficient. The formula of ITPC calculation at a time–frequency bin is shown below

with *N* denoting the total number of the trials, *n* the trial number, *f* the frequency point, *t* the

time point, and *ϕ* the phase value at each time-frequency bin:

$$ITPC_{t,f} = |\sum_{n=1}^{N} e^{\phi_{n,t,f} * i}|/N$$

We used the valid trials across the three blocks after *the EEG preprocessing* to

calculate ITPC. We selected the time range between 0.25 s and 1.75 after the noise onset to

avoid effects of onset and offset responses of the white noise. Then, following the procedures

in *Power modulation spectrum (PMS)*, we conducted modulation spectrum analyses as well

as surrogate tests on ITPC values to determine significant modulation clusters. Only one

salient cluster was shown below Trending bound and, in the theta and delta bands along the

axis of ITPC frequency (Fig. S3). Since no 2-Hz modulation component was found in the

ITPC values, showing that the phase-locked neural responses did not code the temporal

locations, we did not conduct any further analyses on the neural phases.

Train recurrent neural network models to accomplish the tone-detection task

We chose to train continuous-time recurrent neural networks (CRNNs) with mini-batch

gradient descent learned by backpropagation. The structure of CRNNs is defined as the

following equation:

$$\tau \frac{dx_i}{dt} = -x_i + \sum_{j=1}^{N} W_{ij}^{rec} r_j + \sum_{k=1}^{N_{in}} W_{ik}^{in} u_k + \xi_i$$

$$r_i = [x_i]_+$$

$$z_l = \sum_{i=1}^{N} W_{li}^{out} r_{i,T}$$

The number of hidden units, N, was 100. We tested CRNNs with fewer hidden units

(64 and 32) and found out that those CRNNs could not be trained to achieve human-level

performance when the external noise level was high in our settings. The time step size, *Δt*,

was 25 ms (a sampling rate of 40 Hz), which was sufficient to capture the dynamics at 2 Hz

shown in our neural findings and made the training efficiently. τ is the time constant of the

network units and was set to 100 ms. u is the input vector, so k can only be 1. W^{in} is

a $1 \times N$ matrix of connection weights from the inputs to N network units. W^{rec} is

an $N \times N$ matrix of recurrent connection weights between network units. ξ are N independent

Gaussian white noise processes with zero mean and unit variance that represent noise

intrinsic to the network. The rectified linear activation function (ReLU) was applied on

hidden units. W^{out} is an $N \times 2$ matrix of connection weights from N network units to the

outputs (two choices: yes or no). The outputs, z, were generated after the input is over; the

ending time point is indicated by T. The CRNN, a continuous-time formulation of a standard

RNN, was converted to a discrete-time version through the Euler method for implementation.

The connections between hidden units were constraint according to Dale's rule: 80%

of units were fixed to be excitatory units and 20% of units were inhibitory units (H Francis

Song et al., 2016). We initialized the weights by generating random weights using the normal

distribution with the mean of zero and the standard deviation (SD) of square root of 5, taking

absolute values of all weights, multiplying 20% of weights with -1 to set them as inhibitory

units, and equalizing the gain between excitatory units and inhibitory units. The diagonal of

the weight matrix (self-connections of units) was fixed at 0. The internal noise level of ξ was

set to 0.01, so that we could focus on studying the effect of noise level of inputs by varying

the noise level of inputs systemically.

The inputs fed to CRNNs were constructed in NeuroGym

(https://neurogym.github.io/) to mimic the trial structure of the experiment (Fig. 1A&4A).

The magnitude of the fixation period in each trial was set as zero with additive gaussian noise

of the mean of zero and SD of 0.05. The length of the fixation period varied from 1000 ms to

1500 ms in different batches during training in a step size of 25 ms. For example, in one

batch, the fixation period was 1000 ms; in another batch, the fixation period became 1175

ms. The stimulus period was 2 s and the noise was gaussian noise with the mean of 1 and

various levels of SD according to the following training designs. The target magnitude was

set as 1 and was added to the stimulus period at either 500 ms, 1000 ms, or 1500 ms. The

CRNNs made the decision by the end of the stimulus period to answer whether there was a

target in a trial – a yes-no response. We did not simulate the reaction time but only aimed to

simulate the procedure of human participants detecting a tone and indicating their decision by

pressing a button as soon as the white noise was over. The loss was calculated only by the

end of trials using cross entropy.

During training CRNNs, each mini-batch included 16 trials with each trial equally

likely coming from each of four conditions (three tone trial conditions and one no-tone trial

condition) (Fig. 4A). We set the maximum training sessions to 40000 and terminated the

training if the mean accuracy over the last 100 mini batches was above 99 percentage correct.

We used ADAM (Kingma & Ba, 2014) during training and scheduled the learning rate to

drop in three steps: 0.001 for 10000 sessions, then 0.0001 for 20000 following sessions, and

0.00001 for the last 10000 sessions. All the training procedures were implemented in

PyTorch 1.7.0 (Paszke et al., 2019).

The main parameter we varied during training and testing CRNNs was the external

noise level of the stimulus period. The external noise was added to the stimuli and was shared

across all the hidden units of CRNNs. We trained five CRNNs at five external noise levels,

separately: 0.01, 0.05, 0.1, 0.15, and 0.2 SD, and tested each trained CRNN at 7 noise levels:

0.01, 0.05, 0.1, 0.15, 0.2, 0.25, and 0.3 SD.

CRNN behavioral performance, hidden state dynamics, and modulation spectra

We trained five CRNNs at different noise levels, respectively, and tested them across all the 7

testing noise levels. We generated 27 sets of behavioral data at each testing noise level to

match the number of recruited human participants. Following the same procedure in

Behavioral data analysis, we calculated d-prime values from CRNN outputs. Then, we

selected one testing noise level for each of the five CRNNs according to how close the group-

level d-prime value derived at this testing noise level matched the group-level mean of the

human behavioral data averaged over the three temporal locations in Fig. S1C.

We fed each of the five trained CRNNs with 40 no-tone trials at the chosen testing

noise level and extracted activities of their hidden units. As neural signals recorded by EEG

reflect summarized neural activities over both excitatory and inhibitory synapses, similarly,

we averaged the CRNN hidden state activities over both excitatory and inhibitory units. We

analyzed the averaged hidden state activities following the procedure used in calculating the

modulation spectrum of EEG signals. We selected the averaged hidden activity of each trial

from 0.25 s to 1.75 s after noise onset and applied the Fourier transform, and then extracted

the amplitude of the Fourier coefficients for each trial and averaged the modulation spectra

over all the 40 no-tone trials to generate an averaged modulation spectrum of the hidden

activities. We found that the CRNN trained at the noise level of 0.15 SD and tested at the

noise level of 0.2 SD showed the largest 2 Hz modulation strength (Fig. 4C), so we selected

this CRNN for the subsequent investigations as it both matched the human behavioral

performance and the neural findings (Fig. 2). The hidden activities of CRNNs shown in Fig.

4D and Fig. 4E were derived from averaging hidden activities over 40 no-tone trials.

Train and test CRNNs with a uniform probabilistic distribution of tone position from 0.25 s to

1.75 s

We trained a CRNN with a new trial structure in which the target tone can appear any time

from 0.25 s to 1.75 s with an equal probability (Fig. 4F), so that the newly trained CRNN can

be compared with the CRNNs trained with a target tone appearing at one of the three

temporal locations. The motivation here is twofold: 1) If the 2 Hz modulation component was

driven by the specific temporal locations, the CRNN trained with the new trial structure

should not contain the 2 Hz modulation component; 2) We aimed to test tone detection

performance of the CRNN trained with the three temporal locations at other time points.

In the original trial structure for training CRNNs (Fig. 4A), 75 percent of trials had

the target tone imbedded and the other 25 percent of trials were no-tone trials. Accordingly,

here, we fixed 25 percent of trials to be no-tone trials, and, in the other 75 percent of trials, a

target randomly appeared any time from 0.25 s to 1.75 s. In each training batch, the trial type

was randomly sampled with a probability of 0.75 to be tone trials and 0.25 to be no-tone

trials. We chose the training noise level to be 0.15 SD. The other parameters and procedures

for training CRNNs were the same as in the previous CRNN training procedures. After 40000

batches, the training accuracy reached 0.97.

We tested both the newly trained CRNN and the CRNN trained at 0.15 SD with three

temporal locations at a testing noise level of 0.2 SD. We chose a different way to analyze the

tone detection performance, in contrast with the above behavioral analyses (Fig. 4B). We

independently generated 40 no-tone trials and 40 tone trials with the target at one time point,

and asked CRNNs to generate outputs. For each time point, we measured a hit rate from the

tone trials and a false alarm rate from the no-tone trials and calculated a d-prime value. This

d-prime calculation was conducted from 0.25 s to 1.75 s with a step of 0.025 s (Fig. 4G).

Perturb E/I ratio of the CRNN trained at the noise level of 0.15 SD

To account for the neural power precession observed in Fig. 3D, we perturbed excitation-

inhibition balance of trained CRNNs. The perturbed CRNNs may generate new hidden

dynamics that potentially capture the phenomenon of the shortened peak latencies of

Corrected cluster 234.

We conducted the perturbation on the CRNN trained at the noise level of 0.15 SD

with three temporal locations and derived the hidden activities at the testing noise level of 0.2

SD, as the above analyses showed that this CRNN captured the neural dynamics successfully

(Fig. 4C-E). To perturb the excitation-inhibition balance, we changed the weights of

excitatory and inhibitory units. For example, we increased the weights of excitatory units by

10 percent so that the excitatory units sent stronger projections to all units. However, only

increasing excitatory unit weights would disturb the CRNN and potentially drive it to a

chaotic state or to a silent state. Hence, we changed the weights of excitatory and inhibitory

units by the same amount but in opposite directions. For example, when the perturbation was

10 percent, we increased the excitatory weights by 10 percent and, correspondingly,

decreased the inhibitory weights by 10 percent.

We perturbed the excitation-inhibition balance from -40 percent (negative gain for excitatory

weights and positive gain for inhibitory weights) to 40 percent with a step of 5 percent. After

we perturbed CRNNs, we asked the CRNNs to generate behavioral outputs and calculated d-

prime values. Meaningful positive d-prime values were generated when the perturbations

were -10, -5, 5, 10, and 15 percent, so we only analyzed the CRNNs under those five

perturbations. We derived the averaged hidden activities of the perturbed CRNNs from 40

no-tone trials at a noise level of 0.2 SD (Fig. 4H) and calculated d-prime values at three

temporal locations from 27 simulated datasets following the procedure used in analyzing

human behavioral data (Fig. 4I).

1089

References

1090

Arnal, L. H., Doelling, K. B., & Poeppel, D. (2015). Delta-Beta Coupled Oscillations
Underlie Temporal Prediction Accuracy. *Cereb Cortex*, *25*(9), 3077-3085.
https://doi.org/10.1093/cercor/bhu103
Arnal, L. H., & Giraud, A. L. (2012). Cortical oscillations and sensory predictions. *Trends*
Cogn Sci, *16*(7), 390-398. https://doi.org/10.1016/j.tics.2012.05.003
Baayen, R. H., & Milin, P. (2010). Analyzing reaction times. *International journal of*
psychological research, *3*(2), 12-28.
Bauer, M., Kluge, C., Bach, D., Bradbury, D., Heinze, H. J., Dolan, R. J., & Driver, J. (2012).
Cholinergic enhancement of visual attention and neural oscillations in the human
brain. *Curr Biol*, *22*(5), 397-402. https://doi.org/10.1016/j.cub.2012.01.022
Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and
powerful approach to multiple testing. *Journal of the Royal statistical society: series*
B (Methodological), *57*(1), 289-300.
Bolger, D., Coull, J. T., & Schön, D. (2014). Metrical rhythm implicitly orients attention in
time as indexed by improved target detection and left inferior parietal activation.
Journal of cognitive neuroscience, *26*(3), 593-605.
Breska, A., & Deouell, L. Y. (2017). Neural mechanisms of rhythm-based temporal
prediction: Delta phase-locking reflects temporal predictability but not rhythmic
entrainment. *PLoS Biol*, *15*(2), e2001665.
https://doi.org/10.1371/journal.pbio.2001665
Breska, A., & Deouell, L. Y. (2017). Neural mechanisms of rhythm-based temporal
prediction: Delta phase-locking reflects temporal predictability but not rhythmic
entrainment. *PLoS biology*, *15*(2), e2001665.
Breska, A., & Ivry, R. B. (2020). Context-specific control over the neural dynamics of
temporal attention by the human cerebellum. *Sci Adv*, *6*(49).
https://doi.org/10.1126/sciadv.abb1141
Brunel, N., & Wang, X.-J. (2003). What determines the frequency of fast network
oscillations with irregular neural discharges? I. Synaptic dynamics and excitation-
inhibition balance. *Journal of neurophysiology*, *90*(1), 415-430.
Coull, J. T., Cotti, J., & Vidal, F. (2016). Differential roles for parietal and frontal cortices in
fixed versus evolving temporal expectations: Dissociating prior from posterior
temporal probabilities with fMRI. *Neuroimage*, *141*, 40-51.
https://doi.org/10.1016/j.neuroimage.2016.07.036
Coull, J. T., & Nobre, A. C. (1998). Where and when to pay attention: the neural systems for
directing attention to spatial locations and to time intervals as revealed by both PET
and fMRI. *Journal of Neuroscience*, *18*(18), 7426-7435.
Cravo, A. M., Rohenkohl, G., Santos, K. M., & Nobre, A. C. (2017). Temporal Anticipation
Based on Memory. *J Cogn Neurosci*, *29*(12), 2081-2089.
https://doi.org/10.1162/jocn_a_01172
Dale, H. (1935). Pharmacology and Nerve-endings (Walter Ernest Dixon Memorial Lecture):
(Section of Therapeutics and Pharmacology). *Proc R Soc Med*, *28*(3), 319-332.
https://www.ncbi.nlm.nih.gov/pubmed/19990108
Doelling, K. B., Assaneo, M. F., Bevilacqua, D., Pesaran, B., & Poeppel, D. (2019). An
oscillator model better predicts cortical entrainment to music. *Proc Natl Acad Sci U S*
A, *116*(20), 10113-10121. https://doi.org/10.1073/pnas.1816414116
Doelling, K. B., & Poeppel, D. (2015). Cortical entrainment to music and its modulation by
expertise. *Proc Natl Acad Sci U S A*, *112*(45), E6233-6242.
https://doi.org/10.1073/pnas.1508431112

Elliott, T. M., & Theunissen, F. E. (2009). The modulation transfer function for speech intelligibility. *PLoS Comput Biol*, *5*(3), e1000302. https://doi.org/10.1371/journal.pcbi.1000302

Faisal, A. A., Selen, L. P., & Wolpert, D. M. (2008). Noise in the nervous system. *Nat Rev Neurosci*, *9*(4), 292-303. https://doi.org/10.1038/nrn2258

Friston, K. (2005). A theory of cortical responses. *Philos Trans R Soc Lond B Biol Sci*, *360*(1456), 815-836. https://doi.org/10.1098/rstb.2005.1622

Grabenhorst, M., Maloney, L. T., Poeppel, D., & Michalareas, G. (2021). Two sources of uncertainty independently modulate temporal expectancy. *Proc Natl Acad Sci U S A*, *118*(16). https://doi.org/10.1073/pnas.2019342118

Grabenhorst, M., Michalareas, G., Maloney, L. T., & Poeppel, D. (2019). The anticipation of events in time. *Nat Commun*, *10*(1), 5802. https://doi.org/10.1038/s41467-019-13849-0

Haegens, S., Luther, L., & Jensen, O. (2012). Somatosensory anticipatory alpha activity increases to suppress distracting input. *J Cogn Neurosci*, *24*(3), 677-685. https://doi.org/10.1162/jocn_a_00164

Haegens, S., & Zion Golumbic, E. (2018). Rhythmic facilitation of sensory processing: A critical review. *Neurosci Biobehav Rev*, *86*, 150-165. https://doi.org/10.1016/j.neubiorev.2017.12.002

Hautus, M. J., Macmillan, N. A., & Creelman, C. D. (2021). *Detection theory: A user's guide*. Routledge.

Heilbron, M., & Chait, M. (2018). Great expectations: is there evidence for predictive coding in auditory cortex? *Neuroscience*, *389*, 54-73.

Henry, M. J., Herrmann, B., & Obleser, J. (2014). Entrained neural oscillations in multiple frequency bands comodulate behavior. *Proc Natl Acad Sci U S A*, *111*(41), 14935-14940. https://doi.org/10.1073/pnas.1408741111

Henry, M. J., & Obleser, J. (2012). Frequency modulation entrains slow neural oscillations and optimizes human listening behavior. *Proc Natl Acad Sci U S A*, *109*(49), 20095-20100. https://doi.org/10.1073/pnas.1213390109

Herbst, S. K., Fiedler, L., & Obleser, J. (2018). Tracking Temporal Hazard in the Human Electroencephalogram Using a Forward Encoding Model. *eNeuro*, *5*(2). https://doi.org/10.1523/ENEURO.0017-18.2018

Herrmann, B., Henry, M. J., Haegens, S., & Obleser, J. (2016). Temporal expectations and neural amplitude fluctuations in auditory cortex interactively influence perception. *Neuroimage*, *124*(Pt A), 487-497. https://doi.org/10.1016/j.neuroimage.2015.09.019

Hillyard, S. A., Vogel, E. K., & Luck, S. J. (1998). Sensory gain control (amplification) as a mechanism of selective attention: electrophysiological and neuroimaging evidence. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *353*(1373), 1257-1270.

Janssen, P., & Shadlen, M. N. (2005). A representation of the hazard rate of elapsed time in macaque area LIP. *Nat Neurosci*, *8*(2), 234-241. https://doi.org/10.1038/nn1386

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: gating by inhibition. *Front Hum Neurosci*, *4*, 186. https://doi.org/10.3389/fnhum.2010.00186

Jones, M. R., Kidd, G., & Wetzel, R. (1981). Evidence for rhythmic attention. *J Exp Psychol Hum Percept Perform*, *7*(5), 1059-1073. https://doi.org/10.1037//0096-1523.7.5.1059

Kevric, J., & Subasi, A. (2017). Comparison of signal decomposition methods in classification of EEG signals for motor-imagery BCI system. *Biomedical Signal Processing and Control*, *31*, 398-406.

Kingma, D. P., & Ba, J. (2014). Adam: A method for stochastic optimization. *arXiv preprint*
arXiv:1412.6980.
Klimesch, W., Sauseng, P., & Hanslmayr, S. (2007). EEG alpha oscillations: the inhibition-
timing hypothesis. *Brain Res Rev*, *53*(1), 63-88.
https://doi.org/10.1016/j.brainresrev.2006.06.003
Kontsevich, L. L., & Tyler, C. W. (1999). Bayesian adaptive estimation of psychometric
slope and threshold. *Vision Res*, *39*(16), 2729-2737. https://doi.org/10.1016/s0042-
6989(98)00285-5
Lakatos, P., Barczak, A., Neymotin, S. A., McGinnis, T., Ross, D., Javitt, D. C., &
O'Connell, M. N. (2016). Global dynamics of selective attention and its lapses in
primary auditory cortex. *Nat Neurosci*, *19*(12), 1707-1717.
https://doi.org/10.1038/nn.4386
Makeig, S., Debener, S., Onton, J., & Delorme, A. (2004). Mining event-related brain
dynamics. *Trends Cogn Sci*, *8*(5), 204-210. https://doi.org/10.1016/j.tics.2004.03.008
Makeig, S., Debener, S., Onton, J., & Delorme, A. (2004). Mining event-related brain
dynamics. *Trends in cognitive sciences*, *8*(5), 204-210.
Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-and MEG-data.
Journal of neuroscience methods, *164*(1), 177-190.
Martens, S., & Johnson, A. (2005). Timing attention: cuing target onset interval attenuates
the attentional blink. *Mem Cognit*, *33*(2), 234-240.
https://doi.org/10.3758/bf03195312
Mattiesing, R. M., Kruijne, W., Meeter, M., & Los, S. A. (2017). Timing a week later: The
role of long-term memory in temporal preparation. *Psychon Bull Rev*, *24*(6), 1900-
1905. https://doi.org/10.3758/s13423-017-1270-3
Mazaheri, A. (2022). THE RELATIONSHIP BETWEEN EVOKED AND INDUCED
EEG/MEG CHANGES. *The Oxford Handbook of EEG Frequency*, 115.
McClure, S. M., Berns, G. S., & Montague, P. R. (2003). Temporal prediction errors in a
passive learning task activate human striatum. *Neuron*, *38*(2), 339-346.
https://doi.org/10.1016/s0896-6273(03)00154-5
Mello, G. B., Soares, S., & Paton, J. J. (2015). A scalable population code for time in the
striatum. *Curr Biol*, *25*(9), 1113-1122. https://doi.org/10.1016/j.cub.2015.02.036
Merchant, H., & Yarrow, K. (2016). How the motor system both encodes and influences our
sense of time. *Current Opinion in Behavioral Sciences*, *8*, 22-27.
Morillon, B., & Baillet, S. (2017). Motor origin of temporal predictions in auditory attention.
Proceedings of the National Academy of Sciences, *114*(42), E8913-E8921.
Morillon, B., Schroeder, C. E., & Wyart, V. (2014). Motor contributions to the temporal
precision of auditory attention. *Nat Commun*, *5*, 5255.
https://doi.org/10.1038/ncomms6255
Morillon, B., Schroeder, C. E., Wyart, V., & Arnal, L. H. (2016). Temporal Prediction in lieu
of Periodic Stimulation. *J Neurosci*, *36*(8), 2342-2347.
https://doi.org/10.1523/JNEUROSCI.0836-15.2016
Motanis, H., & Buonomano, D. V. (2015). Neural coding: time contraction and dilation in the
striatum. *Curr Biol*, *25*(9), R374-376. https://doi.org/10.1016/j.cub.2015.02.057
Naatanen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity
(MMN) in basic research of central auditory processing: a review. *Clin Neurophysiol*,
118(12), 2544-2590. https://doi.org/10.1016/j.clinph.2007.04.026
Ng, B. S., Schroeder, T., & Kayser, C. (2012). A precluding but not ensuring role of
entrained low-frequency oscillations for auditory perception. *J Neurosci*, *32*(35),
12268-12276. https://doi.org/10.1523/JNEUROSCI.1877-12.2012

Niemi, P., & Näätänen, R. (1981). Foreperiod and simple reaction time. *Psychological*
bulletin, *89*(1), 133.
Nobre, A., Correa, A., & Coull, J. (2007). The hazards of time. *Curr Opin Neurobiol*, *17*(4),
465-470. https://doi.org/10.1016/j.conb.2007.07.006
Nobre, A. C. (2001). Orienting attention to instants in time. *Neuropsychologia*, *39*(12), 1317-
1328. https://doi.org/10.1016/s0028-3932(01)00120-8
Nobre, A. C., & Stokes, M. G. (2019). Premembering Experience: A Hierarchy of Time-
Scales for Proactive Attention. *Neuron*, *104*(1), 132-146.
https://doi.org/10.1016/j.neuron.2019.08.030
Nobre, A. C., & van Ede, F. (2018). Anticipated moments: temporal structure in attention.
Nat Rev Neurosci, *19*(1), 34-48. https://doi.org/10.1038/nrn.2017.141
Nozaradan, S., Schwartze, M., Obermeier, C., & Kotz, S. A. (2017). Specific contributions of
basal ganglia and cerebellum to the neural tracking of rhythm. *Cortex*, *95*, 156-168.
https://doi.org/10.1016/j.cortex.2017.08.015
Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open Source
Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological
Data. *Computational Intelligence and Neuroscience*, *2011*. https://doi.org/Artn
156869
10.1155/2011/156869
Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z.,
Gimelshein, N., & Antiga, L. (2019). Pytorch: An imperative style, high-performance
deep learning library. *Advances in neural information processing systems*, *32*.
Paton, J. J., & Buonomano, D. V. (2018). The Neural Basis of Timing: Distributed
Mechanisms for Diverse Functions. *Neuron*, *98*(4), 687-705.
https://doi.org/10.1016/j.neuron.2018.03.045
Prins, N. (2013). The psi-marginal adaptive method: How to give nuisance parameters the
attention they deserve (no more, no less). *Journal of Vision*, *13*(7).
https://doi.org/Artn 3
10.1167/13.7.3
Prins, N. (2016). *Psychophysics: a practical introduction*. Academic Press.
Reynolds, J. H., & Chelazzi, L. (2004). Attentional modulation of visual processing. *Annu*
Rev Neurosci, *27*, 611-647. https://doi.org/10.1146/annurev.neuro.26.041002.131039
Rimmele, J., Jolsvai, H., & Sussman, E. (2011). Auditory target detection is affected by
implicit temporal and spatial expectations. *Journal of cognitive neuroscience*, *23*(5),
1136-1147.
Rimmele, J. M., Morillon, B., Poeppel, D., & Arnal, L. H. (2018). Proactive Sensing of
Periodic and Aperiodic Auditory Patterns. *Trends Cogn Sci*, *22*(10), 870-882.
https://doi.org/10.1016/j.tics.2018.08.003
Salinas, E., & Sejnowski, T. J. (2001). Correlated neuronal activity and the flow of neural
information. *Nat Rev Neurosci*, *2*(8), 539-550. https://doi.org/10.1038/35086012
Samaha, J., Bauer, P., Cimaroli, S., & Postle, B. R. (2015). Top-down control of the phase of
alpha-band oscillations as a mechanism for temporal prediction. *Proceedings of the*
National Academy of Sciences, *112*(27), 8439-8444.
Samaha, J., & Postle, B. R. (2015). The speed of alpha-band oscillations predicts the
temporal resolution of visual perception. *Current Biology*, *25*(22), 2985-2990.
Schubotz, R. I. (2007). Prediction of external events with our motor system: towards a new
framework. *Trends in cognitive sciences*, *11*(5), 211-218.
Sherman, M. A., Lee, S., Law, R., Haegens, S., Thorn, C. A., Hamalainen, M. S., Moore, C.
I., & Jones, S. R. (2016). Neural mechanisms of transient neocortical beta rhythms:
Converging evidence from humans, computational modeling, monkeys, and mice.

Proc Natl Acad Sci U S A, *113*(33), E4885-4894.
https://doi.org/10.1073/pnas.1604135113

Song, H. F., Yang, G. R., & Wang, X.-J. (2016). Training excitatory-inhibitory recurrent neural networks for cognitive tasks: a simple and flexible framework. *PLoS computational biology*, *12*(2), e1004792.

Song, H. F., Yang, G. R., & Wang, X. J. (2016). Training Excitatory-Inhibitory Recurrent Neural Networks for Cognitive Tasks: A Simple and Flexible Framework. *PLoS Comput Biol*, *12*(2), e1004792. https://doi.org/10.1371/journal.pcbi.1004792

Spaak, E., de Lange, F. P., & Jensen, O. (2014). Local entrainment of alpha oscillations by visual stimuli causes cyclic modulation of perception. *J Neurosci*, *34*(10), 3536-3544. https://doi.org/10.1523/JNEUROSCI.4385-13.2014

Spitzer, B., & Haegens, S. (2017). Beyond the status quo: a role for beta oscillations in endogenous content (re) activation. *eNeuro*, *4*(4).

Strauß, A., Wöstmann, M., & Obleser, J. (2014). Cortical alpha oscillations as a tool for auditory selective inhibition. *Frontiers in human neuroscience*, *8*, 350.

Tal, I., Large, E. W., Rabinovitch, E., Wei, Y., Schroeder, C. E., Poeppel, D., & Zion Golumbic, E. (2017). Neural Entrainment to the Beat: The "Missing-Pulse" Phenomenon. *J Neurosci*, *37*(26), 6331-6341. https://doi.org/10.1523/JNEUROSCI.2500-16.2017

Tavano, A., Schroger, E., & Kotz, S. A. (2019). Beta power encodes contextual estimates of temporal event probability in the human brain. *PLoS One*, *14*(9), e0222420. https://doi.org/10.1371/journal.pone.0222420

Ten Oever, S., Schroeder, C. E., Poeppel, D., van Atteveldt, N., Mehta, A. D., Megevand, P., Groppe, D. M., & Zion-Golumbic, E. (2017). Low-Frequency Cortical Oscillations Entrain to Subthreshold Rhythmic Auditory Stimuli. *J Neurosci*, *37*(19), 4903-4912. https://doi.org/10.1523/JNEUROSCI.3658-16.2017

Teng, X., & Poeppel, D. (2020). Theta and Gamma Bands Encode Acoustic Dynamics over Wide-Ranging Timescales. *Cereb Cortex*, *30*(4), 2600-2614. https://doi.org/10.1093/cercor/bhz263

van Bree, S., Sohoglu, E., Davis, M. H., & Zoefel, B. (2021). Sustained neural rhythms reveal endogenous oscillations supporting speech perception. *PLoS Biol*, *19*(2), e3001142. https://doi.org/10.1371/journal.pbio.3001142

Van Diepen, R. M., & Mazaheri, A. (2018). The caveats of observing inter-trial phase-coherence in cognitive neuroscience. *Scientific Reports*, *8*(1), 2990.

van Ede, F., Chekroud, S. R., Stokes, M. G., & Nobre, A. C. (2018). Decoding the influence of anticipatory states on visual perception in the presence of temporal distractors. *Nat Commun*, *9*(1), 1449. https://doi.org/10.1038/s41467-018-03960-z

van Wassenhove, V., & Lecoutre, L. (2015). Duration estimation entails predicting when. *Neuroimage*, *106*, 272-283.

Wilsch, A., Henry, M. J., Herrmann, B., Maess, B., & Obleser, J. (2015). Slow-delta phase concentration marks improved temporal expectations based on the passage of time. *Psychophysiology*, *52*(7), 910-918. https://doi.org/10.1111/psyp.12413

Wittmann, M. (2013). The inner sense of time: how the brain creates a representation of duration. *Nature Reviews Neuroscience*, *14*(3), 217-223.

Wostmann, M., Alavash, M., & Obleser, J. (2019). Alpha Oscillations in the Human Brain Implement Distractor Suppression Independent of Target Selection. *J Neurosci*, *39*(49), 9797-9805. https://doi.org/10.1523/JNEUROSCI.1954-19.2019

Wöstmann, M., Lim, S.-J., & Obleser, J. (2017). The human neural alpha response to speech is a proxy of attentional control. *Cerebral cortex*, *27*(6), 3307-3317.

Yamins, D. L., & DiCarlo, J. J. (2016). Using goal-driven deep learning models to
understand sensory cortex. *Nat Neurosci*, *19*(3), 356-365.
https://doi.org/10.1038/nn.4244
Yamins, D. L., Hong, H., Cadieu, C. F., Solomon, E. A., Seibert, D., & DiCarlo, J. J. (2014).
Performance-optimized hierarchical models predict neural responses in higher visual
cortex. *Proceedings of the National Academy of Sciences*, *111*(23), 8619-8624.
Yamins, D. L., Hong, H., Cadieu, C. F., Solomon, E. A., Seibert, D., & DiCarlo, J. J. (2014).
Performance-optimized hierarchical models predict neural responses in higher visual
cortex. *Proc Natl Acad Sci U S A*, *111*(23), 8619-8624.
https://doi.org/10.1073/pnas.1403112111
Yang, G. R., Joglekar, M. R., Song, H. F., Newsome, W. T., & Wang, X.-J. (2019). Task
representations in neural networks trained to perform many cognitive tasks. *Nature
Neuroscience*, *22*(2), 297-306.
Yang, G. R., & Wang, X.-J. (2020). Artificial neural networks for neuroscientists: a primer.
Neuron, *107*(6), 1048-1070.
Yekutieli, D., & Benjamini, Y. (1999). Resampling-based false discovery rate controlling
multiple test procedures for correlated test statistics. *Journal of Statistical Planning
and Inference*, *82*(1-2), 171-196.
Zanto, T. P., Snyder, J. S., & Large, E. W. (2006). Neural correlates of rhythmic expectancy.
Advances in cognitive psychology, *2*(2), 221.
Zumer, J. M., Scheeringa, R., Schoffelen, J. M., Norris, D. G., & Jensen, O. (2014). Occipital
alpha activity during stimulus processing gates the information flow to object-
selective cortex. *PLoS Biol*, *12*(10), e1001965.
https://doi.org/10.1371/journal.pbio.1001965

Supplementary Information

Sequential Temporal Anticipation Characterized by Neural Power Modulation and in Recurrent Neural Networks

Xiangbin Teng and Ru-Yuan Zhang

Corresponding Author: Xiangbin Teng, xiangbinteng@cuhk.edu.hk

This file includes:

Figure S1 – S4

Supplementary Figures

Fig. S1. Tone Detection Threshold, Condition Probabilities, and ERP Results. Refer to Figures 1 & 2 for related information.

(A) Tone detection thresholds for individual participants. Each dot signifies a participant's threshold.

(B) Condition probabilities. The left plot illustrates the cumulative probability of the target tone appearing in tone trials over time; the right plot displays the probability of a trial being a tone trial, given that no tone is presented.

(C) Behavioral results. From left to right, the plots show d-prime values, criteria, and reaction time, respectively. Line color denotes block number. Error bars indicate ±1 standard error of the mean across participants. Analysis was conducted on tone detection performance using a signal detection framework and a two-way repeated-measure ANOVA (rmANOVA) on the d-prime values with Temporal location and Block number as main factors. A significant main effect was found for Block number ($F(2,34) = 5.74$, $p = .007$, $\eta_p^2 = 0.252$) with a significant linear trend indicating decreased tone detection performance with increasing block number ($F(1,17) = 9.94$, $p = .006$, $\eta_p^2 = 0.369$). This effect is attributed to participants' decreased effort or fatigue due to the

repetitive trial structure in the tone detection task. Temporal location did not have a significant main effect $(F_{(2,34)} = 3.26, p = .051, \eta_p^2 = 0.166)$, but a significant interaction was found between Temporal location and Block number $(F_{(4,68)} = 3.80, p = .008, \eta_p^2 = 0.183)$. Further testing of Temporal location in each block (one-way rmANOVA) revealed a significant main effect in the second $(F_{(2,34)} = 4.97, p = .020, \eta_p^2 = 0.226)$ and third blocks $(F_{(2,34)} = 7.01, p = .009, \eta_p^2 = 0.292)$, but not in the first $(F_{(2,34)} = 2.94, p = .066, \eta_p^2 = 0.148)$. False Discovery Rate (FDR) adjustment was applied to control false positive rate.

The criteria calculated for the three temporal locations shared the same no-tone trials and the differences between them were dominated by the hit rate in the tone trials. We averaged the hit rates over the three temporal locations in the tone trials and calculated a criterion for each block. No significant main effect was found in a one-way rmANOVA on criteria with Block number as the main factor $(F_{(2,34)} = 3.06, p = .056, \eta_p^2 = 0.105)$.

A two-way rmANOVA was conducted on the raw reaction time with Trial condition (three different temporal locations in the tone trials and one no-tone condition) and Block number as main factors. The main effect of Trial condition was significant $(F_{(3,51)} = 28.68, p < .001, \eta_p^2 = 0.636)$; no significance was found for the main effect of Block number $(F_{(2,34)} = 1.65, p = .207, \eta_p^2 = 0.089)$ nor for the interaction $(F_{(6,102)} = .92, p = .485, \eta_p^2 = 0.051)$. The significant main effect of Trial condition suggests that early tone detection might provide participants more time to prepare for button responses.

(D) ERPs to the target tone averaged across three blocks. Channel Cz was selected for statistical analyses. Two significant temporal regions showed a main effect of Temporal location $(p < 0.05)$: region of interest 1 (ROI 1) from 0.18 s to 0.26 s and ROI 2 from 0.35 s to 0.46 s (left plot). ERP amplitudes within each temporal region were averaged to compare ERP magnitude between temporal locations. In both ROIs, the absolute magnitude of ERPs to the target tone significantly increased with the temporal location with the ERP in the third tone location larger than the first and the second tone locations and the ERP in the second tone location larger than the first tone location $(p < 0.05$; adjusted FDR correction was applied).

Fig. S2. Power Coupling Analysis. Refer to Figure 2 for related information. (A) Power of no-tone trials. The plots from left to right represent spectrograms of power in blocks 1, 2, and 3, respectively. (B) Topographies of routinely defined power frequency bands: delta (1-3 Hz), theta (4-7 Hz), alpha (8-12 Hz), beta1 (13-20 Hz), and beta2 (21-30 Hz). The power in each frequency band was averaged over three blocks from 0.25 s to 1.75 s following noise onset. (C) Power coupling between significant clusters depicted in Fig. 2B & D. Phase-locking values (solid black line in each block) were calculated and corresponding thresholds of a one-sided alpha level of 0.01 were determined using surrogate tests (thin pink lines). A significant coupling of power dynamics was observed between Cluster 1 and 2 after noise onset. Additional significant couplings were found after noise onset between Cluster 2 and 3, and between Cluster 3 and 4, suggesting that Clusters 2, 3, and 4 collectively contribute to a common neural process.

Fig. S3. (A) ITPC Modulation Spectrum. Modulation spectrum analyses were conducted on the neural phase measurement - inter-trial phase coherence (ITPC). However, only one cluster displayed robust modulation strength below the Trend bound. (B) ITPC Cluster Dynamics. The ITPC was filtered according to the cluster defined in (A), and the dynamics of ITPC are shown. The ITPC cluster indicates increased phase coherences at noise onset and offset, without displaying signatures of coding local temporal locations in the white noise or monitoring the elapsed time of the noise. It should be noted that ITPC is closely associated with phase-locked, or event-locked, neural responses that underlie routinely-defined 'evoked power'. The power calculation method we used differs from the 'evoked power' calculation method, but the ITPC modulation result here illustrates that the phase-locked responses, or 'evoked power', do not capture the process of sequential temporal anticipation. Refer to Methods for more details. The shaded areas of color represent ±1 standard error of the mean over participants.

Fig. S4. Analyses of Gaussian Fits to Neural Peaks of Corrected Cluster 234 and Their Correlation with Behavioral Measurements. Refer to Fig. 3 for related information. (A) Gaussian fits were applied to each neural peak of Corrected Cluster 234 in every block, resulting in derived peak latencies and widths (Standard Deviation of the Gaussian fits). (B) Analysis of the correlation between d-prime values and the magnitude of neural peaks per block. The first block showed a negative, non-significant correlation, while the following two blocks demonstrated positive, significant correlations. This suggests that, in the second and third blocks, participants utilized the timing information of temporal locations for tone detection. (C) Examination of the correlation between d-prime values and the magnitude of neural peaks per temporal location revealed all positive correlations. This indicates that the neural coding observed in no-tone trials is indeed relevant to tone detection. (D) Analysis of the correlation between the false alarm rate and the magnitude of neural peaks, after the false alarm rates were transformed using the Fisher-Z transformation. A possible explanation for the emergence of Corrected Cluster 234 could be participants' false perception of a tone in no-tone trials, triggering neural peaks. However, the mostly negative or near-zero correlations suggest that this explanation, based on the false alarm rate, does not account for the emergence of Corrected Cluster 234.