Olfactory-driven beta band entrainment of limbic circuitry during neonatal development

Johanna K. Kostka & Ileana L. Hanganu-Opatz

Institute of Developmental Neurophysiology, Center for Molecular Neurobiology, University Medical Center Hamburg-Eppendorf, 20251 Hamburg, Germany

Corresponding authors: Ileana L. Hanganu-Opatz
                                 hangop@zmnh.uni-hamburg.de

                                 Johanna K. Kostka
                                 johanna.kostka@zmnh.uni-hamburg.de

23   Figures: 6
24   Supplementary Material: 5 Figures, 12 Tables
25   Number of pages: 52
26
27   Number of words in Abstract: 159
28   Number of words in Introduction: 579
29   Number of words in Discussion: 1089
30

ABSTRACT

Cognitive processing relies on the functional refinement of the limbic circuitry during the first

two weeks of life. During this developmental period, when most sensory systems are still

immature, the sense of olfaction acts as "door to the world", providing the main source of

environmental inputs. However, it is unknown whether early olfactory processing shapes the

activity in the limbic circuitry during neonatal development. Here, we address this question by

combining simultaneous *in vivo* recordings from the olfactory bulb (OB), lateral entorhinal

cortex (LEC), hippocampus (HP), and prefrontal cortex (PFC) with opto- and chemogenetic

manipulations of mitral/tufted cells (M/TCs) in the OB of non-anesthetized neonatal mice. We

show that the neonatal OB synchronizes the limbic circuity in beta frequency range. Moreover,

it drives neuronal and network activity in LEC, as well as subsequently, HP and PFC via long-

range projections from mitral cells (MCs) to HP-projecting LEC neurons. Thus, OB activity

shapes the communications within limbic circuits during neonatal development.

56 INTRODUCTION

Coordinated neuronal activity during early development refines the neural circuits that account

for complex processing in the adult brain. During the first two postnatal weeks, when rodents

are still blind, deaf, and perform no active whisking, coordinated activity patterns in the sensory

periphery occur independently of sensory input [1,2]. Spontaneous neuronal discharges in the

retina, cochlea, and whisker pad trigger discontinuous oscillatory bursts in the corresponding

primary sensory cortices [3–7] that are necessary for the development of sensory

discrimination [8]. Similar activity patterns can also be observed in brain areas involved in

higher cognitive processing. Discontinuous theta band oscillations in the LEC entrain similar

activity patterns in the HP, which in turn entrains the prelimbic area (PL) of the PFC [9–12].

Disturbance of these early activity patterns in mouse models of psychiatric risk [13–17] as well

as through pharmacological [18] or optogenetic manipulations [19] led to disruption of adult

circuits and behavioral abilities. However, it is not clear whether stimulus-independent activity

patterns in the sensory periphery impact the limbic network during neonatal development.

Due to the limited or absent functionality of most sensory systems during the first two

postnatal weeks, their contribution to the development of limbic networks has been considered

negligible. This hypothesis has been supported by data showing that the synchrony between

V1 and the HP-PFC network before eye-opening is rather weak [11]. In contrast to other

sensory systems, the olfactory system is functional early in life and newborn mice heavily

depend on it for survival [20]. Correspondingly, the anatomical pathways from OB to cortical

areas are unique among sensory systems. MCs send afferents to the piriform cortex (PiR) and

limbic brain areas such as LEC and amygdala, lacking the relay through the thalamus [21,22].

At adult age, in line with the anatomical connectivity, strong functional coupling during odor

processing has been found between OB and these brain areas. For example, adult olfactory

processing relies on respiration-modulated beta and gamma OB activity [23–25]. Further, beta

oscillations in PiR, LEC, and HP play a critical role in olfactory memory processing [26–28].

Moreover, synchronized beta oscillations between OB-HP and LEC-HP are critically involved

in odor learning [29–32]. Recently, beta oscillations in prefrontal-hippocampal networks have

been identified to support the utilization of odor cues for memory-guided decision making [33].

85   The tight and behaviorally relevant coupling between OB and limbic circuits at adult age

86   leads to the question, which role does olfactory activation early in life play for these circuits.

87   Previously, we showed that discontinuous oscillatory activity in the theta-beta range, emerging

88   as a result of bursting MCs in the neonatal OB, entrains similar oscillatory patterns in LEC

89   [34,35]. However, the role of neuronal and network activity in the OB for the functional

90   entrainment of downstream areas within neonatal limbic circuits is still largely unknown.

91   To address this knowledge gap, we simultaneously monitored single-unit activity (SUA)

and local field potentials (LFP) in OB, LEC, HP, and PFC of non-anesthetized neonatal mice

(postnatal day (P) 8-10) during manipulation of M/TC activity using excitatory opsins and

inhibitory DREADDs. We show that activation of M/TCs triggers action potential firing in LEC

and HP as well as prominent beta oscillations that synchronize the OB with the downstream

cortical areas. Conversely, blocking MC output specifically diminishes OB - limbic network

coupling in beta frequency range. These data document the ability of coordinated activity at

the sensory periphery of newborns to shape the network activity in circuits accounting for adult

cognitive processing.

RESULTS

Oscillatory activity in OB times the network activity in limbic circuits of neonatal mice

To get first insights into the impact of OB activity on developing cortical circuits including LEC,

HP, and PFC, we simultaneously recorded the LFP and multiunit activity (MUA) in all four brain

areas in non-anesthetized neonatal (P8-10) mice (n=56, Figs 1A and B) and assessed the

temporal relationships between network oscillations and neuronal firing. All investigated areas

showed discontinuous oscillatory activity in theta-beta range [11,12,34], accompanied by

continuous low amplitude slow frequency oscillations peaking at 2-4 Hz (respiration rhythm,

RR) (Fig 1B). Half (med: 53.779 %, iqr: 47.681 – 65.104 %, n=20) of the oscillatory events

detected in OB co-occurred in all four brain regions. To quantify the coupling of OB to cortical

areas, we calculated the imaginary coherence (Fig 1C). While a high level of synchrony linked

OB with all investigated cortical areas, the strength of coupling was frequency-dependent,

having the highest magnitude in the beta frequency range for OB-LEC and OB-HP and in the

RR frequency band for OB-LEC and OB-PFC.

To uncover whether OB activity times the neuronal firing of cortical areas, we calculated

the phase-locking of single units (SUA) recorded in OB, LEC, HP, and PFC to beta band

oscillations (12-30 Hz) in OB (Fig 1D). Significantly locked OB units fired shortly before the

trough of the beta cycle, while LEC and HP units were locked to significantly shifted phase

angles (Fig 1D, S1 and S2 Tables). Solely the prefrontal firing showed no phase preference of

locking to the oscillatory phase in OB. Next, we questioned whether the communication

between OB and cortical areas is directed and whether OB acts as a driving force within the

circuit. For this, we assessed the temporal relationship between the firing in cortical regions

and OB by calculating the standardized cross-covariance of unit pairs [36]. For unit pairs

between OB and LEC, OB and HP, and OB and PFC, the peak of cross-covariance was at

negative time-lags, indicating that spiking in OB preceded cortical firing (Fig 1E). Monitoring

the timing of interactions between cortical areas (LEC-HP, LEC-PFC, and HP-PFC) confirmed

the previously reported directionality of communication [12], yet less clear as for the OB-driven

coupling. As spike-dependent methods are strongly biased by the firing rate of investigated

neurons, which is rather low in neonatal mice, we next used the spectral dependency ratio

(SDR), a method that infers causal direction from time-series data [37,38], to confirm the

directed communication between OB and cortical areas. SDR values for OB $\rightarrow$ LEC were

significantly higher than for LEC $\rightarrow$ OB, supporting the drive from OB to LEC. Further, the SDR

analysis revealed a spectral dependency of HP as well as PFC on OB, suggesting the

contribution of OB activity to the oscillatory entrainment of prefrontal and hippocampal circuits

(Fig 1F and S3 Table). Moreover, the analysis confirmed the previously reported directed

interaction from HP to PFC and LEC to PFC [11,12]. No SDR difference was detected for

LEC-HP, indicating that, in line with anatomical data [12], a bidirectional coupling links HP and

LEC (Fig 1F and S3 Table).

Thus, tight directed interactions between OB and cortical areas ensure timed firing and

oscillatory entrainment within downstream LEC-HP circuits.

Fig 1. Functional coupling between neonatal OB, LEC, HP, and PFC

(A) Top, schematic of recording configuration for simultaneous extracellular recordings in OB, LEC, HP, and PFC. The positions of recording sites were displayed superimposed on the corresponding brain areas (Brainrender: [39]). Bottom, digital photomontages displaying the DiI-labeled (red) electrode tracks in DAPI (blue) stained slices of OB, LEC, HP, and PL of a P10 mouse. **(B)** Representative MUA trace

recorded in the mitral cell layer (MCL) displayed together with the wavelet spectra of LFP recorded

simultaneously in OB, LEC, HP, and PFC. **(C)** Spectra of imaginary coherence calculated for OB - LEC

(yellow), OB - HP (green), and OB - PFC (blue). **(D)** Top, polar plots displaying the phase-locking of

significantly locked units in OB (red), LEC (yellow), HP (green), and PFC (blue) to beta oscillations in

OB. Bottom, histograms of mean phase angle for significantly phase-locked OB (red), LEC (yellow), HP

(green), and PFC (blue) units. Histograms are replicated over two OB beta cycles (gray curve). (*** $p <$

0.001, Rayleigh test for non-uniformity). **(E)** Plots of standardized mean spike-spike cross-covariance

for OB - LEC (yellow), OB - HP (green), OB - PFC (blue), LEC - HP (light blue), LEC - PFC (purple), and

HP - LEC (light green). Negative lags indicate that spiking in the first brain area precedes spiking in the

second brain area. **(F)** Spectral dependency ratio (SDR) calculated for OB - LEC (yellow), OB - HP

(green), OB - PFC (blue), LEC - HP (light blue), LEC - PFC (purple), and HP - LEC (light green). Gray

dots and lines correspond to individual animals. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Wilcoxon signed-

rank test).

Activation of M/TCs induces beta oscillations in neonatal OB

To elucidate the mechanisms of directed communication between OB and downstream cortical

areas, we activated ChR2-transfected M/TCs by light and simultaneously monitored the

network and neuronal activity in neonatal LEC, HP, and PFC. Transfection of M/TCs was

achieved using a cre-dependent virus vector (AAV9-Ef1a-DIO-hChR2(E123T/T159C)-EYFP)

that was injected into the right OB of P1 Tbet-cre mice (Fig 2A). ChR2-EYFP expression was

reliably detected in M/TCs and their projections 7 days after injection (Fig 2B). Ramp light

stimuli of increasing intensity (473 nm, total duration 3 s) were used to activate M/TCs in the

OB of P8-10 mice (Fig 2A). The stimulation parameters have been set in line with previous

data [40] to prevent not only firing as a result of tissue heating but also artificially synchronous

firing patterns and large stimulation artifacts. Ramp stimulation led to a sustained increase of

spike discharge and broad-band (4-100 Hz) LFP power augmentation in OB that peaked in

beta frequency range (12-30 Hz) (Figs 2C, 2D, and S1A Fig). In cre[+] mice, the modulation

indices (MI) for theta, beta, and gamma power were significantly increased and different from

those calculated for cre[-] animals (Fig 2Dii and S5 Table). Correspondingly, SUA strongly

augmented during ramp stimulation (Figs 2E and 2F). This activation was not layer-specific

and, mirroring the tight OB wiring, not only M/TCs but also granule cells (GCs) and other OB

interneurons increased their firing in response to light activation of ChR2-transfected M/TCs

(S1B and S1C Figs). Analysis of the firing onset along OB layers confirmed the global

activation. Cells in the MCL and GCL started to fire immediately after the 3 ms-long light pulses,

whereas cells in the extra plexiform layer (EPL) and glomerular layer (GL) responded with a

brief delay (Fig 2G).

To assess the temporal relationship between neuronal firing and beta oscillations in

OB, we calculated the locking of SUA firing to the oscillatory phase before (Pre) and during

(Stim) light stimulation. Ramp stimulation caused a significantly stronger locking of OB units to

beta oscillations (Pre: med: 0.147, iqr: 0.094 – 0.227; Stim: med: 0.180, iqr: 0.103 – 0.294,

n_{units}=176 from 26 mice, p=$9.39*10^{-5}$, LMEM) (Fig 2Hi) and an augmentation of the proportion

of significantly phase-locked units to the beta rhythm during ramp stimulation (Pre: 16.478 %,

29/176 units, Stim: 55.114 %, 97/176 units, p=$3.12*10^{-14}$, Fisher's exact test) (Fig 2Hii). Of

note, the coupling of OB units to the RR phase was weaker (Pre: med: 0.129, iqr: 0.080 –

0.220, Stim: med: 0.097, iqr: 0.057 – 0.157, n_{units}=176 from 26 mice, p=$7.782*10^{-6}$, LMEM)

(S1D Fig) even though the proportion of locked units (Pre: 14.773 %, 26/176 units, Stim: 14.205

%, 25/176 units, p=1, Fisher's exact test) and the power of RR oscillations were not altered

upon light stimulation (S1D(iii) and S2D Figs). In contrast, light stimulation had no effects on

the phase-locking of OB units to oscillatory phase in cre⁻ mice (S1E Fig and S4 Table). The

larger beta power observed during ramp stimulation might result from increased M/TC and

interneuronal firing, since spike-triggered power (STP) analysis revealed that the ability of OB

units to trigger beta power is stronger during ramp stimulation compared to baseline periods

(Pre: med: 6.694 µV², iqr: 2.291 – 16.447 µV²; Stim: med: 18.285 µV², iqr: 4.437 – 58.407 µV²;

n_{units} = 309 from 19 mice, p = $3.16*10^{-13}$, LMEM) (Fig 2I).

These data indicate that the activation of M/TCs recruits the local circuitry in the OB and

thereby organizes the OB network activity in the beta rhythm.

Fig 2. Effects of M/TC manipulation by light on single-unit entrainment and oscillatory activity in OB.

(A) Schematic of the experimental protocol. **(B)** Top, photograph of the dorsal (left) and ventral side (middle) of a brain from a Tbet-cre+ mouse showing EYFP expression in the OB and M/TC axonal projections (LOT) to LEC, piriform transition area (APir), and cortical amygdala (CoA) (right). Bottom, digital photomontages displaying the DiI labeled electrode track in OB (left) and confocal images displaying the mitral cell layer (MCL) of the right OB at different magnifications (middle and right). **(C)** Representative extracellularly recorded LFP in the OB displayed band-pass filtered in different frequency bands and accompanied by the corresponding wavelet spectrum during ramp stimulation, as well as by the simultaneously recorded MUA in the MCL. **(D)** (i) Power spectrum for OB LFP before (orange) and during (red) ramp stimulation. The gray shaded area corresponds to the beta band (12-30 Hz). (ii) Mean MI of LFP power in different frequency bands for cre+ (red) and cre- (black) mice. (red stars for cre+: ***

p < 0.001, Wilcoxon signed-rank test; black stars for comparison cre+ vs. cre-: ** p < 0.01, *** p < 0.001,

Wilcoxon rank-sum test). **(E)** Raster plot of SUA in the OB before, during, and after ramp stimulation.

(F) (i) Z-scored firing rate in response to ramp stimulation of units recorded in the OB of cre+ (red) and

cre- (black) mice. **(ii)** MI of SUA firing in response to ramp stimulation (Significantly activated units are

shown in red, whereas significantly inhibited units in gray, p < 0.01, Wilcoxon signed-rank test). **(G)**

Spiking probability of units located in the granule cell layer (GCL), MCL, external plexiform layer (EPL),

and glomerular layer (GL) after a 3 ms light pulse (blue box, 473 nm) delivered to the OB. **(H) (i)** Phase

locking of OB units to beta oscillations in OB. Left, polar plots displaying phase locking of OB units before

(Pre, orange) and during ramp stimulation (Stim, red). The mean resulting vectors are shown as blue

lines. (*** p < 0.001, Rayleigh test for non-uniformity). Right, violin plots displaying the resulting vector

length (RVL) of OB units before (Pre, orange) and during ramp stimulation (Stim, red). Gray dots and

lines correspond to individual units. (*** p < 0.001, linear mixed-effect model). **(ii)** Percentage of

significantly locked units before (Pre, yellow) and during (Stim, red) stimulation. (*** p < 0.001, Fisher's

exact test). **(I) (i)** Plot of mean MI of spike-triggered power (STP) for cre+ (red) and cre- (black) mice

during ramp stimulation. (black line: p < 0.05, Wilcoxon rank-sum test). **(ii)** Violin plots displaying mean

STP for OB units before (Pre, yellow) and during ramp stimulation (Stim, red). Gray dots and lines

correspond to individual units. (*** p < 0.001, linear mixed-effect model).

M/TC activation drives neuronal firing in LEC and HP

To characterize the downstream effects of beta band entrainment of OB, we firstly analyzed

the organization of OB projections in neonatal mice. In line with morphological investigations

in adult mice [21], we previously showed that MC axons are present in superficial layers of

LEC already at neonatal age [34]. Entorhinal neurons in layer II/III strongly project to HP and

weakly to PFC [12,17]. Here, we performed axonal tracing of M/TCs using the anterograde

virus (AAV9-hSyn-hChR2(H134R)-EYFP) injected into the OB at P8. Simultaneously, we

monitored the entorhinal neurons that project to HP by using the retrograde virus (AAVrg-

CamKIIa-mCherry) injected into the HP at P8 (Figs 3A and 3B). At P18, MC axons expressing

EYFP were present in layer I/II of LEC and PiR (Fig 3B). Additionally, mCherry-expressing HP-

projecting neurons were identified in entorhinal layer II/III. These neurons send their apical

dendrites to layer I of LEC, where they collocate with MC axonal projections (Fig 3B).

243   Since these morphological data suggest that the OB interacts with downstream cortical

areas, in a second step, we monitored the functional impact of direct OB projections on limbic

circuits. For this, we used pulse (3 ms) and ramp (3 s) blue light stimulations (473 nm) of

transfected OB neurons and simultaneously recorded the neuronal activity in LEC, HP, and

PFC. Pulse stimulation of M/TCs induced neuronal firing in all investigated brain areas, except

PFC (Fig 3Ci). While the light-evoked OB firing rate sharply peaked already 7-8 ms post-

stimulus, the responses in the other brain areas were substantially broader and delayed (37

ms in LEC, 45-60 ms in HP). A second firing increase was detected in OB after ~28 ms and

might reflect OB-internal processing or feedback activation from downstream areas. To expand

on these results, we used normalized cross-covariance analysis to uncover the temporal

correlations between light-evoked spike trains in the investigated brain regions. The most

prominent interaction was detected for OB-LEC, with OB firing preceding the entorhinal

discharges (Fig 3Cii). While having a similar directionality, the OB-HP cross-covariance

peaked later and less precisely. The data gives first insights into the communication pathways

relaying the information from M/TCs to LEC and subsequently, to HP.

Ramp stimulation of M/TCs evoked neuronal firing in LEC, HP, and PFC with similar

dynamics: a fast increase in OB followed by a delayed spiking in LEC, and subsequently in HP

and PFC. In OB, SUA abruptly increased with ramp onset (76.157 % of units activated

significantly, 2.847 % units inhibited significantly) and decreased post-stimulus (8.185 % of

units activated significantly, 29.893 % of units inhibited significantly) (Figs 3Di and 3Ei). In

contrast, the average SUA firing rate in LEC, HP, and PFC showed a delayed increase starting

around halfway through the ramp and continuing after the light stimulation (Figs 3Dii-iv).

Analysis of the proportion of activated units during and after ramp revealed that neurons in

downstream areas expressed higher firing rates also after the light was switched off (Fig 3E),

indicating that the activation of M/TCs boosted the cortical network activation. Correspondingly,

this post-stimulus firing increase recruited more neurons than those activated during ramp

stimulation (LEC: 10.959 % during stimulation vs. 21.233 % post-stimulus; HP: 9.167 % vs.

10.833 %, PFC: 9.195 % vs. 19.540 %). In HP, the post-stimulus network effect was not

restricted to activation of neurons but also related to the increase in the proportion of neurons

that were inhibited after the ramp (7.5 % vs. 11.667 %). Light stimulation of control animals did

not change the average firing rate of units in all four investigated brain regions (S2A Fig and

S2B Fig).

Thus, M/TC firing drives the activation of entorhinal, and subsequently, hippocampal

and prefrontal circuits.

Fig 3. Effects of optogenetic manipulation of M/TCs on single-unit activity in LEC, HP, and PFC.

Fig 3. Effects of optogenetic manipulation of M/TCs on single-unit activity in LEC, HP, and PFC.

(A) Schematic of the experimental protocol used to trace MC axons and neurons projecting to HP.

(Brainrender: [39]). **(B)** Top, digital photomontages displaying EYFP (green) and mCherry (red)

fluorescence in coronal slices including OB (left, injection side of AAV9-hSyn-hChR2-EYFP), HP

(middle, injection site of AAVrg-CamKIIα-mCherry), and LEC (right). Note the co-expression of EYFP

and mCherry in LEC. Middle, EYFP (left), mCherry (middle), and their co-expression in the LEC are

shown at larger magnification (dashed box). Bottom, EYFP (left), mCherry (middle), and their co-

expression shown at larger magnification for a HP-projecting entorhinal neuron with dendrites targeting

layer I. **(C) (i)** Spike probability of units in OB (red), LEC (yellow), HP (green), and PFC (blue) after a 3

ms light pulse (473 nm) delivered to the OB. Numbers indicate the delay of the peak spike probability

for each brain area. **(ii)** Spike-spike cross-covariance for OB - LEC (yellow), OB - HP (green), and OB -

PFC (blue). Negative lags correspond to OB activity driving spiking in other brain areas. **(D) (i)** Left, z-

scored firing rate of units recorded in the OB of cre$^+$ (red) mice in response to light stimulation. Right, z-

scored firing rate of significantly activated units during ramp stimulation. **(ii)** Same as (i) for units

recorded in LEC (yellow). **(iii)** Same as (i) for units recorded in HP (green). **(iv)** Same as (i) for units

recorded in PFC (blue). **(E) (i)** Left, volcano plot displaying the MI of SUA firing rates recorded in the OB

before (Pre) vs. during (Stim) ramp stimulation (significant activated units are shown in red and

significant inhibited units in gray, $p < 0.01$, Wilcoxon signed-rank test). Middle, same as the left image

but for SUA firing rates before (Pre) vs. after (Post) ramp stimulation. Right, bar plots depicting the

percentage of activated (red) and inhibited (gray) units during (Stim) and after (Post) ramp stimulation.

(ii) Same as (i) for units recorded in LEC. **(iii)** Same as (i) for units recorded in HP. **(iv)** Same as (i) for

units recorded in PFC.

M/TC activation boosts beta band coupling within downstream limbic circuits

The long-lasting effects of M/TC stimulation on the neuronal firing of downstream areas, LEC,

HP, and PFC suggest that OB activation might act as a driving force for the generation of

network oscillations in neonatal limbic circuits. To test this hypothesis, we paired ramp light

stimulation of ChR2-transfected M/TCs with LFP recordings in LEC, HP, and PFC of P8-10

mice. Ramp stimulation of M/TCs increased the oscillatory power in LEC, HP, and PFC (Figs

4A, 4B, and S5 Table). The most prominent increase was detected for beta band oscillations.

Moreover, we assessed the degree of synchrony between OB and cortical areas during light

stimulation by calculating the imaginary part of coherence, a measure that is insensitive to

false connectivity arising from volume conduction [41]. The imaginary coherence between OB

and LEC, OB and HP as well as OB and PFC increased during light activation of M/TCs, the

most prominent effects being detected in beta band range (Fig 4C and S6 Table).

These results indicate that activation of M/TCs not only induces beta oscillations in OB

but also increases the 12-30 Hz oscillatory coupling between OB and downstream cortical

areas.

Fig 4. Oscillatory entrainment of limbic circuits as a result of M/TC activation by light.

(A) Representative LFP traces recorded in the OB, LEC, HP, and PFC during ramp stimulation of ChR2-transfected M/TCs accompanied by the corresponding wavelet spectra. Vertical black lines correspond to 100 µV. **(B) (i)** Left, plot of MI for power during ramp stimulation of oscillations in LEC for cre+ (yellow) and cre- (black) mice. Right, MI of LFP power averaged for different frequency bands for cre+ (yellow) and cre- (black) mice. **(ii)** Same as (i) for HP. **(iii)** Same as (i) for PFC. (colored stars for cre+, gray stars for cre-, * p < 0.05, ** p< 0.01, *** p < 0.001, Wilcoxon signed-rank test; black stars for comparison cre+ vs. cre-: * p < 0.05, Wilcoxon rank-sum test). **(C) (i)** Left, imaginary coherence between OB and LEC before (gray) and during (yellow) light stimulation. Right, MI of LFP coherence averaged for different frequency bands for cre+ (yellow) and cre- (black) mice. **(ii)** Same as (i) for OB and HP. **(iii)** Same as (i) for OB and PFC. (colored stars for cre+: * p < 0.05, ** p < 0.01, *** p < 0.001, Wilcoxon signed-rank test; black stars for comparison cre+ vs. cre-: * p < 0.05, Wilcoxon rank-sum test).

Inhibition of M/TC output reduces oscillatory power as well as neuronal firing in OB, LEC, and HP

To elucidate whether M/TC activity is necessary for the generation of oscillatory activity in downstream areas, we used inhibitory DREADDs (hM4D(Gi)) that block vesicle release when expressed in M/TCs by cre-dependent virus vector injection (AAV9-EF1a-DIO-hM4D(Gi)-mCherry) at P1 (Fig 5A). At P8, M/TC soma as well as their axons forming the lateral olfactory tract (LOT), which targets the posterior part of the cerebrum, expressed hM4D(Gi)-mCherry (Fig 5B). We performed extracellular recordings of LFP and SUA from OB, LEC, and HP of P8-10 mice (n=35) before (baseline, 20 min) and after (40 min) subcutaneous injection of C21 (3 mg/kg), a synthetic activator of DREADDs [42] (Fig 5A). Since the impact of OB activation on PFC was rather weak, we did not monitor its activity during OB silencing.

C21 caused broadband power reduction in OB that reached a maximum magnitude within 5 min after the injection (Figs 5C, 5D, and S7 Table) and persisted for at least 2 h (S3C Fig). The occurrence of discontinuous oscillatory events was lower after C21 injection in OB (Fig 5F and S8 Table), indicating that M/TC activity is involved in the generation of discontinuous events in OB. Solely, the continuous RR in OB was not affected by the activation of DREADDs (Fig 5E and S7 Table). Moreover, silencing the M/TC output led to a broadband reduction of oscillatory power in LEC and HP (Figs 5C-E and Table S7). Correspondingly, the time spend in oscillatory events in LEC and HP decreased after inhibition of M/TC output (Fig 5F and S8 Table). In contrast, for cre$^-$ mice LFP power and time spend in oscillatory events did not differ before and after C21 injection (S4A and S4B Figs, S7 and S8 Tables).

Next, we monitored the effects of chemogenetic silencing of M/TCs on the neuronal firing of downstream areas. Inhibitory DREADDs have been described to mainly reduce the vesicle release in the expressing neurons, while having little, if any, impact on their ability to generate action potentials [43,44]. Indeed, C21 injection had a weak effect on SUA in OB (cre$^+$: med MI: -0.071, iqr: -0.322 – 0.208, n=512, p=0.003, Wilcoxon signed-rank test; cre$^-$: med MI: -0.028, iqr: -0.259 – 0.207, n=418, p=0.171, Wilcoxon signed-rank test; cre$^+$ vs. cre$^-$: p=0.198, Wilcoxon rank-sum test) (Figs 5G and 5H). In particular, the neuronal firing within the first 10

min after C21 injection decreased (Fig 5G), being most likely the result of weaker network

interactions within the OB. The DREADDs manipulation affected not only the network and

neuronal activity in OB but also the spike timing by oscillations. In line with the results of spike-

triggered power (STP) analysis, C21 injection decreased the ability of SUA to entrain the OB

in theta, beta, and gamma rhythms (S4A Fig and S9 Table). STP for RR was comparable in

the presence and absence of C21 (S9 Table). The temporal relationship between OB spikes

and oscillatory events in OB was also assessed by calculating the phase-locking of SUA to RR

and beta rhythm, respectively. In line with the results of STP analysis, the phase-locking to

beta (Baseline: med: 0.105, iqr: 0.061 – 0.152; C21: med: 0.093, iqr: 0.053 – 0.139; n_{units}=524

from 16 mice, p=0.003, LMEM) was reduced after C21 injection. In contrast, the phase coupling

to RR (Baseline: med: 0.094, iqr: 0.051 – 0.153; C21: med: 0.159, iqr: 0.078 – 0.310; n_{units}=524

from 16 mice, p=2.20*10^{-16}, LMEM) was increased after C21 injection (S4B Fig).

Silencing the M/TC output strongly reduced the LEC firing (cre^{+}: med MI: -0.420, iqr: -

0.598 – -0.108, n=126, p=2.96*10^{-16}, Wilcoxon signed-rank test; cre^{-}: med MI: 0.069, iqr: -

0.144 – 0.364, n=49, p=0.168, Wilcoxon signed-rank test; cre^{+} vs. cre^{-}: p=3.87*10^{-10}, Wilcoxon

rank-sum test), the effects lasting > 1 hour after C21 injection (Figs 5G and 5H). In contrast,

silencing of M/TC output had a shorter (~20 min) and weaker impact on hippocampal firing

(cre^{+}: med MI: -0.102, iqr: -0.438 – 0.230, n=102, p=0.036, Wilcoxon signed-rank test; cre^{-}:

med MI: -0.047, iqr: -0.256 – 0.109, n=74, p=0.119, Wilcoxon signed-rank test; cre^{+} vs. cre^{-}:

p=0.484, Wilcoxon rank-sum test).

These results indicate that silencing the M/TC output decouples neuronal firing from

beta oscillations in OB and decreases the oscillatory power and neuronal firing in LEC, as a

first downstream station of OB projections. On its turn, the weaker drive from LEC leads to

poor oscillatory entrainment of HP, yet without significant change of its neuronal firing.

Fig 5. Effects of silencing M/TC output by inhibitory DREADDs on the oscillatory activity in OB, LEC, and HP.

Fig 5. Effects of silencing M/TC output by inhibitory DREADDs on the oscillatory activity in OB, LEC, and HP.

(A) Top, schematic of the experimental protocol. Bottom, schematic of recording configuration for simultaneous extracellular recordings in OB, LEC, and HP (Brainrender: [39]). **(B) (i)** Photograph of the dorsal (left) and ventral side (right) of a brain from a P8 Tbet-cre+ mouse showing mCherry (red) expression in the OB and M/TC axonal projections (LOT) to PIR and LEC. **(ii)** Digital photomontages displaying the DiI labeled electrode track (red) in DAPI (blue) stained slices including the OB (left), LEC (middle), and HP (right) from a P10 mouse. (iii) Confocal images displaying the MCL of the right OB at different magnifications. MC bodies, as well as dendrites, express mCherry. (C) Color-coded MI of LFP power before and after C21 injection in OB (left), LEC (middle), and HP (right). Vertical red lines correspond to the C21 injection. **(D)** Plots displaying the MI of LFP power averaged for 2 to 50 Hz before and after C21 injection in cre+ (colored) and cre- (black) mice for OB (right, red), LEC (middle, yellow), and HP (right, green). Vertical red lines correspond to the C21 injection. (black line: p < 0.05, Wilcoxon rank-sum test). **(E)** MI of LFP power averaged for different frequency bands for cre+ (colored) and cre- (black) mice for OB (left), LEC (middle), and HP (right). (Wilcoxon signed-rank test, colored stars for cre+: * p < 0.05, ** p < 0.01, *** p < 0.001; black stars for comparison cre+ vs. cre-: * p < 0.05, ** p < 0.01, *** p < 0.001, Wilcoxon rank-sum test). **(F)** Violin plots displaying the percentage of time spend in discontinuous oscillatory events before (Baseline, gray) and after C21 injection (C21, colored). Black dots and lines correspond to individual animals. (* p < 0.05, ** p < 0.01, *** p < 0.001, Wilcoxon signed-rank test). **(G)** Line plots displaying the MI of averaged SUA firing rates before and after C21 injection in cre+ (colored) and cre- (black) mice for OB (left, red), LEC (middle, yellow), and HP (right, green). Vertical red lines correspond to the C21 injection. (black line: p < 0.05, Wilcoxon signed-rank test). **(H)** Violin plots displaying the MI of averaged SUA firing rates after C21 injection for cre- (black) and cre+ mice (colored) recorded in OB (left, red), LEC (middle, yellow), and HP (right, green). Red and black dots correspond to individual units. (colored stars for cre+: * p < 0.05, ** p < 0.01, *** p < 0.001, Wilcoxon signed-rank test; black stars for comparison cre+ vs. cre-: *** p < 0.001, Wilcoxon rank-sum test)

Inhibition of M/TC output reduces the communication between OB and downstream cortical areas

To back up the hypothesis that the M/TC activity controls the developmental entrainment of limbic circuits, we monitored the communication between OB and downstream areas during silencing of M/TC output with DREADDs by using three distinct measures. First, we assessed the synchrony between OB, LEC, and HP by calculating the imaginary coherence in different frequency bands before (baseline) and after C21 injection (C21) (Figs 6A and 6B). MIs for beta coherence between OB and LEC, and OB and HP were significantly reduced after C21

injection. In contrast, the coherence in other frequency bands was not affected by C21 injection

(Figs 6A and 6B, S10 Table). Moreover, the C21-induced changes in the beta band were not

detected in cre⁻ mice (S6A Fig and S10 Table).

Second, we calculated the phase-amplitude coupling (PAC) to elucidate the role of

M/TCs in the modulation of cortical beta oscillations by the RR phase in OB. C21 injection

significantly reduced the z-scored PAC values between the OB RR phase and the amplitude

of beta oscillations in LEC (Baseline: med: 2.499, iqr: 1.624 – 2.883; C21: med: 1.608, iqr:

0.674 – 2.361, n=13, p=0.017, Wilcoxon signed-rank test) and HP (Baseline: med: 2.363, iqr:

2.135 – 2.764; C21: med: 1.907, iqr: 1.319 – 2.248, n=10, p=0.037, Wilcoxon signed-rank test)

(Fig 6C). Additionally, fewer mice showed significant RR-beta PAC values after C21 injection

(z-score > 1.96) in LEC (Baseline: 53.85% vs. C21: 39.77%) and HP (90% vs. 50 %).

Third, we tested the effect of C21 on the directionality of interactions between OB, LEC,

and HP (Fig 6D). We calculate the SDR and found that the prominent drive from OB to LEC

was absent after silencing of M/TC output, the values for OB → LEC and LEC → OB being

comparable (Fig 6Di, S11 and S12 Tables). Similarly, the drive from OB to HP was disrupted

by C21 injection (Fig 6Diii, S11 and S12 Tables). In contrast, the directionality of interactions

between LEC and HP was not affected by C21 injection. As reported for the baseline

conditions, the mutual interactions LEC-HP persisted after M/TC silencing (Fig 6Dii, S11 and

S12 Tables). Moreover, the C21-induced changes in directionality were not detected in cre⁻

mice (S5B Fig, S11 and S12 Tables).

Thus, these results show that the M/TC activity is critical for the communication

between OB and its downstream cortical areas.

Fig 6. Modulation of functional communication within olfactory-cortical networks through silencing the M/TC output by inhibitory DREADDs.

Fig 6. Modulation of functional communication within olfactory-cortical networks through silencing the M/TC output by inhibitory DREADDs.

(A) Imaginary coherence calculated for OB - LEC (left, yellow), LEC - HP (middle, light blue), and OB -
HP (right, green), before (Baseline, gray) and after C21 injection (C21, colored). (black line: p < 0.05,
Wilcoxon rank-sum test). **(B)** MI of coherence averaged for different frequency bands between OB and
LEC (left, yellow), LEC and HP (middle, light blue), and OB and HP (right, green), for cre[+] (colored) and
cre- (black) mice. (colored stars for cre[+], gray stars for cre[-]: * p < 0.05, Wilcoxon signed-rank test; black
stars for comparison cre[+] vs. cre[-]: * p < 0.05, Wilcoxon rank-sum test). **(C) (i)** Z-scored phase-amplitude
coupling (PAC) between OB phase and LEC (top) and HP (bottom) amplitude, before (Baseline) and
after C21 injection (C21). **(ii)** Difference of PAC values after and before C21 injection for OB - LEC (top)
and OB - HP (bottom). **(iii)** PAC averaged for RR-beta coupling (black box in (i)) for OB - LEC (top) and
OB - HP (bottom), before (Baseline, gray) and after C21 injection (colored). Dotted gray line corresponds
to a z-score of 1.96. (* p < 0.05, Wilcoxon signed-rank test). **(D) (i)** SDR calculated for OB and LEC.
Left, SDR values for OB -> LEC and LEC -> OB before (Baseline, gray) and after C21 injection (C21,
yellow). Right, difference of SDR values for both directions for cre[+] and cre[-] mice. **(ii)** Same as (i) for
LEC and HP (blue). **(iii)** Same as (i) for OB and HP (green). Black dots and lines correspond to individual
animals. (** p < 0.01, *** p < 0.001, Wilcoxon signed-rank test).

DISCUSSION

Long-range interactions within limbic circuits emerge early in life [45] yet it is still unknown
whether the coordinated activity patterns underlying the coupling are endogenously generated
or result through the driving force of sensory systems. Besides muscle twitches [46] and
passive tactile sensation, olfactory inputs are likely candidates for the instruction of limbic
circuitry development. Newborn rodents are not only able to smell from birth on but,
importantly, also use olfactory information for learning and cue-directed behaviors such as
localization of the nipples of the dam [20,47]. A first piece of evidence for the critical role of
olfaction for limbic development is the fact that the neonatal OB shows functional coupling with
the LEC, the gatekeeper of the limbic circuitry, during discontinuous network oscillations in the
theta-beta frequency range as well as in the continuous respiration-related rhythm (RR) [34].
Here, we extended these findings and uncovered that MC firing sets a beta band entrainment
also in downstream areas, such as HP and PFC. The temporal dynamics of oscillatory and

firing activity revealed that even in periods without active odor sampling, OB drives the

activation of limbic circuits.

Layer-specific analysis of SUA revealed that M/TC activation leads to a complex

entrainment of the OB microcircuit that results in augmented firing rate also for interneurons in

the GCL, EPL, and GL. Experimental and modeling studies have shown that both beta and

gamma oscillations in the OB rely on dendro-dendritic interactions between M/TCs and GCs

[24,48–51]. In adults, the emergence of gamma and beta oscillations is controlled by different

excitability states of GCs as well as their dependency on centrifugal input, with beta oscillations

relying on a higher GC excitability and centrifugal feedback projections [30,48,50]. However,

gamma oscillations are absent in the neonatal OB, most likely as a result of the late functional

integration of interneurons into local circuits and the different biophysical properties of MCs

and GCs during development [52–54]. Instead, discontinuous beta band oscillations are

present not only in OB but also in other sensory and limbic areas [10]. In the neonatal PFC,

they have been shown to accelerate along development until reaching the gamma band range

at juvenile age [55]. Similarly, acceleration of beta to gamma oscillations takes place in V1

during the critical period for vision [56,57]. Whether the beta band activity in OB undergoes a

similar transition to faster rhythms and how this process is controlled by interactions within OB

and by feedback projections from PiR and LEC remain to be elucidated.

The present data show that the OB network activation entrains downstream cortical

areas in beta oscillations. In adult rodents, the axonal terminals of MCs have been found to

target fan and pyramidal neurons in LII/III of LEC that, on their turn, relay this information to

the HP [58,59]. The axonal projections from layer II/III LEC pyramidal neurons to CA1 are

involved in associative odor learning in adults [60]. Already at neonatal age, MC axons reach

layer I of LEC [34,61]. Here, projections of layer II/III neurons that target the HP were detected

and they might establish synaptic contacts with the MC axons. Optogenetic stimulation

revealed that the activation of M/TCs induced delayed firing of LEC neurons and HP neurons,

indicating that the pathway OB-to-HP is indeed already functional from birth on. CA1 receives

entorhinal input not only via the direct performant path but also through the tri-synaptic path,

spanning DG and CA3 [62]. The long latency (~ 60 ms) in light-induced CA1 firing might,

therefore, be partly mediated by the tri-synaptic path. Of note, it was recently shown that a

distinct but rather small population of LEC layer II neurons directly projects to the neonatal PFC

[17], yet light stimulation of M/TCs did not recruit it.

Coordinated activity patterns in OB organized by MCs promote not only neuronal firing

but also network activation in downstream areas. Ramp light stimulation of M/TCs led to an

increase of beta band power in LEC, HP, and PFC. This power surge was accompanied by

increased long-lasting SUA firing in all three brain areas, indicating that the initial activation of

neurons is followed by activation of the local networks in LEC, HP, and PFC. Conversely,

blocking vesicle release on MC synapses by DREADDs reduced the broadband power as well

as neuronal firing in LEC and HP. Moreover, coherence analysis revealed increased

oscillatory, mainly beta band coupling, between OB and cortical areas during ramp stimulation,

whereas inhibition of M/TCs vesicle release reduced the drive OB → LEC and OB → HP as

well as RR-beta cross-frequency coupling and beta coherence between OB-LEC and OB-HP.

While the artificial activation of MCs might not be entirely comparable to the neural processes

underlying odor sampling and processing during a learning task, these results identify the beta

rhythm as a potential mechanism of long-range communication between OB and downstream

cortical networks.

What might be the relevance of OB-controlled beta band activation of cortical circuits

during early postnatal development? Beta oscillations have been reported to play a key role in

working memory and decision making in adult humans [63]. Further, prominent beta band

synchrony between cortical areas has been identified during olfactory-guided memory and

decision making tasks in rodents [29,31,33,64]. A similar, but sniffing-independent increase in

hippocampal beta oscillations has been observed during an object learning task [65].

Moreover, the firing of beta-entrained CA1 interneurons during an odor-place associative

memory and decision-making task is related to an accurate performance, indicating that beta

oscillations enable temporal coordination and recruitment of neurons within functional cell

assemblies [33,64]. In line with these experimental data, modeling confirmed that beta

oscillations optimally contribute to the coupling of cell assemblies over long axonal

conductance delays [66–68]. During development, discontinuous beta band events that have

been identified in PFC, HP, and LEC might facilitate the formation of initial cell assemblies with

relevance for cognitive performance later in life. We previously showed that interfering with

beta band oscillations during a defined developmental period causes network miswiring and

poor behavioral performance of adult mice [19]. Similarly, in a mouse model of psychiatric risk

reduced beta band activity at neonatal age has been found to correlate with later cognitive

deficits [13,17]. Here, we identified the olfactory activity as a prominent driver of these early

beta oscillations. The results let us hypothesize that transient disturbance of neonatal olfactory

processing precludes the functional refinement of entorhinal-hippocampal-prefrontal circuits,

ultimately leading to cognitive deficits in adulthood. Further research is warranted to directly

test this hypothesis and elucidate the role of early activity patterns in OB for cognitive

development.

MATERIALS AND METHODS

Ethical Approval

524   All experiments were performed in compliance with the German laws and the guidelines of the

525   European Union for the use of animals in research (European Union Directive 2010/63/EU)

526   and were approved by the local ethical committee (Behörde für Gesundheit und

527   Verbraucherschutz Hamburg, ID 15/17).

Animals

Time-pregnant C57Bl/6/J and Tbet-cre mice from the animal facility of the University Medical

Center Hamburg-Eppendorf were housed individually in breeding cages at a 12h light / 12h

dark cycle and fed ad libitum. Offspring (both sexes) where injected with either AAV9-Ef1a-

DIO-hChR2(E123T_T159C)-EYFP (Addgene, Plasmid #35509) or AAV9-EF1a-DIO-

hM4D(Gi)-mCherry (Addgene, Plasmid #50461) virus at postnatal day (P) 0 or 1. Genotypes

were determined using genomic DNA and following primer sequences (Metabion,

Planegg/Steinkirchen, Germany) as described previously [34]: for Cre: PCR forward primer 5′-

ATCCGAAAAGAAAACGTTGA-3′ and reverse primer 5′-ATCCAGGTTACGGATATAGT-3′.

The PCR reactions were as follows: 10 min at 95 °C, 30 cycles of 45 s at 95 °C, 90 s at 54 °C,

and 90 s at 72 °C, followed by a final extension step of 10 min at 72 °C. In addition to

genotyping, EGFP expression in OB was detected using a dual fluorescent protein flashlight

(Electron microscopy sciences, Hatfield, PA, USA) prior to surgery. At P8-10 cre⁻ and cre⁺ mice

underwent light stimulation or Compound 21 injections and *in vivo* multi-side

electrophysiological recordings.

Surgical procedures and recordings

Virus injection for transfection of MTCs with ChR2 and hm4D(Gi)

For transfection of M/TCs with the ChR2 derivate E123T/T159C or inhibitory DREADDs

(hm4D(Gi)), P0-1 pups were fixed into a stereotaxic apparatus and received unilateral

injections of one of two viral constructs (AAV9-Ef1a-DIO hChR2(E123T/T159C)-EYFP, 200 μl

at titer ≥ 1×10^{13} vg/mL, Plasmid, #35509, Addgene, MA, USA; AAV9-EF1a-DIO-hM4D(Gi)-

mCherry, 200 μl at titer ≥ 1×10^{14} vg/mL Plasmid #50461, Addgene, MA, USA). The virus was

produced by Addgene or the Virus Facility of the University Medical Center Eppendorf. A total

volume of 200 nl was slowly (200 nl/min) delivered at a depth of around 0.5 mm into the right

OB using a micropump (Micro4, WPI, Sarasota, FL). Following injection, the syringe was left

in place for at least 30 s to avoid reflux of fluid. Pups were maintained on a heating blanket

until full recovery and returned to the dam.

Virus injection for tracing

For the transfection of M/TC axons with EYFP and the retrograde labeling of HP-projecting

neurons with mCherry, P0-1 pups received the viral construct AAV9-hSyn-hChR2(H134R)-

EYFP (200 μl at titer ≥ 1×10^{13} vg/mL, #26973-AAV9, Addgene, MA, USA) into the OB and the

retrograde virus AAVrg-CamKIIα-mCherry (80 μl at titer ≥ 7×10^{12} vg/mL, #114469-AAVrg,

Addgene, MA, USA) into the HP. Virus injection was performed similarly as for the transfection

of M/TCs with ChR2 or hm4D(Gi). After 10 days, the brains of investigated mice were perfused

with 4% paraformaldehyde (PFA), sliced and MC axons and HP-projecting neurons in LEC and

PIR were imaged using a confocal microscope.

Surgical procedure for electrophysiology

For in vivo recordings, P8-10 mice underwent surgery according to previously described

protocols [11,34,35]. Under isoflurane anesthesia (induction: 5 %, maintenance: 2.5 %,

Forane, Abbott), the skin above the skull was removed and 0.5 % bupivacaine / 1 % lidocaine

was locally applied on the neck muscles. Two plastic bars were mounted on the nasal and

occipital bones with dental cement. The bone above the right OB (0.5-0.8 mm anterior to

frontonasal suture, 0.5 mm lateral to inter-nasal suture), LEC (0 mm posterior to lambda, 6-7.5

mm lateral from the midline), HP (2.5 mm anterior to lambda, 3.5 mm lateral from the midline)

and PFC (0.5 mm anterior to bregma, 0.1-0.5 mm lateral from the midline) was carefully

removed by drilling a hole of < 0.5 mm in diameter. Throughout surgery and recording session

the mice were maintained on a heating blanket at 37°C.

Multi-site electrophysiological recordings in vivo

Three-side or four-side recordings were performed in non-anesthetized P8-10 mice. For this,

one-shank electrodes (NeuroNexus, MI, USA) with 16 recording sites (0.4-0.8 MΩ impedance,

50 µm inter-site spacing for recordings in OB and HP, 100 µm inter-site spacing for recordings

in LEC and PFC) were inserted into OB (0.5-1.8 mm, angle 0°), LEC (for 4-side recordings,

depth: 2 mm, angle: 180°; for 3-side recordings, depth: 2-2.5 mm, angle: 10°), HP (1.3-1.9 mm,

angle 20°) and PFC (1.8-2.1 mm, angle 0°). For light stimulation one-shank optrodes

(NeuroNexus, MI, USA) with the same configuration as the electrodes were inserted in the OB.

Before insertion, the electrodes were covered with DiI (1,1'-Dioctadecyl-3,3,3',3'-

tetramethylindocarbocyanine perchlorate, Molecular Probes, Eugene, OR). A silver wire was

inserted into the cerebellum and served as a ground and reference electrode. Before data

acquisition, a recovery period of 20 min following the insertion of electrodes was provided.

Extracellular signals were band-pass filtered (0.1 Hz-9 kHz) and digitized (32 kHz or 32,556

kHz) by a multichannel amplifier (Digital Lynx SX; Neuralynx, Bozeman, MO; USA) and

Cheetah acquisition software (Neuralynx). Spontaneous activity was recorded for at least 20

min before light stimulation or Compound 21 (C21, Hellobio, Ireland) injection. The position of

recording electrodes in OB, LEC, HP, and PFC was confirmed after histological assessment

post-mortem. For the analysis of LFP in OB, the recording site centered in the EPL was used,

whereas for HP the recording site located in the CA1 was considered. For analysis of LFP in

LEC only recording sites that were histologically confirmed to be located in superficial

entorhinal layers were used. Similarly, only recordings sites confined to the prelimbic sub-

division of PFC were considered. For the analysis of spiking activity, all recording sites

confirmed to be located in the areas of interest (OB, LEC, HP, and PFC) were considered.

When necessary, spikes recorded in OB were assigned to specific layers according to the

location of recording sites.

Morphology

Mice were anesthetized with 10% ketamine (Ketamidor, Richter Pharma AG, Germany) / 2%

xylazine (Rompun, Bayer, Germany) in 0.9% NaCl solution (10 µg/g body weight, i.p.) and

transcardially perfused with Histofix (Carl Roth, Germany) containing 4% PFA. Brains were

postfixed in 4% PFA for 24 h and sliced. Slices (100 µm-thick) were mounted with Fluoromount

containing DAPI (Sigma-Aldrich, MI, USA). The positions of the DiI-labeled extracellular

electrodes in the OB, LEC, HP, and PFC were reconstructed to confirm their location. Virus

expression was verified by EYFP (for ChR2) or mCherry (for hM4D(Gi)) fluorescence in the

right OB. For confocal imaging of EYFP or mCherry fluorescence in M/TCs, HP, and LEC, 50

µm-thick slices mounted with Vectashield (CA, USA) were used.

Light stimulation

Activation of M/TCs was achieved by either ramp or pulse light stimulation applied using a

diode laser (473 nm; Omicron, Austria) which was controlled by an arduino uno (Arduino, Italy).

For ramp stimulation, a light stimulus with linear increasing power (3 s rise time) was presented

30-60 times. For pulse stimulation 3 ms light pulses at 2 Hz were delivered. Laser power was

adjusted for every recording (1.37-5.15 mW) to reliably induce neuronal firing.

Compound 21 injection

Compound 21 (3 mg/kg solved in 0.9% NaCl) was injected subcutaneously after >20 min

recording of baseline activity, while the mouse was fixed in the stereotaxic apparatus. The

activity was recorded for 40-120 min post-injection.

Data Analysis

LFP analysis. Data were analyzed offline using custom-written scripts in the MATLAB

environment (MathWorks, Natick, MA). Data were first low-passed filtered (<100 Hz) using a

third-order Butterworth filter before down-sampling by factor 20 to 1.6 kHz to analyze LFP. All

filtering procedures were performed in a manner preserving phase information.

Detection of oscillatory activity. Discontinuous network oscillations in the LFP recorded from

OB, LEC, and HP before and after C21 injection were detected using a previously developed

unsupervised algorithm [69]. Briefly, deflections of the root mean square of band-pass filtered

(4-100 Hz) signals exceeding a variance-depending threshold (2 times the standard deviation

from the mean) were assigned as oscillatory periods. Only oscillatory periods lasting at least 1

s were considered for analysis.

Power spectral density. Power spectral density was analyzed for either the entire baseline

period, 2 s long periods before (Pre), and during light ramp stimulation (Stim) for recordings

combined with optogenetic manipulation. For recordings paired with DREADD manipulation,

the power was either calculated for every minute or averaged for the entire baseline period (19

min) and post C21 injection period (30 min). Power was calculated using Welch's method with

non-overlapping windows of 2 s (ramp periods) or 3 s length. Time-frequency plots of power

were calculated with a continuous wavelet transform (Morlet wavelet).

Coherence. The imaginary part of coherence, which is insensitive to volume-conduction-based

effects [41], was calculated for the same time periods as the power by taking the absolute

value of the imaginary component of the normalized cross-spectrum:

641
$$C_{XY}(f) = \left| Im \left(\frac{P_{XY}(f)}{\sqrt{P_{XX}(f)P_{YY}(f)}} \right) \right|.$$

642 *Spectral Dependency Ratio.* The Spectral Dependency Ratio (SDR) was calculated according

643 to Shajarisales et al. [38] from the power spectral densities ($S_x(f)$ and $S_y(f)$) of the signals X

644 and Y:

645
$$SDR_{X \to Y} = \frac{mean\left(S_y(f)\right)}{mean\left(S_x(f)\right) * mean\left(\frac{S_y(f)}{S_x(f)}\right)}$$

646
$$SDR_{Y \to X} = \frac{mean\left(S_x(f)\right)}{mean\left(S_y(f)\right) * mean\left(\frac{S_x(f)}{S_y(f)}\right)}$$

The most likely direction of causation is the one having significantly larger SDR values.

(https://github.com/OpatzLab/HanganuOpatzToolbox/tree/master/LFP analysis/getSDR.m)

Spiking analysis. Single units were automatically detected and clustered using the python-

based software klusta [70] and manually curated using phy (https://github.com/cortex-lab/phy).

The firing rate was computed by dividing the total number of spikes by the duration of the

analyzed time window. To assess the spike probability, histograms of spike count using 1 ms

bins were calculated for periods around the light pulse (50 ms before to 150 ms after) and

normalized to the number of delivered light pulses. Cross-covariance of spike trains was

calculated as described previously [34,36]. Briefly, cross-covariance for two spike trains N_i and

N_j, was estimated from the cross-correlation histogram ($J_{ij}^{T,b}(u)$) as follows:

657
$$\hat{q}_{ij}(u) = \frac{J_{ij}^{T,b}(u)}{bT} - \hat{P}_i \hat{P}_j,$$

658   (b = binsize, T observation period, $\hat{P}_i = \frac{N_i(T)}{T}$, $\hat{P}_j = \frac{N_j(T)}{T}$). The standardized cross-covariance

659   was calculated as

$$Q_{ij}(u) = \sqrt{\frac{bT}{P_iP_j}}\,\hat{q}_{ij}(u),$$

661   with P_i, P_j being the mean firing rates. Only pairs of units with firing rates > 0.05 Hz and

662   significant standardized cross variance were considered. The Null hypothesis was rejected

663   when $|Q_{ij}(u)| > Z_\alpha$. ($Z_\alpha = \sqrt{2}\,erf^{-1}\left(\frac{1-\alpha}{N_{lags}}\right)$; two-tailed critical z value at level $\alpha = 0.01$). The

664   standardized mean cross-variance for one unit was calculated as

$$Q_i(u) = \sqrt{\frac{1}{K}\sum_{j=1}^{K} Q_{ij}(u)},$$

666   *(K*=number of units in 2. Region) and the mean for all unit pairs as: $\langle Q_i(u) \rangle = \frac{1}{L^2}\sum_{i=1}^{L} Q_i(u)$.

667   *Modulation index.* The modulation index (MI) of power, coherence, firing rate, and STP for light

668   stimulation or DREADD manipulation was calculated as

$$MI = \frac{Value_{Stim} - Value_{Pre}}{Value_{Stim} + Value_{Pre}}.$$

Spike-LFP coupling. Phase locking of spiking units to network oscillations was assessed using

a previously described algorithm [36]. For this, the LFP signal was bandpass filtered (2-4 Hz

(RR), 4-12 Hz (theta), 12-30 Hz (beta), 30-100 Hz (gamma)) using a third-order Butterworth

filter. The instantaneous phase was extracted using the Hilbert transform on the filtered signal.

The coupling between spikes and network oscillations was tested for significance using the

Rayleigh test for non-uniformity. For analysis of baseline properties (Fig 1) only neurons that

showed significant phase locking were considered for the analysis of the mean phase angle

and the locking strength, which was calculated as mean resulting vector length (RVL). For

paired comparison of RVLs (Fig 2, S1, S4) all units with a firing rate higher than 0.01 Hz during

baseline (18 min) and after C21 injection (18 min, DREADD manipulation) or more than 10

spikes before (Pre) and during (Stim) light ramp pulses were considered.

(https://github.com/OpatzLab/HanganuOpatzToolbox/blob/master/Spikes-LFP

analysis/getPPC_PLV.m)

Spike-triggered power. Spike-triggered power (STP) was calculated for the same time periods

as RVL by taking the mean of the LFP power for 0.4 s long time windows centered on each

spike.

Phase-amplitude coupling. Phase-amplitude coupling (PAC) between RR phase in OB and

beta band amplitude in LEC and HP was calculated as previously described [71]. Briefly, the

LFP signals were bandpass filtered and the Hilbert transform was used to extract the phase

and amplitude, respectively. Subsequently, the amplitude of the beta-filtered signal in LEC or

HP was determined at each phase of the filtered OB signal. The phase was divided into 16

bins and the mean amplitude for each bin was calculated and normalized to the total number

of bins. The normalized modulation index (MI) was calculated as the deviation between an

empirical and uniform amplitude distribution. MI matrices were z-scored and the average was

calculated for RR (2-3 Hz) – beta (12-30 Hz) coupling.

Statistics

Statistical analysis was performed in MATLAB environment or R Statistical Software. As none

of the data sets were normally distributed, data were tested for significance using Wilcoxon

rank-sum test (2 unrelated samples) or Wilcoxon sign-rank test (2 related samples). Data

(except phase values) are presented as median (med) and interquartile range (iqr). Outlier

removal was applied to paired data points if the distance of their difference from the 25^{th} or 75^{th}

percentile exceeds 2.5 times the interquartile interval of their difference. Phase locking was

tested for significance using the Rayleigh test for non-uniformity. Phase angles were compared

using a circular non-parametric multi-sample test for equal medians. Differences in proportions

were tested using Fisher's exact test. Nested data were analyzed with linear mixed-effects

models (LMEM) using animals as a fixed effect. Significance levels $*p<0.05$, $**p<0.01$ and

$***p<0.001$ were considered. If not included in the text, values and corresponding test statistics

of all presented data can be found in the supplementary material (Table S1-12).

Additional Information

Acknowledgments

We thank A. Marquardt, A. Dahlmann, P. Putthoff and K. Titze for excellent technical

assistance, Dr. I. Braren from the Vector Facility of the UKE for the virus production as well as

Drs. M. Chini, S.H. Bitzenhofer, and R.L. van den Brink for helpful discussions.

Funding

This work was funded by grants of the German Research Foundation (Ha4466/11-1 and SFB

936 B5 to I.L.H.-O), European Research Council (ERC-2015-CoG 681577 to I.L.H.-O.),

Horizon 2020 DEEPER (101016787 to I.L.H.-O.), and Landesforschungsförderung Hamburg

(LFF73 and LFF76 to I.L. H.-O.).

Author Contributions

I.L.H.-O. and J.K.K conceived the study and designed the experiments. J.K.K carried out the

experiments and analyzed the data. J.K.K, and I.L.H.-O. interpreted the data. J.K.K. and I.L.H.-

O. wrote the article. All authors discussed and commented on the manuscript.

Declaration of Interests

The authors declare no competing interests.

REFERENCES

1.   Hanganu-Opatz IL. Between molecules and experience: Role of early patterns of
coordinated activity for the development of cortical maps and sensory abilities. Brain Res
Rev. 2010;64: 160–176. doi:10.1016/j.brainresrev.2010.03.005

2.   Leighton AH, Lohmann C. The Wiring of Developing Sensory Circuits—From Patterned
Spontaneous Activity to Synaptic Plasticity Mechanisms. Front Neural Circuits. 2016;10.
doi:10.3389/fncir.2016.00071

3.   Ackman JB, Burbridge TJ, Crair MC. Retinal waves coordinate patterned activity
throughout the developing visual system. Nature. 2012;490: 219–225.
doi:10.1038/nature11529

4.   Hanganu IL, Ben-Ari Y, Khazipov R. Retinal Waves Trigger Spindle Bursts in the Neonatal
Rat Visual Cortex. J Neurosci. 2006;26: 6728–6736. doi:10.1523/JNEUROSCI.0752-
06.2006

5.   Khazipov R, Sirota A, Leinekugel X, Holmes GL, Ben-Ari Y, Buzsáki G. Early motor
activity drives spindle bursts in the developing somatosensory cortex. Nature. 2004;432:
758–761. doi:10.1038/nature03132

6.   Mizuno H, Luo W, Tarusawa E, Saito YM, Sato T, Yoshimura Y, et al. NMDAR-regulated
dynamics of layer 4 neuronal dendrites during thalamocortical reorganization in neonates.
Neuron. 2014;82: 365–379. doi:10.1016/j.neuron.2014.02.026

7.   Wang HC, Bergles DE. Spontaneous activity in the developing auditory system. Cell
Tissue Res. 2015;361: 65–75. doi:10.1007/s00441-014-2007-5

8.   Che A, Babij R, Iannone AF, Fetcho RN, Ferrer M, Liston C, et al. Layer I Interneurons
Sharpen Sensory Maps during Neonatal Development. Neuron. 2018;99: 98-116.e7.
doi:10.1016/j.neuron.2018.06.002

9.   Ahlbeck J, Song L, Chini M, Bitzenhofer SH, Hanganu-Opatz IL. Glutamatergic drive
along the septo-temporal axis of hippocampus boosts prelimbic oscillations in the
neonatal mouse. Bartos M, editor. eLife. 2018;7: e33158. doi:10.7554/eLife.33158

10.   Bitzenhofer SH, Ahlbeck J, Wolff A, Wiegert JS, Gee CE, Oertner TG, et al. Layer-specific
optogenetic activation of pyramidal neurons causes beta-gamma entrainment of neonatal
networks. Nat Commun. 2017;8: 14563. doi:10.1038/ncomms14563

11.   Brockmann MD, Pöschel B, Cichon N, Hanganu-Opatz IL. Coupled Oscillations Mediate
Directed Interactions between Prefrontal Cortex and Hippocampus of the Neonatal Rat.
Neuron. 2011;71: 332–347. doi:10.1016/j.neuron.2011.05.041

12.   Hartung H, Brockmann MD, Pöschel B, Feo VD, Hanganu-Opatz IL. Thalamic and
Entorhinal Network Activity Differently Modulates the Functional Development of
Prefrontal–Hippocampal Interactions. J Neurosci. 2016;36: 3676–3690.
doi:10.1523/JNEUROSCI.3232-15.2016

13.   Chini M, Pöpplau JA, Lindemann C, Carol-Perdiguer L, Hnida M, Oberländer V, et al.
Resolving and Rescuing Developmental Miswiring in a Mouse Model of Cognitive
Impairment. Neuron. 2020;105: 60-74.e7. doi:10.1016/j.neuron.2019.09.042

14.   Domnick N-K, Gretenkord S, De Feo V, Sedlacik J, Brockmann MD, Hanganu-Opatz IL.
Neonatal hypoxia–ischemia impairs juvenile recognition memory by disrupting the

maturation of prefrontal–hippocampal networks. Exp Neurol. 2015;273: 202–214.
doi:10.1016/j.expneurol.2015.08.017

15.   Hartung H, Cichon N, De Feo V, Riemann S, Schildt S, Lindemann C, et al. From
Shortage to Surge: A Developmental Switch in Hippocampal–Prefrontal Coupling in a
Gene–Environment Model of Neuropsychiatric Disorders. Cereb Cortex. 2016;26: 4265–
4281. doi:10.1093/cercor/bhw274

16.   Richter M, Murtaza N, Scharrenberg R, White SH, Johanns O, Walker S, et al. Altered
TAOK2 activity causes autism-related neurodevelopmental and cognitive abnormalities
through RhoA signaling. Mol Psychiatry. 2019;24: 1329–1350. doi:10.1038/s41380-018-
0025-5

17.   Xu X, Song L, Kringel R, Hanganu-Opatz IL. Developmental decrease of entorhinal gate
disrupts prefrontal-hippocampal communication in immune-challenged DISC1
knockdown mice. Res Sq. 2021. doi:10.21203/rs.3.rs-290304/v1

18.   Krüger H-S, Brockmann MD, Salamon J, Ittrich H, Hanganu-Opatz IL. Neonatal
hippocampal lesion alters the functional maturation of the prefrontal cortex and the early
cognitive development in pre-juvenile rats. Neurobiol Learn Mem. 2012;97: 470–481.
doi:10.1016/j.nlm.2012.04.001

19.   Bitzenhofer SH, Pöpplau JA, Chini M, Marquardt A, Hanganu-Opatz IL. A transient
developmental increase in prefrontal activity alters network maturation and causes
cognitive dysfunction in adult mice. Neuron. 2021;109: 1350-1364.e6.
doi:10.1016/j.neuron.2021.02.011

20.   Logan DW, Brunet LJ, Webb WR, Cutforth T, Ngai J, Stowers L. Learned recognition of
maternal signature odors mediates the first suckling episode in mice. Curr Biol CB.
2012;22: 1998–2007. doi:10.1016/j.cub.2012.08.041

21.   Igarashi KM, Ieki N, An M, Yamaguchi Y, Nagayama S, Kobayakawa K, et al. Parallel
Mitral and Tufted Cell Pathways Route Distinct Odor Information to Different Targets in
the Olfactory Cortex. J Neurosci. 2012;32: 7970–7985. doi:10.1523/JNEUROSCI.0154-
12.2012

22.   Luskin MB, Price JL. The topographic organization of associational fibers of the olfactory
system in the rat, including centrifugal fibers to the olfactory bulb. J Comp Neurol.
1983;216: 264–291. doi:10.1002/cne.902160305

23.   Kay LM. Chapter 9 - Circuit Oscillations in Odor Perception and Memory. In: Barkai E,
Wilson DA, editors. Progress in Brain Research. Elsevier; 2014. pp. 223–251.
doi:10.1016/B978-0-444-63350-7.00009-7

24.   Neville KR, Haberly LB. Beta and Gamma Oscillations in the Olfactory System of the
Urethane-Anesthetized Rat. J Neurophysiol. 2003;90: 3921–3930.
doi:10.1152/jn.00475.2003

25.   Lockmann ALV, Laplagne DA, Tort ABL. Olfactory bulb drives respiration-coupled beta
oscillations in the rat hippocampus. Eur J Neurosci. 2018;48: 2663–2673.
doi:https://doi.org/10.1111/ejn.13665

26.   Gourévitch B, Kay LM, Martin C. Directional Coupling From the Olfactory Bulb to the
Hippocampus During a Go/No-Go Odor Discrimination Task. J Neurophysiol. 2010;103:
2633–2641. doi:10.1152/jn.01075.2009

27. Vanderwolf CH, Zibrowski EM. Pyriform cortex β-waves: odor-specific sensitization
following repeated olfactory stimulation. Brain Res. 2001;892: 301–308.
doi:10.1016/S0006-8993(00)03263-7

28. Xu W, Wilson DA. Odor-evoked activity in the mouse lateral entorhinal cortex.
Neuroscience. 2012;223: 12–20. doi:10.1016/j.neuroscience.2012.07.067

29. Igarashi KM, Lu L, Colgin LL, Moser M-B, Moser EI. Coordination of entorhinal–
hippocampal ensemble activity during associative learning. Nature. 2014;510: 143–147.
doi:10.1038/nature13162

30. Martin C, Gervais R, Messaoudi B, Ravel N. Learning-induced oscillatory activities
correlated to odour recognition: a network activity. Eur J Neurosci. 2006;23: 1801–1810.
doi:10.1111/j.1460-9568.2006.04711.x

31. Martin C, Beshel J, Kay LM. An Olfacto-Hippocampal Network Is Dynamically Involved in
Odor-Discrimination Learning. J Neurophysiol. 2007;98: 2196–2205.
doi:10.1152/jn.00524.2007

32. Ravel N, Chabaud P, Martin C, Gaveau V, Hugues E, Tallon-Baudry C, et al. Olfactory
learning modifies the expression of odour-induced oscillatory responses in the gamma
(60–90 Hz) and beta (15–40 Hz) bands in the rat olfactory bulb. Eur J Neurosci. 2003;17:
350–358. doi:https://doi.org/10.1046/j.1460-9568.2003.02445.x

33. Symanski CA, Bladon JH, Kullberg ET, Jadhav SP. Rhythmic coordination of
hippocampal-prefrontal ensembles for odor-place associative memory and decision
making. bioRxiv. 2021; 2020.06.08.140939. doi:10.1101/2020.06.08.140939

34. Gretenkord S, Kostka JK, Hartung H, Watznauer K, Fleck D, Minier-Toribio A, et al.
Coordinated electrical activity in the olfactory bulb gates the oscillatory entrainment of
entorhinal networks in neonatal mice. PLOS Biol. 2019;17: e2006994.
doi:10.1371/journal.pbio.2006994

35. Kostka JK, Gretenkord S, Spehr M, Hanganu-Opatz IL. Bursting mitral cells time the
oscillatory coupling between olfactory bulb and entorhinal networks in neonatal mice. J
Physiol. 2020;598: 5753–5769. doi:10.1113/JP280131

36. Siapas AG, Lubenov EV, Wilson MA. Prefrontal Phase Locking to Hippocampal Theta
Oscillations. Neuron. 2005;46: 141–151. doi:10.1016/j.neuron.2005.02.028

37. Ramirez-Villegas JF, Besserve M, Murayama Y, Evrard HC, Oeltermann A, Logothetis
NK. Coupling of hippocampal theta and ripples with pontogeniculooccipital waves.
Nature. 2021;589: 96–102. doi:10.1038/s41586-020-2914-4

38. Shajarisales N, Janzing D, Shoelkopf B, Besserve M. Telling cause from effect in
deterministic linear dynamical systems. ArXiv150301299 Cs. 2015 [cited 18 May 2021].
Available: http://arxiv.org/abs/1503.01299

39. Claudi F, Tyson AL, Petrucco L, Margrie TW, Portugues R, Branco T. Brainrender: a
python-based software for visualizing anatomically registered data. bioRxiv. 2020;
2020.02.23.961748. doi:10.1101/2020.02.23.961748

40. Bitzenhofer SH, Ahlbeck J, Hanganu-Opatz IL. Methodological Approach for Optogenetic
Manipulation of Neonatal Neuronal Networks. Front Cell Neurosci. 2017;11: 239.
doi:10.3389/fncel.2017.00239

41. Nolte G, Bai O, Wheaton L, Mari Z, Vorbach S, Hallett M. Identifying true brain interaction
from EEG data using the imaginary part of coherency. Clin Neurophysiol. 2004;115:
2292–2307. doi:10.1016/j.clinph.2004.04.029

42. Thompson KJ, Khajehali E, Bradley SJ, Navarrete JS, Huang XP, Slocum S, et al.
DREADD Agonist 21 Is an Effective Agonist for Muscarinic-Based DREADDs in Vitro and
in Vivo. ACS Pharmacol Transl Sci. 2018;1: 61–72. doi:10.1021/acsptsci.8b00012

43. Roth BL. DREADDs for Neuroscientists. Neuron. 2016;89: 683–694.
doi:10.1016/j.neuron.2016.01.040

44. Stachniak TJ, Ghosh A, Sternson SM. Chemogenetic synaptic silencing of neural circuits
localizes a hypothalamus→midbrain pathway for feeding behavior. Neuron. 2014;82:
797–808. doi:10.1016/j.neuron.2014.04.008

45. Chini M, Hanganu-Opatz IL. Prefrontal Cortex Development in Health and Disease:
Lessons from Rodents and Humans. Trends Neurosci. 2021;44: 227–240.
doi:10.1016/j.tins.2020.10.017

46. Rio-Bermudez CD, Blumberg MS. Active Sleep Promotes Functional Connectivity in
Developing Sensorimotor Networks. BioEssays. 2018;40: 1700234.
doi:10.1002/bies.201700234

47. Welker WI. Analysis of Sniffing of the Albino Rat. Behaviour. 1964;22: 223–244.
Available: https://www.jstor.org/stable/4533073

48. David F, Courtiol E, Buonviso N, Fourcaud-Trocmé N. Competing Mechanisms of
Gamma and Beta Oscillations in the Olfactory Bulb Based on Multimodal Inhibition of
Mitral Cells Over a Respiratory Cycle. eNeuro. 2015;2. doi:10.1523/ENEURO.0018-
15.2015

49. Fourcaud-Trocmé N, Courtiol E, Buonviso N. Two distinct olfactory bulb sublaminar
networks involved in gamma and beta oscillation generation: a CSD study in the
anesthetized rat. Front Neural Circuits. 2014;8. doi:10.3389/fncir.2014.00088

50. Osinski BL, Kay LM. Granule cell excitability regulates gamma and beta oscillations in a
model of the olfactory bulb dendrodendritic microcircuit. J Neurophysiol. 2016;116: 522–
539. doi:10.1152/jn.00988.2015

51. Osinski BL, Kim A, Xiao W, Mehta NM, Kay LM. Pharmacological manipulation of the
olfactory bulb modulates beta oscillations: testing model predictions. J Neurophysiol.
2018;120: 1090–1106. doi:10.1152/jn.00090.2018

52. Dietz SB, Markopoulos F, Murthy VN. Postnatal Development of Dendrodendritic
Inhibition in the Mammalian Olfactory Bulb. Front Cell Neurosci. 2011;5.
doi:10.3389/fncel.2011.00010

53. Fletcher ML, Smith AM, Best AR, Wilson DA. High-Frequency Oscillations Are Not
Necessary for Simple Olfactory Discriminations in Young Rats. J Neurosci. 2005;25: 792–
798. doi:10.1523/JNEUROSCI.4673-04.2005

54. Yu Y, Burton SD, Tripathy SJ, Urban NN. Postnatal development attunes olfactory bulb
mitral cells to high-frequency signaling. J Neurophysiol. 2015;114: 2830–2842.
doi:10.1152/jn.00315.2015

55. Bitzenhofer SH, Pöpplau JA, Hanganu-Opatz I. Gamma activity accelerates during
prefrontal development. Vinck M, Colgin LL, Perrenoud Q, Moore CI, editors. eLife.
2020;9: e56795. doi:10.7554/eLife.56795

56. Chen G, Rasch MJ, Wang R, Zhang X. Experience-dependent emergence of beta and
gamma band oscillations in the primary visual cortex during the critical period. Sci Rep.
2015;5: 17847. doi:10.1038/srep17847

57. Hoy JL, Niell CM. Layer-Specific Refinement of Visual Cortex Function after Eye Opening
in the Awake Mouse. J Neurosci. 2015;35: 3370–3383.

58. Schwerdtfeger WK, Buhl EH, Germroth P. Disynaptic olfactory input to the hippocampus
mediated by stellate cells in the entorhinal cortex. J Comp Neurol. 1990;292: 163–177.
doi:https://doi.org/10.1002/cne.902920202

59. Wouterlood FG, Nederlof J. Terminations of olfactory afferents on layer II and III neurons
in the entorhinal area: degeneration-Golgi-electron microscopic study in the rat. Neurosci
Lett. 1983;36: 105–110. doi:10.1016/0304-3940(83)90250-1

60. Li Y, Xu J, Liu Y, Zhu J, Liu N, Zeng W, et al. A distinct entorhinal cortex to hippocampal
CA1 direct circuit for olfactory associative learning. Nat Neurosci. 2017;20: 559–570.
doi:10.1038/nn.4517

61. Walz A, Omura M, Mombaerts P. Development and topography of the lateral olfactory
tract in the mouse: Imaging by genetically encoded and injected fluorescent markers. J
Neurobiol. 2006;66: 835–846. doi:https://doi.org/10.1002/neu.20266

62. Basu J, Zaremba JD, Cheung SK, Hitti FL, Zemelman BV, Losonczy A, et al. Gating of
hippocampal activity, plasticity, and memory by entorhinal cortex long-range inhibition.
Science. 2016;351. doi:10.1126/science.aaa5694

63. Spitzer B, Haegens S. Beyond the Status Quo: A Role for Beta Oscillations in
Endogenous Content (Re)Activation. eNeuro. 2017;4: ENEURO.0170-17.2017.
doi:10.1523/ENEURO.0170-17.2017

64. Rangel LM, Rueckemann JW, Riviere PD, Keefe KR, Porter BS, Heimbuch IS, et al.
Rhythmic coordination of hippocampal neurons during associative memory processing.
Bartos M, editor. eLife. 2016;5: e09849. doi:10.7554/eLife.09849

65. Iwasaki S, Sasaki T, Ikegaya Y. Hippocampal beta oscillations predict mouse object-
location associative memory performance. Hippocampus. 2021;31: 503–511.
doi:10.1002/hipo.23311

66. Bibbig A, Traub RD, Whittington MA. Long-range synchronization of gamma and beta
oscillations and the plasticity of excitatory and inhibitory synapses: a network model. J
Neurophysiol. 2002;88: 1634–1654. doi:10.1152/jn.2002.88.4.1634

67. Kopell N, Ermentrout GB, Whittington MA, Traub RD. Gamma rhythms and beta rhythms
have different synchronization properties. Proc Natl Acad Sci. 2000;97: 1867–1872.
doi:10.1073/pnas.97.4.1867

68. Kopell N, Whittington MA, Kramer MA. Neuronal assembly dynamics in the beta1
frequency range permits short-term memory. Proc Natl Acad Sci. 2011;108: 3779–3784.
doi:10.1073/pnas.1019676108

69.   Cichon NB, Denker M, Grün S, Hanganu-Opatz IL. Unsupervised classification of
neocortical activity patterns in neonatal and pre-juvenile rodents. Front Neural Circuits.
2014;8. doi:10.3389/fncir.2014.00050

70.   Rossant C, Kadir SN, Goodman DFM, Schulman J, Hunter MLD, Saleem AB, et al. Spike
sorting for large, dense electrode arrays. Nat Neurosci. 2016;19: 634–641.
doi:10.1038/nn.4268

71.   Tort ABL, Komorowski R, Eichenbaum H, Kopell N. Measuring Phase-Amplitude
Coupling Between Neuronal Oscillations of Different Frequencies. J Neurophysiol.
2010;104: 1195–1210. doi:10.1152/jn.00106.2010

Supplementary Figures

S1 Fig. (related to Fig 2). OB activity in response to ramp stimulation

(A) Plot of MI for LFP power in OB during ramp stimulation for cre+ (red) and cre- (black) mice. **(B)** Digital photomontages displaying the DiI-labeled electrode track in a coronal slice including the ChR2-EYFP-expressing OB. Gray dots correspond to the individual recording sites. Dashed white lines mark borders between the different OB layers. **(C)** Z-scored firing rate of units recorded in the different OB layers in response to light stimulation. **(D) (i)** Polar plots displaying phase locking of OB units to RR before (Pre, orange) and during ramp stimulation (Stim, red) for cre+ mice. The mean resulting vectors are displayed as a blue line. (*** p < 0.001, Rayleigh test for non-uniformity). **(ii)** Violin plots displaying the RVL of OB units before (Pre, orange) and during ramp stimulation (Stim, red). Gray dots and lines correspond to individual units. (***p < 0.001, linear-mixed model). **(iii)** Bar plots displaying the percentage of significantly locked units before (Pre, yellow) and during (Stim, red) stimulation. (* p < 0.05, Fisher's exact test). **(E) (i)** Violin plots displaying the RVL of OB units locked to RR before stimulation (Pre, gray) and during ramp stimulation (Stim, black) for cre-

mice. Black dots and lines correspond to individual units. (* p < 0.05, linear mixed-effect model). **(ii)** Same as (i) for beta oscillations.

S2 Fig. (related to Figs 2 and 3). Effects of optogenetic manipulation of M/TCs from cre⁻ mice on single-unit activity.

(A) (i) Z-scored firing rate of units recorded in the OB of cre⁻ (black) mice in response to light stimulation. **(ii)** Same as (i) for units recorded in LEC. **(iii)** Same as (i) for units recorded in HP. **(iv)** Same as (i) for units recorded in PFC. **(B)** (i) Left, volcano plot displaying the MI of SUA firing rates recorded in the OB of cre⁻ mice before (Pre) vs. during (Stim) ramp stimulation (significant activated units are shown in red and significant inhibited units in gray, $p < 0.01$, Wilcoxon signed-rank test). Middle, same as the left image but for SUA firing rates before (Pre) vs. after (Post) ramp stimulation. Right, bar plots depicting the percentage of activated (red) and inhibited (gray) units during (Stim) and after (Post) ramp stimulation. **(ii)** Same as (i) for units recorded in LEC. **(iii)** Same as (i) for units recorded in HP. **(iv)** Same as (i) for units recorded in PFC.

S3 Fig. (related to Fig 5). Effects of Compound 21 injection in cre⁻ mice and long-lasting effects of M/TC silencing by inhibitory DREADDs.

(A) Color-coded MI of LFP power before and after C21 injection in OB (left), LEC (middle), and HP (right) of cre⁻ mice. The red line corresponds to the C21 injection. **(B)** Violin plots displaying the percentage of time spend in discontinuous oscillatory events before (Baseline, gray) and after C21 injection (C21, black) for cre⁻ mice. Black dots and lines correspond to individual animals. (* $p < 0.05$, Wilcoxon signed-rank test). **(C)** Color-coded MI of LFP power before and after (120 min) C21 injection in OB (left), LEC (middle) and HP (right) of cre⁺ mice (n=3). The red line corresponds to the C21 injection.

S4 Fig. (related to Fig 5). STP and phase locking of OB firing to LFP oscillations in OB after silencing the MT/C output by inhibitory DREADDs

(A) (i) Plot of MI of STP for cre⁺ (red) and cre⁻ (black) mice before and after C21 injection in OB. (black line: $p < 0.05$, Wilcoxon signed-rank test). **(ii)** Violin plots displaying mean MI for STP for each OB unit for different frequency bands for cre⁺ (red) and cre⁻ (black) mice. Red and black dots correspond to individual units. (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, linear mixed-effect model). **(B) (i)** Right, phase locking of OB units to RR oscillations in OB. Left, polar plots displaying phase locking of OB units before (Baseline, orange) and after C21 injection (C21, red). The mean resulting vectors are represented as a blue line. (*** $p < 0.001$, Rayleigh test for non-uniformity). Right, violin plots displaying the RVL of OB units before (Baseline, orange) and after C21 injection (C21, red). Gray dots and lines correspond to individual units. **(ii)** Same as (i) for beta oscillations in the OB (** $p < 0.01$, *** $p < 0.001$, linear mixed-effect model)

S5 Fig. (related to Fig 6). Effects of Compound 21 injection on oscillatory coupling in cre⁻ mice

(A) Imaginary coherence calculated for OB - LEC (top), LEC - HP (middle) and OB - HP (bottom) in cre⁻ mice, before (Baseline, gray) and after C21 injection (C21, black). (black line: p < 0.05, Wilcoxon rank-sum test). **(B)** SDR for OB and LEC (top): SDR values for OB -> LEC and LEC -> OB before (Baseline, gray) and after C21 injection (C21, black). Black dots and lines correspond to individual animals. Middle, same for LEC and HP. Bottom, same for OB and HP. (* p < 0.05, Wilcoxon signed-rank test).

Supplementary Data Tables

	Phase (circ. mean ± circ. SEM, % of sig. locked units)	**p** (Rayleigh test for non-uniformity)
OB	0.763 π ± 0.006 π 640/1124 units (56.94 %)	p=7.6*10^{-297}
LEC	1.556 π ± 0.038 π 42/384 units (10.94 %)	p=4.19*10^{-11}
HP	0.315 π ± 0.047 π 51/384 units (13.28 %)	p=3.68*10^{-5}
PFC	1.715 π ± 0.082 π 26/318 units (8.18 %)	p=0.605

S1 Table (related to Fig 1D). Phase locking of SUA to OB beta phase.

1. Brain area (circ. mean ± circ. SEM)		**2. Brain area** (circ. mean ± circ. SEM)		**p** (Non-parametric multi-sample test for equal medians)
OB	0.763 π ± 0.006 π	**LEC**	1.556 π ± 0.038 π	p=4.58*10^{-4}
		HP	0.315 π ± 0.047 π	p=2.33*10^{-5}
		PFC	1.715 π ± 0.082 π	p=0.689
LEC	1.556 π ± 0.038 π	**HP**	0.315 π ± 0.047 π	p=9.44*10^{-9}
		PFC	1.715 π ± 0.082 π	p=0.618
HP	0.315 π ± 0.047 π	**PFC**	1.715 π ± 0.082 π	p=0.045

S2 Table (related to Fig 1D). Phase angles of significant locked SUA to OB beta phase.

	Area 1 → Area 2 (median, interquartile range)	Area 2 → Area 1 (median, interquartile range)	p (Wilcoxon signed-rank test)
Area 1: OB **Area 2: LEC**	2.051 [1.666 – 2.580]	0.407 [0.262 – 0.522]	p=5.68*10^{-8} (n=39)
Area 1: OB **Area 2: HP**	1.389 [0.920 – 1.868]	0.518 [0.334 – 0.657]	p=2.52*10^{-5} (n=38)
Area 1: OB **Area 2: PFC**	2.809 [1.925 – 3.760]	0.309 [0.226 – 0.399]	p=8.39*10^{-8} (n=38)
Area 1: LEC **Area 2: HP**	0.650 [0.509 – 1.019]	1.022 [0.831 – 1.372]	p=0.056 (n=28)
Area 1: LEC **Area 2: PFC**	1.188 [0.895 – 1.375]	0.762 [0.635 – 1.045]	p=0.024 (n=30)
Area 1: HP **Area 2: PFC**	1.717 [1.317 – 2.034]	0.481 [0.344 – 0.647]	p=0.0006 (n=24)

S3 Table (related to Fig 1F). SDR for pairs of brain areas

Cre- (n_{units}=100 from 13 mice)	Pre (median, interquartile range)	Stim (median, interquartile range)	p (linear fixed-effect model)
RR [2–4 Hz]	0.131 [0.083 – 0.196]	0.128 [0.085 – 0.208]	p=0.763
Beta [12-30 Hz]	0.155 [0.088 – 0.222]	0.140 [0.087 – 0.205]	p=0.486

S4 Table (related to S1E Fig). Mean resulting vector length for phase locking of OB firing to RR and beta oscillations for cre⁻ mice.

		Cre+ (median, interquartile range, Wilcoxon signed-rank test)	Cre- (median, interquartile range, Wilcoxon signed-rank test)	p (Wilcoxon rank-sum test)
OB	**RR [2-4 Hz]**	-0.055 [-0.202 – 0.039] p=0.073 (n=32)	-0.011 [-0.061 – 0.029] p=0.463 (n=14)	p=0.397
	Theta [4-12 Hz]	0.199 [0.056 – 0.322] p=4.96*10^{-5} (n=32)	-0.003 [-0.023 – 0.047] p=0.952 (n=14)	p=0.001
	Beta [12-30 Hz]	0.586 [0.336 – 0.679] p=7.95*10^{-7} (n=32)	0.004 [-0.036 – 0.018] p=1 (n=14)	p=9.52*10^{-8}
	Gamma [30-100 Hz]	0.555 [0.292 – 0.661] p=7.95*10^{-7} (n=32)	-0.005 [-0.037 – 0.022] p=0.670 (n=14)	p=9.52*10^{-8}
LEC	**RR [2-4 Hz]**	0.111 [-0.032 – 0.225] p=0.019 (n=22)	0.096 [0.008 – 0.184] p=0.064 (n=10)	p=0.984
	Theta [4-12 Hz]	0.097 [-0.019 – 0.201] p=0.004 (n=22)	0.065 [0.032 – 0.117] 0.002 (n=10)	p=0.760
	Beta [12-30 Hz]	0.058 [0.021 – 0.118] p=0.0005 (n=22)	0.030 [-0.013 – 0.054] p=0.275 (n=10)	p=0.044
	Gamma [30-100 Hz]	0.044 [0.013 – 0.101] p=0.0008 (n=22)	0.020 [-0.014 – 0.033] p=0.375 (n=10)	p=0.070
HP	**RR [2-4 Hz]**	0.002 [-0.113 – 0.222] p=0.661 (n=22)	0.073 [-0.052 – 0.110] p=0.206 (n=11)	p=0.580
	Theta [4-12 Hz]	0.043 [0.014 – 0.186] p=0.002 (n=22)	0.003 [-0.022 – 0.098] p=0.413 (n=11)	p=0.152
	Beta [12-30 Hz]	0.061 [0.038 – 0.105] p=0.000983 (n=22)	0.007 [-0.021 – 0.045] p=0.465 (n=11)	p=0.049
	Gamma [30-100 Hz]	0.067 [0.016 – 0.114] p=0.001 (n=22)	0.042 [-0.028 – 0.057] p=0.206 (n=11)	p=0.089
PFC	**RR [2-4 Hz]**	0.020 [-0.055 – 0.252] p=0.211 (n=22)	0.017 [-0.033 – 0.191] p=0.461 (n=8)	p=0.981
	Theta [4-12 Hz]	0.083 [0.014 – 0.172] p=0.006 (n=22)	0.032 [-0.022 – 0.068] p=0.25 (n=8)	p=0.336
	Beta [12-30 Hz]	0.080 [0.025 – 0.116] p=0.0002 (n=22)	0.004 [-0.021 – 0.049] p=0.641 (n=8)	p=0.023
	Gamma [30-100 Hz]	0.061 [-0.007 – 0.084] p=0.002 (n=22)	0.010 [-0.011 – 0.064] p=0.383 (n=8)	p=0.336

S5 Table (related to Figs 2D and 4B). MI of power for cre+ and cre- animals during light stimulation.

		Cre[+] (median, interquartile range, Wilcoxon signed-rank test)	Cre[-] (median, interquartile range, Wilcoxon signed-rank test)	p (Wilcoxon rank-sum test)
OB-LEC	**RR [2-4 Hz]**	0.010 [-0.130 – 0.304] p=0.426 (n=22)	0.019 [-0.146 – 0.194] p=0.695 (n=10)	p=0.984
	Theta [4-12 Hz]	-0.030 [-0.136 – 0.078] p=0.426 (n=22)	0.029 [-0.035 – 0.073] p=0.432 (n=10)	p=0.281
	Beta [12-30 Hz]	0.109 [-0.013 – 0.208] p=0.013 (n=22)	0.010 [-0.028 – 0.041] p=0.922 (n=10)	p=0.033
	Gamma [30-100 Hz]	-0.002 [-0.078 – 0.036] p=0.249 (n=22)	0.010 [-0.025 – 0.033] p=0.846 (n=10)	p=0.529
OB-HP	**RR [2-4 Hz]**	0.062 [-0.093 – 0.230] p=0.223 (n=22)	0.103 [-0.191 – 0.280] p=0.577 (n=11)	p=0.924
	Theta [4-12 Hz]	0.036 [-0.060 – 0.143] p= 0.223 (n=22)	-0.041 [-0.235 – 0.061] 0.365 (n=11)	p=0.122
	Beta [12-30 Hz]	0.081 [-0.018 – 0.139] p=0.042 (n=22)	-0.048 [-0.149 – 0.067] p=0.519 (n=11)	p=0.070
	Gamma [30-100 Hz]	-0.028 [-0.067 – 0.016] p=0.082 (n=22)	0.022 [-0.036 – 0.054] p=0.700 (n=11)	p=0.113
OB-PFC	**RR [2-4 Hz]**	-0.091 [-0.253 – 0.173] p=0.322 (n=22)	-0.187 [-0.244 – 0.013] p=0.383 (n=8)	p=0.439
	Theta [4-12 Hz]	-0.006 [-0.096 – 0.107] p=0.910 (n=22)	-0.017 [-0.200 – 0.099] p=0.742 (n=8)	p=0.725
	Beta [12-30 Hz]	0.213 [0.040 – 0.256] p=0.0004 (n=22)	-0.022 [-0.112 – 0.068] p=0.641 (n=8)	p=0.011
	Gamma [30-100 Hz]	0.038 [0.013 – 0.085] p=0.009 (n=22)	-0.020 [-0.059 – 0.036] p=0.641 (n=8)	p=0.071

S6 Table (related to Fig 4C). MI of coherence for cre[+] and cre[-] animals after C21 injection.

		Cre⁺ (median, interquartile range, Wilcoxon signed-rank test)	Cre⁻ (median, interquartile range, Wilcoxon signed-rank test)	p (Wilcoxon rank-sum test)
OB	**RR [2-4 Hz]**	-0.032 [-0.160 – 0.157] p=0.687 (n=17)	-0.025 [-0.084 – 0.116] p=0.879 (n=18)	p=0.609
	Theta [4-12 Hz]	-0.082 [-0.286 – -0.014] p=0.013 (n=17)	-0.010 [-0.094 – 0.156] p=0.472 (n=18)	p=0.031
	Beta [12-30 Hz]	-0.246 [-0.426 – -0.117] p=0.0003 (n=17)	-0.025 [-0.137 – 0.052] p=0.349 (n=18)	p=0.0003
	Gamma [30-100 Hz]	-0.162 [-0.225 – 0.006] p=0.006 (n=17)	-0.009 [-0.083 – 0.040] p=0.647 (n=18)	p=0.028
LEC	**RR [2-4 Hz]**	-0.337 [-0.492 – -0.109] p=0.002 (n=13)	0.246 [-0.096 – 0.488] p=0.063 (n=12)	p=0.000997
	Theta [4-12 Hz]	-0.293 [-0.435 – -0.10] p= 0.001 (n=13)	0.080 [-0.126 – 0.330] 0.259 (n=12)	p=0.0008
	Beta [12-30 Hz]	-0.195 [-0.424 – -0.115] p=0.0002 (n=13)	-0.043 [-0.106 – 0.025] p=0.226 (n=12)	p=0.0005
	Gamma [30-100 Hz]	-0.106 [-0.151 – -0.044] p=0.0005 (n=13)	-0.015 [-0.050 – 0.323] p=0.663 (n=12)	p=0.004
HP	**RR [2- 4 Hz]**	-0.131 [-0.285 – -0.047] p=0.049 (n=10)	-0.074 [-0.160 – 0.036] p=0.426 (n=9)	p=0.243
	Theta [4-12 Hz]	-0.202 [-0.332 – -0.144] p=0.006 (n=10)	-0.053 [-0.116 – 0.017] p=0.359 (n=9)	p=0.010
	Beta [12-30 Hz]	-0.221 [-0.288 – -0.150] p=0.004 (n=10)	-0.069 [-0.112 – 0.001] p=0.055 (n=9)	p=0.017
	Gamma [30-100 Hz]	-0.072 [-0.119 – -0.013] p=0.027 (n=10)	0.003 [-0.043 – 0.033] p=1 (n=9)	p=0.065

S7 Table (related to Fig 5E). MI of power after C21 injection.

		Baseline (median, interquartile range, in %)	Compound 21 (median, interquartile range, in %)	p (Wilcoxon signed-rank test)
OB	**Cre⁺** (n=17)	26.131 [20.490 – 32.841]	19.083 [2.638 – 24.127]	p=0.031
	Cre⁻ (n=17)	32.310 [20.032 – 37.206]	31.103 [22.888 – 51.465]	p=0.435
LEC	**Cre⁺** (n=13)	48.216 [35.999 – 53.788]	14.889 [6.576 – 28.939]	p=0.0005
	Cre⁻ (n=11)	48.294 [29.988 – 59.585]	41.896 [23.372 – 52.576]	p=0.147
HP	**Cre⁺** (n=10)	40.596 [36.197 – 53.448]	18.349 [9.143 – 33.924]	p=0.002
	Cre⁻ (n=9)	53.911 [44.874 – 64.950]	43.233 [30.020 – 60.254]	p=0.164

S8 Table (related to Fig 5F and S3B Fig). Time in discontinuous events.

		Cre⁺ (median, interquartile range) nunits=415 from 15 mice	Cre⁻ (median, interquartile range) nunits=361 from 16 mice	p (linear mixed-effect model)
OB	**RR [2-4 Hz]**	0.230 [0.096 – 0.367]	0.036 [-0.074 – 0.160]	p=0.351
	Theta [4-12 Hz]	-0.025 [-0.189 – 0.056]	-0.014 [-0.098 – 0.093]	p=0.017
	Beta [12-30 Hz]	-0.184 [-0.368 – -0.075]	-0.041 [-0.121 – 0.009]	p=0.0002
	Gamma [30-100 Hz]	-0.103 [-0.195 – 0.016]	-0.020 [-0.071 – 0.044]	p=0.005

S9 Table (related to S4A Fig). STP for OB after C21 injection.

		Cre⁺ (median, interquartile range, Wilcoxon signed-rank test)	Cre⁻ (median, interquartile range, Wilcoxon signed-rank test)	p (Wilcoxon rank-sum test)
OB-LEC	**RR [2-4 Hz]**	-0.052 [-0.208 – 0.150] p=1 (n=13)	-0.103 [-0.368 – 0.212] p=1 (n=11)	p=0.908
	Theta [4-12 Hz]	-0.278 [-0.476 – 0.213] p=1 (n=13)	-0.074 [-0.218 – 0.083] p=0.549 (n=11)	p=0.862
	Beta [12-30 Hz]	-0.239 [-0.346 – -0.051] p=0.022 (n=13)	0.005 [-0.119 – 0.226] p=0.549 (n=11)	p=0.013
	Gamma [30-100 Hz]	-0.021 [-0.109 – 0.151] p=0.581 (n=13)	-0.021 [-0.124 – 0.330] p =1 (n=11)	p=0.524
LEC-HP	**RR [2-4 Hz]**	-0.072 [-0.268 – 0.528] p=0.727 (n=8)	0.024 [-0.089 – 0.251] p=1 (n=7)	p=0.867
	Theta [4-12 Hz]	-0.082 [-0.436 – 0.092] p= 0.727 (n=8)	-0.176 [-0.304 – -0.056] 0.125 (n=7)	p=0.955
	Beta [12-30 Hz]	-0.095 [-0.214 – -0.026] p=0.070 (n=8)	-0.147 [-0.211 – -0.024] p=0.125 (n=7)	p=0.779
	Gamma [30-100 Hz]	-0.079 [-0.183 – 0.133] p=0.727 (n=8)	-0.022 [-0.053 – 0.090] p=1 (n=7)	p=0.536
OB-HP	**RR [2-4 Hz]**	0.311 [0.214 – 0.501] p=0.109 (n=10)	0.169 [-0.303 – 0.212] p=1 (n=9)	p=0.182
	Theta [4-12 Hz]	0.045 [-0.174 – 0.341] p=0.754 (n=10)	-0.047 [-0.204 – -0.006] p=0.180 (n=9)	p=0.356
	Beta [12-30 Hz]	-0.318 [-0.397 – -0.124] p=0.021 (n=10)	-0.178 [-0.306 – -0.058] p=0.039 (n=9)	p=0.278
	Gamma [30-100 Hz]	0.036 [-0.067 – 0.161] p=1 (n=10)	-0.074 [-0.096 – 0.060] p=0.508 (n=9)	p=0.095

S10 Table (related to Fig 6B). MI coherence for cre⁺ and cre⁻ mice.

		Area 1 → Area 2 (median, interquartile range)	Area 2 → Area 1 (median, interquartile range)	p (Wilcoxon signed-rank test)
Area1: OB **Area2: LEC**	**Baseline Cre+** (n=13)	2.164 [1.769 – 2.777]	0.366 [0.249 – 0.433]	p=0.0002
	C21 Cre+ (n=13)	1.211 [0.871 – 1.666]	0.600 [0.378 – 0.855]	p=0.080
	Baseline Cre- (n=11)	1.616 [0.738 - 2.474]	0.494 [0.345 – 1.228]	p=0.102
	C21 Cre- (n=11)	1.628 [1.062 – 2.802]	0.487 [0.271 – 0.673]	p=0.042
Area1: LEC **Area2: HP**	**Baseline Cre+** (n=8)	0.705 [0.531 – 0.986]	1.064 [0.709 – 1.507]	p=0.195
	C21 Cre+ (n=8)	0.683 [0.534 – 0.896]	1.202 [0.806 – 1.271]	p=0.383
	Baseline Cre- (n=7)	0.607 [0.597 – 0.773]	0.928 [0.563 – 1.171]	p=0.469
	C21 Cre- (n=7)	0.634 [0.436 –0.841]	1.072 [0.510 – 1.528]	p=0.469
Area1: OB **Area2: HP**	**Baseline Cre+** (n=10)	1.422 [1.257 – 2.076]	0.492 [0.367 – 0.618]	p=0.002
	C21 Cre+ (n=10)	1.107 [0.709 – 1.605]	0.531 [0.430 – 1.012]	p=0.625
	Baseline Cre- (n=9)	1.703 [1.181 – 1.752]	0.349 [0.322 – 0.623]	p=0.012
	C21 Cre- (n=9)	1.469 [1.145 – 2.010]	0.304 [0.292 – 0.624]	p=0.012

S11 Table (related to Fig 6D and S5B Fig). SDR before and after C21 injection.

		Baseline (median, interquartile range)	Compound 21 (median, interquartile range)	p (Wilcoxon signed-rank test)
Area1: OB **Area2: LEC**	**Difference Cre+** (n=13)	1.792 [1.300 – 2.512]	0.619 [0.087 – 1.160]	p=0.001
	Difference Cre- (n=11)	1.192 [-0.490 – 2.130]	1.045 [0.479 – 2.531]	p=0.175
Area1: LEC **Area2: HP**	**Difference Cre+** (n=8)	-0.357 [-0.929 – 0.254]	-0.471 [-0.746 – 0.041]	p=0.641
	Difference Cre- (n=7)	-0.326 [-0.582 – 0.210]	-0.438 [-1.092 – 0.331]	p=0.813
Area1: OB **Area2: HP**	**Difference Cre+** (n=10)	0.891 [0.601 – 1.709]	0.627 [-0.303 – 1.175]	p=0.004
	Difference Cre- (n=9)	1.347 [0.557 – 1.413]	1.167 [0.553 – 1.708]	p=0.496

S12 Table (related to Fig 6D). SDR difference for before and after C21 injection.