The maize gene *maternal depression of r1* (*mdr1*) encodes a DNA glycosylase with maternal

and paternal fertility functions

Jonathan I. Gent, Kyle W. Swentowsky, Kaitlin M. Higgins, Fang-Fang Fu, Yibing Zeng, Dong

won Kim, R. Kelly Dawe, Nathan M. Springer, and Sarah N. Anderson

Corresponding author: gent@uga.edu

ABSTRACT

Demethylation of transposons can activate expression of nearby genes and cause

imprinted gene expression in endosperm, and it is hypothesized to lead to expression of

transposon siRNAs that reinforce silencing in the next generation through transfer either into egg

or embryo. Here we describe *maternal derepression of R1* (*mdr1*), a DNA glycosylase with

homology to Arabidopsis *DEMETER* that is partially responsible for demethylation of thousands

of regions in endosperm. Maternally-expressed imprinted genes were enriched strongly enriched

for overlap with demethylated regions, but the majority of genes that overlapped demethylated

regions were not imprinted. Demethylated regions were depleted from the majority of repetitive

DNA in the genome but enriched in a set of transposon families accounting for about a tenth of

the total demethylated regions. Demethylated regions produced few siRNAs and were not

associated with excess CHH methylation in endosperm or other tissues. *mdr1* and its close

homolog *dng102* are essential factors in maternal and paternal fertility in maize, as neither

double mutant microgametophytes nor megagametophytes gave rise to seeds. These data

establish DNA demethylation by glycosylases as essential in maize endosperm and pollen and

suggest that neither transposon regulation nor genomic imprinting are its main function.

INTRODUCTION

In 1970, Jerry Kermicle reported that the *red-color* (*R*) gene required for pigmentation in maize endosperm was preferentially expressed from maternal alleles (Kermicle 1970). Since then, other cases of genomically imprinted genes have been reported for hundreds of genes in maize, other plants, and animals as well [reviewed in (Batista and Köhler 2020)]. Research primarily on *Arabidopsis thaliana* endosperm has revealed that reduction of DNA methylation of the maternal genome can differentiate maternal and paternal alleles. In Arabidopsis, the key enzyme is a DNA glycosylase called DEMETER, which excises 5-methylcytosine from DNA and is primarily active in the central cell of the megagametophyte, which gives rise to the endosperm upon fertilization (Choi et al. 2002; Gehring et al. 2006; Park et al. 2017). In the pollen vegetative nucleus, where DEMETER is also highly expressed, many transposons are demethylated and expressed (He et al. 2019). DNA methylation not only represses transposons, but can also regulate developmental gene expression, presumably through inhibiting or promoting transcription factor binding (O'Malley et al. 2016; Borg et al. 2021; Khouider et al. 2021). Absence of maternal DEMETER causes 100% seed inviability (Choi et al. 2002). Pollen that lack DEMETER give rise to apparently normal seeds, but have reduced transmission frequent relative to wild-type and have pollen tube defects, severity dependent on the ecotype (Xiao et al. 2003; Schoft et al. 2011; Khouider et al. 2021). In rice, maternal glycosylase (ROS1a/OsDNG702) is required for seed development, and paternal glycosylase is required for pollen, such that null mutations cannot be transmitted maternally or paternally. The Arabidopsis genome encodes three other glycosylases whose mutants produce subtle phenotypes in a variety of tissues (Pillot et al. 2010; Yamamuro et al. 2014; Schumann et al. 2017; Schumann et al.

2019; Khouider et al. 2021). Endosperm derived from central cells that lack DEMETER have

excessive DNA methylation at thousands of small regions (Ibarra et al. 2012; Park et al. 2016).

Unifying features of these regions have not been identified, but they exhibit a preference for

short transposons in genic areas and for the ends of long transposons. What guides DEMETER to

these regions is unknown, but involves FACT, a versatile ATP-dependent chromatin remodeling

complex (Frost et al. 2018). DEMETER depends on FACT for activity at about half of the

~10,000 identified target regions, which are distinguished by being more heterochromatic than

the other half. The N-terminal half of DEMETER is likely important for targeting, as truncation

of the protein decreases its activity at normal targets and leads to increased activity in gene

bodies (Zhang et al. 2019).

In addition to maternal demethylation by DEMETER in specific regions, there is a

pronounced and dynamic global reduction in DNA methylation of both parental genomes in

endosperm relative to typical vegetative tissues in Arabidopsis (Ibarra et al. 2012; Moreno-

Romero et al. 2016). A similar phenomenon has been observed in other plants, e.g., rice and in

Brassica napa (Park et al. 2016; Chakraborty et al. 2021). This global reduction in methylation is

not caused by DNA glycosylase activity, but instead by passive demethylation through DNA

replication and reduced expression of multiple components of RNA-directed DNA methylation

(RdDM) and the replication-coupled methyltransferase MET1 in early endosperm or central cell

(Jullien et al. 2012; Belmonte et al. 2013; Kawakatsu et al. 2017). RdDM activity apparently

increases later in endosperm development as evidenced by an increase in DNA methylation in all

sequence contexts during after cellularization (Moreno-Romero et al. 2016). Polycomb

Repressive Complex2 (PRC2) promotes non-CG methylation in endosperm, but even so, non-

CG methylation in endosperm remains at greatly reduced levels relative to other tissues (Ibarra et

al. 2012). Perhaps related, PRC2 functions in imprinting of many genes independently of

DEMETER [reviewed in (Batista and Köhler 2020)]. How much of endosperm demethylation is

determined by the central cell is unclear. The central cell is unusual and distinct from endosperm

in its lack of H3K9me2 and visible chromocenters (Garcia-Aguilar et al. 2010; Pillot et al. 2010;

Yelagandula et al. 2014). It has apparently normal methylation outside of DEMETER target

regions, however (Park et al. 2016).

Maternal demethylation of specific transposons can activate nearby gene expression

[reviewed in (Anderson and Springer 2018; Batista and Köhler 2020)]. Studies in multiple

species have revealed a high level of variation in which genes are imprinted, including within-

species variation, and in some cases imprinting is associated with presence of transposons near

genes (Gehring et al. 2009; Waters et al. 2013; Pignatta et al. 2014; Hatorangan et al. 2016;

Pignatta et al. 2018; Rodrigues et al. 2021). Demethylation of transposons in the central cell is

hypothesized to reinforce transposon silencing in the embryo (Ibarra et al. 2012; Bouyer et al.

2017). In this model, transposon activation in produces siRNAs in central cell or endosperm that

move into the egg cell or embryo. Likewise, DEMETER activity in the pollen vegetative cell

could silence transposons in the sperm cells. (Calarco et al. 2012; Ibarra et al. 2012; Martínez et

al. 2016). Methylation patterns in rice *ros1a* mutant pollen also support this hypothesis (Kim et

al. 2019).

While work on gene regulation and chromatin in Arabidopsis endosperm has provided

many insights into maize as well, maize is different than Arabidopsis in several key aspects.

Maize lacks a CMT2-type chromomethyltransferase, causing CHH methylation (mCHH, where

H is A, T, or C) to be largely depleted from heterochromatin and instead associated specifically

with RdDM (Zemach et al. 2013; Gent et al. 2014). It has a much larger set of transposons,

including ones in and near genes and currently active ones (McCarty et al. 2005; Springer et al.

2018; Dooner et al. 2019). Its endosperm persists past fertilization and has a distinct cellular

morphology (Becraft and Gutierrez-Marcos 2012). Unlike both Arabidopsis and rice endosperm,

maize endosperm in general is highly methylated both early and late in development (Zhang et

al. 2014; Fu et al. 2018). Kermicle's demonstration that certain *R* haplotypes are activated

specifically from the maternal genome in endosperm suggests that similar imprinting

mechanisms function in maize as in rice and Arabidopsis. The phenomenon Kermicle studied is

known as R mottling because of the pigmentation pattern. Maternal *R* is uniformly active and

yields solidly pigmented endosperm. When maternal *R* is absent, non-uniform and weak

expression of paternal *R* causes a mottling pattern. Kermicle also reported that mottling can

occur in the presence of maternal *R* in a mutant he called *mdr* (for *maternal derepression of R*)

(Kermicle 1995) (Figure 1A). He mapped the mutation to the long arm of chromosome 4, found

it to be homozygous fertile, and demonstrated that maternal wild-type *Mdr* alone is required and

sufficient for uniform *R* activation in endosperm (Kermicle 1995). In this study, we mapped *mdr*

to a single gene that encodes one of two glycosylases with endosperm expression in maize and

demonstrated that it is required for full demethylation of thousands of diverse regions in

endosperm.

RESULTS

The *maternal derepression of R1* (*mdr1*) gene encodes a DNA glycosylase.

Consistent with current nomenclature in which the original *red-color* gene *R* was changed

to *R1*, we refer to *mdr* as *mdr1* (https://www.maizegdb.org/nomenclature). To map *mdr1*, we

took advantage of the fact that *mdr1* originated spontaneously in a W22 inbred stock (Kermicle,

personal communication). We crossed an *mdr1* homozygote with the sequenced W22 stock

(Springer et al. 2018) to generate *mdr1*/*Mdr1* heterozygotes. We then crossed heterozygotes as

females with *mdr1* homozygotes as males to produce 50% homozygous and 50% heterozygous

mdr1, which we expected to correspond to mottled and solid kernels. After DNA extraction in

bulk from ~500 plants grown from mottled kernels, we prepared and sequenced Illumina

libraries. For comparison, we also sequenced a single pool of five individuals of the W22 stock.

A search for non-W22 reference mutations that were close to 100% homozygous from the

mottled kernels revealed a single region on the long arm of chromosome 4 spanning

approximately 20 MB (Supplemental Figure S1). Several regions of the genome were not shared

with W22 and had abundant non-W22 variants. The long arm of chromosome 4, however, was

nearly identical to W22 (Supplemental Figure S2). In fact, within the 20 MB region, there were

only 45 SNPs or short indels that passed quality filters, and only one of them was in an exon

(*GRMZM2G422464*). This was a deletion of a single T within a string of seven T's starting at

position 229,617,698 and created a frameshift mutation upstream of predicted Endonuclease III

and RRM-fold domains that are typical of DEMETER and ROS family glycosylases. (Figure

1B). To confirm that the mutation in *GRMZM2G422464* caused the maternal R mottling

phenotype, we also obtained an EMS allele (EMS4-06835d) from the Lu et al (2018) Gene-

Indexed Mutations in Maize collection (Lu et al. 2018), which caused a C to T nucleotide change

in the same exon (at position 229,617,638), producing a premature stop codon. After

backcrossing this allele four times into W22 to create an *EMS4-06835d* heterozygote in a W22

background, we tested for a mottling phenotype by crossing the heterozygote as the female

parent with the *Mdr1* homozygous B73 inbred stock as male. Genotyping of *EMS4-06835d* in

plants grown from these kernels revealed 20 of 20 plants from the strongest mottling kernels

inherited the mutation while only 9 of 28 from solid kernels did (two tailed p-value < 0.0001,

Fishers Exact test). These results provide independent evidence that mutations in

GRMZM2G422464 cause maternal R mottling. Hence we refer to *GRMZM2G422464* as *mdr1*.

Functional *mdr1* or its homolog *dng102* is required for male and female fertility

Gramene paralog analysis (http://www.gramene.org) revealed 4 paralogs of *mdr1,* but

only one with close sequence homology (80% amino acid identity) (Supplemental Table S1),

GRMZM2G123587, also known as *dng102* (for *DNA glycosylase102*). Of the five genes, only

mdr1 and *dng102* have strong expression in endosperm, with *mdr1* higher than *dng102,*

including late in endosperm development [displayed on maizeGDB, using data from prior

expression studies (Stelpflug et al. 2016; Doll et al. 2020)]. Both of these have ~32% amino acid

identity to DEMETER (Supplemental Table S1) and cluster with DEMETER by phylogenetic

analysis (Kapazoglou et al. 2013). The Gramene paralog analysis also revealed a similar pair of

genes in the sorghum genome with one-to-one matches for *mdr1* (*SORBI_3004G149800*) and

dng102 (*SORBI_3008G085300*), indicating conservation of both genes. We obtained two mutant

alleles of *dng102*, the UniformMu insertion *mu1083641* (McCarty et al. 2005) in the C-terminal

RRM-fold domain and a TILLING allele (Till et al. 2004) with a premature stop codon upstream

of both the Endonuclease III domain and the RRM-fold domain. Neither allele produced a

maternal R mottling phenotype, nor any obvious phenotype of any kind. However, *mdr1 dng102*

double heterozygotes failed to transmit either the UniformMu insertion or the TILLING allele of

dng102 simultaneously with the *mdr1* mutant allele, neither maternally or paternally. We took

advantage of the Ds-GFP insertion collection to use GFP fluorescence to screen large numbers of

kernels for mutation transmission frequency (Li et al. 2013; Warman et al. 2020). By creating a

mdr1 dng102 double heterozygote with Ds-GFP insertions closely linked to each wild-type allele

and crossing to non-GFP wild-type plants, we were able to screen thousands of kernels for

paternal Ds-GFP inheritance. All but six of 3,786 resulting kernels were florescent. PCR

genotyping of four of the six revealed wild-type *Dng102*, *Mdr1*, or both. The remaining two

kernels failed to germinate. These six exceptions are likely explained by pollen contamination or

recombination between Ds-GFP insertions and mutations. Evenly filled rows of kernels from

these crosses indicated a pre-fertilization pollen defect (Supplemental Figure S3). In a control

cross with Ds-GFP insertions and wild-type *Dng102* and *Mdr1* alleles, 59 of 249 kernels lacked

florescence. We also verified a requirement for maternal glycosylases by crossing the *mdr1*

dng102 double heterozygotes with both Ds-GFP insertions by wild-type pollen. All of 574

resulting kernels in four crosses were florescent, indicating complete failure to transmit mutant

mdr1 and *dng102* alleles to the seed simultaneously. In all four crosses, a high frequency of gaps

in the ears indicated an early defect in female gametophyte development or seed development

(Supplemental Figure S3).

Identification of high-confidence sets of genomic regions that are demethylated in

endosperm

To determine whether the *mdr1* mutant has detectable changes in DNA methylation

relative to wild-type, we prepared and sequenced EM-seq libraries and searched for differentially

methylated regions (DMRs) between homozygous wild-type and *mdr1* mutant mature

endosperm. In parallel, we also searched for DMRs between wild-type mature endosperm and

wild-type developing tassel, and between wild-type mature endosperm and wild-type developing

leaf. In order to identify regions with high-confidence differential methylation, we used a

conservative approach that required differential methylation of both mCG and mCHG

independently over non-overlapping 200-bp regions. First we identified the set of all regions that

were eligible for differential methylation analysis based on their having at least 3X average

coverage of at least five CGs and five CHGs in both tissues. From this set of eligible regions, we

identified hyperDMR as having a greater than twofold relative increase in percent methylation

(value2/value1 > 2) and a greater than 20% relative increase in methylation (value2 – value1 >

0.20) for both mCG and mCHG. HypoDMRs satisfied the opposite requirements. We used the

entire set of eligible regions as controls for DMRs, since they were subject to the same biases in

terms of ability to map EM-seq reads and containing informative CG and CHGs. In the case of

the wild-type vs. *mdr1* endosperm analyses, these eligible region comprised about a third of the

genome. For endosperm vs. tassel or endosperm vs. leaf, they made up about a fifth of the

genome due to lower sequencing depth.

Consistent with expectations, in the wild-type vs. *mdr1* mutant endosperm analysis,

nearly tenfold more DMRs had lower methylation (hypoDMRs) than higher methylation

(hyperDMRs), comprising about three out of every thousand eligible regions (Figure 2A). The

endosperm vs. tassel comparison yielded the strongest enrichment for hypoDMRs, seventyfold

over hyperDMRs, making up nearly nine of every thousand eligible regions. As the hyperDMRs

are likely indirect effects of endosperm demethylation or experimental noise, we excluded them

from additional analysis, and for brevity we refer to hypoDMRs in endosperm simply as DMRs.

We also compared wild-type tassel with *mdr1* mutant in tassel, which yielded few DMRs, of

which only 0.3 of 1000 eligible regions were hypoDMRs (Figure 2A). While the constraints on

depth of sequencing prevented us from defining an exhaustive list of DMRs even in unique

portions of the genome, DMRs defined by the *mdr1* mutant endosperm should overlap at a high

frequency with DMRs defined by comparison of endosperm with tassel or leaf. Indeed, 78.5% of

the DMRs from the comparison with *mdr1* were within a kilobase of a DMR identified from the

comparison with tassel or leaf. This was a 24-fold increase over the random expectation provided

by eligible regions (3.3% of which were within a kilobase of these DMRs).

Endosperm demethylation is mildly enriched near euchromatin and is distinct from RdDM

To determine what types of genetic elements are demethylated in endosperm, we counted

overlaps (at least 100bp) between different types of annotated features and the three sets of

endosperm DMRs (vs. *mdr1* endosperm WT tassel, WT leaf). As controls we counted overlaps

with all eligible regions. Over one-third of the DMRs overlapped with an annotated transposon,

while approximately two-thirds of eligible regions did. The abundant LTR retrotransposons, both

Gypsy and Copia superfamilies, were the most strongly depleted of DMRs (Figure 2B). Regions

containing centromeric (*CentC*) and knob (*knob180* and *TR-1*) tandem repeats were also strongly

depleted from endosperm DMRs (Figure 2B). This depletion was not because of inability to map

reads to tandem repeat regions, as thousands of the control eligible regions overlapped with

them. No transposon superfamilies were strongly enriched for DMRs, but Helitrons had the

highest enrichment, more than twofold relative to eligible regions. To determine whether

endosperm demethylation is linked to RdDM in other tissues, we identified 200-bp regions of the

genome undergoing RdDM using mCHH in the tassel methylome as the defining feature.

Specifically, we required a minimum of 3X coverage, at least 5 informative cytosines in the

CHH context, and an average mCHH value of greater than 20% over the region. Unlike

endosperm DMRs, RdDM regions exhibited a clear pattern of stronger enrichment near TIR

DNA transposons (Figure 2B). An analysis at the transposon family level identified 75 LTR, 11

Helitron, and 9 TIR families that were enriched (p-value < 0.01, binomial test) for DMRs out of

25,730 families that overlapped with at least one eligible region (Supplemental Figure S4). The

majority of these families (63) were annotated in the RLX grouping, which means they were

retrotransposons but could not be assigned to either Copia or Gypsy. Together the 95 families

accounted for 11.6% of total endosperm vs. tassel DMRs and 1.8% of control eligible regions.

Gene bodies were slightly depleted of DMRs (Figure 2B), but regions flanking genes

were enriched for multiple kilobases upstream and down (Figure 2C). The scarcity of DMRs

from gene bodies, especially near their transcription start sites (TSSs) and polyadenylation sites,

is not surprising given the normal absence of mCG and mCHG from these regions. When we

compared DMRs with only the methylated subset of eligible regions, we found up to 7-fold

enrichment for DMRs internal to both ends of genes (Supplemental Figure S5). We defined

methylated eligible regions as ones with both mCG and mCHG values of at least 0.2. This

indicates that gene ends that are methylated in tassel, in leaf, or in *mdr1* mutant endosperm were

7-fold more likely to overlap DMRs than the rest of the genome was. In contrast, the whole set of

gene ends were about 50% less likely to overlap DMRs than the rest of the genome was. While

endosperm DMRs and RdDM regions were both enriched near genes, RdDM regions were

distinct in being strongly enriched in the first 500 bp upstream of genes (9.5 fold enrichment for

RdDM regions vs. 3 fold for DMRs) (Figure 2C). While LTRs retrotransposons were strongly

depleted of DMRs overall, the large number of LTR retrotransposons in the genome provided an

opportunity to examine locations of DMRs within and near them. On average, DMR enrichment

reached levels expected for a random expectation relative to eligible regions within the first 2 Kb

outside both Gypsy LTRs (Figure 2D) and Copia LTRs (Supplemental Figure S6). RdDM

regions were also strongly depleted within both types of retrotransposons, but had small peaks in

enrichment to approximately the random expectation immediately inside Gypsy LTRs and about

twofold enrichment immediately inside Copia LTRs. Motif analysis revealed several that were

significantly enriched but were only present in a small fraction of endosperm DMRs and none

that were consistently enriched across the three sets of DMRs (Supplemental Figure S7).

Endosperm DMRs were AT rich, on average 58.9% A + T in endosperm DMRs versus 53.9% A

+ T in control eligible regions.

Measuring methylation levels in and flanking endosperm DMRs confirmed their

locations near euchromatin and revealed that kilobase-scale regions extending beyond the 200-bp

regions defined by our method were detectably demethylated (Figure 3 and Supplemental Figure

S8). For these analyses, DMRs were oriented relative to the closest euchromatin, using absence

of mCHG as a proxy for euchromatin. Interestingly, *mdr1* mutant endosperm partially restored

endosperm DMRs to a normal, non-endosperm state of mCG and mCHG. We also confirmed

this by examining the distribution of methylation values for individual DMRs for the wild-type

tissues and the *mdr1* mutant endosperm (Supplemental Figure S9). In terms of mCHH, *mdr1*

caused an increase even beyond that of non-endosperm tissues, though still not much above

background levels. Comparison with methylation of control eligible regions also indicated that

methylation of mature endosperm is similar to somatic tissues at a genome-wide level. Overall,

these data indicate that demethylation in endosperm is partially dependent on MDR1, distinct

from RdDM, occurs in a limited but wide variety of genomic contexts, and has a mild preference

for regions near euchromatin.

Except near euchromatin, endosperm DMR siRNAs are in low abundance in endosperm

and embryo

Even though endosperm DMRs have little mCHH, it is possible that siRNAs produced by

endosperm demethylation could direct other chromatin modifications or function post-

transcriptionally (Parent et al. 2021). In Arabidopsis, demethylation of heterochromatin in pollen

vegetative cells corresponds to production of 21 and 22nt siRNAs from LTR retrotransposons

(Creasey et al. 2014; McCue et al. 2015; Borg et al. 2021). Since the preference for DMRs is to

be near euchromatin, where they have the potential to overlap RdDM regions, we separated

endosperm DMRs into two groups, euchromatin DMRs, which were within 2 Kb of euchromatin

(10142 regions), and heterochromatin DMRs, which were not (10014 regions). For this, we used

DMRs from the endosperm vs. tassel comparison as it provided the largest number of

hypoDMRs (Figure 2A). In theory, demethylation-dependent siRNAs would be more clearly

evident in heterochromatin DMRs, where they would not be masked by siRNAs produced by

canonical RdDM. We found, however, that 21 and 22nt siRNAs were rare in both euchromatin

and heterochromatin DMRs, and 24nt siRNAs were rare in heterochromatin DMRs (Figure 4A).

This was true of both endosperm siRNAs as well as non-endosperm siRNAs (embryo and tassel).

To quantify the number of regions that associated with siRNAs we counted all 200-bp regions

with at least 50 bp spanned by mapped siRNAs). 21, 22, and 24-nt siRNA regions were each

identified independently (Figure 4B). The results do not rule out the possibility of demethylation

promoting 21 or 22-nt siRNA expression, but they do indicate that 21 or 22-nt siRNA expression

is not a typical feature of endosperm DMRs, nor is expression of 24-nt siRNAs except in the

euchromatic subset of DMRs.

Endosperm DMRs imprint expression of a small subset of DMR-containing genes and

transposons

Since endosperm inherits two copies of each maternal genome and one of each paternal genome,

the expression of maternal transcripts would be expected to account for two thirds of the total

transcript in endosperm. Reciprocal hybrid crosses, where maternal and paternal alleles can be

differentiated by SNPs, provide a means to test this. The average reciprocal expression ratio

(RER) of 0.67 in maize endosperm indeed corresponds to twice as high expression of alleles

when inherited maternally vs. paternally (Waters et al. 2011; Zhang et al. 2011; Anderson et al.

2021). Reanalysis of published data in conjunction with endosperm DMRs reveals that the

proportion of MEGs (genes with RER >.9) overlapping endosperm DMRs is more than 10-fold

higher than the proportion of non-imprinted genes that overlap endosperm DMRs, and the

propotion of PEGs (genes with RER < .1) overlapping DMRs is more 2-fold higher than non-

imprinted genes. (Figure 5A, B). An example MEG is the transcription factor *myb155*

(GRMZM2G150680), with a 3-Kb region overlapping its promoter that is demethylated in

endosperm (Figure 5D). DMRs in MEGs were preferentially located towards their 5' ends

(Figure 5E) while DMRs in genes as a whole had a more balanced representation at both ends

(Figure 5F). These analyses revealed that while some genes with DMRs and expression in

endosperm like *myb155* are imprinted, the majority of genes with DMRs and expression in

endosperm do not appear to be imprinted. The majority of genes with DMRs clustered around

RER scores of 0.67 even if they overlapped DMRs (Figure 5A). Genes with DMRs in either their

2-Kb flanking regions, either 5' or 3', were also enriched for MEGs and PEGs, but these

flanking effects were weak and were less reproducible between the three DMR comparisons

(Figure 5A, B). Endosperm expression in general did not predict overlap with DMRs (Figure

5C).

The majority of transposon copies with sufficient expression in endosperm and with

variants to allow distinguishing parent of origin also have RER values near 0.67 (Anderson et al.

2021). However, we found that the 5% of transposons with MEG-type expression (RER > .9 and

passing significance thresholds) are ~fivefold more likely to overlap endosperm DMRs than the

endosperm-expressed but non-imprinted set (Figure 5G). Only three transposon copies (0.16% of

the endosperm-expressed set) had PEG-type expression (RER <.1), too few for a meaningful

analysis. These results indicate that demethylation in endosperm-expressed genes usually has

weak or no effects on genomic imprinting, but in specific situations demethylation can cause

maternal-specific expression of both genes and transposons.

Maternally mutant *mdr1* causes reduced maternal *r1* gene expression. In theory, full

maternal expression of the *r1* gene and other MEGs could depend on maternal *mdr1* expression

either because of a requirement for demethylation initiating before paternal genome activation or

because *mdr1* is itself a MEG. To test this, we looked at prior imprinting data in W22 and B73

(Anderson et al. 2021). This revealed that *mdr1* was biparentally expressed, with a reciprocal

expression ratio of 0.73 and 0.68 calculated from over ~1,000 reads for each library. *dng102* was

also biparentally expressed, but with a paternal preference, with ratios of 0.36 and 0.33

calculated from fewer than 100 reads in most libraries. These numbers indicate that neither *mdr1*

nor *dng102* are MEGs and raise the possibility that the *r1* gene and other MEGs are exceptional

in their requirement for *mdr1* prior to paternal genome activation.

DISCUSSION

In Arabidopsis, DEMETER is required for normal development not just in endosperm,

but in the pollen vegetative cell and non-reproductive cell types as well (Borg et al. 2021;

Khouider et al. 2021; Kim et al. 2021). In the case of pollen vegetative cell, developmental

phenotypes are clearly determined by gene regulation rather than transposon activation. Here we

have demonstrated in maize that it is essential to have a functional copy of at least one of the

DNA glycosylases *mdr1* or *dng102* present prior to fertilization in both the maternal and paternal

gametophytes, based on inability to transmit two mutant alleles simultaneously. Since imprinted

gene expression in maize is also clearly linked to demethylation of maternal alleles (Zhang et al.

2014), we expect that gametophyte defects are a consequence of mis-regulated gene expression

rather than genome instability due to transposon activation. Such developmentally dynamic mCG

and mCHG in and near genes would be exceptions to the general rule of such methylation being

stable (Oka et al. 2017; Crisp et al. 2020). In multiple species, demethylation of transposons near

MEGs is clearly connected to their maternal expression (Gehring et al. 2009; Hatorangan et al.

2016; Pignatta et al. 2018; Rodrigues et al. 2021). A prediction from these observations is that a

genome like maize, with more frequent transposons near genes, would have larger numbers of

imprinted genes. This, however, is not the case. In addition, we found that endosperm DMRs

were only weakly associated with transposons, mainly unclassified retrotransposons. These

transposons that were enriched for endosperm DMRs accounted for about a tenth of total

endosperm DMRs. It is clear both from our findings and prior work on other species that DNA

glycosylases demethylate repetitive elements and that their demethylation can cause imprinted

gene expression in endosperm [reviewed in (Anderson and Springer 2018; Batista and Köhler

2020)]. Our results are consistent with this, but further indicate that demethylation of transposons

is not the major function of DNA glycosylases in maize endosperm. Counter-intuitively,

transposon demethylation would make sense in the context of demethylation being part of a

transposon silencing mechanism, where demethylation promotes siRNA expression that transfer

into egg or embryo. While our siRNA expression data does not rule out this hypothesis, it does

indicate that 21 and 22-nt siRNAs are not a typical feature of endosperm DMRs, and 24-nt

siRNAs are only typical of a subset of DMRs near euchromatin. In addition, recent work in

Brassica napa also found no evidence to support the idea of endosperm siRNAs being

transferred to embryo (Chakraborty et al. 2021). These observations suggest transposon defense,

and transposon-mediated gene regulation, are not the main function of DNA glycosylases in

maize endosperm.

Consistent with what is known about glycosylase activity in other plants, MEGs were

strongly enriched for overlapping endosperm DMRs, especially in the 5' end of genes. However,

the majority of genes that were expressed in endosperm and contained endosperm DMRs were

not imprinted (Figure 5A). This indicates that either the majority of endosperm DMRs do not

have a major effect on gene expression or that the effect is not limited to one parent. This latter

possibility, in which glycosylases act post-fertilization on both parental genomes is consistent

with the biparental expression of both *mdr1* and *dng102* in endosperm long after the two parental

genomes have merged. Testing this will require analysis of hybrid methylomes, where SNPs

allow distinguishing one parental allele from another. If glycosylases act on both genomes in

endosperm, both maternal and paternal alleles would have reduced methylation relative to non-

endosperm control tissue. Future experiments with higher read coverage and more complete

identification of DMRs may also reveal whether DMRs in specific genomic contexts, i.e.

overlapping transcription start sites, may have stronger effects on imprinting. It is also possible

that endosperm demethylation has an additional function independent of gene regulation.

Endosperm is unusual in its rapid rate of cell division, which partly takes place in a syncytium

[reviewed in (Becraft and Gutierrez-Marcos 2012)]. One could imagine, for example, that

removal of DNA methylation from select regions of the genome has a function in facilitating

rapid DNA replication. While *mdr1* null mutants effectively behave as partial-loss-of-function

due to complementation by its homolog *dng102*, the viability of *mdr1* endosperm and pollen will

continue to provide an experimental resource for answering these questions.

METHODS

Candidate *mdr1* mutation mapping

Homozygous *mdr1* plants derived from stock X336J (from the Maize Genetics

Cooperation Stock Center, http://maizecoop.cropsci.uiuc.edu/) were crossed with the W22

sequenced stock (Springer et al. 2018) to create *mdr1/Mdr1* heterozygotes. These were crossed

as females with *mdr1* homozygous males derived from stock X336J to produce progeny

segregating homozygous and heterozygous *mdr1*. All plants were homozygous for the *R1-*

r:standard haplotype which is capable of mottling . Approximately 500 putative *mdr1*

homozygotes were selected based on kernel mottling and planted in five batches of 100 each.

Single leaf tips from each seedling were combined for each of the five batches, and a single

DNA extraction performed on each batch, followed by Illumina library preparation with a KAPA

HyperPrep Kit (KK8502). Three batches were sequenced single-end and two sequenced paired-

end, all with 150-nt reads. A single batch of five W22 leaf tips was also processed in the same

manner as a homozygous *Mdr1* control and sequenced single-end. Adapter sequences were

trimmed from all reads using cutadapt 1.9.1 (Martin 2011) with adapter sequence

AGATCGGAAGAGC for both forward and reverse reads and parameters -q 20 -O 1 -m 100.

Trimmed reads were aligned to the W22 reference genome (Springer et al. 2018) using BWA

mem version 0.7.15 (Li and Durbin 2009) with the -M parameter.

To generate a set of high-confidence variants, duplicates were marked using Picard

MarkDuplicates (v2.16.0 http://broadinstitute.github.io/picard). Downstream variant calling was

performed using software in the GATK package (v3.8.1) and the GATK best practices (Van der

Auwera et al. 2013). In particular, indels were realigned using RealignerTargetCreator and

IndelRealigner, and an initial round of SNP calling was performed using HaplotypeCaller and

GenotypeGVCFs. The resulting vcf files for the W22 bulk and the combined *mdr1* bulks were

filtered to remove variants that were present in the W22 bulk using the BEDTools subtract tool

version 2.26.0 (Quinlan and Hall 2010) and stringently filtered to remove low quality score

variants using VCFtools 0.1.15 (Danecek et al. 2011) with the -minQ 200 parameter. The

resulting vcf was used as a training set for base recalibration of the W22 bulk and *mdr1*

combined bulks using BaseRecalibrator, and HaplotypeCaller/GenotypeGVCFs was used to call

variants in W22 and *mdr1*. VariantsToTable was used to convert vcf data into a readable table

for analysis in RStudio.

The mapping scheme predicted that variants across the genome would have a frequency

of 0.75 non-W22 variants (alternative alleles). Since we were looking specifically for a region

was near homozygous for alternative alleles in the *mdr1* bulks, the following additional criteria

were imposed to identify putative variants linked to *mdr1* in the vcf table: 1) An alternative allele

frequency in the W22 bulk of zero, with a read depth of at least two but no more than 15. 2) A

read depth in the *mdr1* combined bulks of at least five but no more than 50. 3) A normalized

Phred-scaled likelihood for homozygous reference alleles of at least 400 and for homozygous

alternative alleles of zero in the *mdr1* combined bulks. Filtering the vcf table in this way selected

for near-homozygous alternative alleles across the genome in the *mdr1* bulks, with an average of

96% alternative allele frequency. A single ~20 MB region (chr4:216965546-237367197) stood

out, with a ~98.5% alternative allele frequency. A second method of analyzing the GATK

variants was also used to identify the *mdr1* candidate region. The vcf table was filtered to

exclude all variants but those with a GenotypeQuality of at least 20 in both genotypes, depth of

coverage between 5-14 in the W22 bulk, a depth between 6-49 in the combined *mdr1* bulks, and

variant frequencies above 0.75 in the *mdr1* bulks. Based on the mapping scheme, this is expected

to remove the majority of technical artifacts. As with the first method, this singled out the same

region on chromosome 4 as the most homozygous in the genome, with variant frequencies near

1, as shown in Supplemental Figure 1. A comparison of variant density by genomic position

revealed that a large region of chromosome 5 as well as a few small regions elsewhere in the

genome of the *mdr1* mutant stock X336J were not shared with the sequenced W22 genome

(Supplemental Figure S2). The tip of chromosome 4, however, had few variants and was clearly

shared between both genomes.

Mutant genotyping

An 867-bp region of *mdr1* was amplified by PCR with primers W22-Ros1-F1 and W22-

Ros1-R1 and digested with TaqI-v2 (NEB #R0149S). The different *mdr1* alleles could be

distinguished by the pattern of restriction fragments due to the presence of SNPs that created or

removed TaqI restriction sites in different genetic backgrounds (Kermicle's *mdr1* allele or its

progenitor in W22, the *EMS4-06835d* allele or its progenitor in B73, and the allele linked to the

Ds-GFP insertion *R128F11* in A188). A single TaqI restriction site in B73 yields two fragments

of 382 and 485 bp. A second site in A188 yields three fragments, 107, 275, and 485 bp. Absence

of any sites in W22 yields an intact amplicon of 867 bp. In cases where the wild-type and mutant

were in the same genetic background or when there was a possibility of recombination between

R128F11 and *mdr1*, the *mdr1* mutation was genotyped by Sanger sequencing of the amplicon.

Kermicle's *mdr1* allele reduced the string of seven A's to six in the W22 amplicon subsequence

AGTTTACGAAAAAAAGTGCTTTC. Primers are as follows:

>W22-Ros1-F1

TTAGAACAAGGCGACTTTTCACC

>W22-Ros1-R1

AGGAACTCGGTAGCTCTTCT

To genotype the *dng102-Q235 allele*, A 569-bp region of *dng102* was amplified by PCR with

JIG-339 and JIG-340 primers and digested with Hpy188I (NEB # R0617S). This amplicon has

two Hpy188I sites in wild-type, but the mutation that converts Q to a stop codon at amino acid

position 235 removes one of them such that digestion of the wild-type amplicon yields three

fragments of 317, 156, and 96 bp, but the *dng102-Q235* allele yields two fragments of 473 and

96 bp. Primers are as follows:

>JIG-339

TCCAAAGCCTCCCAAAGAAA

>JIG-340

GCATATGATCGACCGAGCTG

To genotype the *dng102 mu1083641* allele, short regions were amplified using EoMumix3

primers along with either of two *dng102* primers, JIG-337 or JIG-338, which were also used

together to amplify the wild-type locus. Primers are as follows:

>EoMumix3

GCCTCYATTTCGTCGAATCCS

>JIG-337

CTGGCAGGAAAATGCAACCT

>JIG-338

GCAATTGCGCAAGCTTAATG

To genotype the Ds-GFP insertion *R153F01* linked to *Dng102*, two primer combinations were

used: the Ds-GFP primer JSR01dsR with the flanking primer JIG-368, and the Ds-GFP primer

JGp3dsL with the flanking primer JIG-369. JIG-368 and JIG-369 were also used as a pair to

amplify the wild-type locus. Primers are as follows:

>JSR01dsR

GGGTTCGAAATCGATCGGGATA

>JGp3dsL

ACCCGACCGGATCGTATCGG

>JIG-368

CGGAGTCATTCTTGCTCTTCT

>JIG-369

AGGACGAGACCTTAGTCCTTAT

Amino acid sequence comparison of maize DNA glycosylases and DEMETER.

To analyze the relationship between DEMETER (AT5G04560.2) and its maize

homologs, the longest putative isoform of each protein was selected. Pairwise alignments were

performed between DEMETER and the five maize homologs and between MDR1 and the

remaining four maize homologs. Alignments were performed using the MUSCLE algorithm

(Edgar 2004) in Geneious Prime v2020.2.4 (http://www.geneious.com). The percentage of

identical sites (%Id) and alignment length are reported. The five maize glycosylase amino acid

sequences are from the B73 v5 reference annotations: Zm00001eb202980_P001,

Zm00001eb241240_P002, Zm00001eb289030_P003, Zm00001eb241310_P001, and

Zm00001eb279940_P003 (Hufford et al. 2021).

Ds-GFP screening for mutation transmission

Stocks containing Ds-GFP alleles (tdsgR128F11) and (tdsgR153F01) were obtained from

the Maize Genetics Cooperation Stock Center. Double heterozygous plants (*R128F11/mdr1*

R153F01/dng102-Q235) were reciprocally crossed with wild-type Oh43 inbred plants that did

not carry Ds-GFP. Presence of Ds-GFP insertions in kernels was determined using a blue light

and orange filter glasses similar to previous methods, but on a smaller scale without automated

kernel counting (Warman et al. 2020). The *dng102-Q235* allele was derived from the TILLING

stock 04IA-B73PS-055_C4 in B73, which was then backcrossed into non-mutagenized B73,

selfed twice, then crossed three times into either W22 or W22-related stocks before crossing with

Ds-GFP stocks.

Preparation of EM-seq libraries

EM-seq libraries (NEBNext® Enzymatic Methyl-seq Kit #E7120S) were prepared from

three tissues. 1) Early developing 2^{nd} leaves were collected before the unfurling of the first

leaves. Three replicates each of the W22-related *mdr1* homozygous stock derived from X336J

and the W22-related wild-type stock X336K were included. Each replicate included leaves from

~5 to 9 plants. 2) Premeiotic tassels were collected from plants at approximately the 15-leaf

stage, when tassels were between 0.8 and 2.3 cm in length. Three replicates each of *mdr1*

homozygous and *Mdr1* homozygous siblings were included. The heterozygous parents that were

crossed to produce these plants were the product of a cross between the sequenced W22 inbred

stock and the W22-related *mdr1* homozygous stock derived from X336J. Each replicate included

two or three individual tassels. 3) Mature endosperm was separated from pericarp with forceps

after soaking in water for 20 minutes and separated from embryo by dissection with a razor

blade. Three replicates each of *mdr1* homozygous and *Mdr1* homozygous siblings were included.

The dissected kernels were siblings of the ones that were planted and grown for tassel collection.

Each replicate consisted of endosperm from a single kernel. DNA was extracted using an IBI

plant DNA extraction kit (#IB47231) after grinding to a fine powder (in liquid nitrogen for tassel

and leaf). The input for each EM-seq library consisted of 200 ng of genomic DNA that had been

combined with 1 pg of control pUC19 DNA and 20 pg of control lambda DNA and sonicated to

fragments averaging ~700 bp in length using a Diagenode Bioruptor. The protocol for large

insert libraries was followed. Libraries were amplified with 4 or 5 PCR cycles. Libraries were

Illumina sequenced using paired-end 150 nt reads. Read counts and accession numbers are listed

in Supplemental Table 2.

Preparation of small RNA sequencing libraries

The same tassel tissues used for preparation of EM-seq libraries were also used for

preparation of small RNA sequencing libraries. RNA was extracted using a mirVana miRNA

isolation kit (Thermo Fisher Scientific #AM1560) using the small RNA enrichment method.

Plant RNA Isolation Aid (Thermo Fisher Scientific #AM9690) was added at the lysis step. Small

RNA sequencing libraries were prepared using the NEXTflex Small RNA-Seq Kit v3

(PerkinElmer #5132-05) using the gel-free size selection & cleanup method. ~500 ng small RNA

enriched RNA was used as input for each library, and libraries were amplified with 13 cycles of

PCR. Read counts and accession numbers are listed in Supplemental Table 3.

Gene, transposon, and tandem repeat annotations

Gene annotations (Zm-W22-REFERENCE-NRGENE-2.0_Zm00004b.1.gff3.gz) and transposon

annotations (W22.allTE.withSolo_10Jan18.gff3.gz) were downloaded from

https://ftp.maizegdb.org/MaizeGDB/FTP/Zm-W22-REFERENCE-NRGENE-2.0/. The

transposon annotations are based on a conservative identification method using intact known

structural features (Springer et al. 2018; Anderson et al. 2019a). Retrotransposons having solo

LTRs were distinguished from those having both LTRs in these annotations by "Zm00004bS" in

their names. Superfamilies were identified by their three-letter codes, e.g., RLG for Gypsy. A

subset of genic analysis (Figure 1B and C, Figure 5E and F, and Supplemental Figure S5) made

use of a more stringently filtered set of gene annotations where genes were annotated in both the

W22 reference and B73 refgen_v4 (Jiao et al. 2017). These constituted 24,133 of the total 40,650

W22 genes. MEG and PEG status of genes in W22 were determined as previously described

(Anderson et al. 2021). To define the endosperm gene expression categories used in Figure 5C, a

B73 developmental expression atlas data (Stelpflug et al. 2016) was analyzed using similar

methods as previously (Anderson et al. 2019b). Genes that are constitutively lowly or not

expressed (not.expressed) genes have an RPM > 1 in fewer than 3 of the 225 total libraries

representing different tissues, developmental stages, and biological replicates. Endosperm

expressed (endo.expressed) and not endosperm expressed (not.endo) were identified from the

rest of the genes based on the following criteria: The "endo.expressed" genes have an RPM > 1

in at least 18 of 21 endosperm libraries. The "not.endo" genes have an RPM value of < 0.5 in at

least 20 of the 21 endosperm libraries. Finally, the W22 genes corresponding to the B73 genes

meeting these criteria were determined with a gene key, which can be found

at https://github.com/SNAnderson/Expression/blob/main/genes.expression.development.categori

es.20Sept21.txt. Locations of tandem repeats in the genome were determined by mapping

dimerized consensus sequences for *CentC*, *knob180*, and *TR-1* (Gent et al. 2017; Liu et al. 2020)

to the genome using Bowtie2 (Langmead and Salzberg 2012) (version 2.4.1) with the --local

parameter. All repeat loci that were within 50 bp of each other were merged with the BEDTools

merge tool with the -d 50 parameter. Following this, all loci less than 500 bp in length were

removed.

Methylome read processing, definition of regions by methylation status

EM-seq reads were trimmed of adapter sequence using cutadapt, parameters -q 20 -a

AGATCGGAAGAGC -A AGATCGGAAGAGC -O. Reads were aligned to each genome and

methylation values called using BS-Seeker2 (version 2.1.5), parameters -m 1 --aligner=bowtie2 -

X 1000 (Guo et al. 2013). To identify DMRs, CGmaps from each group of three individual

replicates were merged into single CGmaps using the merge2 tool of CGmapTools version 0.1.2

(Guo et al. 2018). Intersected CGmaps for each pair of merged CGmaps were produced using the

CGmapTools intersect tool in each sequence context separately (CG, CHG, and CHH). Single-bp

level methylation measurements in Figure 5C were produced using the CGmapTools lollipop

tool on the endosperm-tassel intersected CGmaps. CG and CHG DMRs and control eligible

regions were then identified from the separate CG and CHG intersected maps using

DMR_Finder.py (see Supplemental scripts), parameters as follows: binSize = 200, minCov = 3

minCountC = 5, minAbsDif = .20, minRelDif = .5. The set of CG DMRs that overlapped CHG

DMRs were identified using the BEDTools (version 2.29.2) intersect tool. To finalize the set of

DMRs, all DMRs located on scaffolds rather than chromosomes 1 through 10 were discarded.

RdDM regions were identified from the merged tassel CGmap using MR_Finder.py (see

Supplemental scripts), parameters as follows: binSize = 200, minCov = 3, minCountC = 5,

minMeth = .2, context1 = 'CHH', context2 = 'CHH'. CHG unmethylated regions (UMRs) were

identified from the merged tassel CGmap using UMR_Finder.py (see Supplemental scripts),

parameters as follows: binSize = 200, minCov = 3, minCountC = 5, maxMeth = .2, context1 =

'CHG', context2 = 'CHG'.

Characterization of DMRs in terms of methylation levels, overlap with specific genetic

elements, and sequence motifs

Methylation averages on 100-bp intervals centered on DMRs and control eligible regions were

produced using the CGmapTools mfg tool, requiring a minimum coverage of two with the -c 2

parameter. Prior to the mfg analysis, regions were first divided into two groups based on distance

to closest CHG UMR: greater than two Kb as heterochromatin, less than or equal to two Kb as

euchromatin. These distances were measured using the BEDTools closest tool with -t first -d

parameters. Regions were oriented relative to CHG UMRs such that the nearest CHG UMRs was

always in the downstream direction using region_UMR_orienter.py (see Supplemental scripts).

Orientation for regions that were equidistant from two CHG UMRs was randomized. DMR

methylation averages used for violin plots were obtained using the CGmapTools mtr tool. Only

DMRs with an effective coverage of at least 15 were included. Effective coverage is the number

of cytosines covered by reads in the region times the number of reads that cover each cytosine.

The numbers of overlaps between regions and different genetic elements, requiring at least 100

bp of overlap to count, were obtained using the BEDTools intersect tool with parameters -u -f .5.

To determine whether DMRs contained motifs associated with differential methylation, fasta

sequences were derived from bed files using the BEDTools getfasta tool. Three subsamples of

5000 sequences were input into STREME, each with a different seed number (Bailey 2021). All

motifs identified by STREME were input into FIMO to identify matches throughout the

complete sets of sequences. SpaMo (Whitington et al. 2011) was then used to search for

combinations of motifs that may work together to cause the differential methylation, but was also

inconclusive. AT richness was measured using the EMBOSS program infoseq (Rice et al. 2000).

Characterization of siRNA expression in methylated regions

Small RNA-seq reads were quality filtered, trimmed of adapters, and filtered for lengths

of 20-25 nt using cutadapt (Martin, 2011), parameters -q 20 -a

TGGAATTCTCGGGTGCCAAGG -e .05 -O 5 --discard-untrimmed -m 28 -M 33. The four

random nucleotides at each end were then removed using cutadapt with the -u 4 parameter

followed by cutadapt with the -u -4 parameter. Reads were aligned to the genome with BWA-

backtrack (version 0.7.15), parameters aln -t 8 -l 10. A set of miRNA loci were identified using

ShortStack software (Axtell 2013) (grep 'MIRNA=Y' from the ShortStack_D.gff3 output file).

All wild-type small RNA bam files from embryo, endosperm, and tassel were combined and

used as a single input file for ShortStack to identify miRNA loci. Also included in this

combination were *zmet2* and *zmet5* heterozygous samples SRR6425186, SRR6425229,

SRR6425204, and SRR6425207, which were not included in subsequent DMR analyses. To

produce the final sets of siRNAs, the BEDTools intersect tool with parameters -v -f .9 was used

to filter out reads that overlapped at least 90% of their length with ShortStack miRNA loci.

Multi-mapping reads were included in all analyses. siRNAs that overlapped by at least 90% of

their lengths with DMRs were identified using the BEDTools intersect tool with the -u -wa -f .9

parameters. The number of overlaps with DMRs was obtained using the BEDTools coverage

tool.

Accession numbers

All raw sequencing data generated in this study have been submitted to the NCBI BioProject

database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA759188.

Individual accession numbers for small RNAs are listed in Supplemental Table S2 and for DNA

methylation in Supplemental Table S3.

ACKNOWLEDGMENTS

We thank the Maize Genetics Cooperation Stock Center for providing maize stocks;

Akshay Nair, Ankush Sangra, and Mary Tindall Smith for genotyping stocks; and John Fowler

for advice on use of Ds-GFP insertions for measuring frequency of mutation transmission. This

study was supported in part by resources and technical expertise from the Georgia Advanced

Computing Resource Center, a partnership between the University of Georgia's Office of the

Vice President for Research and Office of the Vice President for Information Technology.

FUNDING

NSF 1744001
673

REFERENCES CITED

Anderson SN, Springer NM. 2018. Potential roles for transposable elements in creating
imprinted expression. *Curr Opin Genet Dev* **49**: 8-14.
Anderson SN, Stitzer MC, Brohammer AB, Zhou P, Noshay JM, O'Connor CH, Hirsch CD,
Ross-Ibarra J, Hirsch CN, Springer NM. 2019a. Transposable elements contribute to
dynamic genome content in maize. *Plant J* **100**: 1052-1065.
Anderson SN, Stitzer MC, Zhou P, Ross-Ibarra J, Hirsch CD, Springer NM. 2019b. Dynamic
Patterns of Transcript Abundance of Transposable Element Families in Maize. *G3*
(Bethesda) **9**: 3673-3682.
Anderson SN, Zhou P, Higgins K, Brandvain Y, Springer NM. 2021. Widespread imprinting of
transposable elements and variable genes in the maize endosperm. *PLoS Genet* **17**:
e1009491.
Axtell MJ. 2013. ShortStack: comprehensive annotation and quantification of small RNA genes.
RNA **19**: 740-751.
Bailey TL. 2021. STREME: Accurate and versatile sequence motif discovery. *Bioinformatics*
doi:10.1093/bioinformatics/btab203.
Batista RA, Köhler C. 2020. Genomic imprinting in plants-revisiting existing models. *Genes Dev*
34: 24-36.
Becraft PW, Gutierrez-Marcos J. 2012. Endosperm development: dynamic processes and cellular
innovations underlying sibling altruism. *Wiley Interdiscip Rev Dev Biol* **1**: 579-593.
Belmonte MF, Kirkbride RC, Stone SL, Pelletier JM, Bui AQ, Yeung EC, Hashimoto M, Fei J,
Harada CM, Munoz MD et al. 2013. Comprehensive developmental profiles of gene
activity in regions and subregions of the Arabidopsis seed. *Proc Natl Acad Sci U S A* **110**:
E435-444.
Borg M, Papareddy RK, Dombey R, Axelsson E, Nodine MD, Twell D, Berger F. 2021.
Epigenetic reprogramming rewires transcription during the alternation of generations in
Arabidopsis. *Elife* **10**.
Bouyer D, Kramdi A, Kassam M, Heese M, Schnittger A, Roudier F, Colot V. 2017. DNA
methylation dynamics during early plant life. *Genome Biol* **18**: 179.
Calarco JP, Borges F, Donoghue MT, Van Ex F, Jullien PE, Lopes T, Gardner R, Berger F, Feijó
JA, Becker JD et al. 2012. Reprogramming of DNA methylation in pollen guides
epigenetic inheritance via small RNA. *Cell* **151**: 194-205.
Chakraborty T, Kendall T, Grover JW, Mosher RA. 2021. Embryo CHH hypermethylation is
mediated by RdDM and is autonomously directed in Brassica rapa. *Genome Biol* **22**: 140.
Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, Jacobsen SE, Fischer RL.
2002. DEMETER, a DNA glycosylase domain protein, is required for endosperm gene
imprinting and seed viability in arabidopsis. *Cell* **110**: 33-42.
Creasey KM, Zhai J, Borges F, Van Ex F, Regulski M, Meyers BC, Martienssen RA. 2014.
miRNAs trigger widespread epigenetically activated siRNAs from transposons in
Arabidopsis. *Nature* **508**: 411-415.
Crisp PA, Marand AP, Noshay JM, Zhou P, Lu Z, Schmitz RJ, Springer NM. 2020. Stable
unmethylated DNA demarcates expressed genes and their cis-regulatory space in plant
genomes. *Proc Natl Acad Sci U S A* **117**: 23991-24000.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST et al. 2011. The variant call format and VCFtools. *Bioinformatics*
27: 2156-2158.
Doll NM, Just J, Brunaud V, Caïus J, Grimault A, Depège-Fargeix N, Esteban E, Pasha A,
Provart NJ, Ingram GC et al. 2020. Transcriptomics at Maize Embryo/Endosperm
Interfaces Identifies a Transcriptionally Distinct Endosperm Subdomain Adjacent to the
Embryo Scutellum. *Plant Cell* **32**: 833-852.
Dooner HK, Wang Q, Huang JT, Li Y, He L, Xiong W, Du C. 2019. Spontaneous mutations in
maize pollen are frequent in some lines and arise mainly from retrotranspositions and
deletions. *Proc Natl Acad Sci U S A* **116**: 10734-10743.
Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res* **32**: 1792-1797.
Frost JM, Kim MY, Park GT, Hsieh PH, Nakamura M, Lin SJH, Yoo H, Choi J, Ikeda Y,
Kinoshita T et al. 2018. FACT complex is required for DNA demethylation at
heterochromatin during reproduction in Arabidopsis. *Proc Natl Acad Sci U S A* **115**:
E4720-E4729.
Fu FF, Dawe RK, Gent JI. 2018. Loss of RNA-Directed DNA Methylation in Maize
Chromomethylase and DDM1-Type Nucleosome Remodeler Mutants. *Plant Cell* **30**:
1617-1627.
Garcia-Aguilar M, Michaud C, Leblanc O, Grimanelli D. 2010. Inactivation of a DNA
methylation pathway in maize reproductive organs results in apomixis-like phenotypes.
Plant Cell **22**: 3249-3267.
Gehring M, Bubb KL, Henikoff S. 2009. Extensive demethylation of repetitive elements during
seed development underlies gene imprinting. *Science* **324**: 1447-1451.
Gehring M, Huh JH, Hsieh TF, Penterman J, Choi Y, Harada JJ, Goldberg RB, Fischer RL.
2006. DEMETER DNA glycosylase establishes MEDEA polycomb gene self-imprinting
by allele-specific demethylation. *Cell* **124**: 495-506.
Gent JI, Madzima TF, Bader R, Kent MR, Zhang X, Stam M, McGinnis KM, Dawe RK. 2014.
Accessible DNA and Relative Depletion of H3K9me2 at Maize Loci Undergoing RNA-
Directed DNA Methylation. *Plant Cell* **26**: 4903-4917.
Gent JI, Wang N, Dawe RK. 2017. Stable centromere positioning in diverse sequence contexts of
complex and satellite centromeres of maize and wild relatives. *Genome Biol* **18**: 121.
Guo W, Fiziev P, Yan W, Cokus S, Sun X, Zhang MQ, Chen PY, Pellegrini M. 2013. BS-
Seeker2: a versatile aligning pipeline for bisulfite sequencing data. *BMC Genomics* **14**:
774.
Guo W, Zhu P, Pellegrini M, Zhang MQ, Wang X, Ni Z. 2018. CGmapTools improves the
precision of heterozygous SNV calls and supports allele-specific methylation detection
and visualization in bisulfite-sequencing data. *Bioinformatics* **34**: 381-387.
Hatorangan MR, Laenen B, Steige KA, Slotte T, Köhler C. 2016. Rapid Evolution of Genomic
Imprinting in Two Species of the Brassicaceae. *Plant Cell* **28**: 1815-1827.
He S, Vickers M, Zhang J, Feng X. 2019. Natural depletion of histone H1 in sex cells causes
DNA demethylation, heterochromatin decondensation and transposon activation. *Elife* **8**.
Hufford MB, Seetharam AS, Woodhouse MR, Chougule KM, Ou S, Liu J, Ricci WA, Guo T,
Olson A, Qiu Y et al. 2021. De novo assembly, annotation, and comparative analysis of
26 diverse maize genomes. *Science* **373**: 655-662.

Ibarra CA, Feng X, Schoft VK, Hsieh TF, Uzawa R, Rodrigues JA, Zemach A, Chumak N,
Machlicova A, Nishimura T et al. 2012. Active DNA demethylation in plant companion
cells reinforces transposon methylation in gametes. *Science* **337**: 1360-1364.
Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X, Chin CS
et al. 2017. Improved maize reference genome with single-molecule technologies. *Nature*
546: 524-527.
Jullien PE, Susaki D, Yelagandula R, Higashiyama T, Berger F. 2012. DNA methylation
dynamics during sexual reproduction in Arabidopsis thaliana. *Curr Biol* **22**: 1825-1830.
Kapazoglou A, Drosou V, Argiriou A, Tsaftaris AS. 2013. The study of a barley epigenetic
regulator, HvDME, in seed development and under drought. *BMC Plant Biol* **13**: 172.
Kawakatsu T, Nery JR, Castanon R, Ecker JR. 2017. Dynamic DNA methylation reconfiguration
during seed development and germination. *Genome Biol* **18**: 171.
Kermicle J. 1995. Location, time of action, and dominance of an imprintor of *R*-mottled maize.
In *Modification of gene expression and non-mendelian inheritance*, (ed. Oono,
Takiawa). NIAR, Japan.
Kermicle JL. 1970. Dependence of the R-mottled aleurone phenotype in maize on mode of
sexual transmission. *Genetics* **66**: 69-85.
Khouider S, Borges F, LeBlanc C, Ungru A, Schnittger A, Martienssen R, Colot V, Bouyer D.
2021. Male fertility in Arabidopsis requires active DNA demethylation of genes that
control pollen tube function. *Nat Commun* **12**: 410.
Kim MY, Ono A, Scholten S, Kinoshita T, Zilberman D, Okamoto T, Fischer RL. 2019. DNA
demethylation by ROS1a in rice vegetative cells promotes methylation in sperm. *Proc*
Natl Acad Sci U S A doi:10.1073/pnas.1821435116.
Kim S, Park JS, Lee J, Lee KK, Park OS, Choi HS, Seo PJ, Cho HT, Frost JM, Fischer RL et al.
2021. The DME demethylase regulates sporophyte gene expression, cell proliferation,
differentiation, and meristem resurrection. *Proc Natl Acad Sci U S A* **118**.
Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:
357-359.
Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**: 1754-1760.
Li Y, Segal G, Wang Q, Dooner HK. 2013. Gene tagging with engineered Ds elements in maize.
Methods Mol Biol **1057**: 83-99.
Liu J, Seetharam AS, Chougule K, Ou S, Swentowsky KW, Gent JI, Llaca V, Woodhouse MR,
Manchanda N, Presting GG et al. 2020. Gapless assembly of maize chromosomes using
long-read technologies. *Genome Biol* **21**: 121.
Lu X, Liu J, Ren W, Yang Q, Chai Z, Chen R, Wang L, Zhao J, Lang Z, Wang H et al. 2018.
Gene-Indexed Mutations in Maize. *Mol Plant* **11**: 496-504.
Martin M. 2011. Cutadapt Removes Adapter Sequences From High-Throughput Sequencing
Reads. *EMBnetjournal* **17**: 3.
Martínez G, Panda K, Köhler C, Slotkin RK. 2016. Silencing in sperm cells is directed by RNA
movement from the surrounding nurse cell. *Nat Plants* **2**: 16030.
McCarty DR, Settles AM, Suzuki M, Tan BC, Latshaw S, Porch T, Robin K, Baier J, Avigne W,
Lai J et al. 2005. Steady-state transposon mutagenesis in inbred maize. *Plant J* **44**: 52-61.
McCue AD, Panda K, Nuthikattu S, Choudury SG, Thomas EN, Slotkin RK. 2015.
ARGONAUTE 6 bridges transposable element mRNA-derived siRNAs to the
establishment of DNA methylation. *EMBO J* **34**: 20-35.

Moreno-Romero J, Jiang H, Santos-González J, Köhler C. 2016. Parental epigenetic asymmetry
of PRC2-mediated histone modifications in the Arabidopsis endosperm. *EMBO J* **35**:
1298-1311.
O'Malley RC, Huang SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A,
Ecker JR. 2016. Cistrome and Epicistrome Features Shape the Regulatory DNA
Landscape. *Cell* **165**: 1280-1292.
Oka R, Zicola J, Weber B, Anderson SN, Hodgman C, Gent JI, Wesselink JJ, Springer NM,
Hoefsloot HCJ, Turck F et al. 2017. Genome-wide mapping of transcriptional enhancer
candidates using DNA and chromatin features in maize. *Genome Biol* **18**: 137.
Parent JS, Cahn J, Herridge RP, Grimanelli D, Martienssen RA. 2021. Small RNAs guide
histone methylation in. *Genes Dev* **35**: 841-846.
Park JS, Frost JM, Park K, Ohr H, Park GT, Kim S, Eom H, Lee I, Brooks JS, Fischer RL et al.
2017. Control of DEMETER DNA demethylase gene transcription in male and female
gamete companion cells in. *Proc Natl Acad Sci U S A* **114**: 2078-2083.
Park K, Kim MY, Vickers M, Park JS, Hyun Y, Okamoto T, Zilberman D, Fischer RL, Feng X,
Choi Y et al. 2016. DNA demethylation is initiated in the central cells of Arabidopsis and
rice. *Proc Natl Acad Sci U S A* **113**: 15138-15143.
Pignatta D, Erdmann RM, Scheer E, Picard CL, Bell GW, Gehring M. 2014. Natural epigenetic
polymorphisms lead to intraspecific variation in Arabidopsis gene imprinting. *Elife* **3**:
e03198.
Pignatta D, Novitzky K, Satyaki PRV, Gehring M. 2018. A variably imprinted epiallele impacts
seed development. *PLoS Genet* **14**: e1007469.
Pillot M, Baroux C, Vazquez MA, Autran D, Leblanc O, Vielle-Calzada JP, Grossniklaus U,
Grimanelli D. 2010. Embryo and endosperm inherit distinct chromatin and transcriptional
states from the female gametes in Arabidopsis. *Plant Cell* **22**: 307-320.
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841-842.
Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European Molecular Biology Open Software
Suite. *Trends Genet* **16**: 276-277.
Rodrigues JA, Hsieh PH, Ruan D, Nishimura T, Sharma MK, Sharma R, Ye X, Nguyen ND,
Nijjar S, Ronald PC et al. 2021. Divergence among rice cultivars reveals roles for
transposition and epimutation in ongoing evolution of genomic imprinting. *Proc Natl
Acad Sci U S A* **118**.
Schoft VK, Chumak N, Choi Y, Hannon M, Garcia-Aguilar M, Machlicova A, Slusarz L,
Mosiolek M, Park JS, Park GT et al. 2011. Function of the DEMETER DNA glycosylase
in the Arabidopsis thaliana male gametophyte. *Proc Natl Acad Sci U S A* **108**: 8042-8047.
Schumann U, Lee J, Kazan K, Ayliffe M, Wang MB. 2017. DNA-Demethylase Regulated Genes
Show Methylation-Independent Spatiotemporal Expression Patterns. *Front Plant Sci* **8**:
1449.
Schumann U, Lee JM, Smith NA, Zhong C, Zhu JK, Dennis ES, Millar AA, Wang MB. 2019.
DEMETER plays a role in DNA demethylation and disease response in somatic tissues of
Arabidopsis. *Epigenetics* **14**: 1074-1087.
Springer NM, Anderson SN, Andorf CM, Ahern KR, Bai F, Barad O, Barbazuk WB, Bass HW,
Baruch K, Ben-Zvi G et al. 2018. The maize W22 genome provides a foundation for
functional genomics and transposon biology. *Nat Genet* **50**: 1282-1288.

Stelpflug SC, Sekhon RS, Vaillancourt B, Hirsch CN, Buell CR, de Leon N, Kaeppler SM. 2016.
An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to
Explore Root Development. *Plant Genome* **9**.
Till BJ, Reynolds SH, Weil C, Springer N, Burtner C, Young K, Bowers E, Codomo CA, Enns
LC, Odden AR et al. 2004. Discovery of induced point mutations in maize genes by
TILLING. *BMC Plant Biol* **4**: 12.
Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan
T, Shakir K, Roazen D, Thibault J et al. 2013. From FastQ data to high confidence
variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr Protoc*
Bioinformatics **43**: 11.10.11-11.10.33.
Warman C, Panda K, Vejlupkova Z, Hokin S, Unger-Wallace E, Cole RA, Chettoor AM, Jiang
D, Vollbrecht E, Evans MMS et al. 2020. High expression in maize pollen correlates with
genetic contributions to pollen fitness as well as with coordinated transcription from
neighboring transposable elements. *PLoS Genet* **16**: e1008462.
Waters AJ, Bilinski P, Eichten SR, Vaughn MW, Ross-Ibarra J, Gehring M, Springer NM. 2013.
Comprehensive analysis of imprinted genes in maize reveals allelic variation for
imprinting and limited conservation with other species. *Proc Natl Acad Sci U S A* **110**:
19639-19644.
Waters AJ, Makarevitch I, Eichten SR, Swanson-Wagner RA, Yeh CT, Xu W, Schnable PS,
Vaughn MW, Gehring M, Springer NM. 2011. Parent-of-origin effects on gene
expression and DNA methylation in the maize endosperm. *Plant Cell* **23**: 4221-4233.
Whitington T, Frith MC, Johnson J, Bailey TL. 2011. Inferring transcription factor complexes
from ChIP-seq data. *Nucleic Acids Res* **39**: e98.
Xiao W, Gehring M, Choi Y, Margossian L, Pu H, Harada JJ, Goldberg RB, Pennell RI, Fischer
RL. 2003. Imprinting of the MEA Polycomb gene is controlled by antagonism between
MET1 methyltransferase and DME glycosylase. *Dev Cell* **5**: 891-901.
Yamamuro C, Miki D, Zheng Z, Ma J, Wang J, Yang Z, Dong J, Zhu JK. 2014. Overproduction
of stomatal lineage cells in Arabidopsis mutants defective in active DNA demethylation.
Nat Commun **5**: 4062.
Yelagandula R, Stroud H, Holec S, Zhou K, Feng S, Zhong X, Muthurajan UM, Nie X,
Kawashima T, Groth M et al. 2014. The histone variant H2A.W defines heterochromatin
and promotes chromatin condensation in Arabidopsis. *Cell* **158**: 98-109.
Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, Harmer SL,
Zilberman D. 2013. The Arabidopsis nucleosome remodeler DDM1 allows DNA
methyltransferases to access H1-containing heterochromatin. *Cell* **153**: 193-205.
Zhang C, Hung YH, Rim HJ, Zhang D, Frost JM, Shin H, Jang H, Liu F, Xiao W, Iyer LM et al.
2019. The catalytic core of DEMETER guides active DNA demethylation in Arabidopsis.
Proc Natl Acad Sci U S A **116**: 17563-17571.
Zhang M, Xie S, Dong X, Zhao X, Zeng B, Chen J, Li H, Yang W, Zhao H, Wang G et al. 2014.
Genome-wide high resolution parental-specific DNA and histone methylation maps
uncover patterns of imprinting regulation in maize. *Genome Res* **24**: 167-176.
Zhang M, Zhao H, Xie S, Chen J, Xu Y, Wang K, Guan H, Hu X, Jiao Y, Song W et al. 2011.
Extensive, clustered parental imprinting of protein-coding and noncoding RNAs in
developing maize endosperm. *Proc Natl Acad Sci U S A* **108**: 20042-20047.
905

Figure 1

Figure 1 Maternal *R1* expression phenotype and endosperm DNA glycosylase alleles
A, Kermicle's *mdr1* allele produces a maternal R mottling phenotype. In crosses where functional *R1* is inherited maternally only, maternal *Mdr1* is required for full *R1* expression. W22 (*mdr1*) is the *mdr1* mutant in a full-color W22 background. All listed alleles are homozygous.
B, Sequence and structure of *mdr1* and *dng102*. Nucleotides highlighted in red indicate positions of mutations. Asterisks in black background indicate positions of stop codons. Black triangle indicates position of Mu insertion.

Figure 2

Figure 2 Genomic locations of endosperm DMRs

A, The majority of differentially methylated regions are hypomethylated in endosperm. Eligible regions are defined in the same way as DMRs in terms of read coverage and numbers of informative cytosines, but with no requirement for differential methylation. Endosperm is from mature kernels segregating homozygous wild-type and homozygous *mdr1* mutant. Tassel is premeiotic stage, ~1.5 cm in length, segregating homozygous wild-type and homozygous *mdr1* mutant. Leaf is 2nd leaf prior to emergence through 1st leaf. The first three comparisons use the same wild-type endosperm. The fourth uses the same wild-type tassel as the endosperm vs. tassel comparison, but compares homozygous wild-type tassel to sibling *mdr1* tassel.

B, Endosperm DMRs are depleted from Gypsy and Copia and weakly enriched in CACTA, Mutator, and Helitron transposon superfamilies. DMRs include only the hypoDMRs of **A**. Enrichment is defined as the number of overlaps between DMRs and different genetic normalized by the number of overlaps between eligible regions and the same genetic elements. RdDM regions are defined by average mCHH values of greater than 0.2 in tassel.

C, Endosperm DMRs are weakly enriched near genes. Enrichment values on the Y-axis are the sum of the DMR coverage or RdDM region coverage normalized by eligible region coverage of each 100-bp interval for each gene up to 3 Kb outside the genes and 2 Kb inside. TSS: transcription start site. polyA: polyadenylation site. Tick marks on the X-axis are on 1-Kb intervals.

D, Endosperm DMRs are strongly depleted in the Gypsy superfamily, as in C, except using edges of intact Gypsy retrotransposons rather than genes.

Figure 3

Figure 3 Mutation of *mdr1* partially reverts endosperm to typical methylation profiles
A, mCG profiles centered on endosperm vs. tassel DMRs. Methylation values are averages of 100-bp intervals within and up to 3 Kb on either side of DMRs. Orientation is determined by direction to the closest CHG unmethylated region in tassel (UMR, defined as mCHG < .2). Values for tassel and mature endosperm are from the same methylomes as described in Figure 2A. Values for developing endosperm and developing embryo are from previously published methylomes (Fu et al. 2018).
B, mCHG profiles centered on endosperm vs. tassel DMRs, as in A.
C, mCHH profiles centered on endosperm vs. tassel DMRs, as in A.
D, mCG profiles centered on endosperm vs. tassel eligible regions, as in A.
E, mCHG profiles centered on endosperm vs. tassel eligible regions, as in A.
F, mCHH profiles centered on endosperm vs. tassel eligible regions, as in A

Figure 4

Figure 4 Endosperm DMRs are depleted of siRNAs relative to RdDM regions.
A, siRNAs length and abundance profiles. The normalized abundance of siRNAs from 20 to 25 nt in length are shown for each type of region. DMRs are from endosperm vs. tassel comparison. Heterochromatin DMRs (10014 total) are ones that are more than 2 Kb from the nearest CHG UMR. Euchromatin DMRs (10142 total) are ones that are 2 Kb or less from the nearest CHG UMR. These number include multimapping siRNAs, where one position is assigned at random among multiple equally valid positions. Developing tassel siRNAs are from the same tissue samples used for tassel methylomes. Developing embryo and developing endosperm siRNAs are from the same tissue types as their respective methylomes (Fu et al. 2018).
B, Frequency of 21-nt siRNA regions. Any region with at least 50 nt spanned by mapped siRNAs (25% of the region length) is defined as an siRNA region. These number include multimapping siRNAs, where one position is assigned at random among multiple equally valid positions. Developing tassel siRNAs are from the same tissue samples used for tassel methylomes. Developing embryo and developing endosperm siRNAs are from the same tissue types as their respective methylomes (Fu et al. 2018).
C, Frequency of 22-nt siRNA regions, as in **B**.
D, Frequency of 24-nt siRNA regions, as in **B**.

Figure 5

Figure 5 Endosperm DMRs are enriched in maternally-imprinted genes and transposons, but the majority of endosperm DMR genes are biparentally expressed

A, Genes that overlap endosperm DMRs are skewed toward maternal expression but still center on RER values of 0.67. RER is reciprocal expression ratio, where a value of 1 means completely maternal and 0 means completely paternal. Expression data include all genes, not just syntenic ones, from (Anderson et al. 2021). Genes are classified according to distance from endosperm DMRs. Endosperm DMRs from each of the three comparisons are shown separately. Boxes are interquartile ranges.

B, MEGs have a high frequency of overlap with endosperm DMRs. Genes are classified according distance from endosperm DMRs (by color) and according to imprinting status (X-axis labels). Y-axis indicates the proportion of all genes in each set. MEGs are defined as an RER value greater than 0.9 (117 genes), PEGs as less than 0.1 (69 genes), and not-imprinted as all intermediate values (14,450 genes).

C, DMRs do not predict endosperm gene expression. Genes are classified according distance from endosperm DMRs (by color) and by expression in endosperm (X-axis labels). Expression categories as follows: "expressed.endo" means expressed in endosperm, without regard to expression in other tissues (12,416 genes); "not.endo" means low or no expression in endosperm, but expressed in other tissues (5,019 genes); and "not expressed" means low or no expression in endosperm and other tissues (1,615 genes).

D, An example methylation profile for a MEG. mCG values are shown at single-base level resolution over a 10-Kb region of chromosome 10 including the gene *myb255* (*GRMZM2G150680*). Both height and color of lollypops indicates mCG values (darkest blue = zero mCG).

E, MEGs are enriched for endosperm DMRs near their 5' ends. DMR enrichment is the number of endosperm DMRs that overlap with a 2-Kb range either centered on transcription start sites (TSS) or polyadenylation sites (polyA) of MEGs normalized by the number of eligible regions that overlap the same range. Endosperm DMRs from each of the three comparisons are shown separately. The absence of an orange bar in WT vs. *mdr1* is because there were no overlaps in the polyA-centered range.

F, Genes are not enriched for endosperm DMRs near their 5' ends. Same as in **D**.

G, Maternal imprinted transposons have a high frequency of overlap with endosperm DMRs. MEG-type transposons (95 total) have an RER value greater than 0.9, PEG as less than 0.1, and non-imprinted (1773 total) as all intermediate values. The Y-axis indicates the proportion of transposons that overlap endosperm DMRs from each of the three comparisons. In all three, MEG-type transposons are significantly enriched for DMRs over non-imprinted ones (p-value < 0.01, binomial test). Only one PEG-type transposon overlapped with DMRs and is not shown.

Parsed Citations

Anderson SN, Springer NM. 2018. Potential roles for transposable elements in creating imprinted expression. Curr Opin Genet Dev 49: 8-14.
  Google Scholar: Author Only Title Only Author and Title

Anderson SN, Stitzer MC, Brohammer AB, Zhou P, Noshay JM, O'Connor CH, Hirsch CD, Ross-Ibarra J, Hirsch CN, Springer NM. 2019a. Transposable elements contribute to dynamic genome content in maize. Plant J 100: 1052-1065.
  Google Scholar: Author Only Title Only Author and Title

Anderson SN, Stitzer MC, Zhou P, Ross-Ibarra J, Hirsch CD, Springer NM. 2019b. Dynamic Patterns of Transcript Abundance of Transposable Element Families in Maize. G3 (Bethesda) 9: 3673-3682.
  Google Scholar: Author Only Title Only Author and Title

Anderson SN, Zhou P, Higgins K, Brandvain Y, Springer NM. 2021. Widespread imprinting of transposable elements and variable genes in the maize endosperm. PLoS Genet 17: e1009491.
  Google Scholar: Author Only Title Only Author and Title

Axtell MJ. 2013. ShortStack: comprehensive annotation and quantification of small RNA genes. RNA 19: 740-751.
  Google Scholar: Author Only Title Only Author and Title

Bailey TL. 2021. STREME: Accurate and versatile sequence motif discovery. Bioinformatics doi:10.1093/bioinformatics/btab203.
  Google Scholar: Author Only Title Only Author and Title

Batista RA, Köhler C. 2020. Genomic imprinting in plants-revisiting existing models. Genes Dev 34: 24-36.
  Google Scholar: Author Only Title Only Author and Title

Becraft PW, Gutierrez-Marcos J. 2012. Endosperm development: dynamic processes and cellular innovations underlying sibling altruism. Wiley Interdiscip Rev Dev Biol 1: 579-593.
  Google Scholar: Author Only Title Only Author and Title

Belmonte MF, Kirkbride RC, Stone SL, Pelletier JM, Bui AQ, Yeung EC, Hashimoto M, Fei J, Harada CM, Munoz MD et al. 2013. Comprehensive developmental profiles of gene activity in regions and subregions of the Arabidopsis seed. Proc Natl Acad Sci U S A 110: E435-444.
  Google Scholar: Author Only Title Only Author and Title

Borg M, Papareddy RK, Dombey R, Axelsson E, Nodine MD, Twell D, Berger F. 2021. Epigenetic reprogramming rewires transcription during the alternation of generations in Arabidopsis. Elife 10.
  Google Scholar: Author Only Title Only Author and Title

Bouyer D, Kramdi A, Kassam M, Heese M, Schnittger A, Roudier F, Colot V. 2017. DNA methylation dynamics during early plant life. Genome Biol 18: 179.
  Google Scholar: Author Only Title Only Author and Title

Calarco JP, Borges F, Donoghue MT, Van Ex F, Jullien PE, Lopes T, Gardner R, Berger F, Feijó JA, Becker JD et al. 2012. Reprogramming of DNA methylation in pollen guides epigenetic inheritance via small RNA. Cell 151: 194-205.
  Google Scholar: Author Only Title Only Author and Title

Chakraborty T, Kendall T, Grover JW, Mosher RA. 2021. Embryo CHH hypermethylation is mediated by RdDM and is autonomously directed in Brassica rapa. Genome Biol 22: 140.
  Google Scholar: Author Only Title Only Author and Title

Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, Jacobsen SE, Fischer RL. 2002. DEMETER, a DNA glycosylase domain protein, is required for endosperm gene imprinting and seed viability in arabidopsis. Cell 110: 33-42.
  Google Scholar: Author Only Title Only Author and Title

Creasey KM, Zhai J, Borges F, Van Ex F, Regulski M, Meyers BC, Martienssen RA. 2014. miRNAs trigger widespread epigenetically activated siRNAs from transposons in Arabidopsis. Nature 508: 411-415.
  Google Scholar: Author Only Title Only Author and Title

Crisp PA, Marand AP, Noshay JM, Zhou P, Lu Z, Schmitz RJ, Springer NM. 2020. Stable unmethylated DNA demarcates expressed genes and their cis-regulatory space in plant genomes. Proc Natl Acad Sci U S A 117: 23991-24000.
  Google Scholar: Author Only Title Only Author and Title

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST et al. 2011. The variant call format and VCFtools. Bioinformatics 27: 2156-2158.
  Google Scholar: Author Only Title Only Author and Title

Doll NM, Just J, Brunaud V, Caïus J, Grimault A, Depège-Fargeix N, Esteban E, Pasha A, Provart NJ, Ingram GC et al. 2020. Transcriptomics at Maize Embryo/Endosperm Interfaces Identifies a Transcriptionally Distinct Endosperm Subdomain Adjacent to the Embryo Scutellum. Plant Cell 32: 833-852.
  Google Scholar: Author Only Title Only Author and Title

Dooner HK, Wang Q, Huang JT, Li Y, He L, Xiong W, Du C. 2019. Spontaneous mutations in maize pollen are frequent in some lines and arise mainly from retrotranspositions and deletions. Proc Natl Acad Sci U S A 116: 10734-10743.
  Google Scholar: Author Only Title Only Author and Title

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32: 1792-1797.
  Google Scholar: Author Only Title Only Author and Title

Frost JM, Kim MY, Park GT, Hsieh PH, Nakamura M, Lin SJH, Yoo H, Choi J, Ikeda Y, Kinoshita T et al. 2018. FACT complex is required for DNA demethylation at heterochromatin during reproduction in Arabidopsis. Proc Natl Acad Sci U S A 115: E4720-E4729.
  Google Scholar: Author Only Title Only Author and Title

Fu FF, Dawe RK, Gent JI. 2018. Loss of RNA-Directed DNA Methylation in Maize Chromomethylase and DDM1-Type Nucleosome Remodeler Mutants. Plant Cell 30: 1617-1627.
  Google Scholar: Author Only Title Only Author and Title

Garcia-Aguilar M, Michaud C, Leblanc O, Grimanelli D. 2010. Inactivation of a DNA methylation pathway in maize reproductive organs results in apomixis-like phenotypes. Plant Cell 22: 3249-3267.
  Google Scholar: Author Only Title Only Author and Title

Gehring M, Bubb KL, Henikoff S. 2009. Extensive demethylation of repetitive elements during seed development underlies gene imprinting. Science 324: 1447-1451.
  Google Scholar: Author Only Title Only Author and Title

Gehring M, Huh JH, Hsieh TF, Penterman J, Choi Y, Harada JJ, Goldberg RB, Fischer RL. 2006. DEMETER DNA glycosylase establishes MEDEA polycomb gene self-imprinting by allele-specific demethylation. Cell 124: 495-506.
  Google Scholar: Author Only Title Only Author and Title

Gent JI, Madzima TF, Bader R, Kent MR, Zhang X, Stam M, McGinnis KM, Dawe RK. 2014. Accessible DNA and Relative Depletion of H3K9me2 at Maize Loci Undergoing RNA-Directed DNA Methylation. Plant Cell 26: 4903-4917.
  Google Scholar: Author Only Title Only Author and Title

Gent JI, Wang N, Dawe RK. 2017. Stable centromere positioning in diverse sequence contexts of complex and satellite centromeres of maize and wild relatives. Genome Biol 18: 121.
  Google Scholar: Author Only Title Only Author and Title

Guo W, Fiziev P, Yan W, Cokus S, Sun X, Zhang MQ, Chen PY, Pellegrini M. 2013. BS-Seeker2: a versatile aligning pipeline for bisulfite sequencing data. BMC Genomics 14: 774.
  Google Scholar: Author Only Title Only Author and Title

Guo W, Zhu P, Pellegrini M, Zhang MQ, Wang X, Ni Z. 2018. CGmapTools improves the precision of heterozygous SNV calls and supports allele-specific methylation detection and visualization in bisulfite-sequencing data. Bioinformatics 34: 381-387.
  Google Scholar: Author Only Title Only Author and Title

Hatorangan MR, Laenen B, Steige KA, Slotte T, Köhler C. 2016. Rapid Evolution of Genomic Imprinting in Two Species of the Brassicaceae. Plant Cell 28: 1815-1827.
  Google Scholar: Author Only Title Only Author and Title

He S, Vickers M, Zhang J, Feng X. 2019. Natural depletion of histone H1 in sex cells causes DNA demethylation, heterochromatin decondensation and transposon activation. Elife 8.
  Google Scholar: Author Only Title Only Author and Title

Hufford MB, Seetharam AS, Woodhouse MR, Chougule KM, Ou S, Liu J, Ricci WA, Guo T, Olson A, Qiu Y et al. 2021. De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes. Science 373: 655-662.
  Google Scholar: Author Only Title Only Author and Title

Ibarra CA, Feng X, Schoft VK, Hsieh TF, Uzawa R, Rodrigues JA, Zemach A, Chumak N, Machlicova A, Nishimura T et al. 2012. Active DNA demethylation in plant companion cells reinforces transposon methylation in gametes. Science 337: 1360-1364.
  Google Scholar: Author Only Title Only Author and Title

Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X, Chin CS et al. 2017. Improved maize reference genome with single-molecule technologies. Nature 546: 524-527.
  Google Scholar: Author Only Title Only Author and Title

Jullien PE, Susaki D, Yelagandula R, Higashiyama T, Berger F. 2012. DNA methylation dynamics during sexual reproduction in Arabidopsis thaliana. Curr Biol 22: 1825-1830.
  Google Scholar: Author Only Title Only Author and Title

Kapazoglou A, Drosou V, Argiriou A, Tsaftaris AS. 2013. The study of a barley epigenetic regulator, HvDME, in seed development and under drought. BMC Plant Biol 13: 172.
  Google Scholar: Author Only Title Only Author and Title

Kawakatsu T, Nery JR, Castanon R, Ecker JR. 2017. Dynamic DNA methylation reconfiguration during seed development and germination. Genome Biol 18: 171.
  Google Scholar: Author Only Title Only Author and Title

Kermicle J. 1995. Location, time of action, and dominance of an imprintor of R-mottled maize. In Modification of gene expression and non-mendelian inheritance, (ed. Oono, Takiawa). NIAR, Japan.
  Google Scholar: Author Only Title Only Author and Title

Kermicle JL. 1970. Dependence of the R-mottled aleurone phenotype in maize on mode of sexual transmission. Genetics 66: 69-85.
  Google Scholar: Author Only Title Only Author and Title

Khouider S, Borges F, LeBlanc C, Ungru A, Schnittger A, Martienssen R, Colot V, Bouyer D. 2021. Male fertility in Arabidopsis requires active DNA demethylation of genes that control pollen tube function. Nat Commun 12: 410.
  Google Scholar: Author Only Title Only Author and Title

Kim MY, Ono A, Scholten S, Kinoshita T, Zilberman D, Okamoto T, Fischer RL. 2019. DNA demethylation by ROS1a in rice vegetative cells promotes methylation in sperm. Proc Natl Acad Sci U S A doi:10.1073/pnas.1821435116.
  Google Scholar: Author Only Title Only Author and Title

Kim S, Park JS, Lee J, Lee KK, Park OS, Choi HS, Seo PJ, Cho HT, Frost JM, Fischer RL et al. 2021. The DME demethylase regulates sporophyte gene expression, cell proliferation, differentiation, and meristem resurrection. Proc Natl Acad Sci U S A 118.
  Google Scholar: Author Only Title Only Author and Title

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9: 357-359.
  Google Scholar: Author Only Title Only Author and Title

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25: 1754-1760.
  Google Scholar: Author Only Title Only Author and Title

Li Y, Segal G, Wang Q, Dooner HK. 2013. Gene tagging with engineered Ds elements in maize. Methods Mol Biol 1057: 83-99.
  Google Scholar: Author Only Title Only Author and Title

Liu J, Seetharam AS, Chougule K, Ou S, Swentowsky KW, Gent JI, Llaca V, Woodhouse MR, Manchanda N, Presting GG et al. 2020. Gapless assembly of maize chromosomes using long-read technologies. Genome Biol 21: 121.
  Google Scholar: Author Only Title Only Author and Title

Lu X, Liu J, Ren W, Yang Q, Chai Z, Chen R, Wang L, Zhao J, Lang Z, Wang H et al. 2018. Gene-Indexed Mutations in Maize. Mol Plant 11: 496-504.
  Google Scholar: Author Only Title Only Author and Title

Martin M. 2011. Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads. EMBnetjournal 17: 3.
  Google Scholar: Author Only Title Only Author and Title

Martínez G, Panda K, Köhler C, Slotkin RK. 2016. Silencing in sperm cells is directed by RNA movement from the surrounding nurse cell. Nat Plants 2: 16030.
  Google Scholar: Author Only Title Only Author and Title

McCarty DR, Settles AM, Suzuki M, Tan BC, Latshaw S, Porch T, Robin K, Baier J, Avigne W, Lai J et al. 2005. Steady-state transposon mutagenesis in inbred maize. Plant J 44: 52-61.
  Google Scholar: Author Only Title Only Author and Title

McCue AD, Panda K, Nuthikattu S, Choudury SG, Thomas EN, Slotkin RK. 2015. ARGONAUTE 6 bridges transposable element mRNA-derived siRNAs to the establishment of DNA methylation. EMBO J 34: 20-35.
  Google Scholar: Author Only Title Only Author and Title

Moreno-Romero J, Jiang H, Santos-González J, Köhler C. 2016. Parental epigenetic asymmetry of PRC2-mediated histone modifications in the Arabidopsis endosperm. EMBO J 35: 1298-1311.
  Google Scholar: Author Only Title Only Author and Title

O'Malley RC, Huang SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR. 2016. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. Cell 165: 1280-1292.
  Google Scholar: Author Only Title Only Author and Title

Oka R, Zicola J, Weber B, Anderson SN, Hodgman C, Gent JI, Wesselink JJ, Springer NM, Hoefsloot HCJ, Turck F et al. 2017. Genome-wide mapping of transcriptional enhancer candidates using DNA and chromatin features in maize. Genome Biol 18: 137.
  Google Scholar: Author Only Title Only Author and Title

Parent JS, Cahn J, Herridge RP, Grimanelli D, Martienssen RA. 2021. Small RNAs guide histone methylation in. Genes Dev 35: 841-846.
  Google Scholar: Author Only Title Only Author and Title

Park JS, Frost JM, Park K, Ohr H, Park GT, Kim S, Eom H, Lee I, Brooks JS, Fischer RL et al. 2017. Control of DEMETER DNA demethylase gene transcription in male and female gamete companion cells in. Proc Natl Acad Sci U S A 114: 2078-2083.
  Google Scholar: Author Only Title Only Author and Title

Park K, Kim MY, Vickers M, Park JS, Hyun Y, Okamoto T, Zilberman D, Fischer RL, Feng X, Choi Y et al. 2016. DNA demethylation is initiated in the central cells of Arabidopsis and rice. Proc Natl Acad Sci U S A 113: 15138-15143.
  Google Scholar: Author Only Title Only Author and Title

Pignatta D, Erdmann RM, Scheer E, Picard CL, Bell GW, Gehring M. 2014. Natural epigenetic polymorphisms lead to intraspecific variation in Arabidopsis gene imprinting. Elife 3: e03198.
  Google Scholar: Author Only Title Only Author and Title

Pignatta D, Novitzky K, Satyaki PRV, Gehring M. 2018. A variably imprinted epiallele impacts seed development. PLoS Genet 14: e1007469.
  Google Scholar: Author Only Title Only Author and Title

Pillot M, Baroux C, Vazquez MA, Autran D, Leblanc O, Vielle-Calzada JP, Grossniklaus U, Grimanelli D. 2010. Embryo and endosperm inherit distinct chromatin and transcriptional states from the female gametes in Arabidopsis. Plant Cell 22: 307-320.
  Google Scholar: Author Only Title Only Author and Title

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26: 841-842.
  Google Scholar: Author Only Title Only Author and Title

Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European Molecular Biology Open Software Suite. Trends Genet 16: 276-277.
  Google Scholar: Author Only Title Only Author and Title

Rodrigues JA, Hsieh PH, Ruan D, Nishimura T, Sharma MK, Sharma R, Ye X, Nguyen ND, Nijjar S, Ronald PC et al. 2021. Divergence among rice cultivars reveals roles for transposition and epimutation in ongoing evolution of genomic imprinting. Proc Natl Acad Sci U S A 118.
  Google Scholar: Author Only Title Only Author and Title

Schoft VK, Chumak N, Choi Y, Hannon M, Garcia-Aguilar M, Machlicova A, Slusarz L, Mosiolek M, Park JS, Park GT et al. 2011. Function of the DEMETER DNA glycosylase in the Arabidopsis thaliana male gametophyte. Proc Natl Acad Sci U S A 108: 8042-8047.
  Google Scholar: Author Only Title Only Author and Title

Schumann U, Lee J, Kazan K, Ayliffe M, Wang MB. 2017. DNA-Demethylase Regulated Genes Show Methylation-Independent Spatiotemporal Expression Patterns. Front Plant Sci 8: 1449.
  Google Scholar: Author Only Title Only Author and Title

Schumann U, Lee JM, Smith NA, Zhong C, Zhu JK, Dennis ES, Millar AA, Wang MB. 2019. DEMETER plays a role in DNA demethylation and disease response in somatic tissues of Arabidopsis. Epigenetics 14: 1074-1087.
  Google Scholar: Author Only Title Only Author and Title

Springer NM, Anderson SN, Andorf CM, Ahern KR, Bai F, Barad O, Barbazuk WB, Bass HW, Baruch K, Ben-Zvi G et al. 2018. The maize W22 genome provides a foundation for functional genomics and transposon biology. Nat Genet 50: 1282-1288.
  Google Scholar: Author Only Title Only Author and Title

Stelpflug SC, Sekhon RS, Vaillancourt B, Hirsch CN, Buell CR, de Leon N, Kaeppler SM. 2016. An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to Explore Root Development. Plant Genome 9.
  Google Scholar: Author Only Title Only Author and Title

Till BJ, Reynolds SH, Weil C, Springer N, Burtner C, Young K, Bowers E, Codomo CA, Enns LC, Odden AR et al. 2004. Discovery of induced point mutations in maize genes by TILLING. BMC Plant Biol 4: 12.
  Google Scholar: Author Only Title Only Author and Title

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J et al. 2013. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics 43: 11.10.11-11.10.33.
  Google Scholar: Author Only Title Only Author and Title

Warman C, Panda K, Vejlupkova Z, Hokin S, Unger-Wallace E, Cole RA, Chettoor AM, Jiang D, Vollbrecht E, Evans MMS et al. 2020. High expression in maize pollen correlates with genetic contributions to pollen fitness as well as with coordinated transcription from neighboring transposable elements. PLoS Genet 16: e1008462.
  Google Scholar: Author Only Title Only Author and Title

Waters AJ, Bilinski P, Eichten SR, Vaughn MW, Ross-Ibarra J, Gehring M, Springer NM. 2013. Comprehensive analysis of imprinted genes in maize reveals allelic variation for imprinting and limited conservation with other species. Proc Natl Acad Sci U S A 110: 19639-19644.
  Google Scholar: Author Only Title Only Author and Title

Waters AJ, Makarevitch I, Eichten SR, Swanson-Wagner RA, Yeh CT, Xu W, Schnable PS, Vaughn MW, Gehring M, Springer NM. 2011. Parent-of-origin effects on gene expression and DNA methylation in the maize endosperm. Plant Cell 23: 4221-4233.
  Google Scholar: Author Only Title Only Author and Title

Whitington T, Frith MC, Johnson J, Bailey TL. 2011. Inferring transcription factor complexes from ChIP-seq data. Nucleic Acids Res 39: e98.
  Google Scholar: Author Only Title Only Author and Title

Xiao W, Gehring M, Choi Y, Margossian L, Pu H, Harada JJ, Goldberg RB, Pennell RI, Fischer RL. 2003. Imprinting of the MEA Polycomb gene is controlled by antagonism between MET1 methyltransferase and DME glycosylase. Dev Cell 5: 891-901.
  Google Scholar: Author Only Title Only Author and Title

Yamamuro C, Miki D, Zheng Z, Ma J, Wang J, Yang Z, Dong J, Zhu JK. 2014. Overproduction of stomatal lineage cells in Arabidopsis mutants defective in active DNA demethylation. Nat Commun 5: 4062.
 Google Scholar: Author Only Title Only Author and Title

Yelagandula R, Stroud H, Holec S, Zhou K, Feng S, Zhong X, Muthurajan UM, Nie X, Kawashima T, Groth M et al. 2014. The histone variant H2A.W defines heterochromatin and promotes chromatin condensation in Arabidopsis. Cell 158: 98-109.
 Google Scholar: Author Only Title Only Author and Title

Zemach A, Kim MY, Hsieh PH, Coleman-Derr D, Eshed-Williams L, Thao K, Harmer SL, Zilberman D. 2013. The Arabidopsis nucleosome remodeler DDM1 allows DNA methyltransferases to access H1-containing heterochromatin. Cell 153: 193-205.
 Google Scholar: Author Only Title Only Author and Title

Zhang C, Hung YH, Rim HJ, Zhang D, Frost JM, Shin H, Jang H, Liu F, Xiao W, Iyer LM et al. 2019. The catalytic core of DEMETER guides active DNA demethylation in Arabidopsis. Proc Natl Acad Sci U S A 116: 17563-17571.
 Google Scholar: Author Only Title Only Author and Title

Zhang M, Xie S, Dong X, Zhao X, Zeng B, Chen J, Li H, Yang W, Zhao H, Wang G et al. 2014. Genome-wide high resolution parental-specific DNA and histone methylation maps uncover patterns of imprinting regulation in maize. Genome Res 24: 167-176.
 Google Scholar: Author Only Title Only Author and Title

Zhang M, Zhao H, Xie S, Chen J, Xu Y, Wang K, Guan H, Hu X, Jiao Y, Song W et al. 2011. Extensive, clustered parental imprinting of protein-coding and noncoding RNAs in developing maize endosperm. Proc Natl Acad Sci U S A 108: 20042-20047.
 Google Scholar: Author Only Title Only Author and Title