RecA and RecB: probing complexes of DNA repair proteins with mitomycin C in live *Escherichia coli*
with single-molecule sensitivity

Alex L. Payne-Dwyer[1,2†], Aisha H. Syeda[1,2†], Jack W. Shepherd[1,2], Lewis Frame[3] and Mark. C. Leake[1,2*]

[1] Department of Physics, University of York, York, UK, YO10 5DD;

[2] Department of Biology, University of York, York, UK, YO10 5DD;

[3] School of Natural Sciences, University of York, York, UK, YO10 5DD.

*Correspondence: mark.leake@york.ac.uk

†These authors contributed equally.

Abstract

The RecA protein and RecBCD complex are key bacterial components for the maintenance and repair
of DNA. RecBCD is a helicase-nuclease that uses homologous recombination to resolve double-
stranded DNA breaks. It also facilitates coating of single-stranded DNA with RecA to form RecA
filaments, a vital step in the double-stranded break DNA repair pathway. However, questions remain
about the mechanistic roles of RecA and RecBCD in live cells. Here, we use millisecond super-resolved
fluorescence microscopy to pinpoint the spatial localization of fluorescent reporters of RecA or RecB
at physiological levels of expression in individual live *Escherichia coli* cells. By introducing the DNA
crosslinker mitomycin C, we induce DNA damage and quantify the resulting steady state changes in
stoichiometry, cellular protein copy number and molecular mobilities of RecA and RecB. We find that
both proteins accumulate in molecular hotspots to effect repair, resulting in RecA stoichiometries
equivalent to several hundred molecules that assemble largely in dimeric subunits before DNA
damage, but form periodic subunits of approximately 3-4 molecules within mature filaments of several
thousand molecules. Unexpectedly, we find that the physiologically predominant forms of RecB are
not only rapidly diffusing monomers, but slowly diffusing dimers.

Keywords: recombination; repair; DNA damage; mitomycin C; super-resolution microscopy; single-
molecule tracking; Slimfield.

1. Introduction

Accurate duplication of the genome is crucial in all organisms, accomplished by a sophisticated
molecular machine known as the replisome [1]. An inability to accurately replicate genetic material
can lead to cell death and/or cancers [2,3]. Mitomycin C (MMC) is a naturally occurring antibiotic that
can be used to controllably disrupt DNA replication, and thus a valuable reagent in studying DNA repair
processes. It is used as a chemotherapeutic in treating several cancers [4] and retinopathies [5], and
acts by targeting DNA deoxyguanosine (dG) residues [6], forming intrastrand or interstrand crosslinks
[7]. If unrepaired, these structures can interfere with cellular processes such as transcription and
replication, potentially leading to genome instability [8]. An encounter between a mitomycin C-
induced crosslink and an approaching replisome may result in replisome disassembly and eventually
a double strand break (DSB) [9]. RecBCD recognises DSBs in *E. coli* [10], processing the ends to
generate 3'-ended single-stranded DNA (ssDNA) as a landing pad for the principal recombination
protein, RecA [10]. Recombination of RecA-ssDNA complexes with the homologous DNA restores the
replication fork, on which the replisome can be reloaded. The replisome may resume replication if the
blocking adduct is repaired [11]. As a complex of individual RecB, RecC and RecD proteins, RecBCD is
a versatile helicase-nuclease and underpins two major pathways for homologous DNA recombination,
essential for DSB repair [10]. RecBCD activities involve several processes - it recognises and binds DSBs,
begins unwinding both DNA strands, and also degrades both [10]. This activity continues unhindered
until it encounters an octameric Chi site that induces a shift in enzyme activity to degrade only the 5'-

ended strand [12,13]. This activity shift results in a 3'-ended ssDNA overhang that facilitates RecA
loading. A key function of RecA is its ability to form nucleoprotein filaments on exposed ssDNA in
response to damage [14]. These filaments can infiltrate an intact duplex and, on finding homology,
recombine with the infiltrated duplex [15,16]. The extension of filaments along the cell accelerates
this homology search in a non-linear fashion [17]. Following further processing of the resulting
structure, primosome proteins establish an intact replisome thereby enabling replication to resume
[18]. Recombination proteins, such as RecBCD, need access to replication-transcription conflict sites
and collapsed forks, but if RecBCD is missing then double-stranded DNA (dsDNA) is degraded by
exonucleases [19,20], possibly resulting from replisome disassembly. However, how RecA stabilizes
blocked forks remains an open question.

The nucleoprotein filaments formed by RecA are both a requisite and a hallmark of the cell-wide SOS
response [21–24]. The SOS response is a regulatory shift that promotes cell survival in adverse
conditions associated with increased rates of interrupted replication and DNA damage [25]. The SOS
response to DNA damage induced by antimicrobials plays a major role in the emergence of persister
cells [26] and wider antimicrobial tolerance on a population level [27].

Given these far-reaching implications of RecA and RecB activity as studied comprehensively with
mutants [12,24,28–32], it is important to establish the number of molecules present in cells, how they
are spatially distributed and organized, and how these are affected by antimicrobials such as MMC.
Here, we use millisecond super-resolved Slimfield microscopy [33] in live *E. coli* containing
genomically-encoded fluorescent fusions RecA-mGFP [34] and RecB-sfGFP [35]. Since RecA fusion
constructs retain only partial function, our approach makes use of a merodiploid RecA fusion that
expresses from one copy of the native gene and one copy of the *recA4155* fusion construct [34]. This
strain rescues approximately wild-type sensitivity with mixed assemblies of the two RecA proteins
[34].

We use Slimfield microscopy to visualise the spatial distribution of RecA and RecB fluorescent proteins
in individual cells. From these quantitative images, we identify diffraction-limited local intensity
maxima (we denote these as *foci* – see Table 1 for a description of technical Slimfield microscopy
nomenclature used in this study) to a lateral spatial precision of 40 nm [36]. Slimfield uses ~millisecond
sampling that is sufficiently rapid to link the moving foci derived from the same emitter sources over
sequential image frames, following appropriate bespoke particle tracking analysis [33,37,38], into
tracks. Each of these tracks implies the presence of a particle containing one or more associated
molecules; typically more than one prior to photobleaching, so more generally, we term each a
molecular assembly. These tracks reveal the detailed diffusion of labelled RecA and RecB assemblies
in the cytoplasm of a living cell. By using the single-molecule sensitivity of Slimfield microscopy, we
are able to quantify single-molecule photobleaching steps in fluorescence intensity, to identify the
characteristic brightness of a single fluorescent protein [33]. Not only does this calibration apply to
the fraction of the fluorescence intensity for each tracked assembly, but also to the GFP fluorescence
in the whole, or part, of each cell. We use this to determine the number of GFP-labelled molecules
within each tracked assembly (the *stoichiometry*), and the total number of fluorescently-labelled
molecules within each cell (the *cellular protein copy number*), or intracellular segment (the *segment
protein copy number*). Those fluorescent molecules which contribute to the copy number above the
cell's autofluorescent background but are not detected as foci (typically due to high, uniform emitter
density and/or excessive mobility) are denoted the *pool*.

Slimfield has some similarities to single particle tracking photoactivation localization microscopy
(sptPALM) [39–41], however, our approach is simpler, requiring only constitutively expressed
fluorescent reporters such as GFP, and trades off the condition of observing exclusively single
molecules in order to measure the stoichiometry of dynamic assemblies far more accurately. This is
a deliberate advantage of our technique over other single-molecule microscopy techniques as
previously used to count RecB content in cells molecule-by-molecule [35].

Prior to MMC treatment, only point-like assemblies of RecA or RecB are detectable. RecA presents far brighter fluorescence in a cell than RecB, indicating both a typical stoichiometry and a cellular protein copy number that are 2-3 orders of magnitude greater. On treatment with MMC, we observe an increase in the average cellular protein copy number of RecA, but not of RecB, in each cell, with up to 20% of cells devoid of RecB assemblies. MMC induces the formation of RecA assemblies larger than can be captured in single foci, and we interpret these as RecA nucleoprotein filaments, or bundles of filaments [24,30,42–47], typically associated with the SOS response.

Between cellular states of SOS readiness and MMC-induced response, the stoichiometries of RecA assemblies increase, and the diffusion coefficients of assemblies decrease correspondingly. We also discover surprisingly consistent intervals between the stoichiometries of different assemblies in each condition. We interpret the average number of molecules in the intervals (the *periodicity*) as indicative of an oligomeric structural repeat unit that comprises assemblies. The periodicity of RecA assemblies changes from dimeric in character to groups of roughly 3-4 molecules in response to MMC, while the periodicity of RecB assemblies is dimeric, and insensitive to MMC treatment.

Our results shed new light on the relations between structure and function for RecA and RecBCD in mediating repair upon DNA damage.

Metric / Object	Definition
Segment	An area of the image defined by a contiguous subset of pixels in a binary mask. This area either corresponds to a whole cell (a *cell mask*), or more typically a region inside the cell (an *intracellular segment*) of high fluorescent intensity. The term "*segment*" refers to an intracellular segment unless otherwise stated.
Cell mask	A segment containing the outline of one cell. These are extracted using a machine learning protocol (Supplementary Methods).
Intracellular segment	A segment inside the cell. These are extracted from the set of foci localized in that cell by rendering a superresolved image, followed by local Otsu thresholding (Materials and Methods 4.3.5), with the intention of isolating RecA objects that resemble nucleoprotein filaments or bundles.
Assembly	A group of labeled molecules physically associated with one another, either directly or indirectly, such that their diffusive movement is strongly correlated, and therefore always detected in the same track.
Focus (foci)	A spot-like local intensity maximum in a single frame, which corresponds to a localized group of labeled molecules (Materials and Methods 4.3.1). Associated properties include centroid location, total intensity, and signal-to-noise ratio.
Track	A set of foci in adjacent frames that are spatially close enough to form a contiguous trajectory (Materials and Methods 4.3.1). Typically associated with a single molecular assembly, or a group of strongly colocalized assemblies.
Diffusion coefficient	Measure of the random microscopic motion of a specific track based on the increase in the mean square displacement of its intensity centroid over time (Materials and Methods 4.3.2).
Characteristic single-molecule brightness	The average sum of pixel values in foci associated with a single fluorescent reporter molecule (e.g. mGFP), under a fixed imaging condition (Materials and Methods 4.3.3). Equivalent to the modal step size in intensity for tracks in the final stage of photobleaching (Figure S1).
Stoichiometry	The number of fluorescently labeled molecules in a specific track. This is estimated by extracting the sequence of foci belonging to that track, then extrapolating the sum of pixel values in each focus backwards along that sequence to get an initial track intensity that is independent of photobleaching (Materials and Methods 4.3.4). The initial track intensity is then divided by the characteristic single-molecule brightness.
Periodicity	The population-averaged number of fluorescently labeled molecules in inferred repeat units within tracked objects. Estimated by averaging the consistent intervals between nearest-neighbor peaks in the population-level stoichiometry distribution (Materials and Methods 4.3.5).
Integrated intensity	The total fluorescence intensity of a segment in pixel counts, normalised by the characteristic single molecule brightness (Materials and Methods 4.3.6-7).
Cellular (or segment) protein copy number	The average number of molecules in a cell (or intracellular segment), as estimated from the increase in integrated intensity above negative control (i.e. subtracting the contribution from autofluorescence (Materials and Methods 4.3.6-7).
Pool	The intracellular fluorescence which is not detected in tracks.
Pool stoichiometry	The number of untracked, labeled molecules within an area of the pool equal to the size of one diffraction-limited focus (Materials and Methods 4.3.6).

115   **Table 1. Definitions of quantitative analysis metrics for Slimfield.**

2. Results

2.1 Abundance of RecA, but not RecB, increases on MMC-induced DNA damage

We first optimised MMC treatment conditions so that they did not cause cellular filamentation in wild type cells (Materials and Methods 4.1, Figure S1) but did induce the SOS response [48], since cells would then be sensitised to MMC if the SOS response is blocked [48]. Filamentation and loss of viability was also minimal for the labeled strains (Figure 1), hence we used the same MMC treatment for all strains. Given that SOS induction in these and related strains typically takes <20 min [34,48], the kinetics of initial SOS induction due to MMC will likely reach steady state within the 180 min MMC exposure that we used. In light of the timescale of the initial SOS induction, RecA or RecB dynamics were not in the scope of our study here, but rather the steady state effect of MMC on the distribution and molecular organization of RecA and RecB.

Figure 1. Brightfield and Slimfield (mean average of 3 initial frames) of live *E. coli* in 56-salts minimal media, labeled at RecA-mGFP or RecB-sfGFP before and after MMC treatment. Inset (C,D) is another cell transplanted from the same acquisition outside the cropped field of view at the same scale. Brightness of RecB-GFP Slimfield panels (F,H) scaled 100× *vs*. RecA-mGFP panels (B,D). Scale bar 1 μm. (I,J) Probability distributions for number of tracks detected per cell. Tracks are identified in post-acquisition analysis (Materials and Methods 4.3.1) by first detecting foci as local fluorescent maxima, then linking nearest-neighbour foci in subsequent frames.

We performed brightfield and Slimfield microscopy in each field of view (Materials and Methods 4.2).
Binary masks for each cell were extracted independently of any fluorescence signal, using a machine
learning segmentation protocol on brightfield images. We then applied these masks to fluorescence
images to eliminate extracellular background and facilitate statistics on a cell-by-cell level (Figure S5
and Supplementary Methods). Using home-written automated particle tracking and analysis software
ADEMScode [49], we identified fluorescent foci from local intensity maxima in each cell and linked
these into tracks (Materials and Methods 4.3.1). We determined the stoichiometry (the number of
molecules present) of each track, from the summed pixel intensity values corresponding to the start
of each track before any photobleaching has occurred (Materials and Methods 4.3.4), normalized by
brightness corresponding to a single molecule of GFP (Materials and Methods 4.3.3).

From the localizations of foci during photobleaching, we also reconstructed superresolved images of
fluorescent RecA structures. We extracted binary masks from highly fluorescent regions of interest in
these images (denoted *intracellular segments,* or simply *segments* for brevity) using a classical
segmentation method (Materials and Methods 4.3.7), which enabled statistics on an intracellular
segment level (Figure 3 and Figure S3).

Separately, from the cell masks (or intracellular segments) we also calculated the cellular (or segment)
protein copy number (Materials and Methods 4.3.6-7); first we summed the pixel values in each cell
or segment area and normalized these by the characteristic brightness of a single GFP to obtain the
total intensity within that region, expressed in molecules [50]. Taking the difference from an
equivalent area of the control strain that does not express GFP, then yields the cellular or segment
protein copy number corrected for any cellular autofluorescence.

Since the RecA-mGFP strain is merodiploid, both the *recA-mgfp* gene fusion construct and the
unlabeled endogenous *recA* gene are expressed simultaneously [34]. However, their expression levels
are not necessarily identical, nor equivalently inducible by MMC. From previous estimations of the
relative *lexA* suppression rates of the relevant *recA* promoters [51], reasonable expectations are that
a majority of the RecA present in the cell will be labelled with mGFP, and that RecA-mGFP is 2-3 fold
less inducible under the SOS response as endogenous RecA [52]. We estimated the different cellular
levels of unlabeled RecA *vs.* RecA-mGFP using Western blotting (Figure S4), which confirmed that the
RecA-mGFP was in excess compared to the endogenous protein before and after treatment. Both
qPCR and Western blots indicated that both endogenous RecA and RecA-mGFP are inducible by MMC
treatment (Materials and Methods 4.4), with the RecA-mGFP indeed about half as inducible (Figure
S4). Therefore, the total (i.e. labelled plus unlabelled) amount of RecA protein present, whether as
stoichiometry, periodicity or protein copy numbers, is higher than that reported for the RecA-mGFP
data directly, by an approximate correction factor of 1.3-fold in the presence of MMC. In the absence
of MMC the relative amount of RecA-mGFP to RecA is large enough that the correction factor is
effectively 1. As these corrections are indicative, we do not apply them in the early stages of the
Results, but present them later only where relevant to interpretations (Results 2.3 and Discussion).

We find that in the absence of MMC, RecA-mGFP has an an approximately uniform distribution in the
cytoplasm that is occasionally punctuated by bright fluorescent foci that can be linked into tracks
(Figure 1B). The cellular protein copy number of RecA-mGFP increases from 11,400 ± 200 molecules
(±SEM) in untreated cells to 19,500 ± 300 molecules in MMC treated cells (Figure S5A). MMC
treatment resulted in the subset of these RecA-mGFP molecules that are localized in tracks (*i.e.*, the
mean summed stoichiometry of all tracks detected in the whole cell) approximately doubling from
510 ± 30 to 1,080 ± 60 molecules per cell (Table S1). We denote the fluorescently detected, but
untracked, molecules of RecA as residing in a *pool*. The pool typically comprises molecules that are
sufficiently dim, out-of-focus, and/or rapidly diffusing to evade direct particle-tracking-based
detection; here the RecA concentration is exceptionally high such that the stochastic fluctuations
corresponding to motion of discrete foci are partly averaged out. During photobleaching, the density
of foci decreases, overlap decreases and tracks become more evident. The proportion of RecA-mGFP
molecules in tracks is relatively low compared to the pool, but remains representative of the
population of assemblies containing RecA-mGFP.

RecB-sfGFP also exhibited fluorescent tracks against a relatively diffuse background, before and after MMC treatment (Figure 1F,H). RecB-sfGFP foci were observed more commonly near the poles of the cell regardless of MMC (Figure 1G-H). Since the RecB-sfGFP fluorescence signal is comparatively small, estimates based on cellular protein copy number must account carefully for autofluorescence due to native components other than GFP. We estimate that the contribution of autofluorescence from the summed pixel intensity values from unlabeled MG1655 parental cells grown and imaged under identical conditions. We find that the mean level of RecB-sfGFP fluorescence was almost three times greater than the cellular autofluorescence (Figure S5B), therefore there is a comparatively large population of the cellular RecB-sfGFP that evades direct particle-tracking-based detection (*c.f.*, slower sampled images from commercial confocal/epifluorescence microscope systems) and thereby comprise a RecB pool.

The cellular protein copy number of RecB-sfGFP does not decrease significantly following MMC-induced DNA damage, comprising 126 ± 11 molecules per cell before treatment and 101 ± 14 molecules following MMC treatment (Figure S5B, Brunner-Munzel (BM) test, n=246, p=0.0216 | NS, not significant at Bonferroni-adjusted α = 0.01). However, the mean number of RecB-sfGFP localized into tracks does decrease with MMC; just 13.6 ± 0.5 molecules per cell in all tracks, decreasing to 9.3 ± 0.3 on MMC treatment (BM test, n=246, p<0.001). This is clearly much smaller absolute number of tracked molecules per cell compared to RecA-mGFP, but a similar proportion of the cellular protein copy numbers (ranging from 6-10% in each case). The complementary fractions of the total RecA-mGFP and RecB-sfGFP molecules assigned to their respective pools are thus consistently high (89-95%). In the respective strains, the total concentration of RecB-sfGFP is much lower than that of RecA-mGFP, and this likely indicates the correspondingly more rapid diffusion of RecB-sfGFP species within the pool.

2.2 RecB forms characteristic puncta which are partially lost on MMC exposure

We detected typically 1-3 tracks of RecA-mGFP or RecB-sfGFP in each cell above the local background fluorescence (Figure 1I,J). However, each showed strongly opposing trends in the number of tracks observed upon MMC treatment. While RecA-mGFP showed no significant change in the mean number of tracks on MMC treatment (from 1.66 ± 0.06 to 1.86 ± 0.16 tracks per cell, BM test, n=60, p=0.50 |NS), MMC reduced the population average number of RecB-sfGFP tracks significantly, from 2.06 ± 0.09 to 1.56 ± 0.06 per cell.

If, however, we set aside the fraction of cells with no detected RecB-sfGFP tracks, the change in the mean number of RecB-sfGFP tracks is marginal, from 2.20 ± 0.07 to 1.98 ± 0.07 tracks (BM test, n=234, p=0.006). We see that the cells which continue to harbor RecB tracks are relatively unchanged by MMC, each containing an average of 12.1 ± 0.3 molecules per cell. The unexpected subset of cells that are devoid of RecB-sfGFP tracks increases from 6% to 21% of the population on MMC treatment. These otherwise resemble the other treated cells; rather than filamenting, they retain 92 ± 3% of the population averaged cell length and retain the same pool level of untracked RecB-sfGFP molecules.

The increase in the fraction of cells lacking RecB-sfGFP foci agrees with a model of random, independent survival of assemblies (Figure 1J, the MMC+ condition is consistent with Poisson distribution with same mean; Pearson χ^2 test, dof=6, n=234, p=0.004).

Figure 2. Stoichiometry distributions of detected foci of A) RecA-mGFP and B) RecB-sfGFP with (blue) or without MMC treatment (black), shown as kernel density estimates [53]. The statistics used for MMC- (MMC+) conditions include N=190 (67) whole RecA-mGFP cells containing n=316 (125) tracks, or whole N=249 (307) whole RecB cells containing n=514 (478) tracks within the cell masks. The use of 'probability density' reflects the fact that each distribution is continuous with a total area equal to 1, such that areas under the curve correspond to the probability that the stoichiometry of a given assembly falls within a range. The kernel width (the width for smoothing the discrete stoichiometry of each track) is 0.7 molecules following the known detection sensitivity to single GFP (A, inset and B, both panels), or 8 molecules for clarity (main panel A). Insets are the distributions of intervals between nearest neighbor stoichiometry peaks (solid curves) whose modal position, or periodicity, indicates the number of GFP-labeled molecules in a repeating subunit within molecular assemblies (Table 1). Overlaid are heuristic Gaussian fits that minimize a reduced χ^2 metric, with components of equal width and whose centers are fixed at integer multiples to account for the detected optical overlap of an integer number of subunit repeats of tracked foci. The resulting fits comprise three components for RecA-mGFP with MMC treatment (blue, Pearson's R^2 = 0.979, 5 degrees of freedom (dof)) and two components for RecA-mGFP without MMC (grey, Pearson's R^2 = 0.961, 4 dof). The mode of the peak interval is indicated ±95% confidence interval, alongside the number of contributing peak pairs in the original stoichiometry distribution.

RecA-mGFP foci were approximately two orders of magnitude brighter than those of RecB-sfGFP, corresponding to a greater apparent stoichiometry. A subset of polar assemblies in untreated cells are especially bright (Figure 2B); we defined this subset quantitatively by thresholding at 2× the mean stoichiometry of all assemblies. The mean stoichiometry of these bright assemblies is itself as high as 760 ± 40 molecules (Figure S3A).

On treating with MMC, the RecA-mGFP mean stoichiometry almost doubled from 310 ± 8 to 580 ± 30 molecules per focus, reflecting further local accumulation of RecA-mGFP protein (Figure 2A). We find that most RecA-mGFP molecules comprise an untracked, diffusive pool, in which there are ~30 RecA-mGFP molecules in an area corresponding to that of a typical diffraction-limited focus (which we denote as the *pool stoichiometry*, Table 1). The fact that the relative increase in pool stoichiometry with MMC treatment to ~50 RecA molecules (Figure S5C) is smaller than the fractional increase in the amount of RecA in tracks (Figure 2), indicates that the MMC-driven upregulation of RecA disproportionately affects tracked assemblies. As such, either i) new assemblies are formed which contain much more RecA than those before MMC treatment, or ii) those assemblies that already

contain local concentrations of RecA accumulate more RecA. Under our treatment protocol, these changes do not deplete the reservoir of RecA in the cytoplasm. This observation of localized accumulation of RecA-mGFP is consistent with prior reports of long nucleoprotein filament formation on single stranded DNA [34]. The increased number of RecA tracks we observe upon MMC treatment may therefore indicate greater occurrence of processed ssDNA.

The RecB-sfGFP mean stoichiometry decreases very slightly from 6.6 ± 0.1 to 6.1 ± 0.2 molecules per focus (BM, n=478, p<10^{-6}) (Figure 2B). A mean of approximately 6 RecB-sfGFP molecules in each case can be explained if the assembly contains 3 identical subunits whose periodicity is 2 molecules (Figure 2B inset). A pool stoichiometry of ~1 molecule of RecB-sfGFP (Figure S5B,D) suggests that the untracked RecB-sfGFP are likely to be monomers irrespective of MMC treatment (BM test, n=243, p=0.27 | NS). We find that the untracked pool of RecB-sfGFP comprises 90 ± 1% of the total RecB-sfGFP molecules in the cell.

2.3. RecA reorganises into filaments with 3-4-mer subunits in response to MMC

RecA-mGFP and RecB-sfGFP stoichiometry distributions show clear and reproducible peaks (Figure 2A and 2B). One explanation is that each detected fluorescent focus has a diffracted-limited width of ~250 nm that may potentially contain more than one 'subunit' of RecA-mGFP or RecB-sfGFP, bound sufficiently to co-track, such that the measured focus stoichiometry may appear as an integer multiple of that subunit, manifest as periodic peaks on the focus stoichiometry distribution. The expected difference between pairs of values on the stoichiometry distribution is thus either zero or an integer multiple of the periodicity within measurement error. The magnitude of the most likely non-zero pairwise difference value corresponds to the periodicity, with less likely values corresponding to harmonic peaks. Our approach uses a modal estimate of the nearest-neighbor peak intervals (Materials and Methods 4.3.5), and therefore produces a continuous, heuristic estimate for the periodicity. We then compare this periodicity metric to realistic models with integer numbers of molecules. RecA-mGFP tracks have a periodicity of 2.2 ± 0.3 molecules before addition of MMC (Fig 2A inset). This is clearly most consistent with a dimeric subunit of RecA in structures before MMC treatment. After MMC treatment, the most likely interval value is 3.1 ± 0.5 RecA-mGFP molecules, and estimating the additional unlabeled RecA content indicates a likely overall periodicity range of 3-4 RecA molecules (see Discussion and Figure S4).

Figure 3. Bundles of RecA-mGFP filaments in MMC treated cells as observed in A,D) brightfield and B,E) The initial Slimfield fluorescent frame (green) overlaid with all super-resolved single-molecule tracks from the acquisition (*ca.* 40 nm spatial precision, with point localizations from foci visualized as a Normalised Gaussian rendering in ThunderSTORM, Materials and Methods 4.3.7), revealing filaments with high spatial precision (magenta); note that the contrast for the green Slimfield channel is set to half to aid the visibility of the superresolution rendering. C,F) Slimfield at full contrast, overlaid with segments derived from each super-resolved bundle by Otsu thresholding and expanding the resulting image masks by the point spread function width of 180 nm, so as to match the diffracted-limited widefield image optical resolution (white overlay); these segments were then used to calculate the segment protein copy number. Scale bar 2 µm.

In MMC-treated cultures we observe strikingly bright, elongated structures (Figures 3B-E). These resemble parallel or intertwined RecA-mGFP nucleoprotein filaments, that we denote as *bundles* following similar observations by others [21,28,34,42,54]. The bundles were identified in a pointillistic manner by overlaying the tracked foci with our measured localization precision of 40 nm. Though it is unclear whether this segmentation is able to distinguish individual filaments or bundles of RecA from one another, the segments reproduce the contiguous morphology of the bright structures at a diffraction-limited optical resolution (Figure 3C,F).

Though single contiguous segments are evident along the full length of some cells (Figure 3C), the mean number of segments is 1.8 ± 0.4 per cell (Figure S3B). We occasionally observed several small segments per cell, in quantitative agreement (Figure S3B) with our expectation that segments occur at random under of a Poisson distribution, albeit conditioned on the presence of at least one segment persistently occurring per cell. Assuming that DNA damage also occurs randomly but under unconditional Poisson statistics, the number of unaffected cells can only be small when there is significantly more than one segment-inducing damage site per cell at any time. It is not clear how many DSBs per cell cycle an *E. coli* culture can sustain without loss of viability, but repeated stalling and collapse of the replisome is common, and cells with single chronic DSBs are known to replicate almost normally within the confines of an elevated SOS response [55]. Under the relatively mild MMC treatment conditions of our study, we observe a comparatively small fraction of the available RecA-mGFP in bundle-associated segments (here the sum of the segment protein copy numbers is <40% of the cell protein copy number, in contrast to 70% [34]). These observations favour an explanation that much of the RecA in elongated MMC-induced structures is bound to a relatively large number of ssDNA nicks as well as a small number of DSBs per cell at any one time.

The high estimated amount of RecA (Figure 2), and the substantial super-resolved breadth of these objects (Figure 3, Figure S3D) above the ~40 nm width of individual filaments [17] suggests that these are bundles comprised of either multiple RecA filaments, and/or multiple windings thereof. We calculate the segment protein copy number within each of these segments in a similar manner to each whole cell. We find the segment protein copy number is $2{,}800 \pm 200$ RecA-mGFP molecules (Figure S3A). That means each segment typically includes about three times as much RecA in total than the brightest polar assemblies detected in untreated cells (Figure S3A). Greater than 95% of these segments contain a track whose stoichiometry exceeds twice the mean stoichiometry of untreated cells. The RecA structures observed after MMC treatment cannot therefore be produced solely from the large RecA assemblies prior to MMC treatment, but most likely recruit additional RecA from the cytoplasmic pool. We cannot measure the ratio of RecA to available ssDNA directly, however, the high measured amount of RecA provides some indication that it occurs in high enough excess to form RecA-rich bundles rather than simple nucleoprotein filaments. The binding site density on each helical filament containing ssDNA was found in previous studies to be 1.5 nm per RecA in the presence of ATP [56,57]. As the individual filaments are known to be undersaturated with RecA under physiological conditions [58], one would expect a longer filament per molecule. In contrast, we find that each

bundle-associated segment typically measures 900 ± 400 µm (mean ± s.d.) in length, 140 ± 40 µm wide
(Figure S3C-E) and no greater than ~0.4 µm deep (based on depth of focus constraints), but contains
a quantity of RecA we estimate sufficient to produce >7 µm total length of individual helical filament
based on known structures [59]. The longest segments have a more efficient packing density of RecA-
mGFP (Figure S3F) which approaches the binding site saturation limit of 1.5 nm / molecule. This link
between length and efficiency could result from the functional alignment and elongation of the
filament along the cell axis, meaning fewer re-entrant windings of any bundles, and exposure of vacant
binding sites to free RecA in cytoplasm.

In contrast, the brightest RecA assemblies in untreated cells occur in isolation, and are never elongated
but reside within diffraction-limited foci (Figure 1B). Defining these as containing RecA exceeding
twice the mean labeled stoichiometry, these occur in 10 ± 3% of untreated cells and have a typical
content of 800 ± 100 RecA molecules (Figure S3). This relatively high density is equivalent to >2 µm of
filament packing inside a sphere <0.4 µm in diameter. While these assemblies resemble RecA storage
bodies, as suggested previously [21], the *recA4155* R28A mutation has been shown to inhibit the
formation of true DNA-independent storage bodies [24]. Despite the presence of wild type RecA, it is
likely that our observations before MMC treatment indicate DNA-bound RecA bodies that are not
filamentous.

367
368

Figure 4. Distributions of instantaneous microscopic diffusion coefficient for tracks of A) RecA-mGFP
and B) RecB-sfGFP obtained from Slimfield. Kernel density estimates were generated with a kernel
width of 0.008 µm²/s corresponding to the lower bound uncertainty in diffusion coefficient, estimated
as the localization precision / (timestep)². Statistics are as shown for Figure 2.

The diffusive dynamics of RecA assemblies are also indicative of their state of condensation into
filaments. Returning to the tracked foci of RecA-mGFP, we noticed that the mean diffusion coefficient
decreases sharply from 0.17 ± 0.02 µm²/s to 0.07 ± 0.01 µm²/s following MMC treatment (Figure 4A).
This initially low diffusivity, and the further drop in diffusivity, likely reflect the proportion of RecA

condensed onto ssDNA. MMC induces formation of filaments and these are relatively static on the
ca. 10 s timescale of the Slimfield acquisition. In contrast, we find that the mean diffusion coefficient
of tracked RecB is not significantly affected by MMC treatment (Figure 4B), with untreated and treated
values of 0.82 ± 0.03 μm^2/s and 0.79 ± 0.03 μm^2/s respectively (BM test, n=478, p=0.48 | NS). The
diffusivity of RecB-sfGFP in tracks is still lower than expected for a single molecule freely diffusing in
bacterial cytoplasm of ~10 μm^2/s, based on simplistic assumptions of a hydrodynamic diameter of
~10 nm, and contrasts with the large amount of pool RecB-sfGFP that diffuse too quickly to be tracked.
This observation hints at the tracked subset of RecB forming larger complexes with other partners not
detected here, such as RecC and RecD.

3. Discussion

We used Slimfield to investigate the stoichiometry and spatial location of fluorescently tagged RecA
and RecB proteins in live *E. coli* upon treatment with the DNA cross-linking and alkylating agent MMC.
RecA and RecB are repair proteins whose involvement in MMC-specific damage repair pathways, as
part of the SOS response or otherwise, is unclear. We probed the steady state effect of MMC on RecA
and RecB at the minimum inhibitory concentration, which is relevant to sub-lethal antimicrobial
exposure. Our results show that the sensitivity and dynamic range of Slimfield is sufficient to quantify
counts, either by stepwise photobleaching of multi-molecular complexes or by direct detection of
single molecules using millisecond sampling.

RecA assembly formation is not solely correlated with induced DNA damage. Before treatment with
MMC, we find that a portion of RecA appears in foci at an average incidence of approximately 2 foci
per cell. In 10% of cells, at least one of these foci is especially bright, circular and localized to one of
the cell poles. A previous study reports that a minority of cells (4-9%) exhibit spontaneous RecA foci
near the poles prior to DSB induction [34]. It has been suggested that wild type RecA foci at the cell
membrane might act as nucleation points for later filament formation across DSBs [60], or that these
are storage bodies outside the nucleoid [21]. However, the RecA-mGFP strain used here (and in [34])
is a *recA4155* (R28A) genotype which abolishes DNA-independent aggregation of RecA [52,61]. In this
strain, we cannot eliminate the possibilities that wild-type RecA forms native storage bodies that are
undetected due to exclusion of RecA-GFP, or indeed visible storage structures which do recruit the
mutant RecA-mGFP (RecA4155), which would account for the resemblance of detected foci to
previous observations of these bodies outside the nucleoid [21,34,61]. In the case where RecA-mGFP
cannot participate in storage bodies and can only aggregate in the presence of DNA, there is an
alternative explanation for the subset of RecA-mGFP foci we observe, distinct from membrane
anchors and storage bodies. These foci do not appear to require RecB for spontaneous assembly [34]
indicating that they are independent of DSBs and instead assembled at incidental sites of ssDNA. The
foci lie consistently at the periphery of the cell, which indicates they are not likely to be associated
with ssDNA within replication forks. These occasionally bright foci may instead simply reflect
stochastic ssDNA nicks in a small proportion of cells of an otherwise healthy culture.

Our findings show that the RecA-mGFP copy number increases upon treatment with MMC. We
observed a modest increase in the number of tracks, but whose stoichiometry per focus is almost
twice those of untreated cultures. This observation of spatially localized RecA and is consistent with
significant assembly formation ultimately leading to formation of long nucleoprotein filaments on
ssDNA as nucleated from polar locations [34]. These filaments are known to accumulate into bundles
as posited by Story et al. [42]. We observed filamentous bundles in MMC treated cultures, possibly
due to increased availability of processed ssDNA from DNA damage sites. RecA-assisted homologous
recombination and RecA* disassembly occur on a timespan between 15 min [17] and 2 hours [34]. We
detect a large increase in RecA stoichiometry (Figure 2) and cellular protein copy number (Figure S5A)
and decrease in diffusivity (Figure 4) even after 3 hours' treatment, indicating that RecA bundles
continue to form in response to constantly accumulating DNA damage.

Our observation of an about 2 intracellular segments per cell (Figure S3B) is consistent with
approximately 2 MMC-induced RecA bundles each extending along opposite halves of a cell (Figure
3F) at any one time in the steady state. This observation may indicate the presence of a double-strand
break (DSB) with nearly-bridged loci. However, according to the schemes in previous work [17,34], the
development and breakdown of filaments [17] and bundles [34] takes typically <20 min, while for
bundles only, recombination is the rate-limiting step, taking up to 90 min [34]. It follows that labeled
bundles associated with DSBs would be expected to be bridged for most of their visible lifetime. It is
therefore possible that either i) multiple DSBs are present and the segments correspond to different
simultaneously bridged DSBs, or that ii) one bridged DSB is present alongside other defects which
support RecA filament binding, such as ssDNA nicks.

Intracellular segments were typically aligned along the cell axis (Figure 3B,E) in agreement with the
observations of filaments and bundles by other authors [17,34]. While some degree of alignment is
expected for all segments much longer than the cell diameter (0.78 ± 0.05 μm), we note that more
than half of the detected segments are shorter than this (Figure S3E), which may suggest an alignment
mechanism that is not solely due to segement length. Moreover, segments appeared to follow the
central axis of the cell, rather than the cell outline (Figure 3B,E), which suggests they fall mostly within
the nucleoid rather than residing at the cell membrane, in keeping with the known DNA repair function
of the filaments. Filament extension along the cell axis is not predicated on the presence of sister
homology [34] but inherently reduces the dimensionality of the search for any homology to one across
the cross-section of the cell, independent of cell length or DNA content [17]. Thus, extension vastly
accelerates the search time [17]. However, the cause of the extension is unclear. It may reflect simple
polymeric elongation under spatial confinement inside the cell, but extension is entropically
unfavourable for a flexible polymer. Stiffening and/or thickening of filaments into bundles [34] would
therefore faciliate extension. The bundle model in [34] suggested a thickened central backbone
flanked by thin filament ends. The bundles observed in our study appear to be thickened with a typical
cross-sectional full-width half-maximum of 140 ± 40 nm (Figure S3C) in agreement with previous
observation, 160 ± 30 nm [34]. Rather than a monolithic central section, our observations resemble
beads on a chain, or a sequence of thick and narrow sections (Figure 3B,E). We find the median width
increases rapidly with segment length (Figure S3D), which in this binary framework, suggests the bulk
of the increase in bundle length is taken up by the thickened portions and that the thin sections are
relatively short. Yet, individual ~40 nm-wide filaments without thickened portions have also been
observed previously to extend dynamically on the scale of minutes or less along the length of the cell
[17]. We speculate that this suggests an active process of pole-to-pole translocation of thin filament
ends (for example, as proposed in [59]), to facilitate the reduced search time.

The observable periodicity of RecA structures could indicate a difference in their macromolecular
organization in response to MMC treatment. We observe a change in the periodicity of RecA
stoichiometry from ~2 molecules in foci in untreated cells, to a ~3-4-mer within spatially extended
filaments following treatment with MMC, after accounting for the unlabelled RecA content per cell
with a correction factor of 1.3 ± 0.1 (Results 2.3 and Figure S4). Previous *in vitro* and *in vivo* studies
indicate that RecA undergoes linear polymerization in a head-to-tail fashion, with dimeric nucleation
points on ssDNA mediated by SSB [62] consistent with our finding of dimeric periodicity prior to
treatment. These also provide evidence for stable trimeric, tetrameric, hexameric and the filamentous
forms when ssDNA is present [63], consistent with our findings post-treatment. Our snapshot
observation of filament stoichiometry cannot shed light directly on models of dynamic nucleation or
stepwise growth, as explored in [64–66]. Rather, it explores molecular details of the characteristic
protein subunits within the mature filament at steady state. The helical geometry of the filament,
with a pitch of 6 RecA molecules per turn, implies that each group of 6 RecA forms a split-ring structure
related to the intact hexameric ring of DNA helicases, but distorted axially such that rings each
complete a single helical turn around ssDNA [67]. Such ring-shaped hexamers have been identified *in*
vitro for both the wild type RecA protein, and the RecA (R28A) mutant [61] that is fused with GFP in
our experiment. Even if isolated oligomers were somehow unstable *in vivo*, a polymeric filament could

conceivably still result from a small, periodic barrier to polymerization corresponding to this split-ring
distortion. This points to the hypothesis that the fundamental building block of RecA filaments is a
factor of 6, if not a hexamer. However, our stoichiometry analysis suggests variability in the total size
of assemblies, with our periodicity results indicating a range of 3-4 molecules per subunit. This could
reflect trimers which form half-turns in the filament, or perhaps tetramers as an intermediate
between preexisting dimers and hexameric rings. Although these data cannot directly establish
whether independent oligomers of wild type RecA occur *in vivo* either on DNA or in the cytosol, it is
conceivable that assembly and rearrangement of RecA subunits on DNA could generate the canonical
ATP-inactive and ATP-active DNA-binding filaments [68,69]. In light of a recent study highlighting the
role of RecN in RecA filament formation and activity [59], it is interesting to pose whether RecA
assemblies with the dimeric subunit may be devoid of RecN and are ATP-inactive, and if these might
then change to a higher oligomeric form upon DNA damage via the involvement of RecN and its
associated ATP-activity.

Our measurements confirm that RecA has a very high concentration in the cytosol of live cells. We
observe that untreated cultures comprise approximately 11,000 molecules of RecA-mGFP per cell,
which increases to 20,000 RecA-mGFP molecules in cells treated with MMC. Of the latter, 28 ± 7%
resides in filamentous bundles large enough to be resolved in millisecond widefield fluorescence
images. Applying the approximate merodiploid correction factors that we estimated of 1.0 ± 0.1 and
1.3 ± 0.1 respectively (Results 2.3 and Figure S4), the total copy number is approximately 11,400 ± 200
RecA molecules in untreated cells, increasing to 25,300 ± 400 molecules in treated cells. Though less
than the 4-5-fold transcriptional increase suggested by qPCR (Figure S4), the more than two-fold
increase of total RecA with MMC resembles the increase detected in western blots (Figure S4). While
the RecA copy number we estimate in untreated cells exceeds the *ca.* 5,000 molecules reported
previously by Lesterlin et al [34], our more direct estimations are of similar order and correlate with
previous work indicating 2,900-10,400 molecules, with the high end of this range obtained from cells
in EZ rich medium using a ribosome profiling method [70]. Approximately 15,000 RecA molecules per
cell in rich medium were reported previously, using semi-quantitative immunoblotting [71]; the same
study found that the RecA copy number increased to 100,000 molecules upon MMC treatment. Large
discrepancies between studies in the increase in RecA due to MMC treatment are not only due to
treatment dose [72] but also arise from differences in *recA* genotype, culture media and growth
conditions, as noted by others [21]. In particular, our study uses a minimally inhibitory treatment with
MMC (Figure S1).

While the RecA-mGFP protein is not identical to native RecA in its enzymatic activity [32,43,52], the
merodiploid strain used in our study also includes a full complement of native RecA from a single
allele. This wild-type protein is expected to mitigate the partial loss of sensitivity in RecA-mGFP in any
given mixed assembly, as shown on a cellular level by the similar SOS response profile [34] and lack of
filamentation under treatment with MMC (Figure 1, Figure S1). The *recA* wild type allele is expressed
under control of the native operator, while the *recA-mGFP* allele is expressed under the *recAo1403*
operator. In the absence of treatment with MMC, this operator is known to result in an increased
transcription rate of the *recA-gfp* allele relative to the wild type *recA* gene under its native promoter
by a factor of 2-3, while both alleles are upregulated to the same level under induction of SOS [52].
Using quantitative Western blotting we estimate that prior to MMC treatment, RecA-mGFP is actually
present at several tenfold more than the unlabeled protein (Figure S4), and that in the presence of
MMC the ratio of RecA-mGFP to RecA is lower, at approximately 3-4 to 1. From these ratios, we
derived our approximate correction factors of 1.0 or 1.3-fold for the total amount of RecA protein, in
the absence or presence of MMC respectively. While RecA-mGFP is known to label RecA assemblies
[34], it cannot form DNA-independent assemblies by itself [52], and is therefore reasonable to
conclude that all labelled sites here represent occupied DNA on which wild type RecA and RecA-GFP
are interchangeable. Even if the binding partition of wild type RecA were higher, for example reflecting
the relative sensitivity (Figure S4A and [52]), the high relative concentration of RecA-mGFP (Figure
S4B) would conceivably result in the majority of RecA sites on DNA being occupied by RecA-mGFP.

Lesterlin *et al*. showed that RecA immunostaining of filaments (agnostic as to GFP labelling) correlates
with the fluorescent distribution of RecA-GFP [34], proving that dark filaments exclusively of wild-type
RecA cannot be present. Though this result could potentially be interpreted that the structure is
entirely RecA-GFP and that the highly sensitive wild-type RecA is excluded, this wild-type RecA would
have to somehow rescue DNA repair function in the cytoplasm rather than in filaments, which has no
known basis. We therefore assume the presence of hybrid filaments. In any case, the effect of
excluding wild-type RecA from filaments would simply mitigate our periodicity correction factor
toward unity, and narrow our estimate of the periodicity within filaments toward a value of 3 RecA
molecules.

Unlike RecA-mGFP, we detected only modest quantities of RecB-sfGFP in untreated cells grown in
minimal medium: 13.6 ± 0.5 molecules in tracks, and 126 ± 11 molecules in total per cell based on
integrated GFP fluorescence corrected for cellular autofluorescence. Several previous reports also
indicate that RecB is very scarce – typically less than 20 molecules per cell [35,73]. One of these studies
estimated that there are just 4.9 ± 0.3 RecB molecules per cell using a HaloTag fusion allele labeled
with HTL-TMR, and 4.5 ± 0.4 molecules per cell using magnetic activated cell sorting of the same RecB-
sfGFP strain that we use here, albeit in M9 medium and restricted to nascent cells for which the
average copy number is approximately halved [35]. An earlier mass spectrometry study used intensity
based absolute quantification to estimate 9-20 RecB molecules per cell across different stages of
growth in M9 minimal media [73]. Ribosome profiling estimated the RecB copy number to be 33-93
molecules per cell in different growth media [70]. However, these techniques are either *ex vivo* or
necessitate significantly perturbed intracellular crowding that may conceivably result in potentially
non-physiological molecular assemblies.

Comparing the number of RecB in tracks in our present study with the number of RecB in distinct foci
per cell reported previously, we find a similar albeit slightly higher estimate, possibly because our
approach is based on fluorescent fusions with a high labelling efficiency in unsynchronized cultures,
as opposed to selecting nascent cells. However, our measurement of RecB copy number exceeds
previous estimates. The large remainder in summed pixel fluorescence intensity may represent two
possible contributions. The first is from RecB that diffuses faster than Slimfield can track. The highest
diffusion coefficients of tracked RecB assemblies approach 3 μm^2/s (95% quantiles, Figure 4). We
estimate the limit of measurement as approximately 5 μm^2/s, though it is conceivable that free
monomeric RecB-sfGFP could exceed this, given that it has been estimated to reach diffusion
coefficients equivalent to approximately 8 μm^2/s in *E. coli* cytosol [74,75]. A second possible source is
an increase in net autofluorescence relative to the parental strain when the real RecB are labeled; it
is unlikely that could account for the discrepancy, since this would require a 3-fold increase in
autofluorescence based on our measurements, and such a drastic increase lacks precedence (for
example, upon treatment with MMC at a high level sufficient to induce widespread RecA
filamentation, our estimation suggests only an increase in autofluorescence of no more than 20%).
Furthermore, the measured rate of photobleaching of the diffuse RecB-sfGFP signal matches that of
RecB-sfGFP tracks and is roughly half the rate of the autofluorescent parental cells (Figure S6, Table
S2). The implication is that untracked RecB-sfGFP is the major contributor to mean cellular
fluorescence, which is then a more accurate reflection of total copy of RecB than simply the number
of molecules in tracks.

The cellular protein copy number of RecB does not change significantly with MMC (Table S1, Figure
S5B,D), suggesting that there may only be a modest regulatory response to DNA damage. Although
MMC is known to induce the SOS response and cell cycle arrest [4,48], *recB* expression is itself not
induced directly as part of the SOS response. RecB-sfGFP foci increase neither in number (Figure 1J)
nor stoichiometry (Figure 2B), which compares with earlier observations that treatment with MMC
under similar concentrations to those used in our study do not significantly change RecB expression
[22]. In fact, the number of observed tracks per cell *dropped* considerably after MMC treatment, due
to a sharp increase in the proportion of cells in which RecB assemblies were absent, from 6% to 21%.
This reduction in RecB assemblies was at odds with our expectation that MMC would eventually

increase the recruitment of RecB in response to damage, if not increase the cellular production of
RecB. MMC treatment is known to increase the occurrence of DSBs and thereby drive demand for
DSB processing [4] that is typically mediated by RecB. Yet, rather than initiating cellular upregulation
of RecB, treatment with MMC acts to partially deplete localized assemblies. Given that DSBs are likely
to occur in the majority of cells under our MMC treatment, as indicated by the ubiquitous induction
of RecA filaments (Figure 3), the fate of RecB assemblies cannot simply reflect the presence or absence
of DSBs. The increase in the fraction of cells lacking RecB-sfGFP tracks is consistent with random,
independent survival or breakdown of assemblies (Figure 1J). This result may indicate a situation
where pre-existing RecB (hetero)complexes at foci are occasionally disassembled while interacting
with sites of MMC-induced DNA damage, such as DSBs. This instability of (presumably heteromeric)
RecB assemblies might result from successfully bridged pairs of RecA filaments, however, we did not
detect any correlated loss of pairs of RecB foci, as might be expected for recombination events.
Notably, the number of tracked foci detected per cell is approximately 2 for both RecA-mGFP and
RecB-sfGFP. Future colocalization studies of RecA and RecB assemblies may offer more direct insight
into the functional interaction and turnover of these repair proteins in regards to whether the average
of 2 be related to the number of replication sites, or perhaps simply reflects a small average number
of severe DNA damage sites per cell.

Independent of MMC treatment, we observed a dimeric periodicity for RecB-sfGFP. This suggests that
RecBCD heterotrimers occur in pairs *in vivo*. Indeed, earlier *in vitro* studies identified the occurrence
of $(RecBCD)_2$ complexes, possibly held together by the nuclease domains of the two RecBCD
monomers [12]. However, the authors concluded that the monomeric form is functional while the
dimeric form is nonfunctional [13]. Furthermore, crystallization of the RecBCD complex for structural
studies contained two RecBCD-DNA complexes in the asymmetric unit [76]. Our observations cannot
determine covalent interactions directly between RecB molecules, but their cotracking is very strongly
correlated. We can infer two details: first, that the dimeric form of the complex, $(RecBCD)_2$, occurs in
live cells, and second, that previous *in vitro* observations of dimers are carried over from their
physiological state. Our findings suggest a hypothesis that assemblies with multiple pairs of RecB have
a greater activity on DSBs than isolated RecB in the pool. Making the distinction between monomeric
RecBCD in tracks and monomeric RecB in the untracked pool suggests that RecB monomers in the pool
could potentially act as a reservoir. One may consider the alternative situation, where the monomeric
pool are the functional RecB elements and the assemblies are reservoirs that disassemble in response
to damage, but this makes less sense, since those monomers would already be in excess. A mean
stoichiometry of ~6 molecules indicates that RecB foci may occur as colocalized assemblies that
comprise roughly three pairs of RecBCD heterotrimers (Figure 5). It would be interesting to estimate
the stoichiometries of RecC and RecD in future studies to understand their association in processing
DSBs in greater detail.

Figure 5. A model of DNA damage caused by treatment with MMC and subsequent repair at the replication site by RecA and RecBCD. A) Intact replication fork; occasional binding of multiple RecA dimers to DNA away from the fork as well as RecA dimers as DNA-free storage bodies in the cytoplasm; B) exposure to MMC and induction of an interstrand crosslink that acts as a barrier to an approaching replication fork; C) replisome dissociates if unable to overcome barrier; dissociated fork is recognised by branched DNA specific endonucleases (filled triangle) that can eventually cause DSBs leading to replication fork collapse; replication fork collapse allows access to repair enzymes to recognise the lesion; D) a newly generated DSB is recognised by RecBCD and processed to generate a 3' single strand end; E) RecA dimers identify the newly generated ssDNA and assemble in groups of 3-4-mers into RecA* filaments; RecA is shown as a short stretch for illustrative purposes but may extend for many thousands of molecules over several hundreds of nm of ssDNA, and these filaments may be twisted and/or grouped into bundles. F) Strand exchange followed by processing of the double strand break, then recombination sufficiently upstream of the lesion and subsequent G) reloading of the replisome. This process allows sufficient time for the repair enzymes to repair the lesion on the template strand,

so that replication may resume. For a detailed overview of the possible pathways to fork restoration,
refer to [11].

While MMC-induced damage constitutes a range of chemical moieties [77], the canonical mechanism
of MMC toxicity is of interstrand crosslinks at dG sites [6,7]. The specific repair of interstrand crosslinks
(implied in Figure 5) can involve several repair pathways, primarily nucleotide excision repair (NER),
which converts the crosslinks into dsDNA breaks [78]. Although NER enzymes such as UvrD typically
degrade RecA filaments, NER is involved in the cleavage of damaged replication forks into suitable
substrates for downstream processing, including RecA-mediated recombination [79]. Repair of the
fork is then completed, for example by PriA-, Rep- and PriC-dependent pathways [11,18,80] on sets of
ssDNA and a dsDNA end (Figure 5). The observation of a greater increase in RecA-mGFP copy numbers
and foci compared to RecB-sfGFP could indicate a significant proportion of single-strand breaks and
single-strand gaps at sites of crosslinks. While two previous studies reported that NER action on
crosslinks also produces ssDNA nicks [78,81], we do not know if this applies strictly to MMC-induced
NER, as our present work does not pertain to genes that *process* ss-gaps. Future analysis of proteins
that process ssDNA breaks may potentially shed light on the relative occurrence of the two types of
breaks by MMC and their relation to repair of replication forks.

While others have shown that *recB* deletion abolishes UV-induced filaments of RecA [34], we do not
know the effect of *recB* deletion and MMC treatment on RecA dynamics. To avoid RecA interference
in 'normal' ssDNA processes such as replication, the cell maintains strict control over filament
nucleation, based on RecA and associated cofactor concentrations. It is therefore likely that the
observed filamentation upon treatment with MMC is dependent on RecBCD, indirectly pointing
towards increased occurrences of DSBs in these cells. Alternatively, if RecA nucleation is independent
of RecBCD, one might anticipate little change in RecA dynamics upon *recB* deletion. However, further
analysis of MMC-dependent RecA stoichiometry and copy number in a strain devoid of RecBCD activity
- and with a controlled RecFOR pathway [19] – is needed to differentiate between these models.

In conclusion, RecA occurs as assemblies located near poles of wild type cells in a dimeric periodicity
consistent with nucleation models. Upon mild treatment with MMC, RecA is upregulated at least two-
fold, and assembles into long filamentous bundles on newly generated ssDNA in effectively all cells
without exhausting the cytoplasmic reservoir. These mature bundles have a much lower diffusivity,
reflecting their aggregation of a few thousand molecules each, with a structural periodicity in the
range of 3-4 RecA molecules. The bundles are typically wider than single filaments, but both forms are
known to facilitate homology search for homologous base-pairing with an intact duplex. Generation
of ssDNA is known to occur at a DSB induced by processing of disassembled forks upon recognition by
RecBCD. We observed RecB as a moderately diffusive set of three associated dimers at two locations
in the cell, providing further evidence that RecBCD predominantly occurs as pairs of heterotrimers
inside the cell at either end of DSBs. Our work implies the existence of a separate, significant reservoir
of highly diffusive RecB monomers. Neither of these forms of RecB are upregulated upon MMC
exposure, nor do they change their mobility. Accordingly, RecB is not a part of the SOS regulon.
Instead, MMC-induced DNA damage impacts the formation – or induces a higher turnover – of these
periodic RecB assemblies potentially associated with further DSB repair.

4. Materials and Methods

4.1. Strains, culture and MMC protocol

Three strains of *E. coli* were used in this study without alteration:

Control:
MG1655
RecA-mGFP [34]:
MG1655 rpsL (Str^R^,lac+) ygaD1::kan recAo1403 recA4155,4136-gfp901, fhuB::recAwt-cm
RecB-sfGFP = MEK706 [35]:
MG1655 recB::sfGFP

The RecA-mGFP strain used here is the same as in [34]. It is a merodiploid, natively promoted

derivative of the SS3085 strain [52,82,83]. It expresses both i) the wild type unlabelled RecA protein

from a single, ectopic wild-type *recA* allele, under a wild-type operator, and ii) a labeled mutant

protein at the native *recA* site with a mutant operator *recAo1403*, which has a higher transcription

rate than wild type [52]. The labelled *recA* is a *recA4155* (R28A) mutant, which complements the wild

type recombination function *in vitro* [34,61] but has important differences in self-assembly. Unlike

wild type RecA, the *recA4155* forms of RecA cannot alone form assemblies independently of DNA as

proven by comparisons *in vitro* [61] and competitive binding studies *in vivo* [52]. The *gfp-901* label is

the same as *mut2* (A206T) which corresponds to a monomeric GFP (mGFP) [84]. The notation

recA4136 refers to the insertion of a linker, as well as the *gfp-901* gene, between the penultimate and

ultimate stop codons of *recA*. The fusion with mGFP impairs the recombinant sensitivity of the labeled

RecA protein; were the strain to include only the fused allele, it would be fully as SOS inducible as wild

type MG1655, but only approximately two-thirds as UV resistant, and would be compromised up to

ca. 10-fold for recombination activity [21,52]. Induction of the SOS response would also take 30 min,

or roughly twice as long as wild type (the R28A mutation prevents this from being an additional 2×

slower) [52]. The merodiploid strain rescues both these functions and their kinetics: functional RecA

filaments labeled with 70% of the total available mutant fusion protein form within just 15 min of DNA

damage and reach steady state within 90 min, similar to wild type MG1655 [34].

The RecB-sfGFP fusion was constructed in [35] by wild-type *recB* replacement under plasmid-mediated

recombination. N-terminal fusions were shown in [35] to be functional using growth curves and tests

of DNA repair, in contrast to C-terminal fusions which may disrupt RecBCD complexation [35].

E. coli strains were grown overnight in 56-salts minimal media at 30°C to mid-log phase in an Innova

44 shaker incubator (New Brunswick). The mid-log phase cultures were concentrated to ~100 cells/ml

(OD_{600} ~0.3) and split into two equal fractions. Aliquots were adjusted to either nil (MMC-) or the

minimum wt inhibitory concentration of 0.5 µg/ml MMC (MMC+) (Figure S6 and [48]) and incubated

at 30°C for a further 3 h (Figure S6). Cells were harvested for microscopy on 1% w/v agarose pads

suffused with the same liquid media and imaged within 1 h.

4.2. Slimfield

A custom-built Slimfield microscope was used for single colour, single-molecule-sensitive imaging with

a bespoke GFP/mCherry emission channel splitter as described previously [11,37]. The GFP channel

was recorded, while the mCherry channel was used only as a negative control. The setup included a

high-magnification objective (NA 1.49 Apo TIRF 100× oil immersion, Nikon) and the detector was a

Prime95B sCMOS camera (Photometrics) operating in 12-bit gain at 180 Hz and 3 ms exposure/frame,

for a total magnification of 53 nm/pixel. The samples were illuminated either in brightfield, or for

Slimfield fluorescence in camera-triggered frames by a collimated 488 nm wavelength continuous

wave OPSL laser (Coherent, Obis LS) in Gaussian TEM_{00} mode at a power density of 5 kW/cm^2. The

number of frames per acquisition was 2,000 for RecA and 300 for RecB strains.

4.3 Quantitative tracking and protein copy number analysis

4.3.1 Identification of Slimfield foci and assignment into tracks

Slimfield image sequences were processed by custom ADEMscode software in MATLAB (Mathworks)

[33,80,85–87]. This pipeline identified foci from local maxima in pixel values within individual frames.

An iterative Gaussian mask algorithm was used to detect the centroids of foci, using a circular region

of interest of radius 5 pixels within a sliding window of 17 pixels. The intensity of each focus was

calculated as the sum of the circular region corrected for the average background in the surrounding

annular region. The prospective foci were accepted if their intensity was >0.4× the standard deviation

in the background region. The nearest neighboring foci in adjacent frames within 8 pixels of each other

were assigned to the same track, with a minimum of 4 foci per track. The typical track duration was

limited by diffusion and/or photobleaching to a mean of >13 foci per track over ~75 ms real time, or
~40 ms cumulative exposure (Table S1).

4.3.2. Diffusion coefficient

The centroids of the foci within each track, as generated from the ADEMScode tracking analysis above,
were used to calculate displacements over the length of each track in chronological sequence. From
these, the mean square displacements (MSDs) of each track were calculated by averaging the square
of the displacements across equal lag times, corresponding to all possible intervals between frames
up to the length of the track. For each track, the MSDs at the four lowest lag times were linearly
interpolated (with a constraint on the fit of passing through a specified intercept on the lag time axis,
equal to the square of the measured localization precision of 40 nm divided by the frame interval of
5.7 ms). The initial slope of this fit (and corresponding error) was then divided by a factor of 4
according to the 2D diffusion equation [88] to yield a diffusion coefficient (and error) for that track.

4.3.3 Characteristic single-molecule brightness

The intensity of each focus was estimated by integrating the local pixel values with a local sliding
window background subtraction. After photobleaching sufficiently to show single photoactive GFP
molecules, the characteristic single-molecule brightness of a single GFP molecule was estimated from
the modal brightness of these foci. These were confirmed to be broadly consistent with estimates of
the signal per GFP in each dataset were determined from the monomeric intervals in total number of
counts due to stepwise photobleaching, as identified by a Chung-Kennedy edge-preserving filter
(15 ms window, 50% weighting, Figure S7) [89]. This integrated intensity is characteristic for each
fluorescent protein under fixed imaging conditions, although mGFP and sfGFP were found to be
indistinguishable in this respect, and hereafter referred to collectively as GFP. To ensure consistent
counts per single-molecule probe, analysis was restricted to the uniformly illuminated area lying
within half of the $1/e^2$ beamwaist of the excitation laser in the sample plane. The integrated intensity
of GFP *in vivo* was found to be within 14% and 9% respective errors in RecA and RecB (88 ± 18 and 177
± 16 pixel grey values per GFP for the respective gain modes). The combined equivalent is 88 ± 7
photoelectrons per GFP per frame, which is precise enough to unequivocally identify groups or steps
of up to 12 GFP molecules.

4.3.4 Stoichiometry

Each track is associated with an assembly that contains a certain number of molecules, or
stoichiometry, at the initial point of acquisition. To estimate this stoichiometry for a given track, the
intensities of the constituent foci were linearly extrapolated using the first 4 datapoints in the track
back to the timepoint of initial laser exposure. This initial intensity of this fit was divided by the
characteristic single-molecule brightness signal associated with one fluorescent protein under a fixed
excitation-detection protocol. The result is a stoichometry expressed as a number of molecules. The
standard error associated with a stoichiometry value of 1 molecule is approximately 0.7 molecules. To
avoid undercounting bias due to photobleaching, only tracks in the first 10 frames after laser exposure
were considered for stoichiometry estimates.

4.3.5 Periodicity

The distributions of track stoichiometry may show periodic peaks, whose smallest reproducible
interval can be interpreted as a physical repeat unit or *periodicity* within assemblies. To calculate
periodicity, first the stoichiometries of all tracks within each acquisition were represented as a kernel
density distribution. The kernel width used was the empirical standard deviation on the characteristic
single molecule brightness of 0.7 molecules [41]. Peaks in this distribution were detected using the
MATLAB *findpeaks* function, and the intervals between nearest neighbor peaks were calculated. These
sets of nearest neighbor intervals for each acquisition were then aggregated across the relevant
population of cells. A second kernel density estimate was calculated over the intervals for a
population, with a kernel width of 0.7 molecules multiplied by the square root of the mean
stoichiometry, divided by the square root of the number of interpolated intervals. The fundamental
value of this interval distribution (corresponding to the center of the leftmost peak in Figure 2 insets)

was refined by fitting the curve with a sum of Gaussian terms centerd at multiples of the fundamental
value. To accommodate the uncertainty in the single molecule characteristic brightness, the
fundamental value of the fit was not constrained to an exact integer value but represents a heuristic
model for the periodicity. The number of terms in the fit was set to minimize the reduced χ^2 metric in
the fit. This modal value was reported with 95% confidence interval as the periodicity of assemblies in
each population. This method of estimating periodicity was verified as independent of the mean
stoichiometry using simulated data drawn from noisy Poisson-distributed multiples of an oligomeric
ground truth (artificial input value). This analysis reproduced the expectation that the minimum
number of tracks required for sufficient peak sampling, and therefore the limit of periodicity
detection, scales with the square root of the mean stoichiometry.

4.3.6 Cellular protein copy numbers and pool stoichiometry

The cellular protein copy numbers as reported in the Results Section 2.1, Table S1, Figure S5 and
Discussion correspond to whole cell masks, as identified using the manual annotated machine learning
segmentation output from brightfield images (Figure S2 and Supplementary Methods). Integrated
intensities of cells (uncorrected cellular protein copy numbers) and pool stoichiometries, were
determined not from tracked foci, but directly from the raw image sequences using the CoPro package
in ADEMscode software following [37] with the characteristic single-molecule brightness of GFP (as
described in 4.3.2), the cell masks, and the camera's dark pixel bias as input. The procedure effectively
adds up all of the pixel values within the mask in question in an initial frame, and accounts for the
convolution of the 3D cell volume with the widefield point spread function, followed by projection
onto a 2D image. To obtain the cellular protein copy number in the labeled strains, and account for
the contribution of autofluorescence, we calculated the difference in mean integrated intensity per
segment between the labeled and parent strains under the corresponding MMC± condition, adjusted
by the ratio of mean segment area. The pool stoichiometry in each cell is a measure of its untracked
molecular concentration. It is calculated in CoPro as the cell's integrated intensity, less the mean
integrated intensity of the parental cells, less the total stoichiometry of tracked foci in the cell, divided
by the area of the cell mask relative to the area within one diffraction limited focus.

4.3.7 Super-resolved images and segment protein copy numbers of RecA-mGFP

The segment protein copy numbers as reported in Results Section 2.3 and Figure S3, were calculated
with ImageJ, using as input the segments corresponding to bundles or foci, instead of whole cells.
These segments were obtained starting from the coordinates of localized, tracked foci in Slimfield
analysis, from the latter stages of photobleaching below a threshold stoichiometry of 2 molecules;
these were imported into ThunderSTORM software [90]. The Visualization module to build a
pointillistic super-resolved image at 40 nm lateral spatial precision (as shown in Figures 3B,E) at 5×
upscaling (11 nm pixel size), which was then smoothed with a Gaussian filter of 4 pixels' width, and
automatically Otsu thresholded to generate a superresolved binary mask. The masks were then
expanded by a distance equal to the widefield resolution of 17 pixels (~180 nm) to match the features
in the Slimfield images. The integrated intensities were extracted, as in [91], from the sum of
fluorescent pixel counts in the Slimfield images (Analyze Particles > Multi-Measure function in ImageJ)
less the area multiplied by the camera pixel dark value. To yield segment protein copy numbers (Figure
S3A), the resulting integrated intensities were corrected for the relative autofluorescence, by
subtracting the integrated intensity of parental cells adjusted by by the ratio of mean segment area.
The Multi-Measure output also included the Feret diameter of each segment which was used as an
estimate of its end-to-end length (Figure 3D-F).

4.3.8 Photobleaching rates

Photobleaching rates were estimated by fitting the decrease in background-subtracted cellular
protein copy number or mean track stoichiometry over the exposure time using MATLAB *cftool*. The
fit consisted of a monoexponential decay to the first 10 frames with variable initial intensity and decay
constant, but with a baseline fixed to the average intensity after 50 frames. Fits were then refined to
include only data within the initial $1/e$ decay time (Table S2). RecA-mGFP and RecB-sfGFP photobleach

decay times were consistently dissimilar at 13 ± 2 and 6 ± 1 frames respectively; sfGFP is typically
several-fold less photostable than comparable enhanced GFPs under high intensity illumination [92].

4.3.9 Statistical tests

We performed multiple statistical comparisons on each set of tracked data (typically ~5: number of
tracks, stoichiometry, periodicity, diffusivity, copy number), which we account for using the standard
Bonferroni correction; the significance level is adjusted downwards by a factor of the number of
comparisons, $\alpha = 0.05/5 = 0.01$).

4.4 Gene expression assays

4.4.1 Quantitative PCR (qPCR)

Treated and untreated cultures were grown as in section 4.1. Total RNA was then isolated using
Monarch Total RNA Miniprep Kit (New England Biolabs). cDNA was synthesised from 350 ng of total
RNA from each sample using Superscript IV reverse transcriptase (Invitrogen) according to the
manufacturer's instructions using random hexamer primer (ThermoScientific).

The cDNA was then subjected to qPCR using Fast SYBR Green Master Mix (ThermoFisher) in a
QuantStudio 3 Real-Time PCR System. The *recA* primer pair amplified *recA* cDNA in the wild type and
both *recA* and *recA-GFP* mRNA in the labeled strain. *recA-GFP* alone in the labeled strain was amplified
using *GFP* primer pair. 16s rRNA was used as a housekeeping control.

Data obtained was analysed using the standard curve method [51]. Standard curves were generated
from serial dilutions of PCR products with known concentrations derived from genomic DNA. Fold
increase in mRNA levels was calculated by dividing the values obtained for treated mRNA with the
untreated. Results are shown in Figure S4A.

Primer	Sequence 5'-3'	Complementary region
oAS216	GCAGGCACTGGAAATCTGTG	*recA* (forward)
oAS217	GCCGATTTCGCCTTCGATTTC	*recA* (reverse)
oAS220	CTACAAGACACGTGCTGAAGTC	*GFP* (forward)
oAS221	AGTTGTATTCCAATTTGTGTCCAAGAATG	*GFP* (reverse)
oAS23	GTAGAATTCCAGGTGTAGCGGTG	16s rRNA (forward)
oAS24	CATCGTTTACGGCGTGGACTACCAG	16s rRNA (reverse)

Table 2. Primers used for qPCR to quantify mRNA of *recA*, *recA-gfp* and housekeeping gene *rrsA*.

4.4.2 Western blots

Six samples of normalised *E. coli* cell cultures were prepared as above (section 4.1) in 1 ml aliquots at
OD_{600} ~0.2. The cells were isolated using centrifugation at 10,000 × *g* for 2 min in a microfuge to
prepare them for SDS-PAGE / immuno-detection. The cell pellets were resuspended in 75 µl of SDS
loading buffer and boiled for 5 min at 95°C before application of 15 µl onto a 4-20 % gradient gel. The
gel was subsequently transferred to nitrocellulose and the membrane was placed in blocking solution
(PBS-T, 5% (w/v) non-fat milk) for 3 h. Primary antibody (anti-RecA) was incubated at 1/500 overnight
in blocking solution before the membrane was washed (4 x 5 min) in blocking solution. Secondary
antibody (goat anti rabbit-HRP) was incubated at 1/2,000 dilution for 4 h in blocking solution before
the membrane was again washed (4 x 5 min) in blocking solution. A final wash in PBS was performed
before development using ECL and image acquisition (iBRIGHT). Results are shown in Figure S4B.

Author Contributions:

A.P-D, A.S, and M.L. designed the research; A.S. cultured and treated cells; A.P-D performed
microscopy and data analysis, visualization and curation, J.S. and L.F. wrote and validated
segmentation software; A.P-D and A.S drafted the paper; A.P-D, A.S, J.S. and M.L. edited the paper;

M.L. supervised and administered the project. This research was funded by BBSRC, grant numbers
BB/P000746/1 and BB/N006453/1, and EPSRC, grant number EP/T002166/1.

Data Availability Statement: The raw imaging data is available on reasonable from
https://doi.org/10.5281/zenodo.6639101; the MATLAB tracking analysis code can be found at
https://github.com/alex-payne-dwyer/single-molecule-tools-alpd. The U-Net image segmentation
architecture originated from code obtained from the NEUBIAS Academy workshop
(http://eubias.org/NEUBIAS/training-schools/neubias-academy-home/).

Acknowledgments:
The authors thank Dr. Christian Lesterlin for the gift of the RecA-mGFP strain and Prof. Meriem El
Karoui for the RecB-sfGFP strain. We thank the Biosciences Technology Facility for technical assistance
with gene expression assays.

Conflicts of Interest:
The authors declare no conflict of interest. The funders had no role in the design of the study; in the
collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to
publish the results.

References

1. Reyes-Lamothe R, Sherratt DJ, Leake MC. 2010 Stoichiometry and architecture of active DNA replication machinery in Escherichia coli. *Science* **328**, 498–501. (doi:10.1126/science.1185757)

2. Surova O, Zhivotovsky B. 2013 Various modes of cell death induced by DNA damage. *Oncogene* **32**, 3789–97. (doi:10.1038/onc.2012.556)

3. Lambert S, Carr AM. 2013 Impediments to replication fork movement: Stabilisation, reactivation and genome instability. *Chromosoma* **122**, 33–45. (doi:10.1007/s00412-013-0398-9)

4. Tomasz M. 1995 Mitomycin C: small, fast and deadly (but very selective). *Chem. Biol.* **2**, 575–9. (doi:10.1016/1074-5521(95)90120-5)

5. Arranz-Marquez E, Katsanos A, Kozobolis VP, Konstas AGP, Teus MA. 2019 A Critical Overview of the Biological Effects of Mitomycin C Application on the Cornea Following Refractive Surgery. *Adv. Ther.* **36**, 786–797. (doi:10.1007/s12325-019-00905-w)

6. Weng M-W, Zheng Y, Jasti VP, Champeil E, Tomasz M, Wang Y, Basu AK, Tang M-S. 2010 Repair of mitomycin C mono-and interstrand cross-linked DNA adducts by UvrABC: a new model. *Nucleic Acids Res.* **38**, 6976–6984. (doi:10.1093/nar/gkq576)

7. Bizanek R, McGuinness BF, Nakanishi K, Tomasz M, McGuinness BF, Nakanishi K. 1992 Isolation and Structure of an Intrastrand Cross-Link Adduct of Mitomycin C and DNA. *Biochemistry* **31**, 3084–3091. (doi:10.1021/bi00127a008)

8. Dronkert MLG, Kanaar R. 2001 Repair of DNA interstrand cross-links. *Mutat. Res.* **486**, 217–247. (doi:10.1016/s0921-8777(01)00092-1)

9. Legerski RJ. 2010 Repair of DNA interstrand cross-links during S phase of the mammalian cell cycle. *Environ. Mol. Mutagen.* **51**, 540–551. (doi:10.1002/em.20566)

10. Smith GR. 2012 How RecBCD Enzyme and Chi Promote DNA Break Repair and Recombination: a Molecular Biologist's View. *Microbiol. Mol. Biol. Rev.* **76**, 217–28. (doi:10.1128/mmbr.05026-11)

11. Syeda AH, Hawkins M, McGlynn P. 2014 Recombination and Replication. *Cold Spring Harb. Perspect. Biol.* **6**, 1–14. (doi:10.1101/cshperspect.a016550)

12. Amundsen SK, Smith GR. 2019 The RecB helicase-nuclease tether mediates Chi hotspot control of RecBCD enzyme. *Nucleic Acids Res.* **47**, 197–209. (doi:10.1093/nar/gky1132)

13. Taylor AF, Smith GR. 1995 Monomeric RecBCD enzyme binds and unwinds DNA. *J. Biol. Chem.* **270**, 24451–8. (doi:10.1074/jbc.270.41.24451)

14. Howard-Flanders P, West SC, Stasiak A. 1984 Role of RecA protein spiral filaments in genetic recombination. *Nature* **309**, 215–220. (doi:10.1038/309215a0)

15. Kowalczykowski SC, Eggleston AK. 1994 Homologous pairing and DNA strand-exchange proteins. *Annu. Rev. Biochem.* **63**, 991–1043. (doi:10.1146/annurev.bi.63.070194.005015)

16. Mazin A V., Kowalczykowski SC. 1999 A novel property of the RecA nucleoprotein filament: Activation of double-stranded DNA for strand exchange in trans. *Genes Dev.* **13**, 2005–16. (doi:10.1101/gad.13.15.2005)

17. Wiktor J, Gynnå AH, Leroy P, Larsson J, Coceano G, Testa I, Elf J. 2021 RecA finds homologous DNA by reduced dimensionality search. *Nature* **597**. (doi:10.1038/s41586-021-03877-6)

18. Windgassen TA, Wessel SR, Bhattacharyya B, Keck JL. 2018 Mechanisms of bacterial DNA replication restart. *Nucleic Acids Res.* **46**, 504–519. (doi:10.1093/nar/gkx1203)

19. Dimude JU, Midgley-Smith SL, Rudolph CJ. 2018 Replication-transcription conflicts trigger extensive DNA degradation in Escherichia coli cells lacking RecBCD. *DNA Repair (Amst).* **70**, 37–48. (doi:10.1016/j.dnarep.2018.08.002)

20.   Repar J, Briški N, Buljubašić M, Zahradka K, Zahradka D. 2013 Exonuclease VII is involved in
'reckless' DNA degradation in UV-irradiated Escherichia coli. *Mutat. Res. - Genet. Toxicol.*
Environ. Mutagen. **750**, 96–104. (doi:10.1016/j.mrgentox.2012.10.005)

21.   Ghodke H *et al.* 2019 Spatial and temporal organization of RecA in the Escherichia coli DNA-
damage response. *Elife* **8**, e42761. (doi:10.7554/eLife.42761)

22.   Khil PP, Camerini-Otero RD. 2002 Over 1000 genes are involved in the DNA damage response
of Escherichia coli. *Mol. Microbiol.* **44**, 89–105. (doi:10.1046/j.1365-2958.2002.02878.x)

23.   Adikesavan AK, Katsonis P, Marciano DC, Lua R, Herman C, Lichtarge O. 2011 Separation of
Recombination and SOS Response in Escherichia coli RecA Suggests LexA Interaction Sites. *Plos*
Genet. **7**, e1002244. (doi:10.1371/journal.pgen.1002244)

24.   Gruenig MC, Renzette N, Long E, Chitteni-Pattu S, Inman RB, Cox MM, Sandler SJ. 2008 RecA-
mediated SOS induction requires an extended filament conformation but no ATP hydrolysis.
Mol. Microbiol. **69**, 1165–1179. (doi:10.1111/j.1365-2958.2008.06341.x)

25.   Butala M *et al.* 2011 Interconversion between bound and free conformations of LexA
orchestrates the bacterial SOS response. *Nucleic Acids Res.* **39**. (doi:10.1093/nar/gkr265)

26.   Podlesek Z, Žgur Bertok D. 2020 The DNA Damage Inducible SOS Response Is a Key Player in
the Generation of Bacterial Persister Cells and Population Wide Tolerance. *Front. Microbiol.* **11**.
(doi:10.3389/fmicb.2020.01785)

27.   Memar MY, Yekani M, Celenza G, Poortahmasebi V, Naghili B, Bellio P, Baghi HB. 2020 The
central role of the SOS DNA repair system in antibiotics resistance: A new target for a new
infectious treatment strategy. *Life Sci.* **262**. (doi:10.1016/j.lfs.2020.118562)

28.   Rehrauer WM, Kowalczykowski SC. 1996 The DNA binding site(s) of the Escherichia coli RecA
protein. *J. Biol. Chem.* **271**, 11996–12002. (doi:10.1074/jbc.271.20.11996)

29.   Handa N, Amitani I, Gumlaw N, Sandler SJ, Kowalczykowski SC. 2009 Single Molecule Analysis
of a Red Fluorescent RecA Protein Reveals a Defect in Nucleoprotein Filament Nucleation That
Relates to Its Reduced Biological Functions. *J. Biol. Chem.* **284**, 18664–18673.
(doi:10.1074/jbc.M109.004895)

30.   Ivancic-Bace I, Vlasic I, Salaj-Smic E, Brcic-Kostic K, Ivančić-Baće I, Vlašić I, Salaj-Šmic E, Brčić-
Kostić K. 2006 Genetic evidence for the requirement of RecA loading activity in SOS induction
after UV irradiation in Escherichia coli. *J. Bacteriol.* **188**, 5024–5032. (doi:10.1128/JB.00130-06)

31.   Anderson DG, Kowalczykowski SC. 1998 Reconstitution of an SOS response pathway:
Derepression of transcription in response to DNA breaks. *Cell* **95**, 975–9. (doi:10.1016/S0092-
8674(00)81721-3)

32.   Adams D *et al.* 2007 Selection of Reca+ Recombinant Cosmids - an Easy Method for Making
Reca Strains Temporarily Rec+, Permitting P1-Mediated Transduction in a Reca- Background
and Transduction of a Reca- Mutation. *J. Biol. Chem.* **8**, 88–94. (doi:10.1074/jbc.M110.194407)

33.   Plank M, Wadhams GH, Leake MC. 2009 Millisecond timescale slimfield imaging and
automated quantification of single fluorescent protein molecules for use in probing complex
biological processes. *Integr. Biol.* **1**, 602–612. (doi:10.1039/b907837a)

34.   Lesterlin C, Ball G, Schermelleh L, Sherratt DJ. 2014 RecA bundles mediate homology pairing
between distant sisters during DNA break repair. *Nature* **506**, 249–253.
(doi:10.1038/nature12868)

35.   Lepore A, Taylor H, Landgraf D, Okumus B, Jaramillo-Riveri S, Mclaren L, Bakshi S, Paulsson J, El
Karoui & M. 2019 Quantification of very low-abundant proteins in bacteria using the HaloTag
and epi-fluorescence microscopy. *Sci. Rep.* **9**, 7902. (doi:10.1038/s41598-019-44278-0)

36.   Lenn T, Leake MC. 2012 Experimental approaches for addressing fundamental biological
questions in living, functioning cells with single molecule precision. *Open Biol.* **2**, 120090.
(doi:10.1098/rsob.120090)

37. Wollman AJM, Leake MC. 2015 Millisecond single-molecule localization microscopy combined with convolution analysis and automated image segmentation to determine protein concentrations in complexly structured, functional cells, one cell at a time. *Faraday Discuss.* **184**, 401–424. (doi:10.1039/C5FD00077G)

38. Wollman AJM *et al.* 2022 Critical roles for EGFR and EGFR-HER2 clusters in EGF binding of SW620 human carcinoma cells. *J. R. Soc. Interface* **19**, 20220088. (doi:10.1098/rsif.2022.0088)

39. Manley S, Gillette JM, Patterson GH, Shroff H, Hess HF, Betzig E, Lippincott-Schwartz J. 2008 High-density mapping of single-molecule trajectories with photoactivated localization microscopy. *Nat. Methods* **5**, 155–157. (doi:10.1038/nmeth.1176)

40. Stracy M, Kapanidis AN. 2017 Single-molecule and super-resolution imaging of transcription in living bacteria. *Methods* **120**, 103–114. (doi:10.1016/j.ymeth.2017.04.001)

41. Jin X *et al.* 2021 Membraneless organelles formed by liquid-liquid phase separation increase bacterial fitness. *Sci. Adv.* **7**, eabh2929. (doi:10.1126/sciadv.abh2929)

42. Story RM, Weber IT, Steitz TA. 1992 The structure of the E. coli recA protein monomer and polymer. *Nature* **355**, 318–325. (doi:10.1038/355318a0)

43. Long JE, Renzette N, Centore RC, Sandler SJ. 2008 Differential Requirements of Two recA Mutants for Constitutive SOS Expression in Escherichia coli K-12. *PLoS One* **3**, e4100. (doi:10.1371/journal.pone.0004100)

44. Lee AM, Singleton SF. 2006 Intersubunit electrostatic complementarity in the RecA nucleoprotein filament regulates nucleotide substrate specificity and conformational activation. *Biochemistry* **45**, 4514–4529. (doi:10.1021/bi052296x)

45. Wang Y, Bortner C, Griffith J. 1993 Reca Binding to Bulge-Containing and Mismatch-Containing Dnas - Certain Single-Base Mismatches Provide Strong Signals for Reca Binding Equal to Multiple Base Bulges. *J. Biol. Chem.* **268**, 17571–17577.

46. Schlacher K, Cox MM, Woodgate R, Goodman MF. 2006 RecA acts in trans to allow replication of damaged DNA by DNA polymerase V. *Nature* **442**, 883–887. (doi:10.1038/nature05042)

47. McGrew DA, Knight KL. 2003 Molecular design and functional organization of the RecA protein. *Crit. Rev. Biochem. Mol. Biol.* **38**, 385–432. (doi:10.1080/10409230390242489)

48. Dapa T, Fleurier S, Bredeche MF, Matic I. 2017 The SOS and RpoS regulons contribute to bacterial cell robustness to genotoxic stress by synergistically regulating DNA polymerase pol II. *Genetics* **206**. (doi:10.1534/genetics.116.199471)

49. Miller H, Zhou Z, Wollman AJM, Leake MC. 2015 Superresolution imaging of single DNA molecules using stochastic photoblinking of minor groove and intercalating dyes. *Methods* **88**, 81–88. (doi:10.1016/j.ymeth.2015.01.010)

50. Leake MC, Chandler JH, Wadhams GH, Bai F, Berry RM, Armitage JP. 2006 Stoichiometry and turnover in single, functioning membrane protein complexes. *Nature* **443**, 355–358. (doi:10.1038/nature05135)

51. Peterson KR, Wertman KF, Mount DW, Marinus MG. 1985 Viability of Escherichia coli K-12 DNA adenine methylase (dam) mutants requires increased expression of specific genes in the SOS regulon. *MGG Mol. Gen. Genet.* **201**, 14–19. (doi:10.1007/BF00397979)

52. Renzette N, Gumlaw N, Nordman JT, Krieger M, Yeh SP, Long E, Centore R, Boonsombat R, Sandler SJ. 2005 Localization of RecA in Escherichia coli K-12 using RecA-GFP. *Mol. Microbiol.* **57**. (doi:10.1111/j.1365-2958.2005.04755.x)

53. Leake MC. 2014 Analytical tools for single-molecule fluorescence imaging in cellulo. *Phys. Chem. Chem. Phys.* **16**, 12635–47. (doi:10.1039/c4cp00219a)

54. Shi W-X, Larson RG. 2007 RecA-ssDNA filaments supercoil in the presence of single-stranded DNA-binding protein. *Biochem. Biophys. Res. Commun.* **357**, 755–760. (doi:10.1016/j.bbrc.2007.04.014)

55. Darmon E, Eykelenboom JK, Lopez-Vernaza MA, White MA, Leach DRF. 2014 Repair on the Go: E. Coli maintains a high proliferation rate while repairing a chronic DNA double-strand break. *PLoS One* **9**. (doi:10.1371/journal.pone.0110784)

56. Ellouze C, Takahashi M, Wittung P, Mortensen K, Schnarr M, Norden B, Nordén B, Norden B. 1995 Evidence for Elongation of the Helical Pitch of the Reca Filament Upon Atp and Adp Binding Using Small-Angle Neutron-Scattering. *Eur. J. Biochem.* **233**, 579–583. (doi:10.1111/j.1432-1033.1995.579_2.x)

57. Yu X, Jacobs SA, West SC, Ogawa T, Egelman EH. 2001 Domain structure and dynamics in the helical filaments formed by RecA and Rad51 on DNA. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 8419–8424. (doi:10.1073/pnas.111005398)

58. Zhao B *et al.* 2017 ATPase activity tightly regulates RecA nucleofilaments to promote homologous recombination. *Cell Discov.* **3**, 16053. (doi:10.1038/celldisc.2016.53)

59. Chimthanawala A, Parmar J, Kumar S, Iyer KS, Rao M, Badrinarayanan A. 2021 SMC protein RecN drives translocation and remodelling of RecA filament for homology search. *BiorXiv* , 2021.08.16.456443. (doi:10.1101/2021.08.16.456443)

60. Rajendram M *et al.* 2015 Anionic Phospholipids Stabilize RecA Filament Bundles in Escherichia coli. *Mol. Cell* **60**, 374–384. (doi:10.1016/j.molcel.2015.09.009)

61. Eldin S, Forget AL, Lindenmuth DM, Logan KM, Knight KL. 2000 Mutations in the N-terminal region of RecA that disrupt the stability of free protein oligomers but not RecA-DNA complexes. *J. Mol. Biol.* **299**, 91–101. (doi:10.1006/jmbi.2000.3721)

62. Roy R, Kozlov AG, Lohman TM, Ha T. 2009 SSB protein diffusion on single-stranded DNA stimulates RecA filament formation. *Nature* **461**, 1092–7. (doi:10.1038/nature08442)

63. Norden B, Takahashi. M, Takahashi M, Nordén B. 1994 Structure of RecA-DNA complex and mechanism of DNA strand exchange reaction in homologous recombination. *Adv. Biophys.* **30**, 1–35. (doi:10.1016/0065-227X(94)90009-4)

64. Galletto R, Amitani I, Baskin RJ, Kowalczykowski SC. 2006 Direct observation of individual RecA filaments assembling on single DNA molecules. *Nature* **443**, 875–8. (doi:10.1038/nature05197)

65. Hernández-Tamayo R, Steube N, Heimerl T, Hochberg G, Graumann PL. 2022 ATPase activity of B. subtilis RecA affects the dynamic formation of RecA filaments at DNA double strand breaks. *BiorXiv* , 2022.02.15.480544. (doi:10.1101/2022.02.15.480544)

66. Danilowicz C, Vietorisz E, Godoy-Carter V, Prévost C, Prentiss M. 2021 Influences of ssDNA-RecA Filament Length on the Fidelity of Homologous Recombination. *J. Mol. Biol.* **433**, 167143. (doi:10.1016/J.JMB.2021.167143)

67. Yu X, Egelman EH. 1997 The RecA hexamer is a structural homologue of ring helicases. *Nat. Struct. Biol.* **4**, 101–104. (doi:10.1038/nsb0297-101)

68. Yu X, VanLoock MS, Yang S, Reese JT, Egelman EH. 2004 What is the structure of the RecA-DNA filament? *Curr. Protein Pept. Sci.* **5**, 73–79. (doi:10.2174/1389203043486883)

69. Chen Z, Yang H, Pavletich NP. 2008 Mechanism of homologous recombination from the RecA-ssDNA/dsDNA structures. *Nature* **453**, 489–494. (doi:10.1038/nature06971)

70. Li GW, Burkhardt D, Gross C, Weissman JS. 2014 Quantifying absolute protein synthesis rates reveals principles underlying allocation of cellular resources. *Cell* **157**, 624–35. (doi:10.1016/j.cell.2014.02.033)

71. Stohl EA, Brockman JP, Burkle KL, Morimatsu K, Kowalczykowski SC, Seifert HS. 2003 Escherichia coli RecX inhibits RecA recombinase and coprotease activities in vitro and in vivo. *J. Biol. Chem.* **278**, 2278–85. (doi:10.1074/jbc.M210496200)

72. Giacomoni PU. 1982 Induction by mitomycin C of recA protein synthesis in bacteria and spheroplasts. *J. Biol. Chem.* **257**, 14932–14936. (doi:10.1016/s0021-9258(18)33373-8)

1082   73.   Soufi B, Krug K, Harst A, Macek B. 2015 Characterization of the E. coli proteome and its
1083   modifications during growth and ethanol stress. *Front. Microbiol.* **6**, 103.
1084   (doi:10.3389/fmicb.2015.00103)

1085   74.   Elowitz MB, Surette MG, Wolf PE, Stock JB, Leibler S. 1999 Protein mobility in the cytoplasm of
1086   Escherichia coli. *J. Bacteriol.* **181**, 197–203. (doi:10.1128/jb.181.1.197-203.1999)

1087   75.   Nenninger A, Mastroianni G, Mullineaux CW. 2010 Size dependence of protein diffusion in the
1088   cytoplasm of Escherichia coli. *J. Bacteriol.* **192**, 4535–40. (doi:10.1128/JB.00284-10)

1089   76.   Singleton MR, Dillingham MS, Gaudier M, Kowalczykowski SC, Wigley DB. 2004 Crystal
1090   structure of RecBCD enzyme reveals a machine for processing DNA breaks. *Nature* **432**, 187–
1091   93.

1092   77.   Burby PE, Simmons LA. 2019 A bacterial DNA repair pathway specific to a natural antibiotic.
1093   *Mol. Microbiol.* **111**, 338–353. (doi:10.1111/mmi.14158)

1094   78.   Peng X, Ghosh AK, Van Houten B, Greenberg MM. 2010 Nucleotide excision repair of a DNA
1095   interstrand cross-link produces single- and double-strand breaks. *Biochemistry* **49**.
1096   (doi:10.1021/bi901603h)

1097   79.   Wood RD. 2010 Mammalian nucleotide excision repair proteins and interstrand crosslink
1098   repair. *Environ. Mol. Mutagen.* **51**. (doi:10.1002/em.20569)

1099   80.   Syeda AH, Wollman AJM, Hargreaves AL, Howard JAL, Brüning J-G, McGlynn P, Leake MC. 2019
1100   Single-molecule live cell imaging of Rep reveals the dynamic interplay between an accessory
1101   replicative helicase and the replisome. *Nucleic Acids Res.* **47**, 6287–6298.
1102   (doi:10.1093/nar/gkz298)

1103   81.   Verhoeven EEAA, Kesteren M Van, Moolenaar GF, Visse R, Goosen N, Van Kesteren M,
1104   Moolenaar GF, Visse R, Goosen N. 2000 Catalytic sites for 3' and 5' incision of Escherichia coli
1105   nucleotide excision repair are both located in UvrC. *J. Biol. Chem.* **275**, 5120–3.
1106   (doi:10.1074/jbc.275.7.5120)

1107   82.   Massoni SC, Leeson MC, Long JE, Gemme K, Mui A, Sandler SJ. 2012 Factors Limiting SOS
1108   Expression in Log-Phase Cells of Escherichia coli. *J. Bacteriol.* **194**, 5325–5333.
1109   (doi:10.1128/JB.00674-12)

1110   83.   Reuter A, Hilpert C, Dedieu-Berne A, Lematre S, Gueguen E, Launay G, Bigot S, Lesterlin C. 2021
1111   Targeted-antibacterial-plasmids (TAPs) combining conjugation and CRISPR/Cas systems
1112   achieve strain-specific antibacterial activity. *Nucleic Acids Res.* **49**, 3584–3598.
1113   (doi:10.1093/nar/gkab126)

1114   84.   Renzette N, Sandler SJ. 2008 Requirements for ATP binding and hydrolysis in RecA function in
1115   Escherichia coli. *Mol. Microbiol.* **67**. (doi:10.1111/j.1365-2958.2008.06130.x)

1116   85.   Llorente-Garcia I *et al.* 2014 Single-molecule in vivo imaging of bacterial respiratory complexes
1117   indicates delocalized oxidative phosphorylation. *Biochim. Biophys. Acta - Bioenerg.* **1837**, 811–
1118   24. (doi:10.1016/j.bbabio.2014.01.020)

1119   86.   Dresser L, Hunter P, Yendybayeva F, Hargreaves AL, Howard JAL, Evans GJO, Leake MC, Quinn
1120   SD. 2021 Amyloid-β oligomerization monitored by single-molecule stepwise photobleaching.
1121   *Methods* **193**, 80–95. (doi:10.1016/j.ymeth.2020.06.007)

1122   87.   Shepherd JW, Payne-Dwyer AL, Lee J-E, Syeda A, Leake MC. 2021 Combining single-molecule
1123   super-resolved localization microscopy with fluorescence polarization imaging to study cellular
1124   processes. *JPhys Photonics* **3**. (doi:10.1088/2515-7647/ac015d)

1125   88.   Robson A, Burrage K, Leake MC. 2013 Inferring diffusion in single live cells at the single-
1126   molecule level. *Philos. Trans. R. Soc. B Biol. Sci.* **368**, 20120029. (doi:10.1098/rstb.2012.0029)

1127   89.   Leake MC, Wilson D, Bullard B, Simmons RM, Bubb MR. 2003 The elasticity of single kettin
1128   molecules using a two-bead laser-tweezers assay. In *FEBS Letters*, pp. 55–60.
1129   (doi:10.1016/S0014-5793(02)03857-7)

90. Ovesný M, Křížek P, Borkovec J, Švindrych Z, Hagen GM. 2014 ThunderSTORM: A comprehensive ImageJ plug-in for PALM and STORM data analysis and super-resolution imaging. *Bioinformatics* **30**, 2389–2390. (doi:10.1093/bioinformatics/btu202)

91. Payne-Dwyer AL, Leake MC. 2022 Single-molecular quantification of flowering control proteins within nuclear condensates in live whole Arabidopsis root. In *Chromosome Architecture: Methods in Molecular Biology* (ed MC Leake), p. 2476. New York, NY: Springer US, Humana. (doi:10.1007/978-1-0716-2221-6)

92. Pédelacq JD, Cabantous S, Tran T, Terwilliger TC, Waldo GS. 2006 Engineering and characterization of a superfolder green fluorescent protein. *Nat. Biotechnol.* **24**, 79–88. (doi:10.1038/nbt1172)