Fluorescence Activation Mechanism and Imaging of Drug Permeation with New Sensors for Smoking-Cessation Ligands

Authors: Aaron L. Nichols*[1], Zack Blumenfeld*[1,2], Chengcheng Fan*[5], Laura Luebbert*[1,3], Annet E. M. Blom[5], Bruce N. Cohen[1], Jonathan S. Marvin[4], Philip M. Borden[4], Charlene H. Kim[1], Anand K. Muthusamy[5], Amol V. Shivange[1], Hailey J. Knox[5], Hugo Rego Campello[6], Jonathan H. Wang[1], Dennis A. Dougherty[5], Loren L. Looger[4], Timothy Gallagher[6], Douglas C. Rees[5,7], Henry A. Lester[1a]

* Co-first authors

[1] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA

[2] Keck School of Medicine, University of Southern California, Los Angeles, CA

[3] Institute of Biology, Leiden University, Leiden, The Netherlands

[4] Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA

[5] Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA

[6] School of Chemistry, University of Bristol, Bristol, United Kingdom

[7] Howard Hughes Medical Institute, California Institute of Technology, Pasadena, CA

[a] Corresponding author
E-mail: lester@caltech.edu

ABSTRACT

Nicotinic partial agonists provide an accepted aid for smoking cessation and thus contribute to
decreasing tobacco-related disease. Improved drugs constitute a continued area of study.
However, there remains no reductionist method to examine the cellular and subcellular
pharmacokinetic properties of these compounds in living cells. Here, we developed new intensity-
based drug sensing fluorescent reporters ("iDrugSnFRs") for the nicotinic partial agonists
dianicline, cytisine, and two cytisine derivatives – 10-fluorocytisine and 9-bromo-10-ethylcytisine.
We report the first atomic-scale structures of liganded periplasmic binding protein-based
biosensors, accelerating development of iDrugSnFRs and also explaining the activation
mechanism. The nicotinic iDrugSnFRs detect their drug partners in solution, as well as at the
plasma membrane (PM) and in the endoplasmic reticulum (ER) of cell lines and mouse
hippocampal neurons. At the PM, the speed of solution changes limits the growth and decay rates
of the fluorescence response in almost all cases. In contrast, we found that rates of membrane
crossing differ among these nicotinic drugs by > 30 fold. The new nicotinic iDrugSnFRs, in
combination with previously described nicotine and varenicline sensors, provide insight into the
real-time pharmacokinetic properties of nicotinic agonists and provide a methodology whereby
iDrugSnFRs can inform both pharmaceutical neuroscience and addiction neuroscience.

INTRODUCTION

Because drugs for smoking cessation partially help to decrease tobacco-related disease, drug discovery efforts have targeted nicotinic acetylcholine receptors (nAChRs). The addictive tobacco alkaloid nicotine itself, via transdermal patches and other devices, remains available for people trying to quit smoking.

Prior work suggests that partial agonists with lower efficacy than nicotine could serve as effective smoking-cessation drugs (Rose, et al., 1994), and efforts continue in that direction (Rollema & Hurst, 2018). Another plant alkaloid, (-)-cytisine (also called cytisinicline and Tabex®), an α4β2 nAChR partial agonist, served as a basis for the synthesis of analogs which have not yet entered the clinic (Chellappan, Xiao, Tueckmantel, Kellar, & Kozikowski, 2006; Houllier, Gouault, Lasne, & Rouden, 2006; Imming, Klaperski, Stubbs, Seitz, & Gundisch, 2001; Kozikowski, et al., 2007; Marcaurelle, Johannes, Yohannes, Tillotson, & Mann, 2009; Philipova, et al., 2015; Rouden, et al., 2002). Varenicline (Chantix®) has four rings, two more than nicotine or cytisine, and is currently the only FDA-approved smoking-cessation drug, but the modest quit rate of ~18% at 12 months invites further investigation (Coe, et al., 2005; Mills, Wu, Spurden, Ebbert, & Wilson, 2009). Dianicline, another tetracyclic compound, was discontinued after unfavorable Phase III clinical trials (Cohen, et al., 2003; Fagerstrom & Balfour, 2006).

A nicotinic ligand for smoking cessation must satisfy at least three criteria (Rollema, et al., 2010; Tashkin, 2015). 1) It must enter the brain, where the most nicotine-sensitive nAChRs (α4β2) occur. It must also 2) activate α4β2 nAChRs with an EC_{50} sufficient to reduce cravings and withdrawal (1–2 μM). It must also 3) block nicotine binding to reduce the reward phase of smoking (2 to 30 min). Varenicline meets these criteria, while cytisine (low brain penetration) and dianicline (EC_{50} =18 μM) each fail one of the criteria (Rollema, et al., 2010).

Membrane permeation is interesting for investigating and treating nicotine addiction in at least two ways. Firstly, note criterion #1 above. Enhancing the membrane permeability of cytisine analogs and probing nAChR subtype selectivity was addressed via direct functionalization of (-)-cytisine within the pyridone ring (Rego Campello, et al., 2018). Two of the resulting derivatives, 10-fluorocytisine and 9-bromo-10-ethylcytisine, have cytisine-like EC_{50} for the α4β2 nAChRs, but more positive calculated $LogD_{pH7.4}$ values, suggesting greater membrane permeability at the nearly neutral pH of the blood, brain, and cytoplasm (Blom, Campello, Lester, Gallagher, & Dougherty, 2019). Estimates of $LogD_{pH7.4}$ are inexact, extrapolated, or rely on algorithmic calculations whose results differ over two log units for individual molecules (Pienko, Grudzien, Taciak, & Mazurek, 2016). These estimates have unknown applicability to biological membranes at the $LogD_{pH7.4}$ values < 0 that characterize varenicline, dianicline, and the cytisine analogs.

Secondly, nicotine dependence involves one or more "inside-out" mechanisms. Nicotine itself ($logD_{pH7.4}$ 0.99) enters the endoplasmic reticulum (ER), binds to nascent nAChRs, becomes a pharmacological chaperone for the nAChRs, and eventually causes selective upregulation of these receptors on the plasma membrane (PM) (Henderson & Lester, 2015). For this reason, it is especially important to understand permeation into the ER.

These two neuroscience aspects of nicotinic ligands—pharmaceutical science and addiction science—call for direct measurements of drug movements in living cells. We previously explored the subcellular pharmacokinetics of nicotine and varenicline in immortalized cell lines and cultured neurons using the iDrugSnFRs iNicSnFR3a and iNicSnFR3b, to visualize that these nicotinic agonists enter the ER within seconds of drug application and exit equally rapidly from the ER upon extracellular washing (Shivange, et al., 2019). That nicotine diffuses across cellular membranes in seconds has been suspected for decades: nicotine crosses six plasma membranes to enter the brain within 20 s, providing a "buzz". That varenicline becomes trapped in acidic vesicles suggests appreciable membrane permeation but may also underlie unwanted effects (Govind, et al., 2017; Le Houezec, 2003).

We sought to generate and to apply additional intensity-based drug-sensing fluorescent reporters ("iDrugSnFRs") for candidate smoking cessation drugs: dianicline, cytisine, 10-

95   fluorocytisine, and 9-bromo-10-ethylcytisine. We hypothesized that a family of newly developed
96   iDrugSnFRs would enable quantifiable fluorescence signals that compare the differences in
97   permeation among these compounds.

RESULTS

Generation of additional nicotinic iDrugSnFRs: structural tactic

To generate iDrugSnFRs for cytisine and dianicline, we followed two converging tactics. In
the "structure-based" tactic, we obtained the first structural data for OpuBC-based SnFRs bound
by nicotinic ligands (nicotine and varenicline) (Fig. 1, Supplementary Table 1). Crystals of
iNicSnFR3adt in the presence of 10 mM nicotine diffracted to 2.95 Å resolution (PDB 7S7U).
Overall, the liganded PBP domain of iNicSnFR3adt adopts a closed conformation (Fig. 1A). In the
binding pocket between the top and bottom lobes of the PBP, we observed an "avocado" shaped
electron density in the nicotine binding site, enclosed by several aromatic residues (Fig. 1B). The
combination of protonation/deprotonation and the rotatable bond of nicotine (Elmore & Dougherty,
2000) vitiate unambiguously localizing it within the binding pocket
We obtained an unambiguous ligand placement for iNicSnFR3adt in the presence of 10 mM
varenicline in the same crystallization condition. Crystals of iNicSnFR3adt with varenicline bound
were isomorphous to those of the nicotine-bound crystals and diffracted to 3.2 Å resolution (PDB
7S7T). While the protein structure (Fig. 1D) is identical to that of the nicotine bound structure (Fig.
1A), the rigidity and additional ring of varenicline allowed us to unambiguously localize it in the
binding pocket. Varenicline is enclosed by the same aromatic residues as nicotine, forming cation-
π interactions with Tyr65 and Tyr357, in addition to other interactions with the pocket residues
(Fig. 1E).
The data confirm that similar ligand-induced conformational changes occur in the periplasmic
binding protein (PBP) for nicotine, varenicline, ACh (Borden, et al., 2019), and choline (Fan, 2020)
(Figure 1-figure supplement 1). These changes resemble those for other OpuBC PBPs
(Schiefner, et al., 2004).
In the full iDrugSnFR, in the apo state, the Glu78 in linker 1 approaches within ~ 2.5 Å of the
oxygen of the tyrosine fluorophore (Figure 1E1) (PDB 7S7V). Figure 1E2 provides structural
details confirming the hypothesis (Barnett, Hughes, & Drobizhev, 2017; Nasu, Shen, Kramer, &
Campbell, 2021) that in the liganded state, Glu78 has moved away, presumably allowing the
fluorescent tyrosinate to form. We term this mechanism the "candle snuffer".

Generation of additional nicotinic iDrugSnFRs: mutational tactic

In the mutational tactic, we screened each drug shown in Figure 1-figure supplement 2
against a panel of biosensors that included iNicSnFR3a and iNicSnFR3b (Shivange, et al., 2019)
and iAChSnFR (Borden, et al., 2019) as well as intermediate constructs from their development
process. From this screen, we chose sensors with the lowest EC_{50} for each drug as our starting
protein for iDrugSnFR evolution.
Because the candle snuffer mechanism explains several details of the agonist- and pH-
sensitivity of both iNicSnFR3a and iSketSnFR (see Discussion), we presume that it represents a
general mechanism for OpuBC-cpGFP SnFRs. We did not mutate residues that lie (in 3D space)
between the binding site and linkers.
For dianicline and cytisine separately, we incrementally applied site-saturation mutagenesis
(SSM) to first and second shell amino acid positions within the binding pocket. We evaluated each
biosensor and drug partner in lysate from *E. coli* and carried forward the biosensor with the highest
S-slope to the subsequent round. S-slope, $\frac{\Delta F}{F_0}/[ligand]$ at the beginning of the dose-response

relation, emphasizes the response to ligand concentrations in the pharmacologically relevant
range (Bera, et al., 2019). Table 1 and Fig. 2 summarize dose-response relations for the optimized
sensors. The dianicline sensor, iDianiSnFR, has EC_{50} 6.7 ± 0.3 µM, $\Delta F_{max}/F_0$ 7.4 ± 0.1, and S-
slope 1.1. The cytisine sensor, iCytSnFR, has EC_{50} 9.4 ± 0.8 µM, $\Delta F_{max}/F_0$ 5.0 ± 0.2, S-slope 0.5
(Table 1, Fig. 2A-B). After generating iCytSnFR, we performed additional SSM to progress from
iCytSnFR to SnFRs for 10-fluorocytisine and 9-bromo-10-ethylcytisine. This optimization gave us
iCyt_F_SnFR (EC_{50} 1.4 ± 0.04 µM, $\Delta F_{max}/F0$ 7.9 ± 0.1, S-slope 5.6) and iCyt_BrEt_SnFR (EC_{50}
5.7 ± 0.1 µM, $\Delta F_{max}/F0$ 4.0 ± 0.03, and S-slope 0.7) (Table 1, Fig. 2C-D).

Figure 1. Apo and ligand-bound structures of iNicSnFR3adt (dt indicates that His$_6$ and Myc tags have been removed to aid crystallization). To form an iDrugSnFR, a circularly permuted GFP molecule, flanked by two 4-residue linking sequences, is inserted into a PBP at a position (77-78, in our numbering system) that changes backbone Φ-Ψ angles between the apo and liganded PBP. **(A)** Overall conformation of iNicSnFR3adt crystallized with nicotine; an electron density appears at the nicotine binding site (PDB 7S7U). **(B)** iNicSnFR3adt binding site residues. **(C)** Overall conformation of iNicSnFR3adt with varenicline bound (PDB 7S7T). (**D**) iNicSnFR3adt binding site with varenicline present. **(E)** Aspects of the PBP-Linker1-cpGFP interface, emphasizing contacts that change upon ligand binding. The Phe76-Pro77-GluE78 cluster (in Linker 1) lies 11 to 16 Å from position 43, which defines the outer rim of the ligand site **(B)**; therefore, the cluster makes no direct contact with the ligand site. **(E1)** In the apo conformation, Glu78 acts as a candle snuffer that prevents fluorescence by the chromophore (PDB 7S7V). **(E2)** In the liganded conformation (PDB 7S7T), the Phe76-Pro77-GluE78 cluster moves Glu78 at least ~ 14 Å away from the fluorophore. Pro77 is flanked by Phe76 and Pro396 (in the top lobe of the PBP moiety). The presumably deprotonated Glu78 forms salt bridges with Lys 97 and Arg99, both facing outward on the β6 strand of the original GFP (within the original Phe165-Lys-Ile-Arg-His sequence).

168   **Figure 1-figure supplement 1.** Conformational change of apo (PDB 7S7V) to the liganded,
169   closed form (PDB 7S7T) of iNicSnFR3adt. The bottom lobe of the PBP is superimposed in the
170   two conformations. With respect to the bottom lobe, the "Venus flytrap" conformational change
171   tilts the top lobe of the PBP but does not change its structure (see Supplementary Data). The
172   conformational change also tilts the cpGFP moiety but does not change its structure.
173
174

kJ/mol
715

-10

nicotine

dianicline

cytisine

10-fluorocytisine

9-bromo-10-ethylcytisine

Figure 1-figure supplement 2

Electrostatic surface potential densities for the nicotinic agonists in this study, calculated by SPARTAN at HF/6-31G** theory level. The display ranges from -10 to 715 kJ/mol. The molecules are shown on the same distance scale.

Table 1.

Informal Name	Drug of interest	ΔF$_{max}$/F$_0$		EC$_{50}$ (µM)		S-slope		11	43	44	68	324	360	391	395
		L	P	L	P	L	P								
iNicSnFR3b	nicotine	ND	10	ND	19	ND	0.5	E	E	N	H	S	T	F	G
iDianiSnFR	dianicline	7.4 ± 0.1	4.7 ± 0.2	6.7 ± 0.3	15 ± 1	1.1	0.3	D	R	N	S	N	G	F	
iAChSnFR	ACh	ND	12	ND	1.3	ND	9.2	I	V	N	H	A	T	F	G
iCytSnFR	cytisine	5.0 ± 0.2	7.3 ± 0.4	9.4 ± 0.8	11 ± 1	0.5	0.7	I	Y	N	H	A	T	W	G
iCyt_F_SnFR	10-fluorocytisine	7.9 ± 0.1	2.3 ± 0.1	1.4 ± 0.04	1.6 ± 0.3	5.6	1.4	I	N	G	H	A	T	W	G
iCyt_BrEt_SnFR	9-bromo-10-ethylcytisine	4.0 ± 0.03	3.6 ± 0.04	5.7 ± 0.1	4.2 ± 0.2	0.7	0.9	I	Q	G	H	A	T	W	G

Nicotinic agonist iDrugSnFR naming, dose-response relations, and residues mutated.
Measurements in *E. coli* lysates (L) or with purified protein (P). ND, Not determined. Data for
iAChSnFR from (Borden, et al., 2019); Data for iNicSnFR3b from (Shivange, et al., 2019).

Figure 2. Nicotinic agonist iDrugSnFR development. Dose-response relations on intermediate constructs using *E. coli* lysate were performed with respective drug partners to identify SSM winners. **(A-D)** The progenitor biosensor is listed in black. Dashed lines indicate data that did not reach saturation at the concentrations tested; therefore, EC_{50} and $\Delta F_{max}/F0$ could not be determined. Development of **(A)** iDianiSnFR, **(B)** iCytSnFR, **(C)** iCyt_F_SnFR, and **(D)** iCyt_BrEt_SnFR.

196
Figure 3. Dose-response relations of iDrugSnFR protein versus a nicotinic agonist panel. Abbreviations: Ch (choline), ACh (acetylcholine), Cyt (cytisine), Diani (dianicline), Nic (nicotine), Var (varenicline), 10FC (10-fluorocytisine), and 9Br10EtC (9-bromo-10-ethyl-cytisine). **(A-D)** Relevant EC_{50} values for each iDrugSnFR are listed in Table 2. Dashed lines indicate dose-response relations that did not approach saturation for the concentration ranges tested; therefore, EC_{50} and $\Delta F_{max}/F0$ could not be determined. **(A)** iDianiSnFR shows preference for dianicline, with some promiscuity for other nicotinic agonists. **(B)** iCytSnFR, **(C)** iCyt_F_SnFR, and **(D)** iCyt_BrEt_SnFR bind their drug partner, but also respond to other nicotinic agonists.

206
207 **Table 2.**
208

Drug Name	iDianiSnFR			iCytSnFR			iCyt_F_SnFR			iCyt_BrEt_SnFR		
	$\Delta F_{max}/F_0$	EC_{50} (µM)	S-slope	$\Delta F_{max}/F_0$	EC_{50} (µM)	S-slope	$\Delta F_{max}/F_0$	EC_{50} (µM)	S-slope	$\Delta F_{max}/F_0$	EC_{50} (µM)	S-slope
choline	2.0 ± 0.1	84 ± 20	<0.1	5.8 ± 0.2	240 ± 30	<0.1	2.6 ± 0.1	18 ± 1	0.1	2.6 ± 0.1	12 ± 1	0.2
acetylcholine	7.4 ± 1.0	660 ± 80	<0.1	2.9 ± 0.1	35 ± 3	<0.1	4.4 ± 0.3	222 ± 50	<0.1	2.5 ± 0.2	73 ± 6	<0.1
cytisine	-	-	<0.1*	7.3 ± 0.4	11 ± 1	0.7	4.4 ± 0.1	2.6 ± 0.3	1.7	4.7 ± 0.1	3.5 ± 0.2	1.3
dianicline	4.7 ± 0.2	15 ± 1	0.3	6.5 ± 0.4	34 ± 4	0.2	2.3 ± 0.3	43 ± 6	<0.1	4-6	>100	<0.1**
nicotine	2.2 ± 0.1	440 ± 100	<0.1	6.4 ± 0.2	14 ± 2	0.5	4.7 ± 0.1	3.8 ± 0.2	1.2	4.8 ± 0.1	5.5 ± 0.2	0.9
varenicline	2.4 ± 2.0	1200 ± 500	<0.1	6.5 ± 0.1	0.06 ± 0.01	110	7.1 ± 0.2	0.09 ± 0.02	79	5.3 ± 0.1	0.06 ± 0.01	88
10-fluorocytisine	ND	ND	ND	ND	ND	ND	2.3 ± 0.1	1.6 ± 0.3	1.4	3.0 ± 0.1	4.7 ± 0.3	0.6
9-bromo-10-ethylcytisine	ND	ND	ND	ND	ND	ND	3.1 ± 0.1	31 ± 2	0.1	3.6 ± <0.1	4.2 ± 0.2	0.9

209
210 **iDrugSnFR dose-response relations versus a selected panel of nicotinic agonists.** ND, Not
211 determined. * , ** EC_{50} and $\Delta F_{max}/F_0$ could not be determined from the data (Fig. 3). Therefore,
212 the upper limit to the S-slope is estimated from the data at the foot of the dose-response relation.

Figure 4. Isothermal titration calorimetry traces, fits, and thermodynamic data. **Top row**: Exemplar heat traces of iCytSnFR, iCyt_F_SnFR, iCyt_BrEt_SnFR, and iDianiSnFR paired with their drug partners obtained by isothermal calorimetry. The heats for iCytSnFR, iCyt_F_SnFR, and iCyt_BrEt_SnFR were exothermic, while that for iDianiSnFR was endothermic. **Middle row**: The resulting fits for each iDrugSnFR:drug pair from the integrated heats comprising each series of injections. **Bottom row:** Energy calculations. All iDrugSnFRs show exergonic reactions, but the relative enthalpic and entropic contributions vary among iDrugSnFRs. Data are from 3 separate runs, Mean ± SD.

225
226 **Table 3.**

Biosensor	K_D (µM)	n	ΔH (kcal/mol)	-TΔS (kcal/mol)	ΔG (kcal/mol)
iCytSnFR	13.7 ± 1.1	0.84 ± 0.05	-2.1 ± 0.1	-4.6 ± 0.2	-6.6 ± 0.1
iCyt_F_SnFR	1.8 ± 0.5	0.83 ± 0.02	-5.5 ± 0.1	-2.4 ± 0.2	-7.9 ± 0.1
iCyt_BrEt_SnFR	5.4 ± 0.8	0.69 ± 0.09	-1.12 ± 0.03	6.1 ± 0.1	-7.2 ± 0.1
iDianiSnFR	7.6 ± 1.4	0.92 ± 0.02	3.2 ± 0.5	10.1 ± 0.4	-7.0 ± 0.2

227 **Affinity, occupancy number, and thermodynamic data calculated from isothermal titration**
228 **calorimetry**. Data are the mean ± SD, n = 3 runs.
229
230

Figure 5. Stopped-flow fluorescence kinetic data for **(A)** iDianiSnFR, **(B)** iCytSnFR, **(C)**
iCyt_F_SnFR, and **(D)** iCyt_BrEt_SnFR over 1 s and 100 s. Fluorescence was activated by mixing
with the agonists as noted. Stopped-flow data shows a departure from first-order kinetics for this
set of iDrugSnFRs. iDianiSnFR and iCyt_F_SnFR are fit to a double exponential; iCytSnFR and
iCyt_BrEt_SnFR are fit to a single exponential. **(E-H)** Plots of the observed apparent rate constant
against [agonist] for the 1 s data obtained in **(A-D).**

Specificity and thermodynamics of nicotinic iDrugSnFRs

We characterized the specificity of purified iDrugSnFRs for their drug partners versus a
panel of related nicotinic agonists (Table 2, Fig. 3). The newly developed iDrugSnFRs showed
some sensitivity to related nicotinic agonists. iDianiSnFR had the greatest fidelity for its drug
partner but also showed an increased EC_{50} (15 μM) as a purified protein versus its EC_{50} in lysate
(6.7 μM), possibly indicating decreased stability in a purified form. iCytSnFR, iCyt_F_SnFR, and
iCyt_BrEt_SnFR showed a greater level of promiscuity for the compounds comprising the nicotinic
agonist panel. Of note, iCytSnFR, iCyt_F_SnFR, and iCyt_BrEt_SnFR have an exceptionally low
(60–90 nM) EC_{50} for varenicline. The newly developed iDrugSnFRs showed negligible binding to
choline or the neurotransmitter acetylcholine, leading one to expect minimal endogenous
interference during future *in vivo* experiments.
We also performed dose-response experiments with iDianiSnFR, iCytSnFR,
iCyt_F_SnFR, and iCyt_BrEt_SnFR against a panel of nine endogenous molecules, including
neurotransmitters (Supplementary Fig. 1). iDianiSnFR showed no response to any of the nine
selected compounds above background. iCytSnFR, iCyt_F_SnFR, and iCyt_BrEt_SnFR showed
no response above background for seven of the compounds. However, they exhibited a $\Delta F/F_0$ of
0.25–0.8 to dopamine at 316 μM/1 mM and a $\Delta F/F_0$ of 0.8–1.5 to serotonin (5-HT) at 316 μM/1
mM. In terms of S-slope, the relevant metric for most cellular or *in vivo* experiments, the SnFRs
are at least 250-fold more sensitive to their eponymous partners than to other molecules we have
tested.
To examine the thermodynamics of the iDrugSnFR:drug interaction, we conducted
isothermal titration calorimetry (ITC) binding experiments (Fig. 4). The experimentally determined
K_D of each iDrugSnFR:drug pair using ITC was within a 1.5 factor of the experimentally
determined EC_{50} for fluorescence in *E. coli* lysate or purified protein (Table 3). We infer that the
EC_{50} for fluorescence is dominated by the overall binding of the ligand for all the iDrugSnFRs.

Kinetics of nicotinic agonist iDrugSnFRs: stopped-flow

In a stopped-flow apparatus, we measured the fluorescence changes of iDrugSnFRs with
millisecond resolution during both multiple 1 s trials and an independent 100 s trial. The stopped-
flow data revealed that iDrugSnFRs do not have pseudo-first-order kinetic behaviors typical of
two state binding interactions. Time courses of iDianiSnFR (both over 1 s and 100 s) were best
fitted by double exponential equations. Most of the fluorescence change occurs within the first 0.1
s of mixing (Fig. 5A), with only minor additional increase by 100 s.
Changes in fluorescence from iCytSnFR during the first 1 s of mixing fit well to a single
exponential (Fig. 5B), and have close to pseudo-first order kinetics (i.e. the observed rate of
fluorescence change is nearly linear with drug concentration). As with iDianiSnFR, most of the
fluorescence change occurs within the first second, with additional fluorescent increase continuing
over the next minute (Fig. 5B, right panel).
Like iDianiSnFR, iCyt_F_SnFR fluorescence changes are best fit by a double exponential
(Fig. 5C), but the time course of fluorescence change is significantly slower. Fluorescence
gradually increases throughout the recording period and beyond. This information was considered
in later *in vitro* and *ex vivo* experiments.
iCyt_BrEt_SnFR fits well to a single exponential (Fig. 5D) for the first 1 s of data collection,
but like the other sensors, continues to increase its fluorescence over longer periods.
We plotted the k_{obs} (s^{-1}) obtained in the 1 s stopped-flow experiments versus concentration
(Fig. 5E-5H). The aberrations from an ideal first-order kinetics vitiate generation of definitive k_{off}
and k_{on} values (Supplementary Table 2), but we can approximate a K_{max} and K_D from our fitting
procedures. Our stopped-flow experiments reinforced previous observations (Unger, et al., 2020)

that the kinetics of iDrugSnFR binding involve complexities beyond a simple first-order kinetic
model governing two binding partners.

Kinetics of nicotinic agonist iDrugSnFRs: millisecond microperfusion

We also studied iCytSnFR_PM expressed in HEK293T cells during fluorescence responses
to ACh, cytisine, or varenicline in a microperfusion apparatus that exchanged solutions near the
cell on a millisecond time scale (Methods). This system directly measures the decay of the
response when ligand is suddenly removed. The rank order of the iCytSnFR steady-state
sensitivities is varenicline > cytisine > ACh. The time constant for decay decreased with increasing
steady-state EC_{50} of the ligands, as though more tightly binding ligands dissociate more slowly
(Fig. 6A).
We measured the decay waveforms after drug pulses at concentrations ≥ the EC_{50} of the
steady-state response to maximize the $\Delta F/F_0$ signal/noise ratio (Fig. 6A-D). Because the decay
phases are measured in zero [ligand], one expects that the decay rate constant(s) (k_{off}) for an
iDrugSnFR do not depend on the pulsed ligand concentration. Decay of the ACh response
followed a single exponential time course (Fig. 6B). The values of the k_{off} for 30, 100, and 200 μM
ACh did not differ significantly (ANOVA, $p = 0.62$, degrees of freedom (df) = 2 (model), 20 (error)).
We pooled them to obtain a mean k_{off} of 1.9 ± 0.1 s^{-1} (mean ± SEM, n = 23 areas (50 cells)). The
corresponding time constant τ_{off} was 530 ± 30 ms. Hence, the temporal resolution of the
CytSnFR_PM sensor for changes in the ACh concentration was in the sub-second range.
The decay of the cytisine and varenicline response was biphasic (Fig. 6C-D): two exponential
decay terms with an additional constant component fitted the cytisine decay significantly better
than a single exponential term (F-test, $p < 0.05$). As expected, neither the faster decay rate
constants (kf_{off}) (ANOVA, $p = 0.30$, $df = 3,32$) nor the slower decay rate constants (ks_{off}) (ANOVA,
$p = 0.54$, $df = 3,31$) differed among the tested cytisine concentrations (5-15 μM). The kf_{off} and ks_{off}
for 5-15 μM cytisine were 0.61 ± 0.04 s^{-1} (n = 36 areas, 105 cells) and 0.146 ± 0.006 s^{-1} (n = 35
areas, n = 103 cells), respectively. The corresponding decay time constants ($\tau f_{off}, \tau s_{off}$) were 1.8 ±
0.1 s and 6.9 ± 0.2 s. Therefore, the temporal resolution of CytSnFR_PM sensor for cytisine was
< 10 s, adequate for the temporal resolution of the live-cell experiments presented below.
Interestingly, the decay waveform of the varenicline response was much slower than that for
cytisine or ACh (Fig. 6A, 6D). We pulsed 2 μM varenicline, >> the EC_{50} of the steady-state
response of the isolated protein (60 ± 10 nM) (Fig. 6D). The values of the kf_{off} and ks_{off} were 0.9
± 0.2 s^{-1} and 0.0065 ± 0.0002 s^{-1}, respectively (n= 4 areas (9 cells)). The slower component
dominated the decay phase, with a fractional amplitude of 85 ± 1%. Thus, the temporal resolution
of the iCytSnFR_PM sensor for varenicline was in the minute range. In the live-cell experiments
described below, it would not be possible to resolve the differences between varenicline at the
PM and in the ER. The relatively high affinity of iCytisineSnFR for varenicline, which presumably
arises in part from the increased lifetime of the varenicline-iDrugSnFR complex, has drawbacks.
The temporal resolution of iNicSnFR3a and iNicSnFR3b, which bind varenicline ~ 100-fold less
tightly, is appropriate for subcellular experiments (Shivange, et al., 2019). The previous
experiments showing ER entry of varenicline used iNicSnFR3a and iNicSnFR3b (Shivange, et
al., 2019). For additional microperfusion data and analyses see Supplementary Figures 2-4.

Figure 6. Decay of the iCytSnFR_PM responses after removal of ACh, cytisine, or varenicline. **(A)** The red, blue, and black traces are mean $\Delta F/F_0$ values for the ACh (200 µM), cytisine (15 µM), and varenicline (2 µM) responses as a function of time (n = 4-10 areas per ligand). The $\Delta F/F_0$ was normalized to the peak response for each ligand. Sampling rate was 5 frames/s. Ligand was applied for 5 s, denoted by the green horizontal bar above the traces. **(B-D)** Examples of the decay phase of the response to ACh (200 µM), cytisine (15 µM), and varenicline (2 µM) in individual areas (black traces in each panel). Red lines are fits to the sum of one or two negative exponential terms and a constant (red lines in each panel) using non-linear least-squares regression. **(B)** The decay of the ACh (200 µM) response (n = 1 area, 3 cells) was monophasic with a single time constant (τ_{Off}) of 0.61 ± 0.02 s (± SE, n = 86 frames, sampling rate of 9.8 frames/s). The red line is a fit to the sum of a negative exponential component (R^2 of 0.98). **(C)** The decay of the cytisine (15 µM) response (n = 1 area, 4 cells) was biphasic with time constants (τf_{off}, τs_{off}) of 1.9 ± 0.2 and 6.6 ± 0.5 s (n = 149 frames, sampling rate of 5 frames/s). The red line is a fit to the sum of two negative exponential components and a constant (R^2 of 0.996). It was significantly better than that of the sum of a single negative exponential term and a constant (F-test, $p < 0.05$). The relative amplitude of the slower decay component ($A_s/(A_s+A_f)$; where A_s is amplitude of the slower component of decay in units of $\Delta F/F_0$ and A_f is amplitude of the faster component) was 61%. **(D)** The decay of the varenicline (2 µM) response (n = 1 area, 3 cells) was also biphasic with a τf_{off} and τs_{off} of 9 ± 1 s and 150 ± 10 s (n = 178 frames, sampling rate of 1 frame/s), respectively. The $A_s/(A_s+A_f)$ was 83%. The red line is a fit to the sum of two negative exponential terms and a constant (R^2 of 0.994) and it was significantly better than that to the sum of a single negative exponential term and a constant (F-test, $p < 0.05$).

Characterization of nicotinic iDrugSnFRS in HeLa cells and primary mouse hippocampal culture

We examined the subcellular pharmacokinetics of the nicotinic agonists in mammalian cell lines and primary mouse hippocampal neurons. The nicotinic iDrugSnFRs were targeted to the plasma membrane (PM) (iDrugSnFR_PM) or the endoplasmic reticulum (ER) (iDrugSnFR_ER) as previously described (Bera, et al., 2019; Shivange, et al., 2019). We then performed a dose-response experiment using wide-field fluorescence imaging with each iDrugSnFR and its drug partner, sampling a range of concentrations covering a log scale surrounding the EC_{50} as determined for the purified protein (Fig. 7, Fig. 8, Supplementary Videos 1-4).

iDianiSnFR showed a robust response to dianicline at the PM and the ER in HeLa cells across a range of concentrations (3.125–100 µM) and the speed was nearly limited by solution exchanges; there was a clear return to baseline fluorescence upon washout on the order of seconds after each drug application. At 100 µM, the PM and ER have a $\Delta F/F_0$ of ~1.2, but at lower concentrations, the ER displayed 30–75% of the signal detected at the PM, which may indicate a difference in membrane crossing (Fig. 7A). Imaging in primary mouse hippocampal neurons demonstrated a similar trend (Fig. 8A).

Cytisine showed slower entry into and exit from the ER of HeLa cells. The iCytSnFR_PM construct detected cytisine at concentrations from 0.078–80 µM and demonstrated a return to baseline fluorescence upon washout on the order of seconds after each drug application, reaching a maximum $\Delta F/F_0$ of ~2 at concentrations above 5 µM (Fig. 7B). In contrast to the _PM construct, the iCytSnFR_ER construct only detected cytisine with a $\Delta F/F_0$ above the buffer control in the range of concentrations from 1.25–80 µM with a $\Delta F/F_0$ which was 25–50% of the maximum $\Delta F/F_0$ detected at the PM. Additionally, in the range of detectable concentrations of cytisine, the washout of cytisine was much slower than solution changes (Fig. 7B). The incomplete washout persists even after several minutes and corresponds with previous suggestions that cytisine has low membrane permeability, as evidenced by its low brain penetration (Rollema, et al., 2010).

In primary mouse hippocampal neurons, iCytSnFR detection of cytisine exhibited the same kinetic trends seen in HeLa cell experiments (Fig. 8B). During cytisine application (60 s) from 0.078-80 µM, the iCytSnFR_PM fluorescence nearly reached a plateau, and during the washout (90-180 s), the fluorescence decayed back to baseline, though the decay slowed after removal of higher [cytisine]. The _PM construct reached a maximum $\Delta F/F_0$ of ~1.25 at 80 µM, which was approximately 60% of the signal observed in HeLa cell experiments (Fig. 8B). The iCytSnFR_ER detection of cytisine in the ER reflected the trends seen in HeLa cells, with incomplete cytisine wash-in phases and prolonged cytisine washout phases. One observable difference was that the maximum $\Delta F/F_0$ (~1.25) of iCytSnFR_ER reached a similar maximum to that of iCytSnFR_PM in neurons, which was not observed in HeLa cell experiments (Fig 7B).

In preliminary HeLa cell experiments with varenicline applied to iCytSnFR, we found much slower kinetics that differed little between the _ER and _PM constructs (data not shown). These findings, which vitiated the use of the iCytSnFR:varencline pair in the cellular experiments, are consistent with the markedly slow kinetics of varenicline-iCytSnFR interactions in the microperfusion experiments (see above).

iCyt_F_SnFR targeted to the PM and ER showed characteristics similar to iCytSnFR in HeLa cells. The _PM construct detected 10-fluorocytisine across a range of concentrations with a return to baseline fluorescence between applications, while the _ER construct detected 10-fluorocytisine with $\Delta F/F_0$ values that were only 25–33% of those detected at the PM (Fig. 7C). Similar to the iCytSnFR_ER detection of cytisine, the iCyt_F_SnFR_ER detection of 10-fluorocytisine was much slower than solution changes and did not return to baseline between applications, though the washout occurs on the order of minutes, rather than tens of minutes as with iCytSnFR_ER (Fig. 7C). The difference in PM and ER detection of 10-fluorocytisine again shows decreased membrane permeability into HeLa cells compared to other drugs we have examined with other

iDrugSnFRs. Overall, the detection of 10-fluorocytisine with iCyt_F_SnFR in primary hippocampal
culture resembled our data with iCyt_F_SnFR in HeLa cells. Nevertheless, there were distinct
differences (Fig. 8C), such as a decreased maximum $\Delta F/F_0$ in the iCyt_F_SnFR_PM construct
and a similar maximum $\Delta F/F_0$ of ~1 for both the _ER and _PM constructs. Additionally, the decay
of the iCyt_F_SnFR responses lasted tens of minutes, resembling the iCytSnFR_ER data in
primary hippocampal culture.
9-bromo-10-ethylcytisine   showed   a   kinetic   profile   resembling   dianicline.
iCyt_BrEt_SnFR_PM responses to 9-bromo-10-ethylcytisine (0.1–31.6 µM) were nearly limited
by solution exchanges with a return to baseline fluorescence on the order of seconds, and a
maximum $\Delta F/F_0$ of ~1.5 at 31.6 µM. iCyt_BrEt_SnFR_ER also detected 9-bromo-10-ethylcytisine
over this range of concentrations and returned to baseline fluorescence between applications
(Fig. 7D). $\Delta F/F_0$ values for iCyt_BrEt_SnFR_ER were 50–75% of the $\Delta F/F_0$ values detected by
iCyt_BrEt_SnFR_PM, which indicated that 9-bromo-10-ethylcytisine crossed into and out of cells
readily (Fig. 7D). Imaging in primary mouse hippocampal neurons revealed the same trend (Fig.
8D).
To more fully examine the membrane-crossing properties of the nicotinic agonists, we
recorded the fluorescence waveforms for several drugs at concentrations between 0.1-3.16 µM
with much longer application times and washout times than in the above experiments
(Supplementary Fig. 5). With these conditions, the fluorescence signals suggested complete
washout of each nicotinic agonist from the ER of HeLa cells. However, it is noteworthy that even
when applied at concentrations as low as 0.1 µM and 0.316 µM, cytisine and 10-fluorocytisine
require washout times of several min from the ER. In contrast, the iDrugSnFR localized to the PM
shows a rapid return to baseline after drug application.
Because the data of Fig. 7B and Fig. 8B indicated that iCytSnFR_PM functions as expected
from stopped-flow and millisecond perfusion, we applied additional experiments to ensure that
our observations of drug entry and exit from the ER were not the result of idiosyncratic biosensor
function or folding in the ER. iCytSnFR and iCyt_F_SnFR both bind nicotine in the same
concentration range as cytisine (though with lower $\Delta F/F_0$). After transfection of _PM and _ER
constructs for each sensor into HeLa cells, we performed time-resolved imaging for pulses of 0.1-
31.6 µM nicotine (Fig. 7-figure supplement 2). These nicotine waveforms resembled those already
published with iNicSnFR3a and iNicSnFR3b (Shivange, et al., 2019), confirming that
iCytSnFR_ER functions as expected with a more permeant nicotinic drug. Thus, the slower
kinetics for iCytSnFR_ER with-cytisine and iCyt_F_SnFR_ER with 10-fluorocytisine arise
because these drugs cross membranes more slowly.
To examine localization of the _PM and _ER constructs at higher optical resolution, we
imaged HeLa cells and primary mouse hippocampal culture using a spinning disk laser scanning
inverted confocal microscope. As previously observed, ER-targeted iDrugSnFR was retained in
the ER (Fig. 7-figure supplement 1, Fig. 8-figure supplement 1) (Shivange, et al., 2019).
IDrugSnFR targeted to the PM showed correct localization, with some iDrugSnFR observed in
the cell interior (most likely as part of the cellular membrane trafficking system (Fig. 7-figure
supplement 1; Fig. 8-figure supplement 1).
Several complexities in the HeLa cell and neuron experiments imposed uncertainties on our
kinetic analyses. These complexities include the limitations of solution changes, diffusion within
cytoplasm, unknown mixing at the surface facing the coverslip, and corrections for baseline drift
due to bleaching. We restrict the quantitative comparisons to the estimate that cytisine and 10-
fluorocytisine cross the membrane > 30 fold more slowly than the other drugs tested.
448

Figure 7. Nicotinic agonist iDrugSnFR dose-response relations in HeLa cells. **(A-D)** Each iDrugSnFR detects its drug partner at the PM and ER of HeLa cells at the concentrations sampled. **(A)** iDianiSnFR detects dianicline with a return to baseline fluorescence between drug applications. **(B)** iCytSnFR detection at the PM returns to baseline fluorescence between applications, while detection at the ER shows incomplete wash-in and washout. **(C)** iCyt_F_SnFR fluorescence response to the presence of 10-fluorocytisine in the ER also shows an incomplete washout between applications. **(D)** iCyt_BrEt_SnFR detects 9-bromo-10-ethylcytisine with wash-in and washout fluorescence similar to the pattern seen in iDianiSnFR.

Figure 7-figure supplement 1. Spinning disk laser scanning confocal inverted microscope
images of nicotinic agonist iDrugSnFRs in HeLa cells. ER-targeted constructs of iDianiSnFR,
iCytSnFR, iCyt_F_SnFR, iCyt_BrEt_SnFR are shown before **(A1-D1)** and during **(A2-D2)**
exposure to each drug partner. ER-targeted iDrugSnFRs show the reticulated ER and dark ovals
corresponding to the nucleus. PM-targeted constructs of the same iDrugSnFRs are shown before
(E1-H1) and after **(E2-H2)** drug introduction. Localization to the PM is robust, with some minimal
puncta that may represent inclusion bodies or internal transport.
469

471
472 **Figure 7-figure supplement 2.** Dose-response relations for iCytSnFR and iCyt_F_SnFR against
473 nicotine in HeLa cells. **(A)** iCytSnFR and **(B)** iCyt_F_SnFR detect nicotine at both the PM and
474 ER. Nicotine enters and exits the ER rapidly over seconds, a direct contrast to the behavior of
475 cytisine and 10-fluorocytisine as detected by their iDrugSnFR partners.
476

Figure 8. Nicotinic agonist iDrugSnFR dose-response experiments in mouse primary hippocampal neurons transduced with AAV9-hSyn iDrugSnFR. Cultured primary mouse hippocampal neurons were transduced with ER- or PM- targeted constructs. **(A-D)** Each iDrugSnFR detects its drug partner at the PM and ER over the concentrations sampled. **(A)** iDianiSnFR detects dianicline with a return to baseline fluorescence between drug applications. **(B)** iCytSnFR detection at the PM returns to baseline fluorescence between applications, while detection at the ER shows an incomplete washout. **(C)** iCyt_F_SnFR fluorescence response to the presence of 10-fluorocytisine in the ER also shows an incomplete washout between applications. **(D)** iCyt_BrEt_SnFR_ER detects 9-bromo-10-ethylcytisine with a wash-in and decay intermediate between iDianiSnFR and the other two cytisine derivatives.

492
Figure 8-figure supplement 1. Spinning disk laser scanning confocal inverted microscope
images of nicotinic agonist iDrugSnFRs in primary mouse hippocampal neurons. ER-targeted
constructs of iDianiSnFR, iCytSnFR, iCyt_F_SnFR, and iCyt_BrEt_SnFR are shown before **(A1-**
D1) and during **(A2-D2)** exposure to each drug partner. ER-targeted iDrugSnFRs show the
reticulated the ER and dark ovals corresponding to the nucleus. PM-targeted constructs of the
same iDrugSnFRs are shown before **(E1-H1)** and after **(E2-H2)** drug introduction. Localization in
the PM is robust, with some minimal puncta that may represent inclusion bodies or internal
transport.
501

Discussion

Membrane permeation of molecules with low logD$_{pH7.4}$

The experiments show, to our knowledge, the first time-resolved measurements of
membrane permeation for drugs in the logD$_{pH7.4}$ range less than -1. Most orally available drugs
have logD$_{pH7.4}$ values between 2 and 4 (Smith, Allerton, Kalgutkar, van de Waterbeemd, & Walker,
2012). Cytisine, varenicline, dianicline, and the cytisine analogs studied here have calculated
membrane partition coefficients some 3 to 6 orders of magnitude lower. These values and their
order vary according to the algorithm, partially because of uncertainties in predicting pK$_a$ (Pienko,
et al., 2016); here we provide values calculated by Chemicalize (see Methods): 10-fluorocytisine,
-2.70; cytisine, -2.64; 9-bromo-10-ethylcytisine, -1.13; varenicline, -1.27, dianicline, -1.29; It is
remarkable that drugs with such low calculated partition coefficients do cross membranes on a
time scale of seconds (9-bromo-10-ethylcytisine, varenicline, dianicline) to minutes (10-
fluorocytisine, cytisine). According to some (but not all) algorithms, the calculated logD$_{pH7.4}$ values
fall in the same two classes as the measured kinetics of membrane permeability: 10-fluorocytisine
and cytisine are the slowest, and only these two agonists have logD$_{pH7.4}$ values < -2. These
observations support previous work suggesting that differences among chemical properties of
nicotinic partial agonists correlate with drug permeation into the cerebrospinal fluid (CSF) after
peripheral administration in mice (Rollema, et al., 2010).

The iDrugSnFR paradigm

The iDrugSnFRs are sensitive enough to allow experiments near the experimentally
determined (or otherwise projected) concentration in the human blood and CSF (Astroug,
Simeonova, Kassabova, Danchev, & Svinarov, 2010; Jeong, Sheridan, Newcombe, & Tingle,
2018; Rollema, et al., 2010). The iDrugSnFRs have the advantage that they measure free
aqueous ligand concentration ("activity"), as sensed by nAChRs. Targeting sequences provide
for visualization within the lumen of organelles—here, the ER.
The experiments do not use radiolabeled drugs, *in vivo* microdialysis or other experiments
on live animals, or mass spectrometry-liquid chromatography instruments. Once protein design
has given an entry into a class of iDrugSnFRs, straightforward optimization at the binding site
produces the desired, selective iDrugSnFRs for individual molecules. The experiments use
standard, modest-power fluorescence microscopes. Cultured cell lines yield data comparable to
cultured neurons.

Structure-function relations for nicotinic and other iDrugSnFRs

This study shows that the amine group of nicotinic ligands makes equidistant cation-π
interactions with two tyrosine residues (Tyr65, Tyr357), and this is confirmed by higher-resolution
(1.5 to 1.7 Å) structures of varenicline, acetylcholine, and choline crystallized with isolated PBP
moieties (PDB 7S7X, S6V1R, 7S7Z, respectively; see also 3R6U, 6EYQ, and 3PPQ). Cation-π
interactions also occur for cholinergic and/or nicotinic ligands in nAChRs (Morales-Perez,
Noviello, & Hibbs, 2016; Post, Tender, Lester, & Dougherty, 2017), the acetylcholine-binding
protein (Celie, et al., 2004), PBPs (Schiefner, et al., 2004), and muscarinic receptors (Haga, et
al., 2012). We also observe that the protonated amine of varenicline makes a hydrogen bond to
a backbone carbonyl group, another similar theme in acetylcholine binding protein (Celie, et al.,
2004) and nAChRs (Xiu, Puskar, Shanata, Lester, & Dougherty, 2009).

This study presents a general step forward in understanding the structure-function relations
of iDrugSnFRs. The chromophore in the cpGFP moiety of most present iDrugSnFRs (this paper,
iGluSnFR, iSeroSnFR) contains a tyrosine in an extended π system (Ormo, et al., 1996; Tsien,
1998). The photophysics of the chromophore depends strongly on the surrounding water
molecules and side chains (Brejc, et al., 1997; Tsien, 1998). We found that Glu78 in Linker 1
changes its orientation: in the liganded state, it interacts with two positively charged residues
(Lys97 and Arg99) on the surface of the cpGFP; and in the apo state, Glu78 has moved ~ 14 Å
to form a hydrogen bonding interaction with the tyrosine moiety of the chromophore. Presumably
the liganded state of iNicSnFR3adt allows for a water molecule to hydrogen bond with the hydroxy
group of the chromophore, promoting its fluorescence; but this water molecule is replaced by
protonated Glu78 in the unliganded state, which leads to nonfluorescent state of cpGFP, as
suggested by (Nasu, et al., 2021).
While we cannot resolve the protonation-deprotonation event, the available functional data
show good support for its occurrence, as follows. (a) The apo form of the iDrugSnFR increases
its F_0 by ten-fold per pH unit (Shivange, et al., 2019), as though when deprotonated, Glu78 leaves
the "candle snuffer" position and moves to make the salt bridges with Lys97 and Arg99. (b) The
EC50 for the ligand decreases by ten-fold per pH unit (Shivange, et al., 2019), as though the
conformation of the linker that forms the salt-bridge form is also the closed, liganded, fluorescent
form of the PBP. Other observations favor the crucial role of the Glu78-chromophore interaction.
(a) Only glutamate functions in position 78 of iSeroSnFR (Unger, et al., 2020). (b) The mTurquoise
variant in iGluSnFR, which has a tryptophan chromophore, requires entirely different linkers
(Marvin, et al., 2018).

Challenges at the intersection of pharmaceutical science and nicotine addiction science

Our measurements show that nicotinic ligands with $\log D_{pH7.4} < \sim$ -2 cross membranes much
more slowly than do ligands with $\log D_{pH7.4} > \sim$ -2. These measurements have two, possibly
opposing, implications for future smoking cessation drugs. On the one hand, $\alpha 4\beta 2$ agonists that
enter the ER, like nicotine and varenicline, upregulate nAChRs (Turner, Castellano, & Blendy,
2011), which may be necessary and sufficient for addiction (Henderson & Lester, 2015); and
maintenance of upregulation by varenicline may help to explain its suboptimal quit rate. On the
other hand, ligands that do not enter the ER are also unlikely to enter the brain and therefore
unlikely to be useful for smoking cessation.
Smoking cessation drugs must also contend with other ER-based processes. (1) Most drug
metabolism takes place in the ER; (2) Upregulation occurs at sustained agonist concentration in
the ER some hundredfold lower than the extracellular concentrations that transiently activate
$\alpha 4\beta 2$ nAChRs (Kuryatov, Luo, Cooper, & Lindstrom, 2005).
Given these challenges, further progress may be possible now that we have two types of
real-time, living cellular preparations. (1) For decades, cellular preparations have been available
to measure nAChR pharmacodynamics and upregulation. (2) Now, the iDrugSnFRs present a
paradigm to measure cellular and subcellular pharmacokinetics.
586

Methods

Crystallography

The gene encoding the full-length biosensor iNicSnFR3a was previously cloned into a bacterial expression vector (Shivange, et al., 2019). To improve crystallization, we deleted the N-terminal tags His$_6$ and HA tags and the N-terminal Myc tag, forming the constructs with the suffix "dt". These deletions were carried out with the Q5 Site-Directed Mutagenesis Kit (New England Biolabs, Ipswich, MA). All proteins were overexpressed in *E. coli* BL21-gold (DE3) cells (Agilent Technologies, Santa Clara, CA) using ZYM-5052 autoinduction media (Studier, 2005). Cells were collected by centrifugation and stored at -80 ˚C until use.

For purification, frozen cell pellets were resuspended in lysis buffer containing 100 mM NaCl, 20 mM Tris, pH 7.5, 20 mM imidazole, pH 7.5, 5 mM β-mercaptoethanol (BME), lysozyme, DNase, and protease inhibitor tablet. The resuspended cells were lysed by freezing and thawing using liquid nitrogen and a room temperature water bath for 3 cycles. Intact cells and cell debris were removed by centrifugation at ~20,000x g for 40 min at 4 ˚C. The supernatant was collected and loaded onto a prewashed Ni NTA column with wash buffer at 4 ˚C. Ni NTA wash buffer contained 100 mM NaCl, 20 mM Tris, pH 7.5, 30 mM imidazole, pH 7.5, and 5 mM BME. Elution was achieved using the same buffer with 300 mM imidazole, pH 7.5. The eluted sample was further purified by size exclusion chromatography using HiLoad 16/60 Superdex 200 in the same buffer without imidazole and BME. Peak fractions were collected and concentrated to ~50 mg/ml with Amicon Ultra 15 filter unit (Millipore, Burlington, MA) with 10kDa cutoff.

For all constructs, initial crystallization screening was carried out with 40 mg/ml protein in the presence and absence of 10 mM nicotine or varenicline. iNicSnFR3adt crystallized separately with 10 mM nicotine and varenicline in PACT premier (Molecular Dimensions, Sheffield, England), condition #96 with 0.2 M sodium malonate dibasic monohydrate, 0.1 M Bis-Tris Propane, pH 8.5, and 20% polyethylene glycol (PEG) 3,350 at 20 °C. Crystals of iNicSnFR3adt grew within two weeks of crystallization in a hexagonal rod shape with dimensions of ~ 80 μm x 80 μm x 300 μm. Crystals were harvested and cryo-protected in 25% ethylene glycol, 0.2 M sodium malonate dibasic monohydrate, 0.1 M BisTrisPropane pH 8.5, and 20% PEG 3350. Phase information was obtained through soaking with KI before cryo-protection. The unliganded iNicSnFR3adt crystallized in Morpheus (Molecular Dimensions), condition #92 with 2.5% PEG 1,000, 12.5% PEG 3,350, 12.5% 2-methyl-2,4-pentanediol, 0.02 M of each amino acid, and 0.1 M MOPS/HEPES-Na, pH 7.5 at 23 °C with no further optimization.

X-ray datasets were collected at Stanford Synchrotron Radiation Laboratory beamline 12-2 and Advanced Light Source beamline 5.0.2 using Pilatus 6M detectors. All datasets were processed and integrated with XDS (Kabsch, 2010) and scaled with Aimless (Winn, et al., 2011). For iNicSnFR3adt, molecular replacement was carried out using domains of the unliganded structure (PDB ID: 6EFR) with Phaser in Phenix (Adams, et al., 2010). The experimental phase information of KI-soaked crystals of iNicSnFR3adt was obtained with MR-SAD using AutoSol in Phenix (Adams, et al., 2010). Molecular replacements of the remaining structures were carried out with the refined model of iNicSnFR3adt. Iterative refinement and model building cycles for all structures were carried out separately with phenix.refine in Phenix (Adams, et al., 2010) and Coot (Emsley, Lohkamp, Scott, & Cowtan, 2010).

Directed evolution of iDrugSnFR proteins using bacterial-expressed protein assays

Starting with iAChSnFR and intermediate biosensor constructs of that sensor, we constructed and optimized iDrugSnFRs for each drug partner during iterative rounds of SSM as previously described (Bera, et al., 2019; Shivange, et al., 2019). We utilized the 22-codon procedure

including a mixture of three primers, creating 22 unique codons encoding the 20 canonical amino
acids (Kille, et al., 2013). The 22-codon procedure yields an estimated > 96% residue coverage
for a collection of 96 randomly chosen clones.
A Tecan Spark M10 96-well fluorescence plate reader (Tecan, Männedorf, Switzerland) was
used to measure baseline and drug-induced fluorescence (F_0 and ΔF, respectively). Bacterial
lysates were tested with excitation at 485 nm and emission at 535 nm. Lysates were also
measured against choline to evaluate potential endogenous intracellular binding. Promising
clones were amplified and sequenced. The optimally responding construct in each round of SSM
was used as a template for the next round of SSM.
S-slope allows for comparison between iDrugSnFRs with differing $\Delta F_{max}/F_0$ values (Bera, et
al., 2019) at the beginning of the dose-response relation, which is usually the pharmacologically
relevant range. With lysates or purified protein, which allow complete dose-response relations,
the Hill coefficient is near 1.0. We therefore calculated
$S_{slope} = \frac{\Delta F_{max}}{F_0}/EC_{50},$
in units of μM^{-1}.
648

Measurements on purified iDrugSnFRs

Biosensors selected for further study were purified using a His$_6$ sequence using an ÄKTA
Start FPLC (GE Healthcare, Chicago, IL) as previously described (Shivange, et al., 2019).
Performance of protein quantification and dose-response relations for drug-sensor partners was
also as previously described (Shivange, et al., 2019). Where appropriate, we corrected for
depletion of the ligand by binding with the equation,

$$\Delta F/\Delta F_{max} = \frac{K_D + [S] + [L] - \sqrt{([S]^2 + [L]^2 + K_D{}^2) - 2[S][L] + 2[S]K_D + 2[L]K_D}}{2[S]},$$

where K_D is the ligand-sensor equilibrium dissociation constant (we assume that $K_D = EC_{50}$), $[S]$
is the iDrugSnFR protein concentration (typically 100 nM), and $[L]$ is the nominal ligand
concentration.

Isothermal titration calorimetry

Experiments were performed on an Affinity ITC (TA instruments, New Castle, DE) at
25 °C. The iDrugSnFR protein was buffer-exchanged into 3x PBS, pH 7.0. The nicotinic agonists
were dissolved in the same buffer. 800 µM cytisine (Sigma Aldrich, Munich, Germany) was titrated
into 80 µM iCytSnFR, 160 µM 10-fluorocytisine was titrated into 16 µM iCyt_F_SnFR. 470 µM 9-
bromo-10-ethylcytisine was titrated into 47 µM iCyt_BrEt_SnFR. 1.5 mM dianicline (Tocris, Bio-
Techne, Minneapolis, MN) was titrated into 150 µM iDianiSnFR. Analysis, including correction for
changes in enthalpy generated from the dilution of the ligands, was performed using a single-site
binding model in the manufacturer's Nanoanalyze software.

Stopped-flow kinetic analysis

Kinetics were determined by mixing equal volumes of 0.2 µM iDrugSnFR protein (in 3x
PBS, pH 7.0) with varying concentrations of cognate ligand in an Applied Photophysics (Surrey,
United Kingdom) SX20 stopped-flow fluorimeter with 490 nm LED excitation and 510 nm long-
pass filter at room temperature (22 °C). "Mixing shots" were repeated five times and averaged
(except for 100 s experiments, which were collected only once). Standard deviations are not

included on the plots, but are nearly the same size as the data markers. The first 3 ms of data
were ignored because of mixing artifacts and account for the dead time of the instrument.

Data were plotted and time courses were fitted, when possible, to a single exponential,
and k_{obs} was plotted as a function of [ligand]. The linear portion of that graph was fit, with the slope
reporting k_1 and the y-intercept reporting k_{-1}. When the time course did not fit well to a single rising
exponential, it was fitted to the sum of two increasing exponentials, and the first rise (k_{obs1}) was
treated as above to determine k_1 and k_{-1}.

Expression in mammalian cells

We constructed two variants of each iDrugSnFR for expression in mammalian cells. The
plasma membrane (suffix _PM) and endoplasmic reticulum (suffix _ER) variants were constructed
by circular polymerase extension cloning (Quan & Tian, 2009). To create the _PM constructs, we
cloned the bacterial constructs into pCMV(MinDis), a variant of pDisplay (ThermoFisher Scientific,
Waltham, MA) lacking the hemagglutinin tag (Marvin, et al., 2013). To generate the _ER
constructs, we replaced the 14 C-terminal amino acids (QVDEQKLISEEDLN, including the Myc
tag) with an ER retention motif, QTAEKDEL (Shivange, et al., 2019).
We transfected the iDrugSnFR cDNA constructs into HeLa and HEK293T cells. Cell lines
were purchased from ATCC (Manassas, VA) and cultured according to ATCC protocols. For
chemical transfection, we utilized either Lipofectamine 2000 or Lipofectamine 3000
(ThermoFisher Scientific), following the manufacturer's recommended protocol. Cells were
incubated in the transfection medium for 24 h and imaged 24 – 48 h after transfection.

Millisecond timescale microperfusion

HEK293T cells were imaged using a Nikon (Tokyo, Japan) DIAPHOT 300 with a Zeiss 63X
objective (1.5 NA). Because the ligand concentration after micro-iontophoretic drug application
(Shivange, et al., 2019) is unknown, we applied drugs with a laminar-flow microperfusion (Model
SS-77B Fast-Step perfusion system (Warner Instruments, Holliston, MA). In an array of three
square glass capillaries (600 μ i.d.), the center capillary contained vehicle (Hanks buffered salt
solution, HBSS) plus drug, while the two outer capillaries contained vehicle only. Vehicle also
flowed from a separate input connected to the bath perfusion system. Solution exchange,
measured by loading the center capillary with dye, had a time constant of 90 ± 20 ms (n = six
trials).
We used Fiji ImageJ and Origin Pro 2018 (OriginLab, Northampton, MA) to fit the rise and
decay of the iCytSnFR_PM drug response to the sum of one or two exponential components. An
F-test determined whether two exponential components fit the data significantly better than one
($p < 0.05$). Statistical comparisons between groups were carried out using ANOVA.

AAV production and transduction in primary mouse hippocampal neuronal culture

The adeno-associated virus plasmid vector AAV9-hSyn was described previously (Challis, et
al., 2019). Virus was purified using the AAVpro Purification Kit (TakaraBio USA). Mouse embryo
dissection and culture were previously described (Shivange, et al., 2019). About 4 days after
dissection, we transduced the _ER construct at an MOI of .5 to 5 x 10^4; and separately, the _PM
construct was transduced at an MOI of 0.5 to 1 x 10^5. Neurons were imaged ~2-3 weeks post-
transduction.

Time-resolved fluorescence measurements in live mammalian cells and primary mouse
hippocampal neuronal culture

Time-resolved dose-response imaging was performed on a modified Olympus IX-81
microscope (Olympus microscopes, Tokyo, Japan), in widefield epifluorescence mode using a
40X lens. Images were acquired at 2 – 4 frames/s with a back-illuminated EMCCD camera (iXon
DU-897, Andor Technology USA, South Windsor, CT), controlled by Andor IQ3 software.
Fluorescence measurements at λ_{ex} = 470 nm and the epifluorescence cube were as previously
described (Shivange, et al., 2019; Srinivasan, et al., 2011).
Solutions were delivered from elevated reservoirs by gravity flow, via solenoid valves
(Automate Scientific, Berkeley, CA), then through tubing fed into a manifold, at a rate of 1-2
ml/min. The vehicle was HBSS. Other details have been described (Shivange, et al., 2019;
Srinivasan, et al., 2011). Data analysis procedures included subtraction of "blank" (extracellular)
areas and corrections for baseline drifts using Origin Pro 2018.

728 *Spinning disk confocal fluorescence images*

HeLa cells and mouse primary hippocampal culture were transfected or transduced as
described above. Live-cell images were collected using a Nikon Ti-E spinning disk laser scanning
confocal inverted microscope equipped with 100X objective, 1.49 NA (oil), 120 µm WD. The laser
wavelength was 488 nm at 15% power. Dishes were imaged in a custom incubator (Okolab,
Ottaviano, Italy) at 37° C and 5% CO_2. Initial images were taken in HBSS. To add drug, we
doubled the bath volume by adding HBSS containing drug, using a hand-held pipette. The final
drug concentrations: dianicline, 15 µM; cytisine, 10 µM; 10-fluorocytisine, 10 µM; 9-bromo-10-
ethylcytisine, 7.5 µM.

737 *LogD calculations*

738 We used Chemicalize (https://chemaxon.com/products/chemicalize). The software uses
739 algorithms to calculate LogP and pK_a. The software then calculates
740 $LogD_{7.4} = logP - log[1 + 10^{7.4 - pKa}]$.

741 *Plasmid availability*

742 We will deposit plasmids with the following cDNAs at Addgene:
743 iDianiSnFR,
744 iCytSnFR,
745 iCyt_F_SnFR,
746 iCyt_BrEt_SnFR
747
We will deposit the following plasmids at Addgene:
pCMV(MinDis)-iDianiSnFR_PM,
pCMV(MinDis)-iCytSnFR_PM
pCMV(MinDis)-iCyt_F_SnFR_PM
pCMV(MinDis)-iCyt_BrEt_SnFR_PM
pCMV(MinDis)-iDianiSnFR_ER,
pCMV(MinDis)-iCytSnFR_ER,
pCMV(MinDis)-iCyt_F_SnFR_ER,
pCMV(MinDis)-iCyt_BrEt_SnFR_ER
pAAV9-hSyn-iDianiSnFR_PM,
pAAV9-hSyn-iCytSnFR_PM,
pAAV9-hSyn-iCyt_F_SnFR_PM,
pAAV9-hSyn iCyt_BrEt_SnFR_PM,
pAAV9-hSyn-iDianiSnFR_ER,

pAAV9-hSyn-iCytSnFR_ER,
pAAV9-hSyn-iCyt_F_SnFR_ER,
pAAV9-hSyn iCyt_BrEt_SnFR_ER,

Acknowledgements

We thank Stefan Petrovic for his stewardship of the isothermal titration calorimeter in the Caltech Center for Molecular Medicine, Jens Kaiser for help with structural studies at the Caltech Molecular Observatory, the Gradinaru lab and Caltech CLOVER Center for help with viral vectors, and Andres Collazo and Giada Spigolon at the Caltech Biological Imaging Facility. We thank Zoe Beatty, Kallol Bera, Eve Fine, Shan Huang, Elaine Lin, Stephen Mayo, Lin Tian, and Elizabeth Unger for advice and guidance. We thank Achieve Life Sciences for a gift of cytisine.

Competing interests

The authors declare no financial or non-financial competing interests.

Funding

California Tobacco-Related Disease Research Program (TRDRP) (27FT-0022), Aaron L. Nichols.

California Tobacco-Related Disease Research Program (TRDRP) (27IP-0057), Henry A. Lester.

California Tobacco-Related Disease Research Program (TRDRP) (T29IR0455), Dennis A. Dougherty.

NIH (GM-123582, DA043829), Henry A. Lester.

NIH (DA049140, GM7616), Anand K. Muthusamy.

Howard Hughes Medical Institute (Loren L. Looger, Jonathan S. Marvin, Douglas C. Rees).

UK Engineering and Physical Sciences Research Council (No. EP/N024117/1), Timothy Gallagher.

Leiden University International Studies Fund (LISF L18020-1-45), Laura Luebbert.

References

Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N., et al. (2010). PHENIX: A comprehensive Python-based system for macromolecular structure solution. *Acta Crystallographica D Biological Crystallography, 66*(Pt 2), 213-221. doi: 10.1107/S0907444909052925

Astroug, H., Simeonova, R., Kassabova, L. V., Danchev, N., & Svinarov, D. (2010). Pharmacokinetics of cytisine after single intravenous and oral administration in rabbits. *Interdiscip Toxicol, 3*(1), 15-20. doi: 10.2478/v10102-010-0003-5

Barnett, L. M., Hughes, T. E., & Drobizhev, M. (2017). Deciphering the molecular mechanism responsible for GCaMP6m's Ca^{2+}-dependent change in fluorescence. *PLoS One, 12*(2), e0170934. doi: 10.1371/journal.pone.0170934

Bera, K., Kamajaya, A., Shivange, A. V., Muthusamy, A. K., Nichols, A. L., Borden, P. M., et al. (2019). Biosensors Show the Pharmacokinetics of S-Ketamine in the Endoplasmic Reticulum. *Front Cell Neurosci, 13*, 499. doi: 10.3389/fncel.2019.00499

Blom, A. E. M., Campello, H. R., Lester, H. A., Gallagher, T., & Dougherty, D. A. (2019). Probing Binding Interactions of Cytisine Derivatives to the α4β2 Nicotinic Acetylcholine Receptor. *J Am Chem Soc.* doi: 10.1021/jacs.9b06580

Borden, P., Shivange, A. V., Marvin, J. S., Cichon, J., Dan, C., Podgorski, K., et al. (2019). A genetically encoded fluorescent sensor for in vivo acetylcholine detection. *bioRxiv*.

Brejc, K., Sixma, T. K., Kitts, P. A., Kain, S. R., Tsien, R. Y., Ormo, M., et al. (1997). Structural basis for dual excitation and photoisomerization of the *Aequorea victoria* green fluorescent protein. *Proceedings of the National Academy of Sciences of the United States of America, 94*(6), 2306-2311.

Celie, P., van Rossum-Fikkert, S., Van Dyke, W., Brejc, K., Smit, A., & Sixma, T. (2004). Nicotine and Carbamylcholine Binding to Nicotinic Acetylcholine Receptors as Studied in AChBP Crystal Structures. *Neuron, 41*, 907-914.

Challis, R. C., Kumar, S. R., Chan, K. Y., Challis, C., Beadle, K., Jang, M. J., et al. (2019). Publisher Correction: Systemic AAV vectors for widespread and targeted gene delivery in rodents. *Nat Protoc, 14*(8), 2597. doi: 10.1038/s41596-019-0155-5

Chellappan, S. K., Xiao, Y., Tueckmantel, W., Kellar, K. J., & Kozikowski, A. P. (2006). Synthesis and pharmacological evaluation of novel 9- and 10-substituted cytisine derivatives. Nicotinic ligands of enhanced subtype selectivity. *J Med Chem, 49*(9), 2673-2676. doi: 10.1021/jm051196m

Coe, J. W., Brooks, P. R., Vetelino, M. G., Wirtz, M. C., Arnold, E. P., Huang, J., et al. (2005). Varenicline: an alpha4beta2 nicotinic receptor partial agonist for smoking cessation. *J Med Chem, 48*(10), 3474-3477. doi: 10.1021/jm050069n

Cohen, C., Bergis, O. E., Galli, F., Lochead, A. W., Jegham, S., Biton, B., et al. (2003). SSR591813, a novel selective and partial alpha4beta2 nicotinic receptor agonist with potential as an aid to smoking cessation. *J Pharmacol Exp Ther, 306*(1), 407-420. doi: 10.1124/jpet.103.049262

Elmore, D. E., & Dougherty, D. A. (2000). A computational study of nicotine conformations in the gas phase and in water. *Journal of Organic Chemistry, 65*(3), 742-747. doi: 10.1021/jo991383q

Emsley, P., Lohkamp, B., Scott, W. G., & Cowtan, K. (2010). Features and development of Coot. *Acta Crystallographica D Biological Crystallography, 66*(Pt 4), 486-501. doi: 10.1107/S0907444910007493

Fagerstrom, K., & Balfour, D. J. (2006). Neuropharmacology and potential efficacy of new treatments for tobacco dependence. *Expert Opin Investig Drugs, 15*(2), 107-116. doi: 10.1517/13543784.15.2.107

Fan, C. (2020). *Structure, Function, and Application of Bacterial ABC Transporters.* Dissertation
(Ph.D.),   California   Institute   of   Technology.   Retrieved   from
https://resolver.caltech.edu/CaltechTHESIS:06012020-160106726
Govind, A. P., Vallejo, Y. F., Stolz, J. R., Yan, J. Z., Swanson, G. T., & Green, W. N. (2017).
Selective and regulated trapping of nicotinic receptor weak base ligands and relevance to
smoking cessation. *Elife, 6.* doi: 10.7554/eLife.25651
Haga, K., Kruse, A. C., Asada, H., Yurugi-Kobayashi, T., Shiroishi, M., Zhang, C., et al. (2012).
Structure of the human M2 muscarinic acetylcholine receptor bound to an antagonist.
Nature, 482(7386), 547-551. doi: 10.1038/nature10753
Henderson, B. J., & Lester, H. A. (2015). Inside-out neuropharmacology of nicotinic drugs.
Neuropharmacol, 96(Pt B), 178-193. doi: 10.1016/j.neuropharm.2015.01.022
Houllier, N., Gouault, S., Lasne, M. C., & Rouden, J. (2006). Regio- and diastereoselective
functionalization   of   (-)-cytisine.   *Tetrahedron,   62*(50),   11679-11686.   doi:
10.1016/j.tet.2006.09.057
Imming, P., Klaperski, P., Stubbs, M. T., Seitz, G., & Gundisch, D. (2001). Syntheses and
evaluation of halogenated cytisine derivatives and of bioisosteric thiocytisine as potent
and selective nAChR ligands. *Eur J Med Chem, 36*(4), 375-388. doi: 10.1016/s0223-
5234(01)01222-3
Jeong, S. H., Sheridan, J., Newcombe, D., & Tingle, M. (2018). Plasma concentrations of cytisine,
a commercially available plant-based alkaloid, in healthy adult smokers taking
recommended doses for smoking cessation. *Xenobiotica, 48*(12), 1245-1248. doi:
10.1080/00498254.2017.1409916
Kabsch, W. (2010). Xds. *Acta Crystallographica D Biological Crystallography, 66*(Pt 2), 125-132.
doi: 10.1107/S0907444909047337
Kille, S., Acevedo-Rocha, C. G., Parra, L. P., Zhang, Z. G., Opperman, D. J., Reetz, M. T., et al.
(2013). Reducing codon redundancy and screening effort of combinatorial protein libraries
created by saturation mutagenesis. *ACS Synth Biol, 2*(2), 83-92. doi: 10.1021/sb300037w
Kozikowski, A. P., Chellappan, S. K., Xiao, Y., Bajjuri, K. M., Yuan, H., Kellar, K. J., et al. (2007).
Chemical medicine: novel 10-substituted cytisine derivatives with increased selectivity for
alpha4beta2 nicotinic acetylcholine receptors. *ChemMedChem, 2*(8), 1157-1161. doi:
10.1002/cmdc.200700073
Kuryatov, A., Luo, J., Cooper, J., & Lindstrom, J. (2005). Nicotine acts as a pharmacological
chaperone to up-regulate human $\alpha4\beta2$ acetylcholine receptors. *Mol Pharmacol, 68*(6),
1839-1851.
Lancet, D., & Pecht, I. (1976). Kinetic evidence for hapten-induced conformational transition in
immunoglobin MOPC 460. *Proc Natl Acad Sci U S A, 73*(10), 3549-3553. doi:
10.1073/pnas.73.10.3549
Le Houezec, J. (2003). Role of nicotine pharmacokinetics in nicotine addiction and nicotine
replacement therapy: a review. *Int J Tuberc Lung Dis, 7*(9), 811-819.
Marcaurelle, L. A., Johannes, C., Yohannes, D., Tillotson, B. P., & Mann, D. (2009). Diversity-
oriented synthesis of a cytisine-inspired pyridone library leading to the discovery of novel
inhibitors   of   Bcl-2.   *Bioorg   Med   Chem   Lett,   19*(9),   2500-2503.   doi:
10.1016/j.bmcl.2009.03.037
Marvin, J. S., Borghuis, B. G., Tian, L., Cichon, J., Harnett, M. T., Akerboom, J., et al. (2013). An
optimized fluorescent probe for visualizing glutamate neurotransmission. *Nat Methods,*
10(2), 162-170. doi: 10.1038/nmeth.2333
Marvin, J. S., Scholl, B., Wilson, D. E., Podgorski, K., Kazemipour, A., Muller, J. A., et al. (2018).
Stability, affinity, and chromatic variants of the glutamate sensor iGluSnFR. *Nat Methods,*
15(11), 936-939. doi: 10.1038/s41592-018-0171-3

Mills, E. J., Wu, P., Spurden, D., Ebbert, J. O., & Wilson, K. (2009). Efficacy of pharmacotherapies for short-term smoking abstinance: a systematic review and meta-analysis. *Harm Reduct J, 6*, 25. doi: 10.1186/1477-7517-6-25

Morales-Perez, C. L., Noviello, C. M., & Hibbs, R. E. (2016). X-ray structure of the human alpha4beta2 nicotinic receptor. *Nature, 538*(7625), 411-415. doi: 10.1038/nature19785

Nasu, Y., Shen, Y., Kramer, L., & Campbell, R. E. (2021). Structure- and mechanism-guided design of single fluorescent protein-based biosensors. *Nat Chem Biol, 17*(5), 509-518. doi: 10.1038/s41589-020-00718-x

Ormo, M., Cubitt, A. B., Kallio, K., Gross, L. A., Tsien, R. Y., & Remington, S. J. (1996). Crystal structure of the *Aequorea victoria* green fluorescent protein. *Science, 273*(5280), 1392-1395.

Philipova, I., Stavrakov, G., Vassilev, N., Nikolova, R., Shivachev, B., & Dimitrov, V. (2015). Cytisine as a scaffold for ortho-diphenylphosphinobenzenecarboxamide ligands for Pd-catalyzed asymmetric allylic alkylation. *Journal of Organometallic Chemistry, 778*, 10-20. doi: 10.1016/j.jorganchem.2014.12.001

Pienko, T., Grudzien, M., Taciak, P. P., & Mazurek, A. P. (2016). Cytisine basicity, solvation, logP, and logD theoretical determination as tool for bioavailability prediction. *J Mol Graph Model, 63*, 15-21. doi: 10.1016/j.jmgm.2015.11.003

Post, M. R., Tender, G. S., Lester, H. A., & Dougherty, D. A. (2017). Secondary Ammonium Agonists Make Dual Cation-π Interactions in α4β2 Nicotinic Receptors. *eNeuro, 4*(2). doi: 10.1523/eneuro.0032-17.2017

Quan, J., & Tian, J. (2009). Circular polymerase extension cloning of complex gene libraries and pathways. *PLoS One, 4*(7), e6441. doi: 10.1371/journal.pone.0006441

Rego Campello, H., Del Villar, S. G., Honraedt, A., Minguez, T., Oliveira, A. S. F., Ranaghan, K. E., et al. (2018). Unlocking Nicotinic Selectivity via Direct C–H Functionalization of (−)-Cytisine. *Chem, 4*(7), 1710-1725. doi: 10.1016/j.chempr.2018.05.007

Rollema, H., & Hurst, R. S. (2018). The contribution of agonist and antagonist activities of α4β2* nAChR ligands to smoking cessation efficacy: a quantitative analysis of literature data. *Psychopharmacology (Berl), 235*(9), 2479-2505. doi: 10.1007/s00213-018-4921-9

Rollema, H., Shrikhande, A., Ward, K. M., Tingley, F. D., 3rd, Coe, J. W., O'Neill, B. T., et al. (2010). Pre-clinical properties of the alpha4beta2 nicotinic acetylcholine receptor partial agonists varenicline, cytisine and dianicline translate to clinical efficacy for nicotine dependence. *Br J Pharmacol, 160*(2), 334-345. doi: 10.1111/j.1476-5381.2010.00682.x

Rose, J. E., Behm, F. M., Westman, E. C., Levin, E. D., Stein, R. M., & Ripka, G. V. (1994). Mecamylamine combined with nicotine skin patch facilitates smoking cessation beyond nicotine patch treatment alone. *Clin Pharmacol Ther, 56*(1).

Rouden, J., Ragot, A., Gouault, S., Cahard, D., Plaquevent, J. C., & Lasne, M. C. (2002). Regio- and diastereoselective functionalization of (-)-cytisine: an unusual N-C acyl migration. *Tetrahedron-Asymmetry, 13*(12), 1299-1305. doi: Pii S0957-4166(02)00271-9

Doi 10.1016/S0957-4166(02)00271-9

Schiefner, A., Breed, J., Bosser, L., Kneip, S., Gade, J., Holtmann, G., et al. (2004). Cation-pi interactions as determinants for binding of the compatible solutes glycine betaine and proline betaine by the periplasmic ligand-binding protein ProX from Escherichia coli. [Research Support, Non-U.S. Gov't]. *J Biol Chem, 279*(7), 5588-5596. doi: 10.1074/jbc.M309771200

Shivange, A. V., Borden, P. M., Muthusamy, A. K., Nichols, A. L., Bera, K., Bao, H., et al. (2019). Determining the pharmacokinetics of nicotinic drugs in the endoplasmic reticulum using biosensors. *J Gen Physiol, 151*(6), 738-757. doi: 10.1085/jgp.201812201

Smith, D., Allerton, C., Kalgutkar, A., van de Waterbeemd, H., & Walker, D. (2012). *Pharmacokinetics and Metabolism in Drug Design* (3rd ed.). Weinheim: Wiley.

Srinivasan, R., Pantoja, R., Moss, F. J., Mackey, E. D. W., Son, C., Miwa, J., et al. (2011). Nicotine
upregulates α4β2 nicotinic receptors and ER exit sites via stoichiometry-dependent
chaperoning. *J. Gen .Physiol., 137*, 59-79. doi: 10.1085/jgp.201010532
Studier, F. W. (2005). Protein production by auto-induction in high-density shaking cultures.
Protein Expression and Purification, 41(1), 207-234. doi: 10.1016/j.pep.2005.01.016
Tashkin, D. P. (2015). Smoking Cessation in Chronic Obstructive Pulmonary Disease. *Semin
Respir Crit Care Med, 36*(4), 491-507. doi: 10.1055/s-0035-1555610
Tsien, R. Y. (1998). The green fluorescent protein. *Annual Review of Biochemistry, 67*, 509-544.
doi: 10.1146/annurev.biochem.67.1.509
Turner, J. R., Castellano, L. M., & Blendy, J. A. (2011). Parallel anxiolytic-like effects and
upregulation of neuronal nicotinic acetylcholine receptors following chronic nicotine and
varenicline. *Nicotine Tob Res, 13*(1), 41-46.
Unger, E. K., Keller, J. P., Altermatt, M., Liang, R., Matsui, A., Dong, C., et al. (2020). Directed
Evolution of a Selective and Sensitive Serotonin Sensor via Machine Learning. *Cell,
183*(7), 1986-2002 e1926. doi: 10.1016/j.cell.2020.11.040
Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., et al. (2011).
Overview of the CCP4 suite and current developments. *Acta Crystallographica D
Biological Crystallography, 67*(Pt 4), 235-242. doi: 10.1107/S0907444910045749
Xiu, X., Puskar, N. L., Shanata, J. A., Lester, H. A., & Dougherty, D. A. (2009). Nicotine binding
to brain receptors requires a strong cation-π interaction. *Nature, 458*(7237), 534-537. doi:
10.1038/nature07768
964