Title

In vivo reprogramming of murine host immune response genes following *Leishmania major* infection.

Gopinath Venugopal[a,1], Jordan T. Bird[b,c,1], Charity L. Washam[b,c], Hayden Roys[a], Anne Bowlin[a], Stephanie D. Byrum[b,c,*], and Tiffany Weinkopff[a,*]

aDepartment of Microbiology and Immunology, and bDepartment of Biochemistry and Molecular Biology, College of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR, USA 72205; cArkansas Children's Research Institute, Little Rock, AR, USA 72202

1 G.V. and J.T.B. contributed equally to this work.

* To whom correspondence may be addressed. Email: TSWeinkopff@uams.edu, SBYRUM@uams.edu

Classification: Biological Sciences

Keywords: *Leishmania major*, RNA sequencing, Transcripts, Differentially Expressed Genes, Macrophages, Blood Endothelial Cells, Lymphatic Endothelial Cells

Abstract

Leishmania parasites cause cutaneous leishmaniasis (CL), a pathologic disease characterized by disfiguring, ulcerative skin lesions. Both parasite and host gene expression following infection with various *Leishmania* species has been investigated in vitro, but global transcriptional analysis following *L. major* infection in vivo is lacking. Thus, we conducted a comprehensive transcriptomic profiling study combining bulk RNA sequencing (RNA-Seq) and single-cell RNA sequencing (scRNA-Seq) to identify global changes in gene expression in vivo following *L. major* infection. Bulk RNA-Seq analysis revealed that host immune response pathways like the antigen processing and presentation pathway were significantly enriched amongst differentially expressed genes (DEGs) upon infection, while ribosomal pathways were significantly downregulated in infected mice compared to naive controls. scRNA-Seq analyses revealed cellular heterogeneity including distinct resident and recruited cell types in the skin following murine *L. major* infection. Within the individual immune cell types, several DEGs indicative of many interferon induced GTPases and antigen presentation molecules were significantly enhanced in the infected ears including macrophages (*Gbp2, H2-K1, H2-Aa, H2-Ab1*), resident macrophages (*H2-K1, H2-D1, Gbp4, Gbp8, Gbp2*), and inflammatory monocytes (*Gbp2, Gbp5, Gbp7, Gbp3*). Ingenuity Pathway Analysis of scRNA-Seq data indicated the antigen presentation pathway was increased with infection, while EIF2 signaling is the top downregulated pathway followed by eIF4/p70S6k and mTOR signaling in multiple cell types including macrophages, BECs, and LECs. Altogether, this transcriptomic profile highlights known recruitment of myeloid cells to lesions and recognizes a previously undefined role for EIF2 signaling in murine *L. major* infection in vivo.

Introduction

Leishmaniasis is a multifaceted disease caused by different species of obligate intracellular protozoan parasites of the genus *Leishmania*, belonging to the Trypanosomatid family. Depending on the complex interaction between the species and the host immune system, the disease can vary in severity resulting in a wide spectrum of clinical outcomes that have been classified into the following categories: cutaneous leishmaniasis (CL), mucocutaneous leishmaniasis (MCL), or diffuse CL (DCL) where symptoms remain localized to skin or mucosal surfaces, and a life-threatening condition called visceral leishmaniasis (VL) where parasites migrate to the internal organs like the liver, spleen and bone marrow[1]. As per the World Health Organization (WHO), leishmaniasis is still considered to be major public health problem due to its annual incidence up to 1.7 million new cases worldwide[1,2]. *Leishmania* spp. possess a variety of virulence mechanisms, which helps parasites to survive and replicate inside the parasitophorous vacuoles of macrophages, and elimination of parasites by macrophages is critical for host resistance[3]. Although, macrophages are the primary host cell for *Leishmania* parasites, recruited neutrophils[4–6] and dendritic cells[7–9] at the site of infection can also harbor parasites, suggesting that these immune cells could play an important role in host-parasite interplay. *Leishmania* parasites use several strategies to evade the host immune response for its intracellular survival including modulating the host immune response by altering T cell responses, impeding antigen display by MHCII, hindering nitric oxide production[10,11]. Importantly, *Leishmania* parasites can escape from oxidative burst and they fail to activate optimal macrophage innate immune responses[12,13]. However, changes at the transcriptional level following *Leishmania* infection within different cell types and especially within the hostile tissue microenvironment is still largely unknown.

Of the species belonging to subgenus *Leishmania*, *L. major* is an important

etiological agent of cutaneous leishmaniasis (CL) and possesses clinical and

epidemiological importance, especially in parts of Asia, the Middle East, Northern

Africa, and Southern Europe[14]. Although CL is not fatal and considered to be a self-

healing disease, the development of nodules or papules followed by ulcerations at the

site of promastigote infection is the hallmark of the disease; importantly, both parasite

replication and the host immune response can contribute to the disease pathology[14].

CL produces permanently disfiguring skin lesions that are associated with massive

immune cell recruitment, across the blood vascular endothelium, and into the skin

where the parasite resides[15–19]. While CD4$^+$ Th1 cells producing IFNγ are required to

activate macrophages to kill parasites, the exacerbated activation and sustained

recruitment of immune cells including neutrophils, NK cells, Ly6C$^+$ inflammatory

monocytes, and CD4$^+$ and CD8$^+$ T lymphocytes induces a chronic inflammatory

response; this chronic inflammation leads to tissue necrosis and skin damage, a

feature of non-healing lesions[20–22].

The *L. major* genome was completed in 2005[23]. Until recently, many studies

examining the host response relied mostly on microarray-based or serial analysis of

gene expression tag approaches. Using these approaches, previous studies

compared the host gene expression profile between infection with promastigotes and

amastigotes or different species of *Leishmania*[24–30]. Transcriptomic studies

dramatically enhanced our understanding of CL. In general, these studies concluded

that human or murine macrophages infected by various *Leishmania* species

downregulate pro-inflammatory gene expression, while concomitantly upregulating the

expression of anti-inflammatory genes[31,32]. However, microarray-based approaches

have technical limitations such as hybridization and cross hybridization artefacts, dye-

based detection issues, certain probes cannot be included on the microarrays, and the inability to detect 5' and 3' UTRs boundaries[33,34]. In recent years, RNA-Seq has emerged as a powerful tool to study transcriptional changes in many disease conditions due to its high sensitivity. Moreover, transcriptomic profiling using RNA-Seq following infection with various species of *Leishmania* has been mostly applied to in vitro experiments and some studies have investigated transcriptional changes in the human or murine host as well as the parasite simultaneously[35–42]. A recent study comparing the gene expression profile of primary cutaneous lesions from *L. braziliensis*-infected patients with or without pentavalent antimony treatment revealed most of the differentially expressed transcripts were correlated with components of cytotoxicity related pathways and parasite load in the skin[39]. While another study using RNA-Seq from the same group revealed a consistent and significant myeloid interferon stimulated gene (ISG) signature in skin lesions from *L. braziliensis*-infected patients[43]. Altogether, these studies revealed that the host immune response upregulates transcripts related to both pro-inflammatory and anti-inflammatory responses during leishmaniasis[44,45]. Furthermore, none of the RNA-Seq studies to date have characterized the global transcriptional reprogramming following *L. major* infection in vivo, which is the most widely used murine model of CL to study disease pathogenesis and parasite-host interactions. The *Leishmania* field also lacks a comprehensive murine transcriptomic profile that applies more recent genomic technologies like scRNA-Seq to identify transcriptomic changes within individual cell types in vivo. Therefore, the purpose of this study was to generate comprehensive transcriptomic profile using a combination of bulk RNA-Seq and scRNA-Seq to identify global changes in gene expression that occur during murine *L. major* infection in vivo.

Our bulk RNA-Seq transcriptomic analyses revealed the antigen presentation pathway was significantly upregulated while ribosomal pathways were significantly downregulated by the host following *L. major* infection. Our scRNA-Seq analyses revealed a cellular heterogeneity including distinct resident and recruited cell types at the site of *L. major* infection. Confirming the bulk RNA-Seq, we found macrophages, blood endothelial cells (BECs), and lymphatic endothelial cells (LECs) display a transcriptomic profile associated with increased antigen presentation and decreased EIF2, eIF4/p70S6K, and mTOR signalling suggesting cells in lesions are undergoing a stress response while they participate in Th1 protective immunity. Overall, this study combines bulk RNA-Seq and scRNA-Seq to assemble a comprehensive dataset that defines how the murine host response reprograms individual cell types following *L. major* infection to combat the infection.

Results

Mouse model of *L. major* infection

To characterize the transcriptomic landscape during *L. major* infection in vivo, we employed next-generation genome sequencing on RNA analysis with ears from mice that were infected with *L. major* promastigotes and uninfected naive control ears. During this experimental murine model of *L. major* infection, lesion volume peaks between 4-6 weeks post-infection (p.i.) and then the lesion resolves spontaneously, and parasites are present in lesions at this time point (Fig. 1A-B). As a result, the host transcriptome was investigated from ear samples collected from experimentally-infected mice at 4 weeks p.i. and compared to naive controls.

Enriched pathways for differentially expressed genes during *L. major* infection by bulk RNA-Seq

To compare the gene expression profiles of infected and naive mice, bulk RNA-Seq was performed on *L. major*-infected ears and naive control ears. Transcriptional analysis revealed that ears from infected mice and naive controls were distinct from one another, as determined by multidimensional scaling (MDS) plot and DEG analysis. MDS plot shows the positions of each sample, with samples from different experimental groups being well separated, and samples from the same experimental group clustering together (Fig. 1C). Therefore, the distance between samples reveals the distinct pattern of gene expression between the infected and naive animals (Fig. 1C). To investigate transcriptomic signatures associated with infection, we carried out DEG analysis between infected mice and naive controls by comparing the RNA-Seq read counts of the various genes and subsequently applying the cut-off criteria. High and low expression genes (logCPM><1) were included in the volcano plot showing transcriptional differences observed between infected and naive ears (Fig. 1D). The gene expression profiles derived from the RNA-Seq data were calculated using the RPKM method and a fold change >2 and $p < 0.05$ were considered statistically significant. Of more than 10,800 genes that were detectable in the infected ears, we observed that 211 genes were upregulated and 34 genes were downregulated, while 10,014 genes did not show any significant differences between naive and *L. major*-infected ears (Fig. 1D).

Gene set enrichment analysis (GSEA) using KEGG pathways revealed a total of 276 enriched pathways which includes pathways involved in both disease conditions and molecular signaling networks. Specifically, the antigen processing and presentation pathway was found to be significantly enriched amongst DEGs, while the ribosomal pathway was significantly downregulated during *L. major* infection (Fig. 1E-F). In addition to antigen processing and presentation, we observed many other host

immune response pathways upregulated with infection including: cytokine-cytokine

receptor interaction, phagosome, chemokine signaling, cell-adhesion molecules

pathway, NK cell mediated cytotoxicity, leukocyte trans-endothelial migration and Fcγ

receptor-mediated phagocytosis (Fig. 1E). Conversely, top 20 downregulated

pathways enriched for DEGs in *L. major* infection were related to ribosomal translation,

mineral absorption, ECM-receptor interaction, biosynthesis of unsaturated fatty acids,

and steroid biosynthesis (Fig. 1F). The top 10 KEGG pathways for both upregulated

and downregulated pathways with the fold change and adjusted '*p*' value are listed in

Table 1. A number of disease-specific KEGG pathways appeared prominent in the

enrichment analysis including *Staphylococcus aureus* infection, autoimmune thyroid

disease, and graft vs. host disease (Table S1). Importantly, leishmaniasis emerged as

one of the top disease pathways highlighting the quality of the input data for the

analysis (Table S1).

Differential gene expression in immune-related pathways during *L. major*

infection by bulk RNA-Seq

Using hierarchical clustering analysis, we found that a large number of genes were

robustly induced in the infected ears compared to the naives (Fig. 2A-D). A heat map

of the DEGs shows the expression profiles of infected and naive mice resulted in

separate clusters (Fig. 2). Hierarchical clustering reveals the host immune response

to *L. major* infection is closely linked with DEGs from the antigen processing and

presentation pathway (*Cd4, H2-Q6, H2-M3, H2-Q4, Ifng, Cd8b1, H2-T22, Rfx5, Tap1,*

H2-Q7), chemokine signaling (*Cxcl9, Ccl5, Ccr7, Cxcl10, Cxcl5, Cxcl16, Cxcl1, Fgr,*

Pik3cd), and cell adhesion molecules (*Cd4, Itgam, Itgal, Ctla4, Icos Itga6, Cd274,*

Cd28, Cd86, Selplg, Vcam1) (Fig. 2A-C). Additionally, other immune network

pathways enriched for DEGs included cytokine-cytokine receptor interaction,

phagosome, toll-like receptor signaling, and leukocyte trans-endothelial migration pathway (Fig. S1A-D). In contrast, the biological processes downregulated with infection include ribosomal biogenesis (*Rpl3, Rpl37, Rps5, Rpl11, Rplp1, Rpl28, Rpl19, Rps28, Rps14*) (Fig. 2D). Of note, infected mice clustered together for antigen processing and presentation, chemokine signaling, and the cell adhesion molecules pathways (Fig. 2A-C), but one mouse in each group clustered with the opposing experimental group for the ribosomal pathway (Fig. 2D). Overall, these results demonstrate the host transcriptome undergoes reprogramming in the skin during *L. major* infection.

scRNA-Seq reveals the cellular heterogeneity and altered transcriptomic profile of individual cell types during murine *L. major* infection in vivo

The bulk RNA-Seq analysis revealed global changes in the transcriptional profile between infected mice and naive controls following *L. major* inoculation. To further investigate transcriptomic changes within individual cell types present in leishmanial lesions, scRNA-Seq was performed to provide a deeper understanding of how individual cells function in the tissue microenvironment. Single cells from the ears of infected and naive mice were bar-coded and sequenced using the droplet-based 10X Genomics Chromium platform (Fig. 3A). After quality control assessment and filtering, the datasets were processed using Cell Ranger software. Unbiased hierarchical clustering using Seurat provides single-cell transcriptional profiling with 26,558 cells and displayed the cellular heterogeneity which includes both resident and recruited cell types. Cell populations from 35 distinct cell types were defined using canonical markers from published literature and online databases (Fig. 3B)[46]. The dot plot representing the cell type-specific canonical markers for each cell lineage used to distinguish the 35 distinct clusters is provided (Fig. 4A). Amongst the 35 cell types, we

identified 16 cell types containing immune cells. Feature plots show the expression of cell type-specific canonical markers (in addition to the cell type-specific canonical markers in Fig. 4A) for 12 clusters with corresponding cell types (Fig. 4B). Additionally, a heatmap shows the canonical cell type markers from all the immune cell types along with blood endothelial cells (BECs) and lymphatic endothelial cells (LECs) (Fig. S2).

Detection of *Leishmania major* transcripts in multiple cell types other than macrophages

Additionally, we aligned the reads to *Leishmania major* (LM) reference genome to detect the presence of *Leishmania* transcripts in 35 different cell type clusters. Interestingly, the differential expression of LM transcripts from scRNA-Seq revealed 20 of 35 cell types have at least one LM transcript within that cell. As predicted, we found macrophages are the top immune cell type expressing LM transcripts with about 10% of cells containing LM transcripts (Fig. 5A-B). We found at least 2% of cells contain LM transcripts in other immune cell types such as resident macrophages, DCs, and neutrophils. At least 1% of the cells in CD4[+] Th cells, CD8[+] cytotoxic T cells, T regulatory cells, and basophils also possessed detectable LM transcripts (Fig. 5A-B). Consistent with previous findings, we found fibroblasts and keratinocytes also harbor LM transcripts[47]. Surprisingly, we also detected LM transcripts in >5% myoblasts but not myocytes and almost 1% of the BECs. Overall, our data shows that multiple other cell types at the infection site harbors LM transcripts suggesting other cell types maybe infected with parasites alongside macrophages, which is the well-established primary host cell for *Leishmania* parasites.

scRNA-Seq confirms the immune cell recruitment at the site of *L. major* infection in vivo

Murine *L. major* infection leads to the recruitment of immune cells such as inflammatory monocytes, neutrophils, and macrophages to the site of infection. Particularly macrophages play roles in housing the parasite as the replicative niche for the pathogen, as well as a role in parasite control by killing the pathogen. BECs mediate immune cell recruitment to the infected and inflamed tissue and LECs promote immune cell migration away from the infected skin. Therefore, we speculate that immune cells and ECs participate in parasite control and/or immunopathology during CL. As a result, the remainder of the study focuses on 7 immune cell types along with the BEC and LEC clusters. Our UMAP projection displays 13,034 cells in naive ears and 13,524 cells in the infected ears (Fig. 6A). Consistent with previous findings in CL, the UMAP plot confirms a significant recruitment of various immune cell types such as inflammatory monocytes, neutrophils, macrophages, dendritic cells, NK cells, and CD4$^+$ and CD8$^+$ T cells in the infected ears that are seen at higher frequencies compared to naive controls (Fig. 6A). Concordant with our scRNA-Seq results, flow cytometric analysis detected a significant increase in the frequency and cell number of macrophages (Fig. 6B-C), Ly6C$^+$ inflammatory monocytes (Fig. 6D-E) and neutrophils (Fig. 6F-G) in infected ears compared to naive controls, while no significant alterations in the BEC or LEC populations were observed (Fig. 6H-J). Altogether, these data confirm the enhanced immune cell migration during *L. major* infection and transcriptional changes within these individual cell types were investigated.

Differential gene expression of immune cell types during *L. major* infection

To explore the transcriptional changes in a cell type-specific manner, the DEGs were compared between infected and naive mice within an individual cell type following *L. major* infection. A volcano plot showing DEGs for macrophages, resident

macrophages, inflammatory monocytes, and neutrophils reveals several markers indicative of an increase in myeloid cells in leishmanial lesions (Fig. 7A-D). For instance, transcripts commonly elevated within the top 10 DEGs in myeloid cells and dendritic cells (DCs) include *B2m, H2-K1, Gbp2, Iigp1,* whereas multiple ribosomal proteins were significantly downregulated within the top 10 DEGs among myeloid cells and DCs (Table 2, 3, and Fig. S3). We found consistent elevation of various interferon-induced GTPases like guanylate binding protein (GBP) transcripts with *L. major* infection in macrophages (*Gbp2*) (Table 2A), resident macrophages (*Gbp4, Gbp8, Gbp2*) (Table 2B), inflammatory monocytes (*Gbp2, Gbp5, Gbp7, Gbp3*) (Table 3A), and DCs (*Gbp2*) (Fig. S3). Many of the transcriptomic differences detected in myeloid cells, such as elevated GBPs, were also found in EC populations. For instance, BECs expressed increased *Gbp4* and *Gbp2*, and LECs expressed increased *Gbp4, Gbp2,* and *Gbp7* upon infection (Table 4A-B). Furthermore, we detected a significant elevation of both MHCI and MHCII molecules in the infected ears in myeloid cells and ECs, which include *H2-K1, H2-Aa, H2-Ab1* in macrophages; *H2-K1, H2-D1* in resident macrophages; *H2-K1* in DCs, *H2-K1, H2-Aa, H2-Ab1* in BECs; and *H2-K1, H2-D1, H2-Q7, H2-Aa, H2-Ab1* in LECs (Table 2, 4, and Fig. S3). In contrast to myeloid cells, DCs, and ECs, we only detected few transcripts that are significantly elevated with infection in T cells including *B2m, Satb1, Gm42418, Gimap6* in CD4[+] T cells and *Gm42418* in CD8[+] T cells (Fig. S4).

In addition, many chemokines were differentially modulated in myeloid cells and ECs with *L. major* infection (Fig. 7). Specifically, *Cxcl9* was significantly elevated in macrophages, resident macrophages, inflammatory monocytes, DCs, BECs, and LECs following *L. major* infection (Fig. 7 and Fig. S3). In contrast, we found significant downregulation of *Ccl24* in resident macrophages, *Cxcl12* in BECs, and *Ccl17* in DCs

following *L. major* infection (Fig. 7B, 7E and Fig S3). Also important in immune cell

recruitment, selectins (*Selp, Sele*) and adhesion molecules (*Vcam1)* were significantly

upregulated in BECs with infection, while tight junction molecules like *Cldn5* were

downregulated (Fig. 7E). Of note, known canonical markers were significantly elevated

with *L. major* infection including *Arg1, Nos2,* and *Pla2g7* in macrophages, *Fcgr4, C3,*

and *Ccl8* in resident macrophages, and *Ifitm1, Syngr2, Cd200, Ccr7* in DCs (data not

shown). The complete list of DEGs that are enriched during *L. major* infection from

other cell type clusters such as fibroblasts, keratinocytes, chondrocytes, sebaceous

glands, basophils, upper hair follicle cells, pericytes, schwann cells, mast cells,

myocytes and myoblasts can be found in Gene Expression Omnibus database with

the GEO accession number - GSE181720.

Characterization of upstream gene regulators and canonical pathways during *L.*

***major* infection in vivo**

To determine the cellular and biological mechanisms at the molecular level during *L.*

major infection, IPA analysis was performed to define the gene signature for each

individual cell type at the site of infection. IPA analysis of our scRNA-Seq data revealed

several known and unknown canonical pathways, upstream regulators, and disease-

based functional networks. Here, we present the genes that are significantly altered

(adj. p value < 0.05) in macrophages, BECs, and LECs from infected ears compared

to naive controls.

Upstream gene regulators: Our IPA analysis on macrophages, BECs, and LECs

revealed potential transcription factors as well as transcriptional targets like anti- and

pro-inflammatory genes. In macrophages, we observed 651 upstream regulators in

total which include 38 upregulated and 17 downregulated gene regulators. We found

cytokines like *IFNγ, IL-4, IL-13, IFNβ1, TNFα*, and transcriptional regulators such as

HIF1α, STAT1, CTCF, TP73, IRF1, MXD1, ATF4, SPI1 mediate macrophage

activation upon infection (Table 5). In contrast, transcriptional regulators like *MLXIPL,*

MYC, TP53, MYCL, CEBPB, GATA1 were inhibited and no cytokines were identified

to downregulate macrophages activation following infection (Table 5). We identified

32 regulators were activated and 64 regulators were inhibited with infection in BECs.

The upregulated cytokines activating BECs included *IFNγ, TNF, IL-2, IL-4,* while *IL-10*

downregulates BEC activation (Table 6). Additionally, we found transcriptional

regulators that are either activated (*IRF3, STAT1, IRF7, MXD1*) or inhibited (*MLXIPL,*

MYC, TRIM24, TP53, SIRT1, HSF1, MYCL, GLIS2, CEBPB, NFE2L2) in BECs

following infection (Table 6). Likewise, 212 upstream regulators were detected of LECs

including 17 activated and 23 inhibited regulators. Corresponding to BECs, *IFNγ*

increases LEC activation, while *IL-10* downregulates LEC activation upon infection

(Table 7). In the LECs, we detected activated transcriptional regulators such as *IRF3,*

STAT1, and *IRF7*; however, *MLXIPL, MYC, TRIM24, TP53, SIRT1, MYCL, TARDBP,*

STAT6, AIRE, LDB1, HNF4A, and *NFE2L2* were inhibited in LECs comparing infected

mice to naive controls (Table 7).

Canonical pathways: IPA analysis highlighted an unknown role for eukaryotic

translation initiation factor 2 (EIF2) signaling which includes large ribosomal proteins

(Rpl) and small ribosomal proteins (Rps) that were significantly downregulated with

infection compared to naive animals (Fig. 8). Remarkably, the EIF2 pathway was the

top downregulated pathway amongst multiple cell types including macrophages

(Ranked as 1 amongst 377), BECs (Ranked as 1 amongst 257), and LECs (Ranked

as 1 amongst 145) (Fig. 8A, C & E). This was followed by an involvement of mTOR

signalling and eIF4/p70S6K signalling in macrophages, BECs, and LECs from infected

ears. Alongside the IPA pathway results, the expression of individual transcripts for

each of the corresponding pathways is provided for macrophages (Fig. 8B), BECs

(Fig. 8D), and LECs (Fig. 8F). These data show the top 10 transcripts in the EIF2

signalling pathway in macrophages, BECs, and LECs, which includes many subunits

of ribosomal proteins and these transcripts are mostly downregulated in the infected

ears compared to naive controls (Fig. 8B, D & F). In contrast, the IPA analysis revealed

the antigen presentation pathway was increased with infection, and this pathway was

also a common feature of infection being the top elevated pathway for macrophages,

BECs, and LECs (Fig. 8B, D & F). The antigen presentation pathway was enriched in

transcripts such as *B2m, Cd74, H2-K1, H2-Aa, H2-Ab1, H2-Eb1* that were elevated

with *L. major* infection in macrophages, BECs, and LECs. In addition to macrophages,

BECs, and LECs, we also noted that an involvement of mTOR signalling pathway is

consistent among the other immune cell types apart of this study such as inflammatory

monocytes, DCs, and CD4$^+$ T cells, as mTOR signalling is top 5 in the list of pathways

(Fig. S5). In summary, EIF2 signaling is the top downregulated pathway in BECs,

LECs, macrophages, as well as other immune cell types from infected ears.

Discussion

We conducted a comprehensive high-resolution transcriptomic analysis using both

bulk and scRNA-Seq approaches to discover the global changes in the gene

expression that occurs following *L. major* infection in vivo. Through our bulk RNA-Seq

analyses, we identified many differentially regulated novel transcripts in immune

compartments that are related to host immune response pathways. We found

significant enrichment of DEGs in the antigen processing and presentation pathway

following *L. major* infection. Specifically, our data indicate that *L. major* infection

upregulates many MHC molecules belonging to the antigen processing and

presentation pathway along with inflammatory cytokines such as *IFNγ* following *L.*

major infection. These findings are consistent with the well-established role of the Th1

immune response in parasite control; we confirm antigen presentation is a critical

process in host defense to *Leishmania* infection[48–53]. Additionally, we noted the

enrichment of DEGs specific for other host immune response pathways such as

chemokine signalling, cell adhesion molecules, and cytokine-cytokine receptor

interactions in infected ears. In contrast, our bulk RNA-Seq results revealed DEGs

associated with the ribosomal pathway were downregulated following *L. major*

infection. While the importance of the downregulation of transcripts that encode 40S

and 60S ribosomal subunits has not been studied in *Leishmania*, these findings

suggest cells at the site of infection are actively controlling translation and/or

ribogenesis or undergoing a stress response similar to bacterial, viral infection and

cancer[54–56]. Altogether, our bulk RNA-Seq results confirm a known role for the

importance of antigen presentation and highlight an unknown feature of

downregulating ribosomal subunits in CL.

Our scRNA-Seq analysis revealed many novel transcripts from various cell

types that remain largely unexplored at the site of *L. major* infection. Through our

scRNA-Seq data, we defined 35 distinct cell type populations by using canonical

markers specific for different cell types including both resident and recruited cells

following *L. major* infection. In agreement with previously published results[10,20], our

scRNA-Seq analysis confirmed a significant increase in various immune cell types at

the site of infection with recruited myeloid cells such as neutrophils, inflammatory

monocytes, and macrophages in lesions (Fig. 6). The transcriptional signature of

interferon-induced GTPases like GBPs were significantly upregulated in

macrophages, resident macrophages, inflammatory monocytes, DCs, BECs, and

LECs following *L. major* infection. These data suggest GBPs may play a protective

role in both immune and nonimmune cells during *L. major* infection. GBPs are involved in controlling intracellular pathogen replication and specifically mediating the protection against intracellular pathogens such as *Listeria monocytogenes* and *Mycobacterium bovis*[57] [58]. Importantly, mice deficient in GBP genes are more susceptible to *Toxoplasma gondii*[59]. Moreover, GBPs restrict *L. donovani* growth in nonphagocytic cells such as murine embryonic fibroblasts in an IFNγ-dependent manner[60].

We detected several transcripts for chemokines (*Cxcl9, Cxcl10*) in myeloid cells that are elevated with infection that may mediate myeloid cell accumulation at the site of infection. Our results revealed a significant downregulation of *Ccl24* transcript in resident macrophages from infected ears compared to naive controls. A recent study demonstrated dermal tissue-resident macrophages shift toward a pro-inflammatory state in *L. major*-infected mice lacking IL-4/IL-13 from eosinophils, which is mainly regulated by CCL24 from resident macrophages[61]. Also, potentially mediating myeloid cell migration, we found BECs in infected skin express elevated transcripts for selectins (*Sele, Selp*) and adhesion molecules (*Vcam1*), while concomitantly downregulating transcripts responsible for junctional stability. Mice doubly deficient in E- and P-selectin develop significantly less inflammation following *L. major* infection[62]. Taken with our findings, these data suggest BECs play an active role during CL to recruit immune cells into the site of infection. Indeed, these scRNA-Seq data also correlate with our bulk RNA-Seq results which indicate a role of leukocyte trans-endothelial migration pathway in infected ears when compared to naive controls. Overall, our scRNA-Seq data reveals cells in leishmanial lesions exist in pro-inflammatory environment, but the actual host protective role of individual DEGs within each cell type requires further investigation.

We detected the presence of *L. major* transcripts in multiple cell types including myeloid cells like macrophages, neutrophils, and DCs which are all known to harbor parasites[47,63,64]. We also detected *L. major* transcripts in stromal cells such as fibroblasts and keratinocytes, which can be infected by parasites[65,66]. Surprisingly, we also found evidence of parasite transcripts in myoblasts, chondrocytes, and BECs which has not previously been reported. Importantly, the molecular techniques used in these studies cannot differentiate between living and dead parasites and we can only report cells harboring parasite transcripts. Therefore, subsequent studies will need to determine if parasites in other non-myeloid cells like myoblasts and BECs are viable and capable of replication and infection. Others have hypothesized *Leishmania* parasites might evade the host immune responses by seeking shelter in different non-macrophage cell types including fibroblasts and keratinocytes in addition to infection in neutrophils, macrophages and DCs which exhibit a more robust pro-inflammatory response against *Leishmania* parasites[67]. However, the presence of *L. major* transcripts from non-myeloid stromal cell types like myoblasts and BECs needs to be further explored to determine whether these cells can serve as safe havens for *L. major* parasites during chronic infection or provide a conduit for metastasis. Regardless, these results shed light on cell types previously not thought to harbor *Leishmania* parasites in the skin and provide an opportunity for new investigation.

Macrophages are the replicative niche for the parasite as well as the major cell type responsible for parasite control. BECs play a crucial role in regulating immune cell entry into inflamed tissues and LECs participate in immune cell migration out of the lesions. As a result of the critical roles of these cell types during *L. major* infection, we focused on identifying the upstream regulators and canonical pathways for macrophages, BECs, and LECs. Surprisingly, the antigen processing and

presentation pathway and EIF2 signaling were the most significant pathways in all three cell types (macrophages, BECs, and LECs) within infected ears. The antigen processing and presentation showed a positive activation z score indicating overall upregulation of the pathway, which is consistent with findings in human CL lesions[68]. Immunoproteasomes play a critical role in the immune response by degrading intracellular proteins to generate MHCI epitopes for effective antigen presentation[69]. While the increased expression of immunoproteasome genes in human lesions caused by *L. braziliensis* infection has been reported[68], our results reveal for the first time that LECs express higher levels of transcripts for immunoproteasomes (*psmb8* and *psmb9*), as well as transcripts involved in the antigen presentation pathway (*Tap1* and *Tapbp*) following *L. major* infection. These data suggest that ECs, and specifically LECs, may play an unknown role in antigen presentation during *L. major* infection, similar to viral infection and vaccination[69,70].

The EIF2 signaling pathway had the highest negative activation z score of all the pathways indicating overall deregulation during CL. To our knowledge, this is the first study highlighting EIF2 signaling as a novel candidate pathway for leishmaniasis. Eukaryotic initiation factor-2 (EIF2) is a GTP-binding protein, which initiates protein translation by delivering charged initiator met-tRNA onto the ribosome[71]. Upon subject to infection-induced cellular stress, EIF2 plays a significant role in attenuating translation initiation by phosphorylation of the alpha subunit of eIF2 leading to immediate shut-off of translation and activation of stress response genes[71]. Phosphorylation of eIF2α plays as a rate limiting step as it reduces active eIF2-GTP levels at translation initiation which ultimately results in a global reduction of protein synthesis[71,72]. Our data indicate that impaired EIF2 signaling is linked to the downregulation of many ribosomal subunits in macrophages, BECs, and LEC in *L.*

major-infected ears (Fig. 8). The known phenomenon of "protein shut off" has been

well described at the molecular level for some viruses[73], but there is little to no

evidence documenting this phenomenon during *L. major* infection. Generally, the

enhanced host response to viral and bacterial infections depends on the upregulation

of EIF2-mediated translational control, thereby reducing general protein

synthesis[55,56,73]. However, we found EIF2 signaling was downregulated with *L. major*

infection. Previously it was demonstrated that *L. major* promotes its survival by

downregulating macrophage protein synthesis, which is mainly mediated by host

translation repressor 4E-BP1[74]. Given the robust production of cytokines and

chemokines in leishmanial lesions, global protein synthesis does not seem impacted

by *L. major* infection in vivo, but careful analysis of EIF2 signaling has not been

performed to date.

Alongside EIF2 signalling, the involvement of mTOR and eIF4/p70S6K

signalling in macrophages, BECs, and LECs in infected mice hints at an important role

for hypoxia-induced oxidative stress at the site of *L. major* infected ears. Our results

indicate the activation of metabolic gene targets like hypoxia-inducible factor (HIF-1α)

in macrophages, which is consistent with our previous results and work by the Jantsch

lab showing leishmanial lesions are hypoxic[75–77]. Therefore, we speculate EIF2

signaling is downregulated as part of the stress response, potentially from hypoxia,

but mechanism by which EIF2 signaling is impaired in multiple cell types and how that

contributes to pathogen control or the pathogenesis of disease in CL is unknown.

Collectively, our transcriptome analysis not only provides the first comprehensive list

of gene expression at single-cell resolution, but also highlights a previously unknown

role of the highly conserved EIF2 signaling pathway in leishmaniasis. Future analysis

by us and others utilizing these datasets will expand our knowledge on the complex

immine networks and pathways participating in the host response to *Leishmania* infection.

Materials and methods

Animals

Female C57BL/6NCr mice were purchased from the National Cancer Institute. Mice were housed in the Division of Laboratory Animal Medicine at University of Arkansas for Medical Sciences (UAMS) under pathogen-free conditions and used for experiments between 6 and 8 weeks of age. All procedures were performed in accordance with the guidelines of the UAMS Institutional Animal Care and Use Committee (IACUC).

Parasite Infection in vivo

Leishmania major (WHO/MHOM/IL/80/Friedlin) strain was used. Parasites were maintained in vitro in Schneider's Drosophila medium (Gibco) supplemented with 20% heat-inactivated FBS (Invitrogen), 2 mM L-glutamine (Sigma), 100 U/mL penicillin, and 100 mg/mL streptomycin (Sigma). Metacyclic stationary phase promastigotes were isolated from 4–5 day cultures by Ficoll density gradient separation (Sigma). For ear dermal infections, 2×10^6 *L. major* promastigote parasites in 10 µL PBS (Gibco) were injected intradermally into the ear. Lesion development was monitored weekly by measuring ear thickness and lesion area with a caliper. Lesion volume was calculated. At 4-week post infection, ears were excised and enzymatically digested using 0.25 mg/mL Liberase (Roche) and 10 mg/mL DNase I (Sigma) in incomplete RPMI 1640 (Gibco) for 90 min at 37°C. After digesting, ears were minced manually to obtain cellular content in single cell suspensions. Parasite burdens were determined by limiting dilution assays (LDAs), as previously described[78].

Bulk RNA-Seq: Sample Preparation

The total RNA was isolated from the cell lysate of naive and infected ears by using Qiagen's RNeasy plus mini kit according to the manufacturer's instructions. The CTPR Genomics and Bioinformatics Core at the Arkansas Children's Research Institute (ACRI) prepared sequencing libraries from RNA samples by use of the Illumina TruSeq Stranded mRNA Sample Preparation Kit v2. for sequencing on the NextSeq 500 platform using Illumina reagents. The quality and quantity of input RNA was determined using the Advanced Analytical Fragment Analyzer (AATI) and Qubit (Life Technologies) instruments, respectively. All samples with RQN (RNA quality number) values of 8.0 or above were processed for sequencing. Sequencing libraries were prepared by use of the TruSeq Stranded mRNA Sample Prep Kit (Illumina). Briefly, total RNA (500 ng) was subjected to polyA selection, then chemically fragmented and converted to single-stranded cDNA using random hexamer primed reverse transcription. The second strand was generated to create double-stranded cDNA, followed by fragment end repair and addition of a single A base on each end of the cDNA. Adapters were ligated to each fragment end to enable attachment to the sequencing flow cell. The adapters also contain unique index sequences that allow the libraries from different samples to be pooled and individually identified during downstream analysis. Library DNA was PCR amplified and enriched for fragments containing adapters at each end to create the final cDNA sequencing library. Libraries were validated on the Fragment Analyzer for fragment size and quantified by use of a Qubit fluorometer. Equal amounts of each library was pooled for sequencing on the NextSeq 500 platform using a high output flow cell to generate approximately 25 million 75 base reads per sample.

Bulk RNA-Seq: Data Analysis

Following demultiplexing, RNA reads were checked for sequencing quality using

FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc) and MultiQC[79]

(version 1.6). The raw reads were then processed according to Lexogen's QuantSeq

data analysis pipeline with slight modification. Briefly, residual 3' adapters, polyA read

through sequences, and low quality (Q < 20) bases were trimmed using BBTools

BBDuk (version 38.52) (https://sourceforge.net/projects/bbmap/). The first 12 bases

were also removed per the manufacture's recommendation. The cleaned reads ($\geq$

20bp) were then mapped to the mouse reference genome (GRCm38/mm10/ensemble

release-84.38/ GCA_000001635.6) using STAR[80] (version 2.6.1a), allowing up to 2

mismatches depending on the alignment length (e.g. 20-29bp, 0 mismatches; 30-

50bp, 1 mismatch; 50-60+bp, 2 mismatches). Reads mapping to > 20 locations were

discarded. Gene level counts were quantified using HTSeq (htseq-counts)[81] (version

0.9.1) (mode: intersection-nonempty).

Genes with unique Entrez IDs and a minimum of ~2 counts-per-million (CPM)

in 4 or more samples were selected for statistical testing. This was followed by scaling

normalization using the trimmed mean of M-values (TMM) method[82] to correct for

compositional differences between sample libraries. Differential expression between

naive and infected ears was evaluated using limma voomWithQualityWeights[83] with

empirical bayes smoothing. Genes with Benjamini & Hochberg[84] adjusted p-values $\leq$

0.05 and absolute fold-changes $\geq$ 1.5 were considered significant.

Gene Set Enrichment Analysis (GSEA) was carried out using Kyoto

Encyclopedia of Genes and Genomes (KEGG) pathway databases and for each

KEGG pathway, a p-value was calculated using hypergeometric test. Cut-off of both *p*

< 0.05 and adjusted p-value/FDR value < 0.05 was applied to identify enriched KEGG

pathways. DEGs that are more than 1.5-fold in *L. major*-infected ears relative to

uninfected controls were used as input, with upregulated and downregulated genes

considered separately. Subsequently, the heat maps were generated using these

genes with complex Heatmap. All analyses and visualizations were carried out using

the statistical computing environment R version 3.6.3, RStudio version 1.2.5042, and

Bioconductor version 3.11. The raw data from our bulk RNA-Seq analysis were

deposited in Gene Expression Omnibus (GEO accession number - GSE185253).

scRNA-Seq Sample Preparation

The Genomics and Bioinformatics Core at the Arkansas Children's Research Institute

(ACRI) prepared NGS libraries from fresh single-cell suspensions using the 10X

Genomics NextGEM 3' assay for sequencing on the NextSeq 500 platform using

Illumina SBS reagents. The quantity and viability of cells input to the assay was

determined using Trypan Blue exclusion under 10X magnification. Library quality was

assessed with the Advanced Analytical Fragment Analyzer (Agilent) and Qubit (Life

Technologies) instruments.

scRNA-Seq Data Analysis

Demultiplexed fastq files generated by the UAMS Genomics Core were analyzed with

the 10x Genomics Cell Ranger alignment and gene counting software, a self-

contained scRNA-Seq pipeline developed by 10X Genomics. The reads are aligned

to the mm10 and *Leishmania major* reference transcriptomes using STAR and

transcript counts are generated[80,85] .

The raw counts generated by *cellranger count* were further processed using the

R package *Seurat*[86,87]. Low quality cells, potential cell doublets, and cells with high

percentage of mitochondrial genes were filtered out of the data. We filtered cells that

have unique feature counts over more the 75^{th} percentile plus 1.5 times the

interquartile range (IQR) or less than the 25^{th} percentile minus 1.5 time the IQR.

Additionally, we filtered cells with mitochondrial counts falling outside the same range

with respect to mitochondrial gene percentage. Following filtering the counts all 8

sequencing runs were merged. The counts are then normalized using the

LogNormalize method, which normalizes the feature expression measurements for

each cell by the total expression, multiplies this by a scale factor (10,000 by default),

and log-transforms the result. Next, the 2000 highest variable features are selected.

The data is then scaled, and linear regression is included to remove variation

associated with percent mitochondria and cell cycle status. Principle component

analysis (PCA) is performed on the scaled data. A JackStraw procedure was

implemented to determine the significant PCA components that have a strong

enrichment of low p-value features.

A graph-based clustering approach is applied[88]. Briefly, these methods embed

cells in a graph structure - for example a K-nearest neighbor (KNN) graph, with edges

drawn between cells with similar feature expression patterns, and then attempt to

partition this graph into highly interconnected 'quasi-cliques' or 'communities'. t-

distributed stochastic neighbor embedding (tSNE) and Uniform Manifold

Approximation and Projection (UMAP)[89] are non-linear dimensional reduction

techniques used to visualize and explore the results and are performed using Seurat.

Seurat *FindNeighbors* and *FindClusters* functions will be optimized to label clusters

based on the visual clustering in the projections. Seurat *FindAllMarkers* function finds

markers that define clusters by differential expression. It identifies positive markers of

a single cluster compared to all other cells and outputs the differential expression

results. These markers will be compared to known markers of expected cell types and

results from previous single-cell transcriptome studies in order to assign appropriate

cell type labels. Cell type determinations were made by manually expecting these

results and some clusters were combined if their expression was found to be similar.

Differential expression analysis will be performed using MAST, a GLM-framework that

treats cellular detection rate as a covariate[90]. The raw data from our scRNA-Seq

analysis were deposited in Gene Expression Omnibus (GEO accession number -

GSE181720)

Ingenuity Pathway Analysis (IPA)

QIAGEN's Ingenuity Pathway Analysis (IPA, QIAGEN Redwood

City, www.qiagen.com/ingenuity) was utilized to investigate the DEGs at the level of

biochemical pathways and molecular functions. We submitted our DEGs to functional

analysis with IPA, and IPA provided canonical pathways, diseases and function, and

upstream regulators based on the experimentally observed cause-effect relationships

related to transcription, expression, activation, molecular modification, etc. Z-score

analyses are used to assess the match between observed and predicted up and down

regulation patterns allowing for Bayesian scoring of the results.

Flow Cytometry Analysis with Statistics

The recruitment of immune cells during *L. major* infection was analyzed by flow

cytometry. Cells from both naive and infected ears were incubated with fixable Aqua

dye (Invitrogen) to assess cell viability, and cells were treated with FcγR blocking

reagent (Bio X Cell) prior to staining for the following markers: anti-CD45-AF700

(clone 30-F11), anti-Ly6C-PerCP-Cy5.5 (clone HK1.4), and anti-Ly6G-eFlour 450

(clone 1A8) were purchased from eBiosciences; anti-CD64-BV711 (clone X54-5/7.1),

anti-CD11b-BV605 (clone M1/70), anti-CD31-AF488 (390), and anti-podoplanin-

PE/Dazzle 594 (clone 8.1.1) were purchased from BioLegend. Cells were acquired

using an LSR II Fortessa flow cytometer (BD Biosciences) and analyzed using FlowJo

software version 10.2 (Tree Star). All statistical analyses were performed using Prism

version 8.0 (GraphPad Software, Inc.). Comparisons between groups were performed

using the two-tailed Students unpaired *t*-test.

Acknowledgements and funding sources

The authors would like to thank Dr. Lu Huang, the Huang lab, Lucy Fry and Conner

Webb for their critical reading of the manuscript. We would like to thank the veterinary

staff from the Division of Laboratory Animal Medicine (DLAM) and Andrea Harris from

Flow cytometry core facility at UAMS for their excellent technical assistance. This work

was supported by the Center for Microbial Pathogenesis and Host Inflammatory

Responses (funded by National Institutes of Health National Institute of General

Medical Sciences Centers of Biomedical Research Excellence Grant P20-GM103625)

and the Oak Ridge Associated Universities (ORAU) 2019 Ralph E. Powe Junior

Faculty Enhancement Award to Dr. Weinkopff. This work was also supported by the

UAMS Translational Research Institute in collaboration with the UAMS Division for

Diversity, Equity and Inclusion (DDEI) Mini Grants for Under-Represented Faculty

Researchers to Dr. Weinkopff as part of the NIH National Center for Advancing

Translational Sciences (NCATS), UL1-TR003107. This study was supported by the

Arkansas Children's Research Institute, the Arkansas Biosciences Institute, and the

Center for Translational Pediatric Research funded under the National Institutes of

Health National Institute of General Medical Sciences (NIH/NIGMS) grant

P20GM121293 and the National Science Foundation Award No. OIA-1946391. The

funders had no role in study design, data analysis, decision to publish or preparation

of the manuscript.

Competing interests:

The authors have declared that no competing interests exist.

References

1. Burza, S., Croft, S. L. & Boelaert, M. Leishmaniasis. *Lancet* **392**, 951–970 (2018).

2. Zhang, W. W. *et al.* A second generation leishmanization vaccine with a markerless attenuated Leishmania major strain using CRISPR gene editing. *Nat. Commun.* **11**, 1–14 (2020).

3. Kaye, P. M., Cruz, I., Picado, A., Van Bocxlaer, K. & Croft, S. L. Leishmaniasis immunopathology—impact on design and use of vaccines, diagnostics and drugs. *Seminars in Immunopathology* (2020) doi:10.1007/s00281-020-00788-y.

4. Laskay, T., Van Zandbergen, G. & Solbach, W. Neutrophil granulocytes - Trojan horses for Leishmania major and other intracellular microbes? *Trends Microbiol.* **11**, 210–214 (2003).

5. Ribeiro-Gomes, F. L. & Sacks, D. The influence of early neutrophil-Leishmania interactions on the host immune response to infection. *Front. Cell. Infect. Microbiol.* **2**, 59 (2012).

6. Peters, N. C. *et al.* In vivo imaging reveals an essential role for neutrophils in leishmaniasis transmitted by sand flies. *Science (80-.).* (2008) doi:10.1126/science.1159194.

7. León, B., López-Bravo, M. & Ardavín, C. Monocyte-Derived Dendritic Cells Formed at the Infection Site Control the Induction of Protective T Helper 1 Responses against Leishmania. *Immunity* **26**, 519–531 (2007).

8. Charmoy, M. *et al.* Neutrophil-derived CCL3 is essential for the rapid recruitment of dendritic cells to the site of Leishmania major inoculation in resistant mice. *PLoS Pathog.* **6**, (2010).

9. Lai, G. N. *et al.* Migratory dermal dendritic cells act as rapid sensors of protozoan parasites. *PLoS Pathog.* **4**, (2008).

10. Sacks, D. & Noben-Trauth, N. The immunology of susceptibility and resistance to Leishmania major in mice. *Nat. Rev. Immunol.* **2**, 845–858 (2002).

11. Scott, P. & Novais, F. O. Cutaneous leishmaniasis: Immune responses in

protection and pathogenesis. *Nat. Rev. Immunol.* **16**, 581–592 (2016).

12.   Olivier, M., Gregory, D. J. & Forget, G. Subversion mechanisms by which
Leishmania parasites can escape the host immune response: A signaling point
of view. *Clin. Microbiol. Rev.* **18**, 293–305 (2005).

13.   Kaye, P. & Scott, P. Leishmaniasis: Complexity at the host-pathogen interface.
Nat. Rev. Microbiol. **9**, 604–615 (2011).

14.   Arenas, R., Torres-Guerrero, E., Quintanilla-Cedillo, M. R. & Ruiz-Esmenjaud,
J. Leishmaniasis: A review. *F1000Research* **6**, 1–15 (2017).

15.   Reithinger, R. Cutaneous leishmaniasis. *Lancet Infect. Dis.* **146**, 581–96 (2007).

16.   Terabe, M. *et al.* CD4+ cells are indispensable for ulcer development in murine
cutaneous leishmaniasis. *Infect. Immun.* (2000) doi:10.1128/IAI.68.8.4574-
4577.2000.

17.   Pereira, L. de O. R. *et al.* Is Leishmania (Viannia) braziliensis parasite load
associated with disease pathogenesis? *Int. J. Infect. Dis.* (2017)
doi:10.1016/j.ijid.2017.01.036.

18.   Gonzalez-Lombana, C. *et al.* IL-17 Mediates Immunopathology in the Absence
of IL-10 Following Leishmania major Infection. *PLoS Pathog.* (2013)
doi:10.1371/journal.ppat.1003243.

19.   Bittencourt, A. L. & Barral, A. Evaluation of the histopathological classifications
of American cutaneous and mucocutaneous leishmaniasis. *Memórias do
Instituto Oswaldo Cruz* vol. 86 51–56 (1991).

20.   Scott, P. & Novais, F. O. Cutaneous leishmaniasis: Immune responses in
protection and pathogenesis. *Nat. Rev. Immunol.* **16**, 581–592 (2016).

21.   Soong, L. *et al.* Role of CD4+ T cells in pathogenesis associated with
Leishmania amazonensis infection. *J. Immunol.* **158**, 5374–83 (1997).

22.   Scorza, B. M., Carvalho, E. M. & Wilson, M. E. Cutaneous manifestations of
human and murine leishmaniasis. *International Journal of Molecular Sciences*
(2017) doi:10.3390/ijms18061296.

23.   Ivens, A. C. *et al.* The genome of the kinetoplastid parasite, Leishmania major.

725   *Science (80-.).* (2005) doi:10.1126/science.1112680.

726 24. Holzer, T. R., McMaster, W. R. & Forney, J. D. Expression profiling by whole-
727   genome interspecies microarray hybridization reveals differential gene
728   expression in procyclic promastigotes, lesion-derived amastigotes, and axenic
729   amastigotes in Leishmania mexicana. *Mol. Biochem. Parasitol.* **146**, 198–218
730   (2006).

731 25. Leifso, K., Cohen-Freue, G., Dogra, N., Murray, A. & McMaster, W. R. Genomic
732   and proteomic expression analysis of Leishmania promastigote and amastigote
733   life stages: The Leishmania genome is constitutively expressed. *Mol. Biochem.*
734   *Parasitol.* **152**, 35–46 (2007).

735 26. McNicoll, F. *et al.* A combined proteomic and transcriptomic approach to the
736   study of stage differentiation in Leishmania infantum. *Proteomics* **6**, 3567–3581
737   (2006).

738 27. Walker, J. *et al.* Identification of developmentally-regulated proteins in
739   Leishmania panamensis by proteome profiling of promastigotes and axenic
740   amastigotes. *Mol. Biochem. Parasitol.* **147**, 64–73 (2006).

741 28. Saxena, A. *et al.* Analysis of the Leishmania donovani transcriptome reveals an
742   ordered progression of transient and permanent changes in gene expression
743   during differentiation. *Mol. Biochem. Parasitol.* **152**, 53–65 (2007).

744 29. Rochette, A. *et al.* Genome-wide gene expression profiling analysis of
745   Leishmania major and Leishmania infantum developmental stages reveals
746   substantial differences between the two species. *BMC Genomics* **9**, 1–26
747   (2008).

748 30. Depledge, D. P. *et al.* Comparative expression profiling of Leishmania:
749   Modulation in gene expression between species and in different host genetic
750   backgrounds. *PLoS Negl. Trop. Dis.* **3**, (2009).

751 31. Chaussabel, D. *et al.* Unique gene expression profiles of human macrophages
752   and dendritic cells to phylogenetically distinct parasites. *Blood* **102**, 672–681
753   (2003).

754 32. Ettinger, N. A. & Wilson, M. E. Macrophage and T-cell gene expression in a

model of early infection with the protozoan Leishmania chagasi. *PLoS Negl. Trop. Dis.* **2**, (2008).

33. Pollack, J. R. A perspective on DNA microarrays in pathology research and practice. in *American Journal of Pathology* (2007). doi:10.2353/ajpath.2007.070342.

34. Akopyants, N. S. *et al.* Expression profiling using random genomic DNA microarrays identifies differentially expressed genes associated with three major developmental stages of the protozoan parasite Leishmania major. *Mol. Biochem. Parasitol.* **136**, 71–86 (2004).

35. Shadab, M. *et al.* RNA-Seq Revealed Expression of Many Novel Genes Associated With Leishmania donovani Persistence and Clearance in the Host Macrophage. *Front. Cell. Infect. Microbiol.* **9**, 17 (2019).

36. Rastrojo, A. *et al.* Analysis by RNA-seq of transcriptomic changes elicited by heat shock in Leishmania major. *Sci. Rep.* **9**, 1–18 (2019).

37. Ruy, P. D. C. *et al.* Comparative transcriptomics in Leishmania braziliensis: disclosing differential gene expression of coding and putative noncoding RNAs across developmental stages. *RNA Biol.* **16**, 639–660 (2019).

38. Christensen, S. M. *et al.* Meta-transcriptome Profiling of the Human-Leishmania braziliensis Cutaneous Lesion. *PLoS Negl. Trop. Dis.* **10**, 1–17 (2016).

39. Amorim, C. F. *et al.* Variable gene expression and parasite load predict treatment outcome in cutaneous leishmaniasis. *Sci. Transl. Med.* **11**, 1–10 (2019).

40. Patino, L. H., Muskus, C. & Ramírez, J. D. Transcriptional responses of Leishmania (Leishmania) amazonensis in the presence of trivalent sodium stibogluconate. *Parasites and Vectors* **12**, 1–15 (2019).

41. Fiebig, M., Kelly, S. & Gluenz, E. Comparative Life Cycle Transcriptomics Revises Leishmania mexicana Genome Annotation and Links a Chromosome Duplication with Parasitism of Vertebrates. *PLoS Pathog.* **11**, 1–28 (2015).

42. Andrade, J. M. *et al.* Comparative transcriptomic analysis of antimony resistant

and susceptible Leishmania infantum lines. *Parasites and Vectors* **13**, 1–15
(2020).

43. Farias Amorim, C. *et al.* Localized skin inflammation during cutaneous
leishmaniasis drives a chronic, systemic ifn-γ signature. *PLoS Negl. Trop. Dis.*
15, 1–19 (2021).

44. Dillon, L. A. L. *et al.* Simultaneous transcriptional profiling of Leishmania major
and its murine macrophage host cell reveals insights into host-pathogen
interactions. *BMC Genomics* **16**, 1–15 (2015).

45. Fernandes, M. C. *et al.* Dual transcriptome profiling of Leishmania-infected
human macrophages reveals distinct reprogramming signatures. *MBio* **7**, 1–16
(2016).

46. Franzén, O., Gan, L. M. & Björkegren, J. L. M. PanglaoDB: A web server for
exploration of mouse and human single-cell RNA sequencing data. *Database*
(2019) doi:10.1093/database/baz046.

47. Walker, D. M. *et al.* Mechanisms of cellular invasion by intracellular parasites.
Cell Mol Life Sci **71**, 1245–1263 (2014).

48. Feijó, D., Tibúrcio, R., Ampuero, M., Brodskyn, C. & Tavares, N. Dendritic cells
and leishmania infection: Adding layers of complexity to a complex disease.
Journal of Immunology Research (2016) doi:10.1155/2016/3967436.

49. Kautz-Neu, K., Schwonberg, K., Fischer, M. R., Schermann, A. I. & von Stebut,
E. Dendritic cells in Leishmania major infections: mechanisms of parasite
uptake, cell activation and evidence for physiological relevance. *Medical*
Microbiology and Immunology (2012) doi:10.1007/s00430-012-0261-2.

50. Liu, D. *et al.* Leishmania major phosphoglycans influence the host early immune
response by modulating dendritic cell functions. *Infect. Immun.* (2009)
doi:10.1128/IAI.01447-08.

51. Woelbing, F. *et al.* Uptake of Leishmania major by dendritic cells is mediated by
Fcγ receptors and facilitates acquisition of protective immunity. *J. Exp. Med.*
(2006) doi:10.1084/jem.20052288.

52.   Axelrod, O., Klaus, S. & Frankenburg, S. Antigen presentation by epidermal
langerhans cells in experimental cutaneous leishmaniasis. *Parasite Immunol.*
(1994) doi:10.1111/j.1365-3024.1994.tb00315.x.

53.   Liu, D. & Uzonna, J. E. The early interaction of Leishmania with macrophages
and dendritic cells and its influence on the host immune response. *Frontiers in*
cellular and infection microbiology (2012) doi:10.3389/fcimb.2012.00083.

54.   Chen, X. *et al.* Transcriptome profiling unveils GAP43 regulates ABC
transporters and EIF2 signaling in colorectal cancer cells. *BMC Cancer* **21**, 1–
10 (2021).

55.   Toribio, R. & Ventoso, I. Inhibition of host translation by virus infection in vivo.
Proc. Natl. Acad. Sci. U. S. A. **107**, 9837–9842 (2010).

56.   Shrestha, N. *et al.* Eukaryotic Initiation Factor 2 (eIF2) signaling regulates
proinflammatory cytokine expression and bacterial invasion. *J. Biol. Chem.*
(2012) doi:10.1074/jbc.M112.375915.

57.   Kim, B. H. *et al.* A family of IFN-γ-inducible 65-kD GTPases protects against
bacterial infection. *Science (80-.).* (2011) doi:10.1126/science.1201711.

58.   Ngo, C. C. & Man, S. M. Mechanisms and functions of guanylate-binding
proteins and related interferon-inducible GTPases: Roles in intracellular lysis of
pathogens. *Cellular Microbiology* (2017) doi:10.1111/cmi.12791.

59.   Yamamoto, M. *et al.* A cluster of interferon-γ-inducible p65 gtpases plays a
critical role in host defense against toxoplasma gondii. *Immunity* (2012)
doi:10.1016/j.immuni.2012.06.009.

60.   Haldar, A. K., Nigam, U., Yamamoto, M., Coers, J. & Goyal, N. Guanylate
binding proteins restrict leishmania donovani growth in nonphagocytic cells
independent of parasitophorous vacuolar targeting. *MBio* (2020)
doi:10.1128/mBio.01464-20.

61.   Lee, S. H. *et al.* M2-like, dermal macrophages are maintained via IL-4/CCL24-
mediated cooperative interaction with eosinophils in cutaneous leishmaniasis.
Sci. Immunol. (2020) doi:10.1126/sciimmunol.aaz4415.

62.   Zaph, C. & Scott, P. Th1 Cell-Mediated Resistance to Cutaneous Infection with
Leishmania major Is Independent of P- and E-Selectins . *J. Immunol.* **171**, 4726–
4732 (2003).

63.   Carlsen, E. D. *et al.* Permissive and protective roles for neutrophils in
leishmaniasis. *Clin. Exp. Immunol.* **182**, 109–118 (2015).

64.   Ribeiro-Gomes, F. L., Peters, N. C., Debrabant, A. & Sacks, D. L. Efficient
capture of infected neutrophils by dendritic cells in the skin inhibits the early anti-
leishmania response. *PLoS Pathog.* **8**, (2012).

65.   Bogdan, C. *et al.* Fibroblasts as host cells in latent leishmaniosis. *J. Exp. Med.*
191, 2121–2129 (2000).

66.   Scorza, B. M. *et al.* Differential Activation of Human Keratinocytes by
Leishmania Species Causing Localized or Disseminated Disease. *J. Invest.*
Dermatol. **137**, 2149–2156 (2017).

67.   Rittig, M. G. & Bogdan, C. Leishmania-host-cell interaction: Complexities and
alternative views. *Parasitol. Today* **16**, 292–297 (2000).

68.   Novais, F. O. *et al.* Genomic Profiling of Human Leishmania braziliensis Lesions
Identifies   Transcriptional   Modules   Associated   with   Cutaneous
Immunopathology. *J. Invest. Dermatol.* **135**, 94–101 (2015).

69.   Santambrogio, L., Berendam, S. J. & Engelhard, V. H. The antigen processing
and presentation machinery in lymphatic endothelial cells. *Frontiers in*
Immunology (2019) doi:10.3389/fimmu.2019.01033.

70.   Tamburini, B. A., Burchill, M. A. & Kedl, R. M. Antigen capture and archiving by
lymphatic endothelial cells following vaccination or viral infection. *Nat. Commun.*
(2014) doi:10.1038/ncomms4989.

71.   Proud, C. G. eIF2 and the control of cell physiology. *Semin. Cell Dev. Biol.*
(2005) doi:10.1016/j.semcdb.2004.11.004.

72.   Jennings, M. D. & Pavitt, G. D. EIF5 has GDI activity necessary for translational
control by eIF2 phosphorylation. *Nature* (2010) doi:10.1038/nature09003.

73.   Liu, Y. *et al.* The role of host eIF2α in viral infection. *Virology Journal* (2020)

doi:10.1186/s12985-020-01362-6.

74.   Jaramillo, M. *et al.* Leishmania repression of host translation through mTOR cleavage is required for parasite survival and infection. *Cell Host Microbe* **9**, 331–341 (2011).

75.   Mahnke, A. *et al.* Hypoxia in leishmania major skin lesions impairs the NO-dependent leishmanicidal activity of macrophages. *J. Invest. Dermatol.* **134**, 2339–2346 (2014).

76.   Schatz, V. *et al.* Myeloid Cell–Derived HIF-1α Promotes Control of Leishmania major . *J. Immunol.* **197**, 4034–4041 (2016).

77.   Weinkopff, T., Roys, H., Bowlin, A. & Scott, P. Leishmania infection induces macrophage vascular endothelial growth factor A production in an ARNT/HIF-dependent manner. *Infect. Immun.* (2019) doi:10.1128/IAI.00088-19.

78.   Titus, R. G., Marchand, M., Boon, T. & Louis, J. A. A limiting dilution assay for quantifying Leishmania major in tissues of infected mice. *Parasite Immunol.* (1985) doi:10.1111/j.1365-3024.1985.tb00098.x.

79.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

80.   Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

81.   Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

82.   Robinson, M. D. & Oshlack, A. A scaling normalization method for differntial expression analysis of RNA-seq data. *Genome Biol.* **11**, 1–9 (2010).

83.   Liu, R. *et al.* Why weight? Modelling sample and observational level variability improves power in RNA-seq analyses. *Nucleic Acids Res.* **43**, (2015).

84.   Benjamini, Y&Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*, **57**, 289–300 (1995).

85. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single
cells. *Nat. Commun.* **8**, 1–12 (2017).

86. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-
cell transcriptomic data across different conditions, technologies, and species.
Nat. Biotechnol. **36**, 411–420 (2018).

87. Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
1902.e21 (2019).

88. Macosko, E.Z., Basu, A., Satija, R., Nemesh, J., Shekhar, K., Goldman, M.,
Tirosh, I., Bialas, A.R., Kamitaki, N., Martersteck, E.M. Highly Parallel Genome-
wide expression Profiling of Individual Cells Using Nanoliter Droplets. *Cell.* 161,
1202-1214 (2017).

89. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. (2018).

90. Finak, G. *et al.* MAST: A flexible statistical framework for assessing
transcriptional changes and characterizing heterogeneity in single-cell RNA
sequencing data. *Genome Biol.* **16**, 1–13 (2015).

Figure 1. *Leishmania* infection is associated with differential regulation of host immune response pathways in vivo. (A) C57BL/6 mice were infected with 2x10⁶ *L. major* metacyclic promastigote parasites intradermally in the ear and lesion development was monitored over time. Data are pooled from 4 experiments (n=30) and shown as +SEM. **(B)** At 4 week post-infection, parasite numbers in the ear were quantified. Data are pooled from 2 experiments (n=20) and shown as +SEM. **(C)** Multi-Dimensional Scaling (MDS) plot showing the gene expression profile between naive and *L. major*-infected ears. **(D)** Volcano plot of all DEGs in naive and infected ears. Red dots represent upregulated DEGs with log2FC >1 and p-value <0.05. Blue represents down-regulated DEGs with log2FC <−1 and p-value <0.05. Only annotated genes are shown in plot. FC, fold-change. **(E-F)** Bulk RNA-Seq analysis indicates DEGs that are highly correlated with signaling pathways. KEGG enrichment analysis of top 20 upregulated (E), and top 20 downregulated (F) pathways enriched among the DEGs between naive and *L. major*-infected ears (pathways selected by significant and FC > 1.5, list includes rank, significance and adjusted average).

Fig. 2

Figure 2: Heat map analysis of host transcriptional responses to *L. major* infection in vivo. Heat maps of DEGs in infected ears compared to naive controls. The DEGs involved in the host immune response pathways by KEGG enrichment analysis whether upregulated **(A,**

938 **B, and C)** or downregulated **(D)** in the infected ears presented as heat maps. Hierarchical

939 clustering of the expression profile was grouped according to functional categories. Heat maps

940 indicate the FC of gene expression in *L. major*-infected ears >2-fold (red) or <2-fold (blue).

Fig. 3

Figure 3. scRNA-Seq reveals cellular heterogeneity including distinct resident and recruited cell types in the skin following *L. major* infection. (A) C57BL/6 mice were infected or not with *L. major* parasites in the ear, and ears were digested to isolate RNA for scRNA-Seq analysis. Schematic of cell isolation, cell processing, capture by droplet-based device. **(B)** Uniform Manifold Approximation and Projection (UMAP) plot revealed cellular heterogeneity with 35 distinct clusters of cells identified and color-coded (both naive and infected groups combined). Seurat's FindClusters function was used to identity each cell cluster and cell type designation is located to the right.

Fig. 4

A

B

Figure 4: Cell type identification and cluster-specific gene expression. (A) Relative

expression of 35 different cell type cluster-specific genes shown as dot plots with two genes

per cluster. Dot size indicates the percentage of cells expressing each gene, the dot color

indicates the expression level, and ordering is performed from low to high expressing cells.

(B) Feature plots of expression distribution for selected cluster-specific genes used to define

the cell types. Expression levels for each marker is color-coded and overlaid onto UMAP plot.

Cells with the highest expression level are colored black.

Figure 5: Presence of *L. major* transcripts in multiple cell types. (A) Differential

expression of *L. major* parasite transcripts in 35 different cell types. Number of cells

containing parasite transcripts was shown in violin plots. **(B)** Table summarizes the

percentages of individual cell that contain *L. major* transcripts.

Fig. 6

Figure 6: scRNA-Seq analysis reveals an enhanced immune cell recruitment to the

inflamed tissue during *L. major* infection. (A) scRNA-Seq UMAP plots of naive and infected

ears showing resident and recruited cell populations/clusters. **(B, D, F)** Representative flow

cytometry plots showing the percentage of CD64+ macrophages (B), Ly6C+ inflammatory

monocytes (D), and Ly6G+ neutrophils (F) of naive and infected ears at 4 week p.i. Cells were

gated on total, live, singlets, CD45+ CD11b+ cells. **(C, E, G)** Cell numbers of CD64+

macrophages (C), inflammatory monocytes (E), neutrophils (G) from naive and infected ears.

(H-J) ECs were gated on total, live, singlets, CD31+ cells. Dermal BECs and LECs were

separated by podoplanin expression during FACS analysis. (H) Representative flow cytometry

dot plots showing the percentages of BECs and LECs from naive and infected ears 4 week

p.i. Corresponding cell numbers of BECs (I) and LECs (J) from naive and infected ears. Data

are representative of at least two independent experiments involving 10-17 mice. Data are

presented as mean +SEM. ***p < 0.005*, ****p < 0.0005*, *****p < 0.0001*, unpaired *t*-test.

Fig. 7

Figure 7: Differentially expressed genes in selected immune cell types during _L. major_

infection. Volcano plot showing the DEGs in macrophages (A), resident macrophages (B),

inflammatory monocyte (C), neutrophils (D), BECs (E), and LECs (F). Colored dots indicate

genes at least 2 (natural log ~0.693) fold increased (red) or decreased (blue) in infected cells

relative to naive cells with an adjusted p-value < 0.05.

Fig. 8

Figure 8: Signaling pathways and molecular networks within individual cell types predicted during *L. major* infection by Ingenuity Pathway Analysis. (A-B) Top 5 differentially regulated canonical pathways and their individual heat maps in macrophages following *L. major* infection. **(C-D)** Top 5 differentially regulated canonical pathways and their individual heat maps in BECs. **(E-F)** Top 5 differentially regulated canonical pathways and their individual heat maps in LECs. The color intensity represents the degree of expression. A red-green color scale was used to reflect the standardized gene expression with red representing high expression and green representing low expression. Cut-off values are adjusted p-value < 0.05.

1000 **Table 1.** List of top 10 KEGG pathways enriched for differentially expressed genes

1001 (DEGs) following *L. major* infection.

Pathway regulation	KEGG enriched pathways	Avg. log-fold change	Adj. p value
Up regulated	Intestinal immune network for IgA production	5.23228	5.08E-07
	Cytokine-cytokine receptor interaction	8.337469	6.97E-06
	Osteoclast differentiation	7.623207	3.31E-05
	Antigen processing and presentation	4.468822	4.21E-05
	Phagosome	6.527853	0.000106
	Chemokine signaling pathway	8.329182	0.000328
	Hematopoietic cell lineage	6.848572	0.000381
	NOD-like receptor signaling pathway	6.638004	0.000744
	Cell adhesion molecules	6.486765	0.000769
	Natural killer cell mediated cytotoxicity	8.022157	0.002282
Down-regulated	Ribosome	-0.696061	0.004705
	Phototransduction	-0.907101	0.007031
	Mineral absorption	-0.992836	0.015538
	ECM-receptor interaction	-0.889139	0.031673
	Oxytocin signaling pathway	-0.948893	0.031673
	Biosynthesis of unsaturated fatty acids	-1.474018	0.042932
	Estrogen signaling pathway	-1.199847	0.055815
	Steroid biosynthesis	-0.908215	0.058351
	Gastric acid secretion	-0.907101	0.072276
	Fatty acid elongation	-0.892548	0.072946

1002

1003

1004

1005 **Table 2.** List of top 10 DEGs enriched in macrophages and resident macrophages
1006 following *L. major* infection.

1007 **A**

1008 <u>**Macrophages**</u>

B

<u>**Resident macrophages**</u>

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
H2-K1	1.203478	2.59E-30
Gm42418	1.013645	3.48E-30
B2m	0.798061	3.33E-28
AW112010	2.205139	1.15E-26
Fth1	1.136905	2.54E-26
Cxcl16	1.778107	5.99E-22
Gbp2	1.141379	1.24E-21
H2-Aa	1.205499	1.62E-20
H2-Ab1	1.264542	9.46E-20
Mmp14	1.784564	6.05E-19
Down regulated		
Rpl32	-1.2065	3.25E-34
Rpl13	-1.23205	2.53E-33
Lyz2	-1.48379	5.03E-32
Rpl18	-1.22252	4.01E-30
S100a4	-1.6294	6.25E-30
Rps18	-1.21541	1.51E-28
Rps14	-1.0183	1.73E-24
Fau	-1.03269	6.54E-23
Ifitm3	-1.30983	8.25E-23
Tpt1	-0.83138	1.52E-22

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
H2-K1	1.441768	6.11E-103
B2m	0.935358	1.97E-81
H2-D1	0.761557	4.47E-53
Stat1	1.637833	1.41E-50
Gbp4	1.457725	1.05E-45
Gbp8	1.043486	2.40E-41
Gbp2	1.43904	3.27E-41
Ly6a	1.716371	2.56E-40
AW112010	1.583531	2.66E-38
Iigp1	1.437197	1.31E-37
Down regulated		
Pf4	-0.93903	7.66E-51
Cd36	-0.8449	1.88E-25
Rpl12	-0.64508	1.92E-25
S100a6	-0.59164	2.65E-24
Rps25	-0.64782	2.88E-24
Rpl10	-0.71332	3.09E-23
Rnf213	0.85734	3.50E-23
Rps12	-0.5271	3.76E-23
Rpl30	-0.60024	4.48E-23
Rps18	-0.56378	9.59E-21

1024

Table 3. List of top 10 DEGs enriched in inflammatory monocytes and neutrophils clusters following *L. major* infection.

A

Inflammatory monocytes

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
Gm42418	0.849004	1.16E-28
Gbp2	2.686188	1.14E-11
Gbp5	2.139579	6.38E-11
Samhd1	1.180168	8.55E-09
Cxcl10	3.062718	2.43E-08
Gbp7	1.761721	1.11E-07
Cxcl9	3.440467	1.79E-07
Gbp3	1.661975	1.62E-06
Iigp1	1.855148	6.54E-06
Nampt	1.57251	1.08E-05
Down regulated		
Fau	-0.67651	8.03E-07
Rpl13	-0.73173	1.00E-06
Rps25	-0.69169	1.18E-06
Lyz2	-0.79057	7.16E-06
Rpl32	-0.66904	1.75E-05
S100a4	-0.85018	0.000375
Tpt1	-0.47099	0.0008
Tmsb4x	-0.51238	0.00088
Rpl9	-0.66084	0.000907
Rpl39	-0.55282	0.001096

B

Neutrophils

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
B2m	1.75856	3.13E-07
Gm42418	0.654377	0.000454
Down regulated		
Sycp2	-1.70554	0.035225

1047   **Table 4.** List of top 10 DEGs enriched in BEC and LEC clusters following *L. major*

1048   infection.

1049

A

1050   <u>BEC</u>

B

<u>LEC</u>

1051

Gene name	Avg. log-fold change	Adj. p value
<u>Upregulated</u>		
B2m	1.330157	3.83E-45
H2-K1	1.134934	4.90E-39
H2-Aa	1.483775	2.02E-27
Cd74	1.574517	1.16E-22
Gbp4	1.127162	1.23E-21
Gbp2	1.442935	3.36E-19
Igtp	1.08015	2.13E-17
H2-Ab1	1.277568	1.37E-16
Ctss	1.372639	3.69E-16
Ubd	1.905291	9.22E-15
<u>Down regulated</u>		
Rpl9	-0.7283	2.97E-16
Cxcl12	-0.84167	6.72E-16
Rpl21	-0.58083	7.74E-14
Rpl13	-0.62302	2.34E-13
Cst3	-0.69511	1.61E-11
Rps14	-0.43341	2.62E-09
Rpl18	-0.563	1.25E-08
Rps15a	-0.51256	1.51E-08
Rpl30	-0.51597	1.72E-08
Rps18	-0.51715	2.05E-08

Gene name	Avg. log-fold change	Adj. p value
<u>Upregulated</u>		
B2m	1.81363722	1.46E-39
H2-K1	1.82143009	6.52E-33
Iigp1	2.05863878	8.72E-22
Gbp4	1.88908626	2.96E-21
H2-D1	1.07551012	1.24E-20
Gbp2	1.89596591	3.54E-18
H2-Q7	1.26192971	3.25E-17
Gbp7	1.31819045	3.86E-17
H2-Aa	1.24429821	3.81E-15
H2-Ab1	1.1254732	1.38E-13
<u>Down regulated</u>		
Rpl21	-0.82606	1.30E-25
Rpl13	-0.86409	6.46E-24
Rpl30	-0.74324	4.82E-19
Rps23	-0.71284	2.88E-18
Rps18	-0.74298	5.32E-18
Rps11	-0.62415	4.34E-14
Rpl9	-0.78012	5.04E-14
Rps25	-0.68713	7.72E-13
Rps4x	-0.54557	1.19E-12
Rpl11	-0.63287	3.41E-12

1070 **Table 5.** List of top 5 upstream regulators identified in macrophages by IPA analysis

1071 for *L. major* infected ears*.*

Predicted activation state	Upstream regulator	Activation z-score	p-value of overlap	No of target molecules in dataset
Cytokines				
Activated	*IFNG*	4.44	9.87E-19	36
	IL4	2.433	1.31E-11	28
	IL13	2.713	8.71E-09	11
	IFNβ1	2.383	0.0000127	10
	TNF	3.364	0.0000372	12
Inhibited	Not detected	-	-	-
Transcription regulators				
Activated	*HIF1A*	2.075	1.62E-08	13
	STAT1	2.396	6.09E-08	12
	CTCF	2.236	5.05E-06	5
	TP73	2.333	5.53E-06	10
	IRF1	2.383	0.0000841	6
Inhibited	*MLXIPL*	-8.062	1.2E-88	65
	MYC	-7.457	3.77E-82	71
	TP53	-2.227	1.77E-19	51
	MYCL	-2.97	7.63E-07	9
	CEBPB	-2.54	0.0000174	16

1072 **Note:** The activation Z-Scores (-8.062 to 4.44) and *p*-values were < 0.05.

1073

1074 **Table 6.** List of top 5 upstream regulators identified in BECs by IPA analysis for *L. major*

1075 infected ears.

Predicted activation state	Upstream regulator	Activation z-score	p-value of overlap	No of target molecules in dataset
Cytokines				
Activated	*IFNG*	3.684	5.19E-14	23
	TNF	2.99	5.79E-10	14
	IL-2	2.449	0.000141	7
	IL-4	2.219	0.00156	10
Inhibited	*IL-10*	-2.01	1.35E-11	14
Transcription regulators				
Activated	*IRF3*	3.065	2.11E-09	10
	STAT-1	2.377	1.37E-08	10
	IRF7	2.573	3.74E-07	7
	MXD1	2.0	0.00129	4
Inhibited	*MLXIPL*	-5.881	3.7E-48	36
	MYC	-5.207	3.54E-43	38
	TRIM24	-2.879	9.58E-12	12
	TP53	-3.877	1.8E-11	28
	SIRT1	-3.357	1.73E-10	15

1076 **Note:** The activation Z-Scores (-5.881 to 3.684) and p-values were < 0.05.

1077

1078 **Table 7.** List of top 5 upstream regulators identified in LECs by IPA analysis for *L. major*

1079 infected ears.

Predicted activation state	Upstream regulator	Activation z-score	p-value of overlap	No of target molecules in dataset
Cytokines				
Activated	*IFNG*	3.644	1.62E-07	14
Inhibited	*IL-10*	-2.219	4.28E-07	9
Transcription regulators				
Activated	*STAT-1*	2.766	3.47E-12	12
	IRF3	3.266	9.08E-12	11
	IRF7	2.795	2.97E-09	8
Inhibited	*MLXIPL*	-6.97	2.47E-80	49
	MYC	-6.714	1.82E-68	49
	SIRT1	-4.338	8.72E-17	19
	TRIM24	-3.564	2.14E-14	13
	MYCL	-2.97	4.91E-10	9

1080 **Note:** The activation Z-Scores (-6.97 to 3.644) and p-values were < 0.05.

1081

1082 **Supplementary Information for**

1083 In vivo reprogramming of murine host immune response genes following *Leishmania major*

1084 infection.

1085

1086 Gopinath Venugopal[a,1], Jordan T. Bird[b,c,1], Charity L. Washam[b,c], Hayden Roys[a], Anne Bowlin[a],

1087 Stephanie D. Byrum[b,c,*], and Tiffany Weinkopff[a,*]

1088

1089 1 G.V. and J.T.B. contributed equally to this work.

1090 * To whom correspondence may be addressed. Email: TSWeinkopff@uams.edu,

1091 SBYRUM@uams.edu

1092

1093

1094 **This PDF file includes:**

1095   Supplementary text

1096   Figures S1 to S5

1097   Tables S1

1098

1099

Fig. S1

A. Cytokine-cytokine receptor interaction

B. Phagosome

C. Toll-like receptor signaling pathway

D. Leukocyte trans-endothelial migration

1100

Figure S1: Heat map analysis showing transcriptional responses from other immune-

related pathways during *L. major* infection in vivo. The DEGs involved in the other host

immune response pathways by KEGG enrichment analysis (A, B, C and D) in the infected

ears compared to naïve mice presented as heat maps. Hierarchical clustering of the

expression profile was grouped according to functional categories. Heat maps indicate the FC

in *L. major* infected ear gene expression >2-fold (red) or <2-fold (blue).

1107

Fig. S2

Figure S2: Differentially expressed genes in selected immune cell types during *L. major* infection. Heat map showing the three highly expressed genes for at least 14 immune cell clusters that were selected along with BECs and LECs. Each column represents a single cell and each row represents an individual gene. Three marker genes per cluster was color-coded and shown on the left. Yellow indicates maximum gene expression and purple indicates no expression in scaled log-normalized unique molecular identifier counts.

Fig. S3. List of top 10 DEGs enriched in DC1 and DC2 clusters between infected vs naïve controls

A

DC1

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
B2m	1.415272	6.77E-22
Gm42418	0.653742	1.85E-20
H2-K1	1.490638	3.31E-18
Samhd1	1.681995	8.18E-14
Ly6a	2.224526	2.03E-11
Ifitm1	2.892357	2.99E-11
Gbp2	2.127184	1.06E-10
Iigp1	1.903031	4.13E-09
Ifitm3	1.338503	6.54E-09
Ptprc	0.760725	1.21E-08
Down regulated		
Rplp1	-0.72544	1.53E-13
S100a10	-1.00579	1.77E-10
Lmna	-1.09686	9.15E-09
Rpl21	-0.6631	9.74E-09
Rpl32	-0.65125	1.65E-08
Tpt1	-0.49815	2.05E-08
Cd207	-1.80546	2.51E-08
Rpl11	-0.65051	4.59E-08
Rplp0	-0.71652	9.95E-08
Emp1	-1.0018	1.03E-07

B

DC2

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
Gm42418	0.953001	7.12E-11
Stat1	2.397145	8.79E-11
AA467197	2.638065	2.18E-09
Syngr2	0.881274	4.46E-07
Gapdh	0.499088	4.63E-05
Fabp5	1.577459	0.000217
Map4k4	1.109246	0.000326
Ramp3	1.877497	0.000514
Sub1	0.825485	0.001213
Gbp2	2.000493	0.001554
Down regulated		
Apol7c	-2.85556	2.94E-17
Rpsa	-1.16891	3.84E-11
H2-M2	-1.8339	9.12E-11
Adam23	-0.68256	7.60E-09
Ankrd33b	-1.40031	5.77E-08
Cetn3	-0.90103	1.97E-07
Ccl17	-2.03132	4.70E-07
Nudt17	-1.0066	6.61E-07
Rpl30	-1.14097	8.43E-07
Zc3h12c	-0.95491	5.56E-06

Figure S3: Differentially expressed genes in DCs during *L. major* infection. Volcano plot showing the DEGs in dendritic cells (DC1 and DC2) and list includes the top DEGs enriched in DCs following *L. major* infection. Colored dots indicate genes at least 2 (natural log ~0.693) fold increased (red) or decreased (blue) in infected cells relative to naïve cells with an adjusted p-value < 0.05.

Fig. S4. List of top 10 DEGs enriched in CD4⁺ and CD8⁺ T cell clusters between infected vs naïve controls

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
B2m	1.13972	4.63E-08
Satb1	2.352451	8.61E-08
Gm42418	0.316364	3.95E-06
Gimap6	1.674671	0.01874
Down regulated		
S100a4	-1.43762	1.36E-08
Crip1	-1.07576	4.72E-05
Rpl21	-0.7473	0.001303
Tmem156	-1.16234	0.001376
Capg	-1.33751	0.002164
Rpl13	-0.70277	0.004808
Rps18	-0.74677	0.005852
Tox	-1.12514	0.006776
Rpl32	-0.64672	0.00741
Mprip	-1.22776	0.010799

Gene name	Avg. log-fold change	Adj. p value
Upregulated		
Gm42418	1.14852	2.20E-05
Down regulated		
Card10	-1.23539	0.013676

Figure S4: Differentially expressed genes in CD4⁺ and CD8⁺ T cells during _L. major_ infection. Volcano plot showing the DEGs in CD4⁺ and CD8⁺ T cells and list includes the top DEGs enriched in CD4⁺ and CD8⁺ T cells following _L. major_ infection. Colored dots indicate genes at least 2 (natural log ~0.693) fold increased (red) or decreased (blue) in infected cells relative to naïve cells with an adjusted p-value < 0.05.

Fig. S5. IPA analysis revealed the role of mTOR signaling pathway in other cell types such as dendritic cells, inflammatory monocytes, CD4⁺ T cell following *L. major* infection.

1127

1128   Figure S5: IPA predicted the role of mTOR signaling in other immune cell types during

1129   *L. major* infection by. (A-C) Differentially regulated canonical pathways in DCs (A),

1130   inflammatory monocytes (B), CD4⁺ T cells following *L. major* infection.

1131

1132

1133

1134

1135

1136
1137

1138 **Table S1.** List of top 20 KEGG disease pathways enriched for differentially expressed genes

1139 (DEGs).

KEGG disease pathway	Adj. p value	Avg. log-fold. change	direction
Staphylococcus aureus infection	1.34E-08	6.60312904	Up
Autoimmune thyroid disease	3.79E-08	6.020025599	Up
Graft-versus-host disease	2.19E-07	5.439582695	Up
Allograft rejection	2.76E-07	5.439582695	Up
Primary immunodeficiency	7.92E-07	8.419604409	Up
Type I diabetes mellitus	4.25E-06	5.439582695	Up
Osteoclast differentiation	3.31E-05	8.133401516	Up
Leishmaniasis	3.31E-05	6.701497553	Up
Systemic lupus erythematosus	0.00016	5.606193223	Up
Rheumatoid arthritis	0.00047	6.580971428	Up
Viral myocarditis	0.001	6.082823244	Up
Tuberculosis	0.0023	6.136724912	Up
Asthma	0.0023	2.73570696	Up
Inflammatory bowel disease (IBD)	0.0034	5.688039923	Up
Prion diseases	0.00458	-0.874425121	Down
Herpes simplex infection	0.00472	6.19895637	Up
Measles	0.00511	7.879610143	Up
Salivary secretion	0.00703	6.634176923	Up
Pertussis	0.00703	5.299789946	Up
Toxoplasmosis	0.00703	6.89131138	Up

1140