Experimental demonstration of prenatal programming of mitochondrial

aerobic metabolism lasting until adulthood

Antoine Stier [a,b,c], Pat Monaghan [a] & Neil B. Metcalfe [a]

[a] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

Glasgow, UK

[b] Department of Biology, University of Turku, Turku, Finland

[c] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622,

Villeurbanne, France

*Corresponding author: antoine.stier@gmail.com / amstie@utu.fi

Abstract

It is increasingly being postulated that among-individual variation in mitochondrial function underlies variation in individual performance (*e.g.* growth rate) and state of health. It has been suggested (but not adequately tested) that environmental conditions experienced before birth have been suggested to programme postnatal mitochondrial function, with persistent effects potentially lasting into adulthood. We tested this hypothesis in an avian model by experimentally manipulating prenatal conditions (incubation temperature and stability), then measuring mitochondrial aerobic metabolism in blood cells from the same individuals during the middle of the growth period and at adulthood. Mitochondrial aerobic metabolism changed markedly across life stages, and part of these age-related changes were influenced by the prenatal temperature conditions. A high incubation temperature induced a consistent and long-lasting increase in mitochondrial aerobic metabolism. Postnatal mitochondrial aerobic metabolism was positively associated with oxidative damage on DNA but not telomere length. While we detected significant within-individual consistency in mitochondrial aerobic metabolism across life-stages, the prenatal temperature regime only accounted for a relatively small proportion (<20%) of the consistent among-individual differences we observed. Our results demonstrate that prenatal conditions can program consistent and long-lasting differences in mitochondrial function, which could potentially underlie among-individual variation in performance and health state.

Keywords: developmental programming, telomeres bioenergetics, mitochondria, oxidative stress, Japanese quail.

Introduction

Mitochondria generate more than 90% of the cellular energy routinely required by animal cells in the form of adenosine triphosphate (ATP), produced through oxidative phosphorylation [1]. Mitochondria are also a major source of reactive oxygen species (ROS), which are likely to be involved in the ageing process if produced in excess of the antioxidant capacity [2]. Consequently, variations in mitochondrial function are likely to play an important role in shaping the life and death of individuals, and to underlie phenotypic variation observed both within and between species [3–6]. Importantly, while mitochondrial function is a very plastic trait (*i.e.* influenced by tissue, temperature, oxygen/food availability or reproductive stage; [7–10]), it has been shown in both human and animal models that adult individuals exhibit some degree of within-individual consistency in mitochondrial traits through time [10,11]. Understanding the origin of such consistent among-individual differences in mitochondrial function (*e.g.* genetics, early-life programming, permanent environmental effects) is key to understand the role of mitochondria in shaping individual performance and health state [6,12].

Scientific evidence from both the biomedical and the ecological fields of research suggest that conditions experienced during early-life have major and persistent effects on adult phenotype and physiology [13,14], a phenomenon termed developmental or early-life programming. For instance, accelerated postnatal growth has been associated with increased metabolic rate at adulthood and reduced lifespan in animal models [15,16], as well as with increased risks of developing various age-related pathologies [17,18]. The underlying molecular and physiological mechanisms are probably numerous, but mitochondrial dysfunction has emerged as one potential candidate linking early-life conditions to both immediate and delayed effects on phenotype and health state[12,14,19,20]. However, much remains to be discovered about the importance of early-

life conditions in determining consistent and long-lasting differences in mitochondrial function

between individuals.

Investigating the long-term programming of physiological function is more amenable

through longitudinal studies (*i.e.* measuring the same individuals over time), especially to avoid

issues linked to the selective disappearance of specific phenotypes [21]. In recent years, measuring

mitochondrial function from blood cells has emerged as an opportunity to conduct longitudinal

studies of mitochondrial biology both in human and animal models [10,11,22].

To the best of our knowledge, only cross-sectional studies have been conducted to date in

the context of early-life effects on mitochondrial function [12]. These studies suggest for instance

that maternal food restriction during pregnancy can alter mitochondrial bioenergetics [12,23]. This

has been framed overall as the 'developmental programming of mitochondrial biology' hypothesis

[12]. However, none of these studies show an unequivocal direct and persistent effect of prenatal

conditions on mitochondrial function. For instance, early-life dietary manipulations usually result in

alterations of body mass (and body fat content) during growth and at adulthood, which could by

itself alter mitochondrial function independently of prenatal conditions. Additionally,

measurements of mitochondrial function are mostly conducted at a single time point, often months

or years after birth, so preventing evaluation of the effect of manipulations on early-life

mitochondrial function, and the potential effects on age-related variation in mitochondrial function.

Oviparous species offer the opportunity to conduct experiments on the direct effects of

prenatal conditions on mitochondrial function, since the developmental conditions of the embryo

can be altered through direct manipulations of the eggs. For instance, it has been shown that higher

incubation temperature of chicken eggs led to higher oxidative capacity (state 3 / *OXPHOS*) and

mitochondrial proton leak (state 4 / *LEAK*) in developing bird embryos [24]. However, this study did

not control for treatment-induced differences in embryo developmental stage and did not assess

the persistence of the effects on mitochondrial function postnatally and over the long term.

Here we investigate the potential prenatal programming of postnatal mitochondrial

function from early-life to adulthood by using manipulations of incubation temperature and stability

in an avian model [25]. Both the absolute incubation temperature and the stability of that

temperature could constrain pre- and postnatal development in non-adaptive ways (*e.g.* lower

and/or unstable incubation temperatures have been shown to slow postnatal growth and increase

metabolic rate; [26,27]). However, they can also convey environmental information that embryos

might use to adjust the postnatal phenotype to suit anticipated environmental conditions (*e.g.* a

higher incubation temperature limits the deleterious impact of post-hatching exposure to heat

stress [28]). Mitochondrial function has been shown to exhibit changes according to age/life-stage

in both animal models and humans [29–31]. Therefore, a second aim was to characterize within-

individual changes in mitochondrial function between the peak of the growth phase and early-

adulthood, and to evaluate potential effects of the prenatal environment on such age-related

changes. Since it has been shown that individuals with more efficient mitochondria (*i.e.* producing

more ATP per unit of O_2 consumed) could grow faster [32], we hypothesize that mitochondrial

efficiency could be maximized during early life to support the costly process of growth. Since we

previously showed that our incubation temperature manipulation led to differences in ageing

biomarkers (*i.e.* DNA damage and telomere length [33]), we also tested for potential relationships

between mitochondrial aerobic metabolism and those ageing markers. Finally, our last objective

was to characterize the extent to which among-individual differences in mitochondrial function are

consistent over time (*i.e.* within-individual consistency), but also the extent to which such consistent

among-individual differences could originate from variation in the prenatal environment. This is

conceptually important for evaluating the scope for permanent environmental effects linked to

early-life programming and heritable variation in mitochondrial aerobic metabolism.

Material and Methods

Experimental design

All procedures were conducted (as previously described for the same experiment in [33]) in 2016 in accordance with UK regulations under the Home Office Project Licence 70/8335 granted to PM and the Home Office Personal Licence ICB1D39E7 granted to AS. We used Japanese quail (*Coturnix japonica*) as a model since they are precocial birds and can be reared successfully without the parent being present, therefore avoiding any confounding effect linked to variation in parental care; in addition, they reach sexual maturity and therefore adulthood very quickly (*ca.* 50 days [34]) making it possible to examine long term effects spanning life history stages within a few months. Japanese quail eggs were bought from Moonridge Farm (Devon, UK) and delivered within 48 hours after collection. The identity of the parents was unfortunately unknown, but given that all the eggs were laid on the same day, it is very unlikely that our study population contained full siblings. We used 164 eggs from which 107 chicks hatched, 6 died in the first 5 days and were excluded from the study, and 25 birds were not used for mitochondrial function assessment due to logistical constraints, giving a sample size of 76 birds.

Eggs were incubated at 3 constant temperatures as described in more detail in [33]: high (H) = 38.4°C, medium (M) = 37.7°C and low (L) = 37.0°C. Additionally a fourth group was incubated under 'unstable' (U) temperature conditions, with an incubation temperature of 37.7°C but five incubation recesses of 30min during the day, leading to a daily average of 37.0°C, similar to the L group (*i.e.* its matched control for developmental speed [33]). Experimental temperature conditions were chosen based on existing literature [35] and pilot experiments, so as to maximize differences in developmental speed and metabolism while minimizing the risks of having differences in hatching success between groups (*i.e.* to avoid

the selective disappearance of embryos in some groups [33]). The conditions for the U group

were chosen based on the natural incubation recesses occurring when females leave the nest

to forage [36]. Temperature and humidity were checked daily within each incubator using a

digital thermo-hygrometer (R-Com DigiLog3) placed in the centre of the incubator, and did not

deviate by more than 0.2°C and 5% from their target values. Our prenatal experimental

treatments affected developmental speed in the predicted direction (*i.e.* lower temperature

leading to slower prenatal growth and metabolism, higher temperature leading to faster

development and metabolism, and unstable temperature treatment leading to growth similar

to low temperature), while having no significant effects on hatching success or mass at

hatching and during growth [33].

Animal husbandry rooms for post-hatching rearing were maintained at 21°C on a 14L:10D

cycle throughout the experiment. After hatching (Day 0) chicks were placed for 24h in a larger

incubator set at 37°C, before being placed into their respective enclosure within each room at Day

1 where an additional heat source was provided (Brinsea Comfort brooder 40, 42W), as well as *ad*

libitum food (Heygates starter crumbs, 22% protein) and water. The additional heat source was

removed at day 15 and chick food was switched to adult pellets (Heygates quail and partridge

pellets, 16% protein) at the same time (*i.e.* 5 days before the first blood sampling). Chicks were

maintained in mixed-sex groups until Day 25 when they can be sexed morphologically. Females

were then kept in groups in enclosures and males were placed by pairs in 0.8m^2 cages to avoid

female exhaustion due to male harassment and to limit male-male conflicts.

Sampling procedures

We used blood cells [22] to measure mitochondrial function of the same individuals at

both Day 20 (peak of the growth phase, see [33]) and Day 60 (early-adulthood). Blood samples

(*ca.* 200µL) were collected at Day 20 and Day 60 by venipuncture of the wing vein with a 26*G*

needle and collection using heparinised capillaries. Blood was centrifuged 10min at 3000*g* and 4°C

to separate plasma from packed blood cells, and 50µL of blood cells (*ca.* 150 to 200.10^6 cells) were

immediately re-suspended in 1mL of Mir05 buffer (0.5 mM EGTA, 3 mM $MgCl_2$, 60 mM K-

lactobionate, 20 mM taurine, 10 mM KH_2PO_4, 20 mM Hepes, 110 mM sucrose, free fatty acid

bovine serum albumin (1 g/L), pH 7.1) and kept on ice until analysis (< 1.5h after blood collection).

High-resolution respirometry analysis of mitochondrial function in intact blood cells

We followed the protocol described in detail for king penguins [22], with some minor

modifications. Briefly, samples were washed before the start of the mitochondrial

measurements by centrifugating the tubes to pellet the blood cells and discarding the

supernatant. Blood cells were then re-suspended in 1mL of respiratory buffer MiR05 pre-

equilibrated at 40.0°C in the chamber of the Oxygraph-2k (Oroboros Instruments, Innsbruck,

Austria), and transferred to the chamber already containing another 1mL of Mir05. After

closing the chamber, baseline O_2 consumption was recorded, followed by the inhibition of

ATP-synthesis using oligomycin (Oligo: 2.5µM), then by the stimulation of maximal uncoupled

respiration using a sequential titration of carbonyl cyanide-p-

trifluoromethoxyphenylhydrazone (FCCP: 0.5µM per step) until maximal stimulation was

reached, and finally by the inhibition of mitochondrial respiration using antimycin A (AA:

2.5µM). Mitochondrial responses of blood cells to this chemical titration are presented in Fig

S1A. We then calculated *ROUTINE* respiration (*i.e.* endogenous cellular respiration = *baseline*

- AA), *OXPHOS* respiration (*i.e.* O_2 consumption linked to ATP synthesis = *baseline - Oligo*),

LEAK respiration (*i.e.* O_2 consumption mostly linked to mitochondrial proton leak = *Oligo - AA*)

and *ETS* respiration (*i.e.* maximal O_2 consumption of the electron transport system = *FCCP -*

AA). We also calculated two mitochondrial *flux control ratios* (FCRs), namely the OXPHOS

coupling efficiency (*OxCE* calculated as *OXPHOS*/*ROUTINE*) indicating the proportion of

endogenous respiration being linked to ATP synthesis, and an index of mitochondrial capacity

usage (FCR$_{R/ETS}$ calculated as *ROUTINE/ETS*) indicating the proportion of maximal capacity

being used under endogenous cellular conditions [22]. To account for potential differences in

cell quantity between samples, we quantified the protein content of the cell suspension using

a BCA protein assay (ThermoScientific) following [22].

Some studies have shown that mitochondrial function measured in blood cells is

correlated to some extent to mitochondrial function in other tissues such as kidneys, heart,

skeletal muscles and brain (*e.g.* [22,37–39], while some other studies did not find any

significant relationship (*e.g.* [40]). Therefore, to evaluate the relevance of using blood cell

mitochondrial function in Japanese quails, we compared mitochondrial function between

blood cells and brain samples (see ESM) in a subsample of adult females (N = 22); and found

a moderate but overall significant correlation between these two tissues (meta-analytic *r* =

0.44, 95% C.I. = [0.27;0.59], p < 0.001; see ESM Fig. S2).

Telomere length and DNA damage

Telomere length and oxidative damage on DNA were measured for a proportion of

the individuals used here (n = 45 at day 20, n = 40 day 60) as part of another study [33]. Briefly,

DNA extracted from blood cells was used to measure absolute telomere length (using *in-gel*

terminal restriction fragment, TRF) and 8-hydroxydeoxyguanosine (8-OHdG), one of the

predominant forms of free radical-induced oxidative lesions in DNA (using a ELISA assay) [33].

Data analysis

We used generalized estimated equations (GEEs) in *SPSS* 24.0 to investigate the effects of

age (*i.e.* repeated effect), prenatal treatment (*i.e.* low, medium, high or unstable incubation

temperature), sex and their interactions on mitochondrial parameters, with the associated post-

hoc tests (non-significant interactions were removed from the final models except the focal *age*

x treatment interaction). We included the cellular protein content as a covariate in the models

to account for potential variations in cell quantity between samples, and the time of day to

account for potential circadian variations in mitochondrial function [41]. Relationships between

telomere length or DNA damage and mitochondrial traits were explored using GEEs, with either

telomere length or DNA damage as the dependent variable, age and treatment plus their

interaction as fixed effects, and one mitochondrial trait at a time (due to strong collinearity

between mitochondrial traits) as a covariate. Mitochondrial respiration rates were expressed

per quantity of blood cell protein for these analyses, and all continuous variables were z-

transformed to provide comparable estimates. To analyse within-individual consistency in

mitochondrial traits, we used the *RptR* package [42] in *R* 3.4.2, including bird identity as a

random (*i.e.* focal) term, as well as age and cellular protein content as fixed effects to account

for changes in mitochondrial traits with age and quantity of cells per sample. We re-ran the

same analyses while including experimental treatment group as an additional random term to

investigate the proportion of within-individual consistency being attributed to our prenatal

treatments. Due to some failed laboratory assays (*e.g.* due to residual inhibition of

mitochondrial respiration in the Oroboros chambers), the final dataset includes n = 136

mitochondrial [measurements from N = 76 individuals (L = 20, M = 20, H = 16 and U = 20)]. A

few (n = 8) *ETS* values being non-biological (*ETS < ROUTINE*) were removed from the final

dataset.

Results

Effect of age and prenatal temperature regime on mitochondrial traits

Fig 1: Effects of age and prenatal temperature regime on blood cells mitochondrial traits: (A) *ROUTINE* **respiration: endogenous mitochondrial O₂ consumption, (B)** *OXPHOS* **respiration: mitochondrial O₂ consumption being linked to ATP synthesis (C)** *LEAK* **respiration: mitochondrial O₂ consumption being mostly linked to mitochondrial proton leak, (D)** *ETS* **respiration: maximal mitochondrial O₂ consumption induced by mitochondrial uncoupling, (E) OXPHOS coupling efficiency (***OxCE***): proportion of endogenous respiration devoted to ATP synthesis and (F) mitochondrial capacity usage (FCR***ROUTINE/ETS***): proportion of maximal mitochondrial respiration being used under endogenous cellular conditions.** Details of statistical tests are given in Tables S1 and S2, means are presented ± SE, letters indicate significant differences between groups according to GEE post-hoc tests, and within-group age effects are presented as * = $p < 0.05$, ** = $p < 0.01$ and *** = $p < 0.001$. Statistical models for A-D included blood cells protein content as a covariate to account for potential variations in blood cell quantity between samples.

Endogenous mitochondrial respiration (*ROUTINE*) was significantly affected by the age

of individuals and the prenatal experimental treatment, as well as by their interaction (Fig 1A,

Table S1A). Specifically, *ROUTINE* decreased overall with age, but in a treatment-specific

manner (H and L decreasing significantly, while M and U only showed a non-significant

decrease). Chicks from the high temperature group had a significantly higher *ROUTINE* at Day

20 than other groups. At adulthood (Day 60), birds from the high temperature and unstable

groups had higher *ROUTINE* than low temperature birds, while birds in the medium

temperature group were intermediate (Fig. 1A). Mitochondrial *OXPHOS* respiration decreased

significantly with the age of individuals (Fig 1B, Table S1B). It also significantly differed

between treatment groups: birds from the high temperature group had the higher *OXPHOS*,

followed by unstable and medium temperature ones, and finally by low temperature ones

(see Fig 4B for significance). *LEAK* respiration increased significantly with age, but was not

significantly affected by treatment (Fig 1C, Table S1C). *ETS* respiration increased overall with

age, but in a treatment-specific manner (significant age x treatment interaction, with M and

U increasing significantly while H and L only exhibited a non-significant increase; Fig 1D, Table

S1D). Chicks from the high temperature group had a significantly higher *ETS* at Day 20 than

those from the medium temperature and unstable groups, while low temperature ones had

an intermediate phenotype (Fig. 1D). At adulthood (Day 60), birds from the low temperature

group had lower *ETS* respiration than all other groups (no significant differences were

observed between H, M and U groups).

Mitochondrial OXPHOS coupling efficiency (*i.e. OxCE*) significantly decreased with age,

but was not significantly affected by the prenatal treatment, either as a main factor or in

interaction with age (Fig 1E, Table S2A). However, this parameter exhibited a sex difference,

with females having slightly less efficient mitochondria than males (Table S2A, LS-mean ± SE:

females = 0.836 ± 0.007 *vs.* males = 0.852 ± 0.005, p = 0.039). The proportion of maximal

respiration being used under endogenous cellular conditions (*i.e.* $FCR_{R/ETS}$) significantly

decreased with age, but was not significantly affected by the prenatal treatment, either as a

main factor or in interaction with age (Fig 1F, Table S2B).

Relationships between telomere length/DNA damage and mitochondrial traits

Mitochondrial respiration rates were positively related to DNA damage levels (Fig. 2A),

significantly so for *ROUTINE* (Wald χ^2 = 4.06, *p* = 0.044) and *LEAK* (Wald χ^2 = 6.27, *p* = 0.012),

but not significantly so for *OXPHOS* (Wald χ^2 = 2.15, *p* = 0.14) and *ETS* (Wald χ^2 = 3.29, *p* =

0.07). *OxCE* (Wald χ^2 = 2.40, *p* = 0.12) and *$FCR_{R/ETS}$* (Wald χ^2 = 0.53, *p* = 0.47) were not

significantly related to DNA damage levels (Fig 2A). Mitochondrial traits were not significantly

related to telomere length (all Wald χ^2 < 0.14 and *p* > 0.71; Fig 2B).

295   <u>**Fig 2**</u>**: Relationships between mitochondrial traits and (A) oxidative damage to DNA (8-OHdG), (B)**
telomere length. Effects are reported as standardized estimates from GEE models (based on z-
296   transformed values) with 95% confidence intervals. Significant effects are shown in bold and details on
statistics are provided in the main text.

Within-individual consistency in mitochondrial phenotype

All mitochondrial respiration rates exhibited a significant within-individual consistency through time between the peak of the growth phase (Day 20) and early adulthood (Day 60) (Fig 3A; *ROUTINE*: R = 0.58, p < 0.001; *OXPHOS*: R = 0.53, p < 0.001; *LEAK*: R = 0.23, p = 0.045; *ETS*: R = 0.48, p < 0.001). While mitochondrial OXPHOS coupling efficiency was not significantly repeatable (*OxCE*: R = 0.12, p = 0.17), the mitochondrial capacity usage was (*FCR$_{R/ETS}$*: R = 0.42, p = 0.001; Fig. 3A). When looking at the variance in mitochondrial traits being explained respectively by bird identity (*i.e.* intrinsic variation) and by incubation temperature (*i.e.* extrinsic variation linked to our treatment), the contribution of the latter was relatively modest (Fig 3B; all < 8%) compared to bird identity (all > 21%, except *OxCE*). Experimental treatment thereby accounted for 7.9-17.5% of the within-individual consistency in mitochondrial respiration rates, and less than 2% of the within-individual consistency in mitochondrial flux ratios.

Fig 3: **(A) Within-individual consistency of mitochondrial respiration rates and flux control ratios measured in intact blood cells at day 20 and 60 in Japanese quail.** Age-adjusted repeatability (*i.e.* consistency) estimates R are presented along with their 95% C.I. The statistical model also contained cellular protein content as a covariate to control for potential variations linked to differences in cell quantity between samples. **(B) Respective variance in mitochondrial traits being explained by bird identity (*i.e.* intrinsic variation), our experimental treatment (incubation temperature, *i.e.* extrinsic variation) and other non-investigated factors as well as measurement error (*i.e.* residual)**.

Discussion

Our results demonstrate that variation in the prenatal environment can influence

mitochondrial function from post-natal life into adulthood, with a high incubation

temperature leading to higher postnatal mitochondrial aerobic metabolism. We found

marked differences in mitochondrial function between the peak of the postnatal growth

phase and early adulthood, but the pattern of this age-related change was partly influenced

by the prenatal temperature. Overall, our results suggest that prenatal conditions can affect

how mitochondria work, but also how mitochondrial function changes with age. The

persistence of pre-natal temperature effects on mitochondrial function into adulthood

suggests that these are long lasting. Higher mitochondrial aerobic metabolism was overall

associated with higher levels of oxidative damage on DNA, but not with shorter telomeres.

We found significant within-individual consistency of mitochondrial respiration rates across

life-stages, with individuals showing consistent relatively high or low values through time.

Age-related variation in mitochondrial function

Mitochondrial parameters differed markedly between the peak of the growth phase

and early adulthood (6 parameters out of 6). The overall pattern suggests that oxidative

phosphorylation decreases with age, while both maximal mitochondrial capacity and proton

leak increase, leading to a higher mitochondrial coupling efficiency but also to a more intense

utilization of the total mitochondrial capacity during growth than at adulthood. These results

are in line with our hypothesis that mitochondrial efficiency should be maximized in early-life

to sustain the growth process, and also indicate that mitochondrial maximal capacity is more

intensively used during growth. Such effects could potentially be mediated by the known

impact of growth hormone on mitochondrial function [43], although this remains to be

properly tested.

Prenatal programming of mitochondrial function by incubation temperature and stability

Our results clearly show that the prenatal environment can affect mitochondrial

respiration rates (*i.e.* endogenous cellular respiration, oxidative phosphorylation and maximal

mitochondrial capacity) in the long-term since we found effects of incubation temperature

regime both during postnatal growth and at adulthood. Specifically, it seems that a high

incubation temperature increased subsequent mitochondrial metabolism at both life-stages

(*i.e.* persistent effect on *ROUTINE*, *OXPHOS* and *ETS* compared to low temperature).

Mitochondrial aerobic metabolism of individuals in the unstable incubation treatment did not

differ from those in the medium temperature group (which shared the same incubation

temperature 90% of the time [33]), but was in some cases (*i.e. OXPHOS*, *ROUTINE* and *ETS* at

day 60) higher than the low temperature group (*i.e.* the group matched for daily average

incubation temperature and developmental speed [33]). This suggests that the temperature

experienced during the majority of the prenatal development is a more likely driver of

mitochondrial programming than average incubation temperature or developmental speed.

While instability in incubation temperature is sufficient to slow down embryo growth and elicit

a prenatal increase in glucocorticoid levels [33], it does not seem sufficient to affect

mitochondrial aerobic metabolism despite the documented effects of glucocorticoids on

mitochondrial biology [44]. Our results demonstrate that prenatal environmental conditions

can have relatively immediate effects (*i.e.* during postnatal growth) as suggested by previous

correlative studies in mammals [45,46], and more importantly persistent effects lasting from

early post-natal life to adulthood. Importantly, unlike previous studies in mammals and

reptiles [47], our results cannot be biased by differences in body mass between experimental

groups at the time of mitochondrial measurement [33]. Mitochondrial flux control ratios were

not affected by the prenatal treatments, suggesting that the differences we observe in

respiration rates between groups were relatively consistent across the different mitochondrial

respiration rates we measured and might be linked to changes in mitochondrial density. The

mechanism(s) by which incubation temperature programmes postnatal mitochondrial aerobic

metabolism on the long-term remain to be investigated, but modifications of the epigenome

could be a key candidate mechanism [48].

Interestingly, age-related changes in mitochondrial function were also partly

influenced by the prenatal environment (*i.e.* age-related decrease in *ROUTINE* for H and L

groups only, as well as age-related increase in *ETS* for M and U groups only), suggesting that

the prenatal environment does not only affect how mitochondria work postnatally, but also

the way mitochondrial aerobic metabolism changes with age. Considering the importance of

mitochondria in the ageing process [29,49], such effects of the prenatal environment on both

mitochondrial aerobic metabolism and its age-related changes could have potential

consequences in influencing ageing trajectories. We partly tested this hypothesis by

investigating the relationships between two biomarkers of ageing and mitochondrial traits,

and found only mixed evidence since high mitochondrial aerobic metabolism was associated

with higher levels of DNA damage, but not with shorter telomeres.

Although we found clear programming effects of mitochondrial aerobic metabolism by

prenatal environmental conditions, we have so far no information about their potential

adaptive or maladaptive value. Further studies investigating the adaptive role of such variation

in mitochondrial function (*e.g.* by testing individual performance under contrasted postnatal

environmental conditions) will be needed to determine the potential adaptive value of

prenatal programming through incubation temperature and stability. It is possible that the

increased aerobic metabolism programmed by high prenatal temperature provides

immediate benefits (*e.g.* higher competitivity and reproductive success) but at the expense of

long-term performance and survival (*i.e.* faster 'pace of life') as suggested by the positive

relationship found between mitochondrial respiration rates and oxidative damage on DNA.

Our results therefore pave the way for further research on the implication of mitochondrial

aerobic metabolism in the 'pace of life' syndrome [50].

Within-individual consistency in mitochondrial function and relative importance of prenatal

conditions

Mitochondrial respiration rates and flux control ratios exhibited a significant within-

individual consistency over time (at the exception of *OxCE*), despite being measured at two

different life stages (peak of growth *vs.* early-adulthood) over a period when there are marked

changes in mitochondrial aerobic metabolism (see above). To the best of our knowledge, this

is the third demonstration that consistent among-individual differences in mitochondrial

aerobic metabolism exist (see [10] in wild adult passerine birds, and [11] in adult humans), but

the first to be conducted under well-controlled environmental conditions (*i.e.* excluding bias

linked to consistent individual differences in environmental conditions) and across life-stages.

These findings of a significant within-individual consistency in mitochondrial function over

time have important implications for the possibility for early-life conditions to programme

mitochondrial function over the life course of individuals [12]. Our experimental treatment

only accounted for less than 20% of the within-individual consistency observed in

mitochondrial respiration rates, meaning that more than 80% of the observed within-

individual consistency must be explained by other prenatal factors (*e.g.* maternal transfer of

nutrients, hormones) and/or by genetic differences between individuals. To the best of our

knowledge, there is no information published about the heritability of mitochondrial function,

but mtDNA copy number, one proxy of mitochondrial density, has been shown to be

significantly heritable (h^2 = 33%) in humans [51]. Therefore, estimating the relative importance

of genetic *vs.* environmental drivers of mitochondrial function appears now fundamental to

evaluate the magnitude to which early-life environment could programme mitochondrial

function and its potential downstream effects such as disease risk and individual performance.

Competing interests

We declare having no competing interests

Data availability

Datasets used in this manuscript are available at: https://doi.org/10.6084/m9.figshare.16708483.v1

Author's contribution

AS designed the study, conducted the experimental work, data analysis and wrote the

manuscript. NBM and PM had input on study design and data analysis, and commented on

the manuscript.

Acknowledgements

We are grateful to two anonymous reviewers for their constructive feedback on an earlier

draft of this manuscript, to Karine Salin for help in setting up mitochondrial analysis of brain

samples, to Pierre Bize for kindly providing access to his O2k-oxygraph, to Norith Eckbo for the

graphic design of Fig. S2, to Franklin Lo, Becky Shaw and Katie Byrne for their help in collecting

blood samples, and to Graham Law and his team for taking care of animal husbandry. Finally,

AS is grateful to the crew of the Marion Dufresnes and the French Polar Institute (IPEV) for

hosting him from 56 to 21° South while writing the first draft of this manuscript. The project

was funded by a Marie Sklodowska-Curie Postdoctoral Fellowship (#658085) to AS, and AS

was supported by a 'Turku Collegium for Science and Medicine' Fellowship and a Marie

Sklodowska-Curie Postdoctoral Fellowship (#894963) at the time of writing. PM was

supported by ERC Advanced Grant #101020037 and NBM by ERC Advanced Grants #322784

and #834653.

References

1. Nicholls DG, Ferguson SJ. 2002 Bioenergetics. *Academic press*

2. Balaban RS, Nemoto S, Finkel T. 2005 Mitochondria, Oxidants, and Aging. *Cell* 120, 483–495.
(doi:10.1016/j.cell.2005.02.001)

3. Stier A, Bize P, Roussel D, Schull Q, Massemin S, Criscuolo F. 2014 Mitochondrial uncoupling as a
regulator of life-history trajectories in birds: an experimental study in the zebra finch. *Journal Of*
Experimental Biology 217, 3579–3589. (doi:10.1242/jeb.103945)

4. Delhaye J, Salamin N, Roulin A, Criscuolo F, Bize P, Christe P. 2016 Interspecific correlation
between red blood cell mitochondrial ROS production, cardiolipin content and longevity in birds. *Age*
(Dordrecht, Netherlands) 38, 433–443. (doi:10.1007/s11357-016-9940-z)

5. Hood WR *et al.* 2018 The Mitochondrial Contribution to Animal Performance, Adaptation, and Life-
History Variation. *Integrative and Comparative Biology* 58, 480–485. (doi:10.1093/icb/icy089)

6. Koch RE *et al.* 2021 Integrating Mitochondrial Aerobic Metabolism into Ecology and Evolution.
Trends in Ecology & Evolution 21, 1–12. (doi:10.1016/j.tree.2020.12.006)

7. Du SNN, Mahalingam S, Borowiec BG, Scott GR. 2016 Mitochondrial physiology and reactive
oxygen species production are altered by hypoxia acclimation in killifish (Fundulus heteroclitus).
Journal Of Experimental Biology 219, 1130–1138. (doi:10.1242/jeb.132860)

8. Pichaud N, m AE x000F6, Breton S, m FS x000F6, Rowinski P, Blier PU, Sandblom E. 2019 Cardiac
mitochondrial plasticity and thermal sensitivity in a fish inhabiting an artificially heated ecosystem.
Scientific Reports 9, 1–11. (doi:10.1038/s41598-019-54165-3)

9. Salin K, Villasevil EM, Anderson GJ, Auer SK, Selman C, Hartley RC, Mullen W, Chinopoulos C,
Metcalfe NB. 2018 Decreased mitochondrial metabolic requirements in fasting animals carry an
oxidative cost. *Funct Ecol* 32, 2149–2157. (doi:10.1111/1365-2435.13125)

10. Stier A, Bize P, Hsu B-Y, Ruuskanen S. 2019 Plastic but repeatable: rapid adjustments of
mitochondrial function and density during reproduction in a wild bird species. *Biology Letters* 15,
20190536. (doi:10.1098/rsbl.2019.0536)

11. Braganza A, Annarapu GK, Shiva S. 2020 Blood-based bioenergetics: An emerging translational
and clinical tool. *Molecular Aspects of Medicine* 71, 100835–12. (doi:10.1016/j.mam.2019.100835)

12. Gyllenhammer LE, Entringer S, Buss C, Wadhwa PD. 2020 Developmental programming of
mitochondrial biology: a conceptual framework and review. *Proceedings of the Royal Society of*
London. Series B: Biological Sciences 287, 20192713–10. (doi:10.1098/rspb.2019.2713)

13. Metcalfe N, Monaghan P. 2001 Compensation for a bad start: grow now, pay later? *Trends in*
Ecology & Evolution 16, 254–260.

14. Tarry-Adkins JL, Ozanne SE. 2017 Nutrition in early life and age-associated diseases. *Ageing*
Research Reviews 39, 96–105. (doi:10.1016/j.arr.2016.08.003)

15. Criscuolo F, Monaghan P, Nasir L, Metcalfe NB. 2008 Early nutrition and phenotypic
development: "catch-up" growth leads to elevated metabolic rate in adulthood. *Proceedings of the*
Royal Society B: Biological Sciences 275, 1565–1570. (doi:10.1098/rsbl.2006.0496)

16. Lee W-S, Monaghan P, Metcalfe NB. 2013 Experimental demonstration of the growth rate–
lifespan trade-off. *Proc Royal Soc B Biological Sci* 280, 20122370. (doi:10.1098/rspb.2012.2370)

17. Crowther NJ. 1998 Association between poor glucose tolerance and rapid post natal weight gain
in seven-year-old children. *Diabetologia* , 1–5.

18. Singhal A. 2010 Does Early Growth Affect Long-Term Risk Factors for Cardiovascular Disease? In
Nestlé Nutrition Institute Workshop Series, pp. 55–69. Nestlé Nutrition Institute Workshop Series.

19. Dillin A. 2002 Rates of Behavior and Aging Specified by Mitochondrial Function During
Development. *Science* 298, 2398–2401. (doi:10.1126/science.1077780)

20. Park KS, Kim SK, Kim MS, Cho EY, Lee JH, Lee K-U, Pak YK, Lee HK. 2003 Fetal and early postnatal
protein malnutrition cause long-term changes in rat liver and muscle mitochondria. *The Journal of*
nutrition 133, 3085–3090.

21. Pol M van de, Verhulst S. 2006 Age-Dependent Traits: A New Statistical Model to Separate
Within- and Between-Individual Effects. *Am Nat* 167, 766–773. (doi:10.1086/503331)

22. Stier A, Romestaing C, Schull Q, Lefol E, Robin J-P, ROUSSEL D, Bize P. 2017 How to measure
mitochondrial function in birds using red blood cells: a case study in the king penguin and
perspectives in ecology and evolution. *Methods in Ecology and Evolution* 8, 1172–1182.
(doi:10.1111/2041-210x.12724)

23. Beauchamp B, Harper M-E. 2016 In utero Undernutrition Programs Skeletal and Cardiac Muscle
Metabolism. *Frontiers in Physiology* 6, 1891–8. (doi:10.3389/fphys.2015.00401)

24. Krischek C, Janisch S, Naraballobh W, Brunner R, Wimmers K, Wicke M. 2015 Altered incubation
temperatures between embryonic Days 7 and 13 influence the weights and the mitochondrial
respiratory and enzyme activities in breast and leg muscles of broiler embryos. *Molecular*
reproduction and development 83, 71–78. (doi:10.1002/mrd.22596)

25. DuRant SE, Hopkins WA, Hepp GR, Walters JR. 2013 Ecological, evolutionary, and conservation
implications of incubation temperature-dependent phenotypes in birds. *Biological Reviews* 88, 499–
509. (doi:10.1111/brv.12015)

26. Nord A, Nilsson J-A. 2011 Incubation Temperature Affects Growth and Energy Metabolism in Blue
Tit Nestlings. *American Naturalist* 178, 639–651. (doi:10.1086/662172)

27. Ben-Ezra N, Burness G. 2016 Constant and Cycling Incubation Temperatures Have Long-Term
Effects on the Morphology and Metabolic Rate of Japanese Quail. *Physiological And Biochemical*
Zoology 90, 96–105. (doi:10.5061/dyad.76qt1)

28. Carvalho AV *et al.* 2020 Embryonic thermal manipulation has short and long-term effects on the
development and the physiology of the Japanese quail. *PLoS ONE* 15, e0227700-20.
(doi:10.1371/journal.pone.0227700)

29. Bratic A, Larsson N-G. 2013 The role of mitochondria in aging. *Journal of Clinical Investigation*
123, 951–957. (doi:10.1172/jci64125)

30. Sirsat SKG, Sirsat TS, Faber A, Duquaine A, Winnick S, Sotherland PR, Dzialowski EM. 2016
Development of endothermy and concomitant increases in cardiac and skeletal muscle mitochondrial
respiration in the precocial Pekin duck (Anas platyrhynchos domestica). *Journal Of Experimental*
Biology (doi:10.1242/jeb.132282)

31. Dawson NJ, Salmon P. 2020 Age-related increase in mitochondrial quantity may mitigate a
decline in mitochondrial quality in red blood cells from zebra finches (Taeniopygia guttata).
Experimental Gerontology 133, 110883. (doi:10.1016/j.exger.2020.110883)

32. Salin K, Villasevil EM, Anderson GJ, Lamarre SG, Melanson CA, McCarthy I, Selman C, Metcalfe NB.
2019 Differences in mitochondrial efficiency explain individual variation in growth performance.
Proceedings of the Royal Society of London. Series B: Biological Sciences 286, 20191466–8.
(doi:10.1098/rspb.2019.1466)

33. Stier A, Metcalfe NB, Monaghan P. 2020 Pace and stability of embryonic development affect
telomere dynamics: an experimental study in a precocial bird model. *Proceedings of the Royal Society*
of London. Series B: Biological Sciences 287, 20201378–9. (doi:10.1098/rspb.2020.1378)

34. Ottinger MA. 2001 Quail and other short-lived birds. *Experimental Gerontology* 36, 859–868.

35. Romao JM, Moraes TGV de, Teixeira RS de C, Buxade CC, Cardoso WM. 2009 INCUBATION OF
JAPANESE QUAIL EGGS AT DIFFERENT TEMPERATURES: HATCHABILITY, HATCH WEIGHT, HATCH TIME
AND EMBRYONIC MORTALITY. *Archives Vet Sci* 14, 155–162. (doi:10.5380/avs.v14i3.14887)

36. Jr FSO, Orcutt AB. 1976 Nesting and parental behavior in domestic common quail. *The Auk* 93,
135–141.

37. Karamercan MA, Weiss SL, Villarroel JPP, Guan Y, Werlin E, Figueredo R, Becker LB, Sims C. 2013
Can Peripheral Blood Mononuclear Cells Be Used as a Proxy for Mitochondrial Dysfunction in Vital
Organs During Hemorrhagic Shock and Resuscitation? *Shock* 40, 476–484.
(doi:10.1097/shk.000000000000026)

38. Tyrrell DJ, Bharadwaj MS, Jorgensen MJ, Register TC, Molina AJA. 2016 Blood cell respirometry is
associated with skeletal and cardiac muscle bioenergetics_ Implications for a minimally invasive
biomarker of mitochondrial health. *Redox Biology* 10, 65–77. (doi:10.1016/j.redox.2016.09.009)

39. Tyrrell DJ *et al.* 2017 Blood-Based Bioenergetic Profiling Reflects Differences in Brain
Bioenergetics and Metabolism. *Oxidative Medicine and Cellular Longevity* 2017, 1–9.
(doi:10.1155/2017/7317251)

40. Hedges CP, Woodhead JST, Wang HW, Mitchell CJ, Cameron-Smith D, Hickey AJR, Merry TL. 2019
Peripheral blood mononuclear cells do not reflect skeletal muscle mitochondrial function or
adaptation to high-intensity interval training in healthy young men. *Journal of applied physiology*
(Bethesda, Md. : 1985) 126, 454–461. (doi:10.1152/japplphysiol.00777.2018)

41. Moorsel D van *et al.* 2016 Demonstration of a day-night rhythm in human skeletal muscle
oxidative capacity. *Molecular Metabolism* 5, 635–645. (doi:10.1016/j.molmet.2016.06.012)

42. Stoffel MA, Nakagawa S, Schielzeth H. 2017 rptR: repeatability estimation and variance
decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution* 8,
1639–1644. (doi:10.1111/2041-210x.12797)

43. Short KR, Moller N, Bigelow ML, Coenen-Schimke J, Nair KS. 2008 Enhancement of Muscle
Mitochondrial Function by Growth Hormone. *Journal of Clinical Endocrinology & Metabolism* 93,
597–604. (doi:10.1210/jc.2007-1814)

44. Manoli I, Alesci S, Blackman MR, Su YA, Rennert OM, Chrousos GP. 2007 Mitochondria as key
components of the stress response. *Trends in Endocrinology & Metabolism* 18, 190–198.
(doi:10.1016/j.tem.2007.04.004)

45. McCurdy CE *et al.* 2016 Maternal obesity reduces oxidative capacity in fetal skeletal muscle of
Japanese macaques. *JCI Insight* 1. (doi:10.1172/jci.insight.86612ds1)

46. Abraham M, Collins CA, Flewelling S, Camazine M, Cahill A, Cade WT, Duncan JG. 2018
Mitochondrial inefficiency in infants born to overweight African- American mothers. *International*
Journal of Obesity 83, 1–11. (doi:10.1038/s41366-018-0051-z)

47. Galli GLJ, Crossley J, Elsey RM, Dzialowski EM, Shiels HA, II DAC. 2016 Developmental plasticity of
mitochondrial function in American alligators, Alligator mississippiensis. *AJP: Regulatory, Integrative*
and Comparative Physiology 311, R1164–R1172. (doi:10.1152/ajpregu.00107.2016)

48. McCaw BA, Stevenson TJ, Lancaster LT. 2020 Epigenetic Responses to Temperature and Climate.
Integr Comp Biol 60, 1469–1480. (doi:10.1093/icb/icaa049)

49. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. 2013 The hallmarks of aging. *Cell*
153, 1194–1217. (doi:10.1016/j.cell.2013.05.039)

50. Chung DJ, Healy TM, Schulte PM. 2018 Mitochondria, temperature, and the pace of life.
Integrative and Comparative Biology (doi:10.1093/icb/icy013/4990407)

51. Curran JE *et al.* 2007 Genetic determinants of mitochondrial content. *Human Molecular Genetics*
16, 1504–1514. (doi:10.1093/hmg/ddm101)

Electronic Supplementary Material (ESM)

Stier *et al.* (2021): **Experimental demonstration of prenatal programming of mitochondrial aerobic metabolism lasting until adulthood**

Correlation between blood cells and brain mitochondrial function

In addition to blood samples (see main text and Fig. S1A for mitochondrial responses of intact blood cells to a standard sequential substrate/inhibitor addition protocol), small brain samples (*ca.* 10-15mg) were collected from a subsample of adult females (Day 90, N = 22) following euthanasia by cervical dislocation and rapid dissection on ice (< 3min after euthanasia). Brain samples were transferred in 1mL of Mir05 buffer and stored on ice until analysis (< 1h after euthanasia). Brain samples were quickly blotted on absorbent paper and weighed (± 0.01mg, Sartorius AC211S®). They were then homogenized during 1min using micro-dissecting scissors and further diluted to 2mg.mL^{-1} in Mir05 (following Salin et al., 2016). One mL of this preparation was then transferred to the Oxygraph-2k chamber already containing 1mL of Mir05 equilibrated at 40°C, giving a final brain content of 1mg.mL^{-1} in the chamber. Mitochondrial respiration was then assessed following a standard sequential substrate/inhibitor addition protocol. First, baseline O$_2$ consumption was recorded (this parameter is not used in further analysis), then substrates of complex I (pyruvate 5mM and malate 2mM) and complex II (succinate 10mM) were added leading to a non-phosphorylating state hereafter referred as *LEAK$_{PMS}$* (*i.e.* equivalent to classical state 2), followed by the addition of a saturating amount of ADP (2mM) to stimulate oxidative phosphorylation (hereafter referred as *OXPHOS$_{PMS+ADP}$*; equivalent to classical state 3). ATP synthesis was then inhibited with 2.5 µM of oligomycin leading a non-phosphorylating state hereafter referred as *LEAK$_{oligo}$* (*i.e.* equivalent to classical state 4), followed by the stimulation of maximal uncoupled respiration (hereafter referred as *ETS*) using a sequential titration of FCCP (0.5µM per step) until maximal stimulation was reached. Finally, we inhibited mitochondrial respiration using antimycin A (2.5µM), and this residual non-mitochondrial O$_2$ consumption was subtracted from the mitochondrial parameters described above. Average mitochondrial responses to this chemical titration is presented in Fig S1B. We verified mitochondrial integrity by adding 20µM

of cytochrome c in preliminary experiments, as well as that the mechanical permeabilization we used was able to fully permeabilize brain samples by testing an additive effect of chemical permeabilization on *OXPHOS* respiration using digitonin (5-25ng.mL^{-1}).

Fig S1: **Mitochondrial O$_2$ consumption responses during high-resolution respirometry assays in intact blood cells (A) and permeabilized brain samples (B) of adult Japanese quails.** Letters indicate significant differences according to GEE post-hoc tests. PMS stands for pyruvate+malate+succinate as substrates to fuel brain mitochondrial respiration. A direct comparison of respiration rates between tissues is unfortunately not possible due to methodological constraints (*i.e.* intact cells *vs.* permeabilized tissue).

Relationships between blood cell and brain mitochondrial function were analyzed using *Pearson* coefficients of correlation *r,* and a meta-analysis on the Z-transformed *r* for the five main focal correlations (*i.e.* between blood cells and brain parameters measuring overall the same mitochondrial responses, see Fig. S2) using OpenMEE software (Wallace et al., 2017). For investigating potential relationships between brain and blood cell mitochondrial functions, blood cell respiration rates were normalized directly by the cellular protein content, and therefore are expressed as pmol O$_2$.s^{-1}.mg protein^{-1}.

Mitochondrial respiration rates were highly correlated within-tissues (Fig S2A), and weakly to moderately correlated between tissues (Fig S2A & S2B). While only two of the focal correlations out of five were significant, the overall meta-analytic correlation (*Zr*) was significant (p < 0.001) and of moderate effect size (Fig 2B).

Fig S2: Relationships between brain and blood cell mitochondrial respiration rates. (A) Correlation matrix between the different mitochondrial parameters in the two tissues. Numbers in right part represent the *Pearson* coefficients of correlation *r*, and significance is indicated by the lines in the scatter plot (solid line: p < 0.05, dashed line: p < 0.10, no line p > 0.10). The 5 focal correlations we were the most interested in between the two tissues (being analysed separately in panel B) are highlighted with a blue background. **(B) Meta-analysis of the focal between-tissues correlations.** Z-transformed *r* are presented along with their 95% confidence intervals. Significant parameters are shown in black and non-significant ones in grey.

Table S1: **Summary of the statistical models (GEEs) testing the effect of age, prenatal treatment, sex and the interaction between age and prenatal treatment on mitochondrial respiration rates of blood cells while controlling for protein content (to correct for variations in cells number) and time of day.** Significant parameters (p ≤ 0.05) are reported in bold and parameters presenting a non-significant trend (p ≤ 0.10) in italic. Estimates are given for age = Day 20, prenatal treatment = medium temperature and sex = females.

(A) *ROUTINE* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	18.39 ± 7.53	**0.007** (7.31)
Repeated effect	**Age**	1.74 ± 0.99	**< 0.001** (16.85)
Fixed effects & covariates	**Prenatal treatment**	-1.80 ± 1.30	**0.005** (12.76)
	Sex	0.28 ± 0.97	0.77 (0.08)
	Time of day	-9.90 ± 3.07	**0.001** (10.42)
	RBC protein content	1.67 ± 0.70	**0.017** (5.70)
	Age x Prenatal treatment	0.35 ± 1.44	**0.047** (7.96)
(B) *OXPHOS* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	11.98 ± 6.48	**0.023** (5.15)
Repeated effect	**Age**	3.56 ± 0.88	**< 0.001** (44.54)
Fixed effects & covariates	**Prenatal treatment**	-1.07 ± 1.07	**0.014** (10.66)
	Sex	-0.08 ± 0.88	0.93 (0.01)
	Time of day	-9.40 ± 2.62	**< 0.001** (12.85)
	RBC protein content	1.70 ± 0.58	**0.004** (8.45)
	Age x Prenatal treatment	0.18 ± 1.33	*0.10* (6.20)

(C) *LEAK* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	6.41 ± 2.79	**0.039** (4.27)
Repeated effect	**Age**	-1.82 ± 0.42	**< 0.001** (13.33)
Fixed effects & covariates	*Prenatal treatment*	-0.73 ± 0.50	*0.09* (10.66)
	Sex	0.36 ± 0.30	0.23 (1.43)
	Time of day	-0.50 ± 0.99	0.61 (0.26)
	RBC protein content	-0.030 ± 0.26	0.91 (0.01)
	Age x Prenatal treatment	0.18 ± 1.33	0.12 (5.82)
(D) *ETS* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	38.07 ± 10.32	**< 0.001** (12.52)
Repeated effect	**Age**	-6.37 ± 1.49	**0.003** (8.95)
Fixed effects & covariates	*Prenatal treatment*	-1.53 ± 1.77	*0.07* (7.14)
	Sex	-0.53 ± 1.32	0.69 (0.16)
	Time of day	5.27 ± 4.23	0.21 (1.56)
	RBC protein content	1.08 ± 0.92	0.24 (1.37)
	Age x Prenatal treatment	-0.06 ± 2.10	**0.004** (13.25)

Table S2: **Summary of the statistical models (GEEs) testing the effect of age, prenatal treatment, sex and the interaction between age and prenatal treatment on mitochondrial flux control ratios while controlling for the time of day.** Significant parameters (p ≤ 0.05) are reported in bold. Estimates are given for age = Day 20, prenatal treatment = medium temperature and sex = females.

(A) Coupling efficiency (*OxCE*)		Estimate ± SE	p-value (χ^2)
	Intercept	0.831 ± 0.016	**< 0.001** (104.37)
Repeated effect	**Age**	0.061 ± 0.011	**< 0.001 (60.94)**
Fixed effects & covariates	Prenatal treatment	0.020 ± 0.013	0.45 (2.66)
	Sex	-0.016 ± 0.008	**0.039 (4.26)**
	Time of day	-0.033 ± 0.026	0.21 (1.57)
	Age x Prenatal treatment	-0.006 ± 0.017	0.29 (3.77)
(B) Mitochondrial capacity usage (FCR$_{R/ETS}$)		Estimate ± SE	p-value (χ^2)
	Intercept	0.716 ± 0.028	**< 0.001** (765.11)
Repeated effect	**Age**	0.110 ± 0.018	**< 0.001 (76.65)**
Fixed effects & covariates	Prenatal treatment	-0.001 ± 0.023	0.16 (5.19)
	Sex	-0.010 ± 0.014	0.49 (0.48)
	Time of day	-0.212 ± 0.051	**< 0.001 (17.05)**
	Age x Prenatal treatment	0.002 ± 0.031	0.31 (3.55)