Experimental demonstration of prenatal programming of mitochondrial

aerobic metabolism lasting until adulthood

Antoine Stier [a,b,c], Pat Monaghan [a] & Neil B. Metcalfe [a]

[a] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,

Glasgow, UK

[b] Department of Biology, University of Turku, Turku, Finland

[c] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622,

Villeurbanne, France

*Corresponding author: antoine.stier@gmail.com / amstie@utu.fi

Abstract

It is increasingly being postulated that among-individual variation in mitochondrial function underlies variation in individual performance (*e.g.* growth rate) and state of health. Environmental conditions experienced before birth have been suggested to programme postnatal mitochondrial biology, but the hypothesis that early-life conditions induce consistent and long-lasting differences in mitochondrial function among individuals remains mostly untested. We tested this hypothesis in an avian model by experimentally manipulating prenatal conditions (incubation temperature and stability), then measuring mitochondrial aerobic metabolism in blood cells from the same individuals during postnatal growth and at adulthood. Mitochondrial aerobic metabolism changed markedly across life stages, and part of these age-related changes were influenced by the prenatal experimental conditions. A high incubation temperature induced a consistent and long-lasting increase in mitochondrial aerobic metabolism, while unstable incubation conditions had a delayed effect, with an increased metabolism only being observed at adulthood. While we detected significant within-individual consistency in mitochondrial aerobic metabolism across life-stages, the prenatal temperature regime only accounted for a relatively small proportion (<20%) of the consistent among-individual differences we observed. Our results demonstrate that prenatal conditions can program consistent and long-lasting differences in mitochondrial function, which could potentially underly among-individual variation in performance and health state.

Keywords: developmental programming, fetal programming, bioenergetics, mitochondria, Japanese quail.

Introduction

Mitochondria produce more than 90% of the cellular energy routinely required by animal cells in the form of adenosine triphosphate (ATP), produced through oxidative phosphorylation [1]. Mitochondria are also a major source of reactive oxygen species (ROS), which are likely to be involved in the ageing process if produced in excess of the antioxidant capacity [2]. Consequently, variations in mitochondrial function are likely to play an important role in shaping the life and death of individuals, and to underlie phenotypic variation observed both within and between species [3-6]. Importantly, while mitochondrial function is a very plastic trait (*i.e.* influenced by tissue, temperature, O_2/food availability or reproductive stage; [7-10]), it has been shown in both human and animal models that adult individuals exhibit some degree of within-individual consistency in mitochondrial traits through time [10,11]. Understanding the origin of such consistent among-individual differences in mitochondrial function (*e.g.* genetics, early-life programming, permanent environmental effects) is key to understand the role of mitochondria in shaping individual performance and health state [6,12].

Scientific evidence from both the biomedical and the ecological fields of research suggest that conditions experienced during early-life have major and persistent effects on adult phenotype and physiology [13,14], a phenomenon termed developmental or early-life programming. For instance, accelerated postnatal growth has been associated with increased metabolic rate at adulthood and reduced lifespan in animal models [15,16], as well as with increased risks of developing various age-related pathologies [17,18]. The underlying molecular and physiological mechanisms are probably numerous, but mitochondrial dysfunction has emerged as one potential candidate linking early-life conditions to both immediate and delayed effects on phenotype and health state [12,14,19,20]. However, much

64 remains to be discovered about the importance of early-life conditions in determining

65 consistent and long-lasting differences in mitochondrial function between individuals.

66   Investigating the long-term programming of physiological function is more amenable

through longitudinal studies (*i.e.* measuring the same individuals over time). Indeed, while

avoiding issues linked to the selective disappearance of specific phenotypes [21], longitudinal

studies allow estimating 1) the transient, persistent or delayed nature of early-life effects at

the within-individual level,   2) within-individual changes through time (*e.g.* age-related

alteration) and how they could be affected by early-life conditions, and 3) if consistent among-

individual differences in mitochondrial function exist (*i.e.* within-individual consistency

through time) and to what extent they could be explained by early-life conditions. In recent

years, measuring mitochondrial function from blood cells has emerged as an opportunity to

conduct longitudinal studies of mitochondrial biology both in human and animal models

[10,11,22].

77   To the best of our knowledge, only cross-sectional studies have been conducted to

date in the context of early-life effects on mitochondrial function [12]. These studies suggest

for instance that maternal food restriction during pregnancy can alter mitochondrial

bioenergetics [12,23]. This has been framed overall as the 'developmental programming of

mitochondrial biology' hypothesis [12]. However, none of these studies show an unequivocal

direct and persistent effect of prenatal conditions on mitochondrial function. For instance,

early-life dietary manipulations usually result in alterations of body mass (and body fat

content) during growth and at adulthood, which could by itself alter mitochondrial function

independently of prenatal conditions. Additionally, measurements of mitochondrial function

are mostly conducted at a single time point, often months or years after birth, so preventing

evaluation of the effect of manipulations on early-life mitochondrial function, and the

potential effects on age-related variation in mitochondrial function.

Oviparous species offer the opportunity to conduct experiments on the direct effects

of prenatal conditions on mitochondrial function, since the developmental conditions of the

embryo can be altered through direct manipulations of the eggs. For instance, it has been

shown that higher incubation temperature of chicken eggs led to higher oxidative capacity

(state 3 / *OXPHOS*) and mitochondrial proton leak (state 4 / *LEAK*) in developing bird embryos

[24]. However, this study did not control for treatment-induced differences in embryo

developmental stage and did not assess the persistence of the effects on mitochondrial

function postnatally and over the long term.

Here we investigate the potential prenatal programming of postnatal mitochondrial

function from early-life to adulthood by using manipulations of incubation temperature and

stability in an avian model. Both the absolute incubation temperature and the stability of

incubation temperature convey environmental information to the embryo that could be used

to adjust the postnatal phenotype to suit anticipated environmental conditions [25].

Mitochondrial function has been shown to exhibit changes according to age/life-stage in both

animal models and humans [26-28]. Therefore, a second aim was to characterize within-

individual changes in mitochondrial function between the peak of the growth phase and early-

adulthood, and to evaluate potential effects of the prenatal environment on such age-related

changes. Since it has been shown that individuals with more efficient mitochondria (*i.e.*

producing more ATP per unit of O_2 consumed) could grow faster [29], we hypothesize that

mitochondrial efficiency could be maximized during early life to support the costly process of

growth. Finally, our last objective was to characterize the extent to which among-individual

differences in mitochondrial function are consistent over time (*i.e.* within-individual

consistency), but also the extent to which such consistent among-individual differences could

originate from variation in the prenatal environment. This is conceptually important for

evaluating the potential for early-life conditions to have effects on mitochondrial function that

persist through adulthood.

Material and Methods

Experimental design

All procedures were conducted (as previously described in [30]) in 2016 in accordance with UK regulations under the Home Office Project Licence 70/8335 granted to PM and the Home Office Personal Licence ICB1D39E7 granted to AS. We used Japanese quail (*Coturnix japonica*) as a model since successful artificial incubation has been widely used in this species (*e.g.* [31]), they are precocial birds and can be reared successfully without the parent being present, therefore avoiding any confounding effect linked to variation in parental care; in addition, they reach sexual maturity and therefore adulthood very quickly (*ca.* 50 days [32]) making it possible to examine long term effects spanning life history stages within a reasonable time. Japanese quail eggs were bought from Moonridge Farm (Devon, UK) and delivered within 48 hours after collection. We used 164 eggs from which 107 chicks hatched, 6 died in the first 5 days and were excluded from the study, and 25 birds were not used for mitochondrial function assessment due to logistical constraints, giving a sample size of 76 birds.

Eggs were incubated at 3 constant temperatures as described in more detail in [30]: high (H) = 38.4°C, medium (M) = 37.7°C and low (L) = 37.0°C. Additionally a fourth group was incubated under 'unstable' (U) temperature conditions, with an incubation temperature of 37.7°C but five incubation recesses of 30min during the day, leading to a daily average of 37.0°C, similar to the L group (*i.e.* its matched control for developmental speed [30]). Our prenatal experimental treatments affected developmental speed in the predicted direction (*i.e.* higher temperature leading to faster development) while having no significant effects on hatching success or mass at hatching and during growth [30].

Animal husbandry rooms for post-hatching rearing were maintained at 21°C on a 14L:10D

cycle throughout the experiment. After hatching (Day 0) chicks were placed for 24h in a larger

incubator set at 37°C, before being placed into their respective enclosure within each room at Day

1 where an additional heat source was provided (Brinsea Comfort brooder 40, 42W), as well as *ad*

libitum food (Heygates starter crumbs, 22% protein) and water. The additional heat source was

removed at day 15 and chick food was switched to adult pellets (Heygates quail and partridge

pellets, 16% protein) at the same time (*i.e.* 5 days before the first blood sampling). Chicks were

maintained in mixed-sex groups until Day 25 when they can be sexed morphologically. Females

were then kept in groups in enclosures and males were placed by pairs in $0.8m^2$ cages to avoid

female exhaustion due to male harassment and to limit male-male conflicts.

Sampling procedures

We used blood cells [22] to measure mitochondrial function of the same individuals at

both Day 20 (peak of the growth phase, see [30]) and Day 60 (early-adulthood). Blood samples

(*ca.* 200μL) were collected at Day 20 and Day 60 by venipuncture of the wing vein with a 26*G*

needle and collection using heparinised capillaries. Blood was centrifuged 10min at 3000*g* and 4°C

to separate plasma from packed blood cells, and 50μL of blood cells (*ca.* 150 to 200.10^6 cells) were

immediately re-suspended in 1mL of Mir05 buffer (0.5 mM EGTA, 3 mM $MgCl_2$, 60 mM K-

lactobionate, 20 mM taurine, 10 mM KH_2PO_4, 20 mM Hepes, 110 mM sucrose, free fatty acid

bovine serum albumin (1 g/L), pH 7.1) and kept on ice until analysis (< 1.5h after blood collection).

High-resolution respirometry analysis of mitochondrial function in intact blood cells

We followed the protocol described in detail for king penguins [22], with some minor

modifications. Briefly, samples were washed before the start of the mitochondrial

measurements by centrifugating the tubes to pellet the blood cells and discarding the

supernatant. Blood cells were then re-suspended in 1mL of respiratory buffer MiR05 pre-

equilibrated at 40.0°C in the chamber of the Oxygraph-2k (Oroboros Instruments, Innsbruck,

Austria), and transferred to the chamber already containing another 1mL of Mir05. After

closing the chamber, baseline O_2 consumption was recorded, followed by the inhibition of

ATP-synthesis using oligomycin (Oligo: 2.5µM), then by the stimulation of maximal uncoupled

respiration using a sequential titration of carbonyl cyanide-p-

trifluoromethoxyphenylhydrazone (FCCP: 0.5µM per step) until maximal stimulation was

reached, and finally by the inhibition of mitochondrial respiration using antimycin A (AA:

2.5µM). Mitochondrial responses of blood cells to this chemical titration is presented in Fig

S1A. We then calculated *ROUTINE* respiration (*i.e.* endogenous cellular respiration = *baseline*

- AA), *OXPHOS* respiration (*i.e.* O_2 consumption linked to ATP synthesis = *baseline - Oligo*),

LEAK respiration (*i.e.* O_2 consumption mostly linked to mitochondrial proton leak = *Oligo - AA*)

and *ETS* respiration (*i.e.* maximal O_2 consumption of the electron transport system = *FCCP -*

AA). We also calculated two mitochondrial *flux control ratios* (FCRs), namely the OXPHOS

coupling efficiency (*OxCE* calculated as *OXPHOS/ROUTINE*) indicating the proportion of

endogenous respiration being linked to ATP synthesis, and an index of mitochondrial capacity

usage ($FCR_{R/ETS}$ calculated as *ROUTINE/ETS*) indicating the proportion of maximal capacity

being used under endogenous cellular conditions [22]. To account for potential differences in

cell quantity between samples, we quantified the protein content of the cell suspension using

a BCA protein assay (ThermoScientific) following [22].

Some studies have shown that mitochondrial function measured in blood cells is

correlated to some extent to mitochondrial function in other tissues such as kidneys, heart,

skeletal muscles and brain (*e.g.* [22,33-35], while some other studies did not find any

significant relationship (*e.g.* [36]). Therefore, to evaluate the relevance of using blood cell

mitochondrial function in Japanese quails, we compared mitochondrial function between

blood cells and brain samples (see ESM) in a subsample of adult females (N = 22); and found

a moderate but overall significant correlation between these two tissues (meta-analytic *r* =

0.44, 95% C.I. = [0.27;0.59], p < 0.001; see ESM Fig. S2).

Data analysis

We used generalized estimated equations (GEEs) in *SPSS* 24.0 to investigate the effects of

age (*i.e.* repeated effect), prenatal treatment (*i.e.* low, medium, high or unstable incubation

temperature), sex and their interactions on mitochondrial parameters, with the associated post-

hoc tests (non-significant interactions were removed from the final models except the focal *age*

x treatment interaction). We included the cellular protein content as a covariate in the models

to account for potential variations in cell quantity between samples, and the time of day to

account for potential circadian variations in mitochondrial function [37]. To analyse within-

individual consistency in mitochondrial traits, we used the *RptR* package [38] in *R* 3.4.2 and

included bird identity as a random (*i.e.* focal) term, as well as age and cellular protein content

as fixed effects to account for changes in mitochondrial traits with age and quantity of cells per

sample. We re-ran the same analyses while including experimental treatment group as an

additional random term to investigate the proportion of within-individual consistency being

attributed to our prenatal treatments. Due to some failed laboratory assays (*e.g.* due to residual

inhibition of mitochondrial respiration in the Oroboros chambers), the final dataset includes n =

136 measurements from N = 76 individuals. A few (n = 8) *ETS* values being non-biological (*ETS* <

ROUTINE) were removed from the final dataset.

Results

Effect of age and prenatal temperature regime on mitochondrial respiration rates

Fig 1: Effects of age and prenatal temperature regime on blood cells mitochondrial respiration rates: (A) *ROUTINE* respiration: endogenous mitochondrial O₂ consumption, **(B)** *OXPHOS* respiration: mitochondrial O₂ consumption being linked to ATP synthesis **(C)** *LEAK* respiration: mitochondrial O₂ consumption being mostly linked to mitochondrial proton leak, and **(D)** *ETS* respiration: maximal mitochondrial O₂ consumption induced by mitochondrial uncoupling. Details of statistical tests are given in Table S1, means are presented ± SE, letters indicate significant differences between groups according to GEE post-hoc tests, and within-group age effects are presented as * = p < 0.05, ** = p < 0.01 and *** = p < 0.001. Statistical models included blood cells protein content as a covariate to account for potential variations in blood cell quantity between samples.

Endogenous mitochondrial respiration (*ROUTINE*) was significantly affected by the age of individuals and the prenatal experimental treatment, as well as by their interaction (Fig 1A, Table S1A). Specifically, *ROUTINE* decreased overall with age, but in a treatment-specific manner (H and L decreasing significantly, while M and U only showed a non-significant decrease). Chicks from the high temperature group had a significantly higher *ROUTINE* at Day 20 than other groups. At adulthood (Day 60), birds from the high temperature and unstable

groups had higher *ROUTINE* than low temperature birds, while birds in the medium temperature group were intermediate (Fig. 1A). *ROUTINE* respiration was significantly affected by the time of day (*i.e.* decreasing through the daylight hours) but not by sex (Table S1A).

Mitochondrial *OXPHOS* respiration decreased significantly with the age of individuals (Fig 1B, Table S1B). It also significantly differed between treatment groups: birds from the high temperature group had the higher *OXPHOS*, followed by unstable and medium temperature ones, and finally by low temperature ones (see Fig 4B for significance). *OXPHOS* respiration decreased significantly with the time of day, but was not significantly affected by sex (Table S1B).

LEAK respiration increased significantly with age, but was not significantly affected by treatment, time of day or sex (Fig 1C, Table S1C). *ETS* respiration increased overall with age, but in a treatment-specific manner (significant age x treatment interaction, with M and U increasing significantly while H and L only exhibited a non-significant increase; Fig 1D, Table S1D). Chicks from the high temperature group had a significantly higher *ETS* at Day 20 than those from the medium temperature and unstable groups, while low temperature ones had an intermediate phenotype (Fig. 1D). At adulthood (Day 60), birds from the low temperature group had lower *ETS* respiration than all other groups (no significant differences were observed between H, M and U groups). *ETS* respiration was not significantly affected by the time of day or sex (Table S1D).

Effect of age and prenatal temperature regime on mitochondrial flux control ratios

Mitochondrial OXPHOS coupling efficiency (*i.e. OxCE*) significantly decreased with age, but was not significantly affected by the prenatal treatment, either as a main factor or in

interaction with age (Fig 2A, Table S2A). However, this parameter exhibited a sex difference,

with females having slightly less efficient mitochondria than males (Table S2A, LS-mean ± SE:

females = 0.836 ± 0.007 *vs.* males = 0.852 ± 0.005, p = 0.039), but there was no significant

effect of the time of day (Table S2A).

The proportion of maximal respiration being used under endogenous cellular

conditions (*i.e.* $FCR_{R/ETS}$) significantly decreased with age, but was not significantly affected by

the prenatal treatment, either as a main factor or in interaction with age (Fig 2B), nor by the

sex of individuals. However, this ratio decreased significantly with the time of day (Table S2B,

p = 0.004).

Fig 2: Effects of age and prenatal temperature regime on blood cells mitochondrial: (A) OXPHOS coupling efficiency (*OxCE*: proportion of endogenous respiration devoted to ATP synthesis) and (B) mitochondrial capacity usage (FCR_ROUTINE/ETS: proportion of maximal mitochondrial respiration being used under endogenous cellular conditions). Details of statistical tests are given in Table S2, means are presented ± SE and letters indicate significant differences.

Within-individual consistency in mitochondrial phenotype

All mitochondrial respiration rates exhibited a significant within-individual consistency

through time between the peak of the growth phase (Day 20) and early adulthood (Day 60)

(Fig 3A; *ROUTINE*: R = 0.58, p < 0.001; *OXPHOS*: R = 0.53, p < 0.001; *LEAK*: R = 0.23, p = 0.045;

ETS: R = 0.48, p < 0.001). While mitochondrial OXPHOS coupling efficiency was not significantly

repeatable (*OxCE*: R = 0.12, p = 0.17), the mitochondrial capacity usage was (*FCR_R/ETS*: R = 0.42,

p = 0.001; Fig. 3A). When looking at the variance in mitochondrial traits being explained

respectively by bird identity (*i.e.* intrinsic variation) and by incubation temperature (*i.e.*

extrinsic variation linked to our treatment), the contribution of the latter was relatively

modest (Fig 3B; all < 8%) compared to bird identity (all > 21%, except *OxCE*). Experimental

treatment thereby accounted for 7.9-17.5% of the within-individual consistency in

mitochondrial respiration rates, and less than 2% of the within-individual consistency in

mitochondrial flux ratios.

Fig 3: **(A) Within-individual consistency of mitochondrial respiration rates and flux control ratios measured in intact blood cells at day 20 and 60 in Japanese quail.** Age-adjusted repeatability (*i.e.* consistency) estimates *R* are presented along with their 95% C.I. The statistical model also contained cellular protein content as a covariate to control for potential variations linked to differences in cell quantity between samples. **(B) Respective variance in mitochondrial traits being explained by bird identity (*i.e.* intrinsic variation), our experimental treatment (incubation temperature, *i.e.* extrinsic variation) and other non-investigated factors as well as measurement error (*i.e.* residual).**

Discussion

Our results demonstrate that variation in the prenatal environment can influence mitochondrial function from post-natal life into adulthood. We found marked differences in mitochondrial function between the peak of the postnatal growth phase and early adulthood, and the pattern of this age-related change was influenced by the environmental temperature regime experienced during prenatal development. Overall, our results suggest that prenatal conditions can have both permanent and delayed effects on mitochondrial function, while also influencing how mitochondrial function changes with age. The persistence of pre-natal temperature effects on mitochondrial function into adulthood suggests that these are long lasting and possibly permanent. We found significant within-individual consistency of mitochondrial respiration rates across life-stages, with individuals showing consistent relatively high or low values through time.

Age-related variation in mitochondrial function

Mitochondrial parameters differed markedly between the peak of the growth phase and early adulthood (6 parameters out of 6). The overall pattern suggests that oxidative phosphorylation decreases with age, while both maximal mitochondrial capacity and proton leak increase, leading to a higher mitochondrial coupling efficiency but also to a more intense utilization of the total mitochondrial capacity during growth than at adulthood. These results are in line with our hypothesis that mitochondrial efficiency should be maximized in early-life to sustain the growth process, and also indicate that mitochondrial maximal capacity is more intensively used during growth. Such effects could potentially be mediated by the known impact of growth hormone on mitochondrial function [39], although this remains to be properly tested.

Prenatal programming of mitochondrial function by incubation temperature and stability

Our results clearly show that the prenatal environment can affect mitochondrial respiration rates (*i.e.* endogenous cellular respiration, oxidative phosphorylation and maximal mitochondrial capacity) in the long-term since we found effects of incubation temperature regime both during postnatal growth and at adulthood. Specifically, it seems that a high incubation temperature increased subsequent mitochondrial metabolism at both life-stages (*i.e.* persistent effect on *ROUTINE*, *OXPHOS* and *ETS* compared to low temperature). Instability in incubation temperature also increased mitochondrial metabolism (compared to its matched control for developmental speed, *i.e.* low temperature), but this was only observed at adulthood (*i.e.* delayed effect on *ROUTINE* and *ETS*). This demonstrates that prenatal environmental conditions can have relatively immediate effects (*i.e.* during postnatal growth) as suggested by previous correlative studies in mammals [40,41], delayed effects as shown for prenatal hypoxia in an experimental reptile model [42], and more importantly, persistent effects lasting from early post-natal life to adulthood. Importantly, unlike previous studies in mammals and reptiles, our results cannot be biased by differences in body mass between experimental groups at the time of mitochondrial measurement [30]. Mitochondrial flux control ratios were not affected by the prenatal treatments, suggesting that the differences we observe in respiration rates between groups were relatively consistent across the different mitochondrial respiration rates we measured.

Interestingly, age-related changes in mitochondrial function were also partly influenced by the prenatal environment (*i.e.* age-related decrease in *ROUTINE* for H and L groups only, as well as age-related increase in *ETS* for M and U groups only), suggesting that the prenatal environment does not only affect how mitochondria work postnatally, but also the way mitochondrial aerobic metabolism changes with age. Considering the importance of

mitochondria in the ageing process [26,43], such effects of the prenatal environment on both

mitochondrial aerobic metabolism and its age-related changes could have potential

consequences in influencing ageing trajectories. Even if this hypothesis remains to be properly

tested, our results on telomere length (a key hallmark of ageing [43,44]) using the same

experimental birds suggest that this might be the case. Indeed, both high temperature and

unstable incubation conditions led to increased aerobic metabolism (see above), while also

leading to shorter telomeres [30], a typical signature of premature ageing.

Although we found clear programming effects of mitochondrial aerobic metabolism by

prenatal environmental conditions, we have so far no information about their potential

adaptive or maladaptive value. In principle, high and unstable incubation temperatures

convey opposite information about environmental quality (*i.e.* high temperature = good

quality environment, unstable = poor quality environment; [25]), yet they were the two

experimental conditions leading to higher mitochondrial respiration rates (compared to their

respective controls) in the present study. Further studies investigating the adaptive role of

such variation in mitochondrial function (*e.g.* by testing individual performance under

contrasted postnatal environmental conditions) will be needed to conclude about the

potential adaptive value of prenatal programming through incubation temperature and

stability.

Within-individual consistency in mitochondrial function and relative importance of prenatal

conditions

Mitochondrial respiration rates and flux control ratios exhibited a significant within-

individual consistency over time (at the exception of *OxCE*), while they have been measured

at two very different life stages (peak of growth *vs.* early-adulthood) and that marked changes

in mitochondrial aerobic metabolism occur during postnatal development (see above). To the best of our knowledge, this is the third demonstration that consistent among-individual differences in mitochondrial aerobic metabolism exist (see [10] in wild adult passerine birds, and [11] in adult humans), but the first to be conducted under well-controlled environmental conditions (*i.e.* excluding bias linked to consistent individual differences in environmental conditions) and across life-stages. These findings of a significant within-individual consistency in mitochondrial function over time have important implications for the possibility for early-life conditions to program mitochondrial function over the life course of individuals [12]. However, our experimental treatment only accounted for less than 20% of the within-individual consistency observed in mitochondrial respiration rates, meaning that more than 80% of the observed within-individual consistency must be explained by other prenatal factors (*e.g.* maternal transfer of nutrients, hormones) and/or by genetic differences between individuals. To the best of our knowledge, there is no information published about the heritability of mitochondrial function, but mtDNA copy number has been shown to be significantly heritable ($h^2 = 33\%$) in humans [45]. Therefore, estimating the relative importance of genetic *vs.* environmental drivers of mitochondrial function appears now fundamental to evaluate the magnitude to which early-life environment could program mitochondrial function and its potential downstream effects such as disease risk and individual performance.

Competing interests

We declare having no competing interests

Data availability

Datasets used in this manuscript are available at: https://doi.org/10.6084/m9.figshare.16708483.v1

Author's contribution

AS designed the study, conducted the experimental work, data analysis and wrote the

manuscript. NBM and PM had input on study design and data analysis, and commented on

the manuscript.

Acknowledgements

We are grateful to Karine Salin for the help in setting-up mitochondrial analysis of brain

samples, to Pierre Bize for kindly providing access to his O2k-oxygraph, to Norith Eckbo for the

graphic design of Fig. S2, to Franklin Lo, Becky Shaw and Katie Byrne for their help in collecting

blood samples, and to Graham Law and his team for taking care of animal husbandry. Finally,

AS is grateful to the crew of the Marion Dufresnes and the French Polar Institute (IPEV) for

hosting him from 56 to 21° South while writing the first draft of this manuscript. The project

was funded by a Marie Sklodowska-Curie Postdoctoral Fellowship (#658085) to AS, and AS

was supported by a 'Turku Collegium for Science and Medicine' Fellowship and a Marie

Sklodowska-Curie Postdoctoral Fellowship (#894963) at the time of writing.

References

1.   Nicholls, D. G. & Ferguson, S. J. 2002 Bioenergetics. *Academic press*

2.   Balaban, R. S., Nemoto, S. & Finkel, T. 2005 Mitochondria, Oxidants, and Aging. *Cell* **120**, 483–
495. (doi:10.1016/j.cell.2005.02.001)

3.   Stier, A., Bize, P., Roussel, D., Schull, Q., Massemin, S. & Criscuolo, F. 2014 Mitochondrial
uncoupling as a regulator of life-history trajectories in birds: an experimental study in the
zebra finch. *J Exp Biol* **217**, 3579–3589. (doi:10.1242/jeb.103945)

4.   Delhaye, J., Salamin, N., Roulin, A., Criscuolo, F., Bize, P. & Christe, P. 2016 Interspecific
correlation between red blood cell mitochondrial ROS production, cardiolipin content and
longevity in birds. *Age (Dordr)* **38**, 433–443. (doi:10.1007/s11357-016-9940-z)

5.   Hood, W. R. et al. 2018 The Mitochondrial Contribution to Animal Performance, Adaptation,
and Life-History Variation. *Integrative and Comparative Biology* **58**, 480–485.
(doi:10.1093/icb/icy089)

6.   Koch, R. E. et al. 2021 Integrating Mitochondrial Aerobic Metabolism into Ecology and
Evolution. *Trends in Ecology & Evolution* **21**, 1–12. (doi:10.1016/j.tree.2020.12.006)

7.   Du, S. N. N., Mahalingam, S., Borowiec, B. G. & Scott, G. R. 2016 Mitochondrial physiology and
reactive oxygen species production are altered by hypoxia acclimation in killifish (Fundulus
heteroclitus). *J Exp Biol* **219**, 1130–1138. (doi:10.1242/jeb.132860)

8.   Pichaud, N., m, A. E. X., Breton, S., m, F. S. X., Rowinski, P., Blier, P. U. & Sandblom, E. 2019
Cardiac mitochondrial plasticity and thermal sensitivity in a fish inhabiting an artificially
heated ecosystem. *Sci. Rep.* **9**, 1–11. (doi:10.1038/s41598-019-54165-3)

9.   Salin, K., Villasevil, E. M., Anderson, G. J., Auer, S. K., Selman, C., Hartley, R. C., Mullen, W.,
Chinopoulos, C. & Metcalfe, N. B. 2018 Decreased mitochondrial metabolic requirements in
fasting animals carry an oxidative cost. **32**, 2149–2157. (doi:10.1111/1365-2435.13125)

10.   Stier, A., Bize, P., Hsu, B.-Y. & Ruuskanen, S. 2019 Plastic but repeatable: rapid adjustments of
mitochondrial function and density during reproduction in a wild bird species. *Biol Letters* **15**,
20190536. (doi:10.1098/rsbl.2019.0536)

11.   Braganza, A., Annarapu, G. K. & Shiva, S. 2020 Blood-based bioenergetics: An emerging
translational and clinical tool. *Molecular Aspects of Medicine* **71**, 100835–12.
(doi:10.1016/j.mam.2019.100835)

12.   Gyllenhammer, L. E., Entringer, S., Buss, C. & Wadhwa, P. D. 2020 Developmental
programming of mitochondrial biology: a conceptual framework and review. *Proceedings of*
the Royal Society of London. Series B: Biological Sciences **287**, 20192713–10.
(doi:10.1098/rspb.2019.2713)

13.   Metcalfe, N. & Monaghan, P. 2001 Compensation for a bad start: grow now, pay later? *Trends*
in Ecology & Evolution **16**, 254–260.

14.   Tarry-Adkins, J. L. & Ozanne, S. E. 2017 Nutrition in early life and age-associated diseases.
Ageing Research Reviews **39**, 96–105. (doi:10.1016/j.arr.2016.08.003)

15.   Criscuolo, F., Monaghan, P., Nasir, L. & Metcalfe, N. B. 2008 Early nutrition and phenotypic
development: 'catch-up' growth leads to elevated metabolic rate in adulthood. *Proceedings of*
the Royal Society B: Biological Sciences **275**, 1565–1570. (doi:10.1098/rsbl.2006.0496)

16.   Lee, W.-S., Monaghan, P. & Metcalfe, N. B. 2013 Experimental demonstration of the growth
rate–lifespan trade-off. *Proceedings of the Royal Society of London. Series B: Biological*
Sciences **280**, 20122370.

17.   Crowther, N. J. 1998 Association between poor glucose tolerance and rapid post natal weight
gain in seven-year-old children. *Diabetologia*, 1–5.

18.   Singhal, A. 2010 Does Early Growth Affect Long-Term Risk Factors for Cardiovascular Disease?
In *Nestlé Nutrition Institute Workshop Series*, pp. 55–69.

19. Dillin, A. 2002 Rates of Behavior and Aging Specified by Mitochondrial Function During
Development. *Science* **298**, 2398–2401. (doi:10.1126/science.1077780)

20. Park, K. S., Kim, S. K., Kim, M. S., Cho, E. Y., Lee, J. H., Lee, K.-U., Pak, Y. K. & Lee, H. K. 2003
Fetal and early postnatal protein malnutrition cause long-term changes in rat liver and muscle
mitochondria. *The Journal of nutrition* **133**, 3085–3090.

21. van de Pol, M. & Verhulst, S. 2006 Age-dependent traits: a new statistical model to separate
within- and between-individual effects. *The American Naturalist* **167**, 766–773.

22. Stier, A., Romestaing, C., Schull, Q., Lefol, E., Robin, J.-P., ROUSSEL, D. & Bize, P. 2017 How to
measure mitochondrial function in birds using red blood cells: a case study in the king penguin
and perspectives in ecology and evolution. *Methods in Ecology and Evolution* **8**, 1172–1182.
(doi:10.1111/2041-210X.12724)

23. Beauchamp, B. & Harper, M.-E. 2016 In utero Undernutrition Programs Skeletal and Cardiac
Muscle Metabolism. *Front. Physiol.* **6**, 1891–8. (doi:10.3389/fphys.2015.00401)

24. Krischek, C., Janisch, S., Naraballobh, W., Brunner, R., Wimmers, K. & Wicke, M. 2015 Altered
incubation temperatures between embryonic Days 7 and 13 influence the weights and the
mitochondrial respiratory and enzyme activities in breast and leg muscles of broiler embryos.
Mol. Reprod. Dev. **83**, 71–78. (doi:10.1002/mrd.22596)

25. DuRant, S. E., Hopkins, W. A., Hepp, G. R. & Walters, J. R. 2013 Ecological, evolutionary, and
conservation implications of incubation temperature-dependent phenotypes in birds.
Biological Reviews **88**, 499–509. (doi:10.1111/brv.12015)

26. Bratic, A. & Larsson, N.-G. 2013 The role of mitochondria in aging. *J. Clin. Invest.* **123**, 951–
957. (doi:10.1172/JCI64125)

27. Sirsat, S. K. G., Sirsat, T. S., Faber, A., Duquaine, A., Winnick, S., Sotherland, P. R. & Dzialowski,
E. M. 2016 Development of endothermy and concomitant increases in cardiac and skeletal
muscle mitochondrial respiration in the precocial Pekin duck (Anas platyrhynchos domestica).
J Exp Biol (doi:10.1242/jeb.132282)

28. Dawson, N. J. & Salmon, P. 2020 Age-related increase in mitochondrial quantity may mitigate
a decline in mitochondrial quality in red blood cells from zebra finches (Taeniopygia guttata).
Experimental Gerontology **133**, 110883. (doi:10.1016/j.exger.2020.110883)

29. Salin, K., Villasevil, E. M., Anderson, G. J., Lamarre, S. G., Melanson, C. A., McCarthy, I.,
Selman, C. & Metcalfe, N. B. 2019 Differences in mitochondrial efficiency explain individual
variation in growth performance. *Proceedings of the Royal Society of London. Series B:*
Biological Sciences **286**, 20191466–8. (doi:10.1098/rspb.2019.1466)

30. Stier, A., Metcalfe, N. B. & Monaghan, P. 2020 Pace and stability of embryonic development
affect telomere dynamics: an experimental study in a precocial bird model. *Proceedings of the*
Royal Society of London. Series B: Biological Sciences **287**, 20201378–9.
(doi:10.1098/rspb.2020.1378)

31. Romao, J. M., Moraes, T. G. V. de, Teixeira, R. S. de C., Buxade, C. C. & Cardoso, W. M. 2010
Incubation of Japanese quail eggs at different temperatures: hatchability, hatch weight, hatch
time and embryonic mortality. *Archives of Veterinary Science* **14**, 155–162.

32.   Ottinger, M. A. 2001 Quail and other short-lived birds. *Experimental Gerontology* **36**, 859–868.

33.   Karamercan, M. A., Weiss, S. L., Villarroel, J. P. P., Guan, Y., Werlin, E., Figueredo, R., Becker, L.
490   B. & Sims, C. 2013 Can Peripheral Blood Mononuclear Cells Be Used as a Proxy for
491   Mitochondrial Dysfunction in Vital Organs During Hemorrhagic Shock and Resuscitation?
492   *Shock* **40**, 476–484. (doi:10.1097/SHK.0000000000000026)

493   34.   Tyrrell, D. J., Bharadwaj, M. S., Jorgensen, M. J., Register, T. C. & Molina, A. J. A. 2016 Blood
494   cell respirometry is associated with skeletal and cardiac muscle bioenergetics_ Implications
495   for a minimally invasive biomarker of mitochondrial health. *Redox Biology* **10**, 65–77.
496   (doi:10.1016/j.redox.2016.09.009)

497   35.   Tyrrell, D. J. et al. 2017 Blood-Based Bioenergetic Profiling Reflects Differences in Brain
498   Bioenergetics and Metabolism. *Oxidative Medicine and Cellular Longevity* **2017**, 1–9.
499   (doi:10.1155/2017/7317251)

500   36.   Hedges, C. P., Woodhead, J. S. T., Wang, H. W., Mitchell, C. J., Cameron-Smith, D., Hickey, A. J.
501   R. & Merry, T. L. 2019 Peripheral blood mononuclear cells do not reflect skeletal muscle
502   mitochondrial function or adaptation to high-intensity interval training in healthy young men.
503   *J. Appl. Physiol.* **126**, 454–461. (doi:10.1152/japplphysiol.00777.2018)

504   37.   van Moorsel, D. et al. 2016 Demonstration of a day-night rhythm in human skeletal muscle
505   oxidative capacity. *Molecular Metabolism* **5**, 635–645. (doi:10.1016/j.molmet.2016.06.012)

506   38.   Stoffel, M. A., Nakagawa, S. & Schielzeth, H. 2017 rptR: repeatability estimation and variance
507   decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*
508   **8**, 1639–1644. (doi:10.1111/2041-210X.12797)

509   39.   Short, K. R., Moller, N., Bigelow, M. L., Coenen-Schimke, J. & Nair, K. S. 2008 Enhancement of
510   Muscle Mitochondrial Function by Growth Hormone. *Journal of Clinical Endocrinology &*
511   *Metabolism* **93**, 597–604. (doi:10.1210/jc.2007-1814)

512   40.   McCurdy, C. E. et al. 2016 Maternal obesity reduces oxidative capacity in fetal skeletal muscle
513   of Japanese macaques. *JCI Insight* **1**. (doi:10.1172/jci.insight.86612DS1)

514   41.   Abraham, M., Collins, C. A., Flewelling, S., Camazine, M., Cahill, A., Cade, W. T. & Duncan, J. G.
515   2018 Mitochondrial inefficiency in infants born to overweight African- American mothers.
516   *International Journal of Obesity* **83**, 1–11. (doi:10.1038/s41366-018-0051-z)

517   42.   Galli, G. L. J., Crossley, J., Elsey, R. M., Dzialowski, E. M., Shiels, H. A. & Crossley, D. A., II 2016
518   Developmental plasticity of mitochondrial function in American alligators, Alligator
519   mississippiensis. *AJP: Regulatory, Integrative and Comparative Physiology* **311**, R1164–R1172.
520   (doi:10.1152/ajpregu.00107.2016)

521   43.   López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. 2013 The hallmarks of
522   aging. *Cell* **153**, 1194–1217. (doi:10.1016/j.cell.2013.05.039)

523   44.   Wilbourn, R. V., Moatt, J. P., Froy, H., Walling, C. A., Nussey, D. H. & Boonekamp, J. J. 2018 The
524   relationship between telomere length and mortality risk in non-model vertebrate systems: a
525   meta-analysis. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **373**, 20160447–9.

526   45.   Curran, J. E. et al. 2007 Genetic determinants of mitochondrial content. *Human Molecular*
527   *Genetics* **16**, 1504–1514. (doi:10.1093/hmg/ddm101)

Electronic Supplementary Material (ESM)

Stier *et al.* (2021): **Experimental demonstration of prenatal programming of mitochondrial aerobic metabolism lasting until adulthood**

Correlation between blood cells and brain mitochondrial function

In addition to blood samples (see main text and Fig. S1A for mitochondrial responses of intact blood cells to a standard sequential substrate/inhibitor addition protocol), small brain samples (*ca.* 10-15mg) were collected from a subsample of adult females (Day 90, N = 22) following euthanasia by cervical dislocation and rapid dissection on ice (< 3min after euthanasia). Brain samples were transferred in 1mL of Mir05 buffer and stored on ice until analysis (< 1h after euthanasia). Brain samples were quickly blotted on absorbent paper and weighed (± 0.01mg, Sartorius AC211S®). They were then homogenized during 1min using micro-dissecting scissors and further diluted to 2mg.mL^{-1} in Mir05 (following Salin et al., 2016). One mL of this preparation was then transferred to the Oxygraph-2k chamber already containing 1mL of Mir05 equilibrated at 40°C, giving a final brain content of 1mg.mL^{-1} in the chamber. Mitochondrial respiration was then assessed following a standard sequential substrate/inhibitor addition protocol. First, baseline O$_2$ consumption was recorded (this parameter is not used in further analysis), then substrates of complex I (pyruvate 5mM and malate 2mM) and complex II (succinate 10mM) were added leading to a non-phosphorylating state hereafter referred as *LEAK$_{PMS}$* (*i.e.* equivalent to classical state 2), followed by the addition of a saturating amount of ADP (2mM) to stimulate oxidative phosphorylation (hereafter referred as *OXPHOS$_{PMS+ADP}$*; equivalent to classical state 3). ATP synthesis was then inhibited with 2.5 µM of oligomycin leading a non-phosphorylating state hereafter referred as *LEAK$_{oligo}$* (*i.e.* equivalent to classical state 4), followed by the stimulation of maximal uncoupled respiration (hereafter referred as *ETS*) using a sequential titration of FCCP (0.5µM per step) until maximal stimulation was reached. Finally, we inhibited mitochondrial respiration using antimycin A (2.5µM), and this residual non-mitochondrial O$_2$ consumption was subtracted from the mitochondrial parameters described above. Average mitochondrial responses to this chemical titration is presented in Fig S1B. We verified mitochondrial integrity by adding 20µM

44   of cytochrome c in preliminary experiments, as well as that the mechanical permeabilization

45   we used was able to fully permeabilize brain samples by testing an additive effect of chemical

46   permeabilization on *OXPHOS* respiration using digitonin (5-25ng.mL^{-1}).

Fig S1: Mitochondrial O$_2$ consumption responses during high-resolution respirometry assays in intact blood cells (A) and permeabilized brain samples (B) of adult Japanese quails. Letters indicate significant differences according to GEE post-hoc tests. PMS stands for pyruvate+malate+succinate as substrates to fuel brain mitochondrial respiration. A direct comparison of respiration rates between tissues is unfortunately not possible due to methodological constraints (*i.e.* intact cells *vs.* permeabilized tissue).

Relationships between blood cell and brain mitochondrial function were analyzed using

Pearson coefficients of correlation *r,* and a meta-analysis on the Z-transformed *r* for the five

main focal correlations (*i.e.* between blood cells and brain parameters measuring overall the

same mitochondrial responses, see Fig. S2) using OpenMEE software (Wallace et al., 2017). For

investigating potential relationships between brain and blood cell mitochondrial functions,

blood cell respiration rates were normalized directly by the cellular protein content, and

therefore are expressed as pmol O$_2$.s^{-1}.mg protein^{-1}.

Mitochondrial respiration rates were highly correlated within-tissues (Fig S2A), and

weakly to moderately correlated between tissues (Fig S2A & S2B). While only two of the focal

correlations out of five were significant, the overall meta-analytic correlation (*Zr*) was

significant (p < 0.001) and of moderate effect size (Fig 2B).

Fig S2: Relationships between brain and blood cell mitochondrial respiration rates. (A) Correlation matrix between the different mitochondrial parameters in the two tissues. Numbers in right part represent the *Pearson* coefficients of correlation *r*, and significance is indicated by the lines in the scatter plot (solid line: p < 0.05, dashed line: p < 0.10, no line p > 0.10). The 5 focal correlations we were the most interested in between the two tissues (being analysed separately in panel B) are highlighted with a blue background. **(B) Meta-analysis of the focal between-tissues correlations.** Z-transformed *r* are presented along with their 95% confidence intervals. Significant parameters are shown in black and non-significant ones in grey.

Table S1: **Summary of the statistical models (GEEs) testing the effect of age, prenatal treatment, sex**
and the interaction between age and prenatal treatment on mitochondrial respiration rates of blood
cells while controlling for protein content (to correct for variations in cells number) and time of day.
Significant parameters (p ≤ 0.05) are reported in bold and parameters presenting a non-significant
trend (p ≤ 0.10) in italic. Estimates are given for age = Day 20, prenatal treatment = medium
temperature and sex = females.
98

(A) *ROUTINE* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	18.39 ± 7.53	**0.007** (7.31)
Repeated effect	Age	1.74 ± 0.99	**< 0.001** (16.85)
Fixed effects & covariates	Prenatal treatment	-1.80 ± 1.30	**0.005** (12.76)
	Sex	0.28 ± 0.97	0.77 (0.08)
	Time of day	-9.90 ± 3.07	**0.001** (10.42)
	RBC protein content	1.67 ± 0.70	**0.017** (5.70)
	Age x Prenatal treatment	0.35 ± 1.44	**0.047** (7.96)
(B) *OXPHOS* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	11.98 ± 6.48	**0.023** (5.15)
Repeated effect	Age	3.56 ± 0.88	**< 0.001** (44.54)
Fixed effects & covariates	Prenatal treatment	-1.07 ± 1.07	**0.014** (10.66)
	Sex	-0.08 ± 0.88	0.93 (0.01)
	Time of day	-9.40 ± 2.62	**< 0.001** (12.85)
	RBC protein content	1.70 ± 0.58	**0.004** (8.45)
	Age x Prenatal treatment	0.18 ± 1.33	*0.10* (6.20)

(C) *LEAK* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	6.41 ± 2.79	**0.039** (4.27)
Repeated effect	**Age**	-1.82 ± 0.42	**< 0.001** (13.33)
Fixed effects & covariates	*Prenatal treatment*	-0.73 ± 0.50	*0.09* (10.66)
	Sex	0.36 ± 0.30	0.23 (1.43)
	Time of day	-0.50 ± 0.99	0.61 (0.26)
	RBC protein content	-0.030 ± 0.26	0.91 (0.01)
	Age x Prenatal treatment	0.18 ± 1.33	0.12 (5.82)

(D) *ETS* respiration		Estimate ± SE	p-value (χ^2)
	Intercept	38.07 ± 10.32	**< 0.001** (12.52)
Repeated effect	**Age**	-6.37 ± 1.49	**0.003** (8.95)
Fixed effects & covariates	*Prenatal treatment*	-1.53 ± 1.77	*0.07* (7.14)
	Sex	-0.53 ± 1.32	0.69 (0.16)
	Time of day	5.27 ± 4.23	0.21 (1.56)
	RBC protein content	1.08 ± 0.92	0.24 (1.37)
	Age x Prenatal treatment	-0.06 ± 2.10	**0.004** (13.25)

99
100

Table S2: **Summary of the statistical models (GEEs) testing the effect of age, prenatal treatment, sex and the interaction between age and prenatal treatment on mitochondrial flux control ratios while controlling for the time of day.** Significant parameters (p ≤ 0.05) are reported in bold. Estimates are given for age = Day 20, prenatal treatment = medium temperature and sex = females.

(A) Coupling efficiency (*OxCE*)		Estimate ± SE	p-value (χ^2)
	Intercept	0.831 ± 0.016	**< 0.001** (104.37)
Repeated effect	**Age**	0.061 ± 0.011	**< 0.001 (60.94)**
Fixed effects & covariates	Prenatal treatment	0.020 ± 0.013	0.45 (2.66)
	Sex	-0.016 ± 0.008	**0.039 (4.26)**
	Time of day	-0.033 ± 0.026	0.21 (1.57)
	Age x Prenatal treatment	-0.006 ± 0.017	0.29 (3.77)
(B) Mitochondrial capacity usage (FCR$_{R/ETS}$)		Estimate ± SE	p-value (χ^2)
	Intercept	0.716 ± 0.028	**< 0.001** (765.11)
Repeated effect	**Age**	0.110 ± 0.018	**< 0.001 (76.65)**
Fixed effects & covariates	Prenatal treatment	-0.001 ± 0.023	0.16 (5.19)
	Sex	-0.010 ± 0.014	0.49 (0.48)
	Time of day	-0.212 ± 0.051	**< 0.001 (17.05)**
	Age x Prenatal treatment	0.002 ± 0.031	0.31 (3.55)