Isotopic signatures induced by upwelling reveal regional fish

populations in Lake Tanganyika

Benedikt Ehrenfels[1,2]*[¶], Julian Junker[3,4][¶], Demmy Namutebi[4,5], Cameron M. Callbeck[1],

Christian Dinkel[1], Anthony Kalangali[6], Ismael A. Kimirei[6,7], Athanasio S. Mbonde[6],

Julieth B. Mosille[6], Emmanuel A. Sweke[6,8], Carsten J. Schubert[1,2], Ole Seehausen[3,4],

Catherine E. Wagner[9], Bernhard Wehrli[1,2]

[1] Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department Surface Waters –

Research and Management, Kastanienbaum, Switzerland

[2] ETH Zurich, Institute of Biogeochemistry and Pollutant Dynamics, Zurich, Switzerland

[3] Eawag, Swiss Federal Institute of Aquatic Science and Technology, Department Fish Ecology and

Evolution, Kastanienbaum, Switzerland

[4] University of Bern, Institute of Ecology & Evolution, Bern, Switzerland

[5] IHE Delft, Institute for Water Education, Delft, Netherlands

[6] TAFIRI, Tanzania Fisheries Research Institute, Kigoma, Tanzania

[7] TAFIRI, Tanzania Fisheries Research Institute, Dar es Salaam, Tanzania

[8] DSFA, Deep Sea Fishing Authority, Zanzibar, Tanzania

[9] University of Wyoming, Department of Botany and Program in Ecology, Laramie, USA

* Corresponding author: Benedikt Ehrenfels (benedikt.ehrenfels@eawag.ch)

[¶] shared first authorship

Keywords: stratification, mixing, isoscape, isotope analysis, genetic structure, diet, lipid

storage, food web, fish migration, pelagic

Abstract

Lake Tanganyika's pelagic fish sustain the second largest inland fishery in Africa and are under pressure from heavy fishing and global warming related increases in stratification. Only little is known about whether basin-scale hydrodynamics – including a more stratified north and an upwelling-driven south – induce ecological and genetic differences among populations of highly mobile, pelagic fish inhabiting these different areas. Here, we examine whether the basin-scale dynamics leave distinct isotopic imprints in the pelagic fish of Lake Tanganyika, which may reveal differences in habitat, diet, or lipid content. We conducted two lake-wide campaigns during different seasons and collected physical, nutrient, chlorophyll, phytoplankton and zooplankton data. Additionally, we analyzed the pelagic fish – the clupeids *Stolothrissa tanganicae*, *Limnothrissa miodon* and four *Lates* species – for their isotopic and elemental carbon (C) and nitrogen (N) compositions. The $\delta^{13}C$ values were significantly higher in the productive south after the upwelling/mixing period across all trophic levels, implying that the fish have regional foraging grounds, and thus record these latitudinal isotope gradients. By combining our isotope data with genetics, we demonstrate that the fish form regional populations on a seasonal to multiannual time scale. Based on $\delta^{15}N$ and C:N ratios, we found no strong evidence for varying diets or lipid contents between those regional populations. Additional analyses revealed that isotopic variations between specimens from the same location are not linked to genetic differences. We suggest that the development of basin-scale ecological differences in response to the prevailing hydrodynamic regimes may be inhibited by lake-wide gene flow on the long term. Our findings show that the pelagic fish species are genetically adapted to the whole lake, but they form regional populations

47   on short time scales. This implies that sustainable management strategies may adopt

48   basin-scale fishing quotas.

1 Introduction

Lake Tanganyika is by volume the second largest freshwater lake in the world, and its

pelagic fish community sustains the second largest inland fishery in Africa (1), providing

important employment opportunities and animal protein for millions of people in the

riparian communities (2,3). The pelagic food web in Lake Tanganyika is composed of a

copepod-dominated zooplankton assemblage, a phyto- and zooplankton grazer

community consisting of two endemic sardine species (*Stolothrissa tanganicae* and

Limnothrissa miodon), and a predator assemblage comprising of four endemic latid

species (genus *Lates*), of which *Lates stappersii* is the most common (4). Today, the

sardines and *Lates stappersii* account for 95 % of the pelagic fish catch in Lake

Tanganyika (5).

The pelagic fish stocks suffer from heavy fishing (2,6) and from a long term decline that

was attributed to climate change (7–9). The increased warming of the surface waters

caused by climate change leads to steep temperature gradients in the water column. These

gradients build physical barriers to vertical mixing, thereby limiting the transfer of

nutrients to surface waters where light is available to drive primary productivity (7,8,10–

13).

Although assessing potential long-term changes in the ecology of the pelagic fish is

impaired by data scarcity, Lake Tanganyika's limnological cycle offers the opportunity to

study the impact of varying levels of stratification on a basin-scale. This annual cycle is

driven by climatic differences between the north and south and is characterized by four

stages (Fig. 1a-d; 14,15): (i) In the warm rainy season (November-March), stagnant and

highly stratified waters lead to an overall nutrient-depleted epilimnion (Fig. 1 a), and

internal waves only cause local nutrient injections into the surface waters (16). (ii) In

March-May, the southeast trade winds initiate the lake circulation in the upper water

column, resulting in strong nutrient upwelling in the southern basin (Fig. 1 b). (iii) The

upwelling in the south transforms into a convective mixing of the upper ~150 m. The

sinking, cool surface waters in the south reverse the lake circulation by initiating a

northward current between 50-100 m and a surface counter current, further weakening

thermal stratification across the lake (Fig. 1 c). (iv) The trade winds cease in October,

slowing down the circulation, while the water column re-stratifies lake-wide. A weaker,

secondary upwelling leads to a nutrient pulse at the northern end of the lake (Fig. 1 d).

Fig. 1: The limnological cycle of Lake Tanganyika with its four major phases according to Plisnier et al. (14) and Verburg et al. (15). (**a**) Stagnant, highly stratified waters during the warm rainy season (November-March) only support low nutrient availability. (**b**) The onset of the cool dry winds in March-May initiates the upwelling in the South leading to high nutrient fluxes in this region. (**c**) The lake circulation reverses during the dry season (May-September). Water column stratification is low and the nutrient availability high across the lake, with a maximum in the convective mixing area in the south. (**d**) The trade winds cease in October slowing down the lake circulation, while the water column re-stratifies. A weaker secondary upwelling leads to a nutrient pulse at the northern end of the lake. During the dry season, wind-driven upwelling and mixing are the dominant driving force behind nutrient injections into the euphotic zone, whereas internal waves are particularly important in the rainy season. The color gradient indicates the level of thermal stratification. Note that this latitudinal cross-section is not to scale and that the outlined mechanism primarily affects the upper water column (<200 m). Our two sampling campaigns were timed at the seasonal transitions in September/October and April/May to compare the effects of the preceding dry and rainy seasons. (**e**) The map shows the nine stations for water column and plankton

sampling. Fish samples representing the pelagic catch were collected from the respective coastal

villages/towns.

The limnological cycle of Lake Tanganyika leads to seasonal and regional gradients in

nutrient availability, causing overall higher primary productivity in the dry season

compared to the rainy season and higher primary productivity in the south than in the

north of the lake (17–21). High densities of both zooplankton and the sardines

Stolothrissa and *Limnothrissa* are coupled to the phytoplankton blooms (22–24), which

also seem to benefit sardine spawning and recruitment (25). The north-south variability

furthermore affects the zooplankton community composition: shrimps and calanoid

copepods prevail in the south (26–28), whereas cyclopoid copepods and jellyfish

dominate in the north (24,29,30). Differences in the zooplankton community may in turn

influence predatory fish. Mannini et al. (31) found that the diet of *Lates stappersii* in the

north is heterogeneous and consists of copepods, shrimps, and sardines, whereas *Lates*

stappersii in the south feed mainly on shrimps.

The spatial variability in Lake Tanganyika's pelagic habitat, driven by the mixing regime,

could additionally impact the life cycle (e.g. spawning phenology, developmental timing

and recruitment success) of the pelagic fish species (23,25,32,33). The spatial variation in

the pelagic environment might generate different fitness optima and drive divergent

adaptation of pelagic fish populations between north and south, if horizontal migration

remains limited. However, recent genetic studies of the sardines (34,35) and the four

Lates species (36) did not find evidence for genetic population differentiation along the

north-south gradient, suggesting that gene flow may overcome any possible effects of

divergent natural selection between the basins. Nonetheless, the fish populations may still

respond to differences in physicochemical conditions and food supply by evolving

ecological adaptations to the regional environments. Such responses might involve

variations in food web interactions (37) or lipid storage for bridging lean periods (38,39).

However, possible effects of the seasonality and spatial variation in the physical mixing

regime of Lake Tanganyika on the distribution and ecology of its pelagic fish have not

been studied.

Characterizing seasonal and spatial gradients in the carbon (C) and nitrogen (N)

elemental and isotopic composition of Lake Tanganyika's food web may provide insight

to the migration distances, diets, and lipid contents of the fish species. Indeed, an isotopic

study of fish and their surrounding food web along a geographical gradient can reveal

regional population isolation if environmental differences among sites translate into

divergent isotopic signatures of regional or local fish populations (40). In this isotopic

framework, the $^{13}C/^{12}C$ ratio or $\delta^{13}C$ increases only little from one trophic level to the

next and therefore reflects the source of primary production (41–43). Differences in

primary productivity can alter the $\delta^{13}C$ of particulate organic matter (POM): high primary

productivity results typically in high values of $\delta^{13}C$, due to the ongoing depletion of the

of the DIC pool and decreasing discrimination against ^{13}C by phytoplankton (44–49).

Changes in $\delta^{13}C$ at the base of the food web can be tracked across trophic levels from

plankton to fish in lake ecosystems (50). In Lake Tanganyika, O'Reilly et al. (7) and

Verburg (12) used the $\delta^{13}C$-POM to reconstruct historical changes in primary

productivity. In addition, previous $\delta^{13}C$ analyses in the northern basin have reported

higher $\delta^{13}C$-POM values in the productive dry season compared to the rainy season (51–

53). The ratio of $^{15}N/^{14}N$, or $\delta^{15}N$, provides insight into the trophic position of an

organism, because it increases significantly with each trophic level. This successive

enrichment allows estimating an organism's trophic position in the food web and is used

in ecology to describe prey and predator relationships (41–43). Lastly, the elemental C:N

ratio was often used to infer the lipid content of fish muscle tissue, with higher C:N

denoting higher lipid contents (54,55).

In this study, we explore the latitudinal and seasonal patterns of δ^{13}C and δ^{15}N in the

pelagic food web of Lake Tanganyika in the context of the lake's limnological variability.

During two lake-wide field campaigns in the final phases of the dry and the rainy

seasons, we measured the C and N isotopic and elemental compositions of the major

pelagic food web members (POM, zooplankton, the bivalve *Pleiodon spekii*, fish). These

samples were collected in concert with limnological data including the physical

properties of the water column, oxygen and nutrient concentrations, chlorophyll, as well

as the phyto- and zooplankton community and abundance (56). Using the extensive data

sets from those two contrasting time points, we first tested to which extend the regional

and seasonal patterns in primary productivity induce systematic differences in the

isotopic signatures of plankton, and then tracked the isotopic signals and C:N ratios

through the food web to the pelagic fish. The results allowed us to assess the extent of

regional isolation and ecological differentiation of the pelagic fish populations. Finally,

we tested whether existing genetic differences (35,36) were linked to dietary differences

in the six major pelagic fish species.

2 Materials & Methods

2.1 Study site and sampling

Our two Lake Tanganyika sampling campaigns, spanning two different hydrological conditions across a north-south transect of ~500 km, were conducted at the end of the dry season (28 September - 8 October 2017) and the end of the following rainy season (27 April - 7 May 2018). Water column and plankton characteristics were sampled during two cruises on *M/V Maman Benita* (56,57). At the end of the dry season, we collected fish samples as described in Junker et al. (2020) at station 1, station 2, station 5, station 7 and station 9 during a land-based excursion prior to the cruise (17-24 September 2017), whereas all nine landing sites, corresponding to our nine pelagic sampling stations, were sampled during the cruise at the end of the rainy season (Fig. 1 e). In addition, we took fish samples in Kigoma in July 2017 (35).

2.2 Physical and chemical parameters

We measured temperature, dissolved oxygen, photosynthetically active radiation, and in-situ chlorophyll fluorescence via CTD profiling (Sea-Bird SBE 19plus) at each station. From these stations, we also collected water with large Niskin bottles (20-30 L) at 5-25 m depth intervals down to 250 m depth. Water column stratification was expressed as buoyancy frequency (N^2) and Schmidt stability (Sc). We interpreted clear peaks in N^2 as thermoclines, whereby the N^2 value at the peak provides a measure of steepness of the thermocline. In addition, we calculated the Sc over 1 m^2 between 50 and 100 m for each

station using the R package 'rLakeAnalyzer' (58). This depth interval extends from the

typical location of the nitrate peak to the bottom of the euphotic zone (56,59,60). For a

more detailed description of the thermal structure of the water column see Ehrenfels et al.

(56).

Water samples to measure nutrients (phosphate, ammonium, nitrate, and nitrite) were

taken directly from the Niskin bottles, filtered sterile through 0.2 μm filters and processed

on-board following standard methods (61–63). On average, the detection limits were

0.22, 0.34, 0.20, and 0.03 μM for phosphate, ammonium, nitrate, and nitrite, respectively.

Water samples to measure dissolved inorganic carbon (DIC) were collected in 12 mL

exetainers directly from the Niskin bottles and filtered sterile (0.2 μm). Samples were

stored at room temperatures and shipped to Switzerland. At Eawag Kastanienbaum, the

DIC concentrations were measured by high temperature combustion catalytic oxidation

using a Shimadzu TOC-L Analyzer (Shimadzu TOC-VCPH/CPN). A 2 mL aliquot of the

DIC sample was used to quantify the isotopic fractionation of δ^{13}C-DIC. The aliquot

subsample was transferred to a new 12 mL exetainer, where it was Helium purged for 2

minutes. The sample was then capped and 50 μL orthophosphorous acid (85 %) was

added. The samples were mixed and stored for ~15 h at room temperature for

equilibration prior to analysis by GC-IRMS (Isoprime). Sample δ^{13}C-DIC was adjusted to

the standard *Carrara marmor* (ETH Zurich).

2.3 CO$_2$ fixation rates

Carbon fixation incubations and rate calculations were done as described in Schunck et

al. (64) and Callbeck et al. (65). Briefly, samples were carefully filled from the Niskin

into 4.5 L polycarbonate bottles capped with polypropylene membranes. Per sampled

depth, we filled off triplicate bottles, including one control (no added label) and duplicate

treatments (with amended ^{13}C-HCO$_3^-$). We added 4.5 mL of ^{13}C-bicarbonate solution (1 g

^{13}C-bicarbonate in 50 ml water; Sigma Aldrich) to each of the treatment bottles. The label

was mixed in the treatment bottles for ~30 min under shaking. Thereafter, a 12 mL

subsample was taken for quantifying the labelling percent (mean 2.8 %). The resulting

headspace was re-filled with water from the same depth, and bottles were then incubated

headspace-free in 60 L incubators covered with shaded light filters (LEE Filters)

mimicking the in-situ irradiance and light spectrum. After 24 h, the samples were filtered

on pre-combusted GF/F filters (Whatman). The filters were oven-dried (60° C for 48 h)

and stored at ambient temperatures. Filter samples were shipped to Switzerland and

further processed as described in 2.7. Due to the small difference between the in-situ and

incubation temperatures (<5° C), the derived CO$_2$ fixation rates were not adjusted for

temperature.

2.4 Chlorophyll, phytoplankton and particulate matter

We measured the chlorophyll-*a* concentrations according to Wasmund, Topp & Schories

(66). Briefly, 2-4 L of lake water were filtered through 47 mm glass fibre filters (GF55,

Hahnemühle), which were directly transferred to 15 mL plastic tubes. Five mL ethanol

(>90 %) were added to the samples, followed by 10 min cold ultrasonification. The

samples were stored at 5 °C overnight and sterile-filtered (0.2 µm) the following

morning. The extracts were measured on-board with a fluorometer (Turner Trilogy) and

calibrated against a chlorophyll-*a* standard (Lot# BCBS3622S, Sigma-Aldrich). Samples

and standards were always handled and processed in the dark. In-situ chlorophyll

fluorescence was calibrated against extracted chlorophyll-*a* samples and then used to

calculate depth-integrated chlorophyll-*a* stocks (0-125 m).

For estimating the phytoplankton abundances, 4-10 L of water were concentrated to 20

mL using a 10 µm plankton net and fixed with alkaline Lugol solution. At TAFIRI

Kigoma, phytoplankton cells were counted from 2 mL subsamples by inverted

microscopy (at ×400 magnification). For particulate organic matter (POM), 2-4 L lake

water was filtered through precombusted GF/F filters (nominal pore size 0.7 µm;

Whatman).

2.5 Zooplankton and Pleiodon spekii

Zooplankton was collected with vertical net hauls across the oxygenated water column

(0-150 m) at each pelagic station. We sampled different size fractions of the zooplankton

community using three different nets. For smaller zooplankton, we used 25 and 95 µm

nets with 0.03 and 0.02 m^2 mouth openings, respectively. A 250 µm net with a 0.28 m^2

mouth opening was used for larger, fast swimming species. We preserved all zooplankton

collected from the first haul in ethanol for taxonomic zooplankton community

assessment, while the individuals from the second haul were designated for stable isotope

analysis (only for samples from the 95 and 250 µm nets). At TAFIRI Kigoma, we

analyzed the zooplankton community composition of the ethanol-preserved samples by

compound microscopy (Leica Wild M3B) at x200 magnification. Additionally, we

picked living individuals of the long-lived, filter-feeding bivalve *Pleiodon spekii* at near-

shore habitats in water depths of 1.5-6 m by snorkelling. Bivalves were first euthanized

with an overdose of MS222. Then we sampled the foot using clean scalpels and forceps

and removed the mucous with tissues and deionized water.

2.6 Fish

At on-shore landing sites adjacent to our sampling stations, we obtained fish specimens

from fishermen, which usually fish within a 20 km radius from their landing sites. We

collected *Stolothrissa tanganicae*, *Limnothrissa miodon*, *Lates stappersii*, *Lates*

microlepis, *Lates mariae*, and *Lates angustifrons* and processed them according to the

standard protocol described in Junker et al. (35). For stable isotope analysis, we sampled

the dorsal muscle using clean scalpels and forceps and removed the skin.

2.7 Isotopic and elemental analysis of solids

All solid isotope samples (POM, zooplankton, *P. spekii*, and fish) were oven-dried at ~60

°C for at least 24 h after collection and then packed in aluminium foil or small sample

tubes. Dried samples were stored at room temperature and shipped to Switzerland. At

Eawag Kastanienbaum, we fumed the POM samples for 48 h under HCl atmosphere to

remove inorganic carbon. Fish and *P. spekii* samples were ground to fine powder using a

Qiagen Tissuelyzer II. We measured the C and N elemental and isotopic compositions

with an EA-IRMS (vario PYRO cube, Elementar coupled with an IsoPrime IRMS, GV

Instruments). *Acetanilide #1* (Indiana University, CAS # 103-84-4) was used as an

internal standard. The isotopic ratios of the samples are reported in the delta notation

VPDB for carbon and air for nitrogen. Standard and sample reproducibility was generally

better than 0.2 ‰ for $\delta^{13}C$ and 0.5 ‰ for $\delta^{15}N$ and highest for fish tissue (0.1 ‰ for $\delta^{13}C$

and 0.2 ‰ for $\delta^{15}N$).

279   2.8 Lipid content of fish muscle tissue

For a subsample of *Stolothrissa* individuals, which exhibited the largest range in C:N

ratios, we measured the total lipid content to test whether a high C:N ratio effectively

translates to a higher amount of lipids in fish tissue. Total lipid content was determined

gravimetrically following Folch, Lees & Sloane-Stanley (67). In brief, ~1 mg of dried

fish muscle powder was weighed into a pre-combusted glass vial, and 1 mL of 2:1

(vol:vol) dichloromethane:methanol solution was added. The sample was then

ultrasonicated for 10 min. The supernatant was transferred to another pre-combusted, pre-

weighed glass vial and evaporated in a heat block. The entire procedure was repeated two

more times, and the resulting dry lipid mass weighed to the nearest 0.001 mg.

2.9 Data analysis

For calculating the depth-integrated isotopic values of POM, we normalized each sample for the phytoplankton abundance at the respective depth. We corrected the $\delta^{13}C$ of non-lipid-extracted animal tissue for its lipid content according to Post et al. (54). We estimated lipid content in fish tissue according to the theoretical model from the same study. For comparing the isotopic composition and the C:N ratios of fish populations between different regions, we selected individuals from stations where samples were available from both campaigns (north: stations 1 and 2; south: stations 7 and 9). From this subset, we additionally selected the individuals from the 50 mm size range with the highest overlap across regions and sampling campaigns for each species to minimize size-specific effects (Figs. S1; S2; 62). The chosen size ranges were 40-90 mm for *Stolothrissa*, 75-125 mm for *Limnothrissa*, and 200-250 mm for *Lates stappersii*. For the same analyses and reasons, we selected *P. spekii* from the exact same sites, whereas for zooplankton we used stations 1-3 (north) and 7-9 (south), because zooplankton samples were available from all stations, but we had only one sample per station. The Bayesian ellipses in the isotopic space were calculated using the R package SIBER (69).

3. Results

3.1 Biogeochemistry and hydrodynamic conditions in Lake Tanganyika during sampling

We observed strong differences in water column stratification and biogeochemistry in Lake Tanganyika between the two sampling campaigns and among basins (Fig. 2). Our sampling campaign at the end of the dry season (September/October) exhibited southerly trade winds that are characteristic for the wind-driven upwelling and mixing period (Fig. 1 c, d). By contrast, the winds were mostly calm and it was rainy at the end of the rainy season (April), which is typical for this time (Fig. 1 b). Compared to Apr/May, we find significantly lower Sc and N^2 values as well as lower surface temperatures ($\Delta\sim0.5$ °C) in Sep/Oct, indicating that the lake was less stratified during that period (Fig. 2 a-d; Sc: 0.4-0.7 versus 0.5-1.1 kJ m^{-2}; N^2: 1.5-2.7 versus 2.4-5.0 · 10^{-4} s^{-2}; Mann-Whitney-U Tests, p < 0.05). However, during both campaigns Sc values of the 50-100 m interval decreased towards the south, indicating some degree of upwelling/mixing (Fig. 2 a, b). At the end of the rainy season, the southern stations were sampled in May during the onset of the trade winds. The thermocline was not as heavily uplifted in the southern basin in Apr/May compared to Sep/Oct, when no thermocline had formed at station 9 (Fig. 2 c, d). This is a good indicator that wind-driven upwelling/mixing was not as pronounced in Apr/May. The oxygen distribution closely followed the thermal structure of the lake (Fig. 2 e, f). In Sep/Oct, the more stratified water column in the northern basin exhibited a shallower oxycline (50-70 m at stations 1-3), defined here as a sharp drop in oxygen concentrations, compared to the wind-driven mixing in the south, which introduced more oxygen to

deeper layers of the water column (up to 5 mg L^{-1} at 113 m at station 9; Fig. 2 e). In

Apr/May, the oxycline was lying relatively shallow across the full length of the lake from

70-120 m; oxygen concentrations were roughly 2-fold lower in the deep waters of the

southern basin compared to the well-mixed conditions in Sep/Oct (~2.3 mg L^{-1} at 113 m

at station 9; Fig. 2 f).

We also observed strong latitudinal changes in the nutrient distribution in Sep/Oct.

Moving from north to south, the position of the nitrate maximum in the water column

deepened from 67 m to 137 m, paralleling variation of the vertical phosphate gradients

(Fig. 2 g, i). The surface waters in the southern basin were also associated with generally

higher nitrate and especially phosphate concentrations. For instance, at some stations in

the south, surface water nitrate and phosphate concentrations reached up to 2.3 µM,

whereas they were <0.5 µM at most other stations. By contrast, in Apr/May, nitrate and

phosphate were more uniformly distributed across the north-south transect, and the nitrate

maxima were positioned at approximately 100 m across the lake (Fig. 2 h, j). A local

nitrate maximum (4.7 µM) at 40 m at station 3 in Apr/May may have been caused by an

overlying cyanobacterial bloom (56). The upward tilting thermocline in the south

enhanced the nutrient transport to the productive surface waters during both campaigns.

However, the tilting of the thermocline as well as surface nitrate and phosphate

concentrations in the south, were lower in Apr/May than in Sep/Oct (Fig. 2 c, d, g, h, i, j).

Fig. 2: Physical and chemical properties of Lake Tanganyika along our north-south transects (from

station 1-9) at the end of the dry season (left) and the end of the rainy season (right). (a,b) Schmidt

stability (Sc) of the 50-100 m depth interval and buoyancy frequency of the primary thermocline (N^2).

Distribution of (**c,d**) temperature (T), (**e,f**) dissolved oxygen, (**g,h**) nitrate, and (**i,j**) phosphate. The solid

white line depicts the thermocline, whereas the dashed white line represents less pronounced secondary

thermoclines. No clear thermocline had formed at station 9 at the end of the dry season. Samples are

indicated by vertical lines (continuous profiles) or points (discrete samples).

3.2 Concentration and isotopic composition of DIC and CO₂ fixation rates in April/May

In Apr/May, we determined the DIC concentration (stations 2, 5 and 7) and C isotopic composition (stations 1, 3 and 8) in the north and the south. Like other biogeochemical parameters, neither the concentration, nor the $\delta^{13}C$ of DIC showed strong latitudinal trends in Apr/May (Fig. 3), but both do exhibit pronounced vertical gradients. The DIC concentration varied between 70 and 72 mg C L^{-1} in the productive upper 50 m, followed by a sharp increase to about 74 mg C L^{-1} that flattened out with increasing depth. Concurrently, the $\delta^{13}C$-DIC values reached their maximum near the surface (2.1 ‰), decreased to 0.8 ‰ at the boundary between the metalimnion and the upper hypolimnion (100-150 m) and showed a further trend to slightly heavier values at greater depth.

Fig. 3: Distribution and isotopic composition of dissolved inorganic carbon (DIC) at the end of the rainy season (Apr/May 2018) in Lake Tanganyika. (**a**) DIC concentration profiles from stations 2, 5, and 7. (**b**) $\delta^{13}C$-DIC profiles from stations 1, 3, and 8.

In addition, we performed CO_2 fixation experiments at station 2 in the north and station 7

in the south (Fig. 4). At both stations, CO_2 fixation rates were highest in the euphotic

zone. CO_2 fixation rates in the top 50 m were higher at station 7 in the upwelling-driven

south (0.58-1.60 µM C d^{-1}), compared to station 2 in the permanently stratified north

basin (0.12-0.48 µM C d^{-1}).

Fig. 4: Experimentally determined CO_2 fixation rates relative to the oxygen and chlorophyll-*a*

distributions. (**a,c**) Oxygen and in-situ chlorophyll-*a* as well as (**b,d**) CO_2 fixation rates from stations 2

(**a,b**) and 7 (**c,d**) in Apr/May.

3.3 Particulate matter, phytoplankton, and zooplankton

The abundance of medium- to large-celled phytoplankton (>10 µm) decreased

significantly from north to south by 2.6 and $4.4 \cdot 10^8$ ind. m^{-2} per degree latitude (linear

regression, $p < 0.05$) in Sep/Oct and Apr/May, respectively (Fig. 5 a, b). Chlorophyll-*a*

was slightly higher in Sep/Oct compared to Apr/May (on average 46 versus 41 mg chl-*a*

m^{-2}), and generally showed the highest values in the south during both campaigns (Fig. 5

a, b).

We observed strong differences in δ^{13}C-POM between the two campaigns (Fig. 5 c, d). In

Sep/Oct, the values of δ^{13}C-POM varied between -26.5 and -24.9 ‰, whereas they

spanned from -32.3 to -30.1 ‰ in Apr/May. During both seasons, we observed an

increase from north to south of ~0.3 ‰ (linear regression, $p < 0.1$) and ~0.7 ‰ (linear

regression, $p < 0.01$) per degree latitude in Sep/Oct and Apr/May, respectively. The δ^{15}N

of POM was substantially higher in Sep/Oct, with an average of 1.5 ± 1.0 ‰ (Fig. 5 c). In

Apr/May, values were lower (mean: 0.2 ± 1.3 ‰) and increased from the north (station 3:

-1.0 ‰) to the south (station 9: 2.7 ‰; Fig. 5 d).

The patterns in zooplankton abundance and δ^{13}C were different from those observed in

phytoplankton and POM (Fig. 5 e, f). The zooplankton abundances were up to one order

of magnitude higher in Sep/Oct versus Apr/May, whereby this temporal change was less

pronounced in the north and center of the lake. In Sep/Oct, zooplankton abundances

reached a lake-wide maximum in the south (29.9 and $1.9 \cdot 10^3$ ind. m^{-2} for the 95 µm and

250 µm size fractions, respectively) and a secondary peak of large zooplankton (250 µm)

Fig. 5: Different plankton parameters sampled across the epi- and metalimnion (0-125 m) along the

north-south transects at (left) the end of the rainy season and (right) the end of the dry season. (a,b)

Depth-integrated phytoplankton abundances (>10 μm) and chlorophyll-*a* stocks. (c,d) Depth-integrated and

phytoplankton abundance-weighed $\delta^{13}C$ and $\delta^{15}N$ of POM. (e,f) Depth-integrated abundances of the 95 μm

and 250 μm size fractions of the zooplankton community. (g,h) $\delta^{13}C$ and (i,j) $\delta^{15}N$ of the 95 μm and 250 μm

size fractions of the zooplankton community.

in the northern part of the lake ($1.3 \cdot 10^3$ ind. m^{-2}). In Apr/May, the abundances from both

net types varied only slightly across the lake, reaching up to 2.0 and $0.5 \cdot 10^3$ ind. m^{-2} for

the 95 µm and 250 µm size fractions, respectively (Fig. 5 f). Zoo- and phytoplankton

abundances were negatively correlated in Sep/Oct, i.e. phytoplankton (>10 µm)

decreased, when zooplankton increased (rho = -0.58, $p \sim 0.1$, Spearman's rank

correlation). No significant correlation between phyto- (>10 µm) and zooplankton was

found in Apr/May (rho = 0.38, $p > 0.3$, Spearman's rank correlation).

The zooplankton δ^{13}C values changed slightly between the seasons. In Apr/May, both

size fractions were ~0.6 ‰ heavier compared to zooplankton from Sep/Oct, opposite of

what we found in POM. Overall, zooplankton δ^{13}C lie between -24.7 and -21.1 ‰ (Fig. 5

g, h). In Sep/Oct, we observed a trend towards higher values in the south (Δ +1 ‰ for

both net types), whereas the highest values in Apr/May occurred towards the northern (-

22.1 ‰) and southern (-21.1 ‰) extremities of the lake, with a minimum of -24.0 ‰ in

the central region. The zooplankton δ^{15}N values tended to be higher in the 250 µm

compared to the 95 µm fraction. Overall, they were similar between the campaigns, but

showed generally lower values in the north (2.5-6.3 ‰) compared to the center and the

south (3.2-10.4 ‰; Fig. 5 i, j).

The zooplankton community composition was dominated by copepods (Fig. S3). In

Sep/Oct, cyclopoids dominated the small size fraction (on average 66 and 78 % for 25

and 95 µm, respectively), whereas calanoids were prevalent in the 250 µm fraction

(mean: 74 %). In Apr/May, calanoid copepods dominated the zooplankton community

throughout all size fraction, but the relative abundance of cyclopoids was highest in the

small size fractions here, too (up to 78 % in the north basin). The freshwater jellyfish

Limnocnida tanganyicae medusa appeared in higher relative abundances in Sep/Oct,

especially in the 250 μm size fraction, reaching up to 19 % at station 7. The contributions

of shrimps to the total zooplankton community was low in our samples, reaching its

highest values in the south during both seasons (max. 16 %). Fish and insect larvae were

rare in all our samples (<1 %).

Overall, we find that the phytoplankton abundances (>10 μm) decreased towards the

south, whereas chlorophyll-*a* and zooplankton reached the highest values in the south.

Zooplankton $\delta^{13}C$ and $\delta^{15}N$ were variable, but also showed the highest values in the

south. The southward increase was more expressed in POM $\delta^{13}C$ and $\delta^{15}N$. In addition,

POM $\delta^{13}C$ and $\delta^{15}N$ were generally heavier in Sep/Oct compared to Apr/May.

3.4 Isotopic and elemental composition of bivalve and fish populations

3.4.1 Variations in $\delta^{13}C$ and $\delta^{15}N$

The $\delta^{13}C$ of *P. spekii* revealed a consistent seasonal and latitudinal dynamic (Fig. 6 a, b).

We found a significant divergence between northern and southern populations in Sep/Oct

(mean -22.5 and -21.4 ‰, respectively; Mann-Whitney U test, $p < 0.01$), whereas the

$\delta^{13}C$ values of the *P. spekii* samples from the two basins overlapped in Apr/May (means

of -22.7 and -22.3 ‰, respectively), with northern samples nesting fully within the range

of southern samples. The samples from the northern basin showed little variation between

the seasons, whereas samples collected in the southern basin differed significantly

(Mann-Whitney U test, $p < 0.001$).

In Sep/Oct, the mean δ^{13}C values of the populations of the largely planktivorous sardines

Stolothrissa and *Limnothrissa* as well as the zooplankti- and piscivorous *Lates stappersii*

diverged significantly (Mann-Whitney U tests, $p < 0.001$) by approximately 0.7 ‰

between the north and the south (Fig. 6 c, e, g). By contrast, the differences in mean

values were completely erased or going slightly into the opposite direction in Apr/May

(Fig. 6 d, f, h).

Moreover, the southern populations of all three species exhibited the highest δ^{13}C with

mean values of -20.7 ‰, -20.3 ‰, and -20.4 ‰ for *Stolothrissa*, *Limnothrissa,* and *Lates*

stappersii, respectively. By contrast, the δ^{13}C of the northern population in Sep/Oct was

similar to the population-wide averages from both basins in Apr/May and never differed

by more than 0.4 ‰. The lowest mean values were consistently observed in Apr/May (-

21.5 ‰, -21.0 ‰, and -21.2 ‰ for *Stolothrissa*, *Limnothrissa,* and *Lates stappersii*,

respectively).

In contrast to *P. spekii* and *Stolothrissa*, the southern populations of *Limnothrissa* and

Lates stappersii showed 0.2 and 0.4 ‰ lower δ^{13}C values compared to the northern one

in Apr/May, respectively. This difference was not significant for *Limnothrissa* (Mann-

Whitney U test, $p > 0.05$) and was significant for *Lates stappersii* (Mann-Whitney U test,

$p < 0.001$). It is worth noting that samples of *Limnothrissa* and *Lates stappersii* from

Apr/May included in our analysis were slightly unbalanced, with samples from the north

being larger compared to samples from the south (Fig. S2). Since larger individuals in

these two species tend to have less depleted δ^{13}C values (Fig. S4 c, e), the δ^{13}C values of

the northern populations, and with them the difference between the basins, are slightly

overestimated here.

Fig. 6: Carbon (normalized for C:N mass ratio according Post et al. (54)) and nitrogen stable isotope

signatures of the major pelagic food web members. (**a,b**) the bivalve *Pleiodon spekii* as well as the fish

(**c,d**) *Stolothrissa tanganicae*, (**e,f**) *Limnothrissa miodon*, and (**g,h**) *Lates stappersii* at the end of the dry

season (left) and the end of the rainy season (right). Orange dots represent the northern basin (stations 1 and

2) and blue dots represent the southern basin (stations 7 and 9). Numbers indicate the mean δ^{13}C (black)

and δ^{15}N (grey) of a population and Δ denotes the δ^{13}C difference between the southern and northern

populations. Significant differences in δ^{13}C are marked by stars. Ellipses encompass approximately 95 % of

the data of each population. Note different y-axis limits.

We had fewer samples of the large predators *Lates microlepis*, *Lates mariae*, and *Lates*

angustifrons, preventing an in-depth statistical analysis, but the results hint at similar

patterns. Across our entire data set, these three species showed the highest $\delta^{13}C$ values,

with most observations being heavier than -21 ‰ (Figs. S4 g, i. k; S5). In Sep/Oct, the

$\delta^{13}C$ of both *Lates microlepis* and *Lates angustifrons* specimens were >0.5 ‰ lighter than

the individuals from the southern populations. In Apr/May, samples from both the north

and south basins were only available for *Lates mariae*. Here, the $\delta^{13}C$ values from both

basins varied within the same range and their averages differed only slightly (north: -20.1

‰; south: -20.3 ‰).

Similar to zooplankton and POM *P. spekii* showed consistently higher $\delta^{15}N$ values in the

southern basin, reaching 1.7 and 2.1 ‰ on average in Sep/Oct and Apr/May, respectively,

compared to the northern basin with averages of 1.5 and 1.3 ‰, for the two campaigns

(Fig. 6 a, b). In contrast to $\delta^{13}C$, we observed no systematic seasonal or regional

differences in fish $\delta^{15}N$ (Figs. 6 c-h; S5). *Stolothrissa* exhibited the lowest values with

population averages spanning from 4.9 to 5.1 ‰. The other sardine, *Limnothrissa*, had

markedly higher values (means: 5.4-5.6 ‰). Most observations ranged between 4-6 ‰

and 4.5-6.5 ‰ for the two species, respectively (Fig. S4 b, d), whereby the variability

appeared to be neither related to site nor season, and only to a small extent to size. By

contrast, the $\delta^{15}N$ of the larger *Lates* species were primarily related to size (Fig. S4 f, h, j,

l). Individuals smaller than 200-250 mm showed an increase from <4 ‰ up to around 6-8

‰, which then flattened out at these high values for specimens >250 mm. In the >250

mm size class, the $\delta^{15}N$ of *Lates stappersii* and *Lates microlepis* never exceeded 7.5 ‰,

whereas *Lates mariae* and *Lates angustifrons* reached values of >8 ‰ (Fig. S4 f, h, j, l).

None of the *Lates* species revealed clear population-wide differences in $\delta^{15}N$ between the

basins or seasons (Figs. 6 g, h; S5).

Overall, we find that the different fish species and *P. spekii* revealed congruent patterns

in $\delta^{13}C$ with diverging northern and southern populations when upwelling/mixing in the

south were strongest, i.e. at the end of the dry season in Sep/Oct. By contrast, $\delta^{13}C$ values

converged, and were generally lower, when the lake was more heavily and homogenously

stratified, i.e. at the rainy season-dry season transition (Apr/May). On the contrary, the

fish populations exhibited no clear basin-scale trends in $\delta^{15}N$, whereas *P. spekii* showed

slightly higher values in the south during both seasons.

3.4.2 C:N ratios and estimated lipid content

We observed strong changes in C:N ratios between Sep/Oct and Apr/May at lower and

middle trophic levels, but not at high trophic levels; there were no clear differences

between north and south in all organisms (Fig. 7). *Pleiodon spekii* and zooplankton

showed consistently lower C:N ratios in Sep/Oct compared to Apr/May (Fig. 7 a, c, e),

but the low sample size (n ≥ 3) ruled out a statistical analysis. We observed the largest

seasonal change in zooplankton (250 µm) from the southern basin. The medians differed

between 5.1 (Sep/Oct) and 6.8 (Apr/May). The differences were smaller in *P. spekii*,

revealing an overall minimum median of 3.9 in the northern basin in Sep/Oct and a

maximum of 4.3 in the southern basin in Apr/May.

Fig. 7: Mass C:N ratios of primary consumers (left) and fish tissue (right) for the different sampling

campaigns and basins of Lake Tanganyika. (**a**) *Pleiodon spekii*, (**c**, **e**) zooplankton, (**b**) *Stolothrissa*

tanganicae, (**d**) *Limnothrissa miodon*, and (**f**) *Lates stappersii*. Stars indicate significant differences

between campaigns and numbers depict the % change in estimated lipid content according to Post et al.

(54). Note varying y-axis scaling.

Contrary to the trends in *P. spekii* and zooplankton, the sampled fish species showed a

tendency towards higher C:N ratios in Sep/Oct (Fig. 7 b, d, f). This trend vanished with

increasing trophic level of the fish species. The C:N ratios of the planktivorous clupeid

Stolothrissa decreased significantly (Mann-Whitney U tests, $p < 0.001$) between Sep/Oct

and Apr/May in both the northern and southern basins (medians of 3.65 versus 3.24 and

3.49 versus 3.19, respectively). *Limnothrissa* revealed significant changes in the north

(Sep/Oct median: 3.23; Apr/May median: 3.18; Mann-Whitney U test, $p < 0.001$),

whereas the pattern was reversed and insignificant in the south (Sep/Oct median: 3.19;

Apr/May median: 3.21; Mann-Whitney U test, $p > 0.1$). By contrast, the C:N ratios of

Lates stappersii varied only within a narrow range, with medians spanning from 3.18 to

3.21 and no significant differences between seasons (Mann-Whitney U tests, $p > 0.1$).

Using the model from Post et al. (54), we estimated the lipid contents in the dorsal

muscle tissue of the investigated fish species from their C:N ratios. This analysis suggests

a reduction of lipid content in *Stolothrissa* from Sep/Oct to Apr/May by 43 and 45 % in

the north and south, respectively. *Limnothrissa* exhibited a 13 % decrease (north) and a 5

% increase (south). *Lates stappersii* showed the lowest differences with a 9 % increase

(north) and 1 % decrease (south). A gravimetric determination of lipid content from

selected *Stolothrissa* samples confirmed that higher C:N ratios translate into higher lipid

contents (linear regression, $R^2 = 0.91$, $p < 0.01$, $n = 5$; Fig. S6).

4 Discussion

4.1 Effect of upwelling and mixing on the isotopic composition of the

planktonic food web

Upwelling and convective mixing moderate the transport of nutrients to the surface

waters, and thus drive biological productivity during the dry season in Lake Tanganyika.

In this study, we compare two contrasting hydrodynamic situations: First, the period of

re-establishing water column stratification at the end of the dry season (Sep/Oct). During

this time, stratification was weaker and the thermocline was still absent at the

southernmost station, enabling particularly high nutrient fluxes in the south. Second, the

period of lake-wide stratification at the rainy season-dry season transition (Apr/May).

Here, the water column experienced stronger stratification from the preceding rainy

season and beginning trade winds initiated the upwelling in the south, resulting in overall

lower nutrient fluxes with a maximum in the south. Our data show that upwelling and

mixing do not only influence biological productivity, but also impact the isotopic

composition of all food web members, likely owing to differences in primary

productivity and N acquisition pathways of phytoplankton. Such systematic changes can

be used to delineate regional fish populations.

In contrast to the southward increase in nutrient availability, we detected a southward

decrease in the abundance of large-sized phytoplankton (>10 μm) a pattern that has

previously been observed in Lake Tanganyika (70), pointing out additional ecological

controls, such as zooplankton grazing or competition within the phytoplankton

community. In Lake Tanganyika, the nano- and pico size fractions (<10 μm) are more

competitive under nutrient-rich conditions and therefore dominate the phytoplankton

community in south (71,72). As a result of their high densities, total phytoplankton

abundance and biomass is generally highest during the dry season upwelling in the south

(18,21,73).

In addition, upwelling/mixing and the subsequent stimulation of primary productivity,

place important bottom-up control on the abundances of zooplankton in Lake

Tanganyika. This was evidenced by high zooplankton abundances in the dry season with

maxima in the southern basin (Fig. 5; 18,70), which in turn sustain the growth of the

pelagic fish populations, especially the sardine species (4,23,24,75). The high

zooplankton abundances can in turn exert top-down control over phytoplankton, which is

indicated by the negative correlation between the phytoplankton (>10 μm) and

zooplankton abundances in Sep/Oct. This grazing effect may have also been responsible

for the absence of strong differences in chlorophyll-*a* between Sep/Oct and Apr/May.

Zooplankton abundance, and thus grazing pressure, exhibited no clear latitudinal trend

during the April/May campaign, when the lake-wide stratification was stronger, and the

north-south gradients in nutrient availability and biological productivity were not as

pronounced.

The varying hydrodynamics were also associated with distinct δ^{13}C-POM signatures that

may reflect differences in primary productivity. The average δ^{13}C was ~5 ‰ heavier in

Sep/Oct compared to Apr/May, and the δ^{13}C increased by ~2 ‰ from north to south

during both campaigns (Fig. 5 c, d). In Lake Tanganyika, previous studies also revealed

heavier δ^{13}C-POM values in the dry season (51–53), even though the differences were

smaller (max. 3.1 ‰) than in our study (max. 6.7 ‰), possibly due to the varying timing

of the sampling. These differences in δ^{13}C-POM likely reflect the well-documented

changes in primary production (19,73,76), where heavier isotopic signatures in POM

mirror the incorporation of a larger ^{13}C fraction by higher photosynthesis and cell growth

rates (44,45) and a stronger drawdown of the DIC pool (46–48,77). The links between

stratification, vertical nutrient supply, and primary productivity in Lake Tanganyika are

well established (10–12,19) and several studies have used the δ^{13}C of sediment POM to

infer primary productivity (7,12). Our own CO_2 fixation rate measurements, done during

Apr/May, show evidence for higher productivity rates in the southern basin at station 7 in

the south compared to station 2 in the north (Fig. 4). On the other hand, upwelling of

intermediate waters will not only supply nutrients, but also isotopically light DIC

(depleted by ~1 ‰; Fig. 3; 59,75). Although this mechanism will slightly dilute ^{13}C

enrichment, our proposed mechanism of higher primary productivity is apparently strong

enough to overcome this depletion in δ^{13}C-POM, ultimately leading to higher δ^{13}C-POM

when upwelling/mixing is stronger.

The analogous pattern in δ^{15}N-POM implies that upwelling and mixing may have

influenced the N sources of primary producers, where lighter values are typically

interpreted as inputs from N fixation (77,79). POM δ^{15}N values increased from -1.0 ‰ at

station 3 to 2.7 ‰ at station 9 in the south in Apr/May, concurrent with a decrease in

filamentous, N-fixing cyanobacteria (56), whereas it fluctuated with slightly higher

values (-0.3-2.6 ‰) in Sep/Oct devoid of a latitudinal or phytoplankton composition

related pattern. We also observed no correlation between the presence of surface nitrate

and δ^{15}N-POM, which may have induced fractionation effects during nitrate-uptake.

When free nitrate remains, the phytoplankton community does not represent a complete

sink of the upward diffusing nitrate, i.e. the residual nitrate should be isotopically heavy

and phytoplankton relatively light. In line with the higher density of N-fixing

cyanobacteria (21,56,72), the generally lighter δ^{15}N-POM in Apr/May (Δ-1.4 ‰) point at

higher inputs from N fixation compared to Sep/Oct, when nutrient fluxes are higher due

to upwelling/mixing.

The isotopic composition of the zooplankton community did not show clear latitudinal

and seasonal patterns, pointing at additional influencing factors than the signal from the

base of the food web, i.e. POM. The δ^{13}C values in our zooplankton samples oscillated

between -24.7 and -21.1 ‰, in accordance with previous isotope surveys. The δ^{15}N

values from the north were also in agreement with other studies, whereas the maxima in

633 the south, where previously no isotopic characterization of the food web was undertaken,

634 exceeded earlier reports by min. 2.7 ‰ (Fig. 5; 45–47,62). The high intra-basin

635 variability in $\delta^{13}C$ and $\delta^{15}N$ as well as the high absolute values relative to other members

636 of the food web, with some zooplankton $\delta^{15}N$ exceeding top predator fish $\delta^{15}N$ values,

637 may be in part attributable to varying zooplankton community compositions (52,68).

638 Such values from a pooled zooplankton sample are not unexpected, because zooplankton

639 communities consist usually of members from several trophic levels (e.g. 46,77,78), and

640 our samples represent batch samples from entire zooplankton communities formed by

641 many different species, genera, and families. In addition, the zooplankton community is

642 notoriously hard to sample and standard netting techniques do not capture fast swimmers

643 such as shrimps efficiently, therefore often underestimating their abundances (26).

644 Indeed, shrimps only made up minor proportions in our samples (Fig. S3) and previous

645 work showed that they have $\delta^{13}C$ and $\delta^{15}N$ values lower than our community isotope

646 values (52,53,68). However, reported $\delta^{15}N$ values of individual zooplankton taxa,

647 including detrivorous jellyfish and fish larvae, do not exceed 5.9 ‰ (52,53,68) and

648 therefore fail at explaining the high $\delta^{15}N$ in our measured community isotope samples

649 from the south (>10 ‰). In line with our results, earlier reports of bulk community

650 samples found high $\delta^{15}N$ values between 6 and 8 ‰ (51), raising questions about the

651 utility of using bulk community samples. Combining the taxonomic assessment of the

652 community with an isotopic characterization of individual zooplankton taxa would thus

653 be valuable in future food web studies.

654 In summary, our results point to a pivotal role of nutrient upwelling and mixing for

655 sustaining the high biological productivity in the south basin during the dry season.

Upwelling-related increases in primary productivity and decreases in N fixation likely

resulted in markedly heavier planktonic δ^{13}C and slightly heavier δ^{15}N values in the south

(Fig. 8 a). The slightly higher zooplankton δ^{13}C and δ^{15}N values in the southern basin

may reflect the isotopic imprint of the upwelling/mixing, but a clear north-south trend

may be masked to some extent by concomitantly shifting community composition effects.

4.2 Isotopic imprints from upwelling and mixing reveal regional fish

populations

The seasonally and regionally varying hydrodynamic conditions also determine the C

isotopic compositions of organisms higher in the food web, through the incorporation of

phyto- and zooplankton prey. Due to the longevity of many organisms higher in the food

web, they integrate the isotopic signals from their food over a longer time span. For

instance, tissue turnover in bivalves is significantly slower than in phyto- and

zooplankton (51,82). Therefore, we used the filter-feeding bivalve *P. spekii* as a reference

organism for the seasonal phytoplankton isotopic signals. *Pleiodon spekii* lives for at

least five years at a stationary location and was successfully used to record upwelling

events in Lake Tanganyika (83). We find that the southern *P. spekii* samples in Sep/Oct

diverge significantly in δ^{13}C from northern samples (Δ1.1 ‰). Lake-wide stratification

over the rainy season on the other hand resulted in converging δ^{13}C values between

samples from both basins in Apr/May (Fig. 6 a, b).

Such seasonal cycles in δ^{13}C signals are common in aquatic food webs and propagate

along the trophic chain (50,84). The typical muscle half-life ranging from a few weeks in

juvenile fish to several months in adults explains the incorporation of seasonal dietary

isotope patterns (85–87). Accordingly, we find similar patterns as in *P. spekii* in the three

major pelagic fish species of Lake Tanganyika, with a latitudinal difference of ~0.7 ‰

heavier fish tissue δ^{13}C in the south (Fig. 6). The aligning δ^{13}C values between northern

and southern samples during the rainy season are again in agreement with the more

uniform primary productivity patterns. Our results point to fish populations confined to

regional foraging grounds in the respective basins, which therefore record the latitudinal

isotope gradients (Fig. 8 b).

However, our previous high resolution population genetic work did not find evidence for

genetic differentiation between the north and south basins in any of the six fish taxa

investigated in this study. Instead, populations in the north and south basins are closely

related (34–36). The limited genetic differentiation in these species is not spatially

restricted, with the exception of a case in *Lates mariae*. In Rick et al. (36), we found one

Lates mariae cluster confined to the extreme south end of the lake with strong genetic

differentiation from individuals elsewhere in the south basin or in the rest of the lake.

Thus, the genetic structure of the fish populations cannot be explained by the basin-scale

dynamics. This implies that the degree of geographical isolation between north and south

basins itself is insufficient to suppress lake-wide gene flow in these pelagic fish species.

In other words, the isotopically distinct fish populations can only be regarded as regional

on rather short seasonal to multiannual time scales (Fig. 8 c).

Fig. 8: Sketch synthesizing the main conclusions and hypotheses of the study. (**a**) Biological

productivity of phyto- and zooplankton based on abundance and δ^{13}C data, which were used to (**b**) infer the

distribution of regional fish populations from their δ^{13}C signatures. (**c**) The regional isolation of the fish

populations is apparently insufficient to suppress lake-wide gene flow. (**d**) The regional fish populations as

well as different genetic clusters did not exhibit systematic differences in δ^{15}N. (**e**) The clupeid *Stolothrissa*

exhibited strong seasonal changes in C:N, i.e. lipid content, indicating lipid storage after the productive dry

season.

*results from Junker et al. (35) and Rick et al. (36)

4.3 Regional fish populations show no differences in δ^{15}N or C:N ratios

Despite the absence of pronounced spatial genetic structure in either of the sardine

(34,35) or *Lates* species (36), regional fish populations may exhibit phenotypic changes

in diet or lipid content in response to regionally different environments, which include a

northern region with a more stable and clear water column and a plankton-rich upwelling

region in the south (15,18,75).

The rather constant average δ^{15}N of the studied fish (deviation < 0.2 ‰) and *P. spekii*

(deviation < 0.8 ‰) among the sampling campaigns and basins indicates no strong

differences in trophic level between the populations in the different lake basins. However,

the difficulty of quantifying the trophic position of the fish species was exacerbated by

the small differences between the trophic levels. Moreover, the zooplankton community

δ^{15}N exhibited strong intra-basin variability, with maxima similar to the highest fish δ^{15}N

values, which raises doubt about the usefulness of comparing bulk community with tissue

samples (88). Both the high variability in δ^{15}N between individual zooplankton taxa

(52,53,68) and unknown trophic discrimination factors (89), which appear to deviate

from the norm in Lake Tanganyika (52,68), further aggravate assessing subtle differences

in the specific diets. Compound specific isotope analyses of amino acids may further help

constraining the trophic relationships (90).

Nonetheless, on a basin-scale, the relatively consistent fish and *P. spekii* δ^{15}N

demonstrate that the isotopic composition of their N sources does not vary substantially

throughout the year and among basins, whereas POM and zooplankton showed some

tendency towards higher values in the south. Instead, larger δ^{15}N variations of up to >2

‰ among individuals of a similar size found at the same location and time indicate that

other factors than the basin-scale hydrodynamics influence the diets of the studied fish

taxa. We found no clear evidence, however, that differences in $\delta^{15}N$ were linked to the

different genetic clusters in *Limnothrissa* or the four *Lates* species (Figs. S7; 8 d).

Fish use lipids to store energy during times of abundant food supply to bridge resource

limited periods (38,39). However, in congruence with the absence of basin-scale genetic

structure, we found no regional differences in C:N ratios as proxy for lipid content (Fig.

S6; 48,49,88), but we did find seasonal changes: the smallest species, *Stolothrissa,*

showed a significantly higher C:N ratio, i.e. lipid content, at the end of the productive dry

season (Fig. 7), which translates to >40 % change in lipid content according to the model

of Post et al. (54). Seasonal lipid cycling is expected to be more pronounced in smaller

fish, due to higher metabolic rates (92) and their planktivorous diet. While predators, i.e.

Lates stappersii and large *Limnothrissa*, feed on both fish and zooplankton, the solely

planktivorous *Stolothrissa* must cope with the strong seasonal fluctuations in plankton

productivity. Thus, *Stolothrissa* may have a life history adapted to building reserves

during the productive dry season for the following rainy season, when resources are less

abundant. Alternatively, the changing C:N may relate to spawning activities (93).

However, spawning peaks were reported to occur in September and April-July (94,95),

i.e. during both our sampling occasions (Sep/Oct and Apr/May), and can thus not explain

the observed changes in C:N between those two time points. The seasonal effect was less

pronounced in the slightly larger *Limnothrissa* and was clearly absent in *Lates stappersii*,

possibly due to their larger sizes and more piscivorous diets (Fig. 8 e). Overall, the $\delta^{15}N$

and C:N values indicate similar diets and lipid contents of the northern and southern fish

populations. We hypothesize that the long term gene flow across the lake may inhibit the

development of ecological differences among the regional fish populations in response to

the basin-scale environmental conditions.

5 Conclusions

In this study, we showed that the seasonal upwelling and mixing in the south basin of

Lake Tanganyika induce distinct isotopic imprints at the primary producer level. These

distinct isotopic signals can be tracked across the entire pelagic food web. Using $\delta^{13}C$ as

tracer, we identified fish populations with regional foraging grounds, implying some

degree of isolation on a seasonal and basin-wide scale. Correspondingly, regional fishery

management strategies may include basin-scale quotas. Our elemental and bulk isotopic

composition data provide no clear evidence for strong physiological or dietary

differences among these regional populations. Although lake-wide gene flow may inhibit

the evolution of clear ecological differences between the regional populations, a different

set of methods may reveal some ecological variation that cannot be resolved with bulk

elemental and isotopic analyses. In the context of assessing the vulnerability of Lake

Tanganyika's pelagic food web in a warming climate, our study indicates that the

economically relevant pelagic fish species are genetically adapted to the whole lake

although they form regional populations at the seasonal time scale.

Acknowledgements

We are grateful for the support from our research collaborators at the Tanzania Fisheries Research Institute, particularly the Directors Rashid Tamatamah and Semvua Mzighani as well as Mary Kishe. Special thanks go to Mupape Mukuli as well as the captain and crew of the *M/V Maman Benita* for their steady toil in organizing and conducting the cruise work with us. We also thank Andreas Brand, Kathrin B.L. Baumann, and Tumaini M. Kamulali for their help during field work, Serge Robert and Fabian Kuhn for assistance in the lab, and Eliane Scharmin for administrative support. Special thanks go to Jessica A. Rick for providing help in the field, the data of the *Lates* genetic clusters, and comments on the manuscript. Thanks to Blake Matthews for insightful discussions. This work was funded by the Swiss National Science Foundation (grant CR23I2-166589). Thanks to the Tanzania Commission for Science and Technology (COSTECH) for granting the research permits.

Author contributions

Conceptualization: Benedikt Ehrenfels, Julian Junker, Ismael A. Kimirei, Ole Seehausen, Catherine E. Wagner, Bernhard Wehrli

Data Curation: Benedikt Ehrenfels, Julian Junker, Demmy Namutebi, Cameron M. Callbeck, Anthony Kalangali, Athanasio S. Mbonde

Formal Analysis: Benedikt Ehrenfels, Julian Junker, Demmy Namutebi, Cameron M. Callbeck

Funding Acquisition: Julian Junker, Ismael A. Kimirei, Carsten J. Schubert, Ole Seehausen, Catherine E. Wagner, Bernhard Wehrli

Investigation: Benedikt Ehrenfels, Julian Junker, Cameron M. Callbeck, Christian Dinkel, Anthony Kalangali, Ismael A. Kimirei, Athanasio S. Mbonde, Julieth B. Mosille, Emmanuel A. Sweke

Methodology: Benedikt Ehrenfels, Julian Junker, Christian Dinkel, Anthony Kalangali, Ismael A. Kimirei, Athanasio S. Mbonde, Julieth B. Mosille, Emmanuel A. Sweke, Carsten J. Schubert, Ole Seehausen, Catherine E. Wagner, Bernhard Wehrli

Project Administration: Benedikt Ehrenfels, Julian Junker, Christian Dinkel, Anthony Kalangali, Ismael A. Kimirei, Athanasio S. Mbonde, Julieth B. Mosille, Emmanuel A. Sweke, Ole Seehausen, Catherine E. Wagner, Bernhard Wehrli

Resources: Ismael A. Kimirei, Emmanuel A. Sweke, Carsten J. Schubert, Ole Seehausen, Catherine E. Wagner, Bernhard Wehrli

Supervision: Carsten J. Schubert, Ole Seehausen, Catherine E. Wagner, Bernhard Wehrli

Validation: Benedikt Ehrenfels, Julian Junker, Demmy Namutebi, Cameron M. Callbeck, Anthony Kalangali, Athanasio S. Mbonde, Julieth B. Mosille, Emmanuel A. Sweke

Visualization: Benedikt Ehrenfels

812   **Writing – Original Draft Preparation:** Benedikt Ehrenfels

813   **Writing – Review & Editing:** Benedikt Ehrenfels, Julian Junker, Cameron M. Callbeck,

814   Ismael A. Kimirei, Carsten J. Schubert, Ole Seehausen, Catherine E. Wagner, Bernhard

815   Wehrli

References

818   1.   Mölsä H, Reynolds J, Coenen E, Lindqvist O. Fisheries research towards resource

819   management on Lake Tanganyika. Hydrobiologia. 1999;407:1–24.

820   2.   Van der Knaap M, Katonda KI, De Graaf GJ. Lake Tanganyika fisheries frame

821   survey analysis: Assessment of the options for management of the fisheries of

822   Lake Tanganyika. Aquat Ecosyst Heal Manag [Internet]. 2014;17(1):4–13.

823   Available from: http://www.scopus.com/inward/record.url?eid=2-s2.0-

824   84896478558&partnerID=tZOtx3y1

825   3.   Kimirei IA, Mgaya YD, Chande AI. Changes in species composition and

826   abundance of commercially important pelagic fish species in Kigoma area, Lake

827   Tanganyika, Tanzania. Aquat Ecosyst Heal Manag. 2008;11(1):29–35.

828   4.   Coulter GW. Lake Tanganyika and its life. British Museum of Natural History,

829   London and Oxford University Press, Oxford; 1991.

830   5.   Mölsä H, Sarvala J, Badende S, Chitamwebwa D, Kanyaru R, Mulimbwa M, et al.

831   Ecosystem monitoring in the development of sustainable sheries in Lake

Tanganyika. Aquat Ecosyst Heal Manag. 2002;5(3):267–81.

6. Sarvala J, Langenberg VT, Salonen K, Chitamwebwa DBR, Coulter GW, Huttula

T, et al. Fish catches from Lake Tanganyika mainly reflect changes in fishery

practices, not climate. Verhandlungen Int Vereinigung für Theor und Angew

Limnol. 2006;29:1182–8.

7. O'Reilly CM, Alin SR, Plisnier P-D, Cohen AS, McKee BA. Climate change

decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. Nature

[Internet]. 2003;424(6950):766–8. Available from:

http://dx.doi.org/10.1038/nature01833%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/

12917682

8. Cohen AS, Gergurich EL, Kraemer BM, McGlue MM, McIntyre PB, Russell JM,

et al. Climate warming reduces fish production and benthic habitat in Lake

Tanganyika, one of the most biodiverse freshwater ecosystems. Proc Natl Acad Sci

U S A. 2016 Aug;113(34):9563–8.

9. McGlue MM, Ivory SJ, Stone JR, Cohen AS, Kamulali TM, Latimer JC, et al.

Solar irradiance and ENSO affect food security in Lake Tanganyika, a major

African inland fishery. Sci Adv. 2020;6(41):1–9.

10. Verburg P, Hecky RE, Kling H. Ecological consequences of a century of warming

in Lake Tanganyika. Science. 2003;301(5632):505–7.

11. Verburg P, Hecky RE, Kling HJ. Climate warming decreased primary productivity

in Lake Tanganyika, inferred from accumulation of dissolved silica and increased

853 transparency. Int Vereinigung für Theor und Angew Limnol Verhandlungen.

854 2006;29(5):2335–8.

855 12. Verburg P. The need to correct for the Suess effect in the application of $\delta^{13}C$ in

856 sediment of autotrophic Lake Tanganyika, as a productivity proxy in the

857 Anthropocene. J Paleolimnol. 2007;37(4):591–602.

858 13. Tierney JE, Mayes MT, Meyer N, Johnson C, Swarzenski PW, Cohen AS, et al.

859 Late-twentieth-century warming in Lake Tanganyika unprecedented since AD 500.

860 Nat Geosci. 2010;3(6):422–5.

861 14. Plisnier PD, Chitamwebwa D, Mwape L, Tshibangu K, Langenberg V, Coenen E.

862 Limnological annual cycle inferred from physical-chemical fluctuations at three

863 stations of Lake Tanganyika. Hydrobiologia. 1999;407:45–58.

864 15. Verburg P, Antenucci JP, Hecky RE. Differential cooling drives large-scale

865 convective circulation in Lake Tanganyika. Limnol Oceanogr [Internet].

866 2011;56(3):910–26. Available from:

867 http://doi.wiley.com/10.4319/lo.2011.56.3.0910

868 16. Naithani J, Deleersnijder E, Plisnier PD. Analysis of wind-induced thermocline

869 oscillations of Lake Tanganyika. Environ Fluid Mech. 2003;3(1):23–39.

870 17. Stenuite S, Pirlot S, Hardy M-A, Sarmento H, Tarbe A-L, Leporcq B, et al.

871 Phytoplankton production and growth rate in Lake Tanganyika: evidence of a

872 decline in primary productivity in recent decades. Freshw Biol [Internet].

873 2007;52(11):2226–39. Available from: http://doi.wiley.com/10.1111/j.1365-

874   2427.2007.01829.x

875   18.   Bergamino N, Horion S, Stenuite S, Cornet Y, Loiselle S, Plisnier PD, et al.

876   Spatio-temporal dynamics of phytoplankton and primary production in Lake

877   Tanganyika using a MODIS based bio-optical time series. Remote Sens Environ.

878   2010;114(4):772–80.

879   19.   Mziray P, Kimirei IA, Staehr PA, Lugomela C V., Perry WL, Trolle D, et al.

880   Seasonal patterns of thermal stratification and primary production in the northern

881   parts of Lake Tanganyika. J Great Lakes Res [Internet]. 2018;44(6):1209–20.

882   Available from: https://doi.org/10.1016/j.jglr.2018.08.015

883   20.   Hecky RE, Fee EJ, Kling H, Rudd JWM. Studies on the planktonic ecology of

884   Lake Tanganyika. Winnipeg, Man West Reg Fish Mar Serv. 1978;(Fisheries and

885   Marine Service Technical Report 816).

886   21.   Hecky RE, Kling HJ. The phytoplankton and protozooplankton Lake Tanganyika:

887   Species composition, biomass, chlorophyll content, and spatio-temporal

888   distribution. Limnol Ocean. 1981;26(3):548–64.

889   22.   Coulter GW. Population changes within a group of fish species in Lake

890   Tanganyika following their exploitation. J Fish Biol. 1970;2:329–53.

891   23.   Kimirei IA, Mgaya YD. Influence of environmental factors on seasonal changes in

892   clupeid catches in the Kigoma area of Lake Tanganyika. African J Aquat Sci

893   [Internet]. 2007;32(3):291–8. Available from:

894   http://www.informaworld.com/openurl?genre=article&doi=10.2989/AJAS.2007.3

2.3.9.308&magic=crossref%7C%7CD404A21C5BB053405B1A640AFFD44AE3

24.   Cirhuza DM, Plisnier P. Composition and seasonal variations in abundance of

Copepod (Crustacea) populations from the northern part of Lake Tanganyika.

Aquat Ecosyst Heal Manag. 2016;4988(December):401–10.

25.   Mulimbwa N, Sarvala J, Joost AM. Reproductive activities of two

zooplanktivorous clupeid fish in relation to the seasonal abundance of copepod

prey in the northern end of Lake Tanganyika. Belgian J Zool. 2014;144(2):77–92.

26.   Kurki H, Mannini P, Vuorinen I, Aro E, Mölsä H, Lindqvist O V.

Macrozooplankton communities in Lake Tanganyika indicate food chain

differences between the northern part and the main basins. Hydrobiologia.

1999;407:123–9.

27.   Narita T, Mulimbwa N, Mizuno T. Vertical Distribution and Seasonal Abundance

of Zooplankters in Lake Tanganyika. Afr Study Monogr. 1986;6:1–16.

28.   Pearce MJ. A note on the importance of the prawns to the ecology of southern

Lake. Ecol Limnol Study Lake Tanganyika Its Adjac Reg Dep Zool Fac Sci Kyoto

Univ Kyoto. 1991;606p.

29.   Kurki H, Vuorinen I, Bosma E, Bwebwa D. Spatial and temporal changes in

copepod zooplankton communities of Lake Tanganyika. Hydrobiologia.

1999;407:105–14.

30.   Mgana HF, Herzig A, Mgaya YD. Diel vertical distribution and life history

characteristics of Tropodiaptomus simplex and its importance in the diet of

Stolothrissa tanganicae, Kigoma, Tanzania. Aquat Ecosyst Heal Manag.

2014;17(1):14–24.

31.   Mannini P, Katonda KI, Kissaka B, Verburg P. Feeding ecology of Lates stappersii

in Lake Tanganyika. In: Lindqvist O V., Mölsä H, Salonen K, Sarvala J, editors.

From Limnology to Fisheries: Lake Tanganyika and Other Large Lakes. Springer

Netherlands; 1999. p. 131–9.

32.   Matthes H. Preliminary investigations into the biology of the Lake Tanganyika

Clupeidae. Fish Res Bull Zambia. 1967;4:39–45.

33.   Mulimbwa N, Raeymaekers JAM, Sarvala J. Seasonal changes in the pelagic catch

of two clupeid zooplanktivores in relation to the abundance of copepod

zooplankton in the northern end of Lake Tanganyika. Aquat Ecosyst Heal Manag.

2014;17(1):25–33.

34.   De Keyzer ELR, De Corte Z, Van Steenberge M, Raeymaekers JAM, Calboli FCF,

Kmentová N, et al. First genomic study on Lake Tanganyika sprat Stolothrissa

tanganicae: A lack of population structure calls for integrated management of this

important fisheries target species. BMC Evol Biol. 2019;19(1):1–15.

35.   Junker J, Rick JA, Mcintyre PB, Kimirei I, Sweke EA, Mosille JB, et al. Structural

genomic variation leads to genetic differentiation in Lake Tanganyika's sardines.

Mol Ecol. 2020;29:3277–3298.

36.   Rick JA, Junker J, Kimirei IA, Sweke EA, Mosille JB, Dinkel C, et al. The Genetic

936   Population Structure of Lake Tanganyika's Lates Species Flock, an Endemic

937   Radiation of Pelagic Top Predators. J Hered [Internet]. 2021;esab072. Available

938   from: https://doi.org/10.1093/jhered/esab072

939 37.   Cornelissen IJM, Vijverberg J, van den Beld AM, Helmsing NR, Verreth JAJ,

940   Nagelkerke LAJ. Heterogeneity in food-web interactions of fish in the Mwanza

941   Gulf, Lake Victoria: a quantitative stable isotope study. Hydrobiologia.

942   2018;805(1):113–30.

943 38.   Arrington DA, Davidson BK, Winemiller KO, Layman CA. Influence of life

944   history and seasonal hydrology on lipid storage in three neotropical fish species. J

945   Fish Biol. 2006;68(5):1347–61.

946 39.   Montgomery WL, Galzin R. Seasonality in gonads, fat deposits and condition of

947   tropical surgeonfishes (Teleostei: Acanthuridae). Mar Biol. 1993;115(4):529–36.

948 40.   Logan JM, Pethybridge H, Lorrain A, Somes CJ, Allain V, Bodin N, et al. Global

949   patterns and inferences of tuna movements and trophodynamics from stable

950   isotope analysis. Deep Res Part II Top Stud Oceanogr. 2020;175:104775.

951 41.   Post DM. Using Stable Isotopes to Estimate Trophic Position: Models, Methods,

952   and Assumptions. Ecology. 2002;83(3):703–18.

953 42.   Fry B. Stable isotope ecology. New York: Springer, New York; 2006. 308 p.

954 43.   Van der Zanden MJ, Cabana G, Rasmussen JB. Comparing trophic position of

955   freshwater fish calculated using stable nitrogen isotope ratios (δ^{15}N) and literature

dietary data. Can J Fish Aquat Sci. 1997;54(5):1142–58.

44.   Bidigare RR, Fluegge A, Freeman KH, Hanson KL, Hayes JM, Hollander DJ, et

al. Consistent fractionation of ^{13}C in nature and in the laboratory: Growth-rate

effects in some haptophyte algae. Global Biogeochem Cycles. 1997;11(2):279–92.

45.   Bidigare RR, Hanson KL, Buesseler KO, Wakeham SG, Freeman KH, Pancost

RD, et al. Iron-stimulated changes in ^{13}C fractionation and export by equatorial

Pacific phytoplankton: Toward a paleogrowth rate proxy. Paleooceanography.

1999;14(5):589–95.

46.   Hollander DJ, McKenzie JA. \ce{CO_{2}} control on carbon-isotope fractionation

during aqueous photosynthesis: A paleo-p\ce{CO_{2}} barometer. Geology.

1991;19:929–32.

47.   Gu B, Schelske CL, Brenner M. Relationship between sediment and plankton

isotope ratios (δ^{13}C and δ^{15}N) and primary productivity in Florida lakes. Can J Fish

Aquat Sci. 1996;53(4):875–83.

48.   Rau GH, Riebesell U, Wolf-Gladrow D. CO$_{2\ (aq)}$-dependent photosynthetic ^{13}C

fractionation in the ocean: A model versus measurements. Global Biogeochem

Cycles. 1997;11(2):267–78.

49.   Close HG, Henderson LC. Open-ocean minima in δ^{13}C values of particulate

organic carbon in the lower euphotic zone. Front Mar Sci. 2020;7:540165.

50.   Perga ME, Gerdeaux D. "Are fish what they eat" all year round? Oecologia.

2005;144(4):598–606.

51.   O'Reilly CM, Hecky RE, Cohen AS, Plisnier P-D. Interpreting stable isotopes in
978   food webs: Recognizing the role of time averaging at different trophic levels.
979   Limnol Oceanogr. 2002;47(1):306–9.

980   52.   Sarvala J, Badende S, Chitamwebwa D, Juvonen P, Mwape L, Mölsä H, et al.
981   Size-fractionated $\delta^{15}N$ and $\delta^{13}C$ isotope ratios elucidate the role of the microbial
982   food web in the pelagial of Lake Tanganyika. Aquat Ecosyst Heal Manag.
983   2003;6(3):241–50.

984   53.   Salonen K, Högmander P, Langenberg V, Mölsä H, Sarvala J, Tarvainen A, et al.
985   Limnocnida tanganyicae medusae (Cnidaria: Hydrozoa): A semiautonomous
986   microcosm in the food web of Lake Tanganyika. Hydrobiologia. 2012;690(1):97–
987   112.

988   54.   Post DM, Layman CA, Arrington DA, Takimoto G, Quattrochi J, Montaña CG.
989   Getting to the fat of the matter: Models, methods and assumptions for dealing with
990   lipids in stable isotope analyses. Oecologia. 2007;152(1):179–89.

991   55.   Logan JM, Jardine TD, Miller TJ, Bunn SE, Cunjak RA, Lutcavage ME. Lipid
992   corrections in carbon and nitrogen stable isotope analyses: Comparison of
993   chemical extraction and modelling methods. J Anim Ecol. 2008;77(4):838–46.

994   56.   Ehrenfels B, Bartosiewicz M, Mbonde AS, Baumann KBL, Dinkel C, Junker J, et
995   al. Diazotrophic cyanobacteria are associated with a low nitrate resupply to surface
996   waters in Lake Tanganyika. Front Environ Sci [Internet]. 2021;9:277. Available

from: https://www.frontiersin.org/article/10.3389/fenvs.2021.716765

57.   Ehrenfels B, Mbonde AS, Dinkel C, Brand A, Baumann KBL, Junker J, et al.

Physical, biogeochemical, and biological data for Lake Tanganyika (2017–2018).

ETH Res Collect. 2020;Dataset.

58.   Winslow L, Read J, Woolway R, Brentrup J, Leach T, Zwart J, et al. Package

'rLakeAnalyzer.' Lake Phys Tools. 2019;

59.   Cocquyt C, Vyverman W. Phytoplankton in Lake Tanganyika: a Comparison of

Community Composition and Biomass off Kigoma with Previous Studies 27 Years

Ago. J Great Lakes Res [Internet]. 2005;31(4):535–46. Available from:

http://dx.doi.org/10.1016/S0380-1330(05)70282-3

60.   Descy JP, Hardy MA, Sténuite S, Pirlot S, Leporcq B, Kimirei I, et al.

Phytoplankton pigments and community composition in Lake Tanganyika. Freshw

Biol. 2005;50(4):668–84.

61.   Holmes RM, Peterson BJ, Aminot A, Kérouel R, Hooker BA. A simple and

precise method for measuring ammonium in marine and freshwater ecosystems.

Can J Fish Aquat Sci. 1999;56(10):1801–8.

62.   Schnetger B, Lehners C. Determination of nitrate plus nitrite in small volume

marine water samples using vanadium(III)chloride as a reduction agent. Mar Chem

[Internet]. 2014;160:91–8. Available from:

http://dx.doi.org/10.1016/j.marchem.2014.01.010

63.   Grasshoff K, Kremling K, Ehrhardt M. Methods of seawater analysis. Wiley, New

1018   York; 1999. 600 p.

1019   64.   Schunck H, Lavik G, Desai DK, Großkopf T, Kalvelage T, Löscher CR, et al.

1020   Giant Hydrogen Sulfide Plume in the Oxygen Minimum Zone off Peru Supports

1021   Chemolithoautotrophy. PLoS One. 2013;8(8):e68661.

1022   65.   Callbeck CM, Ehrenfels B, Baumann KBL, Wehrli B, Schubert CJ. Anoxic

1023   chlorophyll maximum enhances local organic matter remineralization and nitrogen

1024   loss in Lake Tanganyika. Nat Commun. 2021;12(830).

1025   66.   Wasmund N, Topp I, Schories D. Optimising the storage and extraction of

1026   chlorophyll samples. Oceanologia. 2006;48(1):125–44.

1027   67.   Folch J, Lees M, Sloane-Stanley GH. A simple method for the isolation and

1028   purification of total lipids from animal tissues. J Biol Chem. 1957;226(1):497–509.

1029   68.   Jessen HH, Mgana HF, Grønkjær P, Lauridsen TL. Dietary differences among

1030   commercially important fishes in Lake Tanganyika assessed using stable isotope

1031   analysis. J Great Lakes Res [Internet]. 2019;45(6):1205–14. Available from:

1032   https://doi.org/10.1016/j.jglr.2019.08.008

1033   69.   Jackson AL, Inger R, Parnell AC, Bearhop S. Comparing isotopic niche widths

1034   among and within communities: SIBER — Stable Isotope Bayesian Ellipses in R.

1035   J Anim Ecol. 2011;80(3):595–602.

70.   De Wever A, Muylaert K, Van Der Gucht K, Pirlot S, Cocquyt C, Descy J, et al.

Bacterial Community Composition in Lake Tanganyika: Vertical and Horizontal

Heterogeneity. Appl Environ Microbiol. 2005;71(9):5029–37.

71.   De Wever A, Muylaert K, Langlet D, Alleman L, Descy JP, André L, et al.

Differential response of phytoplankton to additions of nitrogen, phosphorus and

iron in Lake Tanganyika. Freshw Biol. 2008;53(2):264–77.

72.   Descy JP, Tarbe AL, Stenuite S, Pirlot S, Stimart J, Vanderheyden J, et al. Drivers

of phytoplankton diversity in Lake Tanganyika. Hydrobiologia. 2010;653(1):29–

44.

73.   Stenuite S, Tarbe AL, Sarmento H, Unrein F, Pirlot S, Sinyinza D, et al.

Photosynthetic picoplankton in Lake Tanganyika: Biomass distribution patterns

with depth, season and basin. J Plankton Res. 2009;31(12):1531–44.

74.   Rufli H, Chapman DW. Preliminary analysis of zooplankton sampling and

estimates of fish abundance in Lake Tanganyika in October 1975. United Nations

Food Agric Organ Report, FI DP URT/71/012/31. 1976;14p, FAO, Rome.

75.   Plisnier PD, Mgana H, Kimirei I, Chande A, Makasa L, Chimanga J, et al.

Limnological variability and pelagic fish abundance (Stolothrissa tanganicae and

Lates stappersii) in Lake Tanganyika. Hydrobiologia. 2009;625(1):117–34.

76.   Horion S, Bergamino N, Stenuite S, Descy JP, Plisnier PD, Loiselle SA, et al.

Optimized extraction of daily bio-optical time series derived from MODIS/Aqua

imagery for Lake Tanganyika, Africa. Remote Sens Environ. 2010;114(4):781–91.

77.   Gu B, Chapman AD, Schelske CL. Factors controlling seasonal variations in stable

1058   isotope composition of particulate organic matter in a soft water eutrophic lake.

1059   Limnol Oceanogr. 2006;51(6):2837–48.

1060   78.   Craig H. Lake Tanganyika geochemical and hydrographic study: 1973 expedition.

1061   SIO Ref Ser. 1974;75(5).

1062   79.   Morana C, Darchambeau F, Roland FAE, Borges A V., Muvundja F, Kelemen Z,

1063   et al. Biogeochemistry of a large and deep tropical lake (Lake Kivu, East Africa:

1064   Insights from a stable isotope study covering an annual cycle. Biogeosciences.

1065   2015;12(16):4953–63.

1066   80.   Brandl Z. Freshwater Copepods and Rotifers: Predators and their Prey.

1067   Hydrobiologia. 2005;546:475–89.

1068   81.   Matthews B, Mazumder A. Distinguishing trophic variation from seasonal and

1069   size-based isotopic δ^{15}N variation of zooplankton. Can J Fish Aquat Sci.

1070   2007;64(1):74–83.

1071   82.   Fukumori K, Oi M, Doi H, Takahashi D, Okuda N, Miller TW, et al. Bivalve

1072   tissue as a carbon and nitrogen isotope baseline indicator in coastal ecosystems.

1073   Estuar Coast Shelf Sci. 2008;79(1):45–50.

1074   83.   Langlet D, Alleman LY, Plisnier P-D, Hughes H, André L. Manganese content

1075   records seasonal upwelling in Lake Tanganyika mussels. Biogeosciences.

1076   2007;4(2):195–203.

84.   Woodland RJ, Magnan P, Glémet H, Rodríguez MA, Cabana G. Variability and

directionality of temporal changes in $\delta^{13}C$ and $\delta^{15}N$ of aquatic invertebrate primary

consumers. Oecologia. 2012;169(1):199–209.

85.   Xia B, Gao QF, Dong SL, Wang F. Carbon stable isotope turnover and

fractionation in grass carp ctenopharyngodon idella tissues. Aquat Biol.

2013;19(3):207–16.

86.   Busst GMA, Britton JR. Tissue-specific turnover rates of the nitrogen stable

isotope as functions of time and growth in a cyprinid fish. Hydrobiologia.

2018;805(1):49–60.

87.   Weidel BC, Carpenter SR, Kitchell JF, Van der Zanden MJ. Rates and components

of carbon turnover in fish muscle: Insights from bioenergetics models and a whole-

lake ^{13}C addition. Can J Fish Aquat Sci. 2011;68(3):387–99.

88.   Mateo MA, Serrano O, Serrano L, Michener RH. Effects of sample preparation on

stable isotope ratios of carbon and nitrogen in marine invertebrates: Implications

for food web studies using stable isotopes. Oecologia. 2008;157(1):105–15.

89.   Bastos RF, Corrêa F, Winemiller KO, Garcia AM. Are you what you eat? Effects

of trophic discrimination factors on estimates of food assimilation and trophic

position with a new estimation method. Ecol Indic [Internet]. 2017;75:234–41.

Available from: http://dx.doi.org/10.1016/j.ecolind.2016.12.007

90.   Chikaraishi Y, Ogawa NO, Kashiyama Y, Takano Y, Suga H, Tomitani A, et al.

Determination of aquaticfood-web structure based on compound-specific nitrogen

isotopic composition of amino acids. Limnol Oceanogr Methods. 2009;7(11):740–

50.

91.   McConnaughey T, McRoy CP. Food-Web structure and the fractionation of

Carbon isotopes in the bering sea. Mar Biol. 1979;53(3):257–62.

92.   Fernandes T, McMeans BC. Coping with the cold: energy storage strategies for

surviving winter in freshwater fish. Ecography (Cop). 2019;42(12):2037–52.

93.   Abdulkarim B, Bwathondi POJ, Benno BL. Seasonal variations in the proximate

compositions of five economically-important fish species from Lake Victoria and

Lake Tanganyika, Tanzania. Bayero J Pure Appl Sci. 2016;9(1):11–8.

94.   Ellis CMA. The size at maturity and breeding seasons of sardines in southern Lake

Tanganyika. African J Trop Hydrobiol Fish. 1971;1:59–66.

95.   Pearce MJ. A description and stock assessment of the pelagic fishery in the South-

east arm of the Zambian waters of Lake Tanganyika. Rep Dep Fish Zambia.

1985;74p.

1112

Supporting information

Fig. S1: C:N mass ratio of (**a,b**) *Stolothrissa tanganicae,* (**c,d**) *Limnothrissa miodon*, and

(**e,f**) *Lates stappersii* versus standard length in the northern and southern basins during

the end of the dry season and the end of the rainy season. Only stations 1, 2 (north) and 7,

9 (south) are depicted. The shaded areas mark the 50 mm cut-off range for the population

comparisons used in Fig. 5.

Fig. S2: C:N corrected (Post et al. 2007) δ^{13}C of (**a,b**) *Stolothrissa tanganicae,* (**c,d**)

Limnothrissa miodon, and (**e,f**) *Lates stappersii* versus standard length for the end of the

dry season and the end of the rainy season. Only stations 1, 2 (north) and 7, 9 (south) are

depicted. The shaded areas mark the 50 mm cut-off range for the population comparisons

used in Fig. 4. The sampled populations of *L. miodon* and *L. stappersii* from the end of

the rainy season (d,f) were characterized by dense clusters of observations within a

narrow size and δ^{13}C range which may have skewed the basin-scale comparisons.

Fig. S3: Zooplankton community compositions per station for various net types (25 μm,

95 μm, 250 μm) during end of the dry season (top) and the end of the rainy season

(bottom).

Fig. S4: C:N corrected according to Post et al. (54) δ^{13}C (left) and δ^{15}N (right) of (**a,b**)

Stolothrissa tanganicae, (**c,d**) *Limnothrissa miodon*, (**e,f**) *Lates stappersii*, (**g,h**) *Lates*

microlepis, (**i,j**) *Lates mariae* and (**k,l**) *Lates angustifrons* versus standard length

including all sampling locations and campaigns. Samples from the central basin and July

2017 were included for completeness, but were not included in the north-south and

seasonal analysis presented in Fig. 6. Note the different y-axis scaling.

Fig. S5: Carbon (normalized for C:N mass ratio according to Post et al. (54)) and

nitrogen stable isotope signatures of the large *Lates* species, namely (**a,b**) *Lates*

microlepis (**c,d**) *Lates mariae*, and (**e,f**) *Lates angustifrons* at the end of the dry season

(left) and the end of the rainy season (right). Orange dots represent the northern basin

(stations 1-3) and blue dots represent the southern basin (stations 7-9). Numbers indicate

the mean δ^{13}C of a population. Only individuals >150 mm were included in this analysis

to reduce ontogenetic effects on the isotope signatures.

Fig. S6: Lipid content versus C:N ratios of *Stolothrissa tanganicae*. Each dot represents a

tissue sample from one specimen.

Fig. S7: Nitrogen stable isotope signatures of different genetic clusters within the studied

fish species in (**left**) July 2017, (**middle**) September/October 2017, and (**right**) April/May

2018. (**a-c**) *Limnothrissa miodon*, (**d-f**) *Lates stappersii*, (**g-i**) *Lates microlepis* (**j-l**), *Lates*

mariae, and (**m-o**) *Lates angustifrons*. Note the different axis scaling between

Limnothrissa and the *Lates* species.