Attractor Decision Networks with Selective Inhibition

Belle Liu[1, 2], Chung-Chuan Lo[2, 3, ✉], and Kuo-An Wu[1, ✉]

[1] *Department of Physics, National Tsing Hua University, Hsinchu City, Taiwan*

[2] *Institute of Systems Neuroscience, National Tsing Hua University, Hsinchu City, Taiwan*

[3] *Brain Research Center, National Tsing Hua University, Hsinchu City, Taiwan*

Abstract

The ability to decide swiftly and accurately in an urgent scenario is crucial for an organism's survival. The neural mechanisms underlying the perceptual decision and trade-off between speed and accuracy have been extensively studied in the past few decades. Among several theoretical models, the attractor neural network model has successfully captured both behavioral and neuronal data observed in many decision experiments. However, a recent experimental study revealed additional details that were not considered in the original attractor model. In particular, the study shows that the inhibitory neurons in the posterior parietal cortex of mice are as selective to decision making results as the excitatory neurons, whereas the original attractor model assumes the inhibitory neurons to be unselective. In this study, we investigate a more general attractor model with selective inhibition, and analyze in detail how the computational ability of the network changes with selectivity. We proposed a reduced model for the selective model, and showed that selectivity adds a time-varying component to the energy landscape. This time dependence of the energy landscape allows the selective model to integrate information carefully in initial stages, then quickly converge to an attractor once the choice is clear. This results in the selective model having a more efficient speed-accuracy trade-off that is unreachable by unselective models.

Keywords. Selectivity, Attractor Model, Decision Making, Energy Landscape

Introduction

Decision making is a central cognitive function for animals, and the ability to integrate information in a swift and accurate manner is of great importance. The decision behavior is typically measured by speed and accuracy of the subject, and several computational or mathematical models have been proposed to account for these two quantities [1, 2, 3, 4, 5, 6, 7, 8]. In addition to the psychometric measures, the neurological basis of decision making has always been a highly sought-out topic of interest. The attractor neural network models [9, 10, 11] are the models of interest because they provides insight into the neuronal interactions during the decision process. Furthermore, they link the mathematical equations and variables to known biological substrates, which is crucial for studying how structural changes of the network influences its functionalities.

The attractor model is composed of two competing excitatory populations and one inhibitory population, and all populations are interconnected with one another (Fig. 1a, rightmost). The two excitatory populations receive input signals, each representing "evidence" in favor of one of the two choices, and eventually one population "wins" by silencing the other (Fig. 1b).

Attractor models have been successful in recreating the decision making process observed in experiments. One commonly used experimental task is the random dot motion coherence test, where several dots randomly drift towards the left or right [12, 13, 14]. The subject, upon observing the motion, has to report the direction in which the majority of dots are moving towards. However, new experimental results have revealed new details regarding the network structure of attractor models in the posterior parietal cortex of mice. Specifically, a recent study by Najafi *et al.* investigated the selectivity of decision making networks [15]. For the attractor model, the excitatory neurons are selective to the results of the decision. The inhibitory neurons, on the other hand, will react similarly in both cases. Therefore, in the attractor model, excitatory neurons are selective to input while the inhibitory neurons are not.

Selectivity is important for our understanding of cortical network architecture. Previous studies lean towards a framework where excitatory neurons are selective, but not inhibitory neurons [16, 17, 18, 19, 20]. This is motivated by inhibitory neurons having denser input than excitatory neurons, thereby having broader tuning curves [20]. It is also observed that excitatory neurons tend to connect more to other excitatory neurons with similar feature preference, and that its selectivity to orientation increases when the connectivity decreases [20].

However, plenty of contradictory results are also reported, where it is found that inhibitory neurons across multiple brain areas respond selectively to different features of the stimulus [21, 22, 23, 24, 25, 26, 27]. The aforementioned study by Najafi *et al.* tested whether selectivity exists within the posterior parietal cortex of

mice, using a two-alternative forced choice task [15]. They showed that the inhibitory neurons are as selective as the excitatory neurons. This result suggests a network architecture wherein the inhibitory neurons are differently activated by the two inputs. One of the proposed alternative is a model in which there consists of two inhibitory populations, each receiving stronger input from one population (Fig. 1a, leftmost). This could be thought of as a natural generalization of the attractor model, where inhibitory neurons are allowed to be heterogeneous. The previous unselective model is thereby a special case where all inhibitory neurons are homogeneous in synaptic properties. That is, in the SM, the synaptic connections from an excitatory population to an inhibitory population (or vice versa) on the same side is not constrained to be the equal to those to the opposite side. In the UM however, they are required to be equal for the inhibitory population to remain unselective, which is an extra restriction.

Our question of interest, following this proposed alternative, is that whether the incorporation of selectivity brings about any computational changes. That is, does the present model (hereby denoted as the Selective Model, or SM) provide computational advantages not present within the previous model (Unselective Model, or UM)? In this study, we demonstrated how selectivity utilizes a time-dependent energy landscape to reach an efficient speed-accuracy trade-off. In particular, the time-variance allows the SM to evaluate evidence carefully in initial stages, converge quickly once the choice is clear, and avoids indecisive trials.

Figure 1: The selective model of decision making can qualitatively reproduce behavioral and physiological observations. (a) The selective model (SM) is a generalized attractor model, with the previous unselective model (UM) being a special case where all inhibitory neurons are homogeneous in synaptic weights. From *left* to *right*: SM, SM with excitatory-to-inhibitory and inhibitory-to-excitatory connections being identical across the two inhibitory populations, which is equivalent to the UM. The use of solid and dashed lines is to highlight the difference in synaptic weights between the populations. (b) Example simulation trial of the two-variable reduced model. The green line represents the activity of the winning population, and red the losing one. The decision is defined as one of the populations' activity reaching a certain threshold. (c) Percentage correct and mean reaction time for experimental data [13], figure adapted from [10] with permission. (d) Same as in (c) but for SM with $\{\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}\} = \{0.5, 0.6, 0.7, 1\}$. The α_p, β_p values are 7.4 and 1.31 respectively, compared to the experimental values of 7.4 and 1.3. An additional 100 ms is added to the reaction time as the non-decision time that represents other neural processes involved but not related to the decision network modeled here. Number of trials ran is described in Methods. (e) Selectivity as a function of time in experimental data, adapted from [15] with permission. (f) Selectivity as a function of time in the full neural network version of the SM and UM ($\gamma_{ee} = 0.5$). Choice selectivity is defined as in (e), which is $2 \times |AUC - 0.5|$ (see [15]); mean $\pm$ SEM by bootstrapping, number of trials $N = 120$. (g) Correlations for EE, II, EI pairs of neurons with either the same (darker color) or opposite (lighter color) selectivity, compared to shuffled data (at the bottom of the graphs). Figure adapted from [15] with permission. (h) Correlations of EE, II, EI pairs of neurons in the full neural network version of the SM. The color schemes mimic those of (g); mean $\pm$ SEM, $N = 12$.

Methods

Population Models

The classic attractor neural network model of decision making consists of hundreds of neurons, and in combination with the synapses, the complete system will contain thousands of differential equations, which complicates both simulation and analysis [9]. In a study by Wong and Wang [28], it is shown that the attractor model can be reduced to a simpler two-variable form. The reduction takes advantage of several facts: (I) the average activity of the population can be represented by one single variable through mean field reduction; (II) the inhibitory population responds linearly to input; and (III) the time scale of AMPA, GABA and the firing rates are much faster than NMDA, and thus can be assumed to have reached steady state instantaneously. We will call this two-variable model the WWM (Wong and Wang's Model). The resulting model contains only equations for the NMDA channels of the two competing populations (see Supplementary Notes S1 and Fig. S1). However, since the time constant of noise in the input (Eq S5) is comparable to the time constant of GABA dynamics (2 ms and 5 ms respectively), the rate of change in NMDA could be much shorter compared to its relaxation time. Therefore, the GABA dynamics is retained in the following discussion, and it is shown that including GABA dynamics could alter the dynamics drastically (Supplementary Notes S2, Fig. S2).

Here we will start with deriving the simpler model, which is the UM, and generalize to the SM. With the inclusion of GABA dynamics alongside NMDA dynamics, the full UM is:

$$
\begin{aligned}
\dot{S}_{E1} &= -\frac{S_{E1}}{\tau_{\mathrm{NMDA}}} + 0.641 \times (1 - S_{E1}) H(I_{syn,1}) \\
\dot{S}_{E2} &= -\frac{S_{E2}}{\tau_{\mathrm{NMDA}}} + 0.641 \times (1 - S_{E2}) H(I_{syn,2}) \\
\dot{S}_{G} &= -\frac{S_{G}}{\tau_{\mathrm{GABA}}} + \phi(I_{syn,G}),
\end{aligned}
\tag{1}
$$

where S_{E1}, S_{E2} are the gating variables of NMDA, and $\tau_{\mathrm{NMDA}} = 0.1$ (s) is the time constant of the NMDA channels. Gating variables reflect the degree in which an ion channel is opened. S_G is the gating variable of GABA, and $\tau_{\mathrm{GABA}} = 0.005$ (s). $H(x)$ is the $f - I$ curve of the excitatory population, and is shaped like the lower end of a sigmoid-like function (Supplementary Notes S1). $\phi(x)$ takes a form similar to $H(x)$, but can be linearized since according to experimental findings, the activity of the inhibitory population never falls into the nonlinear regime (i.e. assumption (II)) [28]. The synaptic current, I_{syn}, is a weighted sum of pre-synaptic inputs.

The terminology for the synaptic weights are shown in Fig. 1a. To briefly summarize, for a synaptic weight g_{mn} where $\{m, n\} \in \{e, i\}$, m represents the pre-synaptic population type (e for excitatory, i for inhibitory) and n the post-synaptic population type. The intra-population excitatory-to-excitatory synaptic weight ($g_{ee,s}$, where s

94   stands for *same side*) is different from the inter-population excitatory-to-excitatory synaptic weight ($g_{ee,o}$, where

95   o stands for *opposite side*). The bias currents to the excitatory and inhibitory populations are denoted as I_e and

96   I_i.

97   For simplicity, the inhibitory populations receive no noise. The equations for the UM synaptic currents are:

$$I_{syn,1} = g_{ee,s}S_{E1} + g_{ee,o}S_{E2} - g_{ie}S_G + I_e + I_n + I_1$$

$$I_{syn,2} = g_{ee,s}S_{E2} + g_{ee,o}S_{E1} - g_{ie}S_G + I_e + I_n + I_2 \tag{2}$$

$$I_{syn,G} = g_{ei}S_{E1} + g_{ei}S_{E2} - g_{ii}S_G + I_i.$$

98   The SM essentially divides the inhibitory population into two, allowing them to take on different synaptic

99   weights. Therefore, the equation for the SM is:

$$\dot{S}_{E1} = -\frac{S_{E1}}{\tau_{\text{NMDA}}} + 0.641 \times (1 - S_{E1})H(I_{syn,1})$$

$$\dot{S}_{E2} = -\frac{S_{E2}}{\tau_{\text{NMDA}}} + 0.641 \times (1 - S_{E2})H(I_{syn,2})$$

$$\dot{S}_{G1} = -\frac{S_{G1}}{\tau_{\text{GABA}}} + \phi(I_{syn,G1}), \tag{3}$$

$$\dot{S}_{G2} = -\frac{S_{G2}}{\tau_{\text{GABA}}} + \phi(I_{syn,G2}),$$

100   The SM synaptic currents are:

$$I_{syn,1} = g_{ee,s}S_{E1} + g_{ee,o}S_{E2} - g_{ie,s}S_{G1} - g_{ie,o}S_{G2} + I_e + I_n + I_1$$

$$I_{syn,2} = g_{ee,s}S_{E2} + g_{ee,o}S_{E1} - g_{ie,s}S_{G2} - g_{ie,o}S_{G1} + I_e + I_n + I_2$$

$$I_{syn,G1} = g_{ei,s}S_{E1} + g_{ei,o}S_{E2} - g_{ii,s}S_{G1} - g_{ii,o}S_{G2} + I_i \tag{4}$$

$$I_{syn,G2} = g_{ei,s}S_{E2} + g_{ei,o}S_{E1} - g_{ii,s}S_{G2} - g_{ii,o}S_{G1} + I_i,$$

To ensure fair comparison across different models that we study, and to isolate the effect of selective inhibition,

we created models whose equilibrium points are the same when projected onto the S_{E1}-S_{E2} plane. That is, upon

activating assumption (III), these model would all collapse back into an identical WWM. This provides an

additional benefit of assuring us that GABA dynamics do in fact make a difference; because if it does not, then

all models should yield the same results since their two-variable form is identical.

A technical description of how this is done is as follows: for the UM, enforcing assumption (III) on Eq 1 yields

$$S_G = \tau_{\text{NMDA}} \times \phi(g_{ei}(S_{E1} + S_{E2}) - g_{ii}S_G). \tag{5}$$

107   That is, since assumption (III) assumes that S_G always reaches the equilibrium state immediately, it assumes

108   that $\frac{dS_G}{dt} = 0$, which is how we were able to arrive at Eq 5. By substituting Eq 5 back into Eq 1, we can

109   once again lower the number of differential equations back to two. Since in Eq 5, S_G is a linear combination of

S_{E1} and S_{E2}, after substitution the synaptic currents $I_{syn,1}$ and $I_{syn,2}$ will remain a linear combination of S_{E1} and S_{E2}. Therefore, we could collect the coefficients for S_{E1}, S_{E2} and the bias term in the synaptic currents – which would consist of the parameters $\{g_{ee,s}, g_{ee,o}, g_{ei}, g_{ie}, g_{ii}, I_e, I_i\}$ – and call them $\{J_e, J_i, I_b\}$, which are the parameters for the WWM (Fig. S1). J_e represents the strength of self-excitation, J_i represents the degree in which the competing populations inhibit one another, and I_b is the bias.

Since the degrees of freedom in the former exceeds the latter, there are infinite sets of UM parameters $\{g_{ee,s}, g_{ee,o}, g_{ei}, g_{ie}, g_{ii}, I_e, I_i\}$ that could be mapped to the same two-variable parameters $\{J_e, J_i, I_b\}$. For convenience purposes, four constraints will be added to the mapping process, so that the mapping becomes one-to-one. This allows us to be able to adjust one single parameter, and have the other parameters be automatically sought out. These four constraints are: (I) $g_{ie} = g_{ei}$, (II) $g_{ii} = 1.2$, (III) $I_i = 1$, and (IV) $\gamma_{ee} := \frac{g_{ee,o}}{g_{ee,s}}$ ($0 < \gamma_{ee} < 1$). The first constraint originates from the observation that excitation and inhibition are often balanced in neural networks [15, 29, 30]. The second and third are chosen so that the resulting UM does not violate assumption (II). Constraint (IV) defines a parameter γ_{ee} for which we can adjust to control the speed in which the activity of the excitatory population ramps up. Note that when γ_{ee} is altered, the synaptic weights $\{g_{ee,s}, g_{ee,o}, g_{ei}, g_{ie}\}$, as well as bias currents $\{I_e, I_i\}$ will all change along in order to satisfy the requirement imposed by assumption (III) and the four constraints.

Under these constraints, we can obtain unique UM parameters (Supplementary Notes S2). A reduction from the SM to the WWM can also be conducted in a similar fashion. That is, the differential equations for S_{G1}, S_{G2} can be simplified in the same manner as Eq 5, and again substituted into Eq 3, which would again lead to a WWM (Supplementary Notes S2).

Similar to how γ_{ee} was defined in the UM to account for the ratio between the inter- and intra-population synaptic weight of the excitatory-to-excitatory connection, we also define $\gamma_{ei} := \frac{g_{ei,o}}{g_{ei,s}}$, $\gamma_{ie} := \frac{g_{ie,o}}{g_{ie,s}}$ and $\gamma_{ii} := \frac{g_{ii,o}}{g_{ii,s}}$ for the other three types of connections that exist in the SM. Note that while γ_{ee} must be strictly between 0 and 1 in order for the network to function properly, no such constraints exist for γ_{ei}, γ_{ie} and γ_{ii}. Intuitively, if $\gamma_{ee} > 1$, then the favored population would excite its opponent more than it excites itself ($g_{ee,o} > g_{ee,s}$), leading to a situation where no decision can be made. However, if say $\gamma_{ie} > 1$, it merely means that the inhibitory population prefers to inhibit the contralateral excitatory population more; if $\gamma_{ie} < 1$, then the inhibitory population inhibits its ipsilateral excitatory population more.

Reduced Population Models

The UM and SM have three and four variables respectively, which makes geometrical interpretations of their dynamics difficult. To better analyze the UM and SM, we reduced the models back to a two-variable form. However, instead of using assumption (III), the reduction is based on empirical observations of the simulations of population models, which allows the reduced models to better capture the dynamics of the full models.

Linear Relation of Average Excitation and Average Inhibition in UM and SM

It is observed that S_G and the average excitation S_+ in the UM, where $S_+ := \frac{S_{E1}+S_{E2}}{2}$, evolve in a linear manner throughout the simulation (Fig. S3a, e). An analogous relation arises in the SM as well, where the linear relation is between S_{G+} ($S_{G+} := \frac{S_{G1}+S_{G2}}{2}$, i.e. average inhibition) and S_+. Therefore, they are fitted to a linear function, $S_G = aS_+ + b$ ($S_{G+} = aS_+ + b$ for the SM). The goal of fitting a linear function to S_G and S_+ is to reduce the UM and SM back into the two variable form, with new parameters that more accurately describe its dynamics than if assumption (III) were in place. That is, we can replace S_G by $aS_+ + b$ in Eq 2, so that the number of variables once again reduces back to two. The synaptic currents after this simplification become:

$$I_{syn,1} = g_{ee,s}S_{E1} + g_{ee,o}S_{E2} - g_{ie}(aS_+ + b) + I_e$$

$$I_{syn,2} = g_{ee,s}S_{E2} + g_{ee,o}S_{E1} - g_{ie}(aS_+ + b) + I_e. \tag{6}$$

By letting $J_{e,lin} = g_{ee,s} - \frac{g_{ie}a}{2}$, $J_{i,lin} = g_{ee,o} - \frac{g_{ie}a}{2}$ and $I_{b,lin} = I_e - g_{ie}b$, we note that this model has the exact same mathematical form as WWM (Supplementary Notes S1). We will refer to the parameters of the linear reduced model as $\{J_{e,lin}, J_{i,lin}, I_{b,lin}\}$, even though they play the same role as $\{J_e, J_i, I_b\}$ in the WWM.

The fitted a, b values vary slightly across each trial; therefore, a more accurate reduction would be to obtain a_{sim} and b_{sim} via averaging over the simulated trials of the UM. However, to avoid having to run numerous trials just to obtain a and b, they are instead approximated by running a single trial in which no noise is present. The noiseless trial starts at the stable equilibrium point in which $S_{E1} = S_{E2}$, and in the absence of noise, the system stays on the separatrix throughout the simulation. The resulting slope a^* and intersect b^* is then used to simulate the linear reduced model of the UM.

a^*, b^* obtained by the noiseless trials are compared to a_{sim}, b_{sim} obtained by simulations to evaluate its validity (Fig. S3b). The data points used for fitting a_{sim}, b_{sim} are the data points beginning from the onset of the input ($t = 0.5$ (s)) up to $t = 1.0$ (s). This is because including data points where the system is in the stable state only adds noise to the fitting process and no information. Therefore, the time in which the system is in the initial stable state is omitted by only including points from $t = 0.5$ (s) onward. Since the time in which each

165 trial reaches the final stable state fluctuates, we compensated by truncating the data points beyond $t = 1.0$ (s).

166 This is because the average reaction times seldom exceed that value.

Quadratic Relation of Excitation Difference and Inhibition Difference in SM

For the SMs, we define excitatory difference $S_- := \frac{S_{E1} - S_{E2}}{2}$ and inhibitory difference $S_{G-} := \frac{S_{G1} - S_{G2}}{2}$.

(Note that an analogous term of S_{G-} does not exist in the UM, since the UM does not contain two inhibitory

populations.) While at first glance, the relation between S_{G-} and S_- appears linear, it is actually quadratic

(Fig. S3c, e). The coefficients of this quadratic relation, $S_{G-} = a_Q^* S_-^2 + b_Q^* S_- + c_Q^*$, cannot be obtained by

running a fair trial (i.e. S_{E1} and S_{E2} starting out with the same value) without noise, since this would result

in having $S_- = 0$ at all times. Instead, an unfair trial without noise is run by having the initial S_{NMDA} values

be biased towards one side (($S_{E1}, S_{E2}) = (0.26, 0.25)$ at $t = 0$). The value in which we choose to start from will

influence the $\{a_Q^*, b_Q^*\}$ values obtained, but they remain roughly similar within an acceptable regime. The data

points used to find $a_{Q,sim}, b_{Q,sim}$ are similarly the ones from the onset up to 1.0 (s), and they are also compared

to a_Q^*, b_Q^* in Fig. S3d. The fact that this quadratic relation exists in a noiseless trial shows that this is not a

noise-induced artifact, but a property inherent to the SM.

For analysis purposes, we model this quadratic relation as a linear relation with a time-varying slope and

intercept. That is, we linearized the quadratic relation within short periods of time. In the noiseless trial, the

data points for every 250 ms are fitted to a line,

$$S_{G-} = a_L(t)S_- + b_L(t), \tag{7}$$

182 where the coefficients $a_L(t), b_L(t)$ change for the different periods of time. Substituting Eq 7 into Eq 4, the

183 synaptic currents become:

$$I_{syn,1} = g_{ee,s}S_{E1} + g_{ee,o}S_{E2} - (g_{ie,s} + g_{ie,o})(aS_+ + b) - (g_{ie,s} - g_{ie,o})(a_L(t)S_- + b_L(t)) + I_e$$

$$I_{syn,2} = g_{ee,s}S_{E2} + g_{ee,o}S_{E1} - (g_{ie,s} + g_{ie,o})(aS_+ + b) + (g_{ie,s} - g_{ie,o})(a_L(t)S_- + b_L(t)) + I_e. \tag{8}$$

184   We call this the time-dependent linear reduced model. The only difference this model has with the WWM or

185 the linear reduced model in terms of mathematical form is the time dependence of its variables $a_L(t)$ and $b_L(t)$.

186 The relations between the WWM, the full UM/SMs, the linear reduced model and the time-dependent linear

187 reduced model are summarized in Fig. S4. The performance of the reduced models in comparison to the full

188 models are summarized in Supplementary Notes S3 and Fig. S3.

Full Scale Neural Network Model

Population models, being a simplification, may not be able to capture certain quantities observed in experiments in which activities of individual neurons were measured. Therefore, in order for us to compare the UM and SMs to experimental data, we constructed a full neural network model that contains 50 neurons in each population. The connections are all-to-all between populations or within a population, and the synaptic weights are identical for the same type of connections. The equations of each neuron are the same as the mean field equations (Eq 1, 2), but with each neuron receiving independent noise. The parameters of the model are $(\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}) = (0.5, 0.6, 0.7, 1)$.

For tuning the full neural network to have similar selectivity trends as the experimental results of [15], we set the input magnitude μ_0 to 0.0052, the excitatory noise standard deviation D to 5.0 and the inhibitory noise standard deviation is $D = 2.5$ (see Eq S5). To account for additional events that might happen after the decision, we added a mechanism that resets the network. 150 ms after the network has reached its decision, the excitatory neurons stop receiving inputs, and the inhibitory neurons begin receiving a small bias current of 0.002. This would cause a decline in the network activity, and thus reset the network.

When tuning the network to have similar correlation trends as those of [15], we set the excitatory noise SD D to be 6.0, and the inhibitory noise SD is $D = 1.0$. We set a weak correlation coefficient ($= 0.05$) between the noise inputs of every pairs of neurons in the network. No input aside from noise is given to the network, so as to mimic spontaneous activities. Note that the full neural network is used only to compare to the experimental data of Najafi et al., and for all other experimental comparisons and mathematical analyses, we used the aforementioned reduced models of UM and SM.

Energy Landscape

To gain a more intuitive understanding of the underlying dynamical system, we calculated an energy-like function U (Lyapunov function; energy landscape), which governs the dynamics of gating variables. In particular, the "valleys" of the energy landscape correspond to the stable attractors of the dynamical system, and the "hilltop" the unstable attractors. The dynamics of the system would seek to minimize its value, which means that the system behaves like a ball that "runs towards" the valleys.

In terms of our models, the energy landscape can be solved semi-analytically for the WWM and linear reduced model. The time-dependent linear reduced model, with its richer dynamics, on the other hand exhibits a time-varying energy landscape. Hence, instead of directly calculating U in these systems, the energy landscape of

all the models will instead be approximated by averaging over simulation trials [28], and used as a qualitative visualization tool for understanding the properties of the dynamical system.

For a dynamical system $\dot{x} = F(x)$, its energy landscape $U(x)$ is associated to $F(x)$ by $F(x) = -dU/dx$ [28, 31]. Letting $x = S_{E1} - S_{E2}$, the value of $F(x)$ and consequentially the value of $U(x)$ can be obtained via simulation. $F(x)$ is simply $(x(t + \Delta) - x(t))/\Delta$, where Δ is the simulation time step. Since the individual trials are noisy, these values are averaged across 2000 trials to obtain a smooth function, $\bar{F}(x)$. $U(x)$ is then an integration across $\bar{F}(x)$ in terms of x. The energy landscape obtained would resemble a double-well potential, where the two "valleys" correspond to the two choices presented to the system. The system will be forced to move towards one of the valleys, which corresponds to it making a decision (Fig. 1b).

Simulation Protocol

The simulations, unless stated otherwise, always start at the stable equilibrium point where $S_{E1} = S_{E2}$. The input signals for the two excitatory populations are

$$
\begin{aligned}
I_1 &= \mu_0(1 + \frac{c}{100}) \\
I_2 &= \mu_0(1 - \frac{c}{100}),
\end{aligned}
\tag{9}
$$

where $\mu_0 = 0.0156$ (nA), and c represents the coherence of the signal. A large c corresponds to a random dots motion test in which the majority of points move in the same direction (the one that favors population 1). The values of c used here are the same as those used in [32]: 0, 3.2, 6.4, 12.8, 25.6 and 51.2%. The lower the coherence, the higher the variance of the accuracy. Therefore, the trials ran N decreases for higher c: $1500, 1500, 1500, 1000, 250, 100$ are run for the respective coherence levels.

At $t = 0.5$ (s), the input signals are turned on, and the network begins to make a decision. A decision is defined as one of the populations' NMDA gating variable exceeding a threshold of 0.35 (unless stated otherwise), and the winner is defined as the first population to do so. The reaction time is then the time it takes for the winner to reach the threshold. The total simulation time of a trial is 3 (s), and if the network is undecided until that point, the winner is randomly assigned and the reaction time is marked as 2.5 (s). These undecided events seldom occur, especially when the threshold is low. For reference, none of the UMs had undecided trials out of a total of 1500 trials when the threshold is set to 0.35. When the threshold is increased to 0.6 (the largest threshold used in this study), excluding these undecided trials caused a change of $\mathcal{O}(\frac{1}{100})$ (s) in the reaction time of incorrect trials, but did not make a difference in the reaction time of correct trials.

All of the ODEs are solved by the fourth-order Runge-Kutta method with a time step of 0.1 ms.

Statistical Analysis

Speed-Accuracy Trade-off

A trade-off between accuracy and reaction time seems inevitable, but not all models are efficient in its trade-off. To quantify whether the trade-off of a model is efficient compared to others, we first parameterized the psychometric function by this form [29]:

$$p = 0.5 + 0.5(1 - e^{-(\frac{c}{\alpha_p})^{\beta_p}}), \tag{10}$$

where p is the accuracy, c the coherence, α_p the discrimination threshold and β_p the psychometric order. Essentially, α_p controls the coherence that hits 81.6% correct, and β_p controls the slope at said threshold. We can then use α_p as a measurement of accuracy for the model: the smaller it is, the better.

The reaction time, since it decreases monotonically as a function of coherence, will be represented by the average reaction time when coherence equals zero (α_{RT}). Here, both accuracy and reaction time are bootstrap sampled 1000 times to obtain its standard error.

Finally, we can plot the accuracy versus reaction time for each parameter set. Any parameter set that is on the upper right corner of another parameter set is less efficient. For example, if we have reaction time as the x-axis and accuracy as the y-axis, and if point B is located in the upper-right direction of point A, this indicates that it has a longer reaction time compared to parameter set A, yet performs worse.

Non-Regret Choice

During the initial stage, the system walks the separatrix, alternating between the basins of attraction of S_{E1} and S_{E2}. For a given deviation from the separatrix, e.g. $S_{E1} - S_{E2} = 0.04$, the system might continue falling into the favored attractor, S_{E1}, or it might end up in the other one. Conceivably, the larger the deviation, the less likely it is for the system to "change its mind". We call the value of the deviation a benchmark, in order to distinguish it from the threshold (the threshold is S_{E1} or S_{E2} reaching some assigned value; a benchmark is $|S_{E1} - S_{E2}|$ reaching some assigned value).

For each benchmark, we can calculate the probability of the system ending up in the attractor it currently favors. We call this the non-regret choice percentage. As a trivial example, consider having coherence $c = 0$ and benchmark equal to zero: the non-regret choice percentage is roughly 0.5, because only half of the trials that start out favoring one side will actually converge to that side. Comparing how the non-regret choice percentage changes as a function of the benchmark would reveal how careful the system is weighing its options. For instance,

if the percentage is high for a small benchmark value, this means that as soon as $|S_{E1} - S_{E2}|$ is exceeds that benchmark, the system seldom changes its choice.

Results

Selective Model Conforms with Experimental Data

To show that the selective model (SM) as a generalization of attractor model is more biologically realistic, we compared the SM to some of the existing experimental data. We first look at the basics of a decision making task, which are the psychometric and chronometric functions. A decision making model can be fit to random dots motion task data [33] by adjusting the standard deviation of the noise, D (see Eq S5). Both the UM and SM can be reasonable fits to the experimental data, but the SM requires lower D values to achieve the experimental values of α_p and β_p, which already hints at the efficiency of the SM. The psychometric and chronometric functions for the experimental data is shown in Fig. 1c, and those for the SM with $(\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}) = (0.5, 0.6, 0.7, 1)$ are shown in Fig. 1d.

Aside from the psychometric and chronometric functions, the other important property for a decision making network its selectivity, as shown by Najafi *et al.* To better fit our selective model to the experimental result, we constructed a larger neural network that contains 50 neurons in each population. The connections are all-to-all, the synaptic weights are identical for all weights of a particular type, and each neuron receives independent noise. An additional bias is given to the inhibitory populations 150 ms after it has reached its decision, to account for the system resetting itself (see Methods). While this is still quite idealistic for a neural network, its results have a similar selectivity trend as those seen in the study of Najafi *et al.* (Fig. 1e, f).

A further prediction of having a SM configuration is that neuron pairs with the same selectivity would have a higher correlation than pairs with opposite selectivity, since their connection weights are stronger. This is shown experimentally by Najafi *et al.*, where correlation of the same selectivity pairs are larger than the opposite pairs, and both are non-negative (Fig. 1g). However, in the simulations, we note that if each neuron received uncorrelated noise, the Pearson correlation of neurons with the opposite selectivity would be negative. This is because in the SM configuration, the two excitatory-inhibitory pairs are naturally in competition with one another, which would result in their activities moving in opposite trends. This argues for the presence of weakly correlated noise between neurons. Indeed, when adding weakly correlated noise, the SM shows the same trend in correlations as the experimental data (Fig. 1h).

An interesting detail to note here is that both in tuning for selectivity and correlation, the inhibitory neurons

must receive less noise than the excitatory ones to produce results that are qualitatively similar to the experimental data. For example, if they are given equal amounts of noise, the EE pairs would end up having a much higher Pearson correlation than the II pairs, while the opposite is shown in experimental data (Fig. 1e). This is because excitatory neurons tend to synchronize with one another, while inhibitory neurons tend to move in an asynchronous manner.

Selectivity Improves Speed-Accuracy Trade-Off

While SMs explain experimental data better than the UM, whether selectivity is valuable in a computational sense is still undetermined. Selectivity can be quantified by the parameters γ_{ei} and γ_{ie}, where smaller γ_{ei} and γ_{ie} values would lead to a larger difference in the inter-population excitation and inter-population inhibition values, and thus to a more selective model. Speed and accuracy are the most commonly used metrics to evaluate a decision making model's performance, as they are the most crucial. An efficient trade-off of speed and accuracy, where the reaction time is low but the accuracy is high, is desirable. To compare the speed-accuracy trade-off of SM with UMs, we parameterized the psychometric and chronometric functions, and used the metric α_p to represent accuracy and α_{RT} to represent the speed (see Methods and Fig. 2a, b). The smaller α_p is, the more accurate the model; the smaller α_{RT} is, the faster the model. Therefore, a smaller $\alpha_p \times \alpha_{RT}$ value would indicate a more efficient model, and thus we define the trade-off performance of a model by the inverse of $\alpha_p \times \alpha_{RT}$. Another metric to compare to would be the reward rate, which is simply the total number of rewards across all trials over the total reaction time (in units of s). The reward is 1 for each correct trial, and the trials consist of randomly selected coherence levels $0, 3.2, 6.4, 12.8, 32.6, 51.2$, where each coherence level is equally likely to be selected. The trend of efficiency of the trade-off are similar across the two metrics (Fig. 2c, d).

Simulation results indicate that selectivity strongly correlates with the efficiency of the trade-off (Fig. 2c, d). We observe that the trade-off performance increases along with selectivity, up until a cut-off value at $\gamma_{ei} = 0.5$ (for simplicity, in Fig. 2c, d γ_{ie} is fixed at 0.7, while only γ_{ei} is varied). This is because the selectivity of a SM cannot be unrestricted – too much selectivity leads to an inability to make decisions. Intuitively, an extreme case of $\gamma_{ei} = \gamma_{ie} = 0$ has no feedback inhibition from the opposing inhibitory population at all, and with insufficient mutual inhibition between inhibitory populations, there will be no "competing" mechanism that allows one population to win over the other.

Figure 2: Speed-accuracy trade-off as a function of selectivity. (a) The psychometric function of a SM with $(\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}) = (0.5, 0.6, 0.7, 1)$, alongside the psychometric function of an UM with parameters $\gamma_{ee} = 0.5$. The definition of α_p, which is the coherence level when the model reaches 81.6% accuracy, is illustrated here. (b) The chronometric function of the SM and UM in (a). Solid circles are the reaction time for the correct trials, and hollow ones are for incorrect trials. The definition of α_{RT}, which is the average reaction time at $c = 0$, is illustrated. (c) Trade-off performance, defined as the inverse of $\alpha_p \times \alpha_{RT}$, increases with selectivity, defined as $1 - \gamma_{ei}$ here. $(\gamma_{ee}, \gamma_{ie}, \gamma_{ii})$ are fixed to be $(0.5, 0.7, 1)$. (d) Reward rate, in units of $\#reward/s$, increases with selectivity as well. The parameters are the same as (c).

Time Varying Energy Landscape

Why does selectivity lead to a more efficient speed-accuracy trade-off? To better analyze the dynamics of our system, we first reduce the three-variable UM and the four-variable SM into a two-variable form once again, without evoking assumption (III). That is, it can be shown that while assumption (III) does not hold true, the dynamics of S_G is largely determined by the average value of S_{E1} and S_{E2}, i.e. "how much" the system is activated. In fact, the relation between S_G and $S_+ := \frac{S_{E1}+S_{E2}}{2}$ is almost linear in all time periods under many parameter sets of the UM (Fig. S3a). For the SM, the relationship between $S_{G+} := \frac{S_{G1}+S_{G2}}{2}$ and S_+ is linear, and the relation between $S_{G-} := \frac{S_{G1}-S_{G2}}{2}$ and $S_- := \frac{S_{E1}-S_{E2}}{2}$ is quadratic (Fig. 3a).

Therefore, substituting S_G by $a^* S_+ + b^*$ for the UM, where a^*, b^* is the slope and intercept of the S_G-S_+ relation (Eq 6), we would again obtain a WWM Only this time, a different synaptic weight set would result in a different reduced . For the SM, we could similarly replace S_{G-} by $a_Q^* S_-^2 + b_Q^* S_-$, where a_Q^*, b_Q^* is the quadratic and linear coefficient of the S_{G-}-S_- relation. For analysis purposes, we linearized the quadratic relation during short time periods, so that the resulting reduced model resembles those of the linear reduced model except for a constant term (Eq 8). The a_Q^*, b_Q^* values are fitted to an unfair, noiseless trial released at $(S_{E1}, S_{E2}) = (0.15, 0.1)$, and the a_L^*, b_L^* values are subsequently fitted to the data points from different time periods (see Methods). The a_L values slowly increase in time, while the b_L value is initially positive then becomes negative, as one would expect from a quadratic function with positive coefficient for the quadratic and linear terms (Fig. S5a, 3b red lines). The effect of the change in these two values can be elucidated by looking at its influence on the synaptic currents (Eq 8, restated here for convenience):

$$I_{syn,1} = g_{ee,s}S_{E1} + g_{ee,o}S_{E2} - (g_{ie,s} + g_{ie,o})(aS_+ + b) - (g_{ie,s} - g_{ie,o})(a_L(t)S_- + b_L(t)) + I_e$$

$$I_{syn,2} = g_{ee,s}S_{E2} + g_{ee,o}S_{E1} - (g_{ie,s} + g_{ie,o})(aS_+ + b) + (g_{ie,s} - g_{ie,o})(a_L(t)S_- + b_L(t)) + I_e. \tag{11}$$

Since $g_{ie,s} - g_{ie,o}$ is positive in the cases of interest $((\gamma_{ei}, \gamma_{ie}) = (0.6, 0.7))$, when a_L increases, it slows down the system by decreasing $I_{syn,1}$ and $I_{syn,2}$. This is confirmed by the energy landscape of different a_L values, where the slope of the hill decreases for increasing a_L (Fig. S5b, where b_L is temporarily set to 0 to highlight the effect of altering a_L alone). Note that in contrary to the SM, the UM has a fixed energy landscape across time since it has no time varying parameters (Fig. 3c).

Unlike a_L, which has a symmetrical effect on both of the synaptic currents, b_L results in an asymmetry, where positive b_L increases $I_{syn,2}$ but decreases $I_{syn,1}$. Therefore, with an initially positive b_L, the energy landscape favors the attractor opposite of the one the system is currently attracted to. For instance, if initially $S_{E1} > S_{E2}$, then the attractor of S_{E2} has a larger basin of attraction (Fig. 3d, red line). However, since b_L eventually becomes

negative, this means that if the system persists in having $S_{E1} > S_{E2}$, the tables turn and the S_{E1} attractor has a larger basin of attraction instead (Fig. 3d, green line). These are also reflected in the psychometric functions of simulating different time-dependent linear reduced models, where initial $\{a_L, b_L\}$ values have accuracy less than 0.5 at coherence $c = 0$, which corresponds to the system favoring the other side (Fig. S5d, red); later $\{a_L, b_L\}$ values have accuracies of 1 for all coherence levels, which signifies a disappearance of the attractor of the less favored side (Fig. S5c,d, purple).

Together, this means that the system carefully weighs the information in the beginning by having a larger basin of attraction at the opposing side (Fig. 3e, top right; Fig. 3f). If the evidence for the currently favored side is insufficient, the system would be attracted to the other side instead. However, if the evidence is sufficient, then the system quickly changes to a configuration that favors the current side, leading to faster convergence to the favored attractor (Fig. 3e, bottom right). The "careful consideration" part is evident in the percentage of non-regret choices, where the non-regret choices for the SM increases slower than the UM (Fig. 4a, also see Methods). That is, if $S_{E1} - S_{E2}$ is, say, 0.04 in the UM, then it has a high chance of having S_{E1} as the winner. In contrast, for the same situation in the SM, the chance of S_{E1} winning is lower, which means that despite S_{E1} having a slight edge on S_{E2}, the system is still considering its options.

Further statistical analysis done on the reaction time supports the conclusion of the "quick convergence" part. We first calculated the time it takes for the system to travel between thresholds (Fig. 4b). Setting the threshold to different values will result in different reaction times, and calculating the difference in reaction time between successive thresholds would allow us to know how much time it takes to reach one threshold from the previous one. Initially, the SM takes more time than some of the faster UMs to reach the next threshold. This is overturn in later stages, where the SM becomes the fastest one.

Lastly, if during the trial, the system keeps switching back and forth between S_{E1} and S_{E2}, then b_L turns negative faster, which corresponds to strengthening the favored side faster (Fig. 3b, green and blue lines). This means that if the evidence does not favor either side, and it is taking too long to make a decision, then the system escalates the process by making it easier for the favored side to win. This can be showed by grouping trials using the S_+ value at the moment it crosses a certain benchmark (set to 0.05 here to be consistent with Fig. 3b). A larger S_+ value at the benchmark signifies that the trial walks the separatrix longer than others. For each group, we can then calculate the time it takes for b_L to turn negative after the trial crosses the benchmark. This value decreases as S_+ increases, meaning that if the trial takes too long, then b_L turns negative faster, which in turns means the system allows the decision to converge faster to its attractors (Fig. 4c). This could prevent the system from stalling.

The combined effect of "quick convergence" and "avoid stalling" is also evident in the distribution of the reaction time (Fig. 4d). The tail of the reaction time distribution of the SM is significantly shorter than the UMs, indicating that it has less trials with longer reaction times.

We note that the "converge quickly" portion seems to have a larger effect than the "choose carefully" portion on the speed-accuracy trade-off. That is, the changes selectivity have on the accuracy (Fig. 5a) is less notable compared to the changes in reaction time (Fig. 5b). Therefore, when evaluated in terms of the trade-off performance ($\frac{1}{\alpha_p \times \alpha_{RT}}$), the more selective the model is, the more efficient it seems to be. However, even if we constrained the drop in accuracy to be within a certain regime, the SMs still out perform the UMs. This is evident by the fact that the SMs lie of the lower-left corner of the α_p-α_{RT} plane, which corresponds to a more efficient speed-accuracy trade-off.

Model Reduction via Empirical Relations

We have shown that the key to the more efficient trade-off performance of the SM is due to its time-varying energy landscape, and that the time dependence arises from the quadratic relation of S_{G-} with S_{-}. To better understand how the linear relation of S_{G+} with S_{+} and the quadratic relation of S_{G-} with S_{-} arises, we turn to the equations governing the GABA dynamics. According to assumption (II), since the f-I curve of the inhibitory population can be linearized, the equations of S_{G1} and S_{G2} are linear. Hence we can write down the equation of motion for S_{G+} as follows:

$$
\begin{aligned}
\dot{S}_{G+} &= -\left(\frac{1}{\tau_{\text{GABA}}} + 307.5 g_{ii,s}(1 + \gamma_{ii})\right) S_{G+} + 307.5 g_{ei,s}(1 + \gamma_{ei}) \, S_{+} \\
&= -c_{+,relax} \, S_{G+} + c_{+,inp} \, S_{+}.
\end{aligned}
\tag{12}
$$

And the equation of motion for S_{G-} is:

$$
\begin{aligned}
\dot{S}_{G-} &= -\left(\frac{1}{\tau_{\text{GABA}}} + 307.5 g_{ii,s}(1 - \gamma_{ii})\right) S_{G-} + 307.5 g_{ei,s}(1 - \gamma_{ei}) \, S_{-} \\
&= -c_{-,relax} \, S_{G-} + c_{-,inp} \, S_{-}.
\end{aligned}
\tag{13}
$$

The number 307.5 comes from linearizing $\phi(x)$ in Eq 1 (see Eq S3). The time constants for S_{G+} and S_{G-} are $\frac{1}{c_{+,relax}}$ and $\frac{1}{c_{-,relax}}$, respectively. Since γ_{ii} is always larger or equal to zero, $c_{+,relax}$ must also be larger or equal to $c_{-,relax}$. In the strictly larger scenario, this means that S_{G+} will always move towards steady state at a faster speed than S_{G-}. This can be shown by plotting the lag of a noiseless trial for a SM, where S_{G-} clearly lags behind more than S_{G+} (Fig. 6a, bottom panel).

The steady state of Eq 12 and Eq 13 yields a linear relation between both S_{G+} and S_{+}, as well as S_{G-} and S_{-}. However, it can be shown that the lag between the true dynamics and the steady state is non-neglectable

Figure 3: Reduced models and energy landscapes of UM and SM. (a) Quadratic relationship between S_- and S_{G-}. The parameters for $(\gamma_{ee}, \gamma_{ei}, \gamma_{ii})$ are $(0.3, 0.6, 1)$. $log(e)$ as a function of $log(S_-)$, where e is the residual of a linear fit between S_{G-} and S_-. The slope of the resulting log-log plot is around 2, indicating that it is a quadratic function. The inset plot shows how $a_L(t)$ and $b_L(t)$ is derived from the quadratic relation. (b) Evolution of the time-varying intercept b_L as a function of the starting point $((S_{E1}, S_{E2}) = (S_+ + 0.05, S_+))$ of the no-noise trial. In the following panels, the linearized quadratic models are indexed by the time period in which they are fitted to. (c) The energy landscape U of a UM with $\gamma_{ee} = 0.5$. (d) The energy landscape U of a SM with $(\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}) = (0.5, 0.6, 0.7, 1)$. (e) Schematic diagram of how U evolves in time for the UM and SM. The orange ball indicates the position of the system. The UM landscape is fixed in time, while for the SM, the landscape favors the opposing attractor in initial stages, leading to careful weighting of options; in later stages the opposing attractor disappears, leading to faster reaction times. (f) A schematic diagram of the initial stages of the phase diagram for a SM.

Figure 4: Statistical results for accuracy and reaction time that elucidates the effects of the time-varying energy landscape. The SM for all panels uses $(\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}) = (0.5, 0.6, 0.7, 1)$. (a) Percent non-regret choice as a function of benchmark. A lower percent non-regret choice in smaller benchmark values means that the model chooses more carefully. (b) The average time it takes to reach the next threshold, starting from the current threshold. The x-axis is the value of the next threshold, and the starting threshold is the next threshold minus 0.1. The SM is the fastest during the later stages. (c) X-axis: S_+ value of the trial when it reaches a benchmark of 0.05. Y-axis: Time until b_L becomes negative after the trial crosses a benchmark of 0.05 (the stalling time); mean $\pm$ SEM. The smaller this value is, the faster the energy landscape turns to favor the current attractor. (d) Cumulative probability density function of the reaction time for coherence $c = 0$ trials. The distribution of the data is fit to a gamma distribution. The SM does not have a long tail, compared to the UMs.

Figure 5: α_p and A_{RT} as a function of selectivity. (a) α_p as a function of selectivity. $(\gamma_{ee}, \gamma_{ii})= (0.5, 1)$. Regions that cannot make decisions are labeled as "no decisions". (b) α_{RT} as a function of selectivity. (c) Trade-off performance as a function of selectivity. (d) The distribution of models on the α_p-α_{RT} plane. *Inset:* The same graph with an extra data point $(\gamma_{ei}, \gamma_{ie})= (0.5, 0.7)$ included.

(defined as $S_{G+} - S_{G+,ss}$ for S_{G+} ($S_{G-} - S_{G-,ss}$ for S_{G-}, where ss stands for steady state; see Supplementary Note S2 and Fig. S2e). This is why assumption (III) doesn't hold in the first place. Therefore, to gain a more accurate description for these relations, we must take into account the modifications caused by the lags. The lag of S_{G+} is roughly a linear decline, which means that S_{G+} and S_+ will more or less retain its linear relation, albeit with a different slope and intercept than the steady state relation. On the other hand, the lag of S_{G-} is bell shaped, which means that the subtraction of its lag from the steady state relation will yield a concave up and increasing curve (Fig. 6b). This is the quadratic relation we observe in Fig. 3b. We can verify this by plotting a_Q as a function of the peak of the lag (Fig. 6c).

The shape of the lags are in turn a result of the velocity flow of S_+ and S_- (Fig. 6a, top panel). While the velocity of S_+ is roughly linear with a slight decline late in the trial, the velocity of S_- is hyperbolic tangent like. This means that as the speed of S_- increases in the initial stages, it becomes more challenging for S_{G-} to keep up, resulting in an increasing lag. As the speed decreases, the lag then becomes less severe. This would

thus result in a bell shaped lag. Therefore, to obtain a more quadratic curve, the peak of the lag must be large.

The magnitude of the lag is controlled by both $c_{-,relax}$ and $c_{-,inp}$. In our models, $g_{ii,s}$ and γ_{ii} are fixed, so every

model has the same $c_{-,relax}$ value. In that case, the lag will be proportional to $c_{-,inp}$, as we indeed see in Fig.

6d.

Figure 6: Relaxation dynamics of inhibitory average S_{G+} and inhibitory difference S_{G-}. Since the inhibitory dynamics are fast, its relation with the excitatory dynamics are dictated by the evolution of S_+ and S_-. The lag between the inhibitory dynamics' true state and its steady state is directly related to the "swiftness" of the relaxation term ($c_{-,relax}$) relative to the input term ($c_{-,inp}$). (a) *Top panel*: S_+ and S_- in a biased, noiseless trial that starts at $(S_{E1}, S_{E2}) = (0.25, 0.26)$. Since it is a noiseless trial, the input starts at $t = 0$ (s) instead of $t = 0.5$ (s). *Bottom panel*: The lag of S_{G+} ($S_{G+} - S_{G+,ss}$, where ss stands for steady state) and S_{G-} ($S_{G-} - S_{G-,ss}$). The evolution of S_+ is linear (in the initial stages), while those of S_- is not. (b) Schematic diagram of how the quadratic relation arises from the bell-shaped lag. (c) a_Q as a function of the peak of the lag of S_{G-}. The models, from left to right, are SMs with: $(\gamma_{ee}, \gamma_{ei}, \gamma_{ie}, \gamma_{ii}) = (0.5, 0.6, 0.7, 1), (0.3, 0.6, 0.5, 1), (0.5, 0.6, 0.8, 1), (0.3, 0.6, 0.6, 1), (0.3, 0.6, 0.7, 1), (0.3, 0.6, 0.5, 1)$ and $(0.3, 0.8, 1.25, 1)$. The rightmost dot is an UM with $\gamma_{ee} = 0.5$. (d) The peak of the lag as a function of $c_{-,inp}$. The order of the models are the same as (b).

Discussion

Our study indicates that selectivity in the inhibitory population creates a time-dependent energy function that is more efficient in its speed-accuracy trade-off. Furthermore, it emphasizes the importance of the role the GABA dynamics play, despite being 20 times faster than NMDA dynamics. This topic can be further extended in several aspects.

First, the SM introduces a rudimentary level of generalization of the attractor model. Specifically, it is found that heterogeneity is commonly observed in brain areas associated with decision making [34, 35], and subdividing the inhibitory populations into two is a first step towards the inclusion of heterogeneity. If this one step of separation elicits different dynamics, we could expect more complicated forms of functionality from an even higher level of heterogeneity. Together with other studies that demonstrates how heterogeneity aids in encoding information regarding different aspects of the stimulus [36], this highlights the importance of incorporating heterogeneity into future decision making models.

Another important concept from decision making literature is excitation-inhibition balance. E/I balance could refer to either balanced top-down input currents, or the balance of excitation-to-inhibition synaptic currents for the neurons. In terms of the former, it is previously shown that a network with balanced inputs is capable of making more accurate decisions via having a time-varying energy landscape [30]. This time-variance is different from those mentioned in our study, as the variations of the landscape occurs symmetrically to both of the attractors in this case, and asymmetrically in our case. Furthermore, this time-variance requires external top-down control [30, 37], while in the SM, this property is naturally embedded in local circuits whose configuration is supported by the latest experimental observations. The interaction between SM and balanced external inputs, both of which are time-varying components on their own, requires further investigation.

The balancing of the synaptic currents too warrant a deeper look. In our simulations, altering synaptic weights result in drastically different dynamics, as can be seen from the results of altering γ_{ee}. In fact, altering a single synaptic weight by 0.5% can significantly speed up or slow down the reaction time of the system. Previous studies have shown that an imbalance in excitation/inhibition ratio changes the system's behavior: over-excited, and the system tends towards impulsive decisions; over-inhibited, and it becomes indecisive [29]. This picture of balanced excitation/inhibition ratio is a widely explored topic in the field of neuroscience, both experimentally and theoretically. The introduction of two inhibitory populations adds another layer of complexity to the balanced picture, as the balance is now controlled by two inhibitory populations and their interactions simultaneously. How E/I balance may come into play after this complication, and how the dynamics may change accordingly, is an

open question.

The process of learning how to perform these decision tasks is another topic that has gained attention [38]. An interesting result in the study of Najafi *et al* is that equal selectivity between excitatory and inhibitory populations is maintained throughout the course of learning. Is selectivity a property that manifests naturally as a consequence of learning? Specifically, given that SMs are more efficient in their speed-accuracy trade-off than the UMs, could selectivity emerge out of a non-selective network automatically as it learns? Could the network self-organize into a more efficient state of selectivity?

Aside from broad concepts of heterogeneity, E/I balance and learning, another piece of literature to integrate or compare the SM to would be other types of decision making models. In particular, we note how the property of the time-varying landscape has an "urgency" component for trials that take a longer time. That is, if the system is indecisive for too long, the landscape would adjust itself so that convergence to an attractor becomes easier. One of the decision making models proposed in the past is the urgency gating model, which shares the same concept of having an urgency signal that rushes the decision making process along in later stages. However, the SM is mechanistically different from the urgency gating model, which bases decisions on momentary information instead of integrated information [39, 40, 41]. In this regard, the SM is more similar to the diffusion model, with differences such as time-shift invariance pointed out in previous studies [10, 11]. A more thorough analysis of experimental evidence for each of these models may shed light on how these models may be related to one another.

Last but not least, the details that are uncovered in this study are also worth pursuing. For instance, the new reduction reduces the UM and SM back to two variables in a more realistic way. How might the linear or quadratic relationship depend on the synaptic weights? Why do such small deviations from linearity impact the dynamics? How does the GABA dimension change the energy landscape of the system? In our studies, the contribution of AMPA is neglected. However, our results suggest that the inclusion of GABA dynamics significantly alter the decision making network's accuracy and reaction time; hence it is entirely probable that an inclusion of AMPA would also impact the dynamics. In conclusion, our study supports the idea of a more complicated picture of decision making models than the widely studied UM, an adjustment that leads to better computational abilities.

Acknowledgements

We thank Alexander J. White, Cheng-Te Wang and Ching-Che Charng for discussion and feedback on the manuscript.

References

[1] Xiao-Jing Wang. Neural dynamics and circuit mechanisms of decision-making. *Current Opinion in Neurobiology*, 22(6):1039–1046, December 2012.

[2] Xiao-Jing Wang. Decision Making in Recurrent Neuronal Circuits. *Neuron*, 60(2):215–234, October 2008.

[3] Rajesh P. N. Rao. Decision Making Under Uncertainty: A Neural Model Based on Partially Observable Markov Decision Processes. *Front. Comput. Neurosci.*, 4, 2010. Publisher: Frontiers.

[4] Jeffrey M. Beck, Wei Ji Ma, Roozbeh Kiani, Tim Hanks, Anne K. Churchland, Jamie Roitman, Michael N. Shadlen, Peter E. Latham, and Alexandre Pouget. Probabilistic Population Codes for Bayesian Decision Making. *Neuron*, 60(6):1142–1152, December 2008.

[5] Christian K. Machens, Ranulfo Romo, and Carlos D. Brody. Flexible control of mutual inhibition: a neural model of two-interval discrimination. *Science*, 307(5712):1121–1124, February 2005.

[6] Chung-Chuan Lo and Xiao-Jing Wang. Cortico-basal ganglia circuit mechanism for a decision threshold in reaction time tasks. *Nat Neurosci*, 9(7):956–963, July 2006.

[7] Rafal Bogacz, Eric Brown, Jeff Moehlis, Philip Holmes, and Jonathan D. Cohen. The physics of optimal decision making: a formal analysis of models of performance in two-alternative forced-choice tasks. *Psychol Rev*, 113(4):700–765, October 2006.

[8] Drew Fudenberg, Whitney Newey, Philipp Strack, and Tomasz Strzalecki. Testing the drift-diffusion model. *PNAS*, 117(52):33141–33148, December 2020.

[9] Xiao-Jing Wang. Probabilistic Decision Making by Slow Reverberation in Cortical Circuits. *Neuron*, 36(5):955–968, December 2002.

[10] Kong-Fatt Wong, Alexander C. Huk, Michael N. Shadlen, and Xiao-Jing Wang. Neural circuit dynamics underlying accumulation of time-varying evidence during perceptual decision making. *Front. Comput. Neurosci.*, 1, 2007. Publisher: Frontiers.

[11] Kong-Fatt Wong and Alexander C. Huk. Temporal dynamics underlying perceptual decision making: Insights from the interplay between an attractor model and parietal neurophysiology. *Front. Neurosci.*, 2, 2008. Publisher: Frontiers.

[12] Praveen K. Pilly and Aaron R. Seitz. What a difference a parameter makes: A psychophysical comparison of random dot motion algorithms. *Vision Research*, 49(13):1599–1612, July 2009.

[13] Mark E. Mazurek, Jamie D. Roitman, Jochen Ditterich, and Michael N. Shadlen. A Role for Neural Integrators in Perceptual Decision Making. *Cerebral Cortex*, 13(11):1257–1269, November 2003.

[14] Alexander C. Huk and Michael N. Shadlen. Neural Activity in Macaque Parietal Cortex Reflects Temporal Integration of Visual Motion Signals during Perceptual Decision Making. *J. Neurosci.*, 25(45):10420–10436, November 2005. Publisher: Society for Neuroscience Section: Behavioral/Systems/Cognitive.

[15] Farzaneh Najafi, Gamaleldin F. Elsayed, Robin Cao, Eftychios Pnevmatikakis, Peter E. Latham, John P. Cunningham, and Anne K. Churchland. Excitatory and Inhibitory Subnetworks Are Equally Selective during Decision-Making and Emerge Simultaneously during Learning. *Neuron*, 105(1):165–179.e8, January 2020.

[16] Kazuhiro Sohya, Katsuro Kameyama, Yuchio Yanagawa, Kunihiko Obata, and Tadaharu Tsumoto. GABAergic neurons are less selective to stimulus orientation than excitatory neurons in layer II/III of visual cortex, as revealed by in vivo functional Ca2+ imaging in transgenic mice. *J Neurosci*, 27(8):2145–2149, February 2007.

[17] Cristopher M. Niell and Michael P. Stryker. Highly Selective Receptive Fields in Mouse Visual Cortex. *J. Neurosci.*, 28(30):7520–7536, July 2008. Publisher: Society for Neuroscience Section: Articles.

[18] Aaron M. Kerlin, Mark L. Andermann, Vladimir K. Berezovskii, and R. Clay Reid. Broadly tuned response properties of diverse inhibitory neuron subtypes in mouse visual cortex. *Neuron*, 67(5):858–871, September 2010.

[19] Davi D. Bock, Wei-Chung Allen Lee, Aaron M. Kerlin, Mark L. Andermann, Greg Hood, Arthur W. Wetzel, Sergey Yurgenson, Edward R. Soucy, Hyon Suk Kim, and R. Clay Reid. Network anatomy and in vivo physiology of visual cortical neurons. *Nature*, 471(7337):177–182, March 2011. Number: 7337 Publisher: Nature Publishing Group.

[20] Sonja B. Hofer, Ho Ko, Bruno Pichler, Joshua Vogelstein, Hana Ros, Hongkui Zeng, Ed Lein, Nicholas A. Lesica, and Thomas D. Mrsic-Flogel. Differential connectivity and response dynamics of excitatory and in-

hibitory neurons in visual cortex. *Nature Neuroscience*, 14(8):1045–1052, August 2011. Number: 8 Publisher: Nature Publishing Group.

[21] Petr Znamenskiy, Mean-Hwan Kim, Dylan R. Muir, Maria Florencia Iacaruso, Sonja B. Hofer, and Thomas D. Mrsic-Flogel. Functional selectivity and specific connectivity of inhibitory neurons in primary visual cortex. *bioRxiv*, page 294835, April 2018. Publisher: Cold Spring Harbor Laboratory Section: New Results.

[22] Caroline A. Runyan, James Schummers, Audra Van Wart, Sandra J. Kuhlman, Nathan R. Wilson, Z. Josh Huang, and Mriganka Sur. Response features of parvalbumin-expressing interneurons suggest precise roles for subtypes of inhibition in visual cortex. *Neuron*, 67(5):847–857, September 2010.

[23] Alexandra K. Moore and Michael Wehr. Parvalbumin-expressing inhibitory interneurons in auditory cortex are well-tuned for frequency. *J Neurosci*, 33(34):13713–13723, August 2013.

[24] Greg Allen, Roderick Mccoll, Holly Barnard, Wendy Ringe, James Fleckenstein, and Munro Cullum. Magnetic resonance imaging of cerebellar–prefrontal and cerebellar–parietal functional connectivity. *NeuroImage*, 28:39–48, November 2005.

[25] Lucas Pinto and Yang Dan. Cell-Type-Specific Activity in Prefrontal Cortex during Goal-Directed Behavior. *Neuron*, 87(2):437–450, July 2015.

[26] Matthew Lovett-Barron, Patrick Kaifosh, Mazen A. Kheirbek, Nathan Danielson, Jeffrey D. Zaremba, Thomas R. Reardon, Gergely F. Turi, René Hen, Boris V. Zemelman, and Attila Losonczy. Dendritic inhibition in the hippocampus supports fear learning. *Science*, 343(6173):857–863, February 2014.

[27] Valérie Ego-Stengel and Matthew A. Wilson. Spatial selectivity and theta phase precession in CA1 interneurons. *Hippocampus*, 17(2):161–174, 2007.

[28] Kong-Fatt Wong and Xiao-Jing Wang. A Recurrent Network Mechanism of Time Integration in Perceptual Decisions. *J. Neurosci.*, 26(4):1314–1328, January 2006.

[29] Norman H. Lam, Thiago Borduqui, Jaime Hallak, Antonio C. Roque, Alan Anticevic, John H. Krystal, Xiao-Jing Wang, and John D. Murray. Effects of Altered Excitation-Inhibition Balance on Decision Making in a Cortical Circuit Model. *bioRxiv*, page 100347, January 2017. Publisher: Cold Spring Harbor Laboratory Section: New Results.

[30] Cheng-Te Wang, Chung-Ting Lee, Xiao-Jing Wang, and Chung-Chuan Lo. Top-Down Modulation on Perceptual Decision with Balanced Inhibition through Feedforward and Feedback Inhibitory Neurons. *PLOS ONE*, 8(4):e62379, April 2013. Publisher: Public Library of Science.

[31] Alex Roxin and Anders Ledberg. Neurobiological Models of Two-Choice Decision Making Can Be Reduced to a One-Dimensional Nonlinear Diffusion Equation. *PLOS Computational Biology*, 4(3):e1000046, March 2008. Publisher: Public Library of Science.

[32] Chung-Chuan Lo. Dynamic tuning of perceptual decision making in a cortical circuit model by balanced synaptic input. *Frontiers*, January 2010.

[33] Jamie D. Roitman and Michael N. Shadlen. Response of Neurons in the Lateral Intraparietal Area during a Combined Visual Discrimination Reaction Time Task. *J. Neurosci.*, 22(21):9475–9489, November 2002. Publisher: Society for Neuroscience Section: Articles.

[34] Mattia Rigotti, Omri Barak, Melissa R. Warden, Xiao-Jing Wang, Nathaniel D. Daw, Earl K. Miller, and Stefano Fusi. The importance of mixed selectivity in complex cognitive tasks. *Nature*, 497(7451):585–590, May 2013.

[35] Joseph K. Jun, Paul Miller, Adrián Hernández, Antonio Zainos, Luis Lemus, Carlos D. Brody, and Ranulfo Romo. Heterogenous Population Coding of a Short-Term Memory and Decision Task. *J. Neurosci.*, 30(3):916–929, January 2010. Publisher: Society for Neuroscience Section: Articles.

[36] Mikio C. Aoi, Valerio Mante, and Jonathan W. Pillow. Prefrontal cortex exhibits multidimensional dynamic encoding during decision-making. *Nature Neuroscience*, 23(11):1410–1420, November 2020. Number: 11 Publisher: Nature Publishing Group.

[37] Chung-Chuan Lo, Cheng-Te Wang, and Xiao-Jing Wang. Speed-accuracy tradeoff by a control signal with balanced excitation and inhibition. *Journal of Neurophysiology*, 114(1):650–661, July 2015. Publisher: American Physiological Society.

[38] Nicholas A. Roy, Ji Hyun Bak, International Brain Laboratory, Athena Akrami, Carlos D. Brody, and Jonathan W. Pillow. Extracting the dynamics of behavior in sensory decision-making experiments. *Neuron*, 109(4):597–610.e6, February 2021.

[39] Jochen Ditterich. Evidence for time-variant decision making. *Eur J Neurosci*, 24(12):3628–3641, December 2006.

[40] Paul Cisek, Geneviève Aude Puskas, and Stephany El-Murr. Decisions in Changing Conditions: The Urgency-Gating Model. *J. Neurosci.*, 29(37):11560–11571, September 2009. Publisher: Society for Neuroscience Section: Articles.

[41] David Thura, Julie Beauregard-Racine, Charles-William Fradet, and Paul Cisek. Decision making by urgency gating: theory and experimental support. *Journal of Neurophysiology*, 108(11):2912–2930, September 2012. Publisher: American Physiological Society.