Title:

Clinical and immunological signatures of severe COVID-19 in previously healthy patients with clonal hematopoiesis

Authors:

Chang Kyung Kang[1,*], Baekgyu Choi[2,*], Sugyeong Kim[3], Seongwan Park[2], Soon Ho Yoon[4], Dohoon Lee[5], Andrew J. Lee[2], Yuji Ko[2], Euijin Chang[1], Jongtak Jung[1,6], Pyoeng Gyun Choe[1], Wan Beom Park[1], Eu Suk Kim[1,6], Hong Bin Kim[1,6], Nam Joong Kim[1], Myoung-don Oh[1], Suk-jo Kang[2], Kyuho Kang[7], Sun Kim[8,9], Hogune Im[3], Joohae Kim[10], Yong Hoon Lee[11], Jaehee Lee[11], Ji Yeon Lee[10,†], Joon Ho Moon[11,†], Kyoung-Ho Song[1,6,†], Youngil Koh[1,3,†], Inkyung Jung[2,†]

Affiliations:

[1]Department of Internal Medicine, Seoul National University College of Medicine, Seoul 03080, Republic of Korea.

[2]Department of Biological Sciences, Korea Advanced Institute of Science and Technology (KAIST), Daejeon 34141, Republic of Korea.

[3]Genome Opinion Inc., Seoul 04799, Republic of Korea

[4]Department of Radiology, Seoul National University College of Medicine, Seoul 03080, Republic of Korea.

[5]Bioinformatics Institute, Seoul National University, Seoul 08826, Republic of Korea

[6]Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam 13620, Republic of Korea.

[7]Department of Biology, Chungbuk National University, Cheongju, 28644, Republic of Korea

[8]Department of Computer Science and Engineering, College of Engineering, Seoul National University, Seoul 08826, Republic of Korea

[9]Interdisciplinary Program in Bioinformatics, College of Natural Sciences, Seoul National University, Seoul 08826, Republic of Korea

[10]Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, National Medical Center, Seoul 04564, Republic of Korea

[11]Department of Internal Medicine, Kyungpook National University Hospital, School of Medicine, Kyungpook National University, Daegu 41944, Republic of Korea

[*] These authors contributed equally to this work

[†] Correspondence to Ji Yeon Lee (fulgeo@nmc.or.kr), Joon Ho Moon (jhmoon@knu.ac.kr), Kyoung-Ho Song (khsongmd@gmail.com), Youngil Koh (snuhgo01@gmail.com), and Inkyung Jung (ijung@kaist.ac.kr)

Abstract

Identifying additional risk factors for COVID-19 severity in numerous previously healthy

patients without canonical clinical risk factors remains challenging. In this study, we

investigate whether clonal hematopoiesis of indeterminate potential (CHIP), a common aging-

related process that predisposes various inflammatory responses, may exert COVID-19 severity.

We examine the clinical impact of CHIP in 143 laboratory-confirmed COVID-19 patients. Both

stratified analyses and logistic regression including the interaction between canonical risk

factors and CHIP show that CHIP is an independent risk factor for severe COVID-19,

especially in previously healthy patients. Analyses of 60,310 single-cell immune transcriptome

profiles identify distinct immunological signatures for CHIP (+) severe COVID-19 patients,

particularly in classical monocytes, with a marked increase in pro-inflammatory cytokine

responses and potent IFN-γ mediated hyperinflammation signature. We further demonstrate

that the enhanced expression of CHIP (+) specific IFN-γ response genes is attributed to the

CHIP mutation-dependent epigenetic reprogramming of poised or bivalent *cis*-regulatory

elements. Our results highlight a unique immunopathogenic mechanism of CHIP in the

progression of severe COVID-19, which could be extended to elucidate how CHIP contributes

to a variety of human infectious diseases.

Main Text:

Introduction

A pandemic of coronavirus disease-19 (COVID-19) , an emerging infectious disease caused by severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2), has been being a global health threat of the century[1,2]. Epidemiologic data revealed that about 20% of patients underwent severe or critical course[3]. Although many clinical risk factors for severe illness including older age, comorbidities such as diabetes mellitus or hypertension, and morbid obesity have been found[4,5], we could still only partly explain the development of severe COVID-19. Indeed, there have been numerous cases of severe COVID-19 from previously healthy adults[6,7].

Clinical deteriorations such as acute respiratory distress syndrome or intensive care unit admission most commonly occur at around the 10th day of illness[8,9], when the viral loads are declining after the early peak[10,11]. This temporal discrepancy suggests immunological phenomenon may play an important role in the progression of COVID-19. High levels of circulating pro-inflammatory cytokines[12], aberrant hyperactivation of cytotoxic lymphocytes[13] or their infiltration in vital organs[14], or dysregulated monocytes and macrophages[15] have been proposed as mechanisms for pathologic immune responses in severe COVID-19.

Clonal hematopoiesis of indeterminate potential (CHIP) refers to a population of immune cells with acquired gene mutations, but without fulfilling diagnostic criteria for a hematologic malignancy[16]. As a majority of the genes associated with CHIP including *DNMT3A*, *TET2*, and *ASXL1* are involved in epigenetic regulation, CHIP may have a wide range of effects on immune function through altered chromatin activities[17]. There is growing evidence supporting a role for the CHIP mutations in altered immune function through effector cells such as monocytes/macrophages and their dysregulated cytokine/chemokine expression,

which account for the increased risk of cardiovascular disease in individuals with CHIP[18-21].

Since immunopathogenesis of such an adverse outcome of CHIP largely shares that of severe

COVID-19, we hypothesized that CHIP might contribute to the progression of COVID-19 with

its unique immune signature. Although recent reports describe the association between

acquired mutations in hematopoietic cells such as CH and severe COVID-19[22,23], there has

been a controversy on the clinical impact of CHIP in COVID-19[23-25]. However, if there exists

a distinct CHIP-related clinical deterioration mechanism of COVID-19, tailored therapeutics

could be of help to salvage these patients.

In this study, to determine the clinical significance of CHIP in COVID-19 severity,

especially in previously healthy adults, we analyzed thorough clinical, radiological, and

laboratory characteristics of patients with COVID-19. In addition, we explored immune

signatures using single-cell RNA expression data according to the presence of CHIP to suggest

how CHIP attributes to the immunologic responses in severe COVID-19. Lastly, since the

majority of mutations in CHIP are related to epigenetic regulators of DNA methylation and

heterochromatin formation, we investigated CHIP mutation dependent dysregulated epigenetic

gene regulation mechanisms involved in CHIP-specific immunopathogenesis.

Results

Impact of CHIP on severe COVID-19 in previously healthy patients

A total of 143 laboratory-confirmed COVID-19 patients were analyzed in this study (Fig. 1a).
Among those, 34 patients had CHIP (23.8%, Supplementary Table 1). *DNMT3A* (13 variants)
was the most common mutated gene, followed by *TET2* (7 variants) and *ASXL1* (5 variants).
Clinical characteristics were retrospectively reviewed using electronic medical record (EMR)
systems of each institution. It included age, sex, body mass index, presence of comorbidities,
details of oxygen and medical therapy, duration of hospital stay, and in-hospital mortality.
Serial laboratory findings including complete blood count with differential count and
chemistries were also collected. Cases with the highest ordinal scale 3 (need for supplemental
oxygen therapy via nasal cannula) or more were classified as severe COVID-19, while others
were classified with mild ones[3].

Baseline characteristics of these patients are shown in Supplementary Table 1. Median
(interquartile range [IQR]) ages were 73 (61—81) and 65 (50—75) years in those with or
without CHIP, respectively (Student's *t*-test, $P < 0.001$), as consistent with a common aging-
related property of CHIP. The presence of CHIP appears to contribute to COVID-19 severity
as severe COVID-19 tended to be more frequent in patients with CHIP than those without,
while the difference was statistically not significant (25/34, 73.5% vs. 65/109, 59.6%, Chi-
squared test, $P = 0.143$).

To precisely examine the clinical impact of CHIP in patients with COVID-19, we
conducted a hierarchical clustering analysis for baseline characteristics and examined the
distribution of CHIP among clusters (Fig. 1b). All continuous clinical information was
transformed into a range of 0 to 1 using a logistic function, and comorbidity status was
dichotomized into 0 and 1 for absence and presence (see Methods). Patients were grouped into

two main clusters, A and B. They mainly comprised severe and mild cases, respectively, since

cluster A had significantly higher ordinal scores, peak serum C-reactive protein level, and peak

chest X-ray score than cluster B (Unpaired t-test, $P < 0.001$) (Fig. 1c). Cluster A was subdivided

into three clusters, A1, A2, and A3 with their own comorbidity status (Fig. 1d). Cluster A1 was

characterized by the presence of hypertension, while cluster A3 was a DM-enriched group.

Interestingly, cluster A2 tended to have a higher rate of CHIP than the others in cluster

A (Fisher's exact test, $P = 0.074$), while it had significantly lower BMI (median [IQR], 20.8

[17.8—23.1] vs. 23.0 [21.7—26.1]; Student's t-test, $P = 0.005$) and lower rate of DM (0/18, 0%

vs. 22/51, 43.1%; Fisher's exact test, $P = 0.001$) or hypertension (0/18, 0% vs. 40/51, 78.4%;

Fisher's exact test, $P < 0.001$) (Fig. 1d and Supplementary Table 2). However, age distribution

was not statistically different (Fisher's exact test, $P = 0.662$).

The presence of a particular type of severe cases uniquely enriched by CHIP without

canonical risk factors such as DM, hypertension, and high BMI led us to hypothesize that CHIP

may contribute to severe COVID-19 in its own way. An additional hierarchical clustering with

DM, hypertension, and CHIP information in all severe patients (Ordinal scale >=3) also

supported our hypothesis since it showed that CHIP was clustered together (Extended Data Fig.

1).

To test the statistical significance of the impact of CHIP on COVID-19 severity in

previously healthy patients, we stratified the entire patients by the presence of any of the

canonical risk factors such as DM, hypertension, and BMI ≥ 30.0. When adjusted with age

and gender, the risk of severe COVID-19 was significantly higher in CHIP (+) patients than in

CHIP (-) ones in canonical risk factor-absent subgroup (adjusted odds ratio [95% confidence

interval], 14.8 [1.3—164.1]; logistic regression $P = 0.028$; Table 1). Multivariate analysis

including the interaction between canonical risk factors and CHIP revealed that CHIP was an

independent risk factor for severe COVID-19 (adjusted odds ratio [95% confidence interval],

10.7 [1.1—100.7], logistic regression $P = 0.038$, Table 2). The results of the patients clustering

and the statistical evidence indicate that CHIP is an independent risk factor for severe COVID-

19, especially in previously healthy patients.

Distinct immune signatures in CHIP (+) severe COVID-19

We next sought to identify distinct immune signatures for severe COVID-19 according to the

presence of CHIP. To this end, a total of 60,310 high-quality single-cell transcriptome

profiles of peripheral blood mononuclear cells (PBMCs) generated by 10x Genomics single-

cell RNA-seq (scRNA-seq) platform were integrated from healthy donors (n=4), severe

influenza (n=5), CHIP (-) mild COVID-19 (n=5), CHIP (-) severe COVID-19 (n=3), and

CHIP (+) severe COVID-19 (n=6) specimens (Supplementary Table 3 and 4, see Methods),

with an average of 6,100 unique molecular identifies (UMIs), representing 1,400 genes. The

reproducibility and quality were ensured (Extended Data Fig. 2a-c)[26]. Based on t-distributed

stochastic neighbor embedding (t-SNE) of transcriptome profiles, 24 subgroups were derived.

Assigning cell types with previously annotated marker genes (Extended Data Fig. 2d-e, see

Methods)[26], we focused on 9 major immune cell types, including IgG$^+$ B cell, IgG$^-$ B cell,

CD4$^+$ T cell, naïve CD4$^+$ T cell, CD8$^+$ T cell, CD8$^+$ memory T cell, natural killer (NK) T cell,

classical monocyte, and non-classical monocyte in subsequent analyses (Fig. 2a). Non-

immune cells such as platelets, red blood cells (RBCs), and uncategorized small cell

populations were excluded. Notably, the proportion of monocytes was markedly increased,

particularly in CHIP (+) severe COVID-19 patients (K–S test, P=3.49e-3) (Extended Data

Fig. 2f-g), consistent with the known impact of CHIP with increased myeloid cell

population[27].

In terms of transcriptome profiles at the cell-type resolution, as expected, most immune

cell types originating from COVID-19 were clustered together when compared to influenza

(Fig. 2b). Interestingly, those from severe COVID-19 were subdivided according to the

presence of CHIP (Fig. 2b). To investigate relevant biological functions that establish such

unique host immune responses in CHIP (+) severe COVID-19, we identified up-regulated

genes specific to CHIP (+) severe COVID-19 compared to healthy normal control, influenza,

CHIP (-) mild COVID-19, and CHIP (-) severe COVID-19, respectively, using MAST

algorithm (Extended Data Fig. 3a)[28]. Regardless of both normal and disease control groups,

tumor necrosis factor (TNF-α)/NF-kB and interferon-gamma (IFN-γ) responses were

commonly enriched in up-regulated genes in CHIP (+) severe COVID-19 (Extended Data Fig.

3b). A noticeable enrichment of CHIP (+) up-regulated genes was observed in classical

monocytes (Extended Data Fig. 3c).

To further investigate the unique immune signatures at the individual immune cell-

type resolution, we conducted gene set enrichment analysis (GSEA) for differentially

expressed genes between CHIP (+) and CHIP (-) severe COVID-19. Based on cytokine-

responsive gene sets originated from each cytokine treated cells (LINC L1000 ligand

perturbation analysis in Enrichr) (see Methods)[29], strong IL-1β and TNF-α responses were

observed in both CHIP (+) and CHIP (-) severe COVID-19 compared to healthy normal control

(Fig. 2c). However, direct comparison between CHIP (+) and CHIP (-) severe COVID-19

revealed that immune responses in classical monocytes were strongly skewed towards CHIP

(+) patients (Fig. 2d), while other cell types such as T cells, B cells or non-classical monocytes

did not show such trend (Fig. 2e-g, Extended Data Fig. 4a-e). Taken together, CHIP (+) patients

under COVID-19 infection present unique host immune responses, particularly in classical

monocytes.

Classical monocyte-mediated hyperinflammation in CHIP (+) severe COVID-19

To examine how the presence of CHIP attributes to the immunologic responses in classical

monocytes, we focused on analyzing 445 up- and 417 down-regulated CHIP (+) specific genes.

CHIP (+) up-regulated genes demonstrated enrichment with inflammation cytokine responses,

such as IL-1β compared to CHIP (-) (Mann-Whitney's U test, P=3.24e-2, IL-1β) (Fig. 3a).

Other COVID-19 representative interleukins such as IL-6, IL-10 and IL-15 were also elevated

in CHIP (+) severe COVID-19. (Fig. 3b)[30]. Notably, CHIP (+) up-regulated genes were

strongly related to both type I and II interferon (IFN) responses (Mann-Whitney's U test,

P=1.08e-3, IFN-γ; P=3.97e-3, IFN-α). We previously proposed that a group of genes involved

in type I IFN-induced TNF-α mediated hyperinflammation by abolishing the tolerance effects

of TNF-α (Class 1 gene in Park *et al.*[31]) in monocyte has a critical role in promoting

hyperinflammation of COVID-19[26]. Consistently, the CHIP (+) up-regulated genes presented

a modest enrichment with the Class I genes (Fig. 3c), partly explaining the hyperinflammation

signature of CHIP (+). However, other inflammatory TLR-induced genes regardless of TNF-α

tolerization (Class II and III genes in Park *et al.*[31]) were extremely biased to CHIP (+) up-

regulated genes, postulating the presence of additional classical monocyte driven responses

exacerbating the inflammation signatures in CHIP (+).

IFN-γ mediated hyperinflammation in CHIP (+) severe COVID-19 revealed by a

pseudotime analysis

Intriguingly, IFN-γ response was notably high in CHIP (+) up-regulated genes (Fig. 3a). As a

high level of IFN-γ has been reported as an indicator of severe COVID-19[8,32,33], we

hypothesized that IFN-γ response could attribute to hyperinflammation signatures of classical

monocytes in CHIP (+) patients. We first determined the association between IFN-γ response

and COVID-19 severity in CHIP (+) by conducting pseudotime analysis using the specimens

collected twice from one patient to exclude innate individual biases (Fig. 4a) (see Methods).

After ordering cells along with the trajectory analysis, we allocated the annotation of high and

low inflammation clusters based on inflammatory signatures termed in MsigDB Hallmark 2020

(Fig. 4b). We found that IFN-γ response genes were significantly enriched in a high

inflammation group of CHIP (+), even more potent than that of CHIP (-) (Fig. 4c).

Recent mouse and scRNA-seq comparison studies have highlighted the immuno-

pathogenic contribution of IFN-γ in severe COVID-19 as demonstrated by IFN-γ induced

inflammatory macrophage phenotype and synergistic effect of IFN-γ and TNF-α[34,35].

Consistently, in our analysis, CHIP (+) specific genes were significantly enriched by pro-

inflammatory M1-like macrophage-specific genes obtained from two independent studies[36]

[37](Fig. 4d and Extended Data Fig. 5a) and associated with up-regulated genes by co-treatment

of TNF-α and IFN-γ (Fig. 4e)[35]. Such enrichment suggests that CHIP (+) severe COVID-19

patients are representative cases that could be explained by previously discovered IFN-γ

mediated disease exacerbating mechanism. For a treatment perspective, we noticed that spleen

tyrosine kinase (Syk) inhibitor, which is known to reduce the expressions of interferon-

stimulated genes[38], may be an effective molecule for the intervention of CHIP (+) up-regulated

genes (Paired t-test, P=0.042) (Extended Data Fig. 5b). Taken together, IFN-γ response is

thought to be a main source of the hyperinflammation immune signature in CHIP (+) severe

COVID-19.

DNMT3A mutation-specific hypo-DMRs are linked to IFN-γ response genes

Next, we sought to identify mechanisms by which IFN-γ response genes are up-regulated

explicitly in CHIP (+) COVID-19 patients. Considering mutations of multiple epigenetic

regulators such as *DNMT3A*, *TET2*, and *ASXL1* in CHIP, we hypothesized that altered

chromatin activity of *cis*-regulatory elements might exert CHIP-specific gene expression. To

test our hypothesis, we identified 2,348 differentially methylated regions (DMRs) in acute

myeloid leukemia (AML) patients carrying *DNMT3A* mutations (see Methods)

(Supplementary Table 5)[39]. In support of our hypothesis, these DMRs were highly overlapped

with putative regulatory elements (Fisher's exact test, *P*<0.001, Proximal to the promoter;

P<0.001, Distal regulatory element) (Fig. 5a) (see Methods). As exemplified in *FOXO3* and

NFIL3, known IFN response genes, CHIP (+) up-regulated genes were also more closely

located to the hypo-DMRs (Fig. 5b, Extended Data fig. 6).

As many *cis*-regulatory elements are known to target genes over large genomic

distances[40], we performed *in situ* Hi-C experiments on CD14^{++}/CD16^{-} classical monocytes of

two healthy donors to precisely annotate target genes of CHIP-dependent DMRs (see Methods).

With ~500M long-range chromatin interactions over 15kb genomic distance, we defined

significant long-range chromatin contacts using covNorm in 10kb resolution (see Methods)[41].

Using this information, as illustrated in *RBPJ* and *CXCL2* genes (Fig. 5c-d), we revealed that,

in total, around 33% of CHIP-specific up-regulated genes, denoted as 'linked genes', were

associated with hypo-DMRs either in proximal (within 15kb) or long-range chromatin

interactions (over 15kb but less than 2Mb) (Fig. 5e). Notably, those linked genes were more

enriched by both type I and II IFN response genes compared to the remaining up-regulated

genes (Mann-Whitney's U test, *P*=2.50e-3, IFN-γ; *P* =3.97e-3, IFN-α) (see Methods) (Fig. 5f).

Regarding high inflammation cluster in pseudotime analysis, we revealed that IFN-γ response

genes largely overlap by hypo-DMRs linked genes (Mann-Whitney's U test, $P =1.08e-3$, IFN-

γ) (Fig. 5g). Our results strongly support that CHIP-dependent altered chromatin activities are

associated with putative *cis*-regulatory elements of IFN-γ response genes, which may establish

unique gene expression profiles in CHIP patients during immune responses.

Activation of poised and bivalent *cis*-regulatory elements primes IFN-γ response genes

To further characterize CHIP-specific hypo-DMRs, we examined the enrichment of four

representative histone modification marks including histone H3 4th lysine mono-methylation

(H3K4me1), tri-methylation (H3K4me3), 27th lysine acetylation (H3K27ac), and tri-

methylation (H3K27me3) of primary human classical monocyte[42,43]. When comparing DMRs

to randomly selected genomic regions, hypo-DMRs were mostly marked by H3K27me3 in the

primary human classical monocytes, while hyper-DMRs were enriched by H3K27ac as an

indicator of active regulatory elements (Fig. 6a-b). However, interestingly, a subset of hypo-

DMRs, linking to CHIP (+) up-regulated genes, were also significantly co-occupied by

H3K4me1 and H3K4me3 peaks compared to the unlinked hypo-DMRs (Fisher's exact test,

$P<0.001$, H3K4me1; $P<0.001$, H3K4me3; $P =2.37e-2$, H3K27ac) (Fig. 6c-d). Such co-

exhibition of inactive and active chromatin signatures indicates that the regulatory elements of

CHIP (+) up-regulated genes have shifted from poised or bivalent status to an active chromatin

state through the process of CHIP dependent hypo-DNA methylation. To test this possibility,

we annotated chromatin states of linked hypo-DMRs according to the combination of the

histone modification marks (see Methods). We found that promoter-distal hypo-DMRs were

significantly enriched by poised enhancer chromatin signatures (H3K4me1 without

H3K27me3) (Fisher's exact test, $P <0.001$) (Fig. 6e). Similarly, promoter-proximal hypo-

DMRs were enriched by bivalent promoters (H3K4me3 and H3K27me3) or active promoters

(H3K4me3) (Fisher's exact test, $P = 3.80e\text{-}3$, Active promoter; $P= 7.36e\text{-}2$, Bivalent) (Fig. 6f).

Thus, CHIP mutants appear to reprogram the epigenetic states including the loss of silent

marker at poised enhancers or bivalent promoters, which prime the IFN associated immune

response genes, thereby driving hyperinflammation and leading to the critical course of

COVID-19 (Fig. 7).

Discussion

By analyzing thorough clinical, radiological, and laboratory characteristics, as well as the presence of CHIP, we showed that CHIP is a novel risk factor for severe COVID-19 in previously healthy population without canonical clinical risk factors. scRNA-seq analysis revealed a distinct IFN-γ mediated immuno-pathogenic mechanism in CHIP (+) severe COVID-19 plausibly attributed by CHIP dependent chromatin reorganization. These results consistently indicate that CHIP may play a critical role in the progression of severe COVID-19, especially in previously healthy patients with its own immunologic pathway.

Exploration of an additional risk factor for severe COVID-19 is clinically valuable in this pandemic to predict the progression of COVID-19 more accurately and to improve our management strategy. Owing to its pro-inflammatory nature, CHIP may contribute to the progression of severe COVID-19. By analyzing more than 500 patients, Bolton *et al*. reported that CHIP is significantly associated with severe COVID-19[23], especially for patients carrying non-putative driver mutations. There have been a few controversial reports, however, they needed to be critically appraised because of methodological considerations. Duployez *et al.* showed a higher prevalence of CHIP in severe COVID-19 patients than age-matched hematologic malignancy-free cohort, but they could not show that CHIP affects clinical outcome[24]. However, they could only analyze severe patients whose clinical outcomes might hardly be differentiated by solely the presence of CHIP. In another study conducted by Hameister *et al.* involving 102 hospitalized patients with COVID-19, the presence of CHIP was not associated with severe COVID-19[25]. But the study lacked a stratified analysis or an adjustment for possible interactions. By introducing a hierarchical clustering, followed by thorough statistical examinations including interaction analysis, we could clearly show that CHIP is an independent risk factor for severe COVID-19.

COVID-19 patients were reported to show heterogeneous symptoms ranging from

asymptomatic to critical illness[3,44]. In line with it, many studies divided COVID-19 patients

into subgroups defined by immunological characteristics, for instance, patterns of sepsis[45],

subpopulations of lymphocytes[44], IFN responses in lung[46], or carrying loss-of-function

variants[47]. Single-cell techniques have been vigorously applied in COVID-19 to dissect

underlying causes of the diverse immune responses[33,48,49] and to elaborately determine the

relationship between immune subtypes and clinical characteristics[50,51]. Despite those important

works, none of the single-cell study has characterized the immunological effects of CHIP in

COVID-19 yet. In this regard, the current study uniquely demonstrated how CHIP-associated

somatic mutations in immune cells could actively establish a novel subgroup in COVID-19

patients. With single-cell immune transcriptome analysis, we could reveal that IFN-γ related

hyperinflammation is a hallmark of CHIP (+) severe COVID-19. Especially, there was an

enrichment of inflammatory signature in classical monocytes, which is compatible with recent

knowledge regarding the effect of CHIP on myeloid askew hematopoietic stem cell

differentiation[27]. From a cytokine perspective, IFN-γ and its synergism with TNF-α were

thought to play a critical role in the pathogenesis of severe COVID-19 in CHIP (+) patients.

This finding aligns with the previous report stating the role of IFN-γ and/or TNF-α in

exacerbating chronic inflammatory disease by CHIP[52]. Our study also implies that not only

type I IFN response but also type II IFN response plays an important role in disease

exacerbation in certain patients with severe COVID-19. Additional interesting immunological

finding reasonably explained with the CHIP biology is the up-regulation of genes related to

inflammatory macrophage in CHIP (+) severe COVID-19. CHIP is well known to drive

hyperinflammation in chronic disease mainly attributable to the altered function of monocyte

and macrophage [27].

Recent studies have revealed the pathological effects of CHIP such as atherosclerosis

and malignancy[18,53,54] and characterized the effect of CHIP mutations on hematopoiesis and

immunological functions regarding epigenetic mechanisms[55,56]. However, there is still an

ambiguity in how physiological pathogenic characteristics are linked to the altered chromatin

activities by CHIP mutations in actual patient cohorts. In our study, such linkage is explained

as regulatory interactions between pathogenic genes and their *cis*-regulatory elements with the

altered chromatin states, which are expected to be mediated by mutations in chromatin

regulators (Fig. 7). Although further investigation of other types of CHIP mutations such as

TET2 and *ASXL1* are required to comprehensively characterize CHIP mutation-dependent

epigenetic reprogramming, the pathogenesis of other CHIP-dependent diseases could be

elucidated by considering the similar mechanisms.

There are several limitations in the present study. First, sample sizes of either entire

cohort or scRNA-seq-analyzed patients were limited. Second, the proportion of severe COVID-

19 was high in the present study since it was conducted mainly in tertiary-care hospitals.

Although we could show an apparent incline to severe COVID-19 in CHIP (+) patients and its

distinctive immune signature in this study, further validations in a larger prospective cohort

representing general COVID-19 patients are warranted. Third, scRNA-seq was performed with

PBMCs instead of lung fluid or infected lung tissues limiting our analysis in exploring systemic

inflammation as a result of COVID-19 pulmonary disease. Lastly, further studies are needed to

reveal the exact epigenetic differences between mutant and wild type monocytes in CHIP

patients and the real time point when CHIP mutant monocytes stir hyperinflammation.

Lastly, in terms of therapeutic strategy, Syk inhibitors have been implicated in

suppressing these pathogenic immune responses (Extended Data Fig. 5b) induced by CHIP. An

in vitro study on Syk inhibitor fostamatinib suggested its therapeutic effect against COVID-

19[57], and we have pending results of a randomized placebo-controlled trial with the drug

(NCT04579393). As the therapeutic efficacy of Syk inhibitors could be more potent in CHIP

(+) patients, it is worth evaluating treatment outcomes involving Syk inhibitors according to

CHIP status.

In summary, despite such limitations, we successfully clarified that there is a distinct

CHIP-driven severe COVID-19 subgroup and elucidated its unique immunological mechanism.

Revealing the underlying epigenetic mechanism for the altered immune function that aligns

with well-known CHIP biology suggests the robustness of our findings. It appears that classical

monocytes in patients with CHIP (+) COVID-19 undergo distinct immune responses, thus

focusing on immunomodulation strategies according to the presence of CHIP is required.

Considering the shared pathogenic host immune response across infections, we postulate that

our findings might bring a better understanding of previously unexplained exacerbation of

clinical conditions by various viruses.

Acknowledgments

We thank all our laboratory members for their support and critical suggestions throughout this

work. Funding: This work was funded by the SUHF Fellowship (to I.J.). We also thank Dr.

Shin for sharing specimens to determine the presence of CHIP.

Author Contribution

CKK, BC, YK, and IJ conceived the study. SP generated scRNA-seq data. AJL and YK

generated Hi-C data. CKK, BC, SK, SHY, DL, and SK performed data analysis. CKK, EC,

JJ, PGC, WBP, ESK, HBK, NJK, MO, SK, HI, JK, YHL, JL, JYL, JHM, and K-HS

contributed to the collection of clinical information and samples. CKK, BC, YK, and IJ

contributed to data interpretation with assistant from SK and KK. CKK and BC prepared the

manuscript with assistance from YK and IJ. All authors read and commented on the

manuscript.

References

1. World Health Organization. Novel Coronavirus (2019-nCoV) situation reports. (2020).

2. Zhu, N., *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733 (2020).

3. Wu, Z. & McGoogan, J.M. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA* **323**, 1239-1242 (2020).

4. Cunningham, J.W., *et al.* Clinical Outcomes in Young US Adults Hospitalized With COVID-19. *JAMA Intern Med* (2020).

5. Williamson, E.J., *et al.* Factors associated with COVID-19-related death using OpenSAFELY. *Nature* **584**, 430-436 (2020).

6. Zhang, S.Y., Zhang, Q., Casanova, J.L., Su, H.C. & Team, C. Severe COVID-19 in the young and healthy: monogenic inborn errors of immunity? *Nat Rev Immunol* **20**, 455-456 (2020).

7. Zhou, C., *et al.* Predictive factors of severe coronavirus disease 2019 in previously healthy young adults: a single-center, retrospective study. *Respir Res* **21**, 157 (2020).

8. Huang, C., *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506 (2020).

9. Yang, X., *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* (2020).

10. Liu, Y., *et al.* Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* (2020).

11. Zou, L., *et al.* SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *N Engl J Med* (2020).

12. Mehta, P., *et al.* COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* **395**, 1033-1034 (2020).

13. Kang, C.K., *et al.* Aberrant hyperactivation of cytotoxic T-cell as a potential determinant of COVID-19 severity. *Int J Infect Dis* **97**, 313-321 (2020).

14. Xu, Z., *et al.* Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med* **8**, 420-422 (2020).

15. Merad, M. & Martin, J.C. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat Rev Immunol* **20**, 355-362 (2020).

16. Steensma, D.P., *et al.* Clonal hematopoiesis of indeterminate potential and its distinction from myelodysplastic syndromes. *Blood* **126**, 9-16 (2015).

17. Rodrigues, C.P., Shvedunova M., Akhtar A. Epigenetic Regulators as the Gatekeepers of Hematopoiesis. *Trends in Genetics* **37**(2021).

18. Jaiswal, S., *et al.* Clonal Hematopoiesis and Risk of Atherosclerotic Cardiovascular Disease. *N Engl J Med* **377**, 111-121 (2017).

19. Sano, S., *et al.* CRISPR-Mediated Gene Editing to Assess the Roles of Tet2 and Dnmt3a in

Clonal Hematopoiesis and Cardiovascular Disease. *Circ Res* **123**, 335-341 (2018).

20.   Yura, Y., Sano, S. & Walsh, K. Clonal Hematopoiesis: A New Step Linking Inflammation to

Heart Failure. *JACC Basic Transl Sci* **5**, 196-207 (2020).

21.   Jaiswal, S. & Libby, P. Clonal haematopoiesis: connecting ageing and inflammation in

cardiovascular disease. *Nat Rev Cardiol* **17**, 137-144 (2020).

22.   Zekavat, S.M., *et al.* Hematopoietic mosaic chromosomal alterations increase the risk for

diverse types of infection. *Nat Med* **27**, 1012-1024 (2021).

23.   Bolton, K.L., *et al.* Clonal hematopoiesis is associated with risk of severe Covid-19. *medRxiv*

(2020).

24.   Duployez, N., *et al.* Clinico-Biological Features and Clonal Hematopoiesis in Patients with

Severe COVID-19. *Cancers (Basel)* **12**(2020).

25.   Hameister, E., *et al.* Clonal Hematopoiesis in Hospitalized Elderly Patients With COVID-19.

Hemasphere **4**, e453 (2020).

26.   Lee, J.S., *et al.* Immunophenotyping of COVID-19 and influenza highlights the role of type I

interferons in development of severe COVID-19. *Sci Immunol* **5**(2020).

27.   Jaiswal, S. & Ebert, B.L. Clonal hematopoiesis in human aging and disease. *Science* **366**(2019).

28.   Finak, G., *et al.* MAST: a flexible statistical framework for assessing transcriptional changes

and characterizing heterogeneity in single-cell RNA sequencing data. *Genome Biol* **16**, 278

(2015).

29.   Duan, Q., *et al.* LINCS Canvas Browser: interactive web app to query, browse and interrogate

LINCS L1000 gene expression signatures. *Nucleic Acids Res* **42**, W449-460 (2014).

30.   Angioni, R., *et al.* Age-severity matched cytokine profiling reveals specific signatures in

Covid-19 patients. *Cell Death Dis* **11**, 957 (2020).

31.   Park, S.H., *et al.* Type I interferons and the cytokine TNF cooperatively reprogram the

macrophage epigenome to promote inflammatory activation. *Nat Immunol* **18**, 1104-1116

(2017).

32.   Lucas, C., *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19.

Nature **584**, 463-469 (2020).

33.   Chua, R.L., *et al.* COVID-19 severity correlates with airway epithelium-immune cell

interactions identified by single-cell analysis. *Nat Biotechnol* **38**, 970-979 (2020).

34.   Karki, R., *et al.* Synergism of TNF-alpha and IFN-gamma Triggers Inflammatory Cell Death,

Tissue Damage, and Mortality in SARS-CoV-2 Infection and Cytokine Shock Syndromes. *Cell*

184, 149-168 e117 (2021).

35.   Zhang, F., *et al.* IFN-gamma and TNF-alpha drive a CXCL10+ CCL2+ macrophage phenotype

expanded in severe COVID-19 lungs and inflammatory diseases with tissue inflammation.

Genome Med **13**, 64 (2021).

36.   Kang, K., *et al.* IFN-γ selectively suppresses a subset of TLR4-activated genes and enhancers

to potentiate macrophage activation. *Nat Commun* **10**, 3320 (2019).

37.   Carvalho K., R.E., Jansen C., Williams K., Dowey A., McGill C., Mortazavi A. Uncovering the

439   Gene Regulatory Networks Underlying Macrophage Polarization Through Comparative

440   Analysis of Bulk and Single-Cell Data. *bioRxiv* (2021).

441 38.   Liu, S.*, et al.* Critical role of Syk-dependent STAT1 activation in innate antiviral immunity. *Cell*

442   *Rep* **34**, 108627 (2021).

443 39.   Li, S.*, et al.* Distinct evolution and dynamics of epigenetic and genetic heterogeneity in acute

444   myeloid leukemia. *Nat Med* **22**, 792-799 (2016).

445 40.   Jung, I.*, et al.* A compendium of promoter-centered long-range chromatin interactions in

446   the human genome. *Nat Genet* **51**, 1442-1449 (2019).

447 41.   Kim, K. & Jung, I. covNorm: An R package for coverage based normalization of Hi-C and

448   capture Hi-C data. *Comput Struct Biotechnol J* **19**, 3149-3159 (2021).

449 42.   Zhang, J.*, et al.* An integrative ENCODE resource for cancer genomics. *Nat Commun* **11**,

450   3696 (2020).

451 43.   Consortium, E.P. An integrated encyclopedia of DNA elements in the human genome. *Nature*

452   **489**, 57-74 (2012).

453 44.   Mathew, D.*, et al.* Deep immune profiling of COVID-19 patients reveals distinct immunotypes

454   with therapeutic implications. *Science* **369**(2020).

455 45.   Giamarellos-Bourboulis, E.J.*, et al.* Complex Immune Dysregulation in COVID-19 Patients with

456   Severe Respiratory Failure. *Cell Host Microbe* **27**, 992-1000 e1003 (2020).

457 46.   Nienhold, R.*, et al.* Two distinct immunopathological profiles in autopsy lungs of COVID-19.

458   *Nat Commun* **11**, 5086 (2020).

459 47.   Zhang, Q.*, et al.* Inborn errors of type I IFN immunity in patients with life-threatening COVID-

460   19. *Science* **370**(2020).

461 48.   Liao, M.*, et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-

462   19. *Nat Med* **26**, 842-844 (2020).

463 49.   Wilk, A.J.*, et al.* A single-cell atlas of the peripheral immune response in patients with severe

464   COVID-19. *Nat Med* **26**, 1070-1076 (2020).

465 50.   Ren, X.*, et al.* COVID-19 immune features revealed by a large-scale single-cell transcriptome

466   atlas. *Cell* **184**, 1895-1913 e1819 (2021).

467 51.   Stephenson, E.*, et al.* Single-cell multi-omics analysis of the immune response in COVID-19.

468   *Nat Med* **27**, 904-916 (2021).

469 52.   Zhang, C.R.C.*, et al.* Inflammatory cytokines promote clonal hematopoiesis with specific

470   mutations in ulcerative colitis patients. *Exp Hematol* **80**, 36-41 e33 (2019).

471 53.   Jaiswal, S.*, et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl*

472   *J Med* **371**, 2488-2498 (2014).

473 54.   Libby, P.*, et al.* Clonal Hematopoiesis: Crossroads of Aging, Cardiovascular Disease, and

474   Cancer: JACC Review Topic of the Week. *J Am Coll Cardiol* **74**, 567-577 (2019).

475 55.   Izzo, F.*, et al.* DNA methylation disruption reshapes the hematopoietic differentiation

476   landscape. *Nat Genet* **52**, 378-387 (2020).

56.   Zhang, Q.*, et al.* Tet2 is required to resolve inflammation by recruiting Hdac2 to specifically

repress IL-6. *Nature* **525**, 389-393 (2015).

57.   Strich, J.R.*, et al.* Fostamatinib Inhibits Neutrophils Extracellular Traps Induced by COVID-19

Patient Plasma: A Potential Therapeutic. *J Infect Dis* **223**, 981-984 (2021).

Figure Legends

Figure 1. Hierarchical clustering analysis for clinical characteristics of severe COVID-

19

a, Overview of study design. **b**, A heatmap shows the hierarchical clustering result of clinical

characteristics. On the top, three different bars represent clusters and the presence of CHIP

mutations for each patient. The first bar represents two main clusters of patients (n=69 for

cluster A and n=74 for cluster B). The second bar represents five sub-clusters of patients

(n=32 for A1, n=18 for A2, n=19 for A3, n=47 for B1, and n=26 for B2). The third bar

represents the presence of CHIP. **c**, Boxplots showing clinical characteristics between cluster

A (orange) and B (green). For the boxplots, the box represents the interquartile range (IQR)

and the whiskers correspond to the highest and lowest points within $1.5 \times$ IQR. Statistical

significance was examined with unpaired t-test (* < p-value 0.05, ** < p-value 0.01, and ***

< p-value 0.001). **d**, Barplots showing fractions of patients with comorbidities or CHIP for

cluster A1, A2, and A3. HTN stands for hypertension. DM stands for diabetes mellitus.

Figure 2. Single-cell transcriptome analyses of COVID-19 according to CHIP status

a, Scatter plots of integrated scRNA-seq data represented with a t-SNE method. Left, immunes

cells are presented according to the presence of CHIP. Others group indicates CHIP (-) COVID-

19 mild, influenza, and healthy normal control groups. Right, nine immune cell types were

plotted by a t-SNE method. **b**, A PCA analysis with transcriptome profiles according to the

immune cell types and disease groups. Colors and shapes represent the respective disease

groups or cell types. **c,** A scatter plot showing combined scores of LINCS L1000 Ligand

Perturbations up gene ontology library between CHIP (+) severe COVID-19 and CHIP (-)

severe COVID-19 compared to the healthy normal control. Identity lines are presented on

diagonal. Square of correlation of combined scores is shown together. **d-g**, Scatter plots

showing combined scores same gene ontology library in Fig. 2c for differentially expressed

genes between CHIP (+) and CHIP (-) severe COVID-19 in classical monocyte (d), CD4+ T

cell (e), non-classical monocyte (f), and IgG + B cell (g). The horizontal axis, up-regulated

genes in CHIP (+); Vertical axis, up-regulated genes in CHIP (-). Identity lines are presented

on diagonal. The color indicates types of perturbed ligand.

Figure 3. CHIP specific immune signatures in classical monocytes

a, Barplots showing combined scores of DEGs in classical monocytes between CHIP (+) and

CHIP (-) severe COVID-19 for same gene ontology library in Fig. 2c-g. The color indicates

up-regulated genes in CHIP (+) (green) and CHIP (-) (blue). Each point indicates one ligand

perturbation term in the library, in total 6 different types of ligands: IFN-α (n=5), IL-1β (n=6),

IFN-γ (n=6), IL-17 (n=2), IL-6 (n=2), TNF-α (n=6). The mean and standard error of the mean

(s.e.m) of each gene set are shown together. One-sided Mann-Whitney's U test was performed

(*: $P<0.05$, **: $P<0.01$). **b**, Barplots showing combined scores of same gene sets in Fig. 3a for

Ligand Perturbations from GEO up gene ontology library. The color indicates up-regulated

genes of CHIP (+) (pink) and CHIP (-) (yellow). **c**, Gene set enrichment analysis (GSEA) on

the same DEGs used in Fig. 3a-b for three classes of TLR induced genes[31]. Each plot shows

the distribution of enrichment of distinct class genes along with the list of the DEGs pre-ranked

with log-fold changes based on the CHIP (-) severe COVID-19. The color under the plot

indicates patient groups of up-regulated genes. Red, up-regulated in CHIP (+) severe COVID-

19. Blue, up-regulated in CHIP (-) severe COVID-19. Normalized enrichment scores (NES)

and FDR are shown together. Top, Class I genes. Middle, Class II genes. Bottom, Class III

genes.

Figure 4. Pseudotime analyses of severe COVID-19 with CHIP in classical monocytes

a-c, Pseudotime analyses for CHIP (+) or CHIP (-) severe COVID-19 patients. **a**, Cells are

aligned according to the pseudotime axis calculated by Monocle2. The color indicates a type

of cluster. Top, CHIP (+) patient. Bottom, CHIP (-) patient. **b**, Heatmaps representing the

expression level of marker genes for the ordered cells. The list of genes is the representative

gene set of each cluster. Barplots showing combined scores of each cluster for inflammatory

response term in MsigDB Hallmark 2020. **c**, Scatter plots of combined scores of marker genes

of inflammation clusters for LINCS L1000 Ligand Perturbations up gene ontology library. The

horizontal axis, high inflammation cluster; the vertical axis, low inflammation cluster. Identity

lines are presented on diagonal. Colors indicate types of perturbed ligand for IFN-α (n=5), IL-

1β (n=6), IFN- γ (n=6), IL-17 (n=2), IL-6 (n=2), TNF-α (n=6). Shapes represent the presence

or absence of CHIP. **d, e**, GSEA plots for marker genes of inflammation clusters in CHIP (+)

and CHIP (-), respectively. Genes are ordered based on log-fold changes between high

inflammation cluster and low inflammation cluster. Normalized enrichment scores (NES) and

FDR are presented for DEGs between M1-like (LPS-IFN-γ stimulated) and M0 macrophage

(untreated) [36](d) and DEGs between TNF-α-IFN-γ co-treatment and untreated condition[35] (e),

respectively. LPS indicates Lipopolysaccharide.

Figure 5. Regulatory potential of CHIP-specific hypo-DMRs in IFN response genes

a, Stacked bar plots showing the proportion of annotated hypo- and hyper-DMRs. In DMR,

Proximal to the promoter, n=1124; Distal regulatory element, n=423; others, n=801. In

random for DMR, Proximal to promoter, n=314; Distal regulatory element, n=81; others,

n=1953. **b**, Examples of CHIP (+) up-regulated genes are presented with hypo-DMRs and

multiple histone modification signatures for *FOXO3* (right) and *NFIL3* (left). **c,** Examples of

analysis of Hi-C data. Whole interaction map of chromosome 4 in 40kb resolution. The color

indicates normalized chromatin contact frequencies. Two example regions containing CHIP

(+) severe COVID-19 up-regulated genes in classical monocytes (region 1 and region 2) are

highlighted by yellow dashed lines. **d,** Hi-C contact maps (heatmap) for region 1 (*RBPJ*) and

region 2 (*CXCL2*) are shown together with significant long-range chromatin interactions with

the promoter regions (arcs on the below of the heatmap) and distribution of hypo-DMRs and

histone modifications signals. **e**, A pie chart showing proportions of hypo-DMR linked up-

regulated genes in CHIP (+) severe COVID-19 classical monocytes. The color indicates the

types of linkage between hypo-DMRs and up-regulated genes. **f, g**, Barplots of relative

combined scores to TNF-α of genes linked to hypo-DMRs and others for same gene ontology

library in Fig. 2c-g. Colors indicate types of perturbed ligands. Each point indicates one

ligand perturbation term. The mean and standard error of the mean (s.e.m) of each gene set

are presented in barplot. One-sided Mann-Whitney's U test was performed (n.s: non-

significant, **: $P<0.01$). Up-regulated genes in CHIP (+) compared to CHIP (-) (f) and

marker gens of high inflammation cluster compared to low inflammation cluster of CHIP (+)

patient in Fig. 4 (g).

Figure 6. Chromatic states of hypo-DMRs linking to CHIP specifically up-regulated

genes

a, b, CHIP-seq signal distribution (a for H3K27ac and b for H3K27me3) of 20k upstream and

downstream surrounding of hypo-DMRs (n=1,693), hyper-DMRs (n=655) and randomly

selected regions. Each randomly selected regions have the same distribution of chromosome

number and length of hypo- and hyper-DMR, respectively. Top, average ChIP-seq signal

distributions. Bottom, heatmaps of CHIP-seq signals of the corresponding regions. **c, d,** CHIP-

seq signal distribution for CHIP (+) up-regulated genes linked (n=209) and unliked (n=1484)

hypo-DMRs for H3K4me1 (c) and H3K4me3 (d). Top, average profiles of CHIP-seq signals

for linked and unlinked hypo-DMRs. Bottom, heatmaps of CHIP-seq signals of the

corresponding regions. **e**, **f**, Stacked barplots of the linked and unlinked hypo-DMRs with

annotation of chromatin states. For statistical significance, one-sided Fisher's exact test was

performed between linked and unlinked for promoter-distal (e) or -proximal (f) hypo-DMRs.

Figure 7. A proposed model for the pathogenesis of severe COVID-19 in patients with
CHIP

A schematic overview shows the progression of disease exacerbation in terms of gene

regulation, cellular level, and clinical/immunological signatures under CHIP (+) severe

COVID-19.

Extended Data Figure 1. A heatmap showing hierarchical clustering of severe COVID-19

patients with comorbidities and CHIP.

The first bar represents two clusters, cyan for non-CHIP enriched (n=65), and orange for CHIP

enriched one (n=25). The second bar represents five distinct types of sub-clusters, namely, red,

for without comorbidity (n=19), orange for DM (n=9), green for HTN (n=26), purple for CHIP

(n=13), and blue for complex (n=23). The color in the heatmap represents the presence of CHIP.

Extended Data Figure 2. Quality-control and cell-type annotation of single-cell RNA-seq

results

a, b, Scatter plots showing CHIP (+) patients' immune cells according to the UMI count and

other features. Percentage of the mitochondrial genes (a) and the number of detected genes (b).

c, Scatter plot of log 10-transformed count data between individual patients. R represents

Pearson's correlation coefficient values. **d,** t-SNE clusters of integrated single-cell data. **e,** Each

cell type marker gene expression on t-SNE plot. **f**, Stacked barplots show cell-type proportions

of each patient group. **g**, Boxplots showing the ratio of the proportion of classical monocytes.

(left) All pairs within CHIP (+) patients (n=30). (right) all pairs between CHIP (+) and CHIP

(-) (n=18). The box represents the interquartile range (IQR) and the whiskers correspond to the

highest and lowest points within $1.5 \times IQR$. Two-sided Kolmogorov-Smirnov test were

performed (** < p-value 0.01).

Extended Data Figure 3. Analyses of differently expressed genes between CHIP (+) severe

COVID-19 and other groups

a, Chow-ruskey venn diagram of the up-regulated genes in CHIP (+) severe COVID-19

compared to other disease groups. The color of the border in diagrams represents five immune

cell types. Monocyte (red), classical monocyte and non-classical monocyte; natural killer T

cells (orange); CD4$^+$ T cell (green), CD4$^+$ T cell and CD4$^+$ naïve T cell; CD8$^+$ T cell (blue),

CD8$^+$ T cell and CD8$^+$ memory T cells; B cell (purple), IgG$^+$ B cell and IgG$^-$ B cell. **b**, Barplots

showing combined scores for gene ontology terms in MsigDB Hallmark 2020 with up-

regulated genes in CHIP(+) severe COVID-19. Top six terms were presented. The color

indicates normal or disease controls. **c**, Ratio of cell-type specific CHIP (+) severe COVID-19

up-regulated genes compared to total up-regulated genes for comparison with other disease

groups. The mean and standard error of the mean (s.e.m) of each gene set are presented in

barplots. The color indicates the type of disease group for comparison.

Extended Data Figure 4. Enriched immune signatures in CHIP (+) and CHIP (-) severe

COVID-19 up-regulated genes compared to healthy normal control

a-e, Scatter plots showing combined scores of differentially expressed genes (DEGs) in

selected cell types between severe COVID-19 patient groups for same gene ontology library in

Fig. 2c. The horizontal axis, up-regulated genes in CHIP (+) patients; the vertical axis, up-

regulated genes in CHIP (-) patients. Identity lines are presented on diagonal. The color

indicates types of perturbed ligand in database terms. **a**, CD8+ T cell. **b**, CD8+ memory T cell.

c, IgG- B cell. **d**, CD4+ naive T cell. **e**, Natural killer T cell.

Extended Data Figure 5. An inflammatory macrophage signature in CHIP (+) classical

monocytes in severe COVID-19

a, GSEA plots for marker genes of inflammation clusters in CHIP (+) and CHIP (-), respectively.

Genes are ordered based on log-fold changes between high inflammation cluster and low

inflammation cluster. Normalized enrichment scores (NES) and FDR are presented for DEGs

between M1-like and M0 macrophage[37]. **b,** the effect of SYK inhibitor to suppress CHIP (+)

up-regulated genes. Comparison between combined scores of marker genes of high

inflammation cluster for Kinase Perturbations from GEO down gene ontology library for

inhibitors or knock out of SYK kinase terms. One-sided Paired t-test was performed (*: $P<$

0.05).

Extended Data Figure 6. The distribution of the nearest distance between differently

expressed gene promoters and DMRs

Boxplots of the nearest distance between differentially expressed genes (right: CHIP (+) up-

regulated genes, left: CHIP (-) up-regulated genes) and DMRs. Hypo-DMRs were randomly

selected to have the same sample size as hyper-DMRs. The box represents the interquartile

range (IQR) and the whiskers correspond to the highest and lowest points within $1.5 \times$ IQR.

The color indicates the types of DMRs. Distance is represented as the log scale, with a 10kb

resolution. For statistical significance test, two-sided Kolmogorov-Smirnov test were

performed between hypo- and hyper-methylated regions ($P<0.001$, CHIP (+); $P=3.46e-3$, CHIP

(-)).

Kang et al Figure 1

Kang et al Figure 2

Kang et al Figure 4

Kang et al Figure 5

Kang et al Extended Data Figure 1

Kang et al Extended Data Fig 2

a

CHIP(+) Severe COVID-19 patient up-regulated genes

vs Healthy donor   vs Flu   vs CHIP (-) Mild COVID-19   vs CHIP (-) Severe COVID-19

Cell types
- Monocyte
- Natural killer T cell
- CD4⁺ T cell
- CD8⁺ T cell
- B cell

b

Top 6 related terms in MsigDB Hallmark 2020

CHIP(+) Severe COVID-19 up-regulated genes
- vs Healthy donor
- vs Flu
- vs CHIP (-) Mild COVID-19
- vs CHIP (-) Severe COVID-19

c

Disease types
- CHIP(-) Severe COVID-19
- CHIP(-) Mild COVID-19
- Flu
- Healthy donor

Kang et al Extended Data Fig 4

Kang et al Extended Data Fig 5

a

Top 200 up-regulated genes M1-like macrophage vs M0 macrophage

b

Downregulated genes of syk inhibition

Kang et al Extended Data Fig 6

Tables

Table 1. Age and gender-adjusted odds ratio of the presence of CHIP for severe COVID-

19 in canonical risk factors-stratified subgroups

Table 2. Final multivariate model for independent risk factors for severe COVID-19

including interaction term between CHIP and canonical risk factors

Supplementary Tables

Supplementary Table 1. Baseline characteristics of COVID-19 patients in this study

according to the presence or absence of clonal hematopoiesis of indeterminate potential

Supplementary Table 2. Clinical characteristics of cluster A2 when compared other cases

in cluster A

Supplementary Table 3. Clinical information of patient used in scRNA seq

Supplementary Table 4. Data quality of scRNA seq

Supplementary Table 5. A list of DMRs

676 **Table 1. Age and gender-adjusted odds ratio of the presence of CHIP for severe COVID-**

677 **19 in canonical risk factors-stratified subgroups**

	Severe COVID-19		aOR (95% CI)	*P*
	CHIP (+)	CHIP (-)		
Canonical risk factor present	13/21 (61.9%)	47/68 (69.1%)	0.4 (0.1—1.4)	0.159
Canonical risk factor absent	12/13 (92.3%)	18/41 (43.9%)	14.8 (1.3—164.1)	0.028
Total	25/34 (73.5%)	65/109 (59.6%)	1.1 (0.4—2.8)	0.829

678 CHIP, clonal hematopoiesis with indeterminate potential; aOR, adjusted odds ratio; CI,

679 confidence interval

680 Canonical risk factors were BMI ≥ 30.0, diabetes mellitus, and hypertension.

691 **Table 2. Final multivariate model for independent risk factors for severe COVID-19**

692 **including interaction term between CHIP and canonical risk factors**

	aOR	*P*
Age, per one year	1.1 (1.0—1.1)	< 0.001
Male gender	2.0 (0.9—4.5)	0.075
CHIP	10.7 (1.1—100.7)	0.038
Canonical risk factors	2.0 (0.8—5.0)	0.120
CHIP * conventional risk factors	0.0 (0.0—0.5)	0.012

693 CHIP, clonal hematopoiesis with indeterminate potential; aOR, adjusted odds ratio

694 Canonical risk factors were BMI $\geq$ 30.0, diabetes mellitus, and hypertension.

704