Large-scale comparative genomics unravels great genomic diversity across the *Rickettsia* and *Ca.* Megaira genera and identifies Torix group as an evolutionarily distinct clade.

Helen R Davison[1], Jack Pilgrim[1], Nicky Wybouw[2], Joseph Parker[3], Stacy Pirro[4], Simon Hunter-Barnett[1], Paul M Campbell[5], Frances Blow[6], Alistair C Darby[1], Gregory D D Hurst[1] and Stefanos Siozios[1]

Affiliations

1. Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological sciences, University of Liverpool, Liverpool, L69 7ZB, United Kingdom

2. Terrestrial Ecology Unit, Department of Biology, Faculty of Sciences, Ghent University, Ghent, Belgium

3. Division of Biology and Biological Engineering, California Institute of Technology, 1200 E California Boulevard, Pasadena, CA 91125, USA

4. Iridian Genomes, Bethesda, MD, USA

5. School of Health and Life Sciences, Faculty of Biology Medicine and Health, the University of Manchester, Manchester, United Kingdom

6. Center for Genomics and Systems Biology, Department of Biology, New York University, New York, New York, USA

Abstract

Rickettsia are intracellular bacteria originally described as arthropod borne pathogens that are emerging as a diverse group of often biologically important, non-pathogenic symbionts of invertebrates and microeukaryotes. However, sparse genomic resources for symbiotic strains and for the sister genus (*Candidatus* Megaira) inhibit our understanding of *Rickettsia* evolution and biology. Here, we present the first closed genomes of *Ca.* Megaira from an alga (*Mesostigma viride),* and Torix *Rickettsia* from midge (*Culicoides impunctatus*) and bed bug (*Cimex lectularius*) hosts. Additionally, we sequenced and constructed draft genomes for *Ca.* Megaira from another alga (*Carteria cerasiformi*s)*,* Transitional group *Rickettsia* from tsetse fly (*Glossina morsitans submorsitans),* and Torix *Rickettsia* from a spider mite (*Bryobia graminum)*. We further extract 22 draft genomes from arthropod genome sequencing projects, including 1 Adalia, 4 Transitional, 1 Spotted Fever, 7 Torix, 7 Belli and the first Rhyzobius and Meloidae *Rickettsia* group genomes. We used new and existing *Rickettsia* genomes to estimate the phylogeny and metabolic potential across groups and reveal transitions in genomic properties. These data reveal Torix as unique amongst currently described *Rickettsia*, with highly distinct and diverse accessory genomes. We confirm the presence of a third subclade of Torix, previously only known from gene marker sequences. Further, Torix share an intact pentose phosphate pathway with *Ca.* Megaira, not observed in other *Rickettsia.* Considering the distinctness and diversity of Torix, we propose that the group be named *Candidatus* Tisiphia. The wide host range of *Ca.* Tisiphia symbionts necessitates onward research to understand the biological and physiological bases of *Ca.* Tisiphia-host interactions.

Importance statement

Members of the genus *Rickettsia* were originally identified as causative agents of mammalian vector-borne

disease. In the last 25 years we have recognised that many *Rickettsia* are arthropod symbionts, and sit

alongside a sister taxon, *Ca.* Megaira, which are symbiotic associates of microeukaryotes. The lack of

genomic information for symbiotic strains affects our ability to determine the evolutionary relationships

between strains and understand the biological underpinnings of the different symbioses. We clarify these

relationships by assembling 26 genomes of *Rickettsia* from understudied groups, and the first two *Ca*.

Megaira, from various insects and microeukaryotes. Of note, the accessory genome diversity and broad host

range of Torix *Rickettsia* parallels all other *Rickettsia* combined. This diversity, alongside the breadth of host

species, make the Torix clade an important hidden player in invertebrate biology and physiology. We argue

this clade should be given its own genus status, for which we propose *Ca.* Tisiphia.

Introduction

Symbiotic bacteria are vital to the function of most living eukaryotes, including microeukaryotes, fungi, plants, and animals (Boettcher et al., 1996; Clay et al., 2005; Douglas, 2011; Fujishima & Kodama, 2012). The symbioses formed are often functionally important to the host with effects ranging from mutualistic to detrimental. Mutualistic symbionts may provide benefits through the biosynthesis of metabolites, or by protecting their hosts against pathogens and parasitoids (Hendry et al., 2014; Oliver et al., 2010). Meanwhile parasitic symbionts can be detrimental to the host due to resource exploitation or through reproductive manipulation that favours its own transmission over the host's (Engelstädter & Hurst, 2009; Leclair et al., 2017). Across these different symbiotic relationships, symbionts are often important determinants of host ecology and evolution.

The Rickettsiales (Alphaproteobacteria) represent an order of obligate intracellular bacteria that form symbioses with a variety of eukaryotes (Weinert et al., 2015). Within Rickettsiales, the family Rickettsiaceae represent a diverse collection of bacteria that infect a wide range of eukaryotic hosts and can act as symbionts, parasites, and pathogens. Perhaps the best-known clade of Rickettsiaceae is the genus *Rickettsia*, which was initially described as the cause of spotted fever and other rickettsioses in vertebrates that are transmitted by ticks, lice, fleas and mites (Angelakis & Raoult, 2017).

Rickettsia have been increasingly recognised as heritable arthropod symbionts. Since the first description of a maternally inherited male-killer in ladybirds (Werren et al., 1994), we now know that heritable *Rickettsia* are common in arthropods (Pilgrim et al., 2021; Weinert et al., 2009). Further, *Rickettsia*-host symbioses are diverse, with the symbiont capable of reproductive manipulation, nutritional and protective symbiosis, as well as influencing thermotolerance and pesticide susceptibility (Bodnar et al., 2018; Brumin et al., 2011; Chiel et al., 2009; Giorgini et al., 2010; Hurst et al., 1994; Kontsedalov et al., 2008; Łukasik et al., 2013).

Our understanding of the evolution and diversity of the genus *Rickettsia* and its allies has increased in recent years. Weinert et al. (2009) defined 13 different groups of *Rickettsia* with two early branching clades that appeared genetically distant from other members of the genus. The first of these was defined from a symbiont of *Hydra* and was named the Hydra group *Rickettsia*, which has since been assigned its own genus status, *Candidatus* Megaira (Schrallhammer et al., 2013). *Ca.* Megaira forms a sister clade to *Rickettsia* and is common in ciliate protists, amoebae, chlorophyte and streptophyte algae, and cnidarians (Lanzoni et al., 2019). Members of this clade are found in

hosts from aquatic, marine and soil habitats which include model organisms (e.g., *Paramecium*,

Volvox) and economically important vertebrate parasites (e.g., *Ichthyophthirius multifiliis*, the ciliate

that causes white spot disease in fish) (Lanzoni et al., 2019). Whilst symbioses between *Ca.* Megaira

and microeukaryotes are pervasive, there is no complete publicly available genome and the impact

of these symbioses on the host are poorly understood.

A second early branching clade was first described from *Torix tagoi* leeches and is commonly coined

Torix *Rickettsia* (Kikuchi & Fukatsu, 2005). Symbionts in the Torix clade have since been found in a

wide range of invertebrate hosts from midges to freshwater snails, and in a fish-parasitic amoeba

(Pilgrim et al., 2021). The documented diversity of hosts is wider than other *Rickettsia* groups,

which are to date only found in arthropods and their associated vertebrate or plant hosts (Weinert

et al., 2009). Torix clade *Rickettsia* are known to be heritable symbionts, but their impact on host

biology is poorly understood, despite the economic and medical importance of several hosts (inc.

bed bugs, black flies, and biting midges). Rare studies have described the potential effects on the

host, which include: larger body size in leeches (Kikuchi & Fukatsu, 2005); a small negative effect on

growth rate and reproduction in bed bugs (Thongprem et al., 2020); and an association with

parthenogenesis in *Empoasca* Leafhoppers (Aguin-Pombo et al., 2021).

Current data seems to suggest an emerging macroevolutionary scenario where the members of

Rickettsia-Megaira clade originated as symbionts of microeukaryotes, before diversifying to infect

invertebrate symbionts. The Torix *Rickettsia* retained a broad range of hosts from microeukaryotes

to arthropods. The remaining members of the genus *Rickettsia* evolved to be arthropod heritable

symbionts and vector-borne pathogens. However, a lack of genomic and functional information for

symbiotic clades limits our understanding of evolutionary transitions within *Rickettsia* and its sister

groups. No *Ca.* Megaira genome sequences are currently publicly available and of the 165 *Rickettsia*

genome assemblies available on the NCBI (as of 29/04/21), only two derive from the Torix clade

and these are both draft genomes. In addition, dedicated heritable symbiont clades of *Rickettsia*,

such as the Rhyzobius group, have no available genomic data, and there is a single representative

for the Adalia clade. Despite the likelihood that heritable symbiosis with microeukaryotes and

invertebrates was the ancestral state for this group of intracellular bacteria, available genomic

resources are heavily skewed towards pathogens of vertebrates.

In this study we establish a richer base of genomic information for heritable symbiont *Rickettsia*

and *Ca.* Megaira*,* then use these resources to clarify the evolution of these groups. We broaden

available genomic data through a combination of targeted sequencing of strains without complete genomes, and metagenomic assembly of *Rickettsia* strains from arthropod genome projects. We report the first closed circular genome of a *Ca.* Megaira symbiont from a streptophyte alga (*Mesostigma viride*) and provide a draft genome for a second *Ca.* Megaira from a chlorophyte (*Carteria cerasiformis*). In addition, we present the first complete genomes of two Torix *Rickettsia* from a midge (*Culicoides impunctatus*) and a bed bug (*Cimex lectularius*) as well as a draft genome for *Rickettsia* from a tsetse fly (*Glossina morsitans submorsitans*, an important vector species), and a new strain from a spider mite (*Bryobia graminum*). A metagenomic approach established a further 22 draft genomes for insect symbiotic strains, including the first Rhyzobius and Meloidae group draft genomes. We utilize these to carry out pangenomic, phylogenomic and metabolic analyses of our extracted genome assemblies, with comparisons to existing *Rickettsia.*

Methods

Genomic data collection and construction

We employed two different workflows to assemble genomes for *Ca.* Megaira and *Rickettsia*

symbionts (Figure 1). A) Targeted sequencing and assembly of focal *Ca.* Megaira and Torix

Rickettsia. B) Assembly from SRA deposits of *Ca.* Megaira from *Mesostigma viride* NIES296 and the

29 arthropods identified in Pilgrim et al (2021) that potentially harbour *Rickettsia.* These were

analysed alongside previously assembled genomes from the genus *Rickettsia*, and the outgroup

taxon *Orientia tsutsugamushi.*

Figure 1. Workflow diagram for extraction, assembly and analyses performed in this study. Purple highlights Torix *Rickettsia and orange highlights* Ca. Megaira *and red highlights Transitional Rickettsia. A full resolution version can be found here:*
https://doi.org/10.6084/m9.figshare.15081975.

DNA preparation, sequencing strategies and symbiont assembly methodologies varied between

species. Methods are summarised in Figure 1 and detailed in supplementary material

https://doi.org/10.6084/m9.figshare.14865582. The exact pipeline used to assemble genomes from

Short Read Archive (SRA) data can be found here: https://figshare.com/s/d1155765b523a6379443.

Sample collection for targeted genome assembly

Cimex lectularius were acquired from the 'S1' isofemale colony maintained at the University of

Bayreuth described in Thongprem et al (2020). *Culicoides impunctatus* females were collected from

a wild population in Kinlochleven, Scotland (56° 42' 50.7''N 4° 57' 34.9''W) on the evenings of the

2nd and 3rd September 2020 by aspiration. *Carteria cerasiformis* strain NIES 425 was obtained from

the Microbial Culture Collection at the National Institute for Environmental Studies, Japan. The

Glossinia morsitans submorsitans specimen Gms8 was collected in Burkina Faso in 2010 and

Rickettsia infection was present alongside other symbionts as described in Doudoumis et al. (2017).

The assembly itself is a result of later thesis work (Blow, 2017).

A *Bryobia* mite community was sampled from herbaceous vegetation in Turku, Finland. The

Moomin isofemale line was established by isolating a single adult female and was maintained on

detached leaves of *Phaseolus vulgaris* L. cv Speedy at 25 °C, 60 % RH, and a 16:8 light:dark

photoperiod. The Moomin spider mite line was morphologically identified as *Bryobia graminum* by

Prof Eddie A. Ueckermann (North-West University).

Previously published Rickettsia genomes
A total of 86 published *Rickettsia* genomes, and one genome from *Orientia tsutsugamushi* were

retrieved from the European Nucleotide Archive and assessed with CheckM v1.0.13 (Parks et al.,

2015). Inclusion criteria for genomes were high completeness (CheckM > 90%), low contamination

(CheckM < 2%) and low strain heterogeneity (Check M < 50%) except in the case of Adalia for which

there is only one genome (87.6% completeness). Filtering identified 76 high quality *Rickettsia*

genomes that were used in all subsequent analyses (S1

https://figshare.com/s/198c88c6e3ea5553192e).

Genome content comparison and pangenome construction
Anvi'o 7 (Eren et al., 2021) was used to construct a pangenome for *Rickettsia.* Included in this were

the 22 MAGs retrieved from SRA data, 2 *Ca.* Megaira genomes and 4 targeted Torix *Rickettsia*

genomes, and one transitional group *Rickettsia* genome acquired in this study. To these were

added the 76 published and 1 Orientia described above, giving a total of 104 genomes. Individual

Anvi'o genome databases were additionally annotated with HMMER, KofamKOALA, and NCBI COG

profiles (Aramaki et al., 2020; Eddy, 2011; Galperin et al., 2021). For the pangenome itself,

orthologs were identified with NCBI blast, mcl inflation was set to 2, and minbit was 0.5. Genomes

were arranged according to cluster presence absence and average nucleotide sequence identity

was calculated using pyANI (Pritchard et al., 2016). See

https://figshare.com/s/d1155765b523a6379443 for the exact code used in this section.

KofamKOALA annotation (Aramaki et al., 2020) in Anvi-o 7 was used to estimate completeness of

metabolic pathways. Then Pheatmap (Kolde, 2019) in R 3.4.4 (R Core Team, 2020) was used to

produce heatmaps of metabolic potential (figure 7). Annotations for function and *Rickettsia* group

were added *post hoc* in Inkscape.

The biotin operon found in the genome Rhyzobius *Rickettsia* Oopac6 was identified from metabolic

prediction (figure 7). To confirm Oopac6 carries a complete biotin pathway that shares ancestry

with the existing *Rickettsia* biotin operon, Oopac6 biotin was compared to biotin pathways from

five other related symbionts: *Cardinium*, *Lawsonia*, *Buchnera aphidicola*, *Rickettsia buchneri*, and

Wolbachia.. A .gbk annotation was generated with PROKKA (Seemann, 2014) for Oopac6, then the

operon and surrounding genes were extracted. Clinker (Gilchrist & Chooi, 2021) with default

options was used to compare and visualise the similarity of genes within the biotin operon region of

all 6 bacteria.

Complete reference genomes were annotated with PROKKA v1.13 (Seemann, 2014). Secondary

metabolite biosynthetic gene clusters were identified using AntiSMASH version 6.0 (Blin et al.,

2021) along with Norine (Flissi et al., 2019) which searched for similarities to predicted non-

ribosomal peptides.

Functional enrichment analyses between the main *Rickettsia* clade and the Torix – *Ca.* Megaira

clades were performed using the Anvi'o program anvi-get-enriched-functions-per-pan-group and

the "COG_FUNCTION" as annotation source. A gene cluster presence - absence table was exported

using the command "anvi-export-tables". This was used to create an UpSet plot using the R package

ComplexUpset (Krassowski et al., 2020; Lex et al., 2014) to visualize unique and shared gene

clusters between different *Rickettsia* groups. A gene cluster was considered unique to a specified

Rickettsia group when it was present in at least one genome belonging to that group. Gene cluster

accumulation curves were performed for the pan-, core- and unique-genomes based on the same

presence-absence matrix using a custom-made R script (Siozios, 2021). In each case the cumulative

number of gene clusters were computed based on randomly sampled genomes using 100

permutations. The analysis was performed separately for each of the five major *Rickettsia* groups as

well as the complete *Rickettsia* dataset. Curves were plotted using the ggplot2 R package

(Wickham, 2016).

All information on extra genomes can be found at https://doi.org/10.6084/m9.figshare.14865582

and the code pipeline employed can be found at https://figshare.com/s/d1155765b523a6379443.

Phylogeny, Network, and recombination

The single-copy core of all 104 genomes was identified in Anvi'o 7 and is made up of 74 single-copy gene (SCG) clusters. Protein alignments from SCG were extracted and concatenated using the command "anvi-get-sequences-for-gene-clusters". Maximum likelihood phylogeny was constructed in IQ-TREE v2.1.2 (Nguyen et al., 2015). Additionally, 43 ribosomal proteins were identified through Anvi'o 7 to test phylogenomic relationships. These gene clusters were extracted from the pangenome and used for an independent phylogenetic analysis SUPPLEMENTARY FIG. The best model according to the Bayesian Information Criterion (BIC) was selected with Model Finder Plus (MFP) (Kalyaanamoorthy et al., 2017) as implemented in IQ-TREE; this was JTTDCMut+F+R6 for core gene clusters and JTTDCMut+F+R3 for ribosomal proteins. Both models were run with Ultrafast Bootstrapping (1000 UF bootstraps) (Hoang et al., 2018) with *Orientia tsutsugamushi* as the outgroup.

The taxonomic placement of Oopac6, Ppec13 and Dallo3 genomes within the Rhyzobius, Meloidae and Belli groups respectively were confirmed in a smaller phylogenetic analysis, performed as detailed in (Pilgrim et al. 2021) using reference MLST sequences (*gltA*, 16s rRNA, 17kDa, *COI*) from other previously identified *Rickettsia* profiles (S1 https://figshare.com/s/198c88c6e3ea5553192e). The selected models used in the concatenated partition scheme were as follows: 16S rRNA: TIM3e+I+G4; 17KDa: GTR+F+I+G4; COI: TPM3u+F+I+G4; gltA: K3Pu+F+I+G4a.

A nearest neighbour network was produced for core gene sets with default settings in Splitstree4 to further assess distances and relationships between *Rickettsia, Ca.* Megaira and Torix clades. All annotation was added post hoc in Inkscape. Furthermore, recombination signals were examined by applying the Pairwise Homoplasy Index (PHI) test to the DNA sequence of each core gene cluster extracted with Anvio-7. DNA sequences were aligned with MUSCLE (Edgar, 2004) and PHI scores calculated for each of the 74 core gene cluster with PhiPack (Bruen et al., 2006).

The taxonomic identity for new and newly expanded groups was established with GTDB-Tk (Chaumeil et al., 2020) to support the designation of new taxa through phylogenetic comparison of marker genes against an online reference database.

Results and Discussion

We have expanded the available genomic data for several *Rickettsia* groups through a combination of draft and complete genome assembly. This includes an eight-fold increase in available Torix-

group genomes, and the first available genomes for Meloidae and Rhyzobius groups. We further report the first reference genomes for *Ca*. Megaira.

Complete and closed reference genomes for Torix *Rickettsia* and *Ca.* Megaira

The use of long-read sequencing technologies produced the first complete genomes for two subclades of the Torix group (RiCimp-limoniae, RiClec-Leech). Sequencing depth of the *Rickettsia* genomes from *C. impunctatus* (RiCimp) and *C. lectularius* (RiClec) were 18X and 52X respectively. The RiCimp genome provides the first evidence of plasmids in the Torix group (pRiCimp001 and pRiCimp002). In addition, we assembled the first complete closed reference genome of *Ca.* Megaira from *Mestostigma viride* (MegNEIS296) from previously published genome sequencing efforts.

General features of both genomes are consistent with previous genomic studies of the Torix group (Table 1). A single full set of rRNAs (16S, 5S and 23S) and a GC content of ~33% was observed. Notably, the two complete Torix group genomes show a distinct lack of synteny (see S2 https://doi.org/10.6084/m9.figshare.14866263), a genomic feature that is compatible with our phylogenetic analyses that placed these two lineages in different subclades (leech/limoniae) (figures 2 and 3). Of note within the closed reference genomes MegNEIS296 and RiCimp, is the presence of a putative non-ribosomal peptide synthetase (NRPS) and a hybrid non-ribosomal peptide/polyketide synthetase (NRPS/PKS) respectively (see S3 https://doi.org/10.6084/m9.figshare.14865570). Although, the exact products of these putative pathways are uncertain, *in silico* prediction by Norine suggests close similarity with both cytotoxic and antimicrobial peptides hinting at a potential defensive role (see S3 https://doi.org/10.6084/m9.figshare.14865570). Putative toxin-antitoxin systems similar to the one associated with cytoplasmic incompatibility in *Wolbachia* have recently been observed on the plasmid of *Rickettsia felis* in a parthenogenetic booklouse. These are thought to be part of an extensive bacterial mobilome network associated with reproductive parasitism (Gillespie et al., 2018). A BLAST search found a very similar protein in Oopac6 to the putative large pLbAR toxin found in *R. felis* (88% aa identity), and a more distantly related protein in the *C. impunctatus* plasmid (25% aa identity).

 Table 1. Summary of the complete Ca. Megaira *and* Torix *Rickettsia genomes*

Group	*Ca.* Megaira	Torix *Rickettsia*	
Sub-group		Leech	Limoniae
Strain Name	MegNIES296	RiCimp	RiClec
Symbiont genome accession	SAMN21822563	SAMN21822561	SAMN21822562
Host	Mesostigma viride NIES-296	Culicoides impunctatus	Cimex lectularius
Host sequence accession	SRR8439255, SRX5120346	SRR16018514, SRR16018513	SRR16018512, SRR16018511
Total nucleotides	1,532,409	1,566,468	1,611,726
Chromosome size (bp)	1,448,425	1,469,631	1, 611,726
Plasmids	1 (83,984 bp)	2 (77550bp + 19287bp)	None
GC content (%)	33.9	32.9	32.8
Number of CDS	1,359	1,397	1,544
Avg. CDS length (bp)	998	900	874
Coding density (%)	88.5	86	84
rRNAs	3	3	3
tRNAs	34	34	35

268 *S2 Whole genome alignment between the complete* Torix *limoniae (RiCIMP) and* Torix *Leech (RiClec) genomes reveals complete lack*
269 *of synteny. Magenta represents forward matches and blue reverse matches* https://doi.org/10.6084/m9.figshare.14866263.

*S3. The circular chromosomes of **A)** a* Torix *group Rickettsia (RiCimp) and **B)** a Ca.* Megaira sp. *(MegNEIS296). From outside to in, the circles represent: forward CDSs (Red), Reverse CDSs (blue), tRNAs (yellow) rRNAs (black), and GC content (green and magenta). Highlighted are the predicted modules formed by non-ribosomal peptide synthase genes (domains) that define individual amino acids in the synthesised peptide and show the catalytic domains within modules https://doi.org/10.6084/m9.figshare.14865570.*

Sequencing and *de novo* assembly of other *Rickettsia* and *Ca*. Megaira genomes.

Our direct sequencing efforts enabled assembly of draft genomes for a second *Ca*. Megaira strain from the alga *Carteria cerasiformis*, and for *Rickettsia* associated with tsetse flies and *Bryobia* spider mites. The Transitional *Rickettsia* from a wild caught Tsetse fly, RiTSETSE, is a potentially chimeric assembly since we identified an excess of biallelic sites when the raw Illumina reads were mapped back to the assembly. It is also likely that RiTSETSE is not a heritable symbiont but comes from transient infection from a recent blood meal.

From the SRA accessions, the metagenomic pipeline extracted 29 full symbiont genomes for Rickettsiales across 24 host species. Five of 29 were identified as *Wolbachia* and discarded from further analysis, one was a *Rickettsia* discarded for low quality, and another was a previously assembled Torix *Rickettsia*, RiCNE (Pilgrim et al., 2017). Thus, 22 high quality *Rickettsia* metagenomes were obtained from 21 host species. One beetle (SRR6004191) carried coinfecting *Rickettsia* Lappe3 and Lappe4 (Table 2). The high-quality *Rickettsia* covered the Belli, Torix, Transitional, Rhyzobius, Meloidae and Spotted Fever Groups (Table 2 and S1 https://figshare.com/s/198c88c6e3ea5553192e).

Beetles, particularly rove beetle (Staphylinidae) species, appear in this study as a possible hotspot

of *Rickettsia* infection. *Rickettsia* has historically been commonly associated with beetles, including

ladybird beetles (*Adalia bipunctata*), diving beetles (*Deronectes* sp.) and bark beetles (*Scolytinae*)

(Hurst et al., 1994; Küchler et al., 2009; Perlman et al., 2006; Weinert et al., 2009; Zchori-Fein et al.,

2006). Though a plausible and likely hotspot, this observation needs be approached with caution as

this could be an artefact of skewed sampling efforts.

All genome metadata and source information can be found here

https://figshare.com/s/198c88c6e3ea5553192e.

Table 2. Brief summary of draft genomes generated during the current project and their associated hosts. Full metadata can be found
in S1 https://figshare.com/s/198c88c6e3ea5553192e.

Strain	Bacteria Biosample Accession	Group	Number of contigs	Total length (bp)	Host name	Order
Blapp1	SAMN21822536	Belli	171	1266633	*Bembidion lapponicum*	Coleoptera
Btrans1	SAMN21822537	Belli	241	1417452	*Bembidion nr. transversale* OSAC:DRMaddison DNA3205	Coleoptera
Choog2	SAMN21822538	Belli	16	1357829	*Columbicola hoogstraali*	Phthiraptera
Cmasu2	SAMN21822539	Transitional	196	1295004	*Ceroptres masudai*	Hymenoptera
Dallo3	SAMN21822540	Belli	196	990679	*Diachasma alloeum*	Hymenoptera
Drufa1	SAMN21822541	Belli	14	1364611	*Degeeriella rufa*	Phthiraptera
Earac4	SAMN21822542	Transitional	96	1350066	*Ecitomorpha arachnoides*	Coleoptera
Econn1	SAMN21822543	Transitional	238	1070326	*Eriopis connexa*	Coleoptera
Gbili3	SAMN21822544	Torix limoniae	171	1188102	*Gnoriste bilineata*	Diptera
Gdoso1	SAMN21822545	Belli	34	1420758	*Graphium doson*	Lepidoptera
Lappe3	SAMN21822558	Torix limoniae	122	1368980	*Labidopullus appendiculatus*	Coleoptera
Lappe4	SAMN21822559	Torix limoniae	154	1332357	*Labidopullus appendiculatus*	Coleoptera
MegCarte-ria	SAMN21822546	*Ca*. Megaira	72	1298707	*Carteria cerasiformis*	Chlamydomonadales
Ofont3	SAMN21822560	Adalia	91	1529137	*Omalisus fontisbellaquei*	Coleoptera
Oopac6	SAMN21822548	Rhyzobius	181	1497231	*Oxypoda opaca*	Coleoptera
Pante1	SAMN21822549	Torix limoniae	70	1472610	*Pseudomimeciton antennatum*	Coleoptera
Pfluc4	SAMN21822550	Spotted Fever Group	7	1251895	*Proechinophthirus fluctus*	Phthiraptera
Ppec13	SAMN21822551	Belli	90	1426047	*Pyrocoelia pectoralis*	Coleoptera
Psono2	SAMN21822552	Torix limoniae	163	1492063	*Platyusa sonomae*	Coleoptera
RiTSETSE	SAMN21822553	Transitional	172	1451997	*Glossina morsitans submorsitans*	Diptera
S2	SAMN21822554	Torix limoniae	103	1251484	*Sericostoma*	Trichoptera
Sanch3	SAMN21822555	Belli	181	1487154	*Stiretrus anchorago*	Hemiptera
Slati1	SAMN21822556	Transitional	109	1301763	*Sceptobius lativentris*	Coleoptera
Ssp4	SAMN21822557	Torix limoniae	87	1231013	*Sericostoma sp.* HW-2014	Trichoptera
moomin	SAMN21822560	Torix moomin	204	1137559	*Bryobia graminum*	Trombidiformes

Phylogenomic analyses and taxonomic placement of newly assembled genomes

Phylogeny, network, and recombination

The network and phylogeny illustrate the distance of Torix from *Ca.* Megaira and other *Rickettsia*, along with an extremely high level of within-group diversity in Torix compared to any other group (Figures 2 and 3). The phylogenies generated using core genomes are consistent with previously identified *Rickettsia* and host associations using more limited genetic markers. For instance, Pfluc4 from *Proechinophthirus fluctus* lice is grouped on the same branch as a previously sequenced *Rickettsia* from a different individual of *P. fluctus*. Four of 22 genomes from the SRA screen are identified as Transitional, 1 is in Spotted Fever Group, 1 is Adalia, 8 are Belli and 7 are Torix limoniae. Targeted sequences were confirmed as: Torix limoniae (RiCimp), Torix leech (RiClec), Transitional (RiTSETSE), *Ca*. Megaira (MegCarteria and MegNEIS296), and a new Torix clade, Moomin (Moomin). The new Torix include one double infection giving a total of 10 new genomes across 9 potential host species. The double infection is found within the rove beetle *Labidopullus appendiculatus*, forming Lappe3 and Lappe4 (Fig 2 and 3).

In addition, the pre-existing *Rickettsia helvetica*, which is typically cited as a member of the Spotted Fever group as a result of its first description in 1993 (Beati et al., 1993; Chisu et al., 2017), seems to form its own group in all trees and networks (figure 2, 3 and https://doi.org/10.6084/m9.figshare.14865606). We conclude from this that *Rickettsia helvetica* is most similar to Scapularis group *Rickettsia,* but because it does not fall into the same clade in any tree or network, it is likely that the strain belongs to a distinct lineage of tick-borne *Rickettsia*.

Core Phylogeny

Figure 2. Rickettsia and Ca. Megaira maximum likelihood (ML) phylogeny constructed from 74 core gene clusters extracted from the pangenome. New genomes are indicated by ▲ *and bootstrap values based on 1000 replicates are indicated with coloured circles. New complete genomes are: RiCimp, RiClec and MegNEIS296. A full resolution version can be found here: https://doi.org/10.6084/m9.figshare.15081975.*

Ribosomal Phylogeny

S4. Rickettsia and Ca. Megaira maximum likelihood (ML) phylogeny constructed from 43 ribosomal protein gene clusters extracted from the pangenome. New genomes are indicated by ▲ and bootstrap values based on 1000 replicates are indicated with coloured circles. New complete genomes are: RiCimp, RiClec and MegNEIS296. https://doi.org/10.6084/m9.figshare.14865606

We also report the first putative Rhyzobius *Rickettsia* genomes extracted from the staphylinid

beetle *Oxypoda opaca* (Oopac6) and Meloidae *Rickettsia* from the firefly *Pyrocoelia pectoralis*

(Ppec13). They have high completeness (S1 https://figshare.com/s/198c88c6e3ea5553192e), low

pseudogenisation, and consistently group away from the other draft and completed genomes

(Figures 2 and 3). MLST analyses demonstrate that these bacteria are most like the Rhyzobius and

Meloidae groups described by Weinert *et al.* (2009) (see S5

https://doi.org/10.6084/m9.figshare.14865600). The pangenome and metabolic profile of this draft

genome suggests that Meloidae is a sister group to Belli and that Rhyzobius *Rickettsia* is

superficially similar to Belli and Transitional groups. The Rhyzobius-group symbiont is

phylogenetically distant from most *Rickettsia* and is potentially a sister clade linking Torix and the

main *Rickettsia* clades. Further genome construction will help clarify this taxon and its relationship

to the rest of *Rickettsia.*

348   *S5 Phylogram of a maximum likelihood (ML) tree of 90 Rickettsia mutilocus profiles. The tree is based on 4 loci, 16S rRNA, 17KDa,*
349   *gltA, and COI, under a partition model (2,781 bp total). https://doi.org/10.6084/m9.figshare.14865600*

The sequencing data for the wasp, *Diachasma alloeum,* used here has previously been described to contain a pseudogenised nuclear insert of *Rickettsia* material, but not a complete *Rickettsia* genome (Tvedte et al., 2019). The construction of a full, non-pseudogenised genome with higher read depth than the insect contigs, low contamination (0.95%) and high completion (93.13%) suggests that these reads likely represent a viable *Rickettsia* infection in *D. alloeum*. However, these data do not exclude the presence of an additional nuclear insert. It is possible for a whole symbiont genome to be incorporated into the host's DNA (Hotopp et al., 2007), and there are recorded partial inserts of *Ca*. Megaira genomes in the *Volvox carteri* genome (Kawafune et al., 2015). The presence of both the insert and symbiont need confirmation through appropriate microscopy methods.

Recombination is low within the core genomes of *Rickettsia* and *Ca.* Megaira, but may occur between closely related clades that are not investigated here. Across all genomes, the PHI score is significant in 6 of the 74 core gene clusters, suggesting putative recombination events. However, it is reasonable to assume that most of these may be a result of systematic error due to the divergent evolutionary processes at work across *Rickettsia* genomes. Patterns of recombination can occur by chance rather than driven by evolution which cannot be differentiated by current phylogenetic methods (Murray et al., 2016). The function of each respective cluster can be found at https://figshare.com/s/198c88c6e3ea5553192e.

Gene content and pangenome analysis
Pangenome
Across all 104 genomes used in the pangenome (figure 2, full data in S6 https://doi.org/10.6084/m9.figshare.14865576), Anvi'o found 74 core gene clusters which were used to construct a phylogeny (Figure 4). Bacterial strains of the different *Rickettsia* groups, especially the neglected symbiotic Rickettsiaceae, seem to have large, open genomes capable of rapid evolution. As expected, the more genomes that are included in analyses, the smaller the core genome extracted.

Torix is a distinctly separate clade sharing less than 70% ANI similarity to any *Rickettsia* or Cand. *Megaira* genomes. It contains at least three groups that reflect its highly diverse niche in the environment (figure 5) (Jain et al., 2018; Pilgrim et al., 2021; Rodriguez-R et al., 2021). Torix has the most unique genes out of all the clades in this study (figure 6) and is the group where each additional genome included increases the pangenome repertoire to the greatest extent (figure 7).

Torix group is thus more diverse in terms of genome content and size of the pangenome than other

Rickettsia groups.

Rickettsia lineages group together based on gene presence/absence and produce repeated patterns

of accessory genes that reliably occur within each group (figure 2). ANI scores are also strongest

within groups, while genomes tend to share lower similarity outside of their group (figure 4). This is

particularly apparent in Torix and *Ca.* Megaira which are divergent from the main *Rickettsia* clade

(figure 3 and 5).

Figure 4. Pangenome of all 104 genomes including Rickettsia, Torix, Ca. Megaira *and the outgroup Orientia. New genomes are*
indicated by ●. Each genome displays gene cluster presence/absence and is organised by gene cluster frequency. Group identity was
assigned from phylogeny. SFG is Spotted Fever Group. A full resolution version can be found here:
https://doi.org/10.6084/m9.figshare.15081975.

Figure 5. Pairwise Average Nucleotide Identity percentage across all genomes. New genomes are indicated by ●. A full resolution version can be found here: https://doi.org/10.6084/m9.figshare.15081975.

Figure 6. Shared and unique gene clusters across A) All Rickettsia and Ca. Megaira genomes used in this study grouped by clade with Orientia as an outgroup B) all individual Torix genomes, and C) all individual Belli genomes. Horizontal grey bars to the right of each plot represent gene cluster size and vertical, coloured bars represent the size of intersections (the number of shared gene clusters) between genomes in descending order with known COG functions displayed in red and unknown in blue. Black dots mean the cluster is present and connected dots represent gene clusters that are present across groups. SFG is Spotted Fever Group and TG is Typhus Group. A full resolution version can be found here: https://doi.org/10.6084/m9.figshare.15081975.

Figure 7. Gene cluster accumulation curves for pangenome (A), core genome (B) and the unique genome (C) of the 5 largest Rickettsia groups as a function of the number of genomes sequenced. The pangenome and the core genome accumulation curves for the complete Rickettsia dataset is shown in panel D. Error bars represent ± standard deviation based on 100 permutations. SFG is Spotted Fever Group and TG is Typhus Group. A full resolution version can be found here: https://doi.org/10.6084/m9.figshare.15081975.

Gene content and metabolic analyses

Rickettsial genomes extracted from SRA samples are generally congruent with the metabolic

potential of their respective groups (Figure 8). Torix and *Ca.* Megaira have complete pentose

phosphate pathways (PPP); a unique marker for these groups which seems to have been lost in the

other *Rickettsia* clades. The PPP generates NADPH, precursors to amino acids, and is known to

protect against oxidative injury in some bacteria (Christodoulou et al., 2018), as well as conversion

of hexose monosaccharides into pentose used in nucleic acid and exopolysaccharide synthesis. The

PPP has also been associated with establishing symbiosis between the Alphaproteobacteria

Sinorhizobium meliloti and its plant host *Medicago sativa* (Hawkins et al., 2018). This pathway has

previously been highlighted in Torix (Pilgrim et al., 2017) and its presence in all newly assembled

Torix and *Ca.* Megaira draft genomes consolidates its importance as an identifying feature for these

groups (Figure 8, S1 https://figshare.com/s/198c88c6e3ea5553192e). The PPP is likely an ancestral

feature that was lost in the main *Rickettsia* clade.

Glycolysis, gluconeogenesis and cofactor and vitamin metabolism are absent or incomplete across

all *Rickettsia*, except the Rhyzobius group member, Oopac6 (Figure 8). Oopac6 has a complete

biotin synthesis pathway that is related to, but distinct from, the biotin pathway found in *Rickettsia*

buchneri (See S7 https://doi.org/10.6084/m9.figshare.14865567). Based on the clinker plot and an

independent blastx search, the *GlyA* gene in *Rickettsia buchneri* appears to be a misidentified *bioF*

gene (see S7 https://doi.org/10.6084/m9.figshare.14865567). Additionally, the insect SRA sample

was not infected with *Wolbachia*, making it unlikely that the presence of the biotin operon is a

result of misassembly. Animals can't synthesize B-vitamins, so they either acquire them from diet or

from microorganisms that can synthesize them. Oopac6 has retained or acquired a complete biotin

operon where this operon is absent in other members of the genus. Biotin pathways in insect

symbionts can be an indicator of nutritional symbioses (Douglas, 2017), so Rhyzobius *Rickettsia*

could contribute to the feeding ecology of the beetle *O. opaca*. However, like other aleocharine

rove beetles, *O. opaca* is likely predaceous, omnivorous or fungivorous (analysis of gut contents

from a related species, *O. grandipennis*, revealed a high prevalence of yeasts: Klimaszewski et al.,

2013) We posit no obvious reason for how these beetles benefit from harbouring a biotin-

producing symbiont. One theory is that this operon could be a 'hangover' from a relatively recent

host shift event and may have been functionally important in the original host. Similarly, if the

symbiont is undergoing genome degradation, a once useful biotin pathway may be present but not

functional (Blow et al., 2020). As this is the only member of this group with a complete genome so

far, further research is required to firmly establish the presence and function of this pathway.

S7 Biotin operon of the Rhyzobius Rickettsia, Oopac6, and its surrounding genes compared with other known biotin pathways in other related symbionts. Similarity scores in the black boxes refer to the percentage identity between the genes of the operon above or

A 75% complete dTDP-L-rhamnose biosynthesis pathway was observed in 4 of the draft belli

assemblies (Gdoso1, Choog2, Drufa1, Blapp1) (figure 8). Two host species are bird lice (*Columbicola*

hoogstraali, *Degeeriella rufa*), one is a butterfly (*Graphium doson*), and one is a ground beetle

(*Bembidion lapponicum*). dTDP-L-rhamnose is an essential component of human pathogenic

bacteria like *Pseudomonas*, *Streptococcus* and *Enterococcus*, where it is used in cell wall

construction (van der Beek et al., 2019). This pathway has also been utilized in the synthesis of

plant cell walls (Jiang et al., 2021), may be involved in the moulting process of *Caenorhabditis*

elegans (Feng et al., 2016), and is a precursor to rhamnolipids that are used in quorum sensing

(Daniels et al., 2004). In the root symbiont *Azospirillium*, disruption of this pathway alters root

colonisation, lipopolysaccharide structure and exopolysaccharide production (Jofré et al., 2004). No

Rickettsia from typically pathogenic groups assessed in figure 8 has this pathway, and the hosts of

these four bacteria are not involved with human or mammalian disease. Presence in feather lice

provides little opportunity for this *Rickettsia* to be pathogenic as feather lice are chewers rather

than blood feeders, and Belli group *Rickettsia* more generally are rarely pathogenic. Further, this

association does not explain its presence in a butterfly and ground beetle; it is most likely that this

pathway, if functional, would be involved in establishing infection in the insect host or host-

symbiont recognition.

Figure 8. Heatmaps of predicted KEGG pathway completion estimated in Anvi'o 7, separated by function and produced with Pheatmap. Pathways of interest are highlighted: A) The pentose phosphate pathway only present in Torix *and Ca.* Megaira*, B) the biotin pathway present only in the Rhyzobius Rickettsia Oopac6, C) NAD biosynthesis only present in Moomin Rickettsia, D) dTDP-L-rhamnose biosynthesis pathway in Gdoso1, Choog2, Drufa1, and Blapp1. SFG is Spotted Fever. A full resolution version can be found here: https://doi.org/10.6084/m9.figshare.15081975.*

Designation of *Ca.* Tisiphia

In all analyses, Torix consistently cluster away from the rest of *Rickettsia* as a sister taxon. Despite the relatively small number of Torix genomes, its within group diversity is greater than any divergence between previously described *Rickettsia* in any other group (figures 2, 3 and 5). Additionally, Torix shares characteristics with both *Ca.* Megaira and *Rickettsia*, but with many of its own unique features (figures 6 and 8). The distance of Torix from other *Rickettsia* and *Ca.* Megaira is confirmed in both the phylogenomic and metabolic function analyses to the extent that Torix should be separated from *Rickettsia* and assigned its own genus. This is supported by GTDB-Tk analysis which places all Torix genomes separate from *Rickettsia* (S1 https://figshare.com/s/198c88c6e3ea5553192e) and ANI less than 70% in all cases. To this end, we propose the name *Candidatus* Tisiphia after the fury Tisiphone, reflecting the genus *Ca.* Megaira being named after her sister Megaera.

Conclusion

The bioinformatics approach has successfully extracted a substantial number of novel *Rickettsia* and *Ca*. Megaira genes from existing SRA data, including the first putative Rhyzobius *Rickettsia* and several *Ca. Tisiphia* (formerly Torix *Rickettsia*). Successful completion of two *Ca*. Megaira and two *Ca.* Tisiphia genomes provide solid reference points for the evolution of *Rickettsia* and its sister groups. From this, we can confirm the presence of a complete Pentose Phosphate Pathway in *Ca.* Tisiphia and *Ca.* Megaira, suggesting that this pathway was lost during *Rickettsia* evolution. We also describe the first Meloidae and Rhyzobius *Rickettsia* and show that Rhyzobius group *Rickettsia* has the potential to be a nutritional symbiont due to the presence of a complete biotin pathway. These new genomes provide a much-needed expansion of available data for symbiotic *Rickettsia* clades and clarification on the evolution of *Rickettsia* from *Ca.* Megaira and *Ca.* Tisiphia.

Supporting information

All original genomes and raw readsets produced in this study can be accessed at Bioproject accession PRJNA763820 and all assemblies produced from previously published third party data can be accessed at Bioproject PRJNA767332.

Supplementary data and full resolution figures can be accessed on figshare here:
https://doi.org/10.6084/m9.figshare.c.5518182.v1

Acknowledgements

HRD was supported by the NERC ACCE Doctoral Training Programme. Grant code: NE/L002450/1 and NW was supported by a BOF post-doctoral fellowship (Ghent University, 01P03420) and by a

Research Foundation - Flanders (FWO) Research Grant (1513719N). Funding for tsetse fly genomics

were to ACD IP BBSRC projects BB/J017698/1 and BB/K501773/1, the materials from which were

provided by P. Solano (Institut de Recherche pour le Développement, Unité Mixte de Recherche

Interactions Hôtes-Vecteurs-Parasites-Environnement dans les Maladies Tropicales Négligées Dues

aux Trypanosomatides, 34398, Montpellier, France) and J.-B. Rayaisse (Centre International de

Recherche-Développement sur l'Élevage en zone Subhumide (CIRDES), N°559, Rue 5-31 Avenue du

Gouverneur Louveau, 01 BP 454, Bobo Dioulasso 01, Burkina Faso). Jean-Baptiste died a few years

ago but he was a fantastic person to work with and a great field entomologist. We also wish to

thank Dr David Montagnes for teaching skills associated with algal culture.

We wish to thank Dr Débora Pires Paula (Embrapa) for granting permission to use SRA data for

sample number SRR5651504, Iridian Genomes for allowing use of their SRA data, and the Microbial

Culture Collection at the National Institute for Environmental Studies, Japan for use of the sample

Carteria cerasiformis NIES-425.

Contributions

Project concept: HRD, SS, JP and GH

Manuscript written by HRD, SS, JP and GDDH

SRA dive and metagenome assembly carried out by HRD with aid from SS.

Assembly of genome from SRA, pangenomics and phylogenomics carried out by HRD with advice from SS,
GH

Metabolic analysis carried out by HRD, JP and SS

Sequencing and assembly of bacteria from Cimex lecticularius and Culicoides impunctatus genomes by SS
and JP.

Sequencing and assembly of symbionts from Carteria by SHB and SS, supervised by PC and GH.

Sequencing and construction of RiTSETSE carried out by FB as part of thesis work supervised by AD.

SP collected and sequenced staphylinid genomes that were released through NCBI by iridian genomics.

NW collected and sequenced the *Bryobia* Moomin strain and performed preliminary metagenomic analyses

References

Aguin-Pombo, D., Rodrigues, M. C. P. A., Voetdijk, B., & Breeuwer, J. A. J. (2021). Parthenogenesis

and Sex-Ratio Distorting Bacteria in Empoasca (Hemiptera: Cicadellidae) Leafhoppers. *Annals*

of the Entomological Society of America. https://doi.org/10.1093/AESA/SAAB025

Angelakis, E., & Raoult, D. (2017). Rickettsia and Rickettsia-Like Organisms. In *Infectious Diseases*

(pp. 1666-1675.e1). Elsevier. https://doi.org/10.1016/b978-0-7020-6285-8.00187-8

Aramaki, T., Blanc-Mathieu, R., Endo, H., Ohkubo, K., Kanehisa, M., Goto, S., & Ogata, H. (2020).

KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score threshold.

Bioinformatics, *36*(7), 2251–2252. https://doi.org/10.1093/bioinformatics/btz859

Beati, L., Peter, O., Burgdorfer, W., Aeschlimann, A., & Raoult, D. (1993). Confirmation that

Rickettsia helvetica sp. nov. Is a Distinct Species of the Spotted Fever Group of Rickettsiae.

International Journal of Systematic Bacteriology, *43*(3), 521–526.

https://doi.org/10.1099/00207713-43-3-521

Blin, K., Shaw, S., Kloosterman, A. M., Charlop-Powers, Z., van Wezel, G. P., Medema, M. H., &

Weber, T. (2021). antiSMASH 6.0: improving cluster detection and comparison capabilities.

Nucleic Acids Research, *49*(W1), W29–W35. https://doi.org/10.1093/nar/gkab335

Blow, F. (2017). *Variation in the structure and function of invertebrate-associated bacterial*

communities [University of Liverpool]. https://doi.org/10.17638/03009325

Blow, F., Bueno, E., Clark, N., Zhu, D. T., Chung, S. H., Güllert, S., Schmitz, R. A., & Douglas, A. E.

(2020). B-vitamin nutrition in the pea aphid-Buchnera symbiosis. *Journal of Insect Physiology*,

126, 104092. https://doi.org/10.1016/j.jinsphys.2020.104092

Bodnar, J. L., Fitch, S., Rosati, A., & Zhong, J. (2018). The folA gene from the Rickettsia

endosymbiont of Ixodes pacificus encodes a functional dihydrofolate reductase enzyme. *Ticks*

and Tick-Borne Diseases, *9*(3), 443–449. https://doi.org/10.1016/j.ttbdis.2017.12.013

Boettcher, K. J., Ruby, E. G., & McFall-Ngai, M. J. (1996). Bioluminescence in the symbiotic squid

Euprymna scolopes is controlled by a daily biological rhythm. *Journal of Comparative*

Physiology A, *179*(1), 65–73. https://doi.org/10.1007/BF00193435

Bruen, T. C., Philippe, H., & Bryant, D. (2006). A Simple and Robust Statistical Test for Detecting the

Presence of Recombination. *Genetics*, *172*(4), 2665–2681.

https://doi.org/10.1534/genetics.105.048975

Brumin, M., Kontsedalov, S., & Ghanim, M. (2011). Rickettsia influences thermotolerance in the

whitefly Bemisia tabaci B biotype. *Insect Science*, *18*(1), 57–66. https://doi.org/10.1111/j.1744-

7917.2010.01396.x

Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2020). GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. *Bioinformatics*, *36*(6), 1925–1927.

https://doi.org/10.1093/BIOINFORMATICS/BTZ848

Chiel, E., Inbar, M., Mozes-Daube, N., White, J. A., Hunter, M. S., & Zchori-Fein, E. (2009).

Assessments of Fitness Effects by the Facultative Symbiont <I>Rickettsia</I> in the

Sweetpotato Whitefly (Hemiptera: Aleyrodidae). *Annals of the Entomological Society of*

America, *102*(3), 413–418. https://doi.org/10.1603/008.102.0309

Chisu, V., Leulmi, H., Masala, G., Piredda, M., Foxi, C., & Parola, P. (2017). Detection of Rickettsia

hoogstraalii , Rickettsia helvetica , Rickettsia massiliae , Rickettsia slovaca and Rickettsia

aeschlimannii in ticks from Sardinia, Italy. *Ticks and Tick-Borne Diseases*, *8*(3), 347–352.

https://doi.org/10.1016/j.ttbdis.2016.12.007

Christodoulou, D., Link, H., Fuhrer, T., Kochanowski, K., Gerosa, L., & Sauer, U. (2018). Reserve Flux

Capacity in the Pentose Phosphate Pathway Enables Escherichia coli's Rapid Response to

Oxidative Stress. *Cell Systems*, *6*, 569–578. https://doi.org/10.1016/j.cels.2018.04.009

Clay, K., Holah, J., & Rudgers, J. A. (2005). Herbivores cause a rapid increase in hereditary symbiosis

and alter plant community composition. *Proceedings of the National Academy of Sciences*,

102(35), 12465–12470. https://doi.org/10.1073/pnas.0503059102

Daniels, R., Vanderleyden, J., & Michiels, J. (2004). Quorum sensing and swarming migration in

bacteria. *FEMS Microbiology Reviews*, *28*(3), 261–289.

https://doi.org/10.1016/j.femsre.2003.09.004

Doudoumis, V., Blow, F., Saridaki, A., Augustinos, A., Dyer, N. A., Goodhead, I., Solano, P., Rayaisse,

J.-B., Takac, P., Mekonnen, S., Parker, A. G., Abd-Alla, A. M. M., Darby, A., Bourtzis, K., &

Tsiamis, G. (2017). Challenging the Wigglesworthia , Sodalis , Wolbachia symbiosis dogma in

tsetse flies: Spiroplasma is present in both laboratory and natural populations. *Scientific*

Reports 2017 7:1, *7*(1), 1–13. https://doi.org/10.1038/s41598-017-04740-3

Douglas, A. E. (2011). Lessons from studying insect symbioses. In *Cell Host and Microbe* (Vol. 10,

Issue 4, pp. 359–367). Cell Press. https://doi.org/10.1016/j.chom.2011.09.001

Douglas, A. E. (2017). The B vitamin nutrition of insects: the contributions of diet, microbiome and

horizontally acquired genes. *Current Opinion in Insect Science*, *23*, 65–69.

https://doi.org/10.1016/j.cois.2017.07.012

Eddy, S. R. (2011). Accelerated Profile HMM Searches. *PLoS Computational Biology*, *7*(10),

e1002195. https://doi.org/10.1371/journal.pcbi.1002195

Edgar, R. C. (2004). MUSCLE: A multiple sequence alignment method with reduced time and space

complexity. *BMC Bioinformatics*, *5*(1), 1–19. https://doi.org/10.1186/1471-2105-5-113

Engelstädter, J., & Hurst, G. D. D. (2009). The Ecology and Evolution of Microbes that Manipulate

Host Reproduction. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 127–149.

https://doi.org/10.1146/annurev.ecolsys.110308.120206

Eren, A. M., Kiefl, E., Shaiber, A., Veseli, I., Miller, S. E., Schechter, M. S., Fink, I., Pan, J. N., Yousef,

M., Fogarty, E. C., Trigodet, F., Watson, A. R., Esen, Ö. C., Moore, R. M., Clayssen, Q., Lee, M.

D., Kivenson, V., Graham, E. D., Merrill, B. D., … Willis, A. D. (2021). Community-led, integrated,

reproducible multi-omics with anvi'o. *Nature Microbiology*, *6*(1), 3–6.

https://doi.org/10.1038/s41564-020-00834-3

Feng, L., Shou, Q., & Butcher, R. A. (2016). Identification of a dTDP-rhamnose biosynthetic pathway

that oscillates with the molting cycle in Caenorhabditis elegans. *Biochemical Journal*, *473*(11),

1507–1521. https://doi.org/10.1042/BCJ20160142

Flissi, A., Ricart, E., Campart, C., Chevalier, M., Dufresne, Y., Michalik, J., Jacques, P., Flahaut, C.,

Lisacek, F., Leclère, V., & Pupin, M. (2019). OUP accepted manuscript. *Nucleic Acids Research*,

48(D1), D465–D469. https://doi.org/10.1093/nar/gkz1000

Fujishima, M., & Kodama, Y. (2012). Endosymbionts in Paramecium. *European Journal of*

Protistology, *48*(2), 124–137. https://doi.org/10.1016/j.ejop.2011.10.002

Galperin, M. Y., Wolf, Y. I., Makarova, K. S., Vera Alvarez, R., Landsman, D., & Koonin, E. V. (2021).

COG database update: focus on microbial diversity, model organisms, and widespread

pathogens. *Nucleic Acids Research*, *49*(D1), D274–D281.

https://doi.org/10.1093/nar/gkaa1018

Gilchrist, C. L. M., & Chooi, Y.-H. (2021). clinker & clustermap.js: automatic generation of gene

cluster comparison figures. *Bioinformatics*. https://doi.org/10.1093/bioinformatics/btab007

Gillespie, J. J., Driscoll, T. P., Verhoeve, V. I., Rahman, M. S., Macaluso, K. R., & Azad, A. F. (2018). A

Tangled Web: Origins of Reproductive Parasitism. *Genome Biology and Evolution*, *10*(9), 2292–2309. https://doi.org/10.1093/gbe/evy159

Giorgini, M., Bernardo, U., Monti, M. M., Nappo, A. G., & Gebiola, M. (2010). Rickettsia symbionts cause parthenogenetic reproduction in the parasitoid wasp pnigalio soemius (hymenoptera: Eulophidae). *Applied and Environmental Microbiology*, *76*(8), 2589–2599. https://doi.org/10.1128/AEM.03154-09

Hawkins, J. P., Ordonez, P. A., & Oresnik, I. J. (2018). Characterization of Mutations That Affect the Nonoxidative Pentose Phosphate Pathway in Sinorhizobium meliloti. *Journal of Bacteriology*, *200*(2). https://doi.org/10.1128/JB.00436-17

Hendry, T. A., Hunter, M. S., & Baltrus, D. A. (2014). The facultative symbiont Rickettsia protects an invasive whitefly against entomopathogenic Pseudomonas syringae strains. *Applied and Environmental Microbiology*, *80*(23), 7161–7168. https://doi.org/10.1128/AEM.02447-14

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q., & Vinh, L. S. (2018). UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution*, *35*(2), 518–522. https://doi.org/10.5281/zenodo.854445

Hotopp, J. C. D., Clark, M. E., Oliveira, D. C. S. G., Foster, J. M., Fischer, P., Torres, M. C. M., Giebel, J. D., Kumar, N., Ishmael, N., Wang, S., Ingram, J., Nene, R. V., Shepard, J., Tomkins, J., Richards, S., Spiro, D. J., Ghedin, E., Slatko, B. E., Tettelin, H., & Werren, J. H. (2007). Widespread Lateral Gene Transfer from Intracellular Bacteria to Multicellular Eukaryotes. *Science*, *317*(5845), 1753–1756. https://doi.org/10.1126/science.1142490

Hurst, G. D. D., Purvis, E. L., Sloggett, J. J., & Majerus, M. E. N. (1994). The effect of infection with male-killing Rickettsia on the demography of female Adalia bipunctata L. (two spot ladybird). *Heredity*, *73*(3), 309–316. https://doi.org/10.1038/hdy.1994.138

Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T., & Aluru, S. (2018). High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature Communications*, *9*(1), 5114. https://doi.org/10.1038/s41467-018-07641-9

Jiang, N., Dillon, F. M., Silva, A., Gomez-Cano, L., & Grotewold, E. (2021). Rhamnose in plants - from biosynthesis to diverse functions. *Plant Science*, *302*, 110687. https://doi.org/10.1016/j.plantsci.2020.110687

Jofré, E., Lagares, A., & Mori, G. (2004). Disruption of dTDP-rhamnose biosynthesis modifies

lipopolysaccharide core, exopolysaccharide production, and root colonization in Azospirillum

brasilense. *FEMS Microbiology Letters*, *231*(2), 267–275. https://doi.org/10.1016/S0378-

1097(04)00003-5

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., Von Haeseler, A., & Jermiin, L. S. (2017).

ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods*,

14(6), 587–589. https://doi.org/10.1038/nmeth.4285

Kawafune, K., Hongoh, Y., Hamaji, T., Sakamoto, T., Kurata, T., Hirooka, S., Miyagishima, S., &

Nozaki, H. (2015). Two Different Rickettsial Bacteria Invading Volvox carteri. *PLOS ONE*, *10*(2),

e0116192. https://doi.org/10.1371/JOURNAL.PONE.0116192

Kikuchi, Y., & Fukatsu, T. (2005). Rickettsia Infection in Natural Leech Populations. *Microbial

Ecology*, *49*(2), 265–271. https://doi.org/10.1007/s00248-004-0140-5

Klimaszewski, J., Morency, M.-J., Labrie, P., Seguin, A., Langor, D., Work, T., Bourdon, C., Thiffault,

E., Pare, D., Newton, A., & Thayer, M. (2013). Molecular and microscopic analysis of the gut

contents of abundant rove beetle species (Coleoptera, Staphylinidae) in the boreal balsam fir

forest of Quebec, Canada. *ZooKeys*, *353*, 1–24. https://doi.org/10.3897/zookeys.353.5991

Kolde, R. (2019). *pheatmap: Pretty Heatmaps. R package* (1.0.12).

Kontsedalov, S., Zchori-Fein, E., Chiel, E., Gottlieb, Y., Inbar, M., & Ghanim, M. (2008). The presence

of Rickettsia is associated with increased susceptibility of Bemisia tabaci (Homoptera:

Aleyrodidae) to insecticides. *Pest Management Science*, *64*(8), 789–792.

https://doi.org/10.1002/ps.1595

Krassowski, M., Arts, M., & CyrilLagger. (2020). *ComplexUpset* (1.3.1).

https://doi.org/10.5281/zenodo.3700590

Küchler, S. M., Kehl, S., & Dettner, K. (2009). Characterization and localization of Rickettsia sp. in

water beetles of genus Deronectes (Coleoptera: Dytiscidae). *FEMS Microbiology Ecology*,

68(2), 201–211. https://doi.org/10.1111/j.1574-6941.2009.00665.x

Lanzoni, O., Sabaneyeva, E., Modeo, L., Castelli, M., Lebedeva, N., Verni, F., Schrallhammer, M.,

Potekhin, A., & Petroni, G. (2019). Diversity and environmental distribution of the

683 cosmopolitan endosymbiont "Candidatus Megaira". *Scientific Reports*, *9*(1), 1179.

684 https://doi.org/10.1038/s41598-018-37629-w

685 Leclair, M., Polin, S., Jousseaume, T., Simon, J.-C., Sugio, A., Morlière, S., Fukatsu, T., Tsuchida, T., &

686 Outreman, Y. (2017). Consequences of coinfection with protective symbionts on the host

687 phenotype and symbiont titres in the pea aphid system. *Insect Science*, *24*(5), 798–808.

688 https://doi.org/10.1111/1744-7917.12380

689 Lex, A., Gehlenborg, N., Strobelt, H., Vuillemot, R., & Pfister, H. (2014). UpSet: Visualization of

690 intersecting sets. *IEEE Transactions on Visualization and Computer Graphics*, *20*(12), 1983–

691 1992. https://doi.org/10.1109/TVCG.2014.2346248

692 Łukasik, P., Guo, H., van Asch, M., Ferrari, J., & Godfray, H. C. J. (2013). Protection against a fungal

693 pathogen conferred by the aphid facultative endosymbionts Rickettsia and Spiroplasma is

694 expressed in multiple host genotypes and species and is not influenced by co-infection with

695 another symbiont. *Journal of Evolutionary Biology*, *26*(12), 2654–2661.

696 https://doi.org/10.1111/jeb.12260

697 Murray, G. G. R., Weinert, L. A., Rhule, E. L., & Welch, J. J. (2016). The Phylogeny of Rickettsia Using

698 Different Evolutionary Signatures: How Tree-Like is Bacterial Evolution? *Systematic Biology*,

699 *65*(2), 265–279. https://doi.org/10.1093/sysbio/syv084

700 Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A Fast and Effective

701 Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and*

702 *Evolution*, *32*(1), 268–274. https://doi.org/10.1093/molbev/msu300

703 Oliver, K. M., Degnan, P. H., Burke, G. R., & Moran, N. A. (2010). Facultative Symbionts in Aphids

704 and the Horizontal Transfer of Ecologically Important Traits. *Annual Review of Entomology*,

705 *55*(1), 247–266. https://doi.org/10.1146/annurev-ento-112408-085305

706 Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015). CheckM:

707 assessing the quality of microbial genomes recovered from isolates, single cells, and

708 metagenomes. *Genome Research*, *25*(7), 1043–1055. https://doi.org/10.1101/gr.186072.114

709 Perlman, S. J., Hunter, M. S., & Zchori-Fein, E. (2006). The emerging diversity of Rickettsia.

710 *Proceedings of the Royal Society B: Biological Sciences*, *273*(1598), 2097–2106.

711 https://doi.org/10.1098/rspb.2006.3541

Pilgrim, J., Ander, M., Garros, C., Baylis, M., Hurst, G. D. D., & Siozios, S. (2017). Torix group

Rickettsia are widespread in Culicoides biting midges (Diptera: Ceratopogonidae), reach high

frequency and carry unique genomic features. *Environmental Microbiology*, *19*(10), 4238–

4255. https://doi.org/10.1111/1462-2920.13887

Pilgrim, J., Thongprem, P., Davison, H. R., Siozios, S., Baylis, M., Zakharov, E. V., Ratnasingham, S.,

deWaard, J. R., Macadam, C. R., Smith, M. A., & Hurst, G. D. D. (2021). Torix Rickettsia are

widespread in arthropods and reflect a neglected symbiosis. *GigaScience*, *10*(3), 1–19.

https://doi.org/10.1093/gigascience/giab021

Pritchard, L., Glover, R. H., Humphris, S., Elphinstone, J. G., & Toth, I. K. (2016). Genomics and

taxonomy in diagnostics for food security: soft-rotting enterobacterial plant pathogens.

Analytical Methods, *8*(1), 12–24. https://doi.org/10.1039/C5AY02550H

R Core Team. (2020). *R: A Language and Environment for Statistical Computing* (3.4.4). R

Foundation for Statistical Computing. https://www.r-project.org/

Rodriguez-R, L. M., Jain, C., Conrad, R. E., Aluru, S., & Konstantinidis, K. T. (2021). Reply to: "Re-

evaluating the evidence for a universal genetic boundary among microbial species." *Nature*

Communications, *12*(1), 4060. https://doi.org/10.1038/s41467-021-24129-1

Schrallhammer, M., Ferrantini, F., Vannini, C., Galati, S., Schweikert, M., Görtz, H.-D., Verni, F., &

Petroni, G. (2013). 'Candidatus Megaira polyxenophila' gen. nov., sp. nov.: Considerations on

Evolutionary History, Host Range and Shift of Early Divergent Rickettsiae. *PLoS ONE*, *8*(8),

e72581. https://doi.org/10.1371/journal.pone.0072581

Seemann, T. (2014). Prokka: rapid prokaryotic genome annotation. *Bioinformatics*, *30*(14), 2068–

2069. https://doi.org/10.1093/bioinformatics/btu153

Siozios, S. (2021). *SioStef/panplots: A small R script for generating pangenome accumulation curves*.

https://github.com/SioStef/panplots

Thongprem, P., Evison, S. E. F., Hurst, G. D. D., & Otti, O. (2020). Transmission, Tropism, and

Biological Impacts of Torix Rickettsia in the Common Bed Bug Cimex lectularius (Hemiptera:

Cimicidae). *Frontiers in Microbiology*, *11*. https://doi.org/10.3389/fmicb.2020.608763

Tvedte, E. S., Walden, K. K. O., McElroy, K. E., Werren, J. H., Forbes, A. A., Hood, G. R., Logsdon, J.

M., Feder, J. L., & Robertson, H. M. (2019). Genome of the Parasitoid Wasp Diachasma

alloeum, an Emerging Model for Ecological Speciation and Transitions to Asexual

Reproduction. *Genome Biology and Evolution*, *11*(10), 2767–2773.

https://doi.org/10.1093/gbe/evz205

van der Beek, S. L., Zorzoli, A., Çanak, E., Chapman, R. N., Lucas, K., Meyer, B. H., Evangelopoulos,

D., de Carvalho, L. P. S., Boons, G., Dorfmueller, H. C., & van Sorge, N. M. (2019). Streptococcal

dTDP-L-rhamnose biosynthesis enzymes: functional characterization and lead compound

identification. *Molecular Microbiology*, *111*(4), 951–964. https://doi.org/10.1111/mmi.14197

Weinert, L. A., Araujo-Jnr, E. V., Ahmed, M. Z., & Welch, J. J. (2015). The incidence of bacterial

endosymbionts in terrestrial arthropods. *Proceedings of the Royal Society B: Biological*

Sciences, *282*(1807), 20150249. https://doi.org/10.1098/rspb.2015.0249

Weinert, L. A., Werren, J. H., Aebi, A., Stone, G. N., & Jiggins, F. M. (2009). Evolution and diversity of

Rickettsiabacteria. *BMC Biology*, *7*(1), 6. https://doi.org/10.1186/1741-7007-7-6

Werren, J. H., Hurst, G. D., Zhang, W., Breeuwer, J. A., Stouthamer, R., & Majerus, M. E. (1994).

Rickettsial relative associated with male killing in the ladybird beetle (Adalia bipunctata).

Journal of Bacteriology, *176*(2), 388–394. https://doi.org/10.1128/jb.176.2.388-394.1994

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

https://ggplot2.tidyverse.org

Zchori-Fein, E., Borad, C., & Harari, A. R. (2006). Oogenesis in the date stone beetle, Coccotrypes

dactyliperda, depends on symbiotic bacteria. *Physiological Entomology*, *31*(2), 164–169.

https://doi.org/10.1111/j.1365-3032.2006.00504.x