Specific sequence of arrival promotes coexistence via spatial niche preemption by the

weak competitor

Inês Fragata[1], Raul Costa-Pereira[2], Mariya Kozak[1], Agnieszka Majer[3], Oscar Godoy[4*], Sara

Magalhães[1*]

1 – Centre for Ecology, Evolution and Environmental Changes; Department of Animal

Biology, Faculty of Sciences, University of Lisbon, Portugal

2 – Department of Animal Biology, Institute of Biology, Universidade Estadual de Campinas,

Brazil ORCID 0000-0003-2370-5866

3 – Population Ecology Lab, Faculty of Biology, Adam Mickiewicz University, Poland ORCID

0000-0002-2425-7885

4 – Department of Biology, Instituto Universitario de Investigación Marina (INMAR),

Universidad de Cádiz, Puerto Real. 11510, Spain. ORCID 0000-0003-4988-6626

*Co-last authorship

Running title: Niche preemption facilitates coexistence

Keywords: Order of arrival, Priority effects, Modern coexistence theory, Spatial segregation,

Tetranychus urticae and *T. evansi*, Herbivores, Niche modification, Competitive ability

Article type: Letter

Number of words in abstract: 150

Number of words main text: 5000

Number of references: 66

Number of figures: 3

Corresponding Author: Inês Fragata, irfragata@fc.ul.pt, cE3c, Centre for Ecology, Evolution and Environmental changes, Faculdade de Ciências da Universidade de Lisboa, Edifício C2, Campo Grande 1749-016 Lisbon, Portugal, Phone. +351 217500577, ext. 22311.

Authorship statement: IF, OG and SM designed the experiment, IF, AM, MK and RCP collected data, IF and OG performed modelling work and analysed data. IF and SM wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

Data accessibility statement: Data and scripts for data analyses will be deposited in a public repository upon acceptance.

Abstract

Historical contingency, such as the order of species arrival, can modify competitive outcomes

via niche modification or preemption. However, how these mechanisms ultimately modify

stabilising niche and average fitness differences remains largely unknown. By experimentally

assembling two congeneric spider mite species feeding on tomato plants during two

generations, we show that order of arrival affects species' competitive ability and changes the

outcome of competition. Contrary to expectations, order of arrival did not cause positive

frequency dependent priority effects. Instead, coexistence was predicted when the inferior

competitor (*Tetranychus urticae*) arrived first. In that case, *T. urticae* colonised the preferred

feeding stratum (leaves) of *T. evansi* leading to spatial niche preemption, which equalised

fitness and reduced niche differences, driving community assembly to a close-to-neutrality

scenario. Our study demonstrates how the order of species arrival and the spatial context of

competitive interactions can jointly determine whether species can coexist.

Introduction

Priority effects are broadly defined as the process by which historical contingencies in community assembly (e.g. order and/or timing of arrival) change the outcome of interspecific interactions (Chase 2003; Fukami 2015). Inhibitory priority effects, when earlier arrival by one species inhibits the growth of the species arriving next, are expected to result in alternative stable states hampering coexistence (Chase 2003; Fukami 2015; Ke & Letten 2018). In turn, facilitative priority effects, when population growth is higher if individuals arrive after the settlement of a first species, do not always promote coexistence. Rather, the outcome depends on the interaction strength among species and on the environmental context in which they interact (Bulleri *et al.* 2016; Bimler *et al.* 2018). These effects have been less often observed in natural communities (Queijeiro-Bolaños *et al.* 2017; Clay *et al.* 2019; Halliday *et al.* 2020).

Two major mechanisms are predicted to cause priority effects: niche preemption, in which early colonisers reduce the amount of resource available to late colonisers, and niche modification, in which the species arriving first modifies the environment, thereby inhibiting or facilitating later colonisation (Kardol *et al.* 2013; Vannette & Fukami 2014; Fukami 2015; Delory *et al.* 2019, 2021; Grainger *et al.* 2019). Niche preemption in plant communities was found to be strong in environments with high nutrient supply, as early arriving plants grew quickly and prevented growth of later colonisers by depleting space and light (Kardol *et al.* 2013). Niche modification was also detected in plants, as early colonisations modified the soil metabolome and inhibited population growth of forb, but not grass species arriving later (Delory *et al.* 2021). Although distinguishing among niche preemption and modification is not always possible (Grainger *et al.* 2018; Boyle *et al.* 2021), recent advances in coexistence theory can serve as a powerful approach to better understand the importance of historical contingencies for species coexistence. Yet the combination of these theoretical tools has seldom been applied in empirical settings.

Modern coexistence theory posits that the long-term persistence of competing species (i.e.,

species coexistence) can be attained by two non-mutually exclusive mechanisms: (i) equalising

mechanisms that reduce average fitness differences, and therefore, dominance between species

and (ii) stabilising mechanisms, which stabilise the interaction between competitors by

increasing the strength of intraspecific competition relative to interspecific competition

(Chesson 2000). Therefore, species will stably coexist if stabilising niche differences are larger

than differences in fitness between competitors. Otherwise the species with higher fitness will

eventually dominate the community (Chesson 2000; Barabás *et al.* 2018; Spaak & De Laender

2021). Under this framework, priority effects are strictly defined as positive frequency

dependence (i.e., via negative niche differences), leading to the dominance of the early-arriving

species (Ke & Letten 2018; Grainger *et al.* 2019; Spaak & De Laender 2021). Hence, species

cannot coexist unless there is spatial variability in the order of arrival. Although recent theory

offers predictions on the outcome of coexistence in systems with historical contingencies,

empirical tests are conspicuously lacking (but see Cardinaux *et al.* 2018; Grainger *et al.* 2019;

Song *et al.* 2020). Therefore, there is as yet scarce knowledge of which species traits interact

with historical contingencies to determine outcomes of interspecific interactions.

For herbivore communities, habitat use and dispersal capacity can affect resource use and

ultimately the spatial distribution of consumers. This may lead to niche preemption, as

herbivores generally have preferred plant strata and the first arriving species may monopolise

that resource (Grainger *et al.* 2018; Godinho *et al.* 2020a). Moreover, herbivores often induce

defences on the plants they colonise, which is expected to entail niche modification for species

arriving later (Erb *et al.* 2011; Moreira *et al.* 2015; Stam *et al.* 2017). For example, Hougen-

Eitzman & Karban (1995) showed that early colonisation of grape vine leaves by Willamette

mites negatively affected the growth of Pacific mites, probably due to systemic induction of

defences. Other herbivore species can instead down-regulate plant defences, improving the

performance of later colonisers (Sarmento *et al.* 2011a; Godinho *et al.* 2016), thereby

potentially causing facilitative priority effects. Overall, given the environmental heterogeneity

that herbivores experience (e.g., variation in leaf quality within and between plants), effects of

the order of arrival on species coexistence are expected to be prevalent in these systems

(Utsumi *et al.* 2010; Erb *et al.* 2011; Moreira *et al.* 2015; Stam *et al.* 2017, 2018; Godinho *et*

al. 2020a). Still, what type of competitive outcome we should expect is unclear. Indeed,

although the order of arrival is linked to priority effects, the interaction between the chronology

of community assembly and the impact of species on the environment (e.g. where they growth

and how they modify the habitat) can result in diverse outcomes, from competitive exclusion

to species coexistence. Applying modern coexistence theory to this open question can shed

light on the proximate mechanisms that allow for species to coexist under varied historical

contingencies.

Here, we investigate the drivers of competitive outcomes by combining theoretical and

empirical tools to test the mechanisms through which order of arrival affects species

coexistence. We use as a model system the two closely-related competing herbivorous species,

the spider mites *Tetranychus urticae* and *T. evansi*. *Tetranychus evansi* generally outcompetes

T. urticae on tomato plants (Sarmento *et al.* 2011b; Orsucci *et al.* 2017; Alzate *et al.* 2020),

although both species are also commonly observed on the same location (Ferragut *et al.* 2013).

Niche modification is expected to be at play in this system, because the two species interact

with plant defences. Indeed, *T. evansi* suppresses plant defences (Sarmento *et al.* 2011a; Alba

et al. 2014), whereas most *T. urticae* populations induce them (Kant *et al.* 2008). This

asymmetrical niche modification is predicted to increase the probability of coexistence by

hampering growth of the stronger competitor and favouring growth of the inferior one, when

the later arrives on plants colonised by the other species. Moreover, niche preemption may

occur, as both *T. evansi* and *T. urticae* prefer the upper, more nutritious leaves of tomato plants,

where their performance is higher (Godinho *et al.* 2020a). Thus, early-arriving species could

occupy the preferred niche and displace the other species to lower, less optimal, plant strata.

We tested this by performing a series of multi-generational experiments where we varied order

of arrival and measured space use by the two competing species. To quantify the magnitude of

niche modification, we tested how these species modify the expression of genes associated

with induced defenses on tomato. We then applied modern coexistence theory to unravel the

conditions favouring coexistence or potentially leading to priority effects.

Material and Methods

Model system, species characteristics, and maintenance of experimental populations

Tetranychus urticae is a generalist herbivore that feeds on many economically important crops

(Helle & Sabelis 1985; Grbić *et al.* 2011; Sousa *et al.* 2019), whereas *T. evansi* is a solanaceous

specialist that has recently invaded Europe (Boubou *et al.* 2012). Both species colonise tomato

plants, although *T. urticae* may shift to other hosts if *T. evansi* is present (Ferragut *et al.* 2013).

All experiments were performed with outbred populations of *T. urticae* and *T. evansi*

spider mites, formed via controlled crosses among four *T. evansi* and three *T. urticae*

populations collected in different locations in Portugal (Godinho *et al.* 2020b). Populations

were maintained in boxes containing leaves detached from five-week-old tomato plants

(*Solanum lycopersicum*, *var* MoneyMaker), with their petiole in a small pot containing water

Twice a week, overexploited leaves were removed, and water and new tomato leaves were

added. Before infestation, tomato plants were kept in a separate climatic chamber and watered

three times per week. Mites and plants were kept under controlled conditions (25 ºC, 70%

humidity, 16 /8 L/D hours).

We created same-age cohorts of mated *T. urticae* and *T. evansi* females for each block. To

this aim, females were placed during 48h in petri dishes (14.5 cm diameter, with a layer of wet

cotton watered twice per week) and two freshly cut tomato leaves. One week later, another

tomato leaf was added. In the experiment, we used females with 13-15 days of age.

Theoretical approach for predicting competitive outcomes: quantifying niche and fitness

differences

Data collected in the experiments were used to parameterise a mathematical model from which

niche and average fitness differences can be quantified to then draw predictions of competitive

outcomes. We assume that the population dynamics in our experiment can be described by a

Beverton-Holt function (Levine & HilleRisLambers 2009; Godoy & Levine 2014):

$$(1)\ N_{i,t+1} = \frac{\lambda_i * N_{i,t}}{(1 + \alpha_{ii} * N_{i,t} + \alpha_{ij} * N_{j,t})}$$

Where $N_{i,t+1}$ is the number of individuals of species i in the next generation, λ_i the intrinsic

growth rate of species i in absence of competitors, α_{ii} the intraspecific competitive interaction

describing the per-capita effect of species i on itself, α_{ij} the interspecific competitive

interactions describing the per-capita effect of species j on species i, and $N_{i,t}$, $N_{j,t}$ the number of

individuals of species i and j in the current generation, respectively. We assume that spider

mites do not have a dormant stage. Thus, λ_i represents the fraction of eggs that hatch and

become females that reproduce in the next generation. One of the predictions of modern

coexistence theory is that, for species to coexist, they must invade the resident species from

rare. Because for our system equilibrium densities are difficult to attain within a time frame

fast enough to study the impact of priority effects on species coexistence, we instead used

experimental gradients of density and relative frequency to estimate intra and interspecific

competitive interactions (the α's) and intrinsic growth rate (λ) for each species, an approach

well established and validated by previous work (Godoy & Levine 2014; Matías *et al.* 2018;

Song *et al.* 2020).

From the above mentioned model, niche overlap (ρ) is defined as follows (see details in

Chesson 2012; Godoy & Levine 2014).

187
$$(2) \quad \rho = \sqrt{\frac{\alpha_{ij}}{\alpha_{jj}} \frac{\alpha_{ji}}{\alpha_{ii}}}$$

This formula reflects the average degree to which species limit individuals of their own

species relative to heterospecific competitors. If species limit population growth of their own

species more strongly than that of their competitors (α_{jj}, α_{ii}, are much greater than α_{ij}, α_{ji}),

then niche overlap will be low, favouring coexistence. Alternatively, niche overlap will

approach one, which hampers stable coexistence. Stabilising niche differences are thus

expressed as 1-ρ.

Average fitness differences ($\frac{\kappa_j}{\kappa_i}$) (Chesson 2012; Godoy & Levine 2014) are defined as:

195
$$(3) \quad \frac{\kappa_j}{\kappa_i} = \frac{\lambda_j - 1}{\lambda_i - 1} * \frac{\sqrt{\alpha_{ij} * \alpha_{ii}}}{\sqrt{\alpha_{ji} * \alpha_{jj}}}$$

196   The greater the ratio, ($\frac{\kappa_j}{\kappa_i}$), the greater the fitness advantage of species j over i. If this ratio

197   is one, species are equivalent competitors. Coexistence requires both species to invade when

198   rare (Chesson 2012), which is satisfied when (Godoy & Levine 2014):

199
$$(4) \quad \rho < \frac{\kappa_j}{\kappa_i} < \frac{1}{\rho}$$

200
Stable coexistence is possible whenever species have either large stabilising niche

differences (corresponding to small niche overlap) that overcome large average fitness

differences, or at the other extreme, via an a close-to-neutral scenario (Scheffer *et al.* 2018),

where, even with weak niche differences, small fitness differences stabilise the interaction

between competitors. If no coexistence is predicted, we can pinpoint if this is due to

competitive exclusion (when fitness differences are larger than niche differences) or to priority

effects, leading to alternative states when niche differences are negative. Negative niche

differences imply that each species limits the growth of the competitor more than their own

(Fukami & Nakajima 2011; Ke & Letten 2018).

We used maximum likelihood techniques to parameterise the population model following

a nested approach. That is, we first created a single model for which we estimate the intrinsic

growth rate in absence of competitors (λ), and then we used this information as prior for

subsequent more complex models that include intra and interspecific competitive interactions

(the α's) (Matías et al. 2018). λ values were considered fixed per species across empirical

treatments, but competition varied across treatments because mite species can differentially

disperse and modify leaf quality and availability (see the full details in the Supplementary

Material and Methods).

Experiments

To test the impact of order of arrival on coexistence, we performed a series of experiments in

which we either manipulated the order of arrival and relative frequency (i.e., relative initial

abundance with a constant density of 20 individuals), or the initial density of each of two

species of competing spider mites. Furthermore, to estimate the effect of order of arrival on

promoting niche preemption, we quantified leaf occupation for both species at the end of the

experiment. Finally, to estimate the effect of order of arrival on promoting niche modification,

we quantified induction of plant defences of both species.

In the first experiment, both species were introduced simultaneously using the following

proportions of *T. evansi* : *T. urticae*: 1:19; 10:10 and 19:1, along with the single-species

controls (20:0 and 0:20). To manipulate the order of arrival, we introduced (i)10 *T. evansi*

females 48h before 10 *T. urticae* females and vice versa and (ii) 19 *T. evansi* females 48h

before 1 *T. urticae* female and vice versa (Figure S1). The experiment was done in two blocks,

one week apart. Each block contained five boxes of each experimental treatment (nine

treatments, n=10), each with a pot filled with water and two freshly cut tomato leaves from

five-week-old tomato plants. Leaf pairs consisted of leaves 2 and 4 or 3 and 5 (leaf number is

inversely proportional to leaf age), to ensure that each box contained a younger and an older

leaf, since both species prefer younger leaves (Godinho *et al.* 2020a). Adult females were

distributed by the two leaves, following the treatments described above. After one generation

(circa 14 days), two more leaves were added to ensure enough resources for the second mite

generation. Boxes that initially received the leaf pair 2-4, received leaves 3-5 and vice versa.

After two generations, we counted the number of adult females of each species on each leaf.

Next, we estimated the growth rate of each species by counting the number of adult

females obtained from the progeny of a single *T. urticae* or *T. evansi* female ovipositing for

48h in two overlapping 18mm leaf disks (n=10). These disks were placed in square petri dishes

with a layer of wet cotton and were watered every two days. The number of adult females

produced was assessed after one generation.

<u>Quantification of niche modification</u>

To quantify the magnitude of niche modification induced by *T.urticae* and *T.evansi*, we

investigated how these two species modified the expression of genes associated with plant

defences. As controls, we quantified the expression of the same genes upon infestation with

spider mites from *T. urticae* Santpoort and *T.evansi* Viçosa populations, known to induce and

suppress tomato defences, respectively (Alba *et al.* 2014). Details of this experiment are given

in the Supplementary Material and Methods and Table S1.

Data Analyses

<u>Effect of order of arrival and initial frequency on species abundance</u>

To test the impact of order of arrival, frequency and their interaction on the proportion of adult

females of each species after two generations, we performed the following general linear mixed

model (lme4 package, Bates et al. 2015), using the binomial family:

260   (5) Y= Treatment + Block + ε

Where Y corresponds to the combination of two vectors with the number of *T. evansi* and

T. urticae females after two generations, Treatment (fixed factor) to the combination of

different orders of arrival and initial frequencies, Block (random factor) to whether the

experiment was performed on week one or two, and ε to the residual error. We then performed

a priori contrasts, using testInteractions from phia package (Rosario-Martinez 2015) as our

experimental design was not orthogonal. To compare the effect of different orders of arrival,

we performed contrasts between the treatments with same initial frequency but different orders

of arrival. To compare the effect of frequency, we performed contrasts between treatments with

same order of arrival but different initial frequencies. Contrasts were corrected for multiple

comparisons using FDR correction (Benjamini & Yekutieli 2001). To test whether the results

were biased by the order in which the leaf pairs were added to the boxes, we repeated these

statistical analyses separately for each leaf pair.

Effect of order or arrival and initial frequency on leaf occupancy and aggregation

To test if coexistence outcomes could be explained by niche preemption, we compared

occupancy patterns of each species across the four leaves. For the single species treatment, we

tested if the number of females differed across leaves (model 6). For the double species

treatment, we tested if the order of arrival, initial frequency, or their interaction changed mite

distribution (model 7), by comparing it to the distribution of the single species treatment.

We applied the following binomial models, with Leaf and/or Treatment and their

interaction as fixed factors, for the control (model 6) and experimental (model 7) treatments:

282   (6) Y= Leaf + ε

283   (7) Y= Leaf + Treatment+ Leaf x Treatment + ε

where Y corresponds to the combination of two vectors with the number of *T. evansi* (or

T. urticae) females on each leaf and the total number of individuals on each box that were not

on that leaf. To test whether the results were biased by the order in which the leaf pairs were

added to the boxes, we repeated these statistical analyses accounting for the preference of each

species for each leaf pair. For the double treatment, a posteriori contrasts were done between

each treatment and the corresponding single species treatment. The initial fitting with Block as

a random factor, indicated no variance in this factor, thus we fitted only fixed factors. We also

tested in changes in order of arrival affected aggregation scores (see details in Supplementary

Material and Methods).

All analyses were done using R (R Core Team 2021). To predict coexistence outcomes we

used the package "cxr" (García-Callejas *et al.* 2020). Plots were done using "ggplot2"

(Wickham 2016) and "cowplot" (Wilke 2020) packages. Data and scripts are available in the

github repository: https://github.com/irfragata/order_arrival_niche_preemption.

Results

Effect of order of arrival and initial frequency on species abundance

The number of individuals of each species on tomato plants were affected by the order of arrival

(contrasts between *T. evansi* arriving first vs. simultaneously: χ^2 = 44.252, df = 1, p-value <

0.0001; or *T. urticae* arriving first vs. simultaneously: χ^2 = 375.860, df = 1, p-value < 0.0001),

and their initial frequency (contrasts between *T. evansi* starting at equal *vs.* higher frequency.

: χ^2 = 784.335, df = 1, p-value < 0.0001; or *T. urticae* starting at equal *vs.* higher frequency*:* χ^2

= 654.903, df = 1, p-value < 0.0001). Specifically, the abundance of *T. evansi* females after

two generations was higher when this species arrived first or simultaneously with *T. urticae*,

independently of initial frequencies. However, the additional advantage provided by arriving

first was much larger in the equal frequency treatment (Table S2, Fig.1). The abundance of *T.*

urticae after two generations was also affected by initial frequency and order of arrival. Indeed,

the final number of *T. urticae* females was higher when this species arrived first and was at

high initial frequency, than in the equal frequency treatment (Table S2, Fig. 1). We observed

the same patterns when performing these analyses per leaf pair (Table S3). Overall, these

results confirm that *T. evansi* is a superior competitor as observed in previous studies

(Sarmento *et al.* 2011b; Ferragut *et al.* 2013; Alzate *et al.* 2020).

Effect of order of arrival on coexistence

The order of arrival modified the outcome of competition between the two species. *Tetranychus*

evansi (the superior competitor) is predicted to exclude *T. urticae* when it arrives first or at the

same time. Under this exclusion scenario, the rate of competitive exclusion is expected to be

faster when *T. evansi* arrives first due to a decrease in niche differences (Fig 2). The small

overlap between the lower confidence interval with the priority effects region suggests that

positive frequency dependence might also emerge in this system. Interestingly, coexistence

was only possible when *T. urticae* arrived first (Fig. 2). This outcome was due to small niche

and fitness differences among competitors, leading to a quasi-neutral scenario. Specifically,

when *T. urticae* arrived first, we observed similar strengths of intra- and interspecific

interactions among species (Fig S2A). Contrary to expectations and previous studies, the order

of arrival was not associated with positive frequency dependence leading to priority effects.

However, since the order of arrival modified the outcome of the interactions between *T. urticae*

and *T. evansi*, we can also interpret these results as priority effects (*sensu* (Chase 2003; Fukami

2015) allowing for coexistence between species in our system.

Effect of order of arrival and initial frequency on leaf occupancy and aggregation

When *T. evansi* was alone, it reached higher abundances on leaves 3 and 4 (Table S4A, Fig

3B), whereas *T. urticae* was less abundant on leaf 2 in comparison to all others (Table S4A,

Fig 3D). Fewer *T. evansi* females were found on leaf 4 when *T. urticae* arrived first, and on

leaf 3 when *T. urticae* started with higher frequency and both species arrived at the same time

(Fig 3, Fig S3A, Table S4B). When *T. evansi* arrived first or started at higher frequency, we

observed fewer changes on its own leaf occupancy (Fig S4A). The distribution of *T. urticae*

showed a slight shift when it arrived first, with a reduction on the prevalence of leaves 2 and 5

and slightly higher occupation of leaves 3 and 4 (Fig. S3B, Table S4B). When *T. evansi* started

at high frequency, there was also a shift in *T. urticae* distribution, with a lower occupancy of

leaves 2 and 5 (Fig S3B). We observed similar shifts in leaf occupation when performing the

analyses accounting for the order in which each leaf pair was added (Fig S4, Table S5).

Spatial aggregation significantly differed among treatments ($\chi^2 = 18.186$, df = 6, p-value

= 0.01279), being higher in treatments with similar initial densities (cf. Fig S5 with Fig 1, Table

S5). We observed a significant difference in C-score, with higher aggregation when both

species arrived at the same time and had equal frequency, and a lower aggregation when both

species arrived at the same time and *T. evansi* started at higher frequency (Table S6). Order of

arrival did not change the C-score (Fig. S5, Table S6).

Quantification of niche modification

353   Plants infested by *T. urticae* or *T. evansi* populations showed patterns of gene expression

354   similar to those of Viçosa, the suppression control, and significant differences with Santpoort,

355   the induction control (Fig. S6; Table S7). We thus conclude that, both populations suppress

356   plant defences.

Discussion

This study shows that order of arrival interacts with competitive ability to determine the probability of coexistence between congeneric species that share common resources. When both species arrive at the same time or the superior competitor (*Tetranychus evansi*) arrived first, *T. urticae* was predicted to be excluded. Coexistence was only predicted when the inferior competitor (*T. urticae*) was the first species colonising the habitat. Analyses of leaf occupancy show that these competition outcomes are linked to a spatial niche preemption process in which *T. evansi* was displaced from its preferred food stratum when *T. urticae* arrived first. As a result of this complex interaction between order of arrival, species competitive ability, and spatial occupancy, we observed a particular configuration that allows species coexistence: both species equalised their fitness differences to the extent that they can coexist despite small niche differences. These multiple lines of evidence challenge the common understanding of the inhibitory role of niche preemption in coexistence between species.

We found that *T. evansi* had higher competitive ability and growth rate, and often excluded *T. urticae* (Fig 1, 2). This is in line with laboratory observations showing that *T. evansi* outcompeted *T. urticae* on tomato plants (Sarmento *et al.* 2011b; Alzate *et al.* 2020, but see Orsucci *et al.* 2017) and with field observations showing a shift in host use in *T. urticae* upon invasion by *T. evansi* (Ferragut et al. 2013). Still, these two species can co-occur in the field in the same plant species (Ferragut *et al.* 2013; Orsucci *et al.* 2017; Zélé *et al.* 2018). The advantage created by the earlier arrival of *T. urticae*, and associated reduction in interspecific competition by *T. evansi*, could be one of the possible mechanisms fostering their coexistence. Indeed, *T. urticae* can withstand colder temperatures than *T. evansi* (Gotoh *et al.* 2010; Khodayari *et al.* 2013; Riahi *et al.* 2013; White *et al.* 2018), hence it is expected to arrive first

in the season. Field surveys that sample both species in the same location across seasons are

needed to further explore this hypothesis.

Historical contingencies emerging from order of arrival can happen through two main

mechanisms: niche modification or niche preemption (Fukami 2015). In our system, niche

modification may arise via interactions between spider mites and plant defences. However, we

observe that both species suppress plant defenses. If suppression would affect species

performance, we would expect higher production of offspring when the competitor arrives first.

We did not observe this, suggesting that this mechanism of niche modification does not affect

the outcome of competition in this system.

Niche preemption can occur through monopolisation of nutrients or space, which can be

particularly important among competitors with similar requirements (Grainger *et al.* 2018;

Holditch & Smith 2020). In our study, we observed a shift in the leaf occupancy pattern of *T.*

evansi females when *T. urticae* arrived first. This displacement of *T. evansi* from the preferred

food stratum (i.e., younger, more nutritious leaves) by early-arriving *T. urticae* can explain the

decreased performance of the superior competitor. Thus, our results indicate that variation in

species performance driven by habitat quality heterogeneity (Orians *et al.* 2000; Orians & Jones

2001) combines with order of arrival to generate niche preemption, providing a mechanism for

the two herbivores to coexist.

Order of arrival is a major determinant of community assembly across diverse taxa, from

microbes to plants (Chase 2003; Erb *et al.* 2011; Kardol *et al.* 2013; Stam *et al.* 2017; Grainger

et al. 2018, 2019; Clay *et al.* 2019, 2020; Halliday *et al.* 2020). Most of these studies show that

early colonisers inhibit growth and decrease performance of late arriving species, especially in

those that occupy very similar niches (Fargione *et al.* 2003; Vannette & Fukami 2014; Delory

et al. 2019, 2021; Grainger *et al.* 2019), although very few concern herbivorous species

competing for the same niche (e.g. Grainger *et al.* 2018; Holditch & Smith 2020). Other studies

found that order of arrival does not affect community assembly (e.g. Delory *et al.* 2021) or that

initial colonisers facilitate later colonisation of other species (e.g. Queijeiro-Bolaños *et al.*

2017; Delory *et al.* 2019). Here, we show that coexistence is promoted by niche preemption

because early colonisation by the inferior competitor leads to increased intraspecific

competition for the superior competitor and reduced interspecific competition for itself. . As a

result, both species can coexist under a quasi-neutral scenario because this equalising effect on

fitness differences is enough to fit within the constraints of small niche differences. Our study

adds a novel perspective to the growing body of evidence that historical contingencies shape

ecological communities, by showing that the probability of coexistence of two competing

herbivores changes due to an interaction between order of arrival and species competitive

ability.

Priority effects were recently incorporated into modern coexistence theory (Ke & Letten

2018; Spaak & De Laender 2021), but empirical tests quantifying the effects of order of arrival

on species coexistence remain very rare. In another study, Grainger *et al.* (2019) documented

that positive frequency dependence, due to strong priority effects, arose from changes in order

or arrival in yeast species feeding on floral nectars. In contrast, our results show that order of

arrival did not lead to alternative states caused by priority effects under positive frequency

dependence. Rather, we predicted either competitive exclusion when *T. evansi* arrived first

because it excluded *T. urticae* or coexistence when *T. urticae* arrived first. Overall, these results

suggest that in this system deterministic expectations, stemming from theory, can be strongly

influenced by small stochastic events, such as changes in order of arrival, because it affects the

timing of dispersal across and within host plants.

Framing priority effects in the modern coexistence theory (Ke & Letten 2018) is

undoubtedly an important step to mechanistically understand how order of arrival affects

community assembly processes. However, in this framework, priority effects are only caused

by positive frequency dependence (i.e., population growth rate is higher as individuals become

relatively more abundant) (Fukami 2015; Song *et al.* 2020). Including other types of

interactions and outcomes into modern coexistence framework is fundamental to improve our

ability to understand how species coexist (Spaak *et al.* 2021). Here we show that order of arrival

can lead to coexistence via niche preemption by the inferior competitor. Thus, our results show

that changes in the order of arrival can produce a wide range of competitive outcomes from

coexistence to competitive exclusion due to positive and negative frequency dependence.

Therefore, it is urgent that ecologists widen the scope of the multiple outcomes that historical

contingency can produce on species coexistence.

Most empirical and theoretical studies emphasize the inhibitory nature of niche preemption

(Fargione *et al.* 2003; Fukami 2015; Vieira *et al.* 2018; Delory *et al.* 2019), with the early

arriving species outcompeting the other. However, recent theory suggests that, in a resource

competition model of two species, niche preemption by the inferior competitor could facilitate

coexistence under a trade-off between order of arrival and the resource levels of zero net growth

(R^*) (Qi et al. 2021). Our study is, to the best of our knowledge, the first empirical study

showing that niche preemption by the weaker competitor promotes coexistence. This striking

change in the outcome of competitive interactions emerges mostly due to a decrease in niche

overlap, shifting niche differences from negative to positive. This suggests that even small

differences in order of arrival can be sufficient for the monopolisation of a resources in plant-

herbivore interactions, which may suffice to allow coexistence between competitor species.

Therefore, our results demonstrate how small temporal differences percolate into small spatial

heterogeneities, fostering coexistence and the maintenance of diversity.

Acknowledgments: This work was financed by an ERC (European Research Council)

consolidator grant COMPCON, GA 725419 attributed to SM and by FCT (Fundação para

Ciência e Tecnologia) with the Junior researcher contract (CEECIND/02616/2018) attributed

to IF. RC-P is supported by grant #2020/11953-2 São Paulo Research Foundation (FAPESP)

and grant R-2011-37572 Instituto Serrapilheira. OG acknowledges financial support provided

by the Spanish Ministry of Economy and Competitiveness (MINECO) and by the European

Social Fund through the Ramón y Cajal Program (RYC-2017-23666). AM was funded by

National Science Centre, Poland (grant no. 2018/28/T/NZ8/00060) and Excellence Initiative -

Research University programme (support for the internationalization of the Adam Mickiewicz

University PhD students, no. 003/13/UAM/0018). The authors acknowledge stimulating

discussion with all members of the Mite Squad, in particular Flore Zélé and Diogo Godinho,

which have significantly improved the experimental design and interpretation of the results,

and Marta Artal for invaluable infrastructure at meetings in Sevilla.

Competing interests: Authors declare no competing interests.

Bibliography

Alba, J.M., Schimmel, B.C.J., Glas, J.J., Ataide, L.M.S., Pappas, M.L., Villarroel, C.A., *et al.*

(2014). Spider mites suppress tomato defenses downstream of jasmonate and salicylate

independently of hormonal crosstalk. *New Phytol.*, 205, 828–840.

Alzate, A., Onstein, R.E., Etienne, R.S. & Bonte, D. (2020). The role of preadaptation,

propagule pressure and competition in the colonization of new habitats. *Oikos*, 129,

820–829.

Barabás, G., D'Andrea, R. & Stump, S.M. (2018). Chesson's coexistence theory. *Ecol.*

Monogr., 88, 277–303.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *J. Stat. Softw.*, 67, 1–48.

Benjamini, Y. & Yekutieli, D. (2001). The control of the false discovery rate in multiple

testing under dependency. *Ann. Stat.*, 29, 1165–1188.

Bimler, M.D., Stouffer, D.B., Lai, H.R. & Mayfield, M.M. (2018). Accurate predictions of

coexistence in natural systems require the inclusion of facilitative interactions and

environmental dependency. *J. Ecol.*, 106, 1839–1852.

Boubou, A., Migeon, A., Roderick, G.K., Auger, P., Cornuet, J.M., Magalhães, S., *et al.*

(2012). Test of colonisation scenarios reveals complex invasion history of the red

tomato spider mite tetranychus evansi. *PLoS One*, 7.

Boyle, J.A., Simonsen, A.K., Frederickson, M.E. & Stinchcombe, J.R. (2021). Priority effects

alter interaction outcomes in a legume-rhizobium mutualism. *Proc. R. Soc. B Biol. Sci.*,

288, 1–8.

Bulleri, F., Bruno, J.F., Silliman, B.R. & Stachowicz, J.J. (2016). Facilitation and the niche:

Implications for coexistence, range shifts and ecosystem functioning. *Funct. Ecol.*, 30,

70–78.

Cardinaux, A., Hart, S.P. & Alexander, J.M. (2018). Do soil biota influence the outcome of

novel interactions between plant competitors? *J. Ecol.*, 106, 1853–1863.

Chase, J.M. (2003). Community assembly: When should history matter? *Oecologia*, 136,

489–498.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.*,

31, 343–366.

Chesson, P. (2012). Species Competition and Predation. In: *Encyclopedia of Sustainability*

Science and Technology (ed. Meyers, R.A.). Springer, New York, pp. 10 061–10 085.

Clay, P.A., Dhir, K., Rudolf, V.H.W. & Duffy, M.A. (2019). Within-host priority effects

systematically alter pathogen coexistence. *Am. Nat.*, 193, 187–199.

Clay, P.A., Duffy, M.A. & Rudolf, V.H.W. (2020). Within-host priority effects and epidemic

timing determine outbreak severity in co-infected populations. *Proc. R. Soc. B Biol. Sci.*,

287.

Delory, B.M., Weidlich, E.W.A., von Gillhaussen, P. & Temperton, V.M. (2019). When

history matters: The overlooked role of priority effects in grassland overyielding. *Funct.*

Ecol., 33, 2369–2380.

Delory, B.M., Schempp, H., Spachmann, S.M., Störzer, L., van Dam, N.M., Temperton,

V.M., *et al.* (2021). Soil chemical legacies trigger species-specific and context-

dependent root responses in later arriving plants. *Plant Cell Environ.*, 44, 1215–1230.

Erb, M., Robert, C.A.M., Hibbard, B.E. & Turlings, T.C.J. (2011). Sequence of arrival

determines plant-mediated interactions between herbivores. *J. Ecol.*, 99, 7–15.

Fargione, J., Brown, C.S. & Tilman, D. (2003). Community Assembly and Invasion: An

Experimental Test of Neutral versus Niche Processes. *Proc. Natl. Acad. Sci. U. S. A.*,

100, 8916–8920.

Ferragut, F., Garzón-Luque, E. & Pekas, A. (2013). The invasive spider mite Tetranychus

522  evansi (Acari: Tetranychidae) alters community composition and host-plant use of

523  native relatives. *Exp. Appl. Acarol.*, 60, 321–341.

524 Fukami, T. & Nakajima, M. (2011). Community assembly: Alternative stable states or

525  alternative transient states? *Ecol. Lett.*, 14, 973–984.

526 Fukami, T. (2015). Historical Contingency in Community Assembly: Integrating Niches,

527  Species Pools, and Priority Effects. *Annu. Rev. Ecol. Evol. Syst.*, 46, 1–23.

528 García-Callejas, D., Godoy, O. & Bartomeus, I. (2020). cxr: A toolbox for modelling species

529  coexistence in R. *Methods Ecol. Evol.*, 11, 1221–1226.

530 Godinho, D.P., Janssen, A., Dias, T., Cruz, C. & Magalhães, S. (2016). Down - regulation of

531  plant defence in a resident spider mite species and its effect upon con - and

532  heterospecifics. *Oecologia*, 180, 161–167.

533 Godinho, D.P., Janssen, A., Li, D., Cruz, C. & Magalhães, S. (2020a). The distribution of

534  herbivores between leaves matches their performance only in the absence of

535  competitors. *Ecol. Evol.*, 10, 8405–8415.

536 Godinho, D.P., Cruz, M.A., Charlery de la Masselière, M., Teodoro-Paulo, J., Eira, C.,

537  Fragata, I., *et al.* (2020b). Creating outbred and inbred populations in haplodiploids to

538  measure adaptive responses in the laboratory. *Ecol. Evol.*, 10, 7291–7305.

539 Godoy, O. & Levine, J.M. (2014). Phenology effects on invasion success: Insights from

540  coupling field experiments to coexistence theory. *Ecology*, 95, 726–736.

541 Gotoh, T., Sugimoto, N., Pallini, A., Knapp, M., Hernandez-Suarez, E., Ferragut, F., *et al.*

542  (2010). Reproductive performance of seven strains of the tomato red spider mite

543  Tetranychus evansi (Acari: Tetranychidae) at five temperatures. *Exp. Appl. Acarol.*, 52,

544  239–259.

545 Grainger, T.N., Rego, A.I. & Gilbert, B. (2018). Temperature-dependent species interactions

546  shape priority effects and the persistence of unequal competitors. *Am. Nat.*, 191, 197–

209.

Grainger, T.N., Letten, A.D., Gilbert, B. & Fukami, T. (2019). Applying modern coexistence

theory to priority effects. *Proc. Natl. Acad. Sci. U. S. A.*, 116, 6205–6210.

Grbić, M., Van Leeuwen, T., Clark, R.M., Rombauts, S., Rouzé, P., Grbić, V., *et al.* (2011).

The genome of Tetranychus urticae reveals herbivorous pest adaptations. *Nature*, 479,

487–492.

Halliday, F.W., Penczykowski, R.M., Barrès, B., Eck, J.L., Numminen, E. & Laine, A.L.

(2020). Facilitative priority effects drive parasite assembly under coinfection. *Nat. Ecol.*

Evol., 4, 1510–1521.

Helle, W. & Sabelis, M.W. (1985). *Spider mites their biology, natural enemies and control.*

World Crop Pests. Elsevier Science Publishers, Amsterdam.

Holditch, Z. & Smith, A.D. (2020). Priority determines Tribolium competitive outcome in a

food-limited environment. *PLoS One*, 15, 1–14.

Hougen-Eitzman, D. & Karban, R. (1995). Mechanisms of interspecific competition that

result in successful control of Pacific mites following inoculations of Willamette mites

on grapevines. *Oecologia*, 103, 157–161.

Kant, M.R., Sabelis, M.W., Haring, M.A. & Schuurink, R.C. (2008). Intraspecific variation in

a generalist herbivore accounts for differential induction and impact of host plant

defences. *Proc. R. Soc. B Biol. Sci.*, 275, 443–452.

Kardol, P., Souza, L. & Classen, A.T. (2013). Resource availability mediates the importance

of priority effects in plant community assembly and ecosystem function. *Oikos*, 122, 84–

94.

Ke, P.J. & Letten, A.D. (2018). Coexistence theory and the frequency-dependence of priority

effects. *Nat. Ecol. Evol.*, 2, 1691–1695.

Khodayari, S., Colinet, H., Moharramipour, S. & Renault, D. (2013). Seasonal changes in the

cold hardiness of the two-spotted spider mite females (Acari: Tetranychidae). *Environ.*

Entomol., 42, 1415–1421.

Levine, J.M. & HilleRisLambers, J. (2009). The importance of niches for the maintenance of

species diversity. *Nature*, 461, 254–257.

Matías, L., Godoy, O., Gómez-Aparicio, L. & Pérez-Ramos, I.M. (2018). An experimental

extreme drought reduces the likelihood of species to coexist despite increasing

intransitivity in competitive networks. *J. Ecol.*, 106, 826–837.

Moreira, X., Abdala-Roberts, L., Hernández-Cumplido, J., Cuny, M.A.C., Glauser, G. &

Benrey, B. (2015). Specificity of induced defenses, growth, and reproduction in lima

bean (Phaseolus lunatus) in response to multispecies herbivory . *Am. J. Bot.*, 102,

1300–1308.

Orians, C.M., Pomerleau, J. & Ricco, R. (2000). Vascular architecture generates fine scale

variation in systemic induction of proteinase inhibitors in tomato. *J. Chem. Ecol.*, 26,

471–485.

Orians, C.M. & Jones, C.G. (2001). Plants as resource mosaics: A functional model for

predicting patterns of within-plant resource heterogeneity to consumers based on

vascular architecture and local environmental variability. *Oikos*, 94, 493–504.

Orsucci, M., Navajas, M. & Fellous, S. (2017). Genotype-specific interactions between

parasitic arthropods. *Heredity (Edinb)*., 118, 260–265.

Qi, M., DeMalach, N., Sun, T. & Zhang, H. (2021). Coexistence under hierarchical resource

exploitation: the role of R*-preemption tradeoff. *bioarXiv*, 1–23.

Queijeiro-Bolaños, M.E., González, E.J., Martorell, C. & Cano-Santana, Z. (2017).

Competition and facilitation determine dwarf mistletoe infection dynamics. *J. Ecol.*,

105, 775–785.

R Core Team. (2021). R: A language and environment for statistical computing.

Riahi, E., Shishehbor, P., Nemati, A.R. & Saeidi, Z. (2013). Temperature effects on

development and life table parameters of Tetranychus urticae (Acari: Tetranychidae). *J.*

Agric. Sci. Technol., 15, 661–672.

Rosario-Martinez, H. De. (2015). phia: Post-Hoc Interaction Analysis.

Sarmento, R.A., Lemos, F., Bleeker, P.M., Schuurink, R.C., Pallini, A., Oliveira, M.G.A., *et*

al. (2011a). A herbivore that manipulates plant defence. *Ecol. Lett.*, 14, 229–236.

Sarmento, R.A., Lemos, F., Dias, C.R., Kikuchi, W.T., Rodrigues, J.C.P., Pallini, A., *et al.*

(2011b). A herbivorous mite down-regulates plant defence and produces web to exclude

competitors. *PLoS One*, 6, 8–14.

Scheffer, M., Van Nes, E.H. & Vergnon, R. (2018). Toward a unifying theory of biodiversity.

Proc. Natl. Acad. Sci. U. S. A., 115, 639–641.

Song, C., Rohr, R.P., Vasseur, D. & Saavedra, S. (2020). Disentangling the effects of

external perturbations on coexistence and priority effects. *J. Ecol.*, 108, 1677–1689.

Sousa, V.C., Zélé, F., Rodrigues, L.R., Godinho, D.P., Charlery de la Masselière, M. &

Magalhães, S. (2019). Rapid host-plant adaptation in the herbivorous spider mite

Tetranychus urticae occurs at low cost. *Curr. Opin. Insect Sci.*, 36, 82–89.

Spaak, J.W., Godoy, O. & Laender, F. (2021). Mapping species niche and fitness differences

for communities with multiple interaction types. *Oikos*, 1–13.

Spaak, J.W. & De Laender, F. (2021). Intuitive and broadly applicable definitions of niche

and fitness differences. *Ecol. Lett.*, 23, 1117–1128.

Stam, J.M., ChrÉtien, L., Dicke, M. & Poelman, E.H. (2017). Response of Brassica oleracea

to temporal variation in attack by two herbivores affects preference and performance of

a third herbivore. *Ecol. Entomol.*, 42, 803–815.

Stam, J.M., Dicke, M. & Poelman, E.H. (2018). Order of herbivore arrival on wild cabbage

populations influences subsequent arthropod community development. *Oikos*, 127,

622   1482–1493.

623 Utsumi, S., Ando, Y. & Miki, T. (2010). Linkages among trait-mediated indirect effects: A

624   new framework for the indirect interaction web. *Popul. Ecol.*, 52, 485–497.

625 Vannette, R.L. & Fukami, T. (2014). Historical contingency in species interactions: Towards

626   niche-based predictions. *Ecol. Lett.*, 17, 115–124.

627 Vieira, E.A., Flores, A.A.V. & Dias, G.M. (2018). Persistence and space preemption explain

628   species-specific founder effects on the organization of marine sessile communities. *Ecol.*

629   *Evol.*, 8, 3430–3442.

630 White, N., Bale, J.S. & Hayward, S.A.L. (2018). Life-history changes in the cold tolerance of

631   the two-spot spider mite Tetranychus urticae: applications in pest control and

632   establishment risk assessment. *Physiol. Entomol.*, 43, 334–345.

633 Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis.

634 Wilke, C. (2020). cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."

635 Zélé, F., Santos, I., Olivieri, I., Weill, M., Duron, O. & Magalhães, S. (2018). Endosymbiont

636   diversity and prevalence in herbivorous spider mite populations in South-Western

637   Europe. *FEMS Microbiol. Ecol.*, 94, 1–11.

Figure 1 – Proportion of spider mites *Tetranychus evansi* females (y-axis) depending on initial

frequency (number of initial females *T. evansi*: *T. urticae*, x-axis) and order of arrival (same

time vs. *T. evansi* or *T. urticae* arriving 48h before its competitor) after two generations.

Tetranychus evansi is the better competitor overall (ratio above 0.5), unless *T. urticae* arrives

first or is at higher initial frequency. A posteriori contrasts show a strong effect of order of

arrival in the proportion of females of the two species (Table S2B). Initial frequency also

impacts the final ratio, with a stronger effect when *T. urticae* arrives first or at the same time

than *T. evansi* (Table S2B). Boxplots represent median and quartiles of the 10 boxes within

treatment.

Figure 2 – Relationship between average fitness differences ($\frac{\kappa_j}{\kappa_i}$, y-axis) and stabilising niche

differences (1- ρ, x-axis) for different orders of arrival (*Tetranychus evansi* first – red, same

time – blue, *T. urticae* first – yellow). Plotting average fitness differences against niche

differences allows mapping different competitive outcomes predicted by modern coexistence

theory. The coexistence condition (eq. 4) and its symmetrical for each competing species,

represented by the two solid black lines, allow defining the space in which species can coexist

due to negative frequency dependence or enter alternative stable states due to positive

frequency dependence, whenever niche differences are greater or smaller than zero

respectively. Otherwise, the species with higher fitness will exclude the other. In our case, the

only scenario in which species are predicted to coexist is when *T. urticae* arrives first (yellow).

Error bars for each outcome indicate the 95% confidence interval from the maximum

likelihood estimates. For the other two cases, it is predicted that the superior competitor *T.*

evansi will exclude *T. urticae*.

Figure 3 – Differences between expected and observed leaf occupancy for *Tetranychus evansi*

(A) and *T. urticae* (C) for a subset of the experimental treatments (when *T. urticae* arrived first

or at the same time as *T. evansi*, note that Figure S3 includes all treatments); leaf occupancy

for *T. evansi* (B) and *T. urticae* (D) in the control, single species, treatments. Leaf 2 corresponds

to the oldest leaf and leaf 5 to the youngest. For each box, we calculated the ratio of females

occupying each leaf in relation to the total number of females present. For the experimental

treatments we calculated the difference between this ratio and the average ratio for the control

treatments. Thus, positive values indicate that there are more females on that leaf than expected

based on the single-species treatment and negative values indicate the reverse pattern. Overall,

we see that. *T. evansi* reduces occupancy on leaf 4 when *T. urticae* arrives first and on leaf 3

when the two species arrive at the same time. In contrasts, *T. urticae* shows a slight increase in

occupancy of leaf 4 when it arrives first and a slight decrease in occupancy of leaves 2 and 5.

677
678

Figure 1 – Proportion of spider mites *Tetranychus evansi* females (y-axis) depending on initial

frequency (number of initial females *T. evansi*: *T. urticae*, x-axis) and order of arrival (same

time vs. *T. evansi* or *T. urticae* arriving 48h before its competitor) after two generations.

Tetranychus evansi is the better competitor overall (ratio above 0.5), unless *T. urticae* arrives

first or is at higher initial frequency. A posteriori contrasts show a strong effect of order of

arrival in the proportion of females of the two species (Table S2B). Initial frequency also

impacts the final ratio, with a stronger effect when *T. urticae* arrives first or at the same time

than *T. evansi* (Table S2B). Boxplots represent median and quartiles of the 10 boxes within

treatment.

Figure 2 – Relationship between average fitness differences ($\frac{\kappa_j}{\kappa_i}$, y-axis) and stabilising niche

differences (1- ρ, x-axis) for different orders of arrival (*Tetranychus evansi* first – red, same

time – blue, *T. urticae* first – yellow). Plotting average fitness differences against niche

differences allows mapping different competitive outcomes predicted by modern coexistence

theory. The coexistence condition (eq. 4) and its symmetrical for each competing species,

represented by the two solid black lines, allow defining the space in which species can coexist

due to negative frequency dependence or enter alternative stable states due to positive

frequency dependence, whenever niche differences are greater or smaller than zero

respectively. Otherwise, the species with higher fitness will exclude the other. In our case, the

only scenario in which species are predicted to coexist is when *T. urticae* arrives first (yellow).

Error bars for each outcome indicate the 95% confidence interval from the maximum

likelihood estimates. For the other two cases, it is predicted that the superior competitor *T.*

evansi will exclude *T. urticae*.

Figure 3 – Differences between expected and observed leaf occupancy for *Tetranychus evansi*

(A) and *T. urticae* (C) for a subset of the experimental treatments (when *T. urticae* arrived first

or at the same time as *T. evansi*, note that Figure S3 includes all treatments); leaf occupancy

for *T. evansi* (B) and *T. urticae* (D) in the control, single species, treatments. Leaf 2 corresponds

to the oldest leaf and leaf 5 to the youngest. For each box, we calculated for each species the

proportion of females occupying each leaf in relation to the total number of females present.

For the experimental treatments we calculated the difference between this proportion and the

average proportion for the control treatments. Thus, positive values indicate that there are more

females on that leaf than expected based on the single-species treatment and negative values

indicate the reverse pattern. Overall, we see that. *T. evansi* reduces occupancy on leaf 4 when

T. urticae arrives first and on leaf 3 when the two species arrive at the same time. In contrasts,

T. urticae shows a slight increase in occupancy of leaf 4 when it arrives first and a slight

decrease in occupancy of leaves 2 and 5.