Title: Predicting selection-response gradients of heat tolerance in a wide-ranging reef-building
coral

RUNNING TITLE: selection-response gradients in corals

Ponchanok Weeriyanun[1,2*], Rachael B. Collins[1,3*], Alex Macadam[1], Hugo Kiff[4], Janna L. Randle[1],
Kate M. Quigley[1,**]

[1]Australian Institute of Marine Science, Townsville, 4810, Australia

[2]Ghent University, Sint-Pietersnieuwstraat 33, 9000 Gent, Belgium

[3]University of Plymouth, Plymouth, PL4 8AA, United Kingdom

[4]Liverpool John Moores University, Liverpool, L3 3AF, United Kingdom

*These authors contributed equally to the manuscript

**Corresponding author: Kate M. Quigley; katemarie.quigley@my.jcu.edu.au

KEYWORDS: thermal tolerance, heritability, coral reefs, larvae, settlement, survival

SUMMARY STATEMENT: Heat stress exerts disruptive selection on adult corals. This likely underpins variability in offspring survival and results in differences in offspring responses to selection.

ABSTRACT

Ocean temperatures continue to rise due to climate change but it is unclear if heat tolerance of marine organisms will keep pace. Understanding how tolerance scales from individuals to species and quantifying adaptive potentials is essential to forecasting responses to warming. We reproductively crossed corals from a globally distributed species (*Acropora tenuis*) on the Great Barrier Reef (Australia) from three thermally distinct reefs to create 85 novel offspring lineages. Individuals were experimentally exposed to temperatures (27.5, 31, and 35.5 - 36 °C) in adult and two critical early life-history stages (larval development and settlement) to assess acquired heat tolerance via introgression on offspring phenotypes by comparing multiple physiological responses (photosynthetic yields, bleaching, necrosis, settlement, and survival). Adaptive potentials and physiological reaction norms were calculated across multiple life-stages to integrate heat tolerance at different biological scales. Selective breeding improved larval survival to heat by 1.5-2.5x but settlement success showed limited improvement. Adult responses to selection at heat were similar but were greater in larvae from warmer reefs compared to the cooler reef. There was also a divergence between adults and offspring mean population responses, likely underpinned by heat stress imposing strong divergent selection on adult colonies. These results have implications for downstream selection during reproduction, as evidenced by variability in a conserved heat tolerance response across offspring lineages. These results inform our ability to forecast the impacts of climate change on wild populations of corals and will aid in developing novel conservation tools like the assisted evolution of at-risk species.

INTRODUCTION

Ecosystems globally continue to suffer from the adverse effects of anthropogenic climate change. Coral reef ecosystems in particular are declining as sea surface temperatures (SSTs) have risen dramatically (Hughes et al., 2017). Coral reefs are some of the most productive habitats and the continued loss of corals will likely impact reef community dynamics by changing ecological functioning via trophic interactions, with downstream impacts on fish and other invertebrate species (Jones et al., 2004; Komyakova et al., 2013; Nelson et al., 2016). Coral reefs are biodiversity hotspots that provide access to food, act as natural barriers against storm damage and benefit the economy through tourism and other recreational activities (O'Mahony et al., 2017). Coral reef ecosystems like the Great Barrier Reef (GBR) contribute, on average, $6.4 billion (AUD) to the Australian economy per annum (O'Mahony et al., 2017). Consequently, the loss of these crucial habitats will have devastating economic and social-ecological effects on coastal communities.

The upper thermal thresholds of reef-building corals generally reside ~1 °C higher than their local summer maxima (Schoepf et al., 2019). Once temperatures exceed this threshold, corals may lose their symbiotic dinoflagellates (Symbiodiniaceae) through a process known as bleaching, which is typically experienced by corals during marine heatwaves if warming persists for extended periods of time (Hoegh-Guldberg, 1999). Extreme bleaching (defined as >60 % of bleached corals present within a reef) often leads to high mortality (Hughes et al., 2017). At least four mass bleaching events have been recorded on the GBR in the last century - 1998, 2002, 2016 and 2017 – generally increasing in intensity and severity (Hughes et al., 2017) and have led to large-scale losses in hard coral cover. Rates of decline vary by region, but overall, declines have rapidly increased to an average of ~1.45 % per year since 2006 (De'Ath et al., 2012), although there have been recent signs of recovery. Therefore, bleaching related stress has not been equal across the GBR. This trend in decline is reflected globally, with SSTs continuing to exceed previously held records (Heron et al., 2016), suggesting that corals may soon reach their physiological limits to cope with increasing heat (Matz et al., 2017).

Changes in temperature have the potential to impact organismal life-stages differently, in which larvae, recruitment stages, and adults can all vary in their thresholds to climate extremes. In corals, sexually mature adults release their gametes within a temperature range of ~28 – 30 °C, with larvae

able to survive 2 – 5 °C above this range (Heyward and Negri, 2010). However, coral spawning

typically occurs during the warmest summer months, when spikes in SSTs are most likely to occur

(Keith et al., 2016). As a result, coral larvae are potentially subjected to much greater temperatures,

increasing their risk of mortality and reducing recruitment and settlement success (Heyward and

Negri, 2010). After fertilisation, larvae undergo a pre-competency period where they develop and

disperse throughout the water column. Once competent, they respond to biophysical and chemical

cues to find optimal settling conditions (Doropoulos et al., 2018). Increased SSTs may induce

premature metamorphosis by increasing larval metabolic activity and development rates, reducing

pre-competency periods (Heyward and Negri, 2010). This reduction may influence subsequent larval

dispersal distances and result in settlement occurring in suboptimal conditions (Edmunds et al., 2001),

with the potential to reducing dispersal, and therefore gene flow, between populations, thereby

accelerating population differentiation and potentially local adaption (Bassim et al., 2002). Therefore,

post-settlement recruits and later-stage juveniles appear to be highly susceptible to temperature-

induced phenotypic influences (e.g. on growth and survival; Kenkel et al., 2015). This information is

largely unknown for coral early life-history stages but should be incorporated into predictive models

of ecosystem change given the flow-on effects into key demographic and population level dynamics.

There is scope for adaptation of corals to increasing temperatures (Matz et al., 2018). Further, the

temperatures at which bleaching is occurring have increased by ~0.5 °C from 1998 to 2017,

suggesting an increase in more heat adapted genotypes within coral populations (Sully et al., 2019),

potentially through processes like selective sweeps (Quigley et al., 2019a). Locally, there have been

reported increases in coral cover in both the central and southern GBR (AIMS, 2020), with evidence

to suggest that warm or "extreme" habitats harbour an increased number of thermally adapted

genotypes with the potential to transmit heat tolerance (Quigley et al., 2020a; Schoepf et al., 2019).

However, current estimates of the rate of SST increases suggest that these increases may exceed the

potential rate of fixation of beneficial genetic variants given factors like currents and reef topology

(Quigley et al., 2019a). Additionally, the annual increase in SSTs has extended the period at which

'summer' temperatures occur, further increasing global bleaching by reducing potential recovery

periods that occur when coral populations are exposed to cooler 'winter' temperatures (Heron et al.,

2016). Taken together, this suggests that corals adaptive potentials may be constrained.

Variability in heat tolerance has been observed at many ecological levels, including between

individuals, populations, species, and by reef region. For example, differences in the heat tolerance of

corals have been observed across reefs globally, with coral populations present along the Persian Gulf having demonstrated very high thermal thresholds of ~4 °C above mean monthly temperatures (Kirk et al., 2018; Moghaddam et al., 2021; Savary et al., 2021). Northern GBR corals also demonstrate higher thermal tolerances compared to some central populations, in which heritable host genetic mechanisms play a role in the thermal resistance of these northern corals (Dixon et al., 2015; Quigley et al., 2020a). However, it is currently unclear whether similar patterns in heat tolerance at the individual coral level scale to the population level, a trend seen in fish but not insect or plant species collected across temperature clines (Payne et al., 2021; Rezende and Bozinovic, 2019). Quantifying the scaling of heat tolerance across different biological levels will help to elucidate the baseline potential of wild populations to adapt. Further, evolutionary models incorporating metrics like narrow-sense heritability, selection coefficients, and responses to selection are commonly measured (Falconer and Makcay, 1996), and allow for the prediction of organismal responses to future stressors. Thus far, models incorporating the "breeders equation" (Falconer and Makcay, 1996) have demonstrated the utility of evolutionary modelling for predicting the potential of the coral's algal symbionts to confer increased survival (Quigley et al., 2018). Similarly, these principles can be applied to the coral host to quantify adaptive potentials, in which this information can then be used to harness the natural adaptive potential for conservation and restoration.

Coral restoration interventions, some of which can be classified as assisted evolution methods (van Oppen et al., 2015), aim to increase the adaptive potential of organisms by targeting the acceleration of adaptation through various genetic mechanisms. This may include selective breeding for the selection of desirable phenotypes (van Oppen et al., 2015). Assisted Gene Flow (AGF) is one intervention that utilises the differing thermal tolerances of parental colonies combined with the intentional movement of recombinant offspring, by reproductively mixing populations sourced from different environmental conditions within the same species to prepare future populations for increased warming. This strategy is based on increasing gene flow between populations that have a beneficial phenotype(s) with those of a target population(s) via the creation and subsequent introduction of new, more resilient genotypes (Aitken and Whitlock, 2013). Selective breeding through the outcrossing of gametes between these reefs of distinct thermal profiles assumes that populations exposed to climates of similar temperatures to those predicted to arise from anthropogenic climate change may be better adapted, and therefore better able to cope with warming (Aitken and Whitlock, 2013). The inheritance of these beneficial alleles by populations that evolutionarily have experienced cooler temperatures should increase the overall resilience of the species via increased population persistence (National Academies of Science, Engineering and Medicine, 2019).

Here, we compare the physiological responses at heat stress of selectively bred *Acropora tenuis* larvae and newly settled recruits and their parental colonies collected from three sites along the Great Barrier Reef – Davies and Esk reefs (central GBR) and in the Keppels (southern GBR). Percentage larval survival and settlement were recorded following exposure to control (27 °C) and heat stress conditions (35.5 - 36 °C) and compared to adult heat tolerance at 31 °C, as measured in bleaching score, photosynthetic quantum yields, percent necrosis, and percent survival. We then compared these responses across multiple biological scales (individuals to populations) to calculate adaptive potentials from data collected across each life-stage. This work contributes to our understanding of the potential to increase heat tolerance and for the transgenerational transfer of this tolerance using the selective breeding of coral populations sourced from along the GBR.

MATERIALS AND METHODS

Coral colony collections

Reproductively mature *Acropora tenuis* colonies were collected in early November 2019 from three sites on the GBR in Australia, including two central reefs: Davies (-18.90, 147.87) and Esk (-18.86, 146.62), and one southern reef in the Keppel islands (23.06, 150.95). Corals were transported by boat to the National Sea Simulator at the Australian Institute of Marine Science (AIMS). Colonies from each location were acclimated in separate outdoor holding tanks under constant 0.2 μm filtered seawater (FSW) flow-through conditions, maintained at 27.5 °C. Esk was on average the warmest reef location, followed by Davies, and then the Keppels (**Fig. 1A**). The mean maximum monthly temperature of each site was as follows: Esk 26.26 ±2.30 °C, Davies 26.06 ±1.80 °C, and the Keppels 24.62 ±2.30 °C (**Fig. 1B**).

Coral spawning, selective breeding, and larval rearing

Prior to spawning, each *A. tenuis* colony was given an identification number (**Table S1**). Individual colonies were isolated in separate bins following signs of spawning imminence, which includes the appearance of egg–sperm bundles under the oral disc and "setting" (polyps extended but tentacles retracted). Gamete bundles were released between 19:00 and 19:40 on the 17th and 18th of November

2019 and collected by gently skimming and collecting the bundles off the surface of the water. The

eggs and sperm in the bundles were separated by washing for ~2 mins with FSW through a 120 μm

mesh filter. The number of sperm were quantified using a Computer Assisted Semen Analyser

(CASA) (software CEROS II from Hamilton Thorne). The sperm from one parental colony was then

added at a concentration of 10^5 mL to isolated and cleaned eggs from another parental colony to create

85 distinct coral families comprising of intrapopulation (within the same reef) and interpopulation

(between different reefs) crosses, where both are generally referred to here as population crosses

(**Table S2**).

Eggs were allowed to fertilize in separate bowls per cross for 3 hours. Every hour aliquots of

developing embryos were taken from each family (hereafter referred to as crosses; **Fig. 1C-D**) to

visually inspect fertilization success and cell division under magnification. Once fertilization was

confirmed, embryos from each separate bowl were transferred into separate 15 L constant flow-

through conical tanks with 0.2 μm FSW in a temperature-controlled room, such that the temperature

of each cone was maintained at 27.5 °C. Each flow-through cross culture had an outflow covered by a

10 μm filter and air curtain of bubbles to prevent larvae from collecting on the outflow filter. By days

three and four post-fertilisation, larvae were ciliated and motile, consistent with the 96-hour stage of

larval development. Population-level crosses are referred to as follows with the maternal colony first

and then paternal colony, in which intrapopulation crosses (filled circles; **Fig. 1D**) are ExE (EskxEsk),

KxK (KeppelsxKeppels), DxD (DaviesxDavies), and interpopulation crosses are ExK (EskxKeppels),

ExD (EskxDavies), KxE (KeppelsxEsk), KxD (KeppelsxDavies), DxE (DaviesxEsk), and DxK

(DaviesxKeppels) (open circles; **Fig. 1D**). No crosses perished post-fertilization before sampling took

place.

Larval heat stress experiment

Thirty larvae from each of the total 85 crosses were placed into floating net-wells (n = 3 replicates per

cross, per temperature), separated into two holding tanks. One tank was set at 27.5 °C (control

treatment) and the other at 35.5 °C (heat treatment). To achieve this temperature, the heat treatment

was ramped up to 35.5 °C in hourly increments of 0.5 °C from 27.5 °C. Once 35.5 °C was reached,

survival assessments began, in which the number of larvae that were alive in each net-well (of the

total n = 30) was counted twice daily to estimate survival rates per replicate. The final survival counts

occurred once ~50 % of the larval crosses reached 50 % survival in the heat treatment.

Settlement experiment

Of the original 85 lineages produced during the reproductive crosses, a subset of 69 crosses were chosen for settlement to cover a range of available inter- and intrapopulation combinations. The selected lineages of larvae were sampled 35 days after spawning and transferred to sterile six-well plates (n = 10 larvae per well, n = 3 wells per cross, n = 2 crosses per sterile six-well plate) containing 10 mL of 0.2 μm FSW. Crustose coralline algae (CCA) rubble was freshly cut into 3 x 3 mm chips using bone cutters, and placed into each well to induce larval settlement. The plates were then placed into plastic bags and sealed to prevent evaporation and replicates were transferred into two incubators with photosynthetic lights (Steridium E-500, lights: Sylvania FHO24W/T5/865 and Innova 4230, light: Sylvania F15W/865) set at 27 - 27.5 °C (control treatment) and 35.5 - 36°C (heat treatment). The number of settled larvae (metamorphosed and attached to substrate) were counted for 17, 24 and 48 hours after incubation at both temperatures and the number of settled larvae was recorded. Settled larvae were defined as those that were attached to the well or CCA and deposited a basal plate that was visible after metamorphosis. This was distinguished from only metamorphosed larvae (metamorphosis but no attachment to substrate), which were excluded from this analysis.

Adult heat stress experiment

Three to five colonies representing different genotypes for each population were fragmented using an "diamond-tipped" bandsaw (**Table S1**). Each colony was divided into a minimum of six fragments, with each placed into one of six experimental tanks (n = 3 control tanks, n = 3 heat tanks). For the Davies population, five genotypes were selected and cut into 82 fragments; three genotypes were used from the Esk population (51 fragments); and three genotypes from Keppels population (28 fragments). Each fragment was glued to a calcium carbonate plug using epoxy glue, placed into PVC plug holders (hereafter referred to as "sticks") and allowed to acclimate at control temperatures before being moved to the experimental tanks. This acclimation period attempts to minimize the effects of sampling and handling before the experiment, where all coral fragments were allowed to acclimate and recover from the fragmentation stress for one week. During this time and over the course of the experiments, corals were fed once a day at 16:30 hours with 0.5 individual/mL of *Artemia* sp. and 2000 cells/mL of microalgae.

Upon moving the corals to experimental tanks, the position of each fragment was randomized in each tank and across all tank replicates, such that each tank included the same genotypes from each

population in all tanks. Experimental tanks were filled with 45 L of either 27.5 °C FSW (control treatment) or 31 °C (heat treatment) on constant flow through. Each tank was equipped with a tungsten aquarium pump for constant aeration and mixture of water. The heat treatment corals were ramped at a rate of 0.5 °C until 31 °C was reached. The light cycle for each tank was set on a 12:12 hour regime and light intensity at 171 PAR. Sunrise was set at 21:00 hours. The coral fragments underwent these experimental conditions for 14 days.

Coral colour, as a proxy for bleaching, was determined using an underwater colour reference card for corals (Coral Watch Card; Siebeck et al., 2008). Images with the bleaching card were taken at least twice a week, and fragments were compared to the brown hue (D1-D6). ImageJ (Schneider et al., 2012) was used to create a Red-Green-Blue standard curve for coral colour as per Quigley et al. (2019b). Survival of each coral fragment was noted from the images. Coral death was determined by the presence of microalgae growing on the bare skeleton on each coral fragment. The percentage of tissue necrosis was measured using the ImageJ surface area tool (Schneider et al., 2012) per coral fragment, per tank, per time point. Effective quantum yield ($\Delta F/FM'$), which is the efficiency of photon absorption, was measured using the Diving PAM (Walz, 2018). PAM measurements of each fragment were taken twice per week at 10:00 h.

Statistical analysis

Larval heat stress experiment

Differences in larval survival were assessed using the non-parametric *"wilcox.test"* from the package "ggpubr" (Kassambaraype, 2020) to compare if the mean values between each cross at the control and heat treatments (two independent groups) were statistically different.

Settlement experiment

The percentage of settled larvae was first analyzed using the base "stats" package (R Core Team, 2021) to test the normality and homogeneity of the values. If non-normal distributions were present, as shown through diagnostic plots, the random factors "cross" and "plate" were tested for their contribution to variability in settlement. The "ggpubr" package (Kassambaraype, 2020) and the

Wilcoxon's test (Wilcoxon, 1945) were used to statistically compare the median percentage settlement
between heat and control treatments at each time point.

Adult heat stress experiment

Several coral traits have been identified as important biometrics for restoration, including partial

mortality and bleaching, in conjunction with classic response measurements like survival (Baums et

al., 2019). Here bleaching was assessed by the change in colour from photographs, where a "6" is

indicative of a non-bleached, healthy fragment and a "0" is a white, bleached fragment. Differences in

the photophysiological responses of the algal symbionts within corals ($\Delta F/FM'$), percent necrosis,

bleaching and survival were evaluated with respect to temperature treatment and coral population

using linear models implemented in the "lme4" package (Bates, 2005).

Bleaching, $\Delta F/FM'$, and the percentage of necrosis were treated as continuous variables, and

temperature treatment and population were set as fixed factors in each model. Replicate tank and

fragment holder ("sticks") were set as random effects. These factors were not significant and were

dropped from the final model (**Table S3**). Once these random effects were dropped for each model

($\Delta F/FM'$, bleaching, percentage of necrosis, survival), the linear model was re-fit, and all model

assumptions were checked (linearity, normality, and homogeneity) using diagnostic plots in the

"stats" package (R Core Team, 2021). The function "*anova*" was used to calculate model p-values,

used for interpreting the significant difference among means. Post-hoc pairwise comparisons of

population and temperature treatment were then run on the model outputs (Tukey, 1977). For survival

data, a binomial distribution was used to see if there was a significant difference between populations

and temperature treatments. For each trait, the statistical difference in median values was assessed

using the packages "ggplot2", and "plyr" (Wickham, 2011; Wickham, 2016), with the Wilcoxon's test

(Wilcoxon, 1945). The percentage change of adult responses by latitude were calculated and plotted

using "ggplot2" package (Wickham, 2016). All analyses were carried out using RStudio, version

2.13.2 (R Core Team, 2021).

Predicting response gradients to selection

The density plots of adult and larval survival were made by using the "ggplot2" package (Wickham,

2016). The breeder's equation (Falconer and Makcay, 1996) was used to calculate expected responses

to selection (R) and potential constraints to the evolution of heat tolerance given selective potentials

(S) and narrow-sense heritability for this trait (h^2). This approach has previously been used in corals

(Quigley et al., 2018) to determine how selection on host symbiont communities may adapt to

different selective environments. Contour plots of R, S and h^2 for larvae grouped by maternal reef or

origin were made using the packages "plotly" (Plotly Technologies Inc., 2015) and "dplyr" (Hadley

Wickham et al., 2020).

RESULTS

Differences in larval survival under heat stress

The influence of both temperature treatments on larval survival varied by population cross, where

larval lineages produced from Davies, Esk and Keppels corals displayed both high and low survival

under control and heat conditions (**Fig. 2A**). Both median survival and the variance around the median

at both control and heat conditions varied across the lineages. Overall, the Keppels intrapopulation

larvae survived better at heat compared to interpopulation larvae, whilst the Esk and Davies larvae

survived better when crossed with either of the other two reefs as interpopulation crosses. The overall

"winners" at heat were DxE and ExK (which also had high survival at control), while the overall

losers at heat were KxD and KxE.

When scaled to the population level, Davies purebred larvae and larvae produced using Davies eggs

were only significantly different in their survival between temperatures for one of the three lineages

(**Fig. 2B**, $P = 0.71$, 2.50×10^{-2}, 9.70×10^{-2} for DxD, DxE, and DxK, respectively). Survival at heat was

higher in the interpopulation crosses compared to the purebred larvae (median survival: DxD = 56.70

%, DxE = 75.00 %, DxK = 62.30 %). Again, Esk purebred larvae and larvae produced with Esk eggs

were only significantly different in their survival between temperatures for one of the three lineages,

with the same (reciprocal) cross significantly different ($P = 4.10 \times 10^{-2}$, 0.53, 0.29 for ExD, ExE, and

ExK, respectively). Survival at heat was higher for interpopulation crosses than the purebred Esk

larvae (ExD = 61.70 %, ExE = 38.30 %, ExK = 81.70 %). The variation in median survival was also

greater in purebred Esk larvae compared to other lineages at heat. Keppels interpopulation larvae all

survived less at heat compared to control conditions ($P = 3.20 \times 10^{-2}$ and 2.30×10^{-8} for KxD and KxE)

whilst purebred larvae survived about equally between treatments ($P = 0.61$). Median survival was

lower in the interpopulation larvae compared to purebred larvae (KxK= 65.60 %, KxD = 46.70 %,

KxE = 36.70 %).

Influence of heat stress on larval settlement rates

When exposed to heat, larvae from all crosses significantly decreased in their settlement behaviour (attachment and metamorphosis) relative to the control temperature after 17 ($F_{1,433}$= 370.30, P = 2.00×10^{-16}), 24 (6/9 crosses with 0.00 % settlement; $F_{1,430}$= 916.50, P = 2.00×10^{-16}), and 48 hours (7/9 crosses with 0.00 % settlement; $F_{1,334}$= 722.80, P = 2.00×10^{-16}; **Fig. 3A-C**). Specifically, larvae from all crosses settled significantly less at heat compared to control temperatures, regardless of population cross (P-values in **Table S4**). DxE and ExK showed the greatest percentage settlement compared to the other crosses (median settlement at heat = both 0.00 %, control = 60.00 and 75.00 %, respectively), in which DxE was the only population cross that settled after 48 hours of heat exposure.

There was a significant effect of temperature treatment ($F_{1,1633}$ = 887.50, P = 2.00×10^{-16}) and heat exposure time ($F_{3,1631}$ = 120.70, P = 2.00×10^{-16}) on the percentage of settled larvae and a significant interaction between temperature treatment and heat exposure time ($F_{3,1564}$ = 225.42, P = 2.00×10^{-16}), temperature treatment and population level cross ($F_{8,1564}$ = 5.30, P = 1.39×10^{-6}). There was no significant interaction between heat exposure time and population-level cross ($F_{24,1564}$ = 1.06, P = 3.86×10^{-1}) or between temperature treatment, heat exposure time and population-level cross ($F_{23,1564}$ = 1.26, P = 1.82×10^{-1}). This significant decrease in settlement at 24 and 48 hours occurred regardless of larval cross ($F_{8,1626}$ = 1.583, P = 1.25×10^{-1}) or reef of origin of the maternal coral ($F_{2,1632}$ = 0.133, P = 8.75×10^{-1}).

After 17 hours of temperature incubation, a median value of 0.00 % larval settlement was observed for all crosses in the heat treatment and was significantly lower compared to the control temperature (median 70.00 %) across all larval lineages (Wilcoxon's test, P < 0.001; **Fig. 3A**). In the control treatment, the median settlement percentage was highest in DxD and KxE, with a median of 80.00 % settlement. The lowest median percentage settlement was observed in the ExE crosses, with 55.50 % settlement and correspondingly high variability.

After 24 hours (**Fig. 3B**), median settlement remained at 70.00 % in the control treatment and ~0.00 % in the heat treatment, with all crosses settling significantly less at heat compared to controls (Wilcoxon's test, P < 0.001). At this time point, KxD showed the highest percentage settlement (80.00

%) at control, followed by DxK and KxK (75.00 %). Finally, after 48 hours (**Fig. 3C**), the median

settlement percentage in control treatment dropped slightly to 66.67 %, where KxD crosses again

exhibited the highest larval settlement at control (75.00 %). At heat, the only cross to settle included

some of the larval replicates within DxE. Finally, some crosses also experienced mortality of recent

recruits from 17 to 48 hours (DxD), whilst others continued to settle (DxE).

Adult responses to heat stress at the population level

There is a significant effect of temperature treatment (**Fig. 4A**; negative binomial generalized linear

model-nbglm, $P = 2.20\text{x}10^{-16}$) and population origin (nbglm, $P = 3.56\text{x}10^{-15}$) on the mean bleaching

score of coral fragments after 16 days. At the control temperature, fragments sourced from Davies

recorded a mean colour score of 4.75 ±0.14 (median = 5.00) whilst fragments sourced from Keppels

and Esk were 1.85 ±0.25 (median = 2.00) and 1.35 ±0.39 (median = 0.00), respectively (**Fig. 4A**). At

heat, fragments from all populations bleached heavily (all mean scores <1.00, all median = 0.00).

Relative to the control temperature, Davies fragments bleached the most (median percentage change

87.41 ±5.32 % decrease in bleaching category), followed by Keppels (62.96 ±37.04 %), and then Esk

fragments (28.57 ±28.57 %). Pairwise comparisons showed that bleaching score was significantly

different between the control and heat treatments in Davies (mean Tukey's test, $P < 0.001$, median

Wilcoxon $P = 1.80\text{x}10^{-15}$), Esk (Tukey's test, $P < 0.001$, median Wilcoxon $P = 0.11$), and Keppels

(Tukey's test, $P = 0.15$, median Wilcoxon $P = 1.50\times10^{-2}$).

The photophysiological responses, as measured by effective quantum yield (ΔF/FM'), showed that

ΔF/FM' significantly decreased in the heat compared to control treatment (**Fig. 4B**, lm, $P = 1.09\text{x}10^{-15}$)

and by population origin (lm, $P = 6.80\text{x}10^{-8}$). Davies and Keppels fragments reported the highest

yields at control temperatures (mean and median ΔF/FM' >0.60), whereas Esk fragments were

extremely low (0.12 ±0.05, median = 0.00). Mean ΔF/FM' recorded from fragments in the heat

treatment were highest in Keppels (0.22 ±0.07, median = 0.00), Davies (0.20 ±0.05, median = 0.00),

and finally Esk (0.12 ±0.05, median = 0.00) but very variable overall. Relative to control

temperatures, pairwise comparisons showed that only Keppels and Davies fragments significantly

decreased ΔF/FM' (Tukey's test, $P < 0.001$ for both, median Wilcoxon $= 2.40\times10^{-4}$ for Keppels and P

$= 2.70\text{x}10^{-12}$ for Davies) but not Esk (Tukey's test, $P = 5.68\times10^{-1}$, median Wilcoxon $= 0.16$).

Partial mortality was assessed as the percentage of necrotic tissue relative to each fragment (**Fig. 4C**).

Population origin had a significant effect on percentage necrosis (lm, $P = 3.25 \times 10^{-8}$). There was no

significant difference in percentage necrosis due to temperature treatment (lm, $P = 5.50 \times 10^{-2}$). At the

control temperature, Davies and Keppels corals showed no to very little necrosis (median = 0.00 %),

whereas Esk fragments were slightly necrotic (median 15.29 %). After 16 days at heat, fragments

sourced from Esk lost on average, approximately 46.04 ±9.89 % (median = 0.00 %) of their tissue per

fragment, compared to Keppels (26.67 ±11.82 %, median = 0.00 %) and Davies fragments (12.20

±5.17 %, median = 0.00 %). Pairwise comparisons showed that at the control temperature, Esk lost,

on average significantly more tissue compared to Davies (Tukey's test, $P < 0.001$) and Keppels

(Tukey's test, $P = 3.39 \times 10^{-3}$). In the heat treatment, Esk also experienced significantly more necrosis

compared to Davies (Tukey's test, $P = 2.76 \times 10^{-3}$). All other population comparisons were not

significantly different (Esk-Keppels, Tukey's test, $P = 5.35 \times 10^{-1}$, Keppels-Davies, Tukey's test, $P =$

7.43×10^{-1}). Relative to control temperatures, the median percent necrosis was not significantly

different for any population (median Wilcoxon $P = 0.09, 0.91, 5.30 \times 10^{-2}$ for Davies, Esk, Keppels).

Temperature showed significant effect on survival (binomial lm, $P = 1.19 \times 10^{-8}$). In the control

treatment, survivorship was highest in the Keppels fragments (median = 1.00), followed by Davies

(median = 1.00) and Esk (median = 0.00) (**Fig. 4D**). In the heat treatment, survivorship was highest in

the Keppels fragments (40.00 ±13.09 %, median = 0.00), followed by Davies (31.71 ±7.36%, median

= 0.00) and Esk (20.00 ±8.16 %, median = 0.00). Compared between heat and control temperatures,

fragments sourced from Davies and Keppels significantly decreased in survivorship (Tukey's test, $P <$

0.001 and 4.98×10^{-3}, median Wilcoxon = 5.10×10^{-3} and 1.40×10^{-10}, respectively) but not Esk

fragments, which also survived poorly at the control temperature treatment (Tukey's test, $P =$

7.55×10^{-1}, median Wilcoxon = 0.25). Given the significant population effect of temperature treatment

for bleaching score, $\Delta F/FM'$, and necrosis (but not survival), the relative differences in responses

were also calculated and compared across the latitudinal gradient of adult origin (**Fig. 4D**). There was

no trend in performance across traits by latitude.

Predicting adult and offspring responses using gradients of selection

417 When survival was averaged at the population level, almost all adult corals collected from Davies

418 exhibited approximately equivalent survival at the control treatment, but when exposed to the heat

419 treatment, the population responses were generally bimodal, where some individuals within each

population exhibited high survival and others low survival (**Fig. 5A**). Corals collected from Esk and the Keppels also exhibited this bimodal response for both the heat and control temperatures. For both Davies and Keppels corals, the mean survival at the control treatment was high and a roughly equal decrease in survival at heat. Although Esk corals had much lower survival at the control temperature, their overall mean survival at heat was roughly equivalent to Davies and Keppels corals, suggesting that the selection differential for survival at heat in each population was roughly equivalent (horizontal dashed black lines), defined as the difference between a selected phenotypic scope (triangle) and the mean percent survival at heat (vertical dashed red line).

Mean percent survival for larvae in the control treatments were higher (56.44 –78.92 %) compared to at heat (41.48 – 78.40 %). As expected, based on similar adult selection differentials (**Fig. 5A**), purebred larvae from DxD, ExE, and KxK responded similarly in the breadth of larval responses between control and heat (specifically, a small difference between treatment responses). Alternatively, interpopulation crosses differed in their responses, in which KxD and KxE had the largest difference in mean survival between control and heat (17.14 % and 32.84 %) and ExK the smallest (0.52 %).

The DxD purebreds exhibited a 56.44 ±8.77 % mean survival at heat, almost 2x greater survival than the Davies adults at heat (31.71 ±7.36 %; **Fig. 5B**). DxD larval crosses demonstrated high variability (e.g. flat distribution) responses in survival at control, compared to a unimodal response in heat, although the mean survival was roughly equal 56.44 ±8.77 % and 55.78 ±4.71 % respectively, dashed lines). When Davies eggs were crossed with sperm from the other central reef, heat tolerance increased by 14.46 % compared to DxE. DxE larvae exhibited unimodal responses for both temperatures with average survival at 78.38 % at control and 70.24 % at heat. Heat tolerance was slightly less for DxK larvae (55.41 %) compared to DxD, where larval responses were more variable but still unimodal.

The ExE larval purebreds exhibited 48.29 ±7.15 % survival at heat, ~2.5x greater survival than the Esk adults at heat (20.00 ±8.16 %; **Fig. 5C**). ExE larvae at control and heat treatments were relatively flat, demonstrating that both treatments consisted of crosses with high and low survival. When Esk eggs were crossed with sperm from the other reefs, heat tolerance increased by 8.40 % and 30.11 % in ExD and ExK, respectively, relative to ExE. ExD larval survival responses were weakly unimodal with mean survival at 70.29 ±3.47 %, and 56.68 ±4.61 % for control and heat. Alternatively, ExK

larvae exhibited strongly unimodal survival responses at 78.92 ±3.82 % at control and 78.40 ±2.59 %

at heat.

The KxK purebreds exhibited 62.84 ±3.73 % survival at heat, ~1.5x greater survival than the Keppels

adults at heat (40.00 ±13.09 %; **Fig. 5D**). KxK larval responses formed relatively flat distributions in

the control treatment to unimodal responses in the heat. When Keppels eggs were crossed with sperm

from the other reefs, heat tolerance decreased by 17.59 % and 21.35 % in KxD and KxE, respectively.

In KxD, the distribution was flat wide, indicating variation in survival. The KxE density plot for

larvae at the control treatment exhibited a unimodal peak in survival, as well as in the heat treatment.

In the heat stress treatment when larvae were grouped by the population identity of the maternal

colony, larvae produced from Davies and Keppels had selective landscapes that were wider compared

to Esk larvae (**Fig. 6A-C**), although Davies and Esk had overall higher selection coefficients

compared to Keppels larvae (S= Esk: 60 –70, Davies: 54 to 70, Keppels 50 – 60; **Fig. 6A-C**).

Combined with estimated narrow-sense heritability estimates, these differences resulted in overall

higher selective responses (R) for Davies and Esk (i.e., more values between R = 40 – 60) compared

to Keppels offspring (**Fig. 6D-F**).

DISCUSSION

Mean maximum sea surface temperatures are expected to increase by between 2 – 4 °C by 2100

globally (IPCC, 2014). Without adaptation, this will likely exceed the thermal thresholds of corals.

Understanding the underlying adaptive capacity of wild populations is therefore critical to forecasting

species persistence. Moreover, various conservation strategies are being considered worldwide to help

corals withstand increasing ocean temperatures whilst carbon emissions are curbed (National

Academies of Sciences, Engineering, 2019). This includes the introduction of more heat tolerant

offspring produced from selective breeding methods onto cooler reefs to prepare them for warming

(Quigley et al., 2018), a method that will also increase the genetic diversity on reefs, fuel for which

natural selection can act. Quantifying the feasibility for enhancing corals' ability to survive further

ocean warming is therefore vital for the conservation of the world's coral reefs.

Little variation in adult physiological responses to heat stress across three GBR populations

Phenotypic variation in organisms drives the capacity for plastic, adaptive responses to environmental pressure. This variation may be underpinned by genetic variation or by responses mediated by non-genetic mechanisms, like changes in the microbiome (e.g. bacteria or Symbiodiniaceae), in which algal symbiont assemblages may shape corals' responses to heat stress (Parkinson et al., 2016). Understanding the scope for phenotypic variation to heat stress at the adult stage is essential to evaluating the scope for heritable diversity of heat tolerance at later life-stages in corals.

Overall, there was no difference in heat tolerance of adult southern Keppels corals compared to central Davies corals, with both populations suffering similar percent necrosis, drops in photophysiology, and lowered survival under heat stress. However, the relative change in bleaching was greater for Keppels corals. The magnitude of bleaching was also more severe in Davies compared to Keppels corals, suggesting that Davies's symbionts or the host may have lower initial tolerances but are able to recover and survive equally well. Taken together, these results suggest that the absolute heat tolerance of both populations was roughly equal. Finally, adult coral fragments from the Esk population in heat treatment exhibited the highest bleaching severity, lowest effective quantum yield, highest percentage necrosis, and lowest survivorship. It should be noted that although fragments sourced from the Esk population lost the greatest overall percentage of tissue (necrosis) per population, fragments in the control treatment were also highly necrotic, suggesting a compromised health state of corals from this population, also reflected in the low survival and photosynthesis in controls.

The high overall fitness of adult Keppels corals under heat stress was surprising. This population reported the lowest bleaching severity, highest effective quantum yield, and highest survival at heat. Although enhanced heat tolerance in corals is generally attributed to corals from warmer reefs exhibiting higher upper thermal thresholds (Berkelmans, 2002; Howells et al., 2012; Ulstrup et al., 2006), the enhanced performance of Keppels corals may be attributed to the greater variability in their local thermal regime. The increase in tolerance may also partly be attributed to the coral symbionts (Howells et al., 2012; Thomas et al., 2018; Ulstrup et al., 2006), and their interaction with host genetics (Dixon et al., 2015; Smith-Keune and van Oppen, 2006; Thomas et al., 2018), in which complex holobiont interactions influence the overall heat stress-responses via gene regulation,

symbiont density control and assemblage shuffling (Cunning and Baker, 2020; Yuyama et al., 2018).

Our results indicate that the control of heat tolerance is complex and that many factors, including local

thermal regime, likely play a role.

Minimal improvement in larval settlement due to selection for heat tolerance

Previous breeding experiments have demonstrated the transfer of increased offspring survival from

parents sourced from warm reefs when reproductively crossed with cooler reefs (Dixon et al., 2015).

However, it is currently unknown whether an increased propensity for settlement at high temperatures

is also transferable using colonies sourced from warmer reefs to achieve an enhancement in settlement

success. Although settlement is a heritable trait under control conditions ($h^2 = 0.49$; Meyer et al.,

2009), the over-all heritability is low relative to other fitness related traits. Moreover, it is well known

that settlement in corals is negatively impacted by heat. For example, *Diploria strigose* larvae

demonstrated a decrease in the settlement behaviour at temperatures exceeding 30 °C compared to

control (Bassim and Sammarco, 2003) and *Acropora palmata* settlement decreased by 25 % at 31.5

°C compared to 28 °C (Randall and Szmant, 2009). Combined, these previous results suggests that

selection should act on this important trait over time if oceans continue to warm.

During periods of increased temperature, the cellular processes within larvae become compromised,

including disruption in the repair of cellular proteins and enzymes (Negri et al., 2007). Consequently,

abnormalities develop, and the rate of cell cleavage rapidly increases. This cellular impairment could

prevent the transition from larvae to recruit by preventing attachment to substrata or an increase in

hypersensitivity during transition phases of larvae to polyp (Randall and Szmant, 2009). The

reduction in larval settlement could also be a consequence of energy deficiency. As cellular proteins

unfold and aggregate, HSP70, a heat shock protein that refolds degrading proteins (Daugaard et al.,

2007) has been found to be upregulated in *Acropora millepora* larvae to maintain normal cell function

(Rodriguez-Lanetty et al., 2009). This requires vast amounts of energy, that could potentially

otherwise be used for settlement. Higher respiration rates at heat also increase metabolic activity

(Edmunds et al., 2001) which, in turn, increases the amount of food required to maintain these

elevated levels. These factors could all contribute to the reduced settlement of larvae measured here.

Overall, although our results showed a significant decrease in larval settlement at heat compared to
control temperatures, this did not correspond to reef of origin. Our findings only weakly allude to the
potential for selectively bred coral larvae to settle at higher temperatures. For example, while our
results do not demonstrate a significant increase, there was a higher percentage of settled larvae
whose mother coral was from either Esk or Davies (the northern, warmer, inshore reefs in this study)
compared to those with a mother coral from the southern, cooler Keppels reef. There was also some
variation in settlement within population crosses under heat, demonstrating the potential for plasticity.
This aligns with information that corals found in warmer environments or with high daily temperature
variability have greater genetic plasticity (Kenkel and Matz, 2017) which can be passed onto
offspring. However, the findings presented here are preliminarily and it appears that the maximum
thermal limits of parental adult corals and the resulting larvae are not indicative of settlement success.
As warming increases in severity, assessing the impacts on processes like recruitment will become
increasingly important as they are essentially at risk of disruption (Radchuk et al., 2019).

The lack of a significant effect of parental colony from warmer reefs to enhance settlement at high
temperatures may be due to either experimental or biological factors. The high temperatures chosen
here may have surpassed the corals' settlement behaviour, in which temperatures exceeding 35.5 °C
were extreme compared to the mean monthly maximums of these sites (all >29 °C). Alternatively,
environmental factors may contribute a greater influence in determining settlement compared to host
genetics. Specifically, settlement deficiency at high temperature may be driven by the disruption of
the microbial biofilm needed to induce metamorphosis and settlement, where it is well established that
settlement is induced by the presence of CCA and microbial biofilms (Webster et al., 2004). During
periods of increased temperature, chemical cues released by CCA can be weakened and microbial
cells present in biofilms (on the CCA) can become damaged (Randall and Szmant, 2009). Therefore,
this reduction in biochemical cues could be the main contributing factor to the reduction in larval
settlement at higher temperatures.

In summary, larvae from intra- and interpopulation crosses demonstrated a general inability to settle at
high temperatures, suggesting that minimal improvement in larval settlement responses due to
selection for heat tolerance. Combined with information that behavioural or morphological traits
generally respond less to selection compared to life-history traits (Mousseau and Roff, 1987) and the
potential governing importance of environmental factors (e.g. bacterial communities), this suggests

that the enhancement of this trait under heat stress may require the selection through microbial

community contribution more than through processes targeting host genetics.

Predicting responses to heat using selection differentials and gradients of selection

Genetic variation underpins the potential and speed for adaptation through natural selection (Falconer

and Makcay, 1996). Warming influences traits differentially, with morphological traits generally less

impacted compared to phenological traits (Radchuk et al., 2019). Survival was chosen as the trait of

interest to examine gradients of selection given the importance of survival and other life-history traits

compared to behavioural or morphological traits (Mousseau and Roff, 1987). Measures such as

narrow-sense heritability (h^2), selection (S), and responses to selection (R) are useful for quantitively

assessing organisms' ability to respond to their environment, especially future stressors.

Measurements derived from corals suggest that they do have a strong underlying capacity to respond

adaptively to heat either through host genetics (h^2 = mean: 0.86 (range: 0.48 - 0.93) (Dixon et al.,

2015; Dziedzic et al., 2019; Kirk et al., 2018; Quigley et al., 2020b)) or through changes to their

symbiont communities (Quigley et al., 2018). It might be expected that heat stress would elicit

directional selection through differential mortality of adults, resulting in the survival of a subset of

phenotypes at one end of the phenotypic distribution. However, the bimodal responses in survival

curves of coral adults suggests that heat stress manifests as disruptive selection, which may explain

the high variability of heat responses across the numerous offspring lineages. Although the underlying

mechanisms is unknown here, the drivers of bimodality may be linked to biochemical complexity

(Rezende and Bozinovic, 2019).

Selection differentials depend on the heritability of the trait, where heritability is generally equal to

the slope of the Response over Selection, as determined by the breeder's equation. Interestingly, the

selection differentials at heat (i.e. the intensity of adaptive responses) were largely similar across the

three populations of adult corals. This largely mirrors the similar physiological responses of the adult

corals to heat stress. The selection differential between survival at high temperatures can be described

as the difference between the mean value measured (Davies and Keppels = ~40 - 30 %, and Esk ~25

%) and the desired mean value (e.g. ~90 %). Hence, the selection differential was about 55 % (90

minus 35 %) across these three populations, which represents a desired 63 % increase in potential trait

enhancement. When translated to estimated adaptive larval responses, these varied by population

cross. Therefore, our comparative analysis between adult and offspring responses suggests a

divergence between adult population mean responses in adults and interpopulation offspring crosses.

The divergence between thermal responses of different life-stages within species has been

demonstrated in other marine organisms like mollusks (Truebano et al., 2018), in which larval forms
are often more or less vulnerable to heat compared to adults. Finally, adaptive response surfaces
revelated that when larvae were grouped by their maternal populations, Davies and Esk offspring
generally had higher selective responses (R) compared to Keppels offspring. Taken together, this
suggests that although adult populations may respond similarly to heat, the overall potential of
offspring responses to selection in warmer populations of corals from Davies and Esk is greater
compared to cooler populations in the Keppels. These findings have important implications for
forecasting the impacts of climate change on wild populations of corals and for the development of
novel conservation tools like the assisted evolution of at-risk populations.

Conclusions

620 As climate change accelerates ecosystem change, critical information on how important fitness traits
621 will vary in the future is essential to move from understanding impacts to predicting and forecasting
622 those impacts. This comparative physiological dataset across different life-history stages in one
623 important coral species provides key mechanistic and adaptive insights into how corals may function
624 under heat stress caused by warming oceans, and in particular, the heritability of heat tolerance.

Acknowledgements

The authors would like to thank the Traditional Owners from whose Sea Country these corals were
collected. Specifically, we would like to thank the Woppaburra, Manbarra, Bindal, and Wulgurukaba
Traditional Owners. Corals were collected under the following permit numbers: G12/35236.1 (Davies
and Esk) and G19/43024.1 (Keppels) to the Australian Institute of Marine Science. We would like to
thank Andrea Severati, Line Bay, Christine Giuliano, and the spawning collections field crews in the
central and southern Great Barrier Reef for assisting in the coral colony collections.

Footnotes

Author contributions

Conceptualization and Methodology: K.M.Q.; Formal analysis: P.W.; K.M.Q.; Investigation: P.W.,
R.C., A.M., H.K., J.R., K.M.Q; Resources: K.M.Q.; Data curation: P.W., K.M.Q. Writing - original

draft: K.M.Q., R. C., J.R.; Writing - review & editing: P.W., R.C., A.M., K.M.Q.; Supervision:

K.M.Q.; Funding acquisition: K.M.Q.

Funding

This study was supported by funding from the Australian Institute of Marine Science. Ponchanok

Weeriyanun was supported by a travel and study grant provided by Ghent University, Erasmus,

Master of Science in Marine Biological Resources (IMBRSea).

Competing interests

647   The authors declare no competing or financial interests.

Data availability

Physiological data and analysis code is available at https://github.com/LaserKate/KeppelsAGF19

References

AIMS (2020). *Initial recovery of the Great Barrier Reef threatened by the third mass bleaching event*

in five years.

Aitken, S. N. and Whitlock, M. C. (2013). Assisted gene flow to facilitate local adaptation to climate

change. *Annu. Rev. Ecol. Evol. Syst.* **44**, 367–388.

Bassim, K. M. and Sammarco, P. W. (2003). Effects of temperature and ammonium on larval

development and survivorship in a scleractinian coral (*Diploria strigosa*). *Mar. Biol.* **142**, 241–

252.

Bassim, K. M., Sammarco, A. P. W. and Snell, A. T. L. (2002). Effects of temperature on success

of (self and non-self) fertilization and embryogenesis in *Diploria strigosa* (Cnidaria,

Scleractinia). *Mar. Biol.* **140**, 479–488.

Bates, D. (2005). Fitting linear mixed models in R. *Obes. Facts* **3**, 27–30.

Baums, I. B., Baker, A. C., Davies, S. W., Grottoli, A. G., Kenkel, C. D., Kitchen, S. A., Kuffner, I. B., LaJeunesse, T. C., Matz, M. V., Miller, M. W., et al. (2019). Considerations for maximizing the adaptive potential of restored coral populations in the western Atlantic. *Ecol. Appl.* **29**, 1–23.

Berkelmans, R. (2002). Time-integrated thermal bleaching thresholds of reefs and their variation on the Great Barrier Reef. *Mar. Ecol. Prog. Ser.* **237**, 309–310.

Cunning, R. and Baker, A. C. (2020). Thermotolerant coral symbionts modulate heat stress-responsive genes in their hosts. *Mol. Ecol.* **29**, 2940–2950.

Daugaard, M., Rohde, M. and Jäättelä, M. (2007). The heat shock protein 70 family: Highly homologous proteins with overlapping and distinct functions. *FEBS Lett.* **581**, 3702–3710.

De'Ath, G., Fabricius, K. E., Sweatman, H. and Puotinen, M. (2012). The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 17995–17999.

Dixon, G., Davies, S., Aglyamova, G., Meyer, E., Bay, L. and Matz, M. (2015). Genomic determinants of coral heat tolerance across latitudes. *Science (80-.).* **348**, 1460–1462.

Doropoulos, C., Gómez-Lemos, L. A. and Babcock, R. C. (2018). Exploring variable patterns of density-dependent larval settlement among corals with distinct and shared functional traits. *Coral Reefs* **37**, 25–29.

Dziedzic, K. E., Elder, H., Tavalire, H. and Meyer, E. (2019). Heritable variation in bleaching responses and its functional genomic basis in reef-building corals (*Orbicella faveolata*). *Mol. Ecol.* **28**, 2238–2253.

Edmunds, P., Gates, R. and Gleason, D. (2001). The biology of larvae from the reef coral *Porites astreoides*, and their response to temperature disturbances. *Mar. Biol.* **139**, 981–989.

Falconer, D. S. and Makcay, T. F. C. (1996). *Introduction to Quantitative Genetics*. 4th Editio. London, UK.: Longman Group Ltd.

Hadley Wickham, Romain François, Lionel Henry and Kirill Müller (2020). A grammar of data manipulation [R package dplyr version 1.0.0]. *Media*.

Heron, S. F., Maynard, J. A., van Hooidonk, R. and Eakin, C. M. (2016). Warming Trends and Bleaching Stress of the World's Coral Reefs 1985-2012. *Sci. Rep.* **6**, 1–14.

Heyward, A. J. and Negri, A. P. (2010). Plasticity of larval pre-competency in response to
temperature: Observations on multiple broadcast spawning coral species. *Coral Reefs* **29**, 631–
636.

Hoegh-Guldberg, O. (1999). Climate change, coral bleaching and the future of the world's coral
reefs. *Mar. Freshw. Res.* **50**, 839–866.

Howells, E. J., Beltran, V. H., Larsen, N. W., Bay, L. K., Willis, B. L. and van Oppen, M. J. H.
(2012). Coral thermal tolerance shaped by local adaptation of photosymbionts. *Nat. Clim.*
Chang. **2**, 116–120.

Hughes, T. P., Barnes, M. L., Bellwood, D. R., Cinner, J. E., Cumming, G. S., Jackson, J. B. C.,
Kleypas, J., van de Leemput, I. A., Lough, J. M., Morrison, T. H., et al. (2017). Coral reefs
in the Anthropocene. *Nature* **546**, 82–90.

IPCC (2014). *Climate Change 2014: Synthesis Report*.

Jones, G. P., Mccormick, M. I., Srinivasan, M. and Eagle, J. V (2004). Coral decline threatens fish
biodiversity in marine reserves. *Popul. Biol.* **101**, 8251–8253.

Kassambaraype, A. (2020). ggplot2' Based Publication Ready Plots Version.

Keith, S. A., Maynard, J. A., Edwards, A. J., Guest, J. R., Bauman, A. G., van Hooidonk, R.,
Heron, S. F., Berumen, M. L., Bouwmeester, J., Piromvaragorn, S., et al. (2016). Coral mass
spawning predicted by rapid seasonal rise in ocean temperature. *Proc. R. Soc. B Biol. Sci.* **283**,.

Kenkel, C. D. and Matz, M. V. (2017). Gene expression plasticity as a mechanism of coral
adaptation to a variable environment. *Nat. Ecol. Evol.* **1**, 1–6.

Kenkel, C. D., Almanza, A. T. and Matz, M. V. (2015). Fine-scale environmental specialization of
reef-building corals might be limiting reef recovery in the Florida Keys. *Ecology* **96**, 3197–3212.

Kirk, N. L., Howells, E. J., Abrego, D., Burt, J. A. and Meyer, E. (2018). Genomic and
transcriptomic signals of thermal tolerance in heat-tolerant corals (*Platygyra daedalea*) of the
Arabian/Persian Gulf. *Mol. Ecol.* **27**, 5180–5194.

Komyakova, V., Munday, P. L. and Jones, G. P. (2013). Relative importance of coral cover, habitat
complexity and diversity in determining the structure of reef fish communities. *PLoS One* **8**,
e83178.

Matz, M. V., Treml, E. A., Aglyamova, G. V. and Bay, L. K. (2017). Potential and limits for rapid

genetic adaptation to warming in a Great Barrier Reef coral. *PLOS Genet.* **14**, 1–19.

Matz, M. V., Treml, E. A., Aglyamova, G. V., van Oppen, M. J. H. and Bay, L. K. (2018).
Potential for rapid genetic adaptation to warming in a Great Barrier Reef coral. *bioRxiv* 1–19.

Meyer, E., Davies, S., Wang, S., Willis, B. L., Abrego, D., Juenger, T. E. and Matz, M. V. (2009).
Genetic variation in responses to a settlement cue and elevated temperature in the reef-building
coral *Acropora millepora*. *Mar. Ecol. Prog. Ser.* **392**, 81–92.

Moghaddam, S., Shokri, M. R. and Tohidfar, M. (2021). The enhanced expression of heat stress-
related genes in scleractinian coral ' *Porites harrisoni* ' during warm episodes as an intrinsic
mechanism for adaptation in 'the Persian Gulf.' *Coral Reefs* 1–16.

Mousseau, T. A. and Roff, D. A. (1987). Natural selection and the heritability of fitness components.
Heredity (Edinb). **59**, 181–197.

National Academies of Sciences, Engineering, and M. (2019). *A research review of interventions*
to increase the persistence and resilience of coral reefs. Washington, DC: The National
Academies Press.

Negri, A. P., Marshall, P. A. and Heyward, A. J. (2007). Differing effects of thermal stress on coral
fertilization and early embryogenesis in four Indo Pacific species. *Coral Reefs* **26**, 759–763.

Nelson, H. R., Kuempel, C. D. and Altieri, A. H. (2016). The resilience of reef invertebrate
biodiversity to coral mortality. *Ecosphere* **7**, e01399.

O'Mahony, J., Simes, R., Redhill, D., Heaton, K., Atkinson, C., Hayward, E. and Nguyen, M.
(2017). At what price? The economic, social and icon value of the Great Barrier Reef. *Deloitte*
Access Econ. 92.

Parkinson, J. E., Baumgarten, S., Michell, C. T., Baums, I. B., LaJeunesse, T. C. and Voolstra,
C. R. (2016). Gene expression variation resolves species and individual strains among coral-
associated dinoflagellates within the genus symbiodinium. *Genome Biol. Evol.* **8**, 665–680.

Payne, N. L., Morley, S. A., Halsey, L. G., Smith, J. A., Stuart-Smith, R., Waldock, C. and
Bates, A. E. (2021). Fish heating tolerance scales similarly across individual physiology and
populations. *Commun. Biol.* **4**, 1–5.

Plotly Technologies Inc. (2015). Collaborative data science.

Quigley, K. M., Bay, L. K. and Willis, B. L. (2018). Leveraging new knowledge of *Symbiodinium*

community regulation in corals for conservation and reef restoration. *Mar. Ecol. Prog. Ser.* **600**,
245–253.

Quigley, K. M., Bay, L. K. and van Oppen, M. J. H. (2019a). The active spread of adaptive
variation for reef resilience. *Ecol. Evol.* **9**, 1–14.

Quigley, K. M., Willis, B. L. and Kenkel, C. D. (2019b). Transgenerational inheritance of shuffled
symbiont communities in the coral Montipora digitata. *Sci. Rep.* **9**, 1–11.

Quigley, K. M., Bay, L. K. and van Oppen, M. J. H. (2020a). Genome-wide SNP analysis reveals
an increase in adaptive genetic variation through selective breeding of coral. *Mol. Ecol.* **29**,
2176–2188.

Quigley, K. M., Randall, C. J., van Oppen, M. J. H. and Bay, L. K. (2020b). Assessing the role of
historical temperature regime and algal symbionts on the heat tolerance of coral juveniles. *Biol.*
Open **9**, 1–11.

R Core Team (2021). R: A Language and Environment for Statistical Computing. *R Found. Stat.*
Comput. https://www.R-project.org.

Radchuk, V., Reed, T., Teplitsky, C., van de Pol, M., Charmantier, A., Hassall, C., Adamík, P.,
Adriaensen, F., Ahola, M. P., Arcese, P., et al. (2019). Adaptive responses of animals to
climate change are most likely insufficient. *Nat. Commun.* **10**, 1–14.

Randall, C. J. and Szmant, A. M. (2009). Elevated temperature reduces survivorship and settlement
of the larvae of the Caribbean scleractinian coral, *Favia fragum* (Esper). *Coral Reefs* **28**, 537–
545.

Rezende, E. L. and Bozinovic, F. (2019). Thermal performance across levels of biological
organization. *Philos. Trans. R. Soc. B Biol. Sci.* **374**,.

Rodriguez-Lanetty, M., Harii, S. and Hoegh-Guldberg, O. (2009). Early molecular responses of
coral larvae to hyperthermal stress. *Mol. Ecol.* **18**, 5101–5114.

Savary, R., Barshis, D. J., Voolstra, C. R., Cárdenas, A., Evensen, N. R., Banc-Prandi, G., Fine,
M. and Meibom, A. (2021). Fast and pervasive transcriptomic resilience and acclimation of
extremely heat-tolerant coral holobionts from the northern Red Sea. *Proc. Natl. Acad. Sci.* **118**,
1–9.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of

image analysis. *Nat. Methods* **9**, 671–675.

Schoepf, V., Carrion, S. A., Pfeifer, S. M., Naugle, M., Dugal, L., Bruyn, J. and McCulloch, M.
T. (2019). Stress-resistant corals may not acclimatize to ocean warming but maintain heat
tolerance under cooler temperatures. *Nat. Commun.* **10**,.

Siebeck, U. E., Logan, D. and Marshall, N. J. (2008). CoralWatch - a flexible coral bleaching
monitoring tool for you and your group. *Proc. 11th Int. Coral Reef Symp.* 16.

Smith-Keune, C. and van Oppen, M. (2006). Genetic structure of a reef-building coral from
thermally distinct environments on the Great Barrier Reef. *Coral Reefs* **25**,.

Sully, S., Burkepile, D. E., Donovan, M. K., Hodgson, G. and van Woesik, R. (2019). A global
analysis of coral bleaching over the past two decades. *Nat. Commun.* **10**, 1–5.

Thomas, L., Rose, N. H., Bay, R. A., López, E. H., Morikawa, M. K., Ruiz-Jones, L. and
Palumbi, S. R. (2018). Mechanisms of Thermal Tolerance in Reef-Building Corals across a
Fine-Grained Environmental Mosaic: Lessons from Ofu, American Samoa. *Front. Mar. Sci.* **4**,
434.

Truebano, M., Fenner, P., Tills, O., Rundle, S. D. and Rezende, E. L. (2018). Thermal strategies
vary with life history stage. *J. Exp. Biol.* **221**,.

Tukey, J. W. (1977). *Exploratory data analysis*. Reading, Mass.

Ulstrup, K. E., Berkelmans, R., Ralph, P. J. and van Oppen, M. J. H. (2006). Variation in
bleaching sensitivity of two coral species across a latitudinal gradient on the Great Barrier Reef:
The role of zooxanthellae. *Mar. Ecol. Prog. Ser.* **314**, 135–148.

van Oppen, M. J. H., Oliver, J. K., Putnam, H. M. and Gates, R. D. (2015). Building coral reef
resilience through assisted evolution. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 2307–2313.

Walz, H. (2018). DIVING-PAM-II. **49**,.

Webster, N. S., Smith, L. D., Heyward, A. J., Watts, J. E. M., Webb, R. I., Blackall, L. L. and
Negri, A. P. (2004). Metamorphosis of a scleractinian coral in response to microbial biofilms.
Appl. Environ. Microbiol. **70**, 1213–1221.

Wickham, H. (2011). The split-apply-combine strategy for data analysis. *J. Stat. Softw.* **40**, 1–29.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wilcoxon, F. (1945). Individual Comparisons by Ranking Methods. *Biometrics Bull.* **1**, 80–83.

Yuyama, I., Ishikawa, M., Nozawa, M., Yoshida, M. and Ikeo, K. (2018). Transcriptomic changes
with increasing algal symbiont reveal the detailed process underlying establishment of coral-
algal symbiosis. *Sci. Rep.* **8**, 16802.

822   **Figures**

Figure 1. Summary of experimental design for determining variability in heat tolerance across coral life-history stages. Map of the Great Barrier Reef showing the three sites of adult coral collection (A), including Esk (central inshore, black), Davies (central mid-shelf, tan) and Keppels (southern inshore, maroon) reefs (A). Mean monthly sea surface temperature profiles at each reef of adult coral collection (B). A graphic showing when the three life-stages were experimentally exposed to heat stress (C). Summary of the population crosses produced from the three coral populations. Filled circles represent intrapopulation crosses, whereas open circles represent interpopulation crosses (D). Figure indicating the ramping time to reach each experimental temperature and the length that it was held (E).

Figure 2. Larval survival under experimental stress. Percentage survival over time of n =85
different *Acropora tenuis* lineages of larvae exposed to control (27.5ºC) and heat stress (35.5 ºC) (A).
The coloured boxes around the crosses indicate the source reef of the maternal colonies: Davies (tan),
Esk (black), and Keppels (maroon) (B).

 Figure 3. Larval settlement under experiential stress. Percentage of median settlement of n = 69

different *Acropora tenuis* lineages of larvae exposed to control (blue) or heat stress (red) conditions

after 17 h (A), 24 h (B) and 48 h (C). The coloured boxes around the crosses indicate the source reef

of the maternal colonies: Davies (tan), Esk (black), and Keppels (maroon).

Figure 4. **Median physiological responses in adult corals exposed to control and heat stress**

temperatures: Bleaching category core (A), Effective quantum yield (B), percent necrosis (C), and

percent survival (D) of adult *Acropora tenuis* fragments. Colonies were collected from three sites on

the Great Barrier Reef: Davies reef (tan outlines in A-D), Esk reef (black) and Keppels reef (maroon).

Mean percent change in each response by latitude of collection site (E).

Figure 5. Percent survival density plot of the number of adult colonies (y) from each population

(A) and offspring of each population cross (B-D). Arrows indicated the selected phenotypic scope.

Horizontal dashed lines represent the selection differential for each population. Vertical dashed lines

represent mean percent survival per population. The coloured boxes around the crosses indicate the

source reef of the maternal colonies: Davies (tan), Esk (black), and Keppels (maroon).

Figure 6. Adaptive landscape of selection for heat tolerance. Modelled selective values (S) and

narrow-sense heritability (h^2) values of larval survival at heat grouped by the population identity of

the maternal colony (A-C). Estimated response to selection (R) for h^2 and S (D-F).