BREAKTHROUGH REPORT

Spatial differences in stoichiometry of EGR phosphatase and Microtubule-Associated
Stress Protein 1 control root meristem activity during drought stress.

Toshisangba Longkumer[a], Chih-Yun Chen[a,1], Marco Biancucci[a,2], Govinal Badiger
Bhaskara[a,3], Paul E. Verslues[a,b]

a. Institute of Plant and Microbial Biology, Academia Sinica, Taipei 11528, Taiwan
1. Present Address: Biotechnology Center, National Chung Hsing University
Taichung 40227, Taiwan
2. Present Address: Department of Biosciences, University of Milan, Milan 20133, Italy
3. Present Address: Department of Integrative Biology, University of Texas, Austin, TX
78712

b. Corresponding Author: paulv@gate.sinica.edu.tw

Short Title: EGR-MASP1 control of meristem activity

One Sentence Summary: Spatial differences in EGR-MASP1 expression and control of
MASP1 phosphorylation adjust root meristem activity to regulate growth during drought
stress.

Abstract

During moderate severity drought and low water potential (ψ_w) stress, poorly understood
signaling mechanisms restrict both meristem cell division and subsequent cell expansion. We
found that the Clade E Growth-Regulating 2 (*EGR2*) protein phosphatase and *Microtubule*
Associated Stress Protein 1 (MASP1) differed in their stoichiometry of expression across the
root meristem and had opposing effects on root meristem activity at low ψ_w. Ectopic MASP1
or EGR expression increased or decreased, respectively, root meristem size and root
elongation during low ψ_w stress. This, along with the ability of phosphomimic MASP1 to
overcome EGR suppression of root meristem size and observation that ectopic EGR
expression had no effect on unstressed plants, indicated that during low ψ_w EGR activation
and attenuation of MASP1 phosphorylation in their overlapping zone of expression
determines root meristem size and activity. Ectopic EGR expression also decreased root cell
size at low ψ_w. Conversely, both the *egr1-1egr2-1* and *egr1-1egr2-1masp1-1* mutants had
similarly increased root cell size; but, only *egr1-1egr2-1* had increased cell division. These
observations demonstrated that EGRs affect meristem activity via MASP1 but affect cell
expansion via other mechanisms. Interestingly, EGR2 was highly expressed in the root
cortex, a cell type important for growth regulation and environmental response.

Even slight or moderate reduction of water availability (quantified as a reduction in

water potential, ψ_w) leads plants to actively restrict their growth. This can be an adaptive

mechanism to conserve soil water during drought periods. However, such negative

regulation of growth can also restrict plant biomass accumulation more than is needed or

desired in agricultural settings (Skirycz and Inzé, 2010; Verslues, 2017). A better

understanding of the mechanisms that regulate plant growth in response to altered water

status could allow growth inhibition to be circumvented when desirable (Skirycz and Inzé,

2010; Claeys et al., 2014).

Root growth is both a key factor in drought resistance and a model for developmental

study because of the relatively simple organization of the root tip. At the root tip, the

quiescent center (QC) and immediately adjacent cells act as a stem cell niche that produces

cells to generate both the root meristem and root cap (Heidstra and Sabatini, 2014). Just

behind the QC and adjacent cells is the meristematic region which supplies the bulk of new

cells during root elongation. Cells in the meristem divide rapidly until they are displaced

from the root tip, lose their cell division activity, and begin rapid expansion. The rate of root

elongation is determined by both cell expansion and the amount of cell production by the

meristem. Meristem cell production is in turn determined by the rate of cell division (cell

cycle time) as well as the duration of time over which cell division continues as cells are

displaced from the root tip. A longer duration of cell division increases the size of the root

meristem (Beemster and Baskin, 1998).

While the root tip can be anatomically divided into distinct zones of cell division and

cell expansion, the size of each zone may be controlled by underlying gradients of key

regulatory factors (Barlow, 1976). Gradients in hormone content and signaling, particularly

auxin, brassinosteroid and cytokinin, as well as reactive oxygen are important regulators of

meristem size and function (Chaiwanon and Wang, 2015; Vragovic et al., 2015; Di Mambro

et al., 2017; Salvi et al., 2020; Yamada et al., 2020). It is also known that hormone-related

regulatory proteins and transcription factors have gradients of expression across the root

meristem or are specifically expressed in the QC and surrounding stem cell area (Heidstra

and Sabatini, 2014; Salvi et al., 2020; Yamada et al., 2020). Many additional genes have

gradients of increasing or decreasing expression across the root meristem and some of these

lesser characterized genes also affect root growth (Wendrich et al., 2017; Dubois et al., 2018;

Slovak et al., 2020). It has been proposed that the balance between opposing gradients of

pro-cell division genes, which are most highly expressed in the proximal meristem region

close to the QC, versus pro-cell expansion genes, which are highly expressed at the distal end of the meristem and in the expansion zone, determines when cells stop dividing and begin rapid expansion (Salvi et al., 2020). Such opposing gradients of expression suggests that, at different positions within the root meristem, the stoichiometry (ratio) of the proteins will vary. Such differences in protein stoichiometry may impart differing outcomes in terms of allowing continued cell division or inhibiting cell division and promoting cell expansion (Wendrich et al., 2017). While appealing, this idea remains largely unproven as the function of most gradient-expressed genes, and whether they produce different levels of protein at different positions in the root meristem, is unclear. It is also unclear how proteins having spatial differences in stoichiometry across the meristem can affect each other's function at the molecular level.

Low ψ_w during drought stress leads to both decreased cell production and decreased cell expansion (Skirycz and Inzé, 2010; Skirycz et al., 2011). Which of these is the predominant cause of reduced growth can vary between different plants and between different types of abiotic stress. In Arabidopsis roots, low ψ_w (-1.28 MPa), imposed using PEG-infused agar plates, led to a nearly 50% decrease in root cell production rate but only minor reduction of cell expansion (van der Weele et al., 2000). Growth kinematic analysis has also shown that drought stress decreases cell production in the root meristem (Sacks et al., 1997; Voothuluru et al., 2020). These observations suggest that stress signaling can interact with cell cycle and developmental regulation to control meristem activity. Consistent with this idea, several types of abiotic stress elicit root cell type-specific responses (Dinneny et al., 2008; Iyer-Pascuzzi et al., 2011) and development regulators can also influence stress response genes (Moubayidin et al., 2013). Despite these indications, relatively little is known of how stress signaling impinges upon developmental mechanisms to control meristem activity and regulate growth (Shimotohno and Scheres, 2019).

Previous work in our laboratory found that Clade E Growth Regulating (EGR) type 2C protein phosphatases (PP2Cs) act as negative regulators of growth during low ψ_w (Bhaskara et al., 2017). Mutants of *EGR1*, *EGR2*, and *EGR3* all had higher than wild type growth (quantified based on fresh weight, dry weight and primary root length increase) and an *egr1-1egr2-1* double mutant had a stronger effect, consistent with EGR1 and EGR2 having redundant function in growth regulation during low ψ_w. This increased growth was observed in both polyethylene glycol (PEG)-agar plate assays and in controlled soil drying

experiments where plants were exposed to moderate severity water limitation.

Phosphoproteomic analysis of *egr1-1egr2-1* indicated that EGRs primarily affect

phosphorylation of plasma membrane, trafficking and cytoskeleton-associated proteins. One

of the EGR-regulated proteins identified by phosphoproteomics was a protein of unknown

function which we designated as Microtubule Associated Stress Protein 1 (MASP1). The

EGR phosphatases interacted with MASP1 (in Bi-molecular Fluorescence Complementation

and co-immunoprecipitation assays) and we could validate (using Phostag gel analysis) that

during low ψ_w stress MASP1 was more phosphorylated in *egr1-1egr2-1* but less

phosphorylated in transgenic plants ectopically expressing EGR1 (*35S:EGR1*). MASP1

promoted growth during low ψ_w in both PEG-agar plate and soil drying assays. This growth

promotion activity was dependent upon MASP1 serine 670 phosphorylation (Bhaskara et al.,

2017). Consistent with EGR regulation of MASP1 phosphorylation, genetic analysis

indicated that MASP1 acted downstream of EGRs; although, the phenotype of the *egr1-*

1egr2-1masp1-1 triple mutant was somewhat intermediate between that of *egr1-1egr2-1* and

masp1-1. MASP1 bound microtubules *in vitro*. MASP1 and EGRs had converse effects on

microtubule stability *in planta* which correlated with MASP1 S670 phosphorylation status

and with growth at low ψ_w: more stable microtubules were associated with phosphorylated

MASP1 and enhanced growth maintenance at low ψ_w. Other than this, nothing is known

about MASP1 cellular function. The only MASP1-related protein which has been studied is

Auxin-Induced in Root Cultures 9 (AIR9), which binds microtubules and localizes at the

preprophase band during cell division (Buschmann et al., 2015). However, AIR9 is

substantially different from MASP1 in that its microtubule binding region is at the N-terminal

portion of the protein rather than the C-terminus as in MASP1 and the remainder of AIR9

outside of the N-terminal LRR-repeat region is divergent from, and much larger than,

MASP1. We have not observed MASP1 localization on spindle fibers or newly forming cell

plate and the cellular function of MASP1 remains unclear.

Interestingly, it has been reported that *EGR*s have a gradient of expression across the

root meristem. Sorting of root meristem cells into proximal (close to QC), medial, and distal

regions based on GFP expression driven by the *Plant U-Box25* (*PUB25*) or *SPATULA* (*SPT*)

promoters found that *EGR1, EGR2* and *EGR3* had low expression in the proximal meristem

region but significantly higher expression in medial and distal meristem regions (Wendrich et

al., 2017). While this is an intriguing observation, it is not known whether such gradient in

EGR gene expression also leads to differences in EGR protein level in the proximal versus

distal meristem regions and whether spatial differences in EGR protein level are functionally

important for growth regulation.

Here we demonstrate that differences in EGR-MASP1 stoichiometry in different

regions of the root meristem controls root meristem size and activity during low ψ_w. These

results show that EGRs and MASP1 link stress signaling with regulation of root meristem

function during drought stress and show how disruption of EGR-MASP1 stoichiometry and

signaling can enhance growth maintenance, or further downregulate growth, during moderate

severity low ψ_w. We also found that EGR2 was most highly expressed in root cortex cells, an

observation that is interesting in light of recent data indicating a key role of the cortex in

regulating growth responses to low ψ_w.

Results

EGRs and MASP1 have opposing effects on root elongation during low ψ_w.

To further determine how EGRs and MASP1 affect root growth responses to low ψ_w, we transferred 5-day-old seedlings to PEG-agar plates of moderate severity low ψ_w stress (-0.7 MPa). This ψ_w was selected as we have previously shown that it reduces growth by a moderate amount (50-70 percent) in both PEG-agar plate assays and soil drying experiments (Bhaskara et al., 2017; Wong et al., 2019) and is within the range of soil ψ_w that occurs during moderate severity drought in many types of field environments. In these experiments, it was visually clear that *egr1-1egr2-1* and *35S:MASP1* maintained higher root and shoot growth compared to wild type at low ψ_w (Fig. 1A), consistent with previous results where whole seedling fresh and dry weights were quantified (Bhaskara et al., 2017). Quantitation of root elongation rates found that in wild type root elongation rate was reduced by nearly 70 percent compared to the unstressed control over six days after transfer to low ψ_w (Fig. 1B). The *egr1-1egr2-1* double mutant maintained significantly higher root elongation rates than wild type at low ψ_w over this period (Fig. 1B). We also checked root elongation rates of *egr* single mutants and found that there was a trend of increased root elongation at low ψ_w for *egr1* and *egr2* mutants (but the difference was only significant for *egr2-1*) while *egr3* mutants had similar or lesser effect (Supplemental Fig. S1A). Together, these data confirmed that *EGR* phosphatases act redundantly to restrict root elongation during low ψ_w stress.

Ectopic *MASP1* expression (*35S:MASP1*) led to higher root elongation rate at low ψ_w compared to wild type while *masp1-1* and *35S:EGR1* had reduced root elongation rate (Fig. 1B). This also was consistent with previous quantitation of seedling dry weights in *masp1-1* (and *masp1-2*) as well as *35S:EGR1* which found that all these genotypes had decreased growth at low ψ_w but had little effect on growth in the unstressed control (Bhaskara et al., 2017). For the analyses presented in Fig. 1 we used transgenic lines with expression of untagged EGR1 or MASP1 and lines with expression of either protein with an N-terminal fusion to YFP. Transgenic lines used in this study are listed in Supplemental Table S1 and additional data showing that individual transgenic lines have similar effect on root elongation is shown in Supplemental Fig. S2. Note that *35S:EGR1* and *35S:MASP1*, as well as constructs with an N-terminal YFP added to either protein, have been previously shown to complement their respective mutants (Bhaskara et al., 2017). Lines expressing either tagged or untagged EGR1 and MASP1 were used in this study to ensure that our results were robust

Figure 1: EGRs and MASP1 affect growth during low ψ_w stress.

A. Representative seedlings of *EGR* and *MASP1* mutants and overexpression lines. Five-day-old seedlings were transferred to either fresh control plates (-0.25 MPa) or PEG-infused agar plates for moderate severity low ψ_w (-0.7 MPa) treatment. Plants were photographed at the end of the experiment (six days after transfer). Scale bars indicate 1 cm.

B. Primary root elongation rates over six days after transfer of seedlings to control or stress (-0.7 MPa) treatments. Blue or red symbols show elongation rates for individual seedlings. Black bars and error bars indicate the mean and standard deviation for each genotype (n = 22 to 67). Lower case letters in the stress panel indicate statistical differences (ANOVA with Tukey's post hoc test, corrected P ≤ 0.05). Genotypes sharing the same letter did not significantly differ from one another. Data are combined from three independent experiments and three independent transgenic lines per construct. Both *35S:EGR1* and *35S:YFP-EGR1* as well as *35S:MASP1* and *35S:YFP-MASP1* lines were used. Data from additional experiments with each individual transgenic line is shown in Supplemental Fig. S2. The lines expressing untagged or YFP-tagged proteins had consistent result and the combined data are labeled as *35S:MASP1* and *35S:EGR1* for clarity of presentation.

across multiple independent transgenic lines and as an extra check that the presence of the

YFP-tag did not influence the results so that later experiments examining the expression

patterns of YFP-tagged EGR1 and MASP1 could be robustly compared to the results of

physiology experiments. As all EGR1 and MASP1 transgenic lines used in this study had

consistent phenotypes, for subsequent experiments combined data of multiple transgenic lines

is presented in the main text figures and the lines referred to as *35S:EGR1* and *35S:MASP1*

for clarity in presentation while data for individual transgenic lines are presented in

Supplemental Figures.

Interestingly, the *egr1-1egr2-1masp1-1* triple mutant was intermediate between *egr1-*

1egr2-1 and *masp1-1* and did not significantly differ from wild type at low ψ_w (Fig. 1B).

This intermediate phenotype of *egr1-1egr2-1masp1-1* was consistent with our previous

observations and our interpretation that MASP1 acts downstream of EGRs; but, EGRs also

have MASP1-independent effect(s) on growth (Bhaskara et al., 2017).

We also measured growth of EGR-MASP1 mutants and transgenic lines after transfer

to media containing NaCl. Salt stress has both an osmotic component and a sodium toxicity

component (Munns and Tester, 2008). To avoid excessive sodium toxicity which may

obscure the osmotic stress response, we used relatively mild salt stress treatments (75 mM

and 125 mM) which reduced root elongation rate of wild type by 30 percent and 50 percent,

respectively (Supplemental Fig. S3A). These experiments showed that the EGR-MASP1

mutant and transgenic lines exhibited similar differences in root elongation during salt stress

as during low ψ_w, albeit that the effect sizes were smaller because root elongation rates of all

genotypes remained relatively high. At 75 mM NaCl, *egr1-1egr2-1* and *35S:MASP1* had

significantly higher root elongation rates than the contrasting *masp1-1* and *35S:EGR1*

genotypes, respectively. At 125 mM NaCl, *egr1-1egr2-1* and *35S:MASP1* had higher root

elongation rates than wild type or other genotypes. Along with previous data showing that

EGRs and MASP1 affect rosette growth during soil drying (Bhaskara et al., 2017), these data

demonstrate that the effects of EGR and MASP1 on growth can be observed in multiple

treatments that restrict water availability.

MASP1 promotes cell division during low ψ_w and is required for the increased cell

division of *egr1-1egr2-1*.

The effect of EGR and MASP1 on root growth at low ψ_w could be caused by a change

in the number of cells produced by cell division, change in cell size or a combination of the

two. To analyze cell division, we first used a reporter line containing β-glucuronidase

(GUS) fused to the Cyclin B1 promoter and cyclin destruction box (CYCB1::GUS) (Colon-

Carmona et al., 1999). This reporter produces foci of GUS activity when a cell goes through

the G2-M phase transition thus allowing mitotic activity to be estimated by counting the

number GUS foci. In wild type, low ψ_w reduced the number of CYCB1::GUS foci by more

Figure 2: EGRs and MASP1 affect root cell division and cell expansion during low ψ_w stress.

  A. Representative images of CYCB1::GUS staining in primary root tips of the indicated mutant and ectopic expression genotypes 4 days after transfer of five-day-old seedlings to either control or low ψ_w (-0.7 MPa). Scale bars indicate 100 μm.

  B. Quantification of CYCB1::GUS foci in the primary root meristem. Symbols indicate counts of GUS foci from individual roots while black bars and error bars indicate the mean and standard deviation for each genotype. Within the control or stress treatments, genotypes sharing the same letter do not significantly differ from one another (ANOVA with Tukey's post hoc test, corrected P ≤ 0.05). In wild type, the number of CYCB1::GUS foci was significantly different between control and stress treatment (T-test, P ≤ 0.001). Data are combined from three or four independent experiments (n = 21-66) for each genotype and at least two independent transgenic lines for *35S:MASP1* and *35S:EGR1*. Data plotted separately for each transgenic line is shown in Supplemental Fig. S4.

  C. Cell length at 2 mm from the root tip measured four days after transfer of mutant or ectopic expression seedlings to control or low ψ_w (-0.7 MPa). Format of data and statistical analysis are the same as in B (n = 133-224). Both *35S:EGR1* and *35S:YFP-EGR1* as well as *35S:MASP1* and *35S:YFP-MASP1* lines were used. The lines expressing untagged or YFP-tagged proteins all had consistent result and the combined data are labeled as *35S:MASP1* and *35S:EGR1* for clarity of presentation. Data plotted separately for each transgenic line is shown in Supplemental Fig. S4.

than half; consistent with many previous reports that low ψ_w decreases cell division (Fig. 2A

and B; data from individual transgenic lines is shown in Supplemental Fig. S4A). In the low

ψ_w treatment, *egr1-1egr2-1* had increased number of GUS foci compared to wild type.

35S:MASP1 had increased numbers of CYCB1::GUS foci in both the low ψ_w and unstressed

control treatments; however, the effect was proportionally much larger in the low ψ_w

treatment. Interestingly, the *egr1-1egr2-1masp1-1* triple mutant had the same low level

CYCB1::GUS as *masp1-1* (Fig. 2A and B). Thus, the increased cell division of *egr1-1egr2-1*

at low ψ_w was dependent upon MASP1. Note that the CYCB1::GUS reporter does not allow

one to determine which cell layer the GUS foci are located in. This does not affect

interpretation of our results as none of our mutants or transgenic lines disrupted the

arrangement of different cell types within the root tip (see below).

The CYCB1::GUS reporter can also be induced by DNA damage (Schnittger and De

Veylder, 2018). This is unlikely to have affected our results as the moderate severity stress

used here is not expected to cause DNA damage and we observed reduced CYCB1::GUS

under low ψ_w rather than increased activity that would be expected if substantial DNA

damage occurred. Nonetheless, we also assessed cell division activity using 5-ethynyl-2′-

deoxyuridine (EdU) staining to label newly synthesized DNA. Compared to the

CYCB1::GUS results, higher numbers of EdU foci were observed in the meristem region

(apical 300 µm of the root) for both control and stress treatments (Supplemental Fig. S5).

This is likely because of the amount of time needed to effectively stain the roots and because

the EdU stain is not actively removed after DNA replication is complete. Despite this

difference, it was clear that the number of EdU foci was reduced by approximately half in

low ψ_w-treated wild type compared to the unstressed control. Also consistent with the

CYCB1::GUS results, *egr1-1egr2-1* and *35S:MASP1* had significantly increased EdU

staining compared to wild type at low ψ_w while *masp1-1, egr1-1egr2-1masp1-1* and

35S:EGR1 were low but not significantly different from wild type (Supplemental Fig. S5).

This further indicated that EGRs suppressed cell division at low ψ_w while MASP1 promoted

it. In EdU staining of the unstressed control, none of the mutant or ectopic expression lines

differed from wild type (Supplemental Fig. S5).

EGRs affect cell size independently of MASP1

In the Arabidopsis primary root, cells are fully expanded once they reach 1.5 to 2 mm

from the root apex (van der Weele et al., 2000; West et al., 2004; Yang et al., 2017).

Therefore, we measured epidermal cell length at 2 mm from the root apex (Fig. 2C). Cell

length of unstressed wild type was similar to previous reports (West et al., 2004; Yang et al.,

2017). Also consistent with previous reports, we found that root cell expansion was less

affected by low ψ_w than cell division. In wild type, low ψ_w reduced cell length by 20 percent

compared to the unstressed control. Interestingly, *egr1-1egr2-1* and *egr1-1egr2-1masp1-1*

had increased cell size in both the stress and control treatments while *masp1-1* did not. Thus,

loss of EGR activity led to enhanced cell expansion and, in contrast to cell division, the effect

of *egr1-1egr2-1* on cell expansion did not depend on MASP1. These data explain the

intermediate growth phenotype of *egr1-1egr2-1masp1-1* at low ψ_w as the triple mutant had

the increased cell expansion of *egr1-1egr2-1* but not the increased cell division. We also

found that the *egr* single mutants tended to have slightly larger cells than wild type under low

ψ_w; however, there was not a significant difference for any of the individual mutants

(Supplemental Fig. S1B). This indicated that *EGR1* and *EGR2* act redundantly to restrict cell

expansion during low ψ_w stress.

Conversely, *35S:EGR1* further decreased cell size by a small amount under low ψ_w

but had no effect in the unstressed control, consistent with the overall strong growth

restriction of *35S:EGR1* compared to wild type at low ψ_w and lack of *35S:EGR1* effect on

unstressed plants (Fig. 1A; data for individual transgenic lines is shown in Supplemental Fig.

S4B). Despite its decreased growth at low ψ_w, *masp1-1* did not have significantly smaller

cells; although, this was due in part to variability of cell size in *masp1-1*, as we also noted in

previous examination of hypocotyl cells (Bhaskara et al., 2017). *35S:MASP1* had slightly

bigger cells in both control and stress treatments, similar to *egr1-1egr2-1*. As the regulation

of cell size and cell division involves both coordination and compensatory mechanisms

(Gonzalez et al., 2012; Clauw et al., 2016), some of the differences in cell size observed may

be indirect effects of changes in cell division or meristem size.

***egr1-1egr2-1* and *35S:MASP1* maintain a larger root meristem at low ψ_w.**

To determine whether the effects of EGRs and MASP1 on cell division may alter

cellular organization within the root meristem, we examined the proximal root meristem

region just behind the QC for all genotypes in control and low ψ_w stress treatments (Fig. 3A).

The QC region as well as epidermal, cortex and endodermal cell layers were intact in all

genotypes and there was no evidence of improperly placed cell divisions. In the stress

treatment, epidermal, cortex and endodermal cells in the proximal meristem region were

larger (longer) than they were in unstressed roots (Fig. 3A). This was consistent with data in

Figure 3: Ectopic EGR or MASP1 expression alters root meristem size during low ψw stress.
   A. Representative images of PI stained root tips to show the cell size and organization of cell
   files in the proximal meristem region. Scale bars indicate 20 μm. The epidermal (Ep.),
   cortical (Cor.) and endodermal (En.) cell layers are labeled on the wild type control image.

B. Meristem cell number and meristem length five days after transfer of five-day-old mutant
   and ectopic expression seedlings to unstressed control or to low ψw (-0.7 MPa). Data are
   combined from 2-3 independent experiments for each genotype (n = 19-31 in the control, n =
   33-65 in the stress treatment). Symbols indicate data from individual roots while black bars
   and error bars indicate the mean and standard deviation for each genotype. Within the stress
   treatments, genotypes sharing the same letter do not significantly differ from one another
   (ANOVA with Tukey's post hoc test, corrected P ≤ 0.05). There were no significant
   differences among genotypes for meristem cell number in the unstressed control. In wild
   type, meristem size was significantly different between control and stress treatment (T-test, P
   ≤ 0.001). In the stress treatment, the increase of meristem length in *35S:MASP1* was
   marginally non-significant compared to wild type (adjusted P = 0.08) and *egr1-1egr2-1* may
   have a slightly longer meristem than *egr1-1egr2-1masp1-1* (adjusted P = 0.06).
   Representative images showing differences in root meristem length are shown in
   Supplemental Fig. S6). For *35S:EGR1* and *35S:MASP1*, three independent transgenic lines
   were used for each gene. Data for each individual transgenic line plotted separately is shown
   in Supplemental Fig. S7.

Figure 2 showing that low ψw inhibited cell division to greater extent than cell expansion.

Cell shape also became more irregular at low ψ_w, perhaps associated with the altered

coordination between cell division and cell expansion. The larger meristem cell size was

particularly apparent in *egr1-1egr2-1masp1-1* but was less pronounced in *egr1-1egr2-1* and

35S:MASP1, consistent with the different effects of these genotypes on cell division in the

CYCB1::GUS assays.

To quantify EGR and MASP1 effects on root meristem size, we counted the number

of cells in a file of epidermal cells from the QC to the first elongated cell of the transition

zone and also measured the distance from the QC to the first elongated cell (Fig. 3B,

representative images to illustrate differences in meristem length are shown in Supplemental

Fig. S6 and data for individual transgenic lines is shown in Supplemental Fig. S7). Meristem

cell number and size was significantly reduced by low ψ_w in wild type, similar to previous

observations in seedlings exposed to moderate severity salt stress (West et al., 2004).

Compared to wild type, the number of cells in the root meristem was increased in

35S:MASP1 but decreased in *35S:EGR1* at low ψ_w (Fig. 3A; Supplemental Fig. S7),

consistent with their effects on growth and cell division. Meristem length was also strongly

decreased in *35S:EGR1* at low ψ_w. Conversely, meristem length of *35S:MASP1* at low ψ_w

was significantly larger than that of *masp1-1* or *egr1-1egr2-1masp1-1*. The *egr* and *masp1*

mutants did not significantly differ from each other or from wild type in meristem cell

number or meristem length at low ψ_w (although *egr1-1egr2-1* may have a slightly longer

meristem than *egr1-1egr2-1masp1-1*, adjusted P = 0.06). Also, there were no significant

differences in meristem size among genotypes in the unstressed control. Similar effects on

root meristem size were seen at four days after transfer to low ψ_w (Supplemental Fig. S8). In

seedlings transferred to 75 mM NaCl, *egr1-1egr2-1* and *35S:MASP1* had higher meristem

cell number and larger meristem size than *masp1-1* or *35S:EGR1* (Supplemental Fig. S3B).

Roots of all genotypes were thinner after 5 days at low ψ_w compared to unstressed

roots (Supplemental Fig S6). This was consistent with previous observations that low ψ_w

prevented the increase in root diameter that occurs over time in unstressed Arabidopsis roots

(van der Weele et al., 2000). *35S:MASP1*, and perhaps *egr1-1egr2-1,* maintained a higher

root diameter, while *masp1-1* was thinner. This may be consistent with the interpretation of

van der Weele et al. (2000) that Arabidopsis root diameter was positively correlated with root

elongation rate. Root thinning in response to low ψ_w has also been observed in a number of

other plant species, such as maize (Sharp et al., 1988).

As the root meristem is small in newly germinated seeds and later increases in size as root growth accelerates (Beemster and Baskin, 1998; Biancucci et al., 2015) we also examined root meristem size in younger seedlings. In 2- and 4-day-old unstressed seedlings, *egr1-1egr2-1masp1-1* had increased meristem size compared to other genotypes (Supplemental Fig. S9). This indicated that EGRs and MASP1 may have complementary roles in controlling root meristem size during early seedling development in addition to their roles in regulating meristem size under low ψ_w stress.

Spatial differences in EGR2 versus MASP1 stoichiometry influence meristem size during low ψ_w stress.

Examination of EGR2 and MASP1 root tip expression pattern illustrated how they can interact to regulate meristem size and also indicated why ectopic EGR or MASP1 expression had such dramatic effects on root meristem cell number and meristem size. Among the three EGR PP2Cs, *EGR2* was selected for *promoter:GUS* analysis because EGR2 has relatively high expression and because *egr2* single mutants had slightly larger effect on growth compared to other *egr* single mutants (Supplemental Fig. S1; Bhaskara et al., 2017; Wendrich et al., 2017). In unstressed plants, *EGR2$_{promoter}$:GUS* expression was high in the mature region but undetectable in the apical 1 mm of the root tip which included the meristem and cell expansion regions (Fig. 4A). Low ψ_w stress led to an increase in *EGR2$_{promoter}$:GUS* expression including detectable expression closer to the root tip. This may be related to a shifting of the cell expansion and cell differentiation zones closer to the root tip, as has been observed for high temperature and other treatments that inhibit root elongation (Yang et al., 2017). However, *EGR2$_{promoter}$:GUS* activity was still not detected in the apical 200-300 µm which contains the root meristem. *MASP1$_{promoter}$:GUS* expression was essentially the converse of *EGR2* with high expression in the proximal meristem region just behind the QC in both control and stress treatments (Fig. 4A). Low ψ_w induced a higher level of *MASP1$_{promoter}$:GUS* staining across the root meristem as well as an enhanced, but still relatively low, level of expression in mature root tissue. This was consistent with our previous observation that *MASP1* gene expression and protein level of whole seedlings increased in plants exposed to a more severe -1.2 MPa low ψ_w treatment (Bhaskara et al., 2017).

To examine patterns of EGR2 and MASP1 protein accumulation, we generated plants transformed with a genomic fragment containing the *EGR2* or *MASP1* promoter and gene

Figure 4: EGR and MASP1 have differing spatial patterns of expression in the root tip.

A. Representative images of *EGR2~promoter~:GUS* and *MASP1~promoter~:GUS* staining in primary root tips four days after transfer to control or low ψ~w~ (-0.7 MPa). Red scale bars indicate 100 μm. Orange bars indicate 1 mm. Images are representative of the expression patterns seen in three independent transgenic lines for each construct.

B. Quantitation of *EGR2~promoter~:EGR2-YFP* and *MASP1~promoter~:MASP1-YFP* fluorescence intensity along the root tip. Roots were imaged with a fluorescence imager and Image J was used to perform line scans of root tips for 6-10 roots (from two independent transgenic lines for each construct assayed in two independent experiments) at four days after transfer of seedlings to control or low ψ~w~ (-0.7 MPa) treatments. Representative root tip images used for the scans are shown in Supplemental Fig. S12C. Dashed line boxes indicate the end of the root meristem for control and stress treatments (based on data in Fig. 3B). Gray error bars (gray shading) indicate the standard error. Note the log scale of the y-axis.

C. Root tip expression pattern of *EGR2~promoter~:EGR2-YFP* and *MASP1~promoter~:MASP1-YFP* in unstressed plants (Control) and low ψ~w~-treated plants (-0.7 MPa, 4 days). The gray dashed lines in the whole root tip images indicate the positions of the radial scans (50, 150, 350 and 450 μm from the quiescent center). The whole root tip images were made by merging individual images using Leica LAS X software. Images are representative of the expression pattern seen in three independent transgenic lines for each construct. In the whole root tip images, scale bars indicate 100 μm. In the radial scan images, scale bars indicate 20 μm.

body with C-terminal fusion to YFP. We demonstrated that both *MASP1~promoter~:MASP1-YFP*

and *EGR2_{promoter}:EGR2-YFP* could complement the growth phenotypes of *masp1-1* or *egr1-*

1egr2-1 at low ψ_w (Supplemental Fig. S10A and S11A). Note that *egr1-1egr2-1* transformed

with *EGR2_{promoter}:EGR2-YFP* still had a small increase in growth compared to wild type,

again consistent with redundant function of EGR1 and EGR2 in growth regulation at low ψ_w

(Supplemental Fig. S11A).

A broad scale analysis using line scans of fluorescence intensity along the root tip for

EGR2_{promoter}:EGR2-YFP and *MASP1_{promoter}:MASP1-YFP* constructs showed that EGR2 was

very low in the apical 50-100 μm of the root but increased nearly 10-fold in more distal parts

of the root tip (Fig. 4B). In contrast MASP1 protein level was highest in the apical 100 μm

but declined thereafter to a low basal level. MASP1 protein was detected at positions farther

from the root tip than *MASP1_{promoter}:GUS* expression, which was highly concentrated in the

proximal root meristem of unstressed plants. This was consistent with cells being displaced

away from the root tip as the MASP1 protein was being synthesized. In low ψ_w-treated roots

both the decrease in MASP1 and increase in EGR2 occurred closer to the root tip

(Supplemental Fig. S12A, Fig. 4B). The point at which MASP1 expression reached a

minimum value corresponded to the end of the root meristem under both control and low ψ_w

conditions (indicated by the dashed boxes in Fig. 4B). Interestingly, MASP1 protein level

did not substantially increase in the root meristem under low ψ_w stress despite the strong

increase in *MASP1_{promoter}:GUS* activity at low ψ_w (compare Fig. 4A and B). The higher

MASP1_{promoter}:GUS activity at low ψ_w, along with reduced root elongation and slower

displacement of cells away from the root tip would be expected to lead to higher MASP1

protein level. Since this was not observed, it is possible that MASP1 protein stability is

affected by low ψ_w to control its abundance and spatially restrict MASP1 protein

accumulation. Conversely EGR2 accumulated to relatively high level in the region 200-400

μm from the root tip despite minimal EGR2 promoter activity in this region (compare Fig.

4A and B), suggesting that EGR2 is a relatively stable (or efficiently translated) protein

compared to MASP1.

We also conducted a similar line scan analysis of root tips from plants expressing

35S:YFP-EGR1 and *35S:YFP-MASP1* (Supplemental Fig. S12B). The *35S* promoter led to a

relatively even protein level across the root tip for both EGR1 and MASP1. This meant that

35S-driven EGR1 expression was more than 10-fold higher than endogenous EGR2 in the

apical 50-80 μm of the root. Conversely, *35S*-driven MASP1 expression was higher than

endogenous MASP1 in the region 200-500 μm from the root tip. These data indicated that

35S:EGR1 and *35S:MASP1* strongly affect root meristem size and growth at low ψ_w because

35S-driven ectopic expression of either gene disrupts spatial patterns of EGR versus MASP1

stoichiometry in the root tip. This was supported by observation that, in contrast to

35S:EGR1, expression of *EGR2_{promoter}:EGR2-YFP* in the wild type background did not inhibit

growth at low ψ_w (Supplemental Fig. S11A) and observation that *MASP1_{promoter}:MASP1-YFP*

in the wild type background did not promote growth at low ψ_w (Supplemental Fig. S10A). It

was also interesting to note that root tip expression of *35S*-driven EGR1 was similar in both

control and stress conditions (Supplemental Fig. S12B) despite the fact that *35S:EGR1* plants

only had reduced root growth, cell division or cell size in the stress treatments (Fig, 1, 2, 3;

Supplemental Fig. S3). This raises the possibility that a stress-related factor is required to

potentiate the effect of ectopic EGR expression.

High resolution imaging of the root tip showed that EGR2 protein level was highest in

the cortical cell layer with much lower level of EGR2 expression in some epidermal cells

(Fig. 4C, Supplemental Fig. S11C). In unstressed roots, EGR2 was absent from the proximal

meristem region but there was a peak of EGR2 in the distal meristem region. Observations in

multiple transgenic lines found that this peak of EGR2 always preceded the end of the cell

division zone, even though its exact position could vary from root to root. A similar pattern

was observed in roots exposed to low ψ_w; however, we consistently observed that low ψ_w

caused EGR2 expression to encroach closer to the QC (Fig. 4C; Supplemental Fig. S11C).

Radial scans at in the proximal meristem region (50 and 150 µm from the QC) and distal

meristem to beginning of the cell expansion zone (350 and 450 µm from the QC) confirmed

the gradient in EGR2 protein level and also confirmed that EGR2 was mainly expressed in

cortical cells (Fig. 4C; positions of the radial scans are indicated by dashed lines on the root

tip images). Note however that EGR2 was sometimes difficult to fully visualize in radial

scans unless the scan was sufficiently close to a transverse side of a cortex cell where EGR2

signal was highest.

Conversely, MASP1 protein level of unstressed roots was high in the region just

behind the QC and was expressed across all cell types. In the distal meristem and transition

to cell elongation (approximately 250 to 450 µm behind the QC), MASP1 protein level

declined in the inner cell layers (including cortex) but could still be detected in epidermal

cells into the cell elongation zone (Fig. 4C; Supplemental Fig. S10B). At low ψ_w, we

consistently observed that MASP1 protein level remained high in the proximal meristem

region but declined substantially in the distal meristem and beginning of the cell expansion

region (Fig. 4C; Supplemental Fig. S10B). Radial scans confirmed this decline of MASP1

protein level and showed that it occurred across all cell types (Fig. 4C). These data

demonstrated that in unstressed plants EGR2 and MASP1 overlapped in the root cortex near

the distal end of the root meristem. Low ψ_w stress increased the EGR2-MASP1 overlapping

region and pushed it closer to the QC, consistent with shortening of the root meristem in

response to low ψ_w.

Phosphomimic MASP1 (S670D), but not wild type MASP1, can overcome *35S:EGR1*

suppression of root meristem size at low ψ_w.

Previously we demonstrated that EGRs (either directly or indirectly) attenuate

MASP1 phosphorylation and that phosphonull MASP1^{S670A} was inactive and unable to

complement the *masp1-1* mutant (Bhaskara et al., 2017). Consistent with these previously

reported growth phenotypes, we found that *35S:MASP1^{S670D}* increased meristem size while

35S:MASP1^{S670A} slightly decreased meristem size under low ψ_w (Fig. 5A and B). Combined

with the results presented above, this suggested that EGR control of MASP1 S670

phosphorylation in their overlapping zone of expression is a key factor controlling root

meristem size during low ψ_w. If this hypothesis is correct, we expect that *35S:EGR1* will be

able to suppress the increased meristem size of *35S:MASP1* plants but would be unable to

suppress meristem size in plants expressing *35S:MASP1^{S670D}*. To test this hypothesis, we

crossed both wild type and *35S:EGR1* to *35S:MASP1*, *35S:MASP1^{S670D}* and *35S:MASP1^{S670A}*

and then examined root meristem size of low ψ_w-treated F$_1$ seedlings. Consistent with our

hypothesis, F$_1$ seedlings of *35S:MASP1* x W.T. had significantly larger root meristem size

than W.T. and larger than F$_1$ seedlings of *35S:MASP1* x *35S:EGR1* (Fig. 5C). In contrast, F1

seedlings of *35S:MASP1^{S670D}* x *35S:EGR1* had the same increase in meristem size as

35S:MASP1^{S670D} x W.T., indicating that *35S:EGR1* was unable to counteract the effect of

phosphomimic MASP1. *35S:MASP1^{S670A}* was again ineffective and did not increase

meristem size in F$_1$ seedlings of the W.T. or *35S:EGR1* crosses (Fig. 5C). These data

supported our hypothesis that EGR control of MASP1 S670 phosphorylation is a key

mechanism to control root meristem size during low ψ_w stress.

EGR-MASP1 shoot expression patterns

EGRs and MASP1 also clearly affect shoot growth (Fig. 1A; Bhaskara et al., 2017)

and EGR2-MASP1 expression patterns suggested that a similar mechanism of opposing

Figure 5: EGR attenuation of MASP1 S670 phosphorylation in their zone of overlapping expression controls root meristem size at low ψ_w.

A. Root meristem cell number at low ψ_w for wild type and two independent transgenic lines expressing phosphonull MASP1 (*35S:MASP1^{S670A}*) or phosphomimic MASP1 (*35S:MASP1^{S670D}*). The MASP1 phosphonull and phosphomimic lines were previously described in Bhaskara et al. (2017). Data are combined from 2-3 independent experiments (n = 35 for wild type and 10-18 for each transgenic line). The transgenic lines used here are in the GFP-TUA6 background (Supplemental Table S1). It was previously shown that presence of GFP-TUA6 did not affect the growth phenotypes of any EGR-MASP1 mutant or transgenic lines (Bhaskara et al., 2017). Symbols indicate data from individual roots while black bars and error bars indicate the mean and standard deviation for each genotype. Different letters indicate significant differences between genotypes (ANOVA with Tukey's post hoc test, corrected P ≤ 0.05).

B. Representative root meristem images of each genotype in A. Scale bars indicate 100 μm.

C. Quantitation of meristem size in F$_1$ seedlings of *35S:EGR1* crossed to lines ectopically expressing non-mutated MASP1, phosphonull MASP1 (S670A) or phosphomimic MASP1 (S670D). All the MASP1 lines express untagged MASP1 in the GFP-TUA6 background. Root meristem size was measured five days after transfer to low ψ_w (-0.7 MPa). Bars indicate the mean, error bars indicate the standard deviation from 2 independent experiments for each genotype (n = 56 for wild type, n = 21-29 for F$_1$ genotypes). Asterisks directly above the bars indicate a significant difference compared to wild type, brackets show the results of comparison between the indicated genotypes (ANOVA

with Tukey's post hoc test, * indicates corrected P ≤ 0.05; ** P ≤ 0.005; **** P ≤ 0.0001). Note that the difference in meristem cell number between wild type (W.T.) and W.T. x *35S:EGR1* was marginally nonsignificant (adjusted P = 0.07).

D. Model of how EGR-MASP1 signaling affects meristem function and root growth at low ψ_w. MASP1 is highly expressed in all cell types of the proximal root meristem where it promotes cell division. MASP1 decreases in expression farther from the QC, especially in the inner cell layers and cortex. Conversely, EGR2 is low in the proximal meristem but high in the distal meristem and elongation zone where it influences cell expansion. EGR2 is highly expressed in cortical cells while MASP1 does not have cell type specific expression. During low ψ_w stress, EGR2 expression encroaches closer to the proximal meristem region. In their overlapping zone of expression, EGRs attenuate phosphorylation of MASP1, particularly in cortex cells, to suppress MASP1 activity and restrict meristem size. The high expression of EGR2 in cortex is consistent with recent data indicating that this cell layer is important for growth responses to low ψ_w. Low ψ_w activates (or de-represses) EGRs such that EGR-MASP1 signaling is a dominating factor controlling root elongation and meristem function during low ψ_w stress but has less or no effect on unstressed plants.

EGR-MASP1 signaling may also operate in shoot and leaf tissues. *EGR2$_{promoter}$:GUS*

expression was observed in young expanding leaves as well as in recently emerged leaf

primordia and was induced by low ψ_w in young leaves (Supplemental Fig. S13A and B).

However, *EGR2$_{promoter}$:GUS* expression was not detected in the shoot meristem under either

condition. Similarly, in *EGR2$_{promoter}$:EGR2-YFP* lines the EGR-YFP fusion protein could not

be detected in the shoot meristem (Supplemental Fig. S14). Conversely, *MASP1$_{promoter}$:GUS*

expression and native promoter driven MASP1-YFP expression (*MASP1$_{promoter}$:MASP1-YFP*)

were detected in the shoot meristem and surrounding leaf primordia (Supplemental Fig. S13

and S14). Low ψ_w reduced the level of MASP1 protein in the shoot meristem despite strong

MASP1 promoter activity in the shoot meristem at low ψ_w (Supplemental Fig. S13 and S14).

We could also clearly see *MASP1$_{promoter}$:GUS* expression in the base (close to the petiole) of

young leaves where cell division to drive leave blade expansion occurs and could see that in

this tissue it overlapped with *EGR2$_{promoter}$:GUS* expression (Supplemental Fig. S13). Thus,

separate and overlapping domains of MASP1 and EGR expression may be involved in

controlling shoot and leaf growth during low ψ_w in a manner similar to their role in

controlling root growth.

Discussion

The rate and duration of cell division and extent of cell expansion are all potential points of control to alter plant growth in response to environmental signals. Which of these control points are most important for modulating growth during abiotic stress and the signaling proteins involved are important questions for plant stress research. Developmental studies have advanced the idea that opposing gradients of regulatory gene expression across the meristem control the transition from cell division to cell expansion (Wendrich et al., 2017; Salvi et al., 2020). Although many such gene expression gradients have been observed, there is scant data to show whether this leads to differences in the stoichiometric ratios of protein abundance at different positions along the root meristem and little data to show how proteins with spatial differences in stoichiometry can influence each other's activity in a manner consequential for growth regulation. Our results bring these two lines of inquiry together by demonstrating that differing spatial patterns of EGR-MASP1 protein abundance are a key factor regulating root meristem function during low ψ_w stress. Disrupting the spatial pattern of EGR-MASP1 protein stoichiometry, either by ectopic expression or by loss of function mutations, changes how root growth responds to low ψ_w. Under low ψ_w, phosphorylated MASP1 promotes continued cell division in the proximal root meristem. However, in more distal locations MASP1 protein level decreases and EGRs inactivate remaining MASP1, particularly in the cortex, a cell type important for controlling growth responses to low ψ_w (Dietrich et al., 2017). This hastens the transition from cell division to cell expansion (Fig. 5D). Ectopic expression of either EGR2 or MASP1 upsets this balance allowing a longer duration of cell division with higher rates of cell division in the distal meristem region (in the case of *35S:MASP1*) or an earlier exit from cell division and smaller meristem (in the case of *35S:EGR1*). Ectopic expression of MASP1 counteracts the effect of low ψ_w to restrict MASP1 expression to a smaller region just behind the QC. Conversely, ectopic expression of EGR1 amplifies the effect of low ψ_w to push EGR expression closer to the QC. Thus, the opposing action of MASP1 and EGRs, and differences in the stoichiometric ratio of EGR-MASP1 protein abundance at different positions along the root meristem, allow cell division in the proximal meristem to be protected while also allowing the plant to down-regulate meristem activity and decrease growth during moderate severity low ψ_w stress (Fig. 5D). This model of how EGR-MASP1 signaling controls root growth is also supported by previous data demonstrating that the EGR phosphatases interact with MASP1 and that changing the EGR expression level changes the

level of MASP1 phosphorylation (Bhaskara et al., 2017). Note that we do not exclude the

possibility that EGRs and MASP1 also affect the rate of cell division (cell cycle time) within

the meristem since *egr1-1egr2-1* had higher cell division activity at low ψ_w while having

similar meristem size as wild type and *35S:MASP1* had a larger effect on meristem cell

number than on meristem length.

35S:EGR1 did not decrease growth, cell division, cell size or meristem size of

unstressed plants despite having strong effect on plants exposed to low ψ_w and despite similar

levels of root tip *35S*-driven EGR1 protein expression in both control and stress treatments.

One possible explanation is that EGR phosphatase activity is inhibited (or not activated) in

the root meristem of unstressed plants. Such post-translational regulation of EGR activity

could occur via interaction with regulatory proteins in a manner similar to inhibition of Clade

D PP2Cs by SAUR proteins (Spartz et al., 2014), which is also important for growth

regulation, or similar to PYL/RCAR inhibition of Clade A PP2C activity which is of central

importance to ABA signaling (Cutler et al., 2010; Raghavendra et al., 2010). As we have

previously discussed, it is possible that additional PP2Cs, including Clade E PP2Cs, are

paired with regulatory proteins that control their activity (Bhaskara et al., 2019). However,

we cannot rule out other explanations such as the possibility that EGRs are active in

unstressed plants but their effect masked by other mechanisms that control meristem activity,

the phosphatase is active but access to its substrate proteins (including MASP1) is restricted

in unstressed plants or, EGR substrate proteins only become phosphorylated in stress-treated

plants. We also note that high expression of EGR1 in cell types other than the cortex could

be a factor in the dramatic effect of *35S:EGR1* on root meristem size during low ψ_w stress.

However, the observation that *egr1-1egr2-1* had increased cell division at low ψ_w, while

egr1-1egr2-1masp1-1 did not, indicates that EGR regulation of MASP1 phosphorylation

status in cortical cells, where EGR2 is most highly expressed, can be sufficient to affect cell

division activity of the meristem as a whole. None of our mutant or ectopic expression lines,

had altered root meristem morphology in terms of organization of cell files and placement of

cell division planes. How expression of EGR2 predominantly in the cortical cell layer affects

root cell division and cell expansion during low ψ_w stress in way that preserves the

coordination between different cell layers is an interesting question for future research.

EGR2 protein level was dramatically higher in cortex compared to other root cell

types. This is consistent with recent root tip single-cell RNA sequencing which found EGR

expression in most cells that clustered with the cortex and some cells in the epidermal cluster

(Wendrich et al., 2020). The same data set showed MASP1 expression across all root cell

types. The cortex-enriched expression of EGR2 is particularly interesting in light of recent

observations that root cortex is the key cell layer for response to a ψ_w gradient during

hydrotropic bending (Dietrich et al., 2017). Dietrich et al. (2017) demonstrated that

expression of SnRK2.2/SnRK2.3 in cortical cells was critical for differential cell elongation

during hydrotropic root bending. Interestingly, EGR2 has been shown to inhibit the kinase

activity of SnRK2.6 (also known as Open Stomata 1, OST1; Ding et al., 2019) which is

closely related to SnRK2.2 and SnRK2.3. Thus, it will be of interest to investigate whether

EGR regulation of SnRK2 activity in the root cortex is one factor that limits cell elongation

during low ψ_w or is involved hydrotropic root bending. Another recent study has also

proposed that inter-cellular signaling originating from the cortex is important for controlling

root growth in response to environmental signals (Mielke et al., 2021). It is similarly

interesting to note that altered hydrotropic response 1 (*ahr1*), which is putatively involved in

hydrotropic root bending, also has increased root elongation and maintains larger root

meristem size than wild type during low ψ_w stress (Salazar-Blas et al., 2017). Unfortunately,

the mutated gene responsible for the *ahr1* phenotype has not yet been identified.

It is also worth noting that *egr1-1egr2-1* did not have detectable effect on root cell

division of unstressed plants but did affect cell size in both control and stress treatments.

This again is consistent with the idea that EGRs affects cell size via a different, MASP1-

independent, mechanism and suggests that EGR activity may also be regulated differently

during cell expansion and maturation than in the meristem. We also observed that,

35S:MASP1 had a similar increased cell size as *egr1-1egr2-1* and *egr1-1egr2-1masp1-1*.

While it is tempting to speculate that this is related to MASP1 effects on microtubule stability

(Bhaskara et al., 2017), a perhaps simpler explanation is that ectopic MASP1 expression in

the cell expansion zone, where endogenous MASP1 is very low, titrates away EGR

phosphatase activity and thus allows greater cell expansion. Our data confirmed that MASP1

S670 phosphorylation is critical for its function. Thus, kinases that phosphorylate MASP1

S670 are also potential regulators of meristem size and function.

Our observations that low ψ_w decreases root cell division; meristem size and cell

expansion are similar to observations of West et al. (2004) who studied the basis for

decreased root elongation in wild type Arabidopsis exposed to mild salt stress. In their case,

salt stress reduced meristem size and cell production (measured both kinematically and using

the CYCB1::GUS reporter) and also reduced final cell length. The increased growth and

increased root meristem size of *egr1-1egr2-1* and *35S:MASP1* were also similar in some

ways to the recently reported effects of β-cyclocitral which increased root meristem size (but

did not affect cell length) when applied to unstressed plants (Dickinson et al., 2019). β-

cyclocitral also stimulated growth during salt stress; although, whether this was due to

increased meristem size and whether β-cyclocitral affects growth during drought stress are

yet to be reported. Nonetheless, a common theme that emerges from these results is that,

whether by genetic or chemical means, inducing plants to maintain a larger population of

dividing cells (larger meristem) can allow enhanced growth maintenance after exposure to

abiotic stress.

Materials and Methods

Plant materials and stress treatment

The *egr1-1egr2-1*, *masp1-1* and *egr1-1egr2-1masp1-1* mutants have been previously

described (Bhaskara et al., 2017). The meristem analysis used previously described *35S:YFP-*

MASP1 line as this construct could complement the *masp1-1* mutant and was functionally

equivalent to *35S:MASP1* (untagged MASP1) in our previous analysis (Bhaskara et al.,

2017). Data presented in Fig. 1-3 use both *35S:MASP1* and *35S:YFP-MASP1* lines as well as

35S:EGR1 and *35S:YFP-EGR1* to ensure that multiple independent transgenic lines were

used for each construct and to ensure that lines expressing the YFP-tagged proteins had the

same phenotypic effects as lines expressing untagged protein. Details of the construction of

these lines and vectors used are given in Bhaskara et al. (2017). To analyze the effects of

MASP1 phosphorylation, the *35S:MASP1^{S670A}/GFP-TUA6* and *35S:MASP1^{S670D}/GFP-TUA6*

lines described by Bhaskara et al. (2017) were used. It had been previously demonstrated

that that the presence of the GFP-TUA6 marker did not affect root elongation under either

control or stress treatments (Bhaskara et al. 2017). For clarity of presentation, these lines are

referred to simply as *35S:MASP1* and *35S:EGR1* in figures and text. The CYCB1::GUS

reporter lines were generated by crossing the mutants, *35S:YFP-MASP1*, *35S:YFP-EGR1* and

35S:EGR1 lines to the CYCB1:GUS reporter line and F$_3$ homozygous lines were used for cell

division assays. Additionally, the CYCB1:GUS reporter line was transformed with untagged

MASP1 (in pEG100 vector) and both lines used for the root elongation and cell division

assays in Fig. 1 and 2. In all cases, lines with a single locus insertion of the transgene (based

on segregation ratio of the selectable marker in the T$_2$ generation) were isolated and T$_3$

homozygous seed stocks used for experiments of for making crosses. Two to four

independent transgenic lines were used for all experiments. As the data were

indistinguishable between transgenic lines, combined data are shown in the main text figures

(Fig. 1, 2, 3) while data for individual transgenic lines is shown in Supplemental Figures S2,

S4, and S7). A list of all transgenic lines used in this study, and the experiments they were

used for, is shown in Supplemental Table S1.

For analysis of native promoter driven MASP1 and EGR2 protein levels and

expression pattern, C-terminal YFP constructs (*MASP1$_{promoter}$:MASP1-YFP* and

EGR2$_{promoter}$:EGR2-YFP) were generated by amplifying the promoter and coding region of

each gene (3663 bp and 3671 bp for *MASP1* and *EGR2,* respectively) from genomic DNA of

Col-0. The PCR products were first cloned in entry vector (pDNOR221) and were then

transferred to expression vector (pGWB540) via Gateway recombination (primer sequences

are shown in Supplemental Table 1). The constructs were subsequently transformed into

masp1-1 and *egr1-1,2-1* mutants (and Col-0 wild type in the case of *EGR2$_{promoter}$:EGR2-YFP*)

through *Agrobacterium*-mediated floral dip method. *Promoter:GUS* constructs were

generated by cloning the promoter region (1347 bp and 1653 bp upstream of start codon for

MASP1 and *EGR2,* respectively, primer sequences used for cloning are shown in

Supplemental Table S2) into pGWB443 by Gateway cloning and transformation into Col-0

wild-type. Homozygous T$_3$ seed stocks of lines having a single locus transgene insertion were

isolated as described above. Two or three independent transgenic lines were used for

subsequent experiments. Lines with *MASP1$_{promoter}$:MASP1-YFP* in the wild type background

were generated by crossing *MASP1$_{promoter}$:MASP1-YFP/masp1-1* to wild type to segregate the

masp1-1 T-DNA insertion away from the *MASP1$_{promoter}$:MASP1-YFP* transgene.

Seedling growth and low ψ_w treatment on PEG-infused agar plates was performed as

described previously (Verslues et al., 2006; Bhaskara et al., 2017). For seedling growth

assays, seeds were plated on unstressed control media, stratified for three days and

transferred to a growth chamber (22 °C, continuous light intensity of 100-120 μmol m^{-2} s^{-1}).

Five-day-old seedlings were transferred to fresh half-strength MS plates (control, -0.25 MPa)

or plates infused with PEG-8000 (-0.7 MPa). Plates were always prepared and infused with

PEG 15 h before use to avoid drying of the media and generate plates of consistent ψ_w. All

media was prepared without addition of sugar. Primary root elongation rates were measured

by marking the position of the root apex on the back of the plate at 48 h intervals for six days

after transfer. Root elongation rates were combined from two or three independent

experiments.

Other experiments quantifying total root elongation and seedling fresh or dry weight

(presented in Supplemental Fig. S10 and S11) were conducted as described in Bhaskara et al.

(2017). Five-day-old seedlings were transferred to fresh control plates or -0.7 MPa PEG-

infused plates. In the control treatment, root elongation was measured over the subsequent

five days (when the seedlings nearly reached the maximum size accommodated by the agar

plates) and seedling fresh and dry weights measured at the end of the experiment. The same

procedure was followed for the stress treatment except that the seedlings were allowed to

grow for ten days after transfer to reach a developmental stage close to that obtained by

unstressed seedlings at five days after transfer. For these experiments, growth of the

transgenic plants was normalized versus wild type seedlings growing in the same plate. Three

or four independent experiments were performed with each experiment containing two plates

for each genotype analyzed. The experiment averages were tested for significant differences

from wild type using one sample T-tests.

GUS, EdU and PI staining

β-Glucuronidase (GUS) activity staining for *promoter:GUS* and CYCB1::GUS lines

were performed using standard procedures. Seedlings were immersed in ice cold 90 %

acetone for 10 min, washed once with water and then put into staining solution (50 mM

phosphate buffer pH 7.2, 100 µM $K_3Fe(CN)_6$, 100 µM $K_4Fe(CN)_6$, 1 mM X-GLUC in

dimethyl formamide) and vacuum infiltrated for 20 min followed by incubation at 37 °C for 2

to 5 h. The samples were then transferred to 80% ethanol and keep at 4 °C for overnight.

Next day, the samples were cleansed with a solution of methanol and acetic acid (3:1) for 5-

10 min and then transferred to 80% ethanol before taking images (Zeis Imager Z1). Whole

rosette images (Supplemental Fig. S13) were obtained using the tiles function of the Zeiss

Zen software package.

For 5-ethynyl-2′-deoxyuridine (EdU) staining, seedlings were grown on 1/2 MS agar

plates for 5 days and transferred to control or stress plates (-0.7 MPa) for four days. The

seedlings were labelled by pouring EdU (Thermo Fisher Scientific) solution (10 µM EdU in

½ MS or ½ MS + -0.7 MPa PEG) into the plates and incubating for 15-30 min in a growth

chamber. Roots were transferred to a 2 mL tube and fixed for 20 min in 4% (w/v)

paraformaldehyde in PBS (pH 7.4), washed twice with PBS (2x10 min) and placed in 0.5%

(v/v) TritonX-100 in PBS. After 20 min, samples were washed with PBS twice and incubated

in 500 µL of Click-iT reaction cocktail (1× PBS, 4 mM CuSO4, 40 mM sodium ascorbate

and 5 µM AF594 azide) for 30 min in the dark. The Click-iT reaction cocktail was removed

and samples were washed with PBS once before observing under confocal microscope.

Fluorescence spots within the apical 300 μm of the root tip were counted.

For Propidium Iodide (PI) staining, seedlings were immersed in PI solution (0.01

mg/mL of PI in water or 0.01 mg/mL of PI in 285 mM mannitol (to match the water potential

of the -0.7 MPa stress and prevent cell swelling) for 60 sec (control) or 90 sec (stress) and

rinsed with water/mannitol solution before mounting on glass slide for confocal microscopy.

For meristem size analysis of seedlings 4 days after transfer to stress or control plates

(Supplemental Fig. S4) or younger 2 or 4 day-old seedlings (Supplemental Fig. S5), roots

were cleared with a 8:3:1 mixture of chloral hydrate:water:glycerol, mounted on a glass slide

and meristem cells counted as previously described (Biancucci et al., 2015)

Confocal microscopy and image analysis

Confocal laser-scanning microscopy was done using a Zeiss LSM510 for

quantification of meristem cell numbers and meristem size as well as EdU staining. For other

high resolution imaging a Zeiss LSM880 or Leica Stellaris 8 (Fig. 4C images) was used.

Low ψ_w-treated seedlings were mounted in solution of the same ψ_w (-0.7 MPa) as the stress

treatment to prevent swelling during imaging. EdU or PI stained samples were visualized

using excitation at 543 nm and emission at 588nm while YFP imaging used excitation at 514

nm and emission at 542 nm. To quantify the protein level of YFP-tagged EGR or MASP1

across the root tip (Fig. 4B, Supplemental Fig. S12), roots were imaged with a fluorescence

imager (Zeiss Axio Imager Z1) and 2 or 3 images covering the apical 1 cm of the root merged

using the tiles function of Zeiss Zen software and analyzed using the line scan function of

Image J. For each root tip, scans of three different lines starting the root tip and extending for

approximately 800 μm of the root were averaged. Scans followed the area between the stele

and epidermis of the root where fluorescence intensity was highest. Line scans of the area

alongside the root were used for background subtraction. The data presented in Fig. 4B, and

Supplemental Fig. S12A and B, are means of 5-10 roots per genotype and treatment

combined from two independent transgenic lines. For higher resolution visualization of the

whole root tip, merged images of the root tip were obtained by using the tiles function of

either Zeiss ZEN software (Supplemental Fig. S10 and S11) or Leica LAS X software (Fig.

4C). Radial images of root were obtained by collecting a Z-stack of root tip images and

processing using the Fiji orthogonal viewer. To image the shoot meristem, seedlings were

first dissected under a stereo microscope and the meristem region mounted and imaged with a

Zeiss LSM880 microscope.

Acknowledgements

We thank Arnould Savoure (UPMC, Sorbonne Universités) for providing seed of the

CYCB1::GUS reporter line, the staff of the imaging core facility of the Institute of Plant and

Microbial Biology (Ji-Ying Huang and Mei-Jane Fang) for microscopy assistance and Shih-

Shan Huang for laboratory assistance. This research was supported by an Academia Sinica

Investigator Award (AS-IA-108-L04) and Taiwan Ministry of Science and Technology grant

(MOST 107-2311-B-001 -037) to P.E.V.

Author contributions

P.E.V. conceived research, T.L., G.B.B and P.E.V. designed experiments, T.L., C.Y.C, M.B.

and G.B.B. performed experiments, T.L., C.-Y.C. and P.E.V. analyzed data, P.E.V. wrote the

manuscript with assistance from T.L. and G.B.B. All authors read and approved the

manuscript.

 Figure Legends

719 **Figure 1: EGRs and MASP1 affect growth during low ψ_w stress.**
720A. Representative seedlings of *EGR* and *MASP1* mutants and overexpression lines. Five-day-
721 old seedlings were transferred to either fresh control plates (-0.25 MPa) or PEG-infused agar
722 plates for moderate severity low ψ_w (-0.7 MPa) treatment. Plants were photographed at the
723 end of the experiment (six days after transfer). Scale bars indicate 1 cm.
724B. Primary root elongation rates over six days after transfer of seedlings to control or stress (-0.7
MPa) treatments. Blue or red symbols show elongation rates for individual seedlings. Black
bars and error bars indicate the mean and standard deviation for each genotype (n = 22 to 67).
Lower case letters in the stress panel indicate statistical differences (ANOVA with Tukey's
post hoc test, corrected P ≤ 0.05). Genotypes sharing the same letter did not significantly
differ from one another. Data are combined from three independent experiments and three
independent transgenic lines per construct. Both *35S:EGR1* and *35S:YFP-EGR1* as well as
35S:MASP1 and *35S:YFP-MASP1* lines were used. Data from additional experiments with
each individual transgenic line is shown in Supplemental Fig. S2. The lines expressing
untagged or YFP-tagged proteins had consistent result and the combined data are labeled as
35S:MASP1 and *35S:EGR1* for clarity of presentation.
735
Figure 2: EGRs and MASP1 affect root cell division and cell expansion during low ψ_w
stress.
A. Representative images of CYCB1::GUS staining in primary root tips of the indicated
mutant and ectopic expression genotypes 4 days after transfer of five-day-old
seedlings to either control or low ψ_w (-0.7 MPa). Scale bars indicate 100 µm.
B. Quantification of CYCB1::GUS foci in the primary root meristem. Symbols indicate
counts of GUS foci from individual roots while black bars and error bars indicate the
mean and standard deviation for each genotype. Within the control or stress
treatments, genotypes sharing the same letter do not significantly differ from one
another (ANOVA with Tukey's post hoc test, corrected P ≤ 0.05). In wild type, the
number of CYCB1::GUS foci was significantly different between control and stress
treatment (T-test, P ≤ 0.001). Data are combined from three or four independent
experiments (n = 21-66) for each genotype and at least two independent transgenic
lines for *35S:MASP1* and *35S:EGR1*. Data plotted separately for each transgenic line
is shown in Supplemental Fig. S4.
C. Cell length at 2 mm from the root tip measured four days after transfer of mutant or
ectopic expression seedlings to control or low ψ_w (-0.7 MPa). Format of data and
statistical analysis are the same as in B (n = 133-224). Both *35S:EGR1* and *35S:YFP-*
EGR1 as well as *35S:MASP1* and *35S:YFP-MASP1* lines were used. The lines
expressing untagged or YFP-tagged proteins all had consistent result and the
combined data are labeled as *35S:MASP1* and *35S:EGR1* for clarity of presentation.
Data plotted separately for each transgenic line is shown in Supplemental Fig. S4.
758
Figure 3: Ectopic EGR or MASP1 expression alters root meristem size during low ψ_w
stress.
A. Representative images of PI stained root tips to show the cell size and organization of
cell files in the proximal meristem region. Scale bars indicate 20 µm. The epidermal
(Ep.), cortical (Cor.) and endodermal (En.) cell layers are labeled on the wild type
control image.
B. Meristem cell number and meristem length five days after transfer of five-day-old
mutant and ectopic expression seedlings to unstressed control or to low ψ_w (-0.7

MPa). Data are combined from 2-3 independent experiments for each genotype (n =
19-31 in the control, n = 33-65 in the stress treatment). Symbols indicate data from
individual roots while black bars and error bars indicate the mean and standard
deviation for each genotype. Within the stress treatments, genotypes sharing the same
letter do not significantly differ from one another (ANOVA with Tukey's post hoc
test, corrected $P \leq 0.05$). There were no significant differences among genotypes for
meristem cell number in the unstressed control. In wild type, meristem size was
significantly different between control and stress treatment (T-test, $P \leq 0.001$). In the
stress treatment, the increase of meristem length in *35S:MASP1* was marginally non-
significant compared to wild type (adjusted $P = 0.08$) and *egr1-1egr2-1* may have a
slightly longer meristem than *egr1-1egr2-1masp1-1* (adjusted $P = 0.06$).
Representative images showing differences in root meristem length are shown in
Supplemental Fig. S6). For *35S:EGR1* and *35S:MASP1*, three independent transgenic
lines were used for each gene. Data for each individual transgenic line plotted
separately is shown in Supplemental Fig. S7.
782

Figure 4: EGR and MASP1 have differing spatial patterns of expression in the root tip.
A. Representative images of *EGR2$_{promoter}$:GUS* and *MASP1$_{promoter}$:GUS* staining in
primary root tips four days after transfer to control or low ψ_w (-0.7 MPa). Red scale
bars indicate 100 µm. Orange bars indicate 1 mm. Images are representative of the
expression patterns seen in three independent transgenic lines for each construct.
B. Quantitation of *EGR2$_{promoter}$:EGR2-YFP* and *MASP1$_{promoter}$:MASP1-YFP* fluorescence
intensity along the root tip. Roots were imaged with a fluorescence imager and Image
J was used to perform line scans of root tips for 6-10 roots (from two independent
transgenic lines for each construct assayed in two independent experiments) at four
days after transfer of seedlings to control or low ψ_w (-0.7 MPa) treatments.
Representative root tip images used for the scans are shown in Supplemental Fig.
S12C. Dashed line boxes indicate the end of the root meristem for control and stress
treatments (based on data in Fig. 3B). Gray error bars (gray shading) indicate the
standard error. Note the log scale of the y-axis.
C. Root tip expression pattern of *EGR2$_{promoter}$:EGR2-YFP* and *MASP1$_{promoter}$:MASP1-*
YFP in unstressed plants (Control) and low ψ_w-treated plants (-0.7 MPa, 4 days). The
gray dashed lines in the whole root tip images indicate the positions of the radial
scans (50, 150, 350 and 450 µm from the quiescent center). The whole root tip
images were made by merging individual images using Leica LAS X software.
Images are representative of the expression pattern seen in three independent
transgenic lines for each construct. In the whole root tip images, scale bars indicate
100 µm. In the radial scan images, scale bars indicate 20 µm.
805

Figure 5: EGR attenuation of MASP1 S670 phosphorylation in their zone of overlapping
expression controls root meristem size at low ψ_w.
A. Root meristem cell number at low ψ_w for wild type and two independent transgenic lines
expressing phosphonull MASP1 (*35S:MASP1^{S670A}*) or phosphomimic MASP1 (*35S:MASP1^{S670D}*).
The MASP1 phosphonull and phosphomimic lines were previously described in Bhaskara et al.
(2017). Data are combined from 2-3 independent experiments (n = 35 for wild type and 10-18 for
each transgenic line). The transgenic lines used here are in the GFP-TUA6 background
(Supplemental Table S1). It was previously shown that presence of GFP-TUA6 did not affect the
growth phenotypes of any EGR-MASP1 mutant or transgenic lines (Bhaskara et al., 2017). Symbols
indicate data from individual roots while black bars and error bars indicate the mean and standard

deviation for each genotype. Different letters indicate significant differences between genotypes

(ANOVA with Tukey's post hoc test, corrected $P \leq 0.05$).

B. Representative root meristem images of each genotype in A. Scale bars indicate 100 μm.

C. Quantitation of meristem size in F_1 seedlings of *35S:EGR1* crossed to lines ectopically expressing

non-mutated MASP1, phosphonull MASP1 (S670A) or phosphomimic MASP1 (S670D). All the

MASP1 lines express untagged MASP1 in the GFP-TUA6 background. Root meristem size was

measured five days after transfer to low ψ_w (-0.7 MPa). Bars indicate the mean, error bars indicate

the standard deviation from 2 independent experiments for each genotype (n = 56 for wild type, n =

21-29 for F_1 genotypes). Asterisks directly above the bars indicate a significant difference compared

to wild type, brackets show the results of comparison between the indicated genotypes (ANOVA

Transgenic line	Construct	Genetic Background	Experiments used for	Data shown in Figure:	Data Shown in Sup Figure:
35S:MASP1 T3-1/CycB1::GUS	untagged	CycB1::GUS	CycB1::GUS assay	2	S3, S4
35S:MASP1 T3-3/CycB1::GUS	untagged	CycB1::GUS	CycB1::GUS assay, Edu staining, root elongation, meristem size, cell length	1, 2, 3	S2, S3, S4, S5, S7
35S:YFP MASP1XCycB1::GUS F3	N-terminal YFP	CycB1::GUS	CycB1::GUS assay, Edu staining, root elongation, cell length	1, 2	S2, S4, S5, S8, S9
35S:YFP-MASP1 T3-2	N-terminal YFP	Col-0	YFP line scan, meristem size	3	S7, S12
35S:YFP-MASP1 T3-3	N-terminal YFP	Col-0	YFP line scan		S12
35S:YFP-MASP1 T4-2	N-terminal YFP	Col-0	YFP line scan, Edu staining, root elongation, meristem size, cell length	1, 3	S2, S4, S7, S12
35S:YFP-MASP1 T4-3	N-terminal YFP	Col-0	YFP line scan		S12
35S:YFP-EGR1 T4-7	N-terminal YFP	Col-0	YFP line scan, root elongation, Edu staining, meristem size, cell length	1, 3, 5	S2, S3, S4, S5, S7, S8, S9, S12
35S:EGR1 T2-2XCycB1::GUS F3	untagged	CycB1::GUS	CycB1::GUS assay, Edu staining, root elongation, meristem size, cell length	1, 2, 3	S2, S3, S4, S5, S7
35S:YFP-EGR1 T2-12XCycB1::GUS F3	N-terminal YFP	CycB1::GUS	YFP line scan, CycB1::GUS assay, Edu staining, root elongation, meristem size, cell length	1, 2, 3	S2, S4, S5, S7, S12
35S:MASP1 T4-1	untagged	TUA6-GFP	meristem size	5	
35S:MASP1 T4-2	untagged	TUA6-GFP	meristem size	5	
35S:MASP1_S670A T4-3	untagged	TUA6-GFP	meristem size	5	
35S:MASP1_S670A T4-4	untagged	TUA6-GFP	meristem size	5	
35S:MASP1_S670D T4-2	untagged	TUA6-GFP	meristem size	5	
35S:MASP1_S670D T4-6	untagged	TUA6-GFP	meristem size	5	
EGR2_pro:EGR2-YFP/egr1-1egr2-1 #6	Genomic clone with native promoter, C-terminal YFP	egr1-1egr2-1	growth assay, microscopy	4	S11, S14
EGR2_pro:EGR2-YFP/egr1-1egr2-1 #11	Genomic clone with native promoter, C-terminal YFP	egr1-1egr2-1	growth assay, microscopy	4	S11, S14
EGR2_pro:EGR2-YFP/egr1-1egr2-1 #13	Genomic clone with native promoter, C-terminal YFP	egr1-1egr2-1	growth assay, microscopy	4	S11, S14
EGR2_pro:EGR2-YFP/W.T. #6	Genomic clone with native promoter, C-terminal YFP	Col-0	growth assay, microscopy		S11
EGR2_pro:EGR2-YFP/W.T. #13	Genomic clone with native promoter, C-terminal YFP	Col-0	growth assay, microscopy		S11
MASP1_pro:MASP1-YFP/masp1-1 #2	Genomic clone with native promoter, C-terminal YFP	masp1-1	growth assay, microscopy	4	S10, S14
MASP1_pro:MASP1-YFP/masp1-1 #17	Genomic clone with native promoter, C-terminal YFP	masp1-1	growth assay, microscopy	4	S10, S14
MASP1_pro:MASP1-YFP/masp1-1 #20	Genomic clone with native promoter, C-terminal YFP	masp1-1	growth assay, microscopy	4	S10, S14
MASP1_pro:MASP1-YFP/W.T. #2	Genomic clone with native promoter, C-terminal YFP	Col-0	growth assay, microscopy		S10
MASP1_pro:MASP1-YFP/W.T #20	Genomic clone with native promoter, C-terminal YFP	Col-0	growth assay, microscopy		S10
EGR2_pro:GUS #1	Native promoter controling GUS reporter expression	Col-0	GUS staining	4	S13
EGR2_pro:GUS #3	Native promoter controling GUS reporter expression	Col-0	GUS staining	4	S13
EGR2_pro:GUS #4	Native promoter controling GUS reporter expression	Col-0	GUS staining	4	S13
MASP1pro:GUS #1	Native promoter controling GUS reporter expression	Col-0	GUS staining	4	S13
MASP1pro:GUS #6	Native promoter controling GUS reporter expression	Col-0	GUS staining	4	S13
MASP1pro:GUS #7	Native promoter controling GUS reporter expression	Col-0	GUS staining	4	S13

Parsed Citations

Barlow PW (1976) Towards an understanding of the behaviour of root meristems. Journal of Theoretical Biology 57: 433-451
Google Scholar: Author Only Title Only Author and Title

Beemster GTS, Baskin TI (1998) Analysis of Cell Division and Elongation Underlying the Developmental Acceleration of Root Growth in Arabidopsis thaliana. Plant Physiology 116: 1515-1526
Google Scholar: Author Only Title Only Author and Title

Bhaskara GB, Wen TN, Nguyen TT, Verslues PE (2017) Protein Phosphatase 2Cs and Microtubule-Associated Stress Protein 1 Control Microtubule Stability, Plant Growth, and Drought Response. Plant Cell 29: 169-191
Google Scholar: Author Only Title Only Author and Title

Bhaskara GB, Wong MM, Verslues PE (2019) The flip side of phospho-signalling: Regulation of protein dephosphorylation and the protein phosphatase 2Cs. Plant Cell and Environment 42: 2913-2930
Google Scholar: Author Only Title Only Author and Title

Biancucci M, Mattioli R, Moubayidin L, Sabatini S, Costantino P, Trovato M (2015) Proline affects the size of the root meristematic zone in Arabidopsis. BMC Plant Biology 15: 263
Google Scholar: Author Only Title Only Author and Title

Buschmann H, Dols J, Kopischke S, Peña EJ, Andrade-Navarro MA, Heinlein M, Szymanski DB, Zachgo S, Doonan JH, Lloyd CW (2015) Arabidopsis KCBP interacts with AIR9 but stays in the cortical division zone throughout mitosis via its MyTH4-FERM domain. Journal of Cell Science 128: 2033-2046
Google Scholar: Author Only Title Only Author and Title

Chaiwanon J, Wang Z-Y (2015) Spatiotemporal Brassinosteroid Signaling and Antagonism with Auxin Pattern Stem Cell Dynamics in Arabidopsis Roots. Current Biology 25: 1031-1042
Google Scholar: Author Only Title Only Author and Title

Claeys H, Van Landeghem S, Dubois M, Maleux K, Inze D (2014) What Is Stress? Dose-Response Effects in Commonly Used in Vitro Stress Assays. Plant Physiology 165: 519-527
Google Scholar: Author Only Title Only Author and Title

Clauw P, Coppens F, Korte A, Herman D, Slabbinck B, Dhondt S, Van Daele T, De Milde L, Vermeersch M, Maleux K, Maere S, Gonzalez N, Inzé D (2016) Leaf Growth Response to Mild Drought: Natural Variation in Arabidopsis Sheds Light on Trait Architecture. Plant Cell 28: 2417-2434
Google Scholar: Author Only Title Only Author and Title

Colon-Carmona A, You R, Haimovitch-Gal T, Doerner P (1999) Spatio-temporal analysis of mitotic activity with a labile cyclin-GUS fusion protein. Plant Journal 20: 503-508
Google Scholar: Author Only Title Only Author and Title

Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR (2010) Abscisic Acid: Emergence of a Core Signaling Network. In S Merchant, WR Briggs, D Ort, eds, Annual Review of Plant Biology, Vol 61, Vol 61, pp 651-679
Google Scholar: Author Only Title Only Author and Title

Di Mambro R, De Ruvo M, Pacifici E, Salvi E, Sozzani R, Benfey PN, Busch W, Novak O, Ljung K, Di Paola L, Maree AFM, Costantino P, Grieneisen VA, Sabatini S (2017) Auxin minimum triggers the developmental switch from cell division to cell differentiation in the Arabidopsis root. Proceedings of the National Academy of Sciences of the United States of America 114: E7641-E7649
Google Scholar: Author Only Title Only Author and Title

Dickinson AJ, Lehner K, Mi J, Jia K-P, Mijar M, Dinneny J, Al-Babili S, Benfey PN (2019) β-Cyclocitral is a conserved root growth regulator. Proceedings of the National Academy of Sciences 116: 10563-10567
Google Scholar: Author Only Title Only Author and Title

Dietrich D, Pang L, Kobayashi A, Fozard JA, Boudolf V, Bhosale R, Antoni R, Nguyen T, Hiratsuka S, Fujii N, Miyazawa Y, Bae TW, Wells DM, Owen MR, Band LR, Dyson RJ, Jensen OE, King JR, Tracy SR, Sturrock CJ, Mooney SJ, Roberts JA, Bhalerao RP, Dinneny JR, Rodriguez PL, Nagatani A, Hosokawa Y, Baskin TI, Pridmore TP, De Veylder L, Takahashi H, Bennett MJ (2017) Root hydrotropism is controlled via a cortex-specific growth mechanism. Nature Plants 3: 17057
Google Scholar: Author Only Title Only Author and Title

Ding YL, Lv J, Shi YT, Gao JP, Hua J, Song CP, Gong ZZ, Yang SH (2019) EGR2 phosphatase regulates OST1 kinase activity and freezing tolerance in Arabidopsis. EMBO Journal 38: e99819
Google Scholar: Author Only Title Only Author and Title

Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J, Benfey PN (2008) Cell identity mediates the response of Arabidopsis roots to abiotic stress. Science 320: 942-945
Google Scholar: Author Only Title Only Author and Title

Dubois M, Selden K, Bediee A, Rolland G, Baumberger N, Noir S, Bach L, Lamy G, Granier C, Genschik P (2018) SIAMESE-RELATED1 Is Regulated Posttranslationally and Participates in Repression of Leaf Growth under Moderate Drought. Plant Physiology 176: 2834-

2850

Google Scholar: Author Only Title Only Author and Title

Gonzalez N, Vanhaeren H, Inze D (2012) Leaf size control: complex coordination of cell division and expansion. Trends in Plant Science 17: 332-340

Google Scholar: Author Only Title Only Author and Title

Heidstra R, Sabatini S (2014) Plant and animal stem cells: similar yet different. Nature Reviews Molecular Cell Biology 15: 301-312

Google Scholar: Author Only Title Only Author and Title

Iyer-Pascuzzi AS, Jackson T, Cui HC, Petricka JJ, Busch W, Tsukagoshi H, Benfey PN (2011) Cell Identity Regulators Link Development and Stress Responses in the Arabidopsis Root. Developmental Cell 21: 770-782

Google Scholar: Author Only Title Only Author and Title

Mielke S, Zimmer M, Meena MK, Dreos R, Stellmach H, Hause B, Voiniciuc C,

Gasperini D (2021) Jasmonate biosynthesis arising from altered cell walls is

prompted by turgor-driven mechanical compression. Science Advances 7: eabf0356.

Moubayidin L, Di Mambro R, Sozzani R, Pacifici E, Salvi E, Terpstra I, Bao DP, van Dijken A, Dello Ioio R, Perilli S, Ljung K, Benfey PN, Heidstra R, Costantino P, Sabatini S (2013) Spatial Coordination between Stem Cell Activity and Cell Differentiation in the Root Meristem. Developmental Cell 26: 405-415

Google Scholar: Author Only Title Only Author and Title

Munns R, Tester M (2008) Mechanisms of salinity tolerance. Annual Review of Plant Biology 59: 651-681

Google Scholar: Author Only Title Only Author and Title

Raghavendra AS, Gonugunta VK, Christmann A, Grill E (2010) ABA perception and signalling. Trends in Plant Science 15: 395-401

Google Scholar: Author Only Title Only Author and Title

Sacks MM, Silk WK, Burman P (1997) Effect of Water Stress on Cortical Cell Division Rates within the Apical Meristem of Primary Roots of Maize. Plant Physiology 114: 519-527

Google Scholar: Author Only Title Only Author and Title

Salazar-Blas A, Noriega-Calixto L, Campos ME, Eapen D, Cruz-Vazquez T, Castillo-Olamendi L, Sepulveda-Jimenez G, Porta H, Dubrovsky JG, Cassab GI (2017) Robust root growth in altered hydrotropic response 1 (ahr1) mutant of Arabidopsis is maintained by high rate of cell production at low water potential gradient. Journal of Plant Physiology 208: 102-114

Google Scholar: Author Only Title Only Author and Title

Salvi E, Rutten JP, Di Mambro R, Polverari L, Licursi V, Negri R, Dello Ioio R, Sabatini S, Ten Tusscher K (2020) A Self-Organized PLT/Auxin/ARR-B Network Controls the Dynamics of Root Zonation Development in Arabidopsis thaliana. Developmental Cell 53: 431-443.e423

Google Scholar: Author Only Title Only Author and Title

Schnittger A, De Veylder L (2018) The Dual Face of Cyclin B1. Trends in Plant Science 23: 475-478

Google Scholar: Author Only Title Only Author and Title

Sharp RE, Silk WK, Hsiao TC (1988) Growth of the maize primary root at low water potentials: I. Spatial distribution of expansive growth. Plant Physiology 87: 50-57

Google Scholar: Author Only Title Only Author and Title

Shimotohno A, Scheres B (2019) Topology of regulatory networks that guide plant meristem activity: similarities and differences. Current Opinion in Plant Biology 51: 74-80

Google Scholar: Author Only Title Only Author and Title

Skirycz A, Claeys H, De Bodt S, Oikawa A, Shinoda S, Andriankaja M, Maleux K, Eloy NB, Coppens F, Yoo SD, Saito K, Inze D (2011) Pause-and-Stop: The Effects of Osmotic Stress on Cell Proliferation during Early Leaf Development in Arabidopsis and a Role for Ethylene Signaling in Cell Cycle Arrest. Plant Cell 23: 1876-1888

Google Scholar: Author Only Title Only Author and Title

Skirycz A, Inzé D (2010) More from less: plant growth under limited water. Current Opinion in Biotechnology 21: 197-203

Google Scholar: Author Only Title Only Author and Title

Slovak R, Setzer C, Roiuk M, Bertels J, Goschl C, Jandrasits K, Beemster GTS, Busch W (2020) Ribosome assembly factor Adenylate Kinase 6 maintains cell proliferation and cell size homeostasis during root growth. New Phytologist 225: 2064-2076

Google Scholar: Author Only Title Only Author and Title

Spartz AK, Ren H, Park MY, Grandt KN, Lee SH, Murphy AS, Sussman MR, Overvoorde PJ, Gray WM (2014) SAUR Inhibition of PP2C-D Phosphatases Activates Plasma Membrane H+-ATPases to Promote Cell Expansion in Arabidopsis. Plant Cell 26: 2129-2142

Google Scholar: Author Only Title Only Author and Title

van der Weele CM, Spollen WG, Sharp RE, Baskin TI (2000) Growth of Arabidopsis thaliana seedlings under water deficit studied by control of water potential in nutrient-agar media. Journal of Experimental Botany 51: 1555-1562

Google Scholar: Author Only Title Only Author and Title

Verslues PE (2017) Time to grow: factors that control plant growth during mild to moderate drought stress. Plant Cell and Environment 40: 177-179
Google Scholar: Author Only Title Only Author and Title

Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu JK (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. Plant Journal 45: 523-539
Google Scholar: Author Only Title Only Author and Title

Voothuluru P, Makela P, Zhu JM, Yamaguchi M, Cho IJ, Oliver MJ, Simmonds J, Sharp RE (2020) Apoplastic Hydrogen Peroxide in the Growth Zone of the Maize Primary Root. Increased Levels Differentially Modulate Root Elongation Under Well-Watered and Water-Stressed Conditions. Frontiers in Plant Science 11: 392.
Google Scholar: Author Only Title Only Author and Title

Vragovic K, Sela A, Friedlander-Shani L, Fridman Y, Hacham Y, Holland N, Bartom E, Mockler TC, Savaldi-Goldstein S (2015) Translatome analyses capture of opposing tissue-specific brassinosteroid signals orchestrating root meristem differentiation. Proceedings of the National Academy of Sciences of the United States of America 112: 923-928
Google Scholar: Author Only Title Only Author and Title

Wendrich JR, Moller BK, Li S, Saiga S, Sozzani R, Benfey PN, De Rybel B, Weijers D (2017) Framework for gradual progression of cell ontogeny in the Arabidopsis root meristem. Proceedings of the National Academy of Sciences of the United States of America 114: E8922-E8929
Google Scholar: Author Only Title Only Author and Title

Wendrich JR, Yang B, Vandamme N, Verstaen K, Smet W, Van de Velde C, Minne M, Wybouw B, Mor E, Arents HE, Nolf J, Van Duyse J, Van Isterdael G, Maere S, Saeys Y, De Rybel B (2020) Vascular transcription factors guide plant epidermal responses to limiting phosphate conditions. Science 370: eaay4970
Google Scholar: Author Only Title Only Author and Title

West G, Inzé D, Beemster GTS (2004) Cell Cycle Modulation in the Response of the Primary Root of Arabidopsis to Salt Stress. Plant Physiology 135: 1050-1058
Google Scholar: Author Only Title Only Author and Title

Wong MM, Bhaskara GB, Wen TN, Lin WD, Nguyen TT, Chong GL, Verslues PE (2019) Phosphoproteomics of Arabidopsis Highly ABA-Induced1 identifies AT-Hook-Like10 phosphorylation required for stress growth regulation. Proceedings of the National Academy of Sciences of the United States of America 116: 2354-2363
Google Scholar: Author Only Title Only Author and Title

Yamada M, Han XW, Benfey PN (2020) RGF1 controls root meristem size through ROS signalling. Nature 577: 85-+
Google Scholar: Author Only Title Only Author and Title

Yang XL, Dong G, Palaniappan K, Mi GH, Baskin TI (2017) Temperature-compensated cell production rate and elongation zone length in the root of Arabidopsis thaliana. Plant Cell and Environment 40: 264-276
Google Scholar: Author Only Title Only Author and Title