# Reservoir host immunology and life history shape virulence evolution in zoonotic viruses

**Authors:**
Cara E. Brook[1], Carly Rozins[2], Sarah Guth[3], Mike Boots[3,4]

**Affiliations:**
[1] Department of Ecology and Evolution, University of Chicago, Chicago, IL, USA
[2] Department of Science and Technology Studies, York University, Toronto, Canada
[3] Department of Integrative Biology, University of California, Berkeley, Berkeley, CA, USA
[4] Biosciences, University of Exeter, Penryn Campus, UK. TR10 9FE

**Corresponding author:** Cara E. Brook
**Email**: cbrook@uchicago.edu

**Author contributions:**
CEB and MB conceptualized the research. CEB carried out the modeling and analysis with contributions from CR and SG. CEB and CR wrote the materials and methods and Supplementary Appendix. CEB wrote the first draft of the main manuscript, which all authors edited and revised.

**Competing interest statement:** The authors have no competing interests to declare.

**Classification:** Biological sciences: Population biology.

**Key Words**
bat virus, spillover, virulence evolution, tradeoff theory

**Abstract**
The management of future pandemic risk requires a better understanding of the mechanisms that
determine the virulence of emerging zoonotic viruses. Meta-analyses suggest that the virulence of
emerging zoonoses is correlated with but not completely predictable from reservoir host phylogeny,
indicating that specific characteristics of reservoir host immunology and life history may drive the
evolution of viral traits responsible for cross-species virulence. In particular, bats host viruses that cause
higher case fatality rates upon spillover to humans than those derived from any other mammal, a
phenomenon that cannot be explained by phylogenetic distance alone. In order to disentangle the
fundamental drivers of these patterns, we develop a nested modeling framework that pinpoints
mechanisms which underpin the evolution of viral traits in reservoir hosts that cause virulence following
cross-species emergence. We apply this framework to generate virulence predictions for viral zoonoses
derived from diverse mammalian reservoirs, successfully recapturing corresponding virus-induced human
mortality rates reported in the literature. Notably, our work offers a mechanistic explanation for the
extreme virulence of bat-borne zoonoses and, more generally, demonstrates how key differences in
reservoir host longevity, viral tolerance, and constitutive immunity impact the evolution of viral traits that
cause virulence following spillover to humans. Our theoretical framework offers a series of testable
questions and hypotheses designed to stimulate future work comparing cross-species virulence evolution
in zoonotic viruses derived from diverse mammalian hosts.
71
72
**Significance Statement**
Bats are the natural reservoir hosts for zoonotic viruses that cause higher case fatality rates in humans
than do viruses derived from any other mammal or bird host. Using a nested population-level and within-
host modelling approach, we generate virulence predictions for viral zoonoses derived from diverse
mammalian reservoirs, successfully reproducing human case fatality rates from zoonoses reported in the
literature. Our model provides a mechanistic explanation for the virulence of bat-borne zoonoses and a
general predictive framework to forecast the virulence of cross-species viral infections from diverse
mammalian hosts.

109    **Main Text**
110
**Introduction**
The devastating impact of the SARS-CoV-2 pandemic highlights the extreme public health outcomes that
can result upon cross-species emergence of zoonotic viruses. Estimating the relative threats posed by
potential future zoonoses is an important but challenging public health undertaking. In particular, efforts to
predict the virulence of emerging viruses can be complicated since chance will always play a role in
dictating the initial spillover that precedes selection (1), virulence upon emergence may be maladaptive in
novel hosts (2, 3), and patterns in available data may be muddled by attainment bias if avirulent infections
go underreported (1). Nonetheless, a growing body of recent work highlights clear associations between
reservoir and spillover host phylogeny and the virulence of a corresponding cross-species infection (4–7).
In many cases, increasing phylogenetic distance between reservoir and spillover hosts is correlated with
higher virulence infections (4–6), suggesting that spillover host immune systems may be poorly equipped
to tolerate viral traits optimized in more distantly-related reservoirs. Still, the effect of phylogeny on
spillover virulence appears to supersede that of phylogenetic distance alone (7, 8), indicating that taxon-
specific reservoir host immunological and life history traits may be important drivers of cross-species virus
virulence. Notably, bats host viruses that cause higher human case fatality rates than zoonoses derived
from other mammals and birds, a phenomenon that cannot be explained by phylogenetic distance alone
(8). Understanding the mechanisms that select for the evolution of unique viral traits in bats compared to
those selected in other mammalian reservoirs should enable us to better predict the virulence of future
zoonotic threats.
130
Although the disproportionate frequency with which the Chiropteran order may source viral zoonoses
remains debated (9, 10), the extraordinary human pathology induced by many bat-borne zoonoses—
including Ebola and Marburg filoviruses, Hendra and Nipah henipaviruses, and SARS, MERS, and
SARS-CoV-2 coronaviruses (11)—is not contested. Remarkably, bats demonstrate limited clinical
pathology from infection with viruses that cause extreme morbidity and mortality in other hosts (12). Bats
avoid pathological outcomes from viral infection via a combination of unique resistance and tolerance
mechanisms, which, respectively, limit the viral load accrued during infection ('resistance') and reduce the
disease consequences of a given viral load ('tolerance') (13–16). Viral resistance mechanisms vary
considerably across bat species; those described to date include: receptor incompatibilities that limit the
extent of infection for certain viruses in certain bats (17–20), constitutive expression of antiviral cytokines
in some bat species (21), and enhanced autophagy (22) and heat-shock protein expression (23) in
others. While these extensive inflammatory responses would be pathological in most mammals, bats—as
the only mammals capable of powered flight—have evolved numerous unique mechanisms of mitigating
inflammation incurred during the intensive physiological process of flying (24–26). These anti-
inflammatory adaptations include loss of PYHIN (27–29) and downregulation of NLRP3 (30)
inflammasome-forming gene families, dampened STING-dependent interferon activation (31), and
diminished caspase-1 inflammatory signaling (32). In addition to facilitating flight, this resilience to
inflammation has yielded the apparent by-products of extraordinarily long bat lifespans (33) and tolerance
of the immunopathology that typically results from viral infection (11). Moreover, recent work
demonstrates how high virus growth rates easily tolerated in constitutively antiviral bat cells cause
significant pathology in cells lacking these constitutive antiviral defenses (34). The extent to which
inflammatory tolerance may modulate the evolution of the viruses that bats host, however, remains
largely unexplored.
154
Modern theory on the evolution of virulence typically assumes, either explicitly or implicitly, that high
pathogen growth rates should both enhance between-host transmission and elevate infection-induced
morbidity or mortality, resulting in a trade-off between virulence and transmission (35–37). Theory further
suggests that because viral 'tolerance' mitigates virulence without reducing viral load, most host
strategies of tolerance should select for higher growth rate pathogens that achieve gains in between-host
transmission without causing damage to the original host (37–39) (Fig. 1). The widely touted viral
tolerance of bats (11, 16, 40–42) should therefore be expected to support the evolution of enhanced viral
growth rates, which—though avirulent to bats—may cause significant pathology upon spillover to hosts
lacking unique features of bat immunology and physiology. Beyond tolerance, other life history
characteristics unique to diverse reservoir hosts likely also impact the evolution of traits in the viruses they

host—with important consequences for cross-species virulence following spillover. To date, we lack a
specific theory that examines the relative impact of reservoir host life history on spillover virulence. Here,
we explore the extent to which the immunological and life-history traits of mammalian reservoirs can
explain variation in the virulence of zoonotic viruses emerging into human hosts.
169

170
**Figure 1. Conceptual mechanistic framework to predict zoonotic virus virulence from reservoir immunology and life**
**history traits.** (**A**) A within-host predator-prey model of leukocyte-virus dynamics is embedded in a population-level transmission
model for reservoir hosts. Between-host rates of transmission ($\beta_r$) and virulence ($\alpha_r$) are expressed as functions of within-host
dynamics to derive optimal virus growth rates in the reservoir ($r_R^*$). (**B**) Virus growth rates ($r_R^*$) evolve to maximize gains in between-
host transmission ($\beta_R^*$) over virulence incurred on the reservoir ($\alpha_R^*$). Viruses evolved in taxa with physiologies similar to humans (i.e.
non-human primates) should optimize at relatively low growth rates ($r_R^*$) that yield minimal pathology for the reservoir host ($\alpha_R^*$) and
are, correspondingly, likely to cause only minimal pathology in the spillover human host ($\alpha_S$). (**C**) As a result of unique bat virus
tolerance, viruses evolved in bat reservoir hosts may optimize at high $r_R^*$ values that maximize bat-to-bat transmission ($\beta_R^*$) but cause
only minimal pathology in the bat host ($\alpha_R^*$) – but are likely to generate extreme virulence upon spillover ($\alpha_S$) to secondary hosts,
including humans, that lack bat life history traits.
181
182
183 **Results**
184
***General trends in the evolution of high growth rate viruses.***
To elucidate how immunological and life-history traits of mammalian hosts combine to drive zoonotic virus
virulence, we adopted a nested modeling approach, embedding a simple within-host model of viral and
leukocyte dynamics within an epidemiological, population-level framework (Fig. 1). We examined how the
life history traits of a primary reservoir drive the evolution of viral traits likely to cause pathology in a
secondary, spillover host—chiefly, a human. Using our nested model in an adaptive dynamics framework
(43), we first expressed the conditions—called the 'invasion fitness'—which permit invasion of an
evolutionarily 'fitter', mutant virus into a reservoir host system in within-host terms. This allowed us to
calculate the optimal within-host virus growth rate ($r^*$) for a virus evolved in reservoir hosts that differ
across a suite of life history traits and corresponding cellular and immunological parameters (Fig. 2; Table
1; *SI Appendix* Fig S1-S2). In part with this optimization, we also calculated the corresponding
transmission ($\beta^*$) and virulence ($\alpha^*$) for the reservoir host at the optimal $r^*$(Fig. 2; Table 1; *SI Appendix*
Fig S1-S2). We derive the equation for the optimal virus growth as follows:
198

$$r^* = \frac{cg_0}{m} + \frac{\sqrt{c^2 m^2 g g_0 \mu T_v T_w (v T_w + g w T_v)}}{m^2 (v T_w + g w T_v)},$$

where $\mu$ signifies the natural mortality rate of the reservoir host, and all other parameters represent within-
host viral and immune dynamics in a persistently infected reservoir. Thus, $c$ corresponds to the rate of
virus clearance by leukocytes, $g_0$ is the magnitude of constitutive immunity, $m$ is the natural leukocyte
mortality rate, and $g$ is the rate of leukocyte activation following infection. The parameters $v$ and $w$
correspond, respectively, to the intrinsic virulence of the virus and the reservoir host's natural
susceptibility to inflammatory damage, while $T_v$ and $T_w$ represent reservoir host tolerance to direct virus
pathology and to immunopathology. We present all main text results under assumptions by which
tolerance manifests as a *constant* reduction of either direct virus-induced pathology ($T_v$) or
immunopathology ($T_w$). See *SI Appendix* for comparable results under assumptions of *complete*
tolerance, whereby tolerance completely eliminates virus pathology and immunopathology up to a
threshold value, beyond which pathology scales proportionally with virus and immune cell growth.
212

**Figure 2. Optimal virus growth rates vary across reservoir host immunological and life history parameters.** Rows (top-down)
indicate the evolutionarily optimal within-host virus growth rate ($r^*$) and the corresponding between-host transmission rate ($\beta^*$), and
virus-induced mortality rate ($\alpha^*$) for a reservoir host infected with a virus at $r^*$. Columns demonstrate the dependency of these
outcomes on variable reservoir host parameters: background mortality ($\mu$), magnitude of constitutive immunity ($g_0$), rate of leukocyte
activation upon viral contact ($g$), rate of virus consumption by leukocytes ($c$), and leukocyte mortality rate ($m$). Darker colored lines
depict outcomes at higher tolerance of direct virus pathology ($T_v$, red) or immunopathology ($T_w$, blue), assuming no tolerance of the
opposing type. Heat maps demonstrate how $T_v$ and $T_w$ interact to produce each outcome. Parameter values are reported in Table 1.
Figure assumes tolerance in the 'constant' form (see *SI Appendix,* Fig. S2 for 'complete' tolerance assumptions).
222

Our analyses highlight several critical drivers of virus evolution likely to generate significant pathology
following spillover to a secondary host (Fig. 2): higher within-host virus growth rates ($r^*$) are selected in
reservoir hosts with higher background mortality ($\mu$), elevated constitutive immune responses ($g_0$), more
rapid leukocyte activation upon infection ($g$), and more rapid virus clearance by the host immune system
($c$). Additionally, higher $r^*$ viruses are selected in hosts exhibiting lower leukocyte mortality rates ($m$),
resulting in longer-lived immune cells. Critically, higher tolerance of both virus-induced pathology ($T_v$) and
immunopathology ($T_w$) also select for higher $r^*$ when virulence due to direct virus damage and to
immunopathology are modeled, respectively, as proportional to virus and immune cell growth rates (*SI*
*Appendix*). In keeping with trade-off theory, changes in the majority of within-host parameters drive
corresponding increases in $r^*$, $\beta^*$, and $\alpha^*$, such that viruses evolved to high growth rates experience high
transmission within the reservoir host population—but also generate high virus-induced mortality (Fig. 1).
By definition, the two modeled mechanisms of tolerance ($T_v$ and $T_w$) decouple the relationship between
transmission and virulence, permitting evolution of high $r^*$ viruses that achieve gains in between-host
transmission in the reservoir population ($\beta^*$), while simultaneously incurring minimal virulence ($\alpha^*$) on
reservoir hosts. By extension, we predict that viruses evolving high optimal $r^*$ values in reservoir hosts

with high $T_v$ or $T_w$ might incur substantial pathology upon spillover to secondary hosts exhibiting reduced
tolerance as compared with the reservoir.
241
***Order-specific estimates for optimal virus growth rates evolved in reservoir hosts.***
We next applied our model to make broad predictions of the evolution of optimal virus growth rates across
diverse mammalian reservoir orders, based on order-specific variation in three key parameters from our
nested model: the reservoir host background mortality rate ($\mu$), the reservoir host tolerance of
immunopathology ($T_w$), and the magnitude of constitutive immunity ($g_0$). Within-host immunological data
needed to quantify these parameters is lacking across most mammalian orders; thus, we applied
generalized additive models (GAMs (44)) to mammalian life history data—in particular, well-described
allometric relationships between mammalian body mass and basal metabolic rate (BMR) with lifespan
and immune cell concentrations (45–49)—to proxy $\mu$, $T_w$, and $g_0$ across diverse mammalian orders (Fig.
3A-C; Table 1; *SI Appendix,* Fig. S3-S4).
252
Fitted GAMs demonstrated a statistically significant positive association between the orders Chiroptera,
Cingulata, Monotremata, and Primates and the response variable of maximum longevity, indicating that
species in these orders exhibit, on average, longer lifespans than predicted for their body sizes (Fig. 3A;
*SI Appendix,* Fig. S3; Table S1-S2). By contrast, six mammalian orders (Afrosoricida, Dasyuromorphia,
Didelphimorphia, Eulipotyphla, Notoryctemorphia, and Peramelemorphia) showed significantly shorter
lifespans than predicted for body size (Fig. 3A; *SI Appendix,* Fig. S3; Table S1-S2). Fitted GAMs further
demonstrated a statistically significant positive association between the orders Chiroptera and
Monotremata and the response variable of baseline neutrophil concentration, indicating that species in
these orders may have more enhanced constitutive immune responses than predicted by mass-specific
BMR. The orders Cetartiodactyla, Dasyuromorphia, Diprotodontia, and Scandentia, by contrast, showed
significant negative associations, representing lower baseline neutrophil concentrations than predicted
per mass-specific BMR (Fig. 3B; *SI Appendix,* Fig. S3; Table S1-S2).
265

266
267  **Figure 3. Reservoir host life history traits predict evolution of zoonotic virus virulence.** (**A**) Variation in log₁₀ maximum
268  lifespan (y-axis, in years) with log₁₀ adult body mass (x-axis, in g) across mammals, with data derived from Jones et al. 2009 (56)
269  and Healy et al. 2014 (57). Points are colored by mammalian order, corresponding to legend. Black line depicts predictions of
270  mammalian lifespan per body mass, summarized from fitted GAM (but excluding random effect of mammalian order), presented in

We generated an estimate of background mortality ($\mu$) for 26 disparate mammalian orders by taking the
inverse of predicted lifespan (in days) per order from our first fitted GAM, excluding the effects of body
size (*SI Appendix,* Fig. S4; Table S1). Since longevity and resilience to inflammation are linked
extensively in the literature (11, 42, 50–53), we represented our $T_w$ parameter as the scaled effect of
these 26 mammalian orders on the predicted lifespan per body size—such that orders longer-lived than
predicted were modeled with high values for tolerance to immunopathology and vice versa. As a result,
order-level estimates for $\mu$ and $T_w$ were mirror inverses of one another: bats, for instance, demonstrated
the lowest predicted values across all orders for background mortality ($\mu$) and the highest predicted
values for tolerance of immunopathology ($T_w$) (*SI Appendix,* Fig. S4; Table S1). Because our nested
model expressed parameters in timesteps most relevant to viral dynamics (days), differences in mortality
rates across mammalian orders—though substantial on multi-year timescales of the host—had limited
influence on downstream predictions of differences in the evolution of virus growth rates. This result
echoed prior observations from Fig. 2, which showed reduced sensitivity of $r^*$ to realistic variation in the
magnitude of $\mu$, as compared with other parameters.

Building from our approach for $T_w$, we next generated order-level estimates for the parameter $g_0$, which
represents the magnitude of constitutive immune response of the reservoir host in our nested modeling
framework. We calculated $g_0$ as the scaled effect of mammalian order on the prediction of neutrophil
concentration per mass-specific metabolic rate from our second fitted GAM for the 19 orders for which we
possessed paired neutrophil and BMR data (*SI Appendix,* Fig. S4; Table S1). Orders Chiroptera and
Monotremata, which showed significantly higher neutrophil concentrations than predicted by mass-
specific BMR, were thus modeled with high $g_0$, while orders Cetartiodactyla, Dasyuromorphia,
Diprotodontia, and Scandentia, which exhibited lower neutrophil concentrations than predicted, were
modeled with low $g_0$ (*SI Appendix,* Fig. S4; Table S1).

Finally, we combined order-specific parameters for $\mu$, $T_w$, and $g_0$ with fixed parameter values for all other
within-host terms to generate an estimate of the optimal growth rate of a virus evolved in the 19
mammalian orders for which we possessed proxy data for all three variable within-host parameters (Table
1; *SI Appendix,* Fig. S5; Table S1). We termed this reservoir-host optimized virus growth rate, $r_R^*$ (Fig. 1B-
C). In keeping with predictions from our general model (Fig. 2), we predicted the evolution of high $r_R^*$
viruses from reservoir orders exhibiting high $\mu$, $T_w$, and/or $g_0$. Since $\mu$ and $T_w$ were inversely correlated
but within-host dynamics more sensitive to changes in $T_w$, we ultimately recovered the highest predicted
optimal growth rates for viruses evolved in the orders Chiroptera and Monotremata, which exhibited both
long lifespans per body size and high baseline neutrophil concentrations. Data were particularly sparse,
however, for order Monotremata, for which complete records were only available for two species (the
platypus and the short-beaked echidna). As a result, predictions for this order should be interpreted with
caution. More data, particularly for baseline neutrophil concentrations, will be needed to evaluate the

extent to which these predictions hold across all five extant species in the Monotremata order.
Additionally, we predicted the evolution of the lowest growth rate viruses in reservoir hosts of the orders
Cetartiodactyla, Dasyuromorphia, Diprotodontia, and Scandentia, largely as a result of significantly low
values estimated for the magnitudes of constitutive immunity in these orders.
334
***Estimating zoonotic virus virulence in spillover human hosts.***
After establishing optimal growth rates for viruses evolved in diverse reservoir host orders ($r_R^*$), we
subsequently modeled the corresponding 'spillover virulence' ($\alpha_S$) of these viruses following emergence
into a human host. Zoonotic spillovers were modeled as acute infections in the human, and virulence was
calculated while varying only the growth rate of the spillover virus ($r_R^*$) and the human tolerance of direct
virus pathology ($T_{vS}$) between viruses evolved in differing reservoir orders. We varied this last parameter,
$T_{vS}$, to account for any differences in virus adaptation to reservoir host immune systems, which were not
already captured in estimation of reservoir host $T_w$ and $g_0$. $T_{vS}$ was thus computed as the inverse of the
scaled time to most recent common ancestor (MRCA) for each mammalian reservoir host order from
Primates (Fig. 3C; *SI Appendix,* Fig. S4, Table S1-S2), such that we estimated low human tolerance to
viruses evolved in phylogenetically distant orders (e.g. monotreme and marsupial orders), and high
human tolerance to viruses evolved in Primate and Primate-adjacent orders. In general, the modulating
effects of $T_{vS}$ did little to alter virulence rankings of zoonotic viruses from those predicted by raw growth
rate ($r_R^*$) alone—with the highest spillover virulence predicted from viruses evolved in orders Monotremata
and Chiroptera and the lowest spillover virulence predicted from viruses evolved in orders Cetartiodactyla
and Scandentia. Notably, modulating $T_{vS}$ enhanced predictions of spillover virulence for the marsupial
clades (Peramelemorphia, Didelphimorphia, Dasyuromorphia, Diprotodontia) relative to eutherian orders
with similar predicted $r_R^*$ values. This resulted in dampened predicted virulence for the eutherian order
Cetartiodactyla as compared with marsupial orders Dasyuromorphia and Diprotodontia, despite lower
predicted $r_R^*$ values (*SI Appendix,* Fig. S5).
355
***Comparing predictions of spillover virulence by reservoir order with estimates from the literature.***
Since parameter magnitudes estimated from life history traits were largely relative in nature, we scaled
predictions of spillover virulence ($\alpha_S$) by reservoir order in relative terms to compare with estimates
gleaned from the zoonosis literature (8) (Fig. 3D). Among the eight reservoir orders for which we were
able to both generate nested modeling predictions and derive estimates from the literature (8), our nested
modeling approach performed exceptionally well. Critically, we successfully recovered the key result from
the zoonosis literature: bat-derived zoonoses yield higher rates of virus-induced mortality upon spillover to
humans ($\alpha_S$) than do viruses derived from all other eutherian mammals (Fig. 3D; *SI Appendix,* Fig. S6-
S7). In general, high estimates for $T_w$, $\mu$, and $g_0$ and low estimates for $T_{vS}$ predicted high spillover
virulence to humans ($\alpha_S$). Bats demonstrate uniquely long lifespans for their body sizes and uniquely
enhanced constitutive immune responses (21); when combined, as in our analysis, to represent high $T_w$
and high $g_0$, these reservoir host traits elevated predicted $r_R^*$ and $\alpha_S$ beyond all other eutherian orders.
Evaluated against the data (8), our model underpredicted cross-order comparative virulence rankings for
Carnivora-derived viruses, though the discrepancy in our predictions vs. those from the literature was
somewhat resolved when rabies lyssavirus was excluded from data comparisons (*SI Appendix,* Fig. S7)
(8). Though most rabies zoonoses are sourced from domestic dogs, lyssaviruses are Chiropteran by
origin (54), and viral traits responsible for rabies' virulence may reflect its bat evolutionary history more
than that of any intermediate carnivore host. Our model also successfully reproduced estimates of
significantly lower virulence incurred by zoonotic viruses evolved in Cetartiodactyla hosts (8). While
previous work suggested that this effect might result from overreporting of avirulent zoonoses in domestic
livestock with frequent human contact (8), our analysis indicates that viral zoonoses emerging from
Cetartiodactylan hosts may truly be less virulent to humans. Our mechanistic framework demonstrates
that limited constitutive immune responses identified in Cetartiodactylan hosts may drive the evolution of
low growth rate viruses that cause correspondingly benign infections following zoonotic emergence.
Further research into the extent to which baseline neutrophil concentrations accurately reflect
Cetartiodactylan innate immunity, as well as quantification of the natural growth rates of Cetartiodactylan
viruses, is needed to evaluate the accuracy of these predictions.
383
384    **Discussion**
385

Our work provides a mechanistic insight into the biological processes that underpin cross-species
patterns in the evolution of virus virulence—a major advance for efforts to evaluate zoonotic risk. Using a
simple model of evolving virus in a reservoir host immune system, we successfully recapture patterns
previously reported in the literature that document both the distinctive virulence of viral zoonoses derived
from bats and the surprising avirulence of viruses derived from Cetartiodactylan hosts (8). Additionally,
our mechanistic approach allows us to make powerful predictions about the virulence of potential future
viral spillovers, using general life history traits across mammals, including from orders for which zoonoses
have not yet been reported (8). Our model indicates that we should anticipate the evolution of high growth
rate viruses likely to cause virulence upon cross-species emergence from mammalian hosts with
protracted lifespans for their body size (which we link to molecular mechanisms of immunopathological
tolerance (49)), as well as from hosts with robust constitutive immune responses (34). While both
immunopathological tolerance and constitutive immunity should drive the evolution of high growth rate
viruses (Fig. 2), only tolerance will do so in the absence of reservoir host pathology, thus highlighting its
selective influence in driving observed variation in the virulence of viral zoonoses. Importantly, while
tolerance reduces pathogen-induced mortality for a single host, tolerant host populations (with limited
checks on virus transmission) may demonstrate high pathogen prevalence. If imperfectly tolerant hosts
still experience some virus-induced pathology, high prevalence can consequently elevate total population-
level mortality for infected hosts—a phenomenon known as the 'tragedy of tolerance' (38, 39). Reports of
virus-induced mortality in bats are rare (55), suggesting that bat virus tolerance is likely very effective.
405
Intriguingly, our independent evaluations of mammalian life history traits associated with enhanced
longevity (a proxy for immunopathological tolerance (56, 57)) and robust constitutive immunity (48, 58)
highlight several mammalian orders—chiefly Chiroptera and Monotremata—which show synergy in both
features, offering support for ongoing efforts to elucidate links between anti-aging molecular pathways
and antiviral immunity (52, 55). The order Cetartiodactyla presents a possible exception to this synergy
that merits further study: on average, Cetartiodactylans demonstrate predictable lifespans per body size
but significantly reduced constitutive immune responses as compared with other mammals. In all cases,
predictions are limited by a dearth of comparative data in baseline immune cell concentrations across
mammalian orders; notably, however, our estimation of enhanced constitutive immune responses in
bats—derived from baseline neutrophil concentration for just ten species—is robustly supported by
independent molecular work describing constitutive interferon expression in bat cells across multiple
species (21, 59, 60).
418
Notably, we are able to recapture broad patterns in the literature while holding constant many basic
immunological parameters—including rates of virus consumption by leukocytes ($c$), leukocyte activation
upon viral infection ($g$), and reservoir host tolerance of direct virus pathology ($T_v$)—which almost certainly
vary across taxa. Host tolerance of direct virus pathology, in particular, will likely also be modulated by
virus tropism, offering another mechanism by which viral adaptation to reservoir hosts could enhance
spillover virulence. For example, coronavirus tropism is thought to be localized largely in the
gastrointestinal tract for bat hosts (61, 62): in our modeling framework, higher tolerance of direct virus
pathology in this tissue should promote the evolution of higher growth rate viruses in bats, which are likely
to cause virulence upon infection of more vulnerable tissues, such as the respiratory tract, in spillover
hosts.
429
By summarizing within-host traits across mammalian orders, we generalize substantial within-clade
diversity that likely contributes to heterogeneous patterns in available data. Bats alone make up more
than 1,400 species and account for some 20% of mammalian diversity (63). Only a subset of bats are
long-lived (64), and the magnitude of constitutive immunity is known to vary across bat species (21, 34,
42), suggesting considerable variation in the parameters $\mu$, $T_w$, and $g_0$, which we here model universally
across the entire Chiropteran order. Thus, we can expect considerable variation in the evolution of
virulence for viruses evolved in diverse species within a given order—which we largely disregard here. As
more data becomes available, our modeling approach could be fine-tuned to make more specific,
species-level predictions of highly virulent disease risk.
439
Our analysis does not consider the probability of cross species viral emergence, or the potential for
onward transmission of a spilled-over virus in a human host—both of which have been shown to correlate

inversely with phylogenetic distance across mammals (4, 8, 9). Indeed, in keeping with trade-off theory,
onward transmission of viruses following spillover is more commonly associated with low virulence
infections (6, 8, 65, 66), suggesting that reservoir orders highlighted here as potential sources for high
virulence pathogens are not necessarily the same orders likely to source future pandemics. Nonetheless,
the possibility for virus adaptation to gain transmission advantages in human hosts following spillover—as
witnessed for Ebola (67) and SARS-CoV-2 (68)—should not be ignored.
448
Currently, our work emphasizes the uniqueness of bats as flying mammals and, in consequence, as long-
lived, tolerant reservoirs for highly virulent viral zoonoses. For the first time, we offer a mechanism for the
evolution of bat-derived viruses that demonstrate significant pathology upon spillover to non-bat,
particularly human, hosts. In providing a theoretical framework to explain this phenomenon, we generate
a series of testable questions and hypotheses for future comparative immunological studies, to be carried
out at *in vitro* and *in vivo* scales. Empirical work should aim to measure rates of immune cell activation,
growth, and mortality across diverse mammalian orders and determine whether natural virus growth rates
are truly higher when evolved in bat hosts. Additional studies should test whether anti-inflammatory
mechanisms in bat cells of different tissues are equally effective at mitigating virus-induced pathology and
immunopathology and whether comparative taxonomic predictions of virus tolerance, resistance, and
virulence evolution apply to non-viral pathogens, too. In light of the emergence of SARS-CoV-2, the field
of bat molecular biology has echoed the call for mechanistic understanding of bat immunology (69), and
the NIH has responded (70), soliciting research on the development of tools needed to test the
predictions outlined here. We offer a bottom-up mechanistic framework enabling the prediction of
emerging virus virulence from the basic immunological and life-history traits of zoonotic hosts.

## Methods

***Evolutionary framework.***

*Population-level dynamics.*

To evaluate the selective pressures that drive the evolution of cross-species virus virulence, we first sought to derive an equation for an optimal virus growth rate expressed in within-host parameters specific to the life history of the reservoir host. To this end, and in keeping with classic examples of viral maintenance in reservoir populations (71–77), we first modeled the dynamics of a persistent infection ($I_1$) in a hypothetical reservoir host, allowing for the introduction of a rare mutant virus strain which generates distinct infections in that same host ($I_2$) (see *SI Appendix* for more detailed methodology and derivations):

$$\frac{dS}{dt} = N(b - qN) - \beta_{r1}SI_1 - \beta_{r2}SI_2 - \mu S \qquad [1a]$$

$$\frac{dI_1}{dt} = \beta_{r1}SI_1 - \mu I_1 - \alpha_{r1}I_1 \qquad [1b]$$

$$\frac{dI_2}{dt} = \beta_{r2}SI_2 - \mu I_2 - \alpha_{r2}I_2 \qquad [1c]$$

Here, $N$ corresponds to the total host population and $S$ to all hosts susceptible to infection, such that $N = S + I_1 + I_2$. We assumed that hosts are born at rate $b$ and die of natural death at rate $\mu$, where $b > \mu$. We further assumed that all hosts are born susceptible and that population density is regulated via a crowding term ($q$). Because we aimed to model the evolution of rates that link to within-host dynamics for the reservoir host, we represented transmission and virulence as functions of the virus causing the infection (respectively, $\beta_r$ and $\alpha_r$) where $r$ denotes the intrinsic virus growth rate and is represented distinctly for both the endemic ($r_1$) and mutant ($r_2$) virus strains.

With the use of model [1], we performed an evolutionary invasion analysis (see *SI Appendix)* and concluded that the virus should evolve to maximize the ratio of transmission over infection duration. We refer to this ratio as the invasion fitness:

$$\frac{\beta_r}{\mu + \alpha_r} \qquad [2]$$

*Within-host dynamics.*

Next, to evaluate the within-host selective conditions underpinning the evolution of the virus growth rate ($r$), we sought to express the between-host parameters of transmission ($\beta_r$) and virulence ($\alpha_r$) in within-host terms, using a nested modeling approach. To this end, we established a simple within-host model representing the dynamics of infection *within* each $I_1$ and $I_2$ host as outlined above. We adapted a class of Lotka-Volterra predator-prey-like within-host models (reviewed in Alizon and van Baalen 2008 (78)), to overcome some constraints of basic predator-prey models of the immune system, chiefly *(i)* by allowing leukocytes to circulate in the absence of infection and *(ii)* by scaling leukocyte growth with virion density, independent of direct leukocyte-virion contact (79–81). This resulted in the following model, which demonstrates interactions between the virus population (V) and the leukocyte population (L) within each infected host:

$$\frac{dV}{dt} = rV - cVL \qquad [3a]$$

$$\frac{dL}{dt} = g_0 + grV - mL \qquad [3b]$$

Here, $r$ corresponds to the intrinsic virus growth rate, $c$ corresponds to the attack efficacy of the immune system upon contact with the virus, and $g$ signifies the recruitment rate of immune cells scaled to the virus growth rate. The parameter, $g_0$, describes the constitutive, baseline leukocyte recruitment in the absence of infection, and $m$ gives the natural leukocyte death rate.

Building from above, we expressed rates of population-level transmission and virulence known to depend on within-host dynamics ($\beta_{r1/r2}$ and $\alpha_{r1/r2}$) in terms of their within-host components, assuming that within-host dynamics are fast relative to host population-level dynamics and, therefore, converge to the endemic equilibrium (i.e. $mr > cg_0$).

549     In line with previous work, we assumed transmission to be a linear function of viral load (79), which we
550     represented as:
551
552     $\beta_r = \zeta V^*$.                                                                                           [4]
553
554     where $\zeta$ corresponds to a scaling term equating viral load to transmission. We additionally assumed that:
555
556     $\alpha_r = \frac{vrV^*}{T_v} + \frac{wgrV^*}{T_w}$,                                                             [5]
557
by which infection-induced host mortality ('virulence') is modeled to result from both virus-induced
pathology (a function of the intrinsic virulence of the parasite, $v$, and the parasite growth rate $rV^*$) and
immunopathology—which we modeled as proportional to the leukocyte growth rate in response to virus
expansion, multiplied by the host's susceptibility to immunopathology and inflammation ($w$). The terms $T_v$
and $T_w$ correspond to host tolerance of virus-induced pathology and immunopathology, respectively,
under assumptions of "constant" tolerance, by which both virus pathology and immunopathology are
reduced by a constant proportion across the course of infection. For constant tolerance, we assumed that
$T_v > 1$ and $T_w > 1$.  See *SI Appendix* for detailed derivations under assumptions of "complete" tolerance,
whereby virus pathology and immunopathology are completely eliminated up to a threshold value, beyond
which pathology scales proportionally with virus and immune cell growth.
568
569     Now, we rewrote the equations for transmission (*[4]*) and virulence (*[5]*) in purely within-host terms:
570
571     $\beta_r = \frac{\zeta(mr - cg_0)}{cgr}$                                                                         [6]
572     $\alpha_r = \frac{r(mr - cg_0)}{cgT_v} + \frac{w(mr - cg_0)}{cT_w}$                                             [7]
573
574     Using these within-host expressions, we then recomputed the expression for invasion fitness (*[2]*) in
575     within host terms and determined the optimal intrinsic virus growth rate, which is an evolutionarily stable
576     strategy (ESS) (see *SI Appendix)*:
577
578     $r^* = \frac{cg_0}{m} + \frac{\sqrt{c^2 m^2 g g_0 \mu T_v T_w (vT_w + gwT_v)}}{m^2(vT_w + gwT_v)}$            [8]
579
580     In Fig. 2, we explore the sensitivity of $r^*$, and the corresponding reservoir host population-level
581     transmission ($\beta^*$) and virulence ($\alpha^*$) at that growth rate, across varied values for its within-host component
582     parameters.
583
*Cross-species dynamics.*
We next sought to derive an expression to explore the consequences of spillover of a virus evolved to its
optimal growth rate in a reservoir host population – which we termed $r_R^*$ – following cross-species
emergence into the human population. Contrary to its established persistent infection in the reservoir
host, we assumed that such a virus would produce an acute infection in the spillover host. To model the
dynamics of this acute spillover, we borrowed from Gilchrist and Sasaki 2002 (82), who developed a
within-host parasite-leukocyte model in which an immortal leukocyte successfully eradicates the parasite
population to near-zero. We modified their acute model to be more comparable to our chronic infection
model (*[3a]/[3b])* and to reflect our differing notation for within-host dynamics.
593
594     $\frac{dV_S}{d\tau} = r_R^* V_S - c_S V_S L_S$                                                                   [9a]
595     $\frac{dL_S}{d\tau} = r_R^* g_S V_S$                                                                             [9b]
596
Here, $r_R^*$ represents the reservoir-evolved virus growth rate, but all other terms reflect within-host
conditions of the spillover host: $V_S$ and $L_S$ correspond, respectively, to the spillover host virus and
leukocyte populations, $c_S$ is the virus consumption rate upon contact with leukocytes in the spillover host,

and $g_S$ is the growth rate of the spillover host leukocyte population in response to virus. We expressed the
model in units of $\tau$, which we assume to be short in comparison to the $t$ time units of dynamics in the
reservoir host population. Rather than solving for virus and leukocyte populations at equilibrium (as in the
reservoir host population), we instead derived an expression for the virus population at the peak of
infection ($V_{S_{max}}$), where we anticipated maximum pathology for the spillover host.

To achieve this, we divided equation *[9a]* by equation *[9b]* to derive a simple, time-independent solution
demonstrating the relationship between virus and leukocyte density:

$$\frac{dV_S}{dL_S} = \frac{1}{g_S}\left(1 - \frac{c_S L_S}{r_R^*}\right)$$  *[10]*

Integration of *[10]* established the following relationship between virus and leukocyte:

$$V_S = \frac{1}{g_S}L_S - \frac{c_S}{2r_R^* g_S}(L_S)^2 + 1 - \frac{1}{g} + \frac{c_S}{2r_S^* g_S}$$  *[11]*

We let $L_S(0) = 1, V_S(0) = 1$, since both virus and leukocyte populations were assumed to be small at $\tau =$
0. Using equation *[11]*, we were then able to derive an expression for $V_{S_{max}}$, the maximum viral load,
which should precede the end of acute infection and the point of host recovery at the maximum age of
infection:

$$V_{S_{max}} = \frac{r_R^*}{g_S c_S} - \frac{r_R^*}{2g_S c_S} + 1 - \frac{1}{g_S} + \frac{c_S}{2r_R^* g_S}$$  *[12]*

Now that we had an established expression for $V_{S_{max}}$, we adapted equation *[7]* to reflect the acute within-
host dynamics of "spillover virulence", which we modeled (as before) as a combination of mortality
induced from direct virus pathology and from immunopathology. The virulence of the virus in the spillover
host will vary with time, $\tau$, tracking the within-host viral load in that same host ($V_S(\tau)$) but should be
highest at peak viral load ($V_{S_{max}}$). As in the reservoir host system, we also modeled the mitigating impact
of tolerance on the two mechanisms of virulence:

$$\alpha_S = V_{S_{max}}\left(\frac{r_R^* v}{T_{vS}(\eta_R)} + \frac{g_S w_S r_R^*}{T_{wS}}\right)$$  *[13]*

This yielded the above expression for 'spillover virulence'. Here, the growth rate of the virus is expressed
at its evolutionary optimum evolved in the reservoir host ($r_R^*$), and the virus's intrinsic virulence ($v$) is
unchanged from one host to the next. Inspired by findings in the literature that report higher virulence in
cross-species infections between hosts separated by larger phylogenetic distances (8–12), we modeled
spillover host tolerance of virus-induced pathology ($T_{vS}$) as a decreasing function of increasing
phylogenetic distance between the reservoir and secondary host, here given as $\eta_R$. All other immune-
related parameters took on characteristics of the spillover host: $g_S$ is the spillover host's leukocyte growth
rate, $w_S$ is the spillover host's susceptibility to immunopathology and inflammation, and $T_{wS}$ corresponds
to the spillover host's tolerance of immunopathology.

***Order-specific estimates for optimal virus growth rates evolved in reservoir hosts.***
Next, we sought to develop estimates for the optimal virus growth rate ($r_R^*$) expected to evolve across a
suite of diverse mammalian reservoir orders. Because within-host immunological data needed to quantify
within-host parameters underpinning the expression for $r_R^*$ are largely lacking, we chose to proxy a few
key within-host parameters from well-described allometric relationships for mammalian life history data.
As such, we here focus on three key parameters for which data could be gleaned from the literature: the
reservoir host mortality rate ($\mu$), the reservoir host tolerance of immunopathology ($T_w$), and the magnitude
of reservoir host constitutive immunity ($g_0$).

To generate order-level summary terms for $\mu$ and $T_w$, we fit a generalized additive model (44) using the
'mgcv' package in R to the relationship between $\log_{10}$ adult body mass (in g) and $\log_{10}$ maximum life span

(in years). We fit our GAM to data from Jones et al. 2009 (56) and Healy et al. 2014 (57), incorporating a
random effect of mammalian order (*SI Appendix,* Table S2). Mass and lifespan data (Fig. 3A) spanned 26
mammalian orders and 1055 individual species. In lieu of phylogeny, we incorporated a random effect of
mammalian host order in all GAMs, which enabled us to produce order-level summary terms for each
parameter. To estimate $\mu$, we predicted maximum lifespan by mammalian order from the fitted GAM using
the predict.gam function in 'mgcv', while excluding the effects of adult body mass, then took the inverse of
that prediction to obtain an estimate of order-level annual mortality (*SI Appendix,* Table S1). To estimate
$T_w$, we extracted the relative partial effects of mammalian order on maximum lifespan, then rescaled
these effects between 1 and 2 for assumptions of constant tolerance and between 0 and 1 for
assumptions of complete tolerance. We justify this approach based on literature that highlights links
between anti-aging molecular pathways that promote longevity and those that mitigate immunopathology
(11, 42, 50–53). Thus, order-level estimates for $\mu$ and $T_w$ are relative inverses of one another, as
highlighted in *SI Appendix*, Fig. S4 (Table S1).

665
To generate order-level summary terms for the magnitude of constitutive immunity ($g_0$), we fit a GAM to
the relationship between the predictor variables of $\log_{10}$ mass (in g) and basal metabolic rate (BMR, in W)
and the response variable of $\log_{10}$ baseline neutrophil concentration (in $10^9$ cells/L), which offers an
approximation of a mammal's constitutive innate immune response. BMR data for this model were
derived from Jones et al. 2009 (56) and Healy et al. 2014 (57), while neutrophil concentrations were
obtained from zoo animal data presented in the Species360 database (49); prior work using this database
has demonstrated scaling relationships between body size and immune cells across mammals (48).
Paired neutrophil and BMR data were limited to just 19 mammalian orders and 144 species. Our GAM
included a random effect of host order (in lieu of 'phylogeny', the significant predictor used in the original
authors' models) and revealed a statistically significant effect of host order modulating the relationship
between body mass, BMR, and neutrophil concentration. Using a similar approach as that employed
above for $T_w$, we estimated $g_0$ by extracting the relative partial effects of mammalian order on neutrophil
concentration, then rescaled these effects between 0 and 1. Because we estimated significantly positive
partial effects between the orders Chiroptera and Monotremata and baseline neutrophil concentration
(Fig. S3), this generated correspondingly high estimates of $g_0$ for these two orders (Fig. S4, Table S3).
681
682    Using these order-level summary terms for $\mu$, $T_w$, and $g_0$, we then generated an order-level prediction for
683    $r_R^*$ across all mammalian host orders, following equation *[8]* (*SI Appendix,* Fig. S5). All other parameters
684    ($c$, $g$, $m$, $w$, $v$, and $T_v$) were held constant across all taxa at values listed in Table 1.
685
***Estimating zoonotic virus virulence in spillover human hosts.***
Once we had constructed an order-level prediction for $r_R^*$, we finally sought to explore the effects of these
reservoir-evolved viruses upon spillover to humans, following equation *[13]*. As when computing virulence
for the reservoir host population, we held immunological parameters, $c$, $g$, $m$, and $w$, constant in humans
(at the same values listed above) due to a lack of informative data to the contrary. Then, to generate an
order-level estimate for virus virulence incurred on humans ($\alpha_S$), we combined our order-level predictions
for $r_R^*$ with order-specific values for $T_{vS}$, the human tolerance of an animal-derived virus, which we scaled
such that viruses derived from more closely-related orders to Primates were more easily tolerated in
humans. Specifically, we represented $T_{vS}$ as the scaled inverse of $\eta_R$, the cophenetic phylogenetic
distance of each mammalian order from Primates. No summarizing was needed for $\eta_R$ because, following
Mollentze and Streicker 2020 (10), we used a composite time-scaled reservoir phylogeny derived from
the TimeTree database (83), which produced a single mean divergence date for all clades. Thus, all
species within a given mammalian order were assigned identical times to most recent common ancestor
with the order Primates. To convert $\eta_R$ to reasonable values for $T_{vS}$, we divided all order-level values of $\eta_R$
by the largest (to generate a fraction), then subtracted that fraction from 2 for assumptions of constant
tolerance (yielding $T_{vS}$ estimates ranging from 1-2) and from 1 for assumptions of complete tolerance
(yielding $T_{vS}$ estimates ranging from 0-1).
703
704    We then combined reservoir host order-level predictions for $r_R^*$ with these estimates for human tolerance
705    of virus pathology ($T_{vS}$) to generate predictions of the expected spillover virulence of viral zoonoses
706    emerging from the 19 mammalian orders for which we possessed complete data across the four variable

707      parameters $(\mu,\ T_w,\ g_0,\ \text{and}\ T_{vS})$. We scaled these $\alpha_S$ estimates—and their 95% predicted confidence
708      intervals—in relative terms (from 0 to 1) to compare with estimates from the literature.
709
***Comparing predictions of spillover virulence by reservoir order with estimates from the literature.***
To compare estimates of spillover virulence generated from our nested modeling approach ($\alpha_S$) with
estimates from the literature, we next followed Day 2002 (84) to convert case fatality rates of viral
zoonoses in spillover human hosts ($CFR_S$) reported in the literature (8) to empirical estimates of $\alpha_S$ for
each mammalian order, using data on the duration of human infection ($D_{IS}$) for each viral zoonosis. For
this purpose, we collected $D_{IS}$ ourselves by searching the primary literature; raw data for infection
durations and associated references are reported in our publicly available GitHub repository. Briefly, Day
2002 (13) notes that the equation for case fatality rate in the spillover host ($CFR_S$) takes the form:
718
719      $CFR_S = \frac{\alpha_S}{\alpha_S + \sigma_S}$                                                        [14]
720
721      where $\sigma_S$ corresponds to the recovery rate from the spilled-over-virus in the human host. We also note
722      that the total duration of infection, $D_{IS}$, is given by:
723
724      $D_{IS} = \frac{1}{\alpha_S + \sigma_S}$                                                        [15]
725
726      Therefore, we translated human case fatality rates of viral zoonoses ($CFR_S$) into estimates of spillover
727      virulence ($\alpha_S$) using the following equation:
728
729      $\alpha_S = \frac{CFR_S}{D_{IS}}$                                                         [16]
730
To obtain a composite order-level prediction for $\alpha_S$ from $D_{IS}$ and $CFR_S$ as reported in (8), we adopted the
best-fit GAM used by the authors in (8) to summarize $CFR_S$ by order and, here, applied it to $\alpha_S$ estimates
converted from $CFR_S$ reported in the literature (*SI Appendix,* Table S2). We then summarized $\alpha_S$ at the
order-level from the fitted GAM, excluding the effects of viral family. This yielded a literature-derived value
for $\alpha_S$ across disparate mammalian orders, against which to compare predictions from our nested
modeling approach. Because the majority of the within-host parameters underpinning $\alpha_S$ predictions in
our nested modeling approach ($\mu,\ T_w,\ g_0,\ \text{and}\ T_{vS}$) were quantified only in a relative fashion, we were
unable to compare direct magnitudes of virulence (e.g. in terms of host death per unit time). To account
for this, we rescaled $\alpha_S$ estimates from both our nested modeling approach and from the empirical
literature from 0 to 1 and compared the relative rank of virulence by mammalian order instead. We
predicted $\alpha_S$ for 19 discrete mammalian orders, though data from the literature were available for only 8
orders against which to compare.
743
Finally, because bats are the most likely ancestral hosts of all lyssaviruses (54), and it may be more
appropriate to class carnivores as 'bridge hosts' for rabies, rather than reservoirs (as the authors discuss
in Guth et al. 2022 (8)), we recomputed literature-derived estimates of relative $\alpha_S$ by fitting the same
GAMs (*SI Appendix,* Table S2) to a version of the dataset excluding all entries for rabies lyssavirus. We
rescaled these new estimates of spillover virulence between 0 and 1 and compared them again to
predictions from our nested modeling approach in *SI Appendix,* Fig. S6-S7.

**Data availability:**
All raw data and corresponding code for analysis and modeling are publicly available at the following GitHub repository: github.com/brooklabteam/spillover-virulence

**Acknowledgments**

The authors thank the Boots Lab at UC Berkeley for helpful comments on this manuscript and acknowledge funding from a National Institutes of Health grant (1R01AI129822-01) to CEB, an Adolph C. and Mary Sprague Miller Institute for Basic Research fellowship to CEB, a Branco Weiss Science in Society fellowship to CEB, a AAAS/Loréal-USA for Women in Science fellowship to CEB, and a Defense Advanced Research Projects Agency PREEMPT Program subgrant (D18AC00031) to CEB.

**References**

1. C. Bonneaud, B. Longdon, Emerging pathogen evolution. *EMBO Reports* **21**, e51374 (2020).
2. J.-B. André, M. E. Hochberg, Virulence evolution in emerging infectious diseases. *Evolution* **59**, 1406 (2005).
3. B. M. Bolker, A. Nanda, D. Shah, Transient virulence of emerging pathogens. *Journal of the Royal Society Interface* **7**, 811–822 (2010).
4. S. Guth, E. Visher, M. Boots, C. E. Brook, Host phylogenetic distance drives trends in virus virulence and transmissibility across the animal – human interface. *Phil. Trans. R. Soc B* **374**, 20190296 (2019).
5. M. J. Farrell, T. J. Davies, Disease mortality in domesticated animals is predicted by host evolutionary relationships. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 7911–7915 (2019).
6. N. Mollentze, D. G. Streicker, P. R. Murcia, K. Hampson, R. Biek, Virulence mismatches in index hosts shape the outcomes of cross-species transmission. *Proceedings of the National Academy of Sciences of the United States of America* (2020) https:/doi.org/10.1073/pnas.2006778117.
7. B. Longdon, *et al.*, The causes and consequences of changes in virulence following pathogen host shifts. *PLoS Pathog* **11**, e1004728 (2015).
8. S. Guth, *et al.*, Bats host the most virulent—but not the most dangerous—zoonotic viruses. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2113628119 (2022).
9. K. J. Olival, *et al.*, Host and viral traits predict zoonotic spillover from mammals. *Nature* (2017) https:/doi.org/10.1038/nature22975.
10. N. Mollentze, D. G. Streicker, Viral zoonotic risk is homogenous among taxonomic orders of mammalian and avian reservoir hosts. *Proceedings of the National Academy of Sciences*, 1–8 (2020).
11. A. T. Irving, M. Ahn, G. Goh, D. E. Anderson, L.-F. Wang, Lessons from the host defences of bats, a unique viral reservoir. *Nature* **589**, 363–370 (2021).
12. T. Schountz, M. L. Baker, J. Butler, V. Munster, Immunological control of viral infections in bats and the emergence of viruses highly pathogenic to humans. *Frontiers in Immunology* **8**, 1098 (2017).
13. M. Boots, Fight or learn to live with the consequences. *Trends in Ecology and Evolution* **23**, 245–248 (2008).
14. L. Råberg, D. Sim, A. F. Read, Disentangling genetic variation for resistance and tolerance to infectious diseases in animals. *Science* **318**, 812–814 (2007).
15. L. Råberg, A. L. Graham, A. F. Read, Decomposing health: tolerance and resistance to parasites in animals. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **364**, 37–49 (2009).
16. S. Mougari, C. Gonzalez, O. Reynard, B. Horvat, Fruit bats as natural reservoir of highly pathogenic henipaviruses: balance between antiviral defense and viral tolerance. *Current Opinion in Virology* **54**, 101228 (2022).
17. M. Ng, *et al.*, NPC1 contributes to species-specific patterns of Ebola virus infection in bats. *eLife* **4**, e11785 (2015).
18. T. N. Starr, *et al.*, ACE2 binding is an ancestral and evolvable trait of sarbecoviruses. *Nature* **603**, 913–918 (2022).
19. M. Letko, *et al.*, Adaptive evolution of MERS-CoV to species variation in DPP4. *Cell Reports* **24**, 1730–1737 (2018).
20. M. Letko, A. Marzi, V. Munster, Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nature Microbiology* **5**, 562–569 (2020).
21. P. Zhou, *et al.*, Contraction of the type I IFN locus and unusual constitutive expression of IFN-α in bats. *Proceedings of the National Academy of Sciences* **113**, 2696–2701 (2016).
22. E. D. Laing, *et al.*, Enhanced autophagy contributes to reduced viral infection in black flying fox cells. *Viruses* **11** (2019).
23. Y. T. Chionh, *et al.*, High basal heat-shock protein expression in bats confers resistance to cellular heat/oxidative stress. *Cell Stress and Chaperones* **24**, 835–849 (2019).
24. T. J. O'Shea, *et al.*, Bat flight and zoonotic viruses. *Emerging Infectious Diseases* **20**, 741–745 (2014).

25.   B. Y. S. P. Thomas, R. A. Suthers, The physiology and energetics of bat flight. *Journal of*
*Experimental Biology* **57**, 317–335 (1972).

26.   J. Speakman, D. Thomas, "Physiological ecology and energetics of bats" in *Bat Ecology*, T. Kunz,
M. Fenton, Eds. (University of Chicago Press, 2003), pp. 430–490.

27.   G. Zhang, *et al.*, Comparative analysis of bat genomes provides insight into the evolution of flight
and immunity. *Science* **339**, 456–60 (2013).

28.   D. M. Santillan, *et al.*, Large-scale genome sampling reveals unique immunity and metabolic
adaptations in bats. *Molecular Ecology* (2021) https:/doi.org/10.1111/mec.16027.

29.   M. Ahn, J. Cui, A. T. Irving, L.-F. Wang, Unique loss of the PYHIN gene family in bats amongst
mammals: Implications for inflammasome sensing. *Scientific Reports* **6**, 21722 (2016).

30.   M. Ahn, *et al.*, Dampened NLRP3-mediated inflammation in bats and implications for a special viral
reservoir host. *Nature Microbiology* **4** (2019).

31.   J. Xie, *et al.*, Dampened STING-dependent interferon activation in bats. *Cell Host and Microbe* **23**,
1–5 (2018).

32.   G. Goh, *et al.*, Complementary regulation of caspase-1 and IL-1 β reveals additional mechanisms of
dampened inflammation in bats. *Proceedings of the National Academy of Sciences* (2020)
https:/doi.org/10.1073/pnas.2003352117.

33.   S. N. Austad, K. E. Fischer, Mammalian aging, metabolism, and ecology: evidence from the bats
and marsupials. *Journal of gerontology* **46**, B47-53 (1991).

34.   C. E. Brook, *et al.*, Accelerated viral dynamics in bat cell lines, with implications for zoonotic
emergence. *eLife* **9**, e48401 (2020).

35.   D. Ebert, J. J. Bull, Challenging the trade-off model for the evolution of virulence: is virulence
management feasible? *Trends in Microbiology* **11**, 15–20 (2003).

36.   S. Alizon, A. Hurford, N. Mideo, M. Van Baalen, Virulence evolution and the trade-off hypothesis:
History, current state of affairs and the future. *Journal of Evolutionary Biology* **22**, 245–259 (2009).

37.   R. M. Anderson, R. M. May, Coevolution of hosts and parasites. *Parasitology* **85**, 411–426 (1982).

38.   M. R. Miller, A. White, M. Boots, The evolution of parasites in response to tolerance in their hosts:
The good, the bad, and apparent commensalism. *Evolution* **60**, 945 (2006).

39.   A. Best, A. White, M. Boots, The coevolutionary implications of host tolerance. *Evolution* **68**, 1426–
1435 (2014).

40.   H. E. Randolph, L. B. Barreiro, Holy Immune Tolerance, Batman! *Immunity* **48**, 1074–1076 (2018).

41.   D. T. S. Hayman, Bat tolerance to viral infections. *Nature Microbiology* **4**, 728–729 (2019).

42.   S. S. Pavlovich, *et al.*, The Egyptian Rousette genome reveals unexpected features of bat antiviral
immunity. *Cell* **173**, 1–13 (2018).

43.   S. A. H. Geritz, É. Kisdi, G. Meszéna, J. A. J. Metz, Evolutionarily singular strategies and the
adaptive growth and branching of the evolutionary tree. *Evolutionary Ecology* **12**, 35–57 (1998).

44.   S. N. Wood, mgcv: GAMs and Generalized Ridge Regression for R. *R News* **1/2**, 20–24 (2001).

45.   Brown, James, West, Geoffrey, Enquist, Brian, "Scaling in biology: Patterns and processes, causes
and consequences" in *Scaling in Biology*, (Oxford University Press, 2000), pp. 87–112.

46.   J. H. Brown, J. F. Gillooly, A. P. Allen, V. M. Savage, G. B. West, Towards a metabolic theory of
ecology. *Ecology* **85**, 1771–1789 (2004).

47.   G. B. West, W. H. Woodruff, J. H. Brown, Allometric scaling of metabolic rate from molecules and
mitochondria to cells and mammals. *Proceedings of the National Academy of Sciences of the*
*United States of America* **99**, 2473–2478 (2002).

48.   C. J. Downs, N. A. Dochtermann, R. Ball, K. C. Klasing, L. B. Martin, The effects of body mass on
immune cell concentrations of mammals. *American Naturalist* **195**, 107–114 (2020).

49.   Species360 Zoological Information Management System (ZIMS) (August 25, 2022).

50.   G. S. Wilkinson, *et al.*, DNA methylation predicts age and provides insight into exceptional longevity
of bats. *Nature Communications* **12** (2021).

51.   J. Dhinaut, A. Balourdet, M. Teixeira, M. Chogne, Y. Moret, A dietary carotenoid reduces
immunopathology and enhances longevity through an immune depressive effect in an insect model.
*Scientific Reports* **7** (2017).

52.   C. Caruso, D. Lio, L. Cavallone, Aging, longevity, inflammation, and cancer. *Annals of New York*
*Academiy of Sciences* **1028**, 1–13 (2004).

53.   D. K. Fabian, *et al.*, Evolution of longevity improves immunity in *Drosophila*. *Evolution Letters* **2**,
567–579 (2018).

54.    C. E. Rupprecht, A. Turmelle, I. V. Kuzmin, A perspective on lyssavirus emergence and
928           perpetuation. *Current Opinion in Virology* **1**, 662–670 (2016).
929    55.    C. E. Brook, A. P. Dobson, Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends in*
930           *Microbiology* **23**, 172–80 (2015).
931    56.    K. E. Jones, *et al.*, PanTHERIA: a species-level database of life history, ecology, and geography of
932           extant and recently extinct mammals. *Ecology* **90**, 2648–2648 (2009).
933    57.    K. Healy, *et al.*, Ecology and mode-of-life explain lifespan variation in birds and mammals.
934           *Proceedings of the Royal Society B* (2014).
935    58.    R. Hamilton, M. Siva-Jothy, M. Boots, Two arms are better than one: Parasite variation leads to
936           combined inducible and constitutive innate immune responses. *Proceedings of the Royal Society B:*
937           *Biological Sciences* **275**, 937–945 (2008).
938    59.    A. E. Shaw, *et al.*, Fundamental properties of the mammalian innate immune system revealed by
939           multispecies comparison of type I interferon responses. *PLoS Biol* **15**, e2004086 (2017).
940    60.    E. Clayton, M. Munir, Fundamental characteristics of bat interferon systems. *Front. Cell. Infect.*
941           *Microbiol.* **10**, 527921 (2020).
942    61.    W. Widagdo, *et al.*, Tissue distribution of the MERS-coronavirus receptor in bats. *Sci Rep* **7**, 1193
943           (2017).
944    62.    M. Ruiz-Aravena, *et al.*, Ecology, evolution and spillover of coronaviruses from bats. *Nature*
945           *Reviews Microbiology* (2021) https:/doi.org/10.1038/s41579-021-00652-2.
946    63.    N. Simmons, A. Cirranello, Bat species of the world: A taxonomic and geographic database (2020)
947           (July 11, 2020).
948    64.    G. S. Wilkinson, D. M. Adams, Recurrent evolution of extreme longevity in bats. *Biology Letters* **15**,
949           20180860 (2019).
950    65.    J. L. Geoghegan, A. M. Senior, F. Di, E. C. Holmes, Virological factors that increase the
951           transmissibility of emerging human viruses. *Proceedings of the National Academy of Sciences* **113**,
952           4170–4175 (2016).
953    66.    B. Longdon, J. D. Hadfield, C. L. Webster, D. J. Obbard, F. M. Jiggins, Host phylogeny determines
954           viral persistence and replication in novel hosts. *PLoS Pathogens* **7** (2011).
955    67.    R. A. Urbanowicz, *et al.*, Human adaptation of Ebola virus during the West African Outbreak. *Cell*
956           **167**, 1079–1087 (2016).
957    68.    J. A. Plante, *et al.*, Spike mutation D614G alters SARS-CoV-2 fitness. *Nature* (2020)
958           https:/doi.org/10.1038/s41586-020-2895-3.
959    69.    L.-F. Wang, A. M. Gamage, W. O. Y. Chan, Decoding bat immunity: the need for a coordinated
960           research approach. *Nature Reviews Immunology* **0123456789**, 0–2 (2021).
961    70.    National Institutes of Health, Research on Bat Immunology.
962    71.    M. Viana, *et al.*, Assembling evidence for identifying reservoirs of infection. *Trends in Ecology and*
963           *Evolution* **29**, 270–9 (2014).
964    72.    K. Hampson, D. T. Haydon, Persistent pathogens and wildlife reservoirs. *Science* **374**, 35–36
965           (2021).
966    73.    D. T. Haydon, S. Cleaveland, L. H. Taylor, M. K. Laurenson, Identifying reservoirs of infection: A
967           conceptual and practical challenge. *Emerging infectious diseases* **8**, 1468–73 (2002).
968    74.    A. Jolles, *et al.*, Endemic persistence of a highly contagious pathogen: Foot-and-mouth disease in
969           its wildlife host. *Science* **374**, 104–109 (2021).
970    75.    R. K. Plowright, *et al.*, Transmission or within-host dynamics driving pulses of zoonotic viruses in
971           reservoir-host populations. *PLoS Neglected Tropical Diseases* **10**, e0004796 (2016).
972    76.    T. Schountz, *et al.*, Regulatory T cell-like responses in deer mice persistently infected with Sin
973           Nombre virus. *Proceedings of the National Academy of Sciences* **104**, 15496–15501 (2007).
974    77.    T. Schountz, J. Prescott, Hantavirus immunology of rodent reservoirs: Current status and future
975           directions. *Viruses* **6**, 1317–1335 (2014).
976    78.    S. Alizon, M. van Baalen, Acute or Chronic? Within-host models with immune dynamics, infection
977           outcome, and parasite evolution. *The American Naturalist* **172**, E244–E256 (2008).
978    79.    S. Alizon, M. van Baalen, Emergence of a convex trade-off between transmission and virulence.
979           *The American Naturalist* **165**, E155–E167 (2005).
980    80.    S. Alizon, M. van Baalen, Multiple infections, immune dynamics, and the evolution of virulence. *The*
981           *American Naturalist* **172**, E150–E168 (2008).

981    81.   A. Fenton, J. Lello, M. B. Bonsall, Pathogen responses to host immunity: The impact of time delays and memory on the evolution of virulence. *Proceedings of the Royal Society B: Biological Sciences* **273**, 2083–2090 (2006).

82.   M. A. Gilchrist, A. Sasaki, Modeling host – parasite coevolution: A nested approach based on mechanistic models. *Journal of Theoretical Biology* **218**, 289–308 (2002).

83.   S. Kumar, G. Stecher, M. Suleski, S. B. Hedges, TimeTree: a resource for timelines, timetrees, and divergence times. *Molecular Biology and Evolution* **34**, 1812–1819 (2017).

84.   T. Day, On the evolution of virulence and the relationship between various measures of mortality. *Proceedings of the Royal Society B* **269**, 1317–1323 (2002).

85.   M. Boots, R. Donnelly, A. White, Optimal immune defence in the light of variation in lifespan. *Parasite Immunology* **35**, 331–338 (2013).

**Table 1. Default parameter values for population-level and within-host models.**

| Parameter | Definition | Units | Value* | Scale |
|:---:|:---:|:---:|:---:|:---:|
| $b$ | host natural birth rate | days$^{-1}$ | 0.2 | population-level |
| $q$ | crowding term to regulate density-dependent growth | days$^{-1}$ | 0.0002 | population-level |
| $\mu^{\dagger}$ | host background mortality rate | days$^{-1}$ | $\dfrac{1}{20*365}$ | population-level |
| $g_0^{\dagger}$ | magnitude of constitutive immunity *(rate of baseline neutrophil supply)* | days$^{-1}$ | .3 | within-host |
| $g$ | rate of leukocyte activation by virus | days$^{-1}$ | 0.9 | within-host |
| $c$ | rate of virus consumption by leukocyte | days$^{-1}$ | 0.5 | within-host |
| $m$ | background mortality rate for leukocytes | days$^{-1}$ | $\dfrac{1}{21}$ | within-host |
| $\zeta$ | maximum between-host transmission rate | days$^{-1}$ | 0.2 | within-host |
| $w$ | host susceptibility to immunopathology | leukocytes$^{-1}$ | 1 | within-host |
| $v$ | intrinsic virus virulence | virions$^{-1}$ | 1 | within-host |
| $T_w^{\dagger}$ | tolerance of immunopathology (constant or complete) | unitless | *variable* constant: $1-2$ complete: $0-1$ | within-host |
| $T_{vS}^{\dagger}$ | human tolerance of viral pathology (constant or complete) | unitless | *variable* constant: $1-2$ complete: $0-1$ | between-host spillover |

*Value column gives parameter values fixed in Fig. 2 and *SI Appendix,* Fig. S2, excepting instances when parameter was modulated along the x-axis or internally ($T_w$, $T_v$) as indicated in the figure legend. Parameter values are comparable to the ranges explored in Miller et al. 2006 (38) and Boots et al. 2013 (85).
$^{\dagger}$Parameters $\mu$, $g_0$, $T_w$, and $T_{vS}$ were modeled as functions of mammalian order-specific life history traits in Fig. 3 and in *SI Appendix,* Fig. S6-S7, as outlined in the Methods.

1037