Reservoir host immunology and life history shape virulence evolution in zoonotic viruses

Cara E. Brook[1]*, Carly Rozins[2], Sarah Guth[3], Mike Boots[3,4]

[1] Department of Ecology and Evolution, University of Chicago, Chicago, IL, USA

[2] Department of Science and Technology Studies, York University, Toronto, Canada

[3] Department of Integrative Biology, University of California, Berkeley, Berkeley, CA, USA

[4] Biosciences, University of Exeter, TR10 9FE Penryn, UK

* Corresponding author: Cara E. Brook
Email: cbrook@uchicago.edu (CEB)

Abstract

The management of future pandemic risk requires a better understanding of the mechanisms that determine the virulence of emerging zoonotic viruses. Meta-analyses suggest that the virulence of emerging zoonoses is correlated with but not completely predictable from reservoir host phylogeny, indicating that specific characteristics of reservoir host immunology and life history may drive the evolution of viral traits responsible for cross-species virulence. In particular, bats host viruses that cause higher case fatality rates upon spillover to humans than those derived from any other mammal, a phenomenon that cannot be explained by phylogenetic distance alone. In order to disentangle the fundamental drivers of these patterns, we develop a nested modeling framework that highlights mechanisms which underpin the evolution of viral traits in reservoir hosts that cause virulence following cross-species emergence. We apply this framework to generate virulence predictions for viral zoonoses derived from diverse mammalian reservoirs, successfully recapturing corresponding virus-induced human mortality rates reported in the literature. Notably, our work offers a mechanistic explanation for the extreme virulence of bat-borne zoonoses and, more generally, demonstrates how key differences in reservoir host longevity, viral tolerance, and constitutive immunity impact the evolution of viral traits that cause virulence following spillover to humans. Our theoretical framework offers a series of testable questions and hypotheses designed to stimulate future work comparing cross-species virulence evolution in zoonotic viruses derived from diverse mammalian hosts.

Introduction

The devastating impact of the SARS-CoV-2 pandemic highlights the extreme public health outcomes that can result upon cross-species emergence of zoonotic viruses. Estimating the relative threats posed by potential future zoonoses is an important but challenging public health undertaking. In particular, efforts to predict the virulence of emerging viruses can be complicated since chance will always play a role in dictating the initial spillover that precedes selection (1), virulence upon emergence may be maladaptive in novel hosts (2,3), and patterns in available data may be muddled by attainment bias if avirulent infections go underreported (1). Nonetheless, a growing body of recent work highlights clear associations between reservoir and spillover host phylogeny and the virulence of a corresponding cross-species infection (4–7). In many cases, increasing phylogenetic distance between reservoir and spillover hosts is correlated with higher virulence infections (4–6), suggesting that spillover host immune systems may be poorly equipped to tolerate viral traits optimized in more distantly-related reservoirs. Still, the effect of phylogeny on spillover virulence appears to supersede that of simple phylogenetic distance (7,8), indicating that taxon-specific reservoir host immunological and life history traits may be important drivers of cross-species virus virulence. Notably, bats host viruses that cause higher human case fatality rates than zoonoses derived from other mammals and birds, a phenomenon that cannot be explained by phylogenetic distance alone (8). Understanding the mechanisms that select for the evolution of unique viral traits in bats compared to those selected in other mammalian reservoirs should enable us to better predict the virulence of future zoonotic threats.

Although the disproportionate frequency with which the Chiropteran order may source viral zoonoses remains debated (9,10), the extraordinary human pathology induced by many bat-borne zoonoses—including Ebola and Marburg filoviruses, Hendra and Nipah henipaviruses, and SARS, MERS, and SARS-CoV-2 coronaviruses (11)—is not contested. Remarkably, bats demonstrate limited clinical pathology from infection with viruses that cause extreme morbidity and mortality in other hosts (12). Bats avoid pathological outcomes from viral infection via a combination of unique resistance and tolerance mechanisms, which, respectively, limit the viral load accrued during infection ('resistance') and reduce the disease consequences of a given viral load ('tolerance') (13–16). Viral resistance mechanisms vary across bat species; those described to date include: receptor incompatibilities that limit the extent of infection for certain viruses in certain bats (17–20), constitutive expression of antiviral cytokines in some bat species (21), and enhanced autophagy (22) and heat-shock protein expression (23) in others. Expansion of anti-viral APOBEC3 genes has also been documented in a few well-studied bat genomes (24,25). While such robust antiviral immunity would result in widespread immunopathology in most mammals, bats—as the only mammals capable of powered flight—have evolved numerous unique mechanisms of mitigating inflammation incurred during the intensive physiological process of flying (26–28). These anti-inflammatory adaptations include loss of PYHIN (29–31) and downregulation of NLRP3 (32) inflammasome-forming gene families, loss of pro-inflammatory genes in the NF-κB pathway (24), dampened interferon activation in the STING pathway (33), and diminished caspase-1 inflammatory signaling (34). In addition to facilitating flight, this resilience to inflammation has yielded the apparent by-products of extraordinarily long bat lifespans (35) and tolerance of the immunopathology that typically results from viral infection (11). Moreover, recent work demonstrates how high virus growth rates easily tolerated in constitutively antiviral bat cells cause significant pathology in cells lacking these unique antiviral defenses (36). The extent to which inflammatory tolerance may modulate the evolution of the viruses that bats host, however, remains largely unexplored.

Modern theory on the evolution of virulence typically assumes, either explicitly or implicitly, that high pathogen growth rates should both enhance between-host transmission and elevate infection-induced morbidity or mortality, resulting in a trade-off between virulence and transmission (37–39). Theory further suggests that because viral 'tolerance' mitigates virulence without reducing viral load, most host strategies of tolerance should select for higher growth rate pathogens that achieve gains in between-host transmission without causing damage to the original host (39–41). The widely touted viral tolerance of

bats (11,16,42–44) should therefore be expected to support the evolution of enhanced viral growth rates,
which—though avirulent to bats—may cause significant pathology upon spillover to hosts lacking unique
features of bat immunology and physiology. Beyond tolerance, other life history characteristics unique to
diverse reservoir hosts should also impact the evolution of traits in the viruses they host—with important
consequences for cross-species virulence following spillover. However, to date, we lack a specific theory
that examines the relative impact of reservoir host life history on spillover virulence. Here, we explore the
extent to which the immunological and life-history traits of mammalian reservoirs can explain variation in
the virulence of zoonotic viruses emerging into human hosts.

Results

General trends in the evolution of high growth rate viruses

To elucidate how immunological and life-history traits of mammalian hosts combine to drive zoonotic virus
virulence, we adopted a nested modeling approach (45), embedding a simple within-host model of viral
and leukocyte dynamics within an epidemiological, population-level framework (Fig 1). We examined how
the life history traits of a primary reservoir drive the evolution of viral traits likely to cause pathology in a
secondary, spillover host—chiefly, a human. Using our nested model in an adaptive dynamics framework
(46), we first expressed the conditions—called the 'invasion fitness'—which permit invasion of an
evolutionarily 'fitter', mutant virus into a reservoir host system in within-host terms. From this, we derived
an expression for r_R^*, the optimal within-host growth rate of a persistent virus evolved at equilibrium
prevalence in its reservoir host; our modeling framework allowed us to express r_R^* as a function of within-
host traits that we might expect to vary across mammalian reservoirs with divergent life histories. We
deduced that r_R^* can, consequently, also be expected to vary as a result of these life history differences,
which modulate the optimization process by which a virus maximizes gains in between-host transmission
($\beta_{r_R}^*$) while mitigating virulence incurred on its reservoir ($\alpha_{r_R}^*$; Fig 1). From this framework, we next derived
an expression for α_S, the virulence incurred by a reservoir-optimized virus immediately following spillover
to a novel host. We expressed α_S as a function of the reservoir-optimized virus growth rate (r_R^*), combined
with spillover host tolerance of virus pathology (T_{vS}), which we modeled as proportional to the
phylogenetic distance between reservoir and spillover host (Fig 1).

Fig 1. Conceptual mechanistic framework to predict zoonotic virus virulence from reservoir immunology and life history traits. (A) A within-host predator-prey-like model of leukocyte-virus dynamics is embedded in a population-level transmission model for reservoir hosts. Between-host rates of transmission (β_{r_R}) and virulence (α_{r_R}) are expressed as functions of within-host dynamics to derive optimal virus growth rates in the reservoir (r_R^*). See S1 File and Table 1 for parameter definitions and values. (B) Virus growth rates (r_R^*) evolve to maximize gains in between-host transmission ($\beta_{r_R}^*$) over virulence incurred on the reservoir host ($\alpha_{r_R}^*$) where both transmission and virulence are expressed as functions of r_R. Viruses evolved in taxa with immunological and physiological environments similar to humans (i.e. non-human primates) should optimize at relatively low growth rates (r_R^*) that yield minimal pathology for the reservoir host (α_R^*) and are, correspondingly, likely to cause only minimal pathology in the spillover human host (α_S). In our model, the virulence of a virus in its spillover host (α_S) is expressed as a function of both the reservoir host-evolved virus growth rate (r_R^*) and the spillover host tolerance of virus pathology (T_{vS}), which we model as proportional to phylogenetic distance between reservoir and spillover host. (C) As a result of unique bat virus tolerance, viruses evolved in bat reservoir hosts may optimize at high r_R^* values that maximize bat-to-bat transmission ($\beta_{r_R}^*$) but cause only minimal pathology in the bat host ($\alpha_{r_R}^*$). Such viruses are likely to generate extreme virulence upon spillover (α_S) to secondary hosts, including humans, that lack bat life history traits. The virulence of a virus in its spillover host is amplified (α_S) in cases where large phylogenetic distance between reservoir and spillover host results in minimal spillover host tolerance of virus pathology (T_{vS}).

Our nested modeling approach first followed Alizon and van Baalen (2005) (45) in derivation of an expression for r_R^*, the within-host virus growth rate optimized based on endemic circulation in a reservoir host. The equation for r_R^* can be captured as follows:

$$r_R^* = \frac{c_R g_{0R}}{m_R} + \frac{\sqrt{c_R^2 m_R^2 g_R g_{0R} \mu_R T_{vR} T_{wR}(v T_{wR} + g_R w T_{vR})}}{m_R^2(v T_{wR} + g_R w T_{vR})},$$

where μ_R signifies the natural mortality rate of the reservoir host, and all other parameters represent
within-host viral and immune dynamics in a persistently infected reservoir. Thus, c_R corresponds to the
rate of virus clearance by leukocytes, g_{0R} is the magnitude of constitutive immunity, m_R is the natural
leukocyte mortality rate, and g_R is the rate of leukocyte activation following infection, all in the reservoir
host. The parameters v and w correspond, respectively, to the intrinsic virulence of the virus and its
propensity to elicit a damaging inflammatory response from its host's immune system—while T_{vR} and T_{wR}
respectively represent reservoir host tolerance to direct virus pathology and to immunopathology. From
the above expression, we explored a range of optimal within-host virus growth rates (r_R^*) for viruses
evolved in reservoir hosts with diverse cellular and immunological parameters (Fig 2; Table 1; S1-S2
Figs). We subsequently calculated the corresponding transmission ($\beta_{r_R}^*$) and virulence ($\alpha_{r_R}^*$) incurred by
viruses evolved to r_R^* in their reservoir hosts, then, following Gilchrist and Sasaki (2002) (47), modeled the
nascent spillover of a reservoir-evolved virus as acute infection in a secondary host. In this spillover
infection, we assumed that the virus retains its reservoir-optimized growth rate (r_R^*) while replicating in the
physiological and immunological environment of its novel spillover host. We express this spillover
virulence (α_S) as:

$$\alpha_S = V_{S_{avg}} \left(\frac{r_R^* v}{T_{vS}} + \frac{g_S w r_R^*}{T_{wS}} \right)$$

where $V_{S_{avg}}$ corresponds to the average viral load experienced across the timecourse of an acute spillover
host infection, and g_S, T_{vS}, and T_{wS} represent the spillover host analogues of previously-described within-
host parameters in the reservoir host equations (Fig 2; Table 1; S1-S2 Figs). We present all main text
results under assumptions by which tolerance manifests as a *constant* reduction of either direct virus-
induced pathology (T_v) or immunopathology (T_w). See S1 File for comparable results under assumptions
of *complete* tolerance, whereby tolerance completely eliminates virus pathology and immunopathology up
to a threshold value, beyond which pathology scales proportionally with virus and immune cell growth.

Fig 2. Optimal virus growth rates—and subsequent spillover virulence—vary across reservoir host immunological and life history parameters. Rows (top-down) indicate the evolutionarily optimal within-host virus growth rate (r_R^*) and the corresponding between-host transmission rate ($\beta_{r_R}^*$), and virus-induced mortality rate ($\alpha_{r_R}^*$) for a reservoir host infected with a virus at r_R^*. The bottom row then demonstrates the resulting virulence (α_S) of a reservoir-optimized virus evolved to r_R^* upon nascent spillover to a novel, secondary host. Columns demonstrate the dependency of these outcomes on variable within-host parameters in the reservoir host: background mortality (μ_R), magnitude of constitutive immunity (g_{0R}), rate of leukocyte activation upon viral contact (g_R), rate of virus consumption by leukocytes (c_R), and leukocyte mortality rate (m_R). Darker colored lines depict outcomes at higher reservoir host tolerance of direct virus pathology (T_{vR}, red) or immunopathology (T_{wR}, blue), assuming no tolerance of the opposing type. Heat maps demonstrate how T_{vR} and T_{wR} interact to produce each outcome. Parameter values are reported in Table 1. Figure assumes tolerance in the 'constant' form. See S2 Fig for 'complete' tolerance assumptions and S3 Fig for changes in α_S across a variable range of parameter values for *spillover* host tolerance of direct virus pathology, T_{vS}, and immunopathology, T_{wS}.

Table 1. Default parameter values for population-level and within-host models.

Parameter	Definition	Units	Value*	Scale
b_R	reservoir host natural birth rate	days^{-1}	0.2	population-level
q_R	reservoir host crowding term to regulate density-dependent growth	days^{-1}	0.0002	population-level
$\mu_R^{\dagger}$	reservoir host background mortality rate	days^{-1}	$\dfrac{1}{20*365}$	population-level
$g_{0R}^{\dagger}$	reservoir host magnitude of constitutive immunity *(rate of baseline neutrophil supply)*	days^{-1}	.3	within-host
g_R	rate of leukocyte activation by virus	days^{-1}	0.9	within-host
c_R	rate of virus consumption by leukocyte	days^{-1}	0.5	within-host
m_R	background mortality rate for leukocytes	days^{-1}	$\dfrac{1}{21}$	within-host
ζ	scaling factor translating within-host viral load to between-host transmission rate	days^{-1}	0.2	within-host
w	intrinsic virus propensity to elicit inflammatory immune response from host	leukocytes^{-1}	1	within-host
v	intrinsic virus virulence	virions^{-1}	1	within-host
$T_{wR}^{\dagger}$	reservoir host tolerance of immunopathology (constant or complete)	unitless	*variable* constant: $1-2$ complete: $0-1$	within-host
T_{vR}	reservoir host tolerance of virus pathology (constant or complete)	unitless	*variable* constant: $1-2$ complete: $0-1$	within-host
$T_{vS}^{\dagger}$	human tolerance of viral pathology (constant or complete)	unitless	*variable* constant: $1-2$ complete: $0-1$	between-host spillover

Our analyses highlight several critical drivers of virus evolution likely to generate significant pathology following spillover to a secondary host (Fig 2): higher within-host virus growth rates (r_R^*) are selected in reservoir hosts with higher background mortality (μ_R), elevated constitutive immune responses (g_{0R}), more rapid leukocyte activation upon infection (g_R), and more rapid virus clearance by the host immune system (c_R). Additionally, higher r_R^* viruses are selected in hosts exhibiting lower leukocyte mortality rates (m_R), resulting in longer-lived immune cells. Critically, higher reservoir host tolerance of both virus-induced pathology (T_{vR}) and immunopathology (T_{wR}) also select for higher r_R^*. In keeping with trade-off theory, changes in the majority of within-host parameters drive corresponding increases in r_R^*, β_R^*, and α_R^*, such that viruses evolved to high growth rates experience high transmission within the reservoir host population—but also generate high virus-induced mortality (Fig 2). By definition, the two modeled mechanisms of tolerance (T_{vR} and T_{wR}) decouple the relationship between transmission and virulence, permitting evolution of high r_R^* viruses that achieve gains in between-host transmission ($\beta_{r_R}^*$), while simultaneously incurring minimal virulence ($\alpha_{r_R}^*$) on reservoir hosts. By extension, we demonstrate that viruses evolving high optimal r_R^* values in reservoir hosts incur substantial pathology upon spillover to secondary hosts (α_S). Intriguingly, the virulence that a virus incurs on its spillover host (α_S) accelerates substantially faster than that incurred on its reservoir host (α_R^*) at higher values for certain parameters, chiefly, g_{0R}, c_R as well as, most critically, T_{vR} and T_{wR}. This underscores the important capacity of these within-host traits to drive cross-species virulence in emerging viruses.

Order-specific estimates for optimal virus growth rates evolved in reservoir hosts

We next applied our model to make broad predictions of the evolution of optimal virus growth rates (r_R^*) across diverse mammalian reservoir orders, based on order-specific variation in three key parameters from our nested model: the reservoir host background mortality rate (μ_R), the reservoir host tolerance of immunopathology (T_{wR}), and the reservoir host magnitude of constitutive immunity (g_{0R}). Within-host immunological data needed to quantify these parameters is lacking for most taxa; thus, we used regression analyses to summarize these terms across mammalian orders from publicly available life history data. In particular, we used well-described allometric relationships between mammalian body mass and basal metabolic rate (BMR) with lifespan and immune cell concentrations (49–53)—to proxy μ_R, T_{wR}, and g_{0R} across mammalian orders (Fig 3A-C; Table 1; S4-S5 Figs). From here, we used our nested modeling framework to predict optimal virus growth rates (r_R^*) across diverse mammalian reservoirs (Fig 3D). Then, we estimated T_{vS}, the spillover host tolerance of direct virus pathology, as proportional to the time to most recent common ancestor (MRCA) between the human primate order and each mammalian reservoir order, now focusing our analysis on zoonotic spillover (Fig 3E-F; Table 1). Finally, we combined estimates for r_R^* and T_{vS} to generate a prediction for α_S, the virulence of a reservoir-optimized virus in a

human spillover host, which we can compare against case fatality rates for mammalian zoonoses
reported in the literature (4,8) (Fig 3G; S5-S9 Figs).

Fig 3. Reservoir host life history traits predict evolution of zoonotic virus virulence. (**A**) Variation in $\log_{10}$
maximum lifespan (y-axis, in years) with $\log_{10}$ adult body mass (x-axis, in grams) across mammals, with data derived
from Jones et al. 2009 (54) and Healy et al. 2014 (55). Points are colored by mammalian order, corresponding to
legend. Black line depicts predictions of mammalian lifespan per body mass, summarized from fitted model (but
excluding random effect of mammalian order), presented in S2 Table. (**B**) Baseline neutrophil concentrations (y-axis,
in 10^9 cells/L) per mass-specific metabolic rate (x-axis, in W/g) across mammals, with data from Jones et al. 2009
(54) and Healy et al. 2014 (55) combined with neutrophil concentrations from Species360 (53). Black line projects
neutrophil concentration per mass-specific metabolic rate (excluding random effects of mammalian order), simplified

from fitted model presented in S2 Table. (**C**) Order-level parameters for nested modeling framework were derived
from fitting of linear models and linear mixed models visualized here and presented in S4 Fig and S2 Table to data
from (A) and (B). Average annual mortality rate (μ_R) was predicted from a linear regression of species-level annual
mortality (the inverse of maximum lifespan), as described by a predictor variable of host order; tolerance of
immunopathology (T_{w_R}) was derived from the scaled effect of host order on the linear mixed effects regression of
$\log_{10}$ maximum lifespan (in years) by $\log_{10}$ mass (in grams), incorporating a random effect of order. The magnitude of
constitutive immunity (g_{0_R}) was derived from the scaled effect of order on the regression of $\log_{10}$ neutrophil
concentration per $\log_{10}$ body mass (in grams), combined with BMR (in W) (S4 Fig; S1-S2 Tables). Panels shown here
give numerical estimates for μ_R and order-level effects from fitted models that were scaled to numerical values for
T_{w_R} and g_{0_R}, as presented in S4 Fig (S1 Table). Red and blue colors correspond to, respectively, significantly positive
or negative order-level partial effects from these regressions. (**D**) Reservoir-host estimates for μ_R, T_{wR}, and g_{0R} were
combined in our modeling framework to generate a prediction of optimal growth rate for a virus evolved in a host of
each mammalian order (r_R^*). Here, point size corresponds to the average number of species-level datapoints used to
generate each of the three variable parameters impacting r_R^*, as indicated in legend. (**E**) Phylogenetic distance from
Primates (in millions of years, indicated by color) on a timescaled phylogeny, using data from TimeTree (56). (**F**) An
order-level estimate for the nested model parameter, T_{vS}, the spillover human host tolerance of pathology induced by
a virus evolved in a different reservoir order, was estimated as the scaled inverse of the phylogenetic distance shown
in (E) (S4 Fig; S1 Table). (**G**) Reservoir-host predictions of optimal virus growth rates (r_R^*) from (D) were combined
with human spillover host estimates of tolerance for direct virus pathology (T_{vS}) from (F) in our nested modeling
framework to generate a prediction of the relative spillover virulence (α_S) of a virus evolved in a given reservoir host
order immediately following spillover into a secondary, human host. Here, the left panel visualizes predictions from
our nested modeling framework, using order-specific parameters for μ_R, T_{wR}, g_{0R}, and T_{vS} (S1 Table). The right panel
depicts relative human α_S estimates derived from case fatality rates and infection duration reported in the zoonotic
literature (8). For the left panel, point size corresponds to the average number of species-level datapoints used to
generate each of the four variable parameters impacting α_S. For the right panel, point size indicates the total number
of independent host-virus associations from which virulence estimates were determined. In (C), (D), and (G), 95%
confidence intervals were computed by standard error; in (G) for the left panel, these reflect the upper and lower
confidence intervals of the optimal virus growth rate in (D). See S1 Table for order-level values for r_R^*, μ_R, T_{wR}, g_{0R},
and T_{vS} and Table 1 for all other default parameters involved in calculation of α_S. Sensitivity analyses for zoonotic
predictions are summarized in S5-S9 Figs and S3 Table.
317

We first fit a simple linear regression to the response variable of the inverse of maximum lifespan across
diverse mammalian hosts with a single predictor of reservoir host order (Fig 3A; S2 Table). From this
simple model, we easily made projections that summarized μ_R, the reservoir host annual mortality rate, to
an average value for each of 26 mammalian orders. We expressed μ_R in units of days^{-1} on a timescale
most relevant to viral dynamics; as a result, even multi-year differences in maximum longevity did little to
drive differences in μ_R across mammalian orders. Nonetheless, eight orders (Afrosoricida,
Dasyuromorphia, Didelphimorphia, Eulipotyphla, Macroscelidea, Notoryctemorphia, Peramelemorphia,
Rodentia) demonstrated significantly elevated annual mortality rates using the median order mortality rate
(Diprotodontia) as a reference. By contrast, Carnivora, Cetartiodactyla, Primates, and Perissodactyla
demonstrated significantly reduced annual mortality rates as compared to the same reference (Fig 3C;
S1-S2 Tables).
329

Drawing on well-described allometric relationships between mass and lifespan (49) and more recent
literature that links longevity with resilience to inflammation (11,44,57–61), we next sought to characterize
variation in T_{wR}, reservoir host tolerance of immunopathology, across mammalian orders. Smaller-bodied
organisms are hypothesized to be shorter-lived due to higher metabolic rates, more rapid energy
expenditure, and faster accumulation of oxidative damage, which can manifest as inflammation (62). We
hypothesized, then, that organisms that are longer-lived than predicted for their body size might be more
resilient to inflammatory stressors—and, therefore, by extension, more tolerant of immunopathology.
Building on this hypothesis, we scaled T_{wR} after order-level deviations in lifespan as predicted by body
mass. To this end, we fit a linear mixed effect regression with a random effect of host order to the $\log_{10}$
relationship of lifespan (in years) as predicted by body size (in grams), again spanning data representing
26 diverse mammalian orders. From this, we identified five orders (Carnivora, Chiroptera, Cingulata,
Monotremata, and Primates) with significantly longer lifespans than predicted by body size—which scaled

to enhanced estimates for T_{wR}. By contrast, we identified eight orders (Afrosoricida, Cetartiodactyla,
Dasyuromorphia, Didelphimorphia, Eulipotyphla, Notoryctemorphia, Peramelemorphia, and Rodentia)
with significantly shorter lifespans than predicted by body size, which scaled to lower value estimates for
T_{wR} (Fig 3A,C; S4 Fig; S1-S2 Tables). Because the same data on host lifespan were included in both
estimation of μ_R and T_{wR}, order-level estimates for T_{wR} largely mirrored those for μ_R—such that longer-
lived orders were modeled with low values for annual mortality rate and high values for tolerance to
immunopathology (and vice versa). However, because mass was not factored into estimation of μ_R, these
parameters could diverge in select cases: for example, we estimated mid-range mortality rates for order
Chiroptera, as compared with other mammals, but very high values for T_{wR} becase Chiropteran
lifespans—though not remarkably long at face value—far exceed those predicted by body size. By
contrast, we estimated low values for both μ_R and T_{wR} for order Cetartiodactyla, as hosts in this order are
long-lived but less long-lived than predicted for body size.

Finally, we fit another linear mixed effect regression with a random effect of host order to the $\log_{10}$
relationship of baseline circulating neutrophil concentration (in 10^9 cells/L) as predicted by mass (in
grams) and basal metabolic rate (W); these data spanned only 19 mammalian orders (Fig 3B). From this,
we identified a significant positive association between the orders Chiroptera and Monotremata and the
response variable of baseline neutrophil concentration, indicating that species in these orders may have
more enhanced constitutive immune responses than predicted by mass-specific BMR (Fig 3C). The
orders Cetartiodactyla, Dasyuromorphia, Diprotodontia, and Scandentia, by contrast, showed significant
negative associations, representing lower baseline neutrophil concentrations than predicted per mass-
specific BMR. We scaled these order-level effects to correspondingly high and low estimates for g_{0R}, the
within-host parameter representing the magnitude of reservoir-host constitutive immunity in our nested
modeling framework (Fig 3B,C; S4 Fig; S1-S2 Tables).

From life history-derived estimates for μ_R, T_{wR}, and g_{0R}, we next generated a prediction of the optimal
growth rate of a virus evolved in a reservoir host (r_R^*) for each of 19 distinct mammalian orders for which
we possessed the full suite of proxy data for all three variable within-host parameters (Fig 3D). In keeping
with results from our general model (Fig 2), we predicted the evolution of high r_R^* viruses from reservoir
orders exhibiting high μ_R, T_{wR}, and/or g_{0R}.Because our nested model expressed parameters in timesteps
most relevant to viral dynamics (days), differences in mortality rates across mammalian orders—though
substantial on multi-year timescales of the host—had limited influence on downstream predictions of
differences in the evolution of virus growth rates. This result echoed prior observations from Fig 2, which
showed reduced sensitivity of r_R^* to realistic variation in the magnitude of μ_R, as compared with other
parameters. As a result, we ultimately recovered the highest predicted optimal growth rates for viruses
evolved in the orders Chiroptera and Monotremata, which exhibited both long lifespans per body size
(corresponding to high estimates for T_{wR}) and high baseline neutrophil concentrations (corresponding to
high estimates for g_{0R}). Data were particularly sparse, however, for order Monotremata, for which
complete records were only available for two species (the platypus and the short-beaked echidna). As a
result, predictions for this order should be interpreted with caution. More data, particularly for baseline
neutrophil concentrations, will be needed to evaluate the extent to which these predictions hold across all
five extant species in the Monotremata order. Additionally, we predicted the evolution of the lowest growth
rate viruses in reservoir hosts of the orders Scandentia, Cetartiodactyla, Diprotodontia, and
Dasyuromorphia, all of which demonstrated significantly low estimates for the magnitudes of constitutive
immunity and two of which (Cetartiodactyla, Dasyuromorphia) also demonstrated significantly low
estimates for tolerance of immunopathology.

Estimating zoonotic virus virulence in spillover human hosts

After establishing optimal growth rates for viruses evolved in diverse reservoir host orders (r_R^*), we
subsequently modeled the corresponding 'spillover virulence' (α_S) of these viruses following emergence
into a human host. Zoonotic spillovers were modeled as acute infections in the human, and virulence was
calculated while varying only the growth rate of the spillover virus (r_R^*) and the human tolerance of direct
virus pathology (T_{vS}) between viruses evolved in differing reservoir orders. We varied this last parameter,
T_{vS}, to account for any differences in virus adaptation to reservoir host immune systems that were not
already captured in estimation of reservoir host T_{wR} and g_{0R}. T_{vS} was thus computed as the inverse of the
scaled time to MRCA for each mammalian reservoir host order from Primates (Fig 3E; S4 Fig, S1-S2
Tables), such that we estimated low human tolerance to viruses evolved in phylogenetically distant orders
(e.g. monotreme and marsupial orders), and high human tolerance to viruses evolved in Primate and
Primate-adjacent orders. In general, the modulating effects of T_{vS} did little to alter virulence rankings of
zoonotic viruses from those predicted by raw growth rate (r_R^*) alone—with the highest spillover virulence
predicted from viruses evolved in orders Monotremata and Chiroptera and the lowest spillover virulence
predicted from viruses evolved in orders Cetartiodactyla and Scandentia (Fig 3D,G). Notably, modulating
T_{vS} enhanced predictions of spillover virulence for some marsupial clades (Peramelemorphia,
Dasyuromorphia, Diprotodontia) relative to eutherian orders with similar predicted r_R^* values. This resulted
in dampened predicted spillover virulence for the eutherian order Perissodactyla as compared with
marsupial orders Dasyuromorphia and Diprotodontia, despite lower predicted r_R^* values for the latter two
clades (Fig 3D,G; S5 Fig). Similarly, this elevated spillover virulence predictions for Peramelemorphia
above Eulipotyphla, Afrosoricida, and Hyrocoidea, despite higher predicted r_R^* in the three eutherian
orders (Fig 3D,G; S5 Fig).

Comparing predictions of spillover virulence by reservoir order with estimates from the literature

Since parameter magnitudes estimated from life history traits were largely relative in nature, we scaled
predictions of spillover virulence (α_S) by reservoir order in relative terms to compare with estimates
gleaned from the zoonosis literature (8) (Fig 3G, S5-S9 Figs). For eight reservoir orders (Chiroptera,
Eulipotyphla, Primates, Carnivora, Rodentia, Diprotodontia, Perissodactyla, Cetartiodactyla), we were
able to fit a simple linear regression comparing the relative virulence of zoonoses derived from each order
as reported from case fatality rates in the literature (4,8) vs. those predicted by our nested modeling
approach (S6,S8 Figs; S3 Table). Our nested modeling framework recovered available case fatality rate
data well, yielding an R^2 value of 0.57 in the corresponding regression of observed vs. predicted values
(S6 Fig; S3 Table). Critically, we successfully recovered the key result from the zoonosis literature: bat-
derived zoonoses yield higher rates of virus-induced mortality upon spillover to humans (α_S) than do
viruses derived from all other eutherian mammals (Fig 3G; S5-S8 Fig). In general, high estimates for T_{wR},
μ_R, and g_{0R}, and low estimates for T_{vS} predicted high spillover virulence to humans (α_S). Bats demonstrate
uniquely long lifespans for their body sizes and uniquely enhanced constitutive immune responses (21)
as compared with other taxa; when combined, as in our analysis, to represent high T_{wR} and high g_{0R},
these reservoir host traits elevated predicted r_R^* and α_S beyond all other eutherian orders (Fig 3D,G).

Evaluated against the data (4,8), our model overpredicted cross-order comparative virulence rankings for
viruses evolved in order Eulipotyphla (Fig 3G; S6 Fig), largely as a result of our parameterization of high
values for annual mortality rate (μ_R) and correspondingly low values for tolerance of immunopathology
(T_{wR}) in this order. In addition, our model underpredicted virulence for Carnivora-derived viruses, based
on within-host parameter estimates largely inverse to those recovered for Eulipotyphla (e.g. low μ_R and
high T_{wR}). We were able to resolve this underprediction of Carnivora virulence when excluding rabies
lyssavirus from data comparisons (4,8) (S8 Fig); though most rabies zoonoses are sourced from domestic

dogs, lyssaviruses are Chiropteran by origin (63), and viral traits responsible for rabies' virulence may
reflect its bat evolutionary history more than that of any intermediate carnivore host. Nonetheless, while
excluding rabies from comparison improved recovery of literature-estimated relative virulence of
Carnivora-derived viruses, it somewhat destabilized predictions for other orders, such that the overall
performance of the model was largely equivalent as to when comparing against all available data (S8 Fig;
R^2 = 0.57).

In all cases, our model successfully reproduced estimates of significantly lower virulence incurred by
zoonotic viruses evolved in Cetartiodactyla hosts than in all other orders considered for comparison (8).
While previous work suggested that low observed virulence for Cetartiodactylan zoonoses might result
from overreporting of avirulent zoonoses in domestic livestock with frequent human contact (8), our
analysis indicates that viral zoonoses emerging from Cetartiodactylan hosts may truly be less virulent to
humans. Our mechanistic framework demonstrates that reduced tolerance of immunopathology (manifest
as shorter lifespans than predicted by body size) and limited constitutive immune responses identified in
Cetartiodactylan hosts may drive the evolution of low growth rate viruses that cause correspondingly
benign infections following zoonotic emergence. Further research into the extent to which
Cetartiodactylans are impacted by immunopathology, as well as the degree to which low reported
baseline neutrophil concentrations accurately reflect their innate immunity, is needed to evaluate these
predictions. Quantification of the natural growth rates of Cetartiodactylan-evolved viruses could offer one
means of testing this modeling framework.

Sensitivity analyses demonstrated that, as previously observed in Fig 2, parameters had unequal impacts
on the resulting prediction of spillover virulence in our nested modeling framework (S9 Fig; S3 Table).
Indeed, changes in g_{0R}—followed by changes in T_{vS}, and then T_{wR}—had the greatest influence on the
resulting predictions for α_S. Likelihood optimization of g_{0R} while holding μ_R , T_{wR}, and T_{vS} at order-specific
values estimated from regression analyses of publicly available life history data greatly improved our
recovery of spillover virulence as reported in the literature—yielding an R^2 value of 0.99 from the
corresponding regression of observed vs. predicted values (S9 Fig; S4 Table).

465
466

Discussion

Our work provides a mechanistic insight into the biological processes that underpin cross-species
patterns in the evolution of virus virulence—a major advance for efforts to evaluate zoonotic risk. Using a
simple model of evolving virus in a reservoir host immune system, we successfully recapture patterns
previously reported in the literature that document both the extreme virulence of viral zoonoses derived
from bats and the surprising avirulence of viruses derived from Cetartiodactylan hosts (8). Notably, our
nested modeling approach produces rank-ordered predictions of spillover virulence recovered in the
literature (8), which are distinct from those that would result from a scaled inversion of phylogenetic
distance alone. Additionally, our mechanistic approach allows us to make powerful predictions about the
virulence of potential future viral spillovers, using general life history traits across mammals, including
from orders for which zoonoses have not yet been reported (8). Our model indicates that we should
anticipate the evolution of high growth rate viruses likely to cause virulence upon cross-species
emergence from mammalian hosts with protracted lifespans for their body size (which we link to
molecular mechanisms of immunopathological tolerance (49)), as well as from hosts with robust
constitutive immune responses (36). While both immunopathological tolerance and constitutive immunity
should drive the evolution of high growth rate viruses (Fig 2), only tolerance will do so in the absence of
reservoir host pathology, thus highlighting its importance in driving observed variation in the virulence of
viral zoonoses. Notably, while tolerance reduces pathogen-induced mortality for a single host, tolerant
host populations (with limited checks on virus transmission) may demonstrate high pathogen prevalence.
If imperfectly tolerant hosts still experience some virus-induced pathology, high prevalence can
consequently elevate total population-level mortality for infected hosts—a phenomenon known as the

'tragedy of tolerance' (40,41). Reports of virus-induced mortality in bats are rare (64), suggesting that bat
virus tolerance is likely very effective.
490
Intriguingly, our independent evaluations of mammalian life history traits associated with enhanced
longevity (which we model as a proxy for immunopathological tolerance) and robust constitutive immunity
highlight several mammalian orders—chiefly Chiroptera and Monotremata—which show synergy in both
features, offering support for ongoing efforts to elucidate links between anti-aging molecular pathways
and antiviral immunity (59,64). Further bolstering these hypotheses, we demonstrate the inverse synergy
in order Cetartiodactyla, which exhibits shorter than predicted lifespans per body size but significantly
reduced constitutive immune responses as compared with other mammals. Though exciting, our
predictions should nonetheless be regarded with considerable caution, as taxon-specific insights resulting
from our modeling framework are limited by a dearth of comparative data. For example, we characterize
host tolerance of immunopathology based on deviations in lifespan as predicted for body size because
more directly representative measures of, for example, mammalian antioxidant capacity, are not
universally available. In addition, we model variation in reservoir host immunology almost exclusively
based on variation in baseline neutrophil concentrations across mammalian orders, while holding
constant many basic immunological parameters that almost certainly vary across taxa. Though crude,
baseline neutrophil data nonetheless produce an estimate of enhanced constitutive immune responses
for order Chiroptera, which is robustly supported by independent molecular work describing constitutive
interferon expression in bat cells across multiple species (21,65,66).
508
All told, this analysis highlights a critical need for compilation of a more complete comparative
immunological database that would enable quantification of the many additional within-host parameters
(e.g. leukocyte activation rate, virus consumption rate, leukocyte mortality rate, and host tolerance of
direct virus pathology) represented in our model. Our relative success in recapturing broad patterns in
spillover virulence, despite data constraints, suggests that improvements in parameter estimation will
almost certainly yield gains in our nuanced understanding of the process of cross-species virus
emergence. For example, we model host tolerance of direct virus pathology as proportional to
phylogenetic distance between reservoir and spillover host, but this term will likely also be modulated by
virus tropism—presenting yet another mechanism by which viral adaptation to reservoir hosts could
enhance spillover virulence. Indeed, coronavirus tropism is thought to be localized largely in the
gastrointestinal tract for bat hosts (67,68): in our modeling framework, higher tolerance of direct virus
pathology in this tissue should promote the evolution of higher growth rate viruses in bats, which are likely
to cause virulence upon infection of more vulnerable tissues, such as the respiratory tract, in spillover
hosts.
523
By summarizing within-host traits across mammalian orders, we generalize substantial within-clade
diversity that likely also contributes to heterogeneous patterns in available data. Bats alone make up
more than 1,400 species and account for some 20% of mammalian diversity (69). Only a subset of bats
are long-lived (70), and the magnitude of constitutive immunity is known to vary across bat species
(21,36,44), suggesting considerable variation in the parameters μ_R , T_{wR}, and g_{0R}, which we here model
universally across the entire Chiropteran order. Thus, we can expect considerable variation in the
evolution of virulence for viruses evolved in diverse species within a given order—which we largely
disregard here. As more data becomes available, our modeling approach could be fine-tuned to make
more specific, species-level predictions of highly virulent disease risk.
533
Our analysis does not consider the probability of cross species viral emergence, or the potential for
onward transmission of a spilled-over virus in a human host—both of which have been shown to correlate
inversely with phylogenetic distance across mammals (4,8,9). Indeed, in keeping with trade-off theory,
onward transmission of viruses following spillover is more commonly associated with low virulence
infections (6,8,71,72), suggesting that reservoir orders highlighted here as potential sources for high
virulence pathogens are not necessarily the same orders likely to source future pandemics. Nonetheless,
the possibility for virus adaptation to gain transmission advantages in human hosts following spillover—as
witnessed for Ebola (73) and SARS-CoV-2 (74)—should not be ignored.
542

Currently, our work emphasizes the uniqueness of bats as flying mammals and, in consequence, as long-
lived, tolerant reservoirs for highly virulent viral zoonoses. For the first time, we offer a mechanism for the
evolution of bat-derived viruses that demonstrate significant pathology upon spillover to non-bat,
particularly human, hosts. In providing a theoretical framework to explain this phenomenon, we generate
a series of testable questions and hypotheses for future comparative immunological studies, to be carried
out at *in vitro* and *in vivo* scales. Empirical work should aim to measure rates of immune cell activation,
growth, and mortality across diverse mammalian orders and determine whether natural virus growth rates
are truly higher when evolved in bat hosts. Additional studies should test whether anti-inflammatory
mechanisms in bat cells of different tissues are equally effective at mitigating virus-induced pathology and
immunopathology and whether comparative taxonomic predictions of virus tolerance, resistance, and
virulence evolution apply to non-viral pathogens, too. In light of the emergence of SARS-CoV-2, the field
of bat molecular biology has echoed the call for mechanistic understanding of bat immunology (75), and
the NIH has responded (76), soliciting research on the development of tools needed to test the
predictions outlined here. We offer a bottom-up mechanistic framework enabling the prediction of
emerging virus virulence from the basic immunological and life-history traits of zoonotic hosts.

558
559

Methods

Evolutionary framework

Population-level dynamics

To evaluate the selective pressures that drive the evolution of cross-species virus virulence, we first
sought to derive an equation for an optimal virus growth rate expressed in within-host parameters specific
to the life history of the reservoir host. To this end, and in keeping with classic examples of viral
maintenance in reservoir populations (77–83), we first modeled the dynamics of a persistent infection (I_1)
in a hypothetical reservoir host, allowing for the introduction of a rare mutant virus strain which generates
distinct infections in that same host (I_2) (see S1 File for more detailed methodology and derivations):

$$\frac{dS}{dt} = N(b_R - q_R N) - \beta_{r_{1R}} S I_1 - \beta_{r_{2R}} S I_2 - \mu_R S \qquad \text{[1a]}$$

$$\frac{dI_1}{dt} = \beta_{r_{1R}} S I_1 - \mu_R I_1 - \alpha_{r_{1R}} I_1 \qquad \text{[1b]}$$

$$\frac{dI_2}{dt} = \beta_{r_{2R}} S I_2 - \mu_R I_2 - \alpha_{r_{2R}} I_2 \qquad \text{[1c]}$$

Here, N corresponds to the total host population and S to all hosts susceptible to infection, such that $N =$
$S + I_1 + I_2$. We assumed that hosts are born at rate b_R and die of natural death at rate μ_R, where $b_R > \mu_R$.
We further assumed that all hosts are born susceptible and that population density is regulated via a
crowding term (q_R) applied to the birth rate. The subscript "R" on the birth, death, and crowding terms
emphasizes that these rates are specific to the reservoir host. Because we aimed to model the evolution
of rates that link to within-host dynamics for the reservoir host, we further represented transmission and
virulence as functions of the virus causing the infection (respectively, β_{r_R} and α_{r_R}) where r_R denotes the
intrinsic virus growth rate and is represented distinctly for both the endemic (r_{1R}) and mutant (r_{2R}) virus
strains.

With the use of model [1], we performed an evolutionary invasion analysis (see S1 File) and concluded
that the virus should evolve to maximize the ratio of transmission over infection duration. We refer to this
ratio as the invasion fitness:

$$\frac{\beta_{r_R}}{\mu_R + \alpha_{r_R}} \qquad \text{[2]}$$

Within-host dynamics

Next, to evaluate the within-host selective conditions underpinning the evolution of the virus growth rate
(r_R), we sought to express the between-host parameters of transmission (β_{r_R}) and virulence (α_{r_R}) in
within-host terms, using a nested modeling approach. To this end, we established a simple within-host
model representing the dynamics of infection *within* each I_1 and I_2 host as outlined above. We followed
Alizon and Baalen 2005 (45) to adapt a class of Lotka-Volterra predator-prey-like within-host models
(reviewed in (84)), to overcome some constraints of basic predator-prey models of the immune system,
chiefly *(i)* by allowing leukocytes to circulate in the absence of infection and *(ii)* by scaling leukocyte
growth with virion density, independent of direct leukocyte-virion contact (45,85,86). This resulted in the
following model, which demonstrates interactions between the virus population (V_R) and the leukocyte
population (L_R) within each infected reservoir host:

$$\frac{dV_R}{dt} = r_R V_R - c_R V_R L_R \qquad \text{[3a]}$$

$$\frac{dL_R}{dt} = g_{0R} + g_R r_R V_R - m_R L_R \qquad \text{[3b]}$$

Here, r_R corresponds to the intrinsic virus growth rate, c_R corresponds to the attack efficacy of the immune
system upon contact with the virus, and g_R signifies the recruitment rate of immune cells scaled to the
virus growth rate. The parameter, g_{0R}, describes the constitutive, baseline leukocyte recruitment in the
absence of infection, and m_R gives the natural leukocyte death rate, all in the reservoir host.

Building from above, we expressed rates of population-level transmission and virulence known to depend
on within-host dynamics ($\beta_{r_{1R}/r_{2R}}$ and $\alpha_{r_{1R}/r_{2R}}$) in terms of their within-host components, assuming that
within-host dynamics are fast relative to host population-level dynamics and, therefore, converge to the
endemic equilibrium (i.e. $m_R r_R > c_R g_{0R}$).

In line with previous work, we assumed transmission to be a linear function of viral load (45), which we
represented as:

$$\beta_{r_R} = \zeta V_R^*. \qquad \text{[4]}$$

where ζ corresponds to a scaling term equating viral load to transmission. We additionally assumed that:

$$\alpha_{r_R} = \frac{v r_R V_R^*}{T_{vR}} + \frac{w g_R r_R V_R^*}{T_{wR}}, \qquad \text{[5]}$$

by which infection-induced host mortality ('virulence') is modeled to result from both virus-induced
pathology (a function of the intrinsic virulence of the parasite, v, multiplied by the parasite growth rate
$r_R V_R^*$) and immunopathology—which we modeled as proportional to the leukocyte growth rate (g_R)
multiplied by the virus growth rate ($r_R V_R^*$) multiplied by the parasite's intrinsic propensity for host immune
antagonism (w). The terms T_{vR} and T_{wR} correspond to host tolerance of virus-induced pathology and
immunopathology, respectively, under assumptions of "constant" tolerance, by which both virus pathology
and immunopathology are reduced by a constant proportion across the course of infection. For constant
tolerance, we assumed that $T_{vR} > 1$ and $T_{wR} > 1$. See S1 File for detailed derivations under
assumptions of "complete" tolerance, whereby virus pathology and immunopathology are completely
eliminated up to a threshold value, beyond which pathology scales proportionally with virus and immune
cell growth.

Now, we rewrote the equations for transmission (*[4]*) and virulence (*[5]*) in purely within-host terms:

$$\beta_{r_R} = \frac{\zeta(m_R r_R - c_R g_{0R})}{c_R g_R r_R} \qquad\qquad [6]$$

$$\alpha_{r_R} = \frac{v(m_R r_R - c_R g_{0R})}{c_R g_R T_{vR}} + \frac{w(m_R r_R - c_R g_{0R})}{c_R T_{wR}} \qquad [7]$$

Using these within-host expressions, we then recomputed the expression for invasion fitness (*[2]*) in
within host terms and determined the optimal intrinsic virus growth rate, which is an evolutionarily stable
strategy (ESS) (see S1 File*)*:

$$r_R^* = \frac{c_R g_{0R}}{m_R} + \frac{\sqrt{c_R^2 m_R^2 g_R g_{0R} \mu_R T_{vR} T_{wR}(v T_{wR} + g_R w T_{vR})}}{m_R^2(v T_{wR} + g_R w T_{vR})}$$

[8]

In Fig 2, we explore the sensitivity of r_R^*, and the corresponding reservoir host population-level
transmission ($\beta_{r_R}^*$) and virulence ($\alpha_{r_R}^*$) at that growth rate, across varied values for its within-host
component parameters.

Cross-species dynamics

We next sought to derive an expression to explore the consequences of spillover of a virus evolved to its
optimal growth rate in a reservoir host population – which we termed r_R^* – following cross-species
emergence into the human population. Contrary to its established persistent infection in the reservoir
host, we assumed that such a virus would produce an acute infection in the spillover host. To model the
dynamics of this acute spillover, we borrowed from Gilchrist and Sasaki 2002 (47), who developed a
within-host parasite-leukocyte model in which an immortal leukocyte successfully eradicates the parasite
population to near-zero. We modified their acute model to be more comparable to our chronic infection
model (*[3a]/[3b])* and to reflect our differing notation for within-host dynamics.

$$\frac{dV_S}{d\tau} = r_R^* V_S - c_S V_S L_S \qquad\qquad [9a]$$

$$\frac{dL_S}{d\tau} = r_R^* g_S V_S \qquad\qquad [9b]$$

Here, r_R^* represents the reservoir-evolved virus growth rate, but all other terms reflect within-host
conditions of the spillover host: V_S and L_S correspond, respectively, to the spillover host virus and
leukocyte populations, c_S is the virus consumption rate upon contact with leukocytes in the spillover host,
and g_S is the growth rate of the spillover host leukocyte population in response to virus. We expressed the
model in units of τ, which we assumed to be short in comparison to the t time units of dynamics in the
reservoir host population. Rather than solving for virus and leukocyte populations at equilibrium (as in the
reservoir host population), we followed Gilchrist and Sasaki 2002 (47) to instead derive an expression for
the virus population at the peak of infection ($V_{S_{max}}$), where we anticipated maximum pathology for the
spillover host. We then extended this prior work to generate an expression for the average viral load
($V_{S_{avg}}$) in the spillover host across the timecourse of acute infection, which we can use in comparisons of
spillover host virulence with reported case fatality rates for zoonoses in the literature.

If we divide equation *[9a]* by equation *[9b]* we derive a simple time-independent relationship between
virus and leukocyte density:

$\frac{dV_S}{dL_S} = \frac{1}{g_S}\left(1 - \frac{c_S L_S}{r_R^*}\right)$ 　　　　　　　　　　　　　　　　　　　*[10]*

If we then let $L_S(0) = 1, V_S(0) = 1$, assuming that both virus and leukocyte populations will be small at
$\tau = 0$, we can integrate *[10]* to establish the following relationship between virus and leukocyte:

$V_S = \frac{1}{g_S}L_S - \frac{c_S}{2r_R^* g_S}\left(L_S\right)^2 + 1 - \frac{1}{g_S} + \frac{c_S}{2r_S^* g_S}$ 　　　　　　　　　*[11]*

Notice that the above *[11]* is simply a quadratic equation (see *[12]* below):

$V_S(L_S) = -\frac{c_S}{2r_R^* g_S}\left(L_S\right)^2 + \frac{1}{g_S}L_S + \left(1 - \frac{1}{g_S} + \frac{c_S}{2r_S^* g_S}\right)$ 　　　　　*[12]*

Now, we can take the derivative of equation *[12]* to formulate an expression for $V_{S_{max}}$, the maximum viral
load, which should precede the end of acute infection and the point of host recovery at the maximum
duration of infection:

$V_{S_{max}} = \frac{r_R^*}{g_S c_S} - \frac{r_R^*}{2g_S c_S} + 1 - \frac{1}{g_S} + \frac{c_S}{2r_R^* g_S}$ 　　　　　　　　　　　*[13]*

Then, extending previous work (47), we calculated the average viral load by taking the integral of
equation *[12]* from $V_S(0)$ to $V_{S_{max}}$ and dividing by the duration of that interval. From this exercise, we
expressed the average value of equation *[12]* as:

$$V_{S_{avg}} = \frac{r_R^* \sqrt{c_S^2 + 2r_R^*(g_S - 1)c + (r_R^*)^2} + (r_R^*)^2 + 4r_R^*(g-1)c + 2c_S^2}{6c_S g_S r_R^*}$$ [14]

Now, with this established expression for $V_{S_{avg}}$, we adapted equation *[7]* to reflect the acute within-host
dynamics of "spillover virulence", which we modeled (as before) as a combination of mortality induced
from direct virus pathology and from immunopathology. As in the reservoir host system, we also modeled
the mitigating impact of tolerance on the two mechanisms of virulence:

$$\alpha_S = V_{S_{avg}} \left(\frac{r_R^* v}{T_{vS}} + \frac{g_S w r_R^*}{T_{wS}} \right)$$ [15]

This yielded the above expression for 'spillover virulence'. Here, the growth rate of the virus is expressed
at its evolutionary optimum evolved in the reservoir host (r_R^*), and the virus's intrinsic virulence (v) and
propensity to elicit an inflammatory immune response (w) remained unchanged from one host to the next.
Inspired by findings in the literature that report higher virulence in cross-species infections between hosts
separated by larger phylogenetic distances (8–12), we modeled spillover host tolerance of virus-induced
pathology (T_{vS}) as a decreasing function of increasing phylogenetic distance between the reservoir and
secondary host. All other immune-related parameters assumed characteristics of the spillover host: g_S is
the spillover host's leukocyte growth rate, and T_{wS} corresponds to the spillover host's tolerance of
immunopathology.

Order-specific estimates for optimal virus growth rates evolved in reservoir hosts

Next, we sought to develop estimates for the optimal virus growth rate (r_R^*) expected to evolve across a
suite of diverse mammalian reservoir orders. Because within-host immunological data needed to quantify
within-host parameters underpinning the expression for r_R^* are largely lacking, we chose to proxy a few
key within-host parameters from well-described allometric relationships for mammalian life history data.
As such, we here focused on three key parameters for which data could be gleaned from the literature:
the reservoir host mortality rate (μ_R), the reservoir host tolerance of immunopathology (T_{wR}), and the
magnitude of reservoir host constitutive immunity (g_{0R}).

To generate order-level summary terms for μ_R, we fit a simple linear regression to the response variable
of the inverse of maximum lifespan (in days) with a single categorical predictor of host order, using data
from Jones et al. 2009 (54) and Healy et al. 2014 (55) that spanned 26 mammalian orders and 1060
individual species (Fig 3A). This first model thus took the form:

$$Y_{ij} = \alpha_0 + \sum_{j=1}^{k-1} \beta_j \delta_{ij} + \varepsilon_{ij}$$ [16]

where Y_{ij} corresponds to natural mortality rate observations for i species belonging to j orders; α_0 is the
overall intercept; k indicates the total number of available orders (here, 26); β_j is an order-specific slope;
δ_{ij} indicates observations of i species grouped into j orders; and ε_{ij} is a normally distributed error term

corresponding to species i within order j. We estimated μ_R by simply generating predictions from this fitted
model across all 26 mammalian orders represented in our dataset (Fig 3C; S2 Table).

We next used the same dataset of 1060 species grouped into 26 mammalian orders to generate order-
level estimates for the reservoir host tolerance of immunopathology (T_{wR}). Here, we fit a linear mixed
effects regression to the $\log_{10}$ relationship of lifespan (in years) as predicted by body mass (in grams).
This second model took the form:

$$Y_{ij} = \alpha_0 + \beta_1 X_1 + u_{0j} + \varepsilon_{ij} \qquad\qquad [17]$$

where Y_{ij} corresponds to the $\log_{10}$ value of record of maximum lifespan (in years) for i species belonging
to j orders; α_0 is the overall intercept; β_1 indicates the slope of the fixed predictor of $\log_{10}$ mass (in
grams), here represented as X_1; u_{0j} is the order-specific random intercept; and ε_{ij} is a normally distributed
error term corresponding to species i within order j. To estimate T_{wR}, we extracted the relative partial
effects of mammalian order on maximum lifespan per body size, then rescaled these effects between 1
and 2 for assumptions of constant tolerance and between 0 and 1 for assumptions of complete tolerance
(S1-S2 Tables). We justified this approach based on literature that highlights links between anti-aging
molecular pathways that promote longevity and those that mitigate immunopathology (11,44,57–61).

Finally, to generate order-level summary terms for the magnitude of constitutive immunity (g_0), we fit
another linear mixed effects regression to the relationship between the predictor variables of $\log_{10}$ mass
(in g) and basal metabolic rate (BMR, in W) and the response variable of $\log_{10}$ baseline neutrophil
concentration (in 10^9 cells/L), which offers an approximation of a mammal's constitutive innate immune
response. BMR data for this model were derived from Jones et al. 2009 (54) and Healy et al. 2014 (55),
while neutrophil concentrations were obtained from zoo animal data presented in the Species360
database (53); prior work using this database has demonstrated scaling relationships between body size
and immune cells across mammals (52). Paired neutrophil and BMR data were limited to just 19
mammalian orders and 144 species. Our model also included a random effect of host order, resulting in
the following form:

768
$$Y_{ij} = \alpha_0 + \beta_1 X_1 + \beta_2 X_2 + u_{0j} + \varepsilon_{ij} \qquad\qquad [18]$$

where Y_{ij} corresponds to the $\log_{10}$ baseline neutrophil concentrations (in 10^9 cells/L) for i species
belonging to j orders; α_0 is the overall intercept; β_1 indicates the slope of the fixed predictor of $\log_{10}$ mass
(in grams), here represented as X_1; β_2 indicates the slope of the fixed predictor of BMR (in W), here
represented as X_2; u_{0j} is the order-specific random intercept; and ε_{ij} is a normally distributed error term
corresponding to species i within order j. Using a similar approach as that employed above for T_{wR}, we
estimated g_0 by extracting the relative partial effects of mammalian order on neutrophil concentration per
mass-specific BMR, then rescaled these effects between 0 and 1. Because we estimated significantly
positive partial effects between the orders Chiroptera and Monotremata and baseline neutrophil
concentration (Fig 3C), this generated correspondingly high estimates of g_{0R} for these two orders (S4 Fig,
S1 Table).

780
781   Using these order-level summary terms for μ_R, T_{wR}, and g_{0R}, we then generated an order-level prediction
782   for r_R^* across all mammalian host orders, following equation [8] (Fig 3D). All other parameters (c_R, g_R, m_R,

783   w, v, and T_{vR}) were held constant across all taxa at values listed in Table 1.
784

Estimating zoonotic virus virulence in spillover human hosts

Once we had constructed an order-level prediction for r_R^*, we finally sought to explore the effects of these
reservoir-evolved viruses upon spillover to humans, following equation *[15]*. As when computing virulence
for the reservoir host population, we held immunological parameters, c_S, g_S, and m_S constant in humans
(at the same values listed above) due to a lack of informative data to the contrary. Then, to generate an
order-level estimate for virus virulence incurred on humans (α_S), we combined our order-level predictions
for r_R^* with order-specific values for T_{vS}, the human tolerance of an animal-derived virus, which we scaled
such that viruses derived from more closely-related orders to Primates were more easily tolerated in
humans. Specifically, we represented T_{vS} as the scaled inverse of the cophenetic phylogenetic distance
of each mammalian order from Primates. No summarizing was needed for T_{vS} because, following
Mollentze and Streicker 2020 (10), we used a composite time-scaled reservoir phylogeny derived from
the TimeTree database (56), which produced a single mean divergence date for all clades. Thus, all
species within a given mammalian order were assigned identical times to most recent common ancestor
with the order Primates. To convert cophenetic phylogenetic distance into reasonable values for T_{vS}, we
divided all order-level values for this distance by the largest observed (to generate a fraction), then
subtracted that fraction from 2 for assumptions of constant tolerance (yielding T_{vS} estimates ranging from
1-2) and from 1 for assumptions of complete tolerance (yielding T_{vS} estimates ranging from 0-1).
802

803   We then combined reservoir host order-level predictions for r_R^* with these estimates for human tolerance
804   of virus pathology (T_{vS}) to generate predictions of the expected spillover virulence of viral zoonoses
805   emerging from the 19 mammalian orders for which we possessed complete data across the four variable
806   parameters (μ_R, T_{wR}, g_{0R}, and T_{vS}). We scaled these α_S estimates—and their 95% predicted confidence
807   intervals—in relative terms (from 0 to 1) to compare with estimates from the literature.
808

Comparing predictions of spillover virulence by reservoir order with estimates from the literature

To compare estimates of spillover virulence generated from our nested modeling approach (α_S) with
estimates from the literature, we followed Day 2002 (87) to convert case fatality rates of viral zoonoses in
spillover human hosts (CFR_S) reported in the literature (8) to empirical estimates of α_S for each
mammalian order, using data on the duration of human infection (D_{IS}) for each viral zoonosis. For this
purpose, we collected data on D_{IS} by searching the primary literature; raw data for infection durations and
associated references are reported in our publicly available GitHub repository
(github.com/brooklabteam/spillover-virulence). Briefly, Day 2002 (13) notes that the equation for case
fatality rate in the spillover host (CFR_S) takes the form:
819

820    $$CFR_S = \frac{\alpha_S}{\alpha_S + \sigma_S}$$ *[19]*
821

822   where σ_S corresponds to the recovery rate from the spilled-over-virus in the human host. We also note
823   that the total duration of infection, D_{IS}, is given by:
824

$D_{IS} = \frac{1}{\alpha_S + \sigma_S}$ [20]

Therefore, we translated human case fatality rates of viral zoonoses (CFR_S) into estimates of spillover
virulence (α_S) using the following equation:

$\alpha_S = \frac{CFR_S}{D_{IS}}$ [21]

To obtain a composite order-level prediction for α_S from D_{IS} and CFR_S as reported in (8), we adopted the
best-fit generalized additive model (GAM) (88) used by the authors in (8) to summarize CFR_S by order
and, here, applied it to α_S estimates converted from CFR_S reported in the literature (S2 Table). We then
summarized α_S at the order-level from the fitted GAM, excluding the effects of viral family. This yielded a
literature-derived value for α_S across disparate mammalian orders, against which to compare predictions
from our nested modeling approach. Because the majority of the within-host parameters underpinning α_S
predictions in our nested modeling approach (μ_R, T_{wR}, g_{0R}, and T_{vS}) were quantified only in a relative
fashion, we were unable to compare direct magnitudes of virulence (e.g. in terms of host death per unit
time). To account for this, we rescaled α_S estimates from both our nested modeling approach and from
the empirical literature from 0 to 1 and compared the relative rank of virulence by mammalian order
instead. We predicted α_S for 19 discrete mammalian orders, though data from the literature were available
for only 8 orders against which to compare.

Finally, because bats are the most likely ancestral hosts of all lyssaviruses (63), and it may be more
appropriate to class carnivores as 'bridge hosts' for rabies, rather than reservoirs (as the authors discuss
in Guth et al. 2022 (8)), we recomputed literature-derived estimates of relative α_S by fitting the same
GAMs (S2 Table) to a version of the dataset excluding all entries for rabies lyssavirus. We rescaled these
new estimates of spillover virulence between 0 and 1 and compared them again to predictions from our
nested modeling approach.

To quantitatively evaluate the extent to which our nested model accurately recaptured estimates of
relative spillover virulence (α_S) recovered from the literature, we fit a simple linear regression to the
relationship between observed and predicted virulence across the eight mammalian orders for which we
possessed comparative data (S6, S8 Figs; S3 Table). We then conducted a sensitivity analysis to
determine the extent to which our predictions of α_S across reservoir orders could be modulated by
changes in the parameters we estimated from the literature. Because reservoir host natural mortality rate
(μ_R) had minimal influence on optimal virus growth rates (r_R^*) – and because we assumed mortality data
to be the most likely to be accurately reported in the literature – we focused our analysis on the extent to
which variation in estimation of reservoir host tolerance of immunopathology (T_{wR}), reservoir host
magnitude of constitutive immunity (g_{0R}), and spillover host tolerance of reservoir-derived virus pathology
(T_{vS}) impacted the resulting prediction for spillover virulence (α_S) (S9 Fig; S3 Table). To this end, we
quantified changes in the maximum likelihood fit of our nested model's predicted spillover virulence to the
literature-derived data while profiling each of these three parameters (T_{wR}, g_{0R}, T_{vS}) in turn across a range
of reasonable values—both while holding all other parameters at constant values across all taxa and at
heterogenous values estimated from regression analysis presented here in the main text (S9 Fig; S3
Table). To quantify the impact of this profiling on the overall recovery of relative spillover across our
dataset, we refit linear regressions of observed vs. predicted spillover virulence to each new data subset
generated from individual profiling of variable parameters (S9 Fig; S3 Table).

870
871

Acknowledgments

873 The authors thank the Boots Lab at UC Berkeley for helpful comments on this manuscript.
874
875
876

References

878
1. Bonneaud C, Longdon B. Emerging pathogen evolution. EMBO Reports. 2020;21:e51374.

2. André JB, Hochberg ME. Virulence evolution in emerging infectious diseases. Evolution.
2005;59(7):1406.

3. Bolker BM, Nanda A, Shah D. Transient virulence of emerging pathogens. Journal of the
Royal Society Interface. 2010;7(46):811–22.

4. Guth S, Visher E, Boots M, Brook CE. Host phylogenetic distance drives trends in virus
virulence and transmissibility across the animal – human interface. Phil Trans R Soc B.
2019;374(1782):20190296.

5. Farrell MJ, Davies TJ. Disease mortality in domesticated animals is predicted by host
evolutionary relationships. Proceedings of the National Academy of Sciences of the United
States of America. 2019;116(16):7911–5.

6. Mollentze N, Streicker DG, Murcia PR, Hampson K, Biek R. Virulence mismatches in index
hosts shape the outcomes of cross-species transmission. Proceedings of the National
Academy of Sciences of the United States of America [Internet]. 2020; Available from:
http://www.ncbi.nlm.nih.gov/pubmed/33122433

7. Longdon B, Hadfield JD, Day JP, Smith SCL, McGonigle JE, Cogni R, et al. The causes and
consequences of changes in virulence following pathogen host shifts. Schneider DS, editor.
PLoS Pathog. 2015 Mar 16;11(3):e1004728.

8. Guth S, Mollentze N, Renault K, Streicker DG, Visher E, Boots M, et al. Bats host the most
virulent—but not the most dangerous—zoonotic viruses. Proc Natl Acad Sci USA. 2022 Apr
5;119(14):e2113628119.

9. Olival KJ, Hosseini PR, Zambrana-Torrelio C, Ross N, Bogich TL, Daszak P. Host and viral traits
predict zoonotic spillover from mammals. Nature [Internet]. 2017; Available from:
http://www.nature.com/doifinder/10.1038/nature22975

10. Mollentze N, Streicker DG. Viral zoonotic risk is homogenous among taxonomic orders of
mammalian and avian reservoir hosts. Proceedings of the National Academy of Sciences.
2020;1–8.

11. Irving AT, Ahn M, Goh G, Anderson DE, Wang LF. Lessons from the host defences of bats, a
unique viral reservoir. Nature. 2021;589(842):363–70.

12. Schountz T, Baker ML, Butler J, Munster V. Immunological control of viral infections in bats
and the emergence of viruses highly pathogenic to humans. Frontiers in Immunology.
2017;8:1098.

13. Boots M. Fight or learn to live with the consequences. Trends in Ecology and Evolution.
2008;23(5):245–8.

14. Råberg L, Sim D, Read AF. Disentangling genetic variation for resistance and tolerance to
infectious diseases in animals. Science. 2007;318(5851):812–4.

15. Råberg L, Graham AL, Read AF. Decomposing health: tolerance and resistance to parasites
in animals. Philosophical transactions of the Royal Society of London Series B, Biological
sciences. 2009;364(1513):37–49.

16. Mougari S, Gonzalez C, Reynard O, Horvat B. Fruit bats as natural reservoir of highly
pathogenic henipaviruses: balance between antiviral defense and viral tolerance. Current
Opinion in Virology. 2022 Jun;54:101228.

17. Ng M, Ndungo E, Kaczmarek M, Herbert AS, Binger T, James R, et al. NPC1 contributes to
species-specific patterns of Ebola virus infection in bats. eLife. 2015;4:e11785.

18. Starr TN, Zepeda SK, Walls AC, Greaney AJ, Alkhovsky S, Veesler D, et al. ACE2 binding is an
ancestral and evolvable trait of sarbecoviruses. Nature. 2022 Mar 31;603(7903):913–8.

19. Letko M, Miazgowicz K, McMinn R, Seifert SN, Sola I, Enjuanes L, et al. Adaptive evolution of
MERS-CoV to species variation in DPP4. Cell Reports. 2018;24(7):1730–7.

20. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for
SARS-CoV-2 and other lineage B betacoronaviruses. Nature Microbiology. 2020;5(4):562–9.

21. Zhou P, Tachedjian M, Wynne JW, Boyd V, Cui J, Smith I, et al. Contraction of the type I IFN
locus and unusual constitutive expression of IFN-α in bats. Proceedings of the National
Academy of Sciences. 2016 Mar 8;113(10):2696–701.

22. Laing ED, Sterling SL, Weir DL, Beauregard CR, Smith IL, Larsen SE, et al. Enhanced
autophagy contributes to reduced viral infection in black flying fox cells. Viruses.
2019;11(3).

23. Chionh YT, Cui J, Koh J, Mendenhall IH, Ng JHJ, Low D, et al. High basal heat-shock protein
expression in bats confers resistance to cellular heat/oxidative stress. Cell Stress and
Chaperones. 2019;24(4):835–49.

24. Jebb D, Huang Z, Pippel M, Hughes GM, Lavrichenko K, Devanna P, et al. Six reference-
quality genomes reveal evolution of bat adaptations. Nature. 2020 Jul 23;583(7817):578–
84.

25. Hayward JA, Tachedjian M, Cui J, Cheng AZ, Johnson A, Baker ML, et al. Differential
evolution of antiretroviral restriction factors in Pteropid bats as revealed by APOBEC3 gene
complexity. Teeling E, editor. Molecular Biology and Evolution. 2018 Jul 1;35(7):1626–37.

26. O'Shea TJ, Cryan PM, Cunningham AA, Fooks AR, Hayman DTS, Luis AD, et al. Bat flight and
zoonotic viruses. Emerging Infectious Diseases. 2014 May;20(5):741–5.

27. Thomas BYSP, Suthers RA. The physiology and energetics of bat flight. Journal of
Experimental Biology. 1972;57:317–35.

28. Speakman J, Thomas D. Physiological ecology and energetics of bats. In: Kunz T, Fenton M,
editors. Bat ecology. Chicago: University of Chicago Press; 2003. p. 430–90.

29. Zhang G, Cowled C, Shi Z, Huang Z, Bishop-Lilly KA, Fang X, et al. Comparative analysis of bat
genomes provides insight into the evolution of flight and immunity. Science. 2013 Jan
25;339(6118):456–60.

30. Santillan DM, Lama T, Guti Y, Brown A, Zhao H, Rossiter S, et al. Large-scale genome
sampling reveals unique immunity and metabolic adaptations in bats. Molecular Ecology.
2021;

31. Ahn M, Cui J, Irving AT, Wang LF. Unique loss of the PYHIN gene family in bats amongst
mammals: Implications for inflammasome sensing. Scientific Reports. 2016;6(August
2015):21722.

32. Ahn M, Anderson DE, Zhang Q, Tan CW, Lim BL, Luko K, et al. Dampened NLRP3-mediated
inflammation in bats and implications for a special viral reservoir host. Nature Microbiology
[Internet]. 2019;4(May). Available from: http://dx.doi.org/10.1038/s41564-019-0371-3

33. Xie J, Li Y, Shen X, Xie J, Li Y, Shen X, et al. Dampened STING-dependent interferon
activation in bats. Cell Host and Microbe. 2018;23:1–5.

34. Goh G, Ahn M, Zhu F, Lee L, Luo D, Irving AT. Complementary regulation of caspase-1 and
IL-1 β reveals additional mechanisms of dampened inflammation in bats. Proceedings of the
National Academy of Sciences. 2020;

35. Austad SN, Fischer KE. Mammalian aging, metabolism, and ecology: evidence from the bats
and marsupials. Journal of gerontology. 1991 Mar;46(2):B47-53.

36. Brook CE, Boots M, Chandran K, Dobson AP, Drosten C, Graham AL, et al. Accelerated viral
dynamics in bat cell lines, with implications for zoonotic emergence. eLife. 2020;9:e48401.

37. Ebert D, Bull JJ. Challenging the trade-off model for the evolution of virulence: is virulence
management feasible? Trends in Microbiology. 2003 Jan;11(1):15–20.

38. Alizon S, Hurford A, Mideo N, Van Baalen M. Virulence evolution and the trade-off
hypothesis: History, current state of affairs and the future. Journal of Evolutionary Biology.
2009;22(2):245–59.

39. Anderson RM, May RM. Coevolution of hosts and parasites. Parasitology. 1982;85:411–26.

40. Miller MR, White A, Boots M. The evolution of parasites in response to tolerance in their
hosts: The good, the bad, and apparent commensalism. Evolution. 2006;60(5):945.

41. Best A, White A, Boots M. The coevolutionary implications of host tolerance. Evolution.
2014 May;68(5):1426–35.

42. Randolph HE, Barreiro LB. Holy Immune Tolerance, Batman! Immunity. 2018;48(6):1074–6.

43. Hayman DTS. Bat tolerance to viral infections. Nature Microbiology. 2019;4(5):728–9.

44. Pavlovich SS, Lovett SP, Koroleva G, Guito JC, Arnold CE, Nagle ER, et al. The Egyptian
Rousette genome reveals unexpected features of bat antiviral immunity. Cell. 2018;173:1–
13.

45. Alizon S, van Baalen M. Emergence of a convex trade-off between transmission and
virulence. The American Naturalist. 2005;165(1537-5323 (Electronic)):E155–67.

46. Geritz SAH, Kisdi É, Meszéna G, Metz JAJ. Evolutionarily singular strategies and the adaptive
growth and branching of the evolutionary tree. Evolutionary Ecology. 1998;12(1):35–57.

47. Gilchrist MA, Sasaki A. Modeling host – parasite coevolution: A nested approach based on
mechanistic models. Journal of Theoretical Biology. 2002;218:289–308.

48. Boots M, Donnelly R, White A. Optimal immune defence in the light of variation in lifespan.
Parasite Immunology. 2013;35(11):331–8.

49. Brown, James, West, Geoffrey, Enquist, Brian. Scaling in biology: Patterns and processes,
causes and consequences. In: Scaling in biology. Oxford University Press; 2000. p. 87–112.

50. Brown JH, Gillooly JF, Allen AP, Savage VM, West GB. Towards a metabolic theory of
ecology. Ecology. 2004;85(7):1771–89.

51. West GB, Woodruff WH, Brown JH. Allometric scaling of metabolic rate from molecules and
mitochondria to cells and mammals. Proceedings of the National Academy of Sciences of
the United States of America. 2002;99(SUPPL. 1):2473–8.

52. Downs CJ, Dochtermann NA, Ball R, Klasing KC, Martin LB. The effects of body mass on
immune cell concentrations of mammals. American Naturalist. 2020;195(1):107–14.

53. Species360 Zoological Information Management System (ZIMS) [Internet]. [cited 2022 Aug
25]. Available from: https://zims.species360.org/

54. Jones KE, Bielby J, Cardillo M, Fritz SA, O'Dell J, Orme CDL, et al. PanTHERIA: a species-level
database of life history, ecology, and geography of extant and recently extinct mammals.
Ecology. 2009;90(9):2648–2648.

55. Healy K, Guillerme T, Finlay S, Kane A, Kelly BA, Mcclean D, et al. Ecology and mode-of-life
explain lifespan variation in birds and mammals. Proceedings of the Royal Society B. 2014;

56. Kumar S, Stecher G, Suleski M, Hedges SB. TimeTree: a resource for timelines, timetrees,
and divergence times. Molecular Biology and Evolution. 2017;34:1812–9.

57. Wilkinson GS, Adams DM, Haghani A, Lu AT, Zoller J, Breeze CE, et al. DNA methylation
predicts age and provides insight into exceptional longevity of bats. Nature
Communications [Internet]. 2021;12(1). Available from: http://dx.doi.org/10.1038/s41467-
021-21900-2

58. Dhinaut J, Balourdet A, Teixeira M, Chogne M, Moret Y. A dietary carotenoid reduces
immunopathology and enhances longevity through an immune depressive effect in an
insect model. Scientific Reports. 2017 Dec 1;7(1).

59. Caruso C, Lio D, Cavallone L. Aging, longevity, inflammation, and cancer. Annals of New York
Academiy of Sciences. 2004;1028:1–13.

60. Fabian DK, Garschall K, Klepsatel P, Santos-Matos G, Sucena É, Kapun M, et al. Evolution of
longevity improves immunity in *Drosophila*. Evolution Letters. 2018 Dec;2(6):567–79.

61. Kolora SRR, Owens GL, Vazquez JM, Stubbs A, Chatla K, Jainese C, et al. Origins and
evolution of extreme life span in Pacific Ocean rockfishes. Science. 2021 Nov
12;374(6569):842–7.

62. Speakman JR. Body size, energy metabolism and lifespan. Journal of Experimental Biology.
2005;208:1717–30.

63. Rupprecht CE, Turmelle A, Kuzmin IV. A perspective on lyssavirus emergence and
perpetuation. Current Opinion in Virology. 2016;1(6):662–70.

64. Brook CE, Dobson AP. Bats as 'special' reservoirs for emerging zoonotic pathogens. Trends
in Microbiology. 2015;23(3):172–80.

65. Shaw AE, Hughes J, Gu Q, Behdenna A, Singer JB, Dennis T, et al. Fundamental properties of
the mammalian innate immune system revealed by multispecies comparison of type I
interferon responses. Malik H, editor. PLoS Biol. 2017 Dec 18;15(12):e2004086.

66. Clayton E, Munir M. Fundamental characteristics of bat interferon systems. Front Cell Infect
Microbiol. 2020 Dec 11;10:527921.

67. Widagdo W, Begeman L, Schipper D, Run PR van, Cunningham AA, Kley N, et al. Tissue
distribution of the MERS-coronavirus receptor in bats. Sci Rep. 2017 Dec;7(1):1193.

68. Ruiz-Aravena M, McKee C, Gamble A, Lunn T, Morris A, Snedden CE, et al. Ecology,
evolution and spillover of coronaviruses from bats. Nature Reviews Microbiology. 2021;

69. Simmons N, Cirranello A. Bat species of the world: A taxonomic and geographic database
[Internet]. 2020 [cited 2020 Jul 11]. Available from: https://batnames.org/

70. Wilkinson GS, Adams DM. Recurrent evolution of extreme longevity in bats. Biology Letters.
2019;15:20180860.

71. Geoghegan JL, Senior AM, Di F, Holmes EC. Virological factors that increase the
transmissibility of emerging human viruses. Proceedings of the National Academy of
Sciences. 2016;113(15):4170–5.

72. Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. Host phylogeny determines
viral persistence and replication in novel hosts. PLoS Pathogens. 2011;7(9).

73. Urbanowicz RA, Mcclure CP, Sakuntabhai A, Sall AA, Kobinger G, Muller MA, et al. Human
adaptation of Ebola virus during the West African Outbreak. Cell. 2016;167:1079–87.

74. Plante JA, Liu Y, Liu J, Xia H, Johnson BA, Lokugamage KG, et al. Spike mutation D614G alters
SARS-CoV-2 fitness. Nature [Internet]. 2020;(September). Available from:
http://dx.doi.org/10.1038/s41586-020-2895-3

75. Wang LF, Gamage AM, Chan WOY. Decoding bat immunity: the need for a coordinated
research approach. Nature Reviews Immunology. 2021;0123456789:0–2.

76. National Institutes of Health. Research on Bat Immunology [Internet]. Available from:
https://grants.nih.gov/grants/guide/pa-files/PAR-21-184.html

77. Viana M, Mancy R, Biek R, Cleaveland S, Cross PC, Lloyd-Smith JO, et al. Assembling
evidence for identifying reservoirs of infection. Trends in Ecology and Evolution. 2014
May;29(5):270–9.

78. Hampson K, Haydon DT. Persistent pathogens and wildlife reservoirs. Science.
2021;374(6563):35–6.

79. Haydon DT, Cleaveland S, Taylor LH, Laurenson MK. Identifying reservoirs of infection: A
conceptual and practical challenge. Emerging infectious diseases. 2002 Dec;8(12):1468–73.

80. Jolles A, Gorsich E, Gubbins S, Beechler B, Buss P, Juleff N, et al. Endemic persistence of a
highly contagious pathogen: Foot-and-mouth disease in its wildlife host. Science. 2021
Oct;374(6563):104–9.

81. Plowright RK, Peel AJ, Streicker DG, Gilbert A, McCallum H, Wood J, et al. Transmission or
within-host dynamics driving pulses of zoonotic viruses in reservoir-host populations. PLoS
Neglected Tropical Diseases. 2016;10(8):e0004796.

82. Schountz T, Prescott J, Cogswell AC, Lauren O, Mirowsky-Garcia K, Galvez AP, et al.
Regulatory T cell-like responses in deer mice persistently infected with Sin Nombre virus.
Proceedings of the National Academy of Sciences. 2007;104(39):15496–501.

83. Schountz T, Prescott J. Hantavirus immunology of rodent reservoirs: Current status and
future directions. Viruses. 2014;6:1317–35.

84. Alizon S, van Baalen M. Acute or Chronic? Within-host models with immune dynamics,
infection outcome, and parasite evolution. The American Naturalist. 2008;172(6):E244–56.

85. Alizon S, van Baalen M. Multiple infections, immune dynamics, and the evolution of
virulence. The American Naturalist. 2008;172(4):E150–68.

86. Fenton A, Lello J, Bonsall MB. Pathogen responses to host immunity: The impact of time
delays and memory on the evolution of virulence. Proceedings of the Royal Society B:
Biological Sciences. 2006;273(1597):2083–90.

87. Day T. On the evolution of virulence and the relationship between various measures of
mortality. Proceedings of the Royal Society B. 2002;269:1317–23.

88. Wood SN. mgcv: GAMs and Generalized Ridge Regression for R. R News. 2001;1/2:20–4.

Supporting Information

S1 Fig. Pairwise invasibility plots demonstrate optimal viral growth rate, under low and high tolerance conditions. Invading growth rates (r_{R2}) will displace resident growth rates (r_{R1}) at values indicated by the shaded regions. Reservoir host tolerance of immunopathology (T_{wR}) and tolerance of direct virus pathology (T_{vR}) are both modeled as low in left column (0.5 and 10 for row 1 and 2, respectively) and high in right column (0.97 and 100 for row 1 and 2), assuming a complete (row 1) or a constant form (row 2). For this visualization, r_{R1} and r_{R2} span from 3.18 to 3.5. All other parameters involved in computation of r_R^* (see S1 File equations *[25]* and *[26]*) were fixed at values listed in Table 1 (main text).

S2 Fig. Optimal virus growth rates—and subsequent spillover virulence—vary across reservoir host immunological and life history parameters. Figure replicates Fig 2 (main text) under assumptions of complete tolerance. Rows (top-down) indicate the evolutionarily optimal within-host virus growth rate (r_R^*) and the corresponding transmission rate ($\beta_{r_R}^*$), and virus-induced mortality rate ($\alpha_{r_R}^*$) for a reservoir host infected with a virus at r_R^*. The bottom row then demonstrates the resulting virulence (α_S) of a reservoir-optimized virus evolved to r_R^* upon nascent spillover to a novel, secondary host. Columns demonstrate the dependency of these outcomes on variable reservoir host parameters: background mortality rate (μ_R), extent of constitutive immunity (g_{0R}), leukocyte activation rate upon viral contact (g_R), virus consumption rate by leukocytes (c_R), leukocyte mortality rate (m_R). Darker colored lines depict outcomes at higher values for reservoir host tolerance of virus pathology (T_{vR},red) or immunopathology (T_{wR},blue), assuming no tolerance of the opposing type. Heat maps demonstrate how T_{vR} and T_{wR} interact to produce each outcome. Outcome ranges differ between lineplots (y-axes) and heatmaps (scalebars). Parameter values are listed in Table 1 (main text).

S3 Fig. Spillover host tolerance of direct virus pathology and immunopathology modulate resulting virulence of spilled-over viruses. Virulence of spilled-over virus (α_S) across a range of values for two mechanisms of tolerance in the spillover host: tolerance of direct virus pathology (T_{vS}, left column) and tolerance of immunopathology (T_{wS}, right column). Results are expressed under assumptions of constant tolerance (top panels) and complete tolerance (bottom panels). In main text results, T_{wS} is held constant for all predictions of spillover virulence but T_{vS} is varied proportionally to the inverse time to MRCA between reservoir and spillover host.

S4 Fig. Parameter estimates for life history traits across mammalian orders. Model parameter estimates for (A) reservoir-host background mortality (μ_R), (B) tolerance of immunopathology (T_{wR}) (left y-axis: constant tolerance assumptions; right y-axis: complete tolerance assumptions), (C) magnitude of constitutive immunity (g_{0R}), and (D) magnitude of human tolerance of virus pathology for a virus evolved in a disparate mammalian reservoir (T_{vS}). Estimates are derived from (A) linear model predictions of maximum lifespan at the order level, (B) the scaled effect of order on a linear mixed model prediction of lifespan per body size, (C) the scaled effect of order on a linear mixed model prediction of neutrophil concentration for mass-specific BMR, and (D) the magnitude of human tolerance of virus pathology for a virus evolved in a disparate mammalian reservoir (T_{vS}), corresponding to data presented in Fig 3E (main text). Default parameter values involved in the estimation process are summarized in Table 1 (main text), and estimated parameters and corresponding 95% confidence intervals by standard error are presented in S1 Table. See main text Methods and our open-source GitHub repository for a detailed walk-through of the parameter estimation process.

S5 Fig. Optimal virus growth rates (r_R^*) and subsequent spillover virulence (α_S) for viruses evolved in diverse mammalian reservoirs. Figure replicates Fig 3D and Fig 3G from the main text, here under assumptions of complete tolerance. Panel (A) depicts optimal r_R^* across 19 mammalian orders, for which we were able to estimate order-level specific values for the three within-host reservoir parameters which we varied in our analysis (μ_R, T_{wR}, and g_{0R}; visualized in S4 Fig), while panel (B) depicts the resulting estimation of relative spillover virulence (α_S), which also relies on order-specific values for the spillover host tolerance of direct virus pathology (T_{vS}). Taxa in panels (A) and (B) are arranged in descending order from highest to lowest predicted values for, respectively r_R^* and α_S. This order varies slightly from panel (A) to (B), as highlighted by alluvial flows and discussed in the main text. See S1 Table for order-level values for r_R^*, μ_R, T_{wR}, g_{0R}, and T_{vS} and Table 1 (main text) for all other parameters involved in calculation of r_R^* and α_S.

S6 Fig. Comparison of human spillover virulence (α_S) as observed in the literature vs. predicted from nested modeling framework. Figure plots observed vs. predicted spillover virulence for eight orders from Fig 3G (main text) for which case fatality rates from corresponding zoonoses are reported in the literature (8). Panel (A) compares nested modeling predictions under assumptions of constant tolerance with those from the literature, while panel (B) does the same under assumptions for complete tolerance. In both cases, a fitted linear regression and corresponding R^2 value is shown as a quantitative evaluation of model fit to the data. Dashed lines give the residual of each data point from the regression line.

S7 Fig. Comparison of relative human spillover virulence (α_S) predictions for zoonoses from nested life history model with estimates from the literature, excluding rabies. Figure replicates Fig 3G (main text), respectively, under assumptions of (A) constant and (B) complete tolerance but excluding rabies lyssavirus from the zoonotic data (right-half of panels). Rank-order predictions of virulence are more consistent with order Carnivora further down in the rankings. As in Fig 3G, order-specific parameter values for r_R^*, μ_R, T_{wR}, g_{0R}, and T_{vS} are listed in S1 Table; all other parameters involved in calculation of α_S are listed in Table 1 (main text).

S8 Fig. Comparison of human spillover virulence (α_S) as observed in the literature (excluding rabies) vs. predicted from nested modeling framework. Plot recapitulates S6 Fig exactly, but comparisons are drawn from case fatality rates reported in the literature but excluding rabies lyssavirus, which is often classed as a Carnivora-derived virus, though its evolutionary origins are found in bats. Removal of rabies improves estimates of virulence for Carnivora-derived zoonoses as compared with the complete dataset, but resulting linear regression offers no better fit to the entire dataset than previously shown in S6 Fig.

S9 Fig. Sensitivity analysis of individual parameter influence on nested model fit to observed data. Figure replicates S6 Fig in part with observed spillover virulence (α_S) from case fatality rates reported in the literature (8) depicted on the x-axis and predictions from nested modeling framework on the y-axis. In all panels, circles correspond to nested modeling predictions of spillover virulence using parameter values recovered from regression analysis of publicly available life history data as presented in the main text, replicating points from S6 Fig. Projections from nested modeling approach assuming constant tolerance are shown in the top panels and complete tolerance in the bottom. In lieu of life history-derived parameter values, squares show α_S estimates from nested model using universal values across order for all parameters excepting the profiled optimum of the parameter corresponding to the column in question (T_{wR}, T_{vS}, or g_{0R}). When not profiled, T_{wR}=1.5 (constant) and 0.5 (complete); T_{vS}=1.5 (constant) and 0.5 (complete); and g_{0R}=0.5 for simulations resulting in square points. Finally, triangles give α_S estimates from nested model approach using parameters generated by profiling the optimum of the parameter corresponding to the column in question (T_{wR}, T_{vS}, or g_{0R}) and using regression analysis values from the literature for other variable parameters (S1 Table). Lines and corresponding R^2 values signify the fit of a

1201 simple linear regression of observed vs. predicted α_S across all mammalian orders, where predicted
1202 values are generated from nested modeling approach using: linear regression analysis of life history data
1203 for T_{wR}, T_{vS}, and g_{0R} (solid line, red); profiling T_{wR}, T_{vS}, or g_{0R} while holding constant all other parameters
1204 across orders (thin dashed line); and profiling T_{wR}, T_{vS}, or g_{0R} while using linear regression estimates from
1205 life history data for parameters not being profiled (thick dashed line).
1206
1207 **S1 Table. Order-specific parameter values for within-host nested model.**
1208
1209 **S2 Table. Model summary outputs for within-host parameter estimation.**
1210
1211 **S3 Table. Sensitivity analysis and comparison of influence of individual parameter estimates on**
1212 **overall fit of nested model to data.**
1213
1214 **S1 File. Supplementary Information Text.**
1215
1216
1217

A

max lifespan (yrs) vs mass (g)

Legend:
- Afrosoricida
- Carnivora
- Cetartiodactyla
- Chiroptera
- Cingulata
- Dasyuromorphia
- Dermoptera
- Didelphimorphia
- Diprotodontia
- Eulipotyphla
- Hyracoidea
- Lagomorpha
- Macroscelidea
- Microbiotheria
- Monotremata
- Notoryctemorphia
- Peramelemorphia
- Perissodactyla
- Pholidota
- Pilosa
- Primates
- Proboscidea
- Rodentia
- Scandentia
- Sirenia
- Tubulidentata

B

neutrophil concentrations (10^9 cells/L) vs mass-specific BMR (W/g)

E

phylogenetic distance from primates (Myr)
- 0
- 100
- 200
- 300

Primates, Dermoptera, Scandentia, Rodentia, Lagomorpha, Carnivora, Pholidota, Perissodactyla, Cetartiodactyla, Chiroptera, Eulipotyphla, Macroscelidea, Afrosoricida, Tubulidentata, Sirenia, Hyracoidea, Proboscidea, Pilosa, Cingulata, Dasyuromorphia, Peramelemorphia, Notoryctemorphia, Diprotodontia, Microbiotheria, Didelphimorphia, Monotremata

C

μ_R , reservoir host annual mortality rate

T_{wR} , reservoir host tolerance of immunopathology

g_{0R} , reservoir host constitutive immunity

F

T_{vS} , spillover host tolerance of direct virus pathology

estimate

Taxa (C, F): Tubulidentata, Sirenia, Scandentia, Rodentia, Proboscidea, Primates, Pilosa, Pholidota, Perissodactyla, Peramelemorphia, Notoryctemorphia, Monotremata, Microbiotheria, Macroscelidea, Lagomorpha, Hyracoidea, Eulipotyphla, Diprotodontia, Didelphimorphia, Dermoptera, Dasyuromorphia, Cingulata, Chiroptera, Cetartiodactyla, Carnivora, Afrosoricida

r_R^* in reservoir host

α_S in spillover host

D

r_R^*, optimal virus growth rate in reservoir host

Taxa: Monotremata, Chiroptera, Eulipotyphla, Afrosoricida, Hyracoidea, Peramelemorphia, Didelphimorphia, Primates, Proboscidea, Pilosa, Carnivora, Rodentia, Cingulata, Tubulidentata, Perissodactyla, Dasyuromorphia, Diprotodontia, Cetartiodactyla, Scandentia

Size legend: 50, 100, 150

G

From nested model   **From zoonotic literature**

α_S, relative virulence in spillover host

Taxa: Monotremata, Chiroptera, Peramelemorphia, Eulipotyphla, Afrosoricida, Hyracoidea, Didelphimorphia, Proboscidea, Pilosa, Carnivora, Primates, Rodentia, Cingulata, Dasyuromorphia, Tubulidentata, Diprotodontia, Perissodactyla, Cetartiodactyla, Scandentia

Size legend: 50, 100, 150