1   Dramatic consequences of reducing erythrocyte membrane cholesterol on *Plasmodium falciparum*

2   Avantika I. Ahiya[a], Suyash Bhatnagar[a], Joanne Morrisey[a], Josh R. Beck[b], Akhil B. Vaidya[a*]

3   a.   *Center for Molecular Parasitology, Institute for Molecular Medicine and Infectious*

4   *Disease, Department of Microbiology and Immunology, Drexel University College of*

5   *Medicine, Philadelphia, Pennsylvania 19129*

6   b.   *Department of Biomedical Sciences, Iowa State University, Ames, Iowa 50011*

8   *To whom correspondence should be addressed at Center for Molecular Parasitology, Dept. of

9   Microbiology and Immunology, Drexel University College of Medicine, 2900 Queen Lane,

10   Philadelphia, PA 19129. Tel.: 215-991- 8557; Fax: 215-848-2271;

11   E-mail: av27@drexel.edu

Abstract

Cholesterol is the most abundant lipid in the erythrocyte. During its blood stage development, the malaria parasite establishes an active cholesterol gradient across the various membrane systems within the infected erythrocyte. Interestingly, some antimalarial compounds have recently been shown to disrupt cholesterol homeostasis in intraerythrocytic stages of *Plasmodium falciparum*. These studies point to the importance of cholesterol for parasite growth. Previously, reduction of cholesterol from the erythrocyte membrane by treatment with methyl-ß-cyclodextrin (MßCD) was shown to inhibit parasite invasion and growth. In addition, MßCD treatment of trophozoite stage *P. falciparum* was shown to result in parasite expulsion from the host cell. We have revisited these phenomena by using live video microscopy, ultrastructural analysis, and response to antimalarial compounds. By using time-lapse video microscopy of fluorescently tagged parasites, we show that MßCD treatment for just 30 min causes dramatic expulsion of the trophozoite stage parasites. This forceful expulsion occurs within 10 sec. Remarkably, the plasma membrane of the host cell from which the parasite has been expelled does not appear to be compromised. The parasitophorous vacuolar membrane (PVM) continued to surround the extruded parasite, but the PVM appeared damaged. Treatment with antimalarial compounds targeting PfATP4 or PfNCR1 prevented MßCD-mediated extrusion of the parasites, pointing to a potential role of cholesterol dynamics underlying the expulsion phenomena. We also confirmed the essential role of erythrocyte plasma membrane cholesterol for invasion and growth of *P. falciparum*. This defect can be partially complemented by cholesterol and desmosterol but not with epicholesterol, revealing stereospecificity underlying cholesterol function. Overall, our studies advance previous observations and reveal unusual cell biological features underlying cholesterol depletion of infected erythrocyte plasma membrane.

Keywords:

Malaria, *Plasmodium falciparum*, parasite extrusion, cholesterol dynamics, PfATP4 and PfNCR1 inhibitors, vomocytosis

42 **Importance:**
43
Malaria remains a major challenge in much of the world. Symptoms of malaria are caused by the

growth of parasites belonging to *Plasmodium* spp. inside the red blood cells (RBC), leading to their

destruction. The parasite depends upon its host for much of its nutritional need. Cholesterol is a

major lipid in the RBC plasma membrane, which is the only source of this lipid for malarial parasites.

We have previously shown that certain new antimalarial compounds disrupt cholesterol homeostasis

in *P. falciparum*. Here we use live time-lapse video microscopy to show dramatic expulsion of the

parasite from the host RBC when the cholesterol content of the RBC is reduced. Remarkably, this

expulsion is inhibited by the antimalarials that disrupt lipid homeostasis.   We also show

stereospecificity of cholesterol in supporting parasite growth inside RBC. Overall, these results point

to a critical role of cholesterol in physiology of malaria parasites.

Introduction:

A salient feature of malaria parasites is their dependence on the host to fulfil their nutrient requirements. In addition to various nutrients, *Plasmodium* salvages lipids and fatty acids from the host. While the parasite possesses pathways for synthesizing and modifying lipids (1, 2), it lacks the machinery for *de novo* cholesterol synthesis (3-6). Unlike the intrahepatic stages where the parasites have access to copious amounts of cholesterol (7), blood stage parasites can only access cholesterol present in the erythrocyte membrane (8). Recent studies from our laboratory have identified antimalarials that inhibit two parasite plasma membrane (PPM) transporters, PfATP4 and PfNCR1, which disrupt cholesterol and lipid homeostasis in the PPM (9, 10). Upon treatment with these compounds, there is a rapid accumulation of cholesterol in the PPM rendering the parasites sensitive to the cholesterol dependent detergent saponin. This was observed as the loss of cytosolic proteins in drug treated parasites exposed to saponin. This effect was reversible indicating an active mechanism of cholesterol dynamics within the parasite (9). These unexpected consequences of exposure to novel antimalarials suggest the presence of mechanisms that influence cholesterol transport within intraerythrocytic *P. falciparum*. Previously, it was shown that intra-erythrocytic *P. falciparum* growth requires a supply of fatty acids that cannot be substituted with lipids and cholesterol from serum derived lipoproteins (11). Furthermore, studies from the Haldar laboratory have shown that cholesterol in the erythrocyte plasma membrane is essential for invasion and growth of *P. falciparum*. Interestingly, reduction of cholesterol from the erythrocyte plasma membrane by treatment with MßCD resulted in extrusion of the trophozoite stage parasites (12).

Given the observations regarding effects of new antimalarials on cholesterol dynamics and the consequences of cholesterol reduction from infected erythrocyte plasma membrane, we wished to examine a potential link between these phenomena. We have used live fluorescence video microscopy, transmission electron microscopy and treatment with antimalarials to revisit previous findings regarding the consequences of MßCD treatment of intra erythrocytic *P. falciparum*. Results from our experiments confirm and extend these observations and raise important mechanistic questions regarding the role of cholesterol in *P. falciparum* biology.

83   **Results:**

85   ***Dramatic expulsion of trophozoite stage parasites upon treatment with*** MβCD

86   A previous study showed that treatment of trophozoite stage parasites with MβCD releases

the parasites from the erythrocyte. In this study, the authors used filipin and ethidium bromide

staining of glutaraldehyde-fixed parasites and did not assess disposition of PVM and PPM in extruded

parasites (13). We aimed to examine this effect in real-time using time lapse video microscopy as well

as to assess the disposition of PVM and PPM. For this purpose, we used two different transgenic *P.*

falciparum lines: in one line, the gene encoding PPM-localized PfVP1 was tagged with mNeonGreen

at its endogenous locus [kindly provided by Dr. Hangjun Ke; (14)] . The second line expressed two

different fluorescently tagged proteins generated by using the CRISPR/Cas9 approach as described in

Fig. 1 A and B. The gene encoding PVM-localized EXP2 was tagged with mRuby3 at its endogenous

location, and the gene encoding the RhopH complex protein RhopH2 was tagged with mNeonGreen

(Fig. 1 A and B). As shown in Fig. 1C, tagging these proteins with fluorescent markers did not affect

their proper expression and localizations. EXP2 was expressed all stages and localized to the PVM,

whereas RhopH2 was expressed at the highest level in mature stages of the parasite and localized to

the punctate rhoptries in the schizont stage. RhopH2 is initially stored in merozoite rhoptries and

secreted into the parasitophorous vacuole (PV) during invasion, eventually trafficking to the host cell

membrane (15), as seen in ring stages (Fig. 1C) . The use of these parasite lines permitted us to

monitor the PPM, PVM and rhoptries of the parasite when exposed to MßCD using live fluorescent

microscopy. The trophozoites and early schizont stage-infected erythrocytes were attached to poly-L-

lysine coated plates, stained with fluorescent probes for the RBC membrane and nuclei followed by

treatment with MßCD for 30 mins. MßCD was removed by washing with the culture medium and the

parasites were observed by live time-lapse fluorescent video microscopy. As shown in Fig. 2A and

Supplementary Video 1, the parasite extruded out of the host erythrocyte in a dramatic fashion. This

expulsion occurred 10-15 min after MβCD was washed off, but the transition from being intracellular

to being extracellular occurred in less than 10 seconds. The extruded parasites were surrounded by

PPM (in green, Fig. 2B) as well as the PVM (in red, Fig. 2C; Supplementary Fig. 1). Depending upon

the plane from which the parasite emerges from the erythrocyte, parts of the PVM were observed still

tethered to the host cell. In contrast to the late stage parasites, cholesterol depletion of ring stage

parasites did not result in extrusion; RhopH2 was partially localized to the erythrocyte plasma

membrane in the ring stage parasites, but the parasite remained inside the host cell (Fig. 2D,

Supplementary video 3). Quantification of this phenomenon showed that about 70% of the late stage

parasites extruded upon MßCD treatment (Fig. 2E).

The exclusion phenomenon was also observed by phase contrast microscopy of the infected

erythrocytes treated with 5mM MßCD, which also showed that the late stages extrude out of the

erythrocyte without causing lysis of the erythrocytes (Supplementary Fig. 2).   We assessed integrity

of the erythrocyte membrane by staining MβCD treated cultures with fluorescent phalloidin.

Phalloidin stains F-actin underlying the erythrocyte membrane when able to enter the cell. Parasite

cultures not exposed to MβCD showed minimal phalloidin staining of either infected or uninfected

erythrocytes. On the other hand, 60-70% of both uninfected and infected erythrocyte cytoskeletons

were stained with phalloidin in MβCD-treated culture (Supplementary Fig. 3). These observations

suggest that MβCD causes a breach in the erythrocyte membrane in a manner that allows phalloidin to

enter the cell but without causing catastrophic lysis that would release most of its cytosolic content.

This breach in erythrocyte membrane was seen in ring stage infected cells as well, yet this did not

result in their expulsion.

The forceful expulsion of the parasite from the host cell remains mechanistically unexplained.

MßCD treatment of mammalian cells has been shown to affect actin polymerization (16). Thus, one

possibility could be that rearrangement and polymerization of RBC cytoskeletal actin may underlie

the parasite extrusion. To test this, we treated trophozoite-infected erythrocytes with the actin

polymerization inhibitor, Cytochalasin D (CytD) (17). Treatment with CytD prior to MßCD treatment

did not inhibit MßCD mediated extrusion of the parasites. Treatment with CytD alone did not cause

the parasites to extrude out; however, we did observe defects in the PVM of the CytD treated parasites

(Fig. 2F, Supplementary video 2). Although CytD did affect the morphology of the PVM, it failed to

prevent extrusion of the parasites when exposed to MßCD.

To assess the disposition of membranes of extruded parasites in some detail, we performed

transmission electron microscopy of mature stage parasites treated with MßCD. The extruded

parasites were still surrounded by PVM, but the PVM was compromised at multiple places (Fig. 2G).

The PPM on the other hand remained intact with minimal changes in the intracellular structures.

Images shown here are representative of multiple electron micrographs.

Prior Treatment with novel antimalarials abrogates MßCD mediated parasite extrusion

We have previously shown that inhibition of PPM transporters, PfATP4 or PfNCR1 alter

cholesterol dynamics in the PPM of late stage parasites (9, 10, 18). We therefore examined the

effects of these cholesterol homeostasis disruptors on MßCD-mediated extrusion phenomenon. We

treated late stage parasites for 2 hours with KAE609 (a PfATP4 inhibitor), MMV009108 (a PfNCR1

inhibitor) or chloroquine (a heme detoxification inhibitor) followed by cholesterol depletion with

MßCD. Interestingly, the parasite extrusion was greatly reduced by prior exposure to PfATP4 and

PfNCR1 inhibitors but not by chloroquine (Fig. 3A). Quantitation of this phenomenon showed that

80% of parasites treated with DMSO or chloroquine extruded following MßCD treatment, whereas

only about 25% did so following treatment with PfATP4 and PfNCR1 inhibitors (Fig. 3B). This

quantitation was also supported by observation of intracellular and extracellular parasites in Giemsa-

stained thin blood smear (Fig. 3C and D). These results suggest that disruption of normal cholesterol

dynamics in mature stage parasite results in inhibition of MßCD mediated extrusion of parasites.

Role of erythrocyte plasma membrane cholesterol in invasion and the intra-erythrocytic

development of P. falciparum

Previous studies have shown that reduction of erythrocyte plasma membrane cholesterol by

treatment with MßCD resulted in inhibition of parasite growth (12). Since we found that the ring stage

parasites were not extruded from the erythrocyte by MßCD treatment, we went on to examine the

effects of MßCD treatment of ring stage parasites for their intra-erythrocytic developmental cycle

(IDC). In addition, we also examined the consequence of complementing cholesterol depleted ring

stage parasites with different sterols such as cholesterol, desmosterol or epicholesterol (structures

shown in Fig. 4A). We treated the ring stage infected erythrocytes with MßCD for 30 mins followed

by washes with normal culture medium. Development of ring stage parasites was assessed by

examination of Giemsa-stained thin smears. As shown in Fig. 4B, parasites in the control cells

proceeded normally through the IDC and formed trophozoites and rings. In contrast, ring stage

parasites growing in cholesterol depleted erythrocytes (MßCD treated) progressed to form what

appeared to be trophozoites but failed to mature. Reconstitution with MßCD saturated with

cholesterol (CD/Cho) or desmosterol (CD/Des) appeared to restore normal IDC progression. On the

other hand, reconstitution with epicholesterol (CD/Epi) did not restore normal IDC, suggesting the

importance of stereospecificity of cholesterol polar moiety in supporting parasite development. We

also attempted to reconstitute with lanosterol or ß-sitosterol but could not assess it since these resulted

in lysis of erythrocytes. To assess the viability of the ring stage parasites growing in MßCD-treated or

sterol-reconstituted erythrocytes, we split the culture 1:10 into untreated normal erythrocytes. This

was followed by measuring parasitemia over a subsequent four-day period. As shown in Fig 4C, ring

stage parasites growing in cholesterol-depleted or epicholesterol-reconstituted cells failed to

propagate over the next two cycles. On the other hand, cholesterol- or desmosterol-reconstituted ring

stage parasites were able to propagate at a level of about 50% compared to the control ring stage

parasites. To assess progression of ring stage parasites, we measured hypoxanthine incorporation

over a 24-hour period. As shown in Fig 4D, there was significant reduction in hypoxanthine

incorporation in MßCD treated and epicholesterol reconstituted parasites, whereas ring stage parasites

in cholesterol or desmosterol reconstituted cells were able to incorporate hypoxanthine.

We examined the parasites by transmission electron microscopy at 24 and 48 hours in control

and following MßCD treatment. As expected at 24 hours the trophozoites in the control group

displayed normal morphology with distinctive internal structures and normal hemozoin formation and

went on to form ring stages at 48 hours (Fig 4E, top panel; Supplementary Fig. 1). In contrast MßCD

treated ring stages showed abnormal morphology with diminished food vacuoles at 24 hours. These

parasites failed to develop into normal schizonts and did not egress at 48 hours after MßCD treatment.

(Fig 4E, lower panel; Supplementary Fig 4). These results suggest a defect in normal trophozoite

development in erythrocytes with depleted cholesterol content.

We also assessed the ability of merozoites to invade and establish infection in MßCD treated

erythrocytes. MßCD treated erythrocytes were added to Percoll enriched schizonts. As shown in Fig

5A, reduced cholesterol content in uninfected erythrocytes prevented merozoite invasion. The

merozoites appeared to remain attached to the surface of MßCD treated erythrocytes but failed to

penetrate. Importantly, reconstitution of MßCD treated RBCs with cholesterol complemented this

invasion defect (Fig. 5B). Furthermore, the parasites were able to grow in these reconstituted

erythrocytes and undergo normal IDC progression. Splitting the cultures 1:10 with addition of normal

erythrocytes resulted in continued normal IDC progression of parasites from control and cholesterol

reconstituted cells but not from cholesterol deficient cells (Fig. 5C).

Different *P. falciparum* strains invade using different pathways: a sialic acid dependent or a

sialic acid independent pathway (19, 20). To see if cholesterol in the erythrocyte membrane is

selective for a given pathway, we added schizonts of either NF54 (sialic acid dependent) or Dd2

(sialic acid independent) strains to MßCD treated erythrocytes. Merozoites from neither strain were

able to invade MßCD treated erythrocytes (Fig. 5D).

Discussion:

Using live video microscopy of fluorescently tagged parasites, we have extended previous

observations showing expulsion of trophozoite stage *P. falciparum* upon treatment with MßCD. We

also showed that this expulsion is inhibited by new antimalarial compounds that disrupt cholesterol

homeostasis in the parasite. The dramatic expulsion of the trophozoite stage parasites following

MßCD treatment remains unexplained. It is hard to perceive how an intracellular parasite can escape

out of the host erythrocyte without causing catastrophic lysis of the host membrane in response to

reduction in the cholesterol content in the erythrocyte plasma membrane. Expulsion of the parasite

occurs after 40-45 min exposure to MβCD and after MβCD is washed off. Actual expulsion,

however, occurs within seconds. The force with which the expulsion appears to occur suggest an

active process mediated either by the host or the parasite. Our observation that Cytochalasin D did not

prevent expulsion of trophozoites indicates that actin polymerization is not required for this process.

Osmotic pressure build-up through solute uptake in trophozoites resulting in their release from the

host cells has been widely used for synchronization of blood stage *P. falciparum*. However, this

requires incubation with high (about 0.5 M) concentration of solutes such as sorbitol. At 5 mM

concentration, MßCD exposure does not pose an osmotic challenge, and at ~1300 Da molecular

weight MßCD would not be taken up by the trophozoite-infected erythrocyte through its new

permeability pathway. Furthermore, sorbitol-mediated expulsion of the parasite occurs within a few

minutes and is accompanied by catastrophic lysis of the infected erythrocyte. In contrast, expulsion

caused by MβCD-mediated cholesterol depletion occurs after 40-45 min when MβCD is no longer

present, and the infected erythrocyte does not undergo lysis. Also, the MβCD-mediated expulsion is

inhibited by prior treatment with PfATP4 and PfNCR1 inhibitors. These observations seem to argue

against an osmotic imbalance as the force underlying the expulsion phenomenon described here. We

speculate the possibility of a hitherto unrecognized process through which the parasite might sense

cholesterol content of the erythrocyte membrane. It is of interest to note that MßCD-mediated

expulsion of the parasite resembles expulsion of fungal pathogens from the mammalian cells through

a process termed vomocytosis wherein the expulsion does not result in lysis of the host cell (21-23).

Inhibition of the parasite expulsion by compounds that inhibit PfATP4 or PfNCR1was

unexpected; this inhibition was not a result of parasite demise since chloroquine failed to do so. We

have previously shown that both these inhibitors induce cholesterol accumulation in the PPM in a

reversible manner (10). PfATP4 inhibition results in Na^+ influx into the parasite and a collapse of Na^+

gradient across the PPM. We propose that PfNCR1 requires a Na^+ gradient to maintain cholesterol

homeostasis, and its collapse by PfATP4 inhibition results in indirect inhibition of PfNCR1. The net

result of both PfATP4 and PfNCR1 inhibition, therefore, is a disruption of cholesterol homeostasis in

late-stage *P. falciparum*. We envision a dynamic movement of cholesterol between various

membranes of infected erythrocytes. Recent investigations have also supported the notion of

cholesterol gradients in infected erythrocytes (24-28). Reduction of the plasma membrane cholesterol

content, thus, appears to be sensed by the parasite in a manner that results in the expulsion of

trophozoites.

The erythrocyte plasma membrane contains large amounts of cholesterol (50 mole% of total

lipids). Upon infection by *Plasmodium*, the PVM acquires cholesterol but the PPM remains highly

deficient in cholesterol. Indeed, this fact is the basis for widely used saponin mediated freeing of

intra-erythrocytic parasites. Previous studies have suggested the importance of cholesterol for *P.*

falciparum growth and invasion (12, 13, 25). A recent study using lattice light scattering microscopy

demonstrated the importance erythrocyte membrane remodeling for the formation of the PVM, a

process in which cholesterol is likely to play a significant role [11]. We observed that that ring stage

parasites growing in cholesterol deficient erythrocytes fail to fully progress. Furthermore, the

digestive vacuoles of trophozoite-like parasites at 24 hours after MßCD treatment of ring stages had

unusual morphology (Fig.3E and Supplementary figure 1). At 48 hours after MßCD treatment of ring

stages, parasites appeared to attempt segmentation without being able to form mature merozoites (Fig.

3E and Supplementary figure 1). One possible explanation could be that cholesterol dynamics assist

lipid and/or fatty acid import to sustain parasite growth. The fact that cholesterol and desmosterol (but

not epicholesterol) could substantially rescue these defects suggest stereospecificity in the role of

cholesterol to sustain parasite growth. Further studies will be required to gain mechanistic insights

into cholesterol dynamics and their contribution to parasite growth.

Methods:

Parasite lines and culture conditions

Experiments were carried out using different *P. falciparum* tagged lines. 1) the 3D7 strain in

which endogenous PfVP1 gene was tagged with mNeon green (mNG) at the C-terminal end

(PfVP1-mNG) (14). This line was used for monitoring the dynamics of the parasite plasma

membrane; 2) the NF54 strain in which endogenous RhopH2 gene was tagged with mNG and

the endogenous Exp2 gene tagged with mRuby3 (NF54 Rhoph2/Exp2). Cloning was carried

out with Infusion (Clontech) and NEBuilder HiFi (NEB). To generate an endogenous EXP2-

mRuby3 fusion, 5' and 3' homology flanks targeting the 3' end of *exp2* were PCR amplified

from plasmid pyPM2GT-EXP2-mNG (29) using primers

CACTATAGAACTCGAGGGAGAAACAATCTTTTATATAAAATGTACAGAGTTTGA

AAG and

TCCTCCACTTCCCCTAGGTTCTTTATTTTCATCTTTTTTTTCATTTTTAAATAAATC

TCCAC and inserted between *XhoI* and *AvrII* sites in the plasmid pbPM2GT (30). The

mRuby3 coding sequence was then amplified from plasmid pLN-HSP101-SP-mRuby3(31)

using primers GATGAAAATAAAGAACCTAGGGGAAGTGGAGGAGTG and

TAACTCGACGCGGCCGTCACTTGTACAGCTCGTCCATGCC and inserted between

XhoI and *AvrII*, resulting in the plasmid pbEXP2-mRuby3. This plasmid was linearized at the

AflII site between the 3' and 5' homology flanks and co-transfected with pUF-Cas9-EXP2-

CT-gRNA into NF54^{attB}::HSP101^{DDD} (31) . Parasites were maintained in 10 µM trimethoprim

(TMP) to stabilize the HSP101^{DDD} fusion and selection with 2.5 µg/ml Blasticidin S was

applied 24 hours post transfection. A clonal line bearing the EXP2-mRuby3 fusion was

derived by limiting dilution after parasite returned from selection and designated

NF54^{attB}::HSP101^{DDD}+EXP2-mRuby3.

For generation of an endogenous RhopH2-mNG fusion, a gRNA target site was chosen

upstream of the *rhoph2* stop codon (TCTTCACTGATTTCTTTGTA) and the gRNA seed

sequence was synthesized as a sense and anti-sense primer pair (sense shown)

TAAGTATATAATATTTCTTCACTGATTTCTTTGTAGTTTTAGAGCTAGAA, annealed

and inserted into the *AflII* site of the plasmid pAIO3 (32), resulting in the plasmid pAIO3-

RhopH2-CT-gRNA. To integrate mNG at the 3' end of *rhoph2*, a 5' homology flank (up to

but not including the stop codon) was amplified from NF54^attB genomic DNA using primers

AATTCATCATTATGAAAGTTCTCAGCTTAAGAAGCATATATTAAGAATATAGTT

TCAGA and

CCTCCACTTCCCCTAGGACTGCTCTTCAGAATATACAGGTTTTTTATAAGATCCTC

CGATATCTCCTTATATGGATCAGATATATCTGAGAAA, incorporating several

synonymous mutations in the seed sequence of the gRNA target site within the *rhoph2*

coding sequence. A 3' homology flank (beginning at the endogenous stop codon) was

amplified using primers

GTGACACTATAGAACTCGAGTAAACGTTAAAAAAAAAATATATATAAGGAGAAA

GCACTG and

TCTGAAACTATATTCTTAATATATGCTTCTTAAGCTGAGAACTTTCATAATGATGA

AATT, assembled in a second PCR reaction using primers

GTGACACTATAGAACTCGAGTAAACGTTAAAAAAAAAATATATATAAGGAGAAA

GCACTG and

CCTCCACTTCCCCTAGGACTGCTCTTCAGAATATACAGGTTTTTTATAAGATCCTC

CGATATCTCCTTATATGGATCAGATATATCTGAGAAA and inserted between *XhoI*

and *AvrII* sites in pyPM2GT-EXP2-mNG (29) resulting in the plasmid pyPM2GT-RhopH2-

mNG. This plasmid was linearized at the *AflII* site between the 3' and 5' homology flanks

and co-transfected with pAIO3-RhopH2-CT-gRNA into NF54^attB::HSP101^DDD+EXP2-

mRuby3. Selection with 2 μM DSM-1 was applied 24 hours post transfection (along with 10

μM trimethoprim). After parasites returned from selection, a clonal line containing the EXP2-

mRuby3 and RhopH2-mNG fusions was obtained by limiting dilution and designated

NF54^attB::HSP101^DDD+EXP2-mRuby3+RhopH2-mNG. Correct integration of the transgenes

was confirmed by PCR amplifications using the following primers (indicated in Fig. 1):

A:CCAGAATGTTTCGGACCATGTAC; B: TGGTATCCGGAGCCATCTACCATG; C:

CACTTTGTAACTTCATTTTCTAAAATGACCTTGTTC; D: GCAACAAGTGCCTTAACCACCG; E:

CGATGACTTTGATCCTCATGGTTTGC; and F: TCACTTATGTTGTATAGAGACACAATTCGT

This line was used for monitoring the parasitophorous vacuolar membrane and to mark the

parasite. *P. falciparum* parasites were cultured in O^+ human blood from Interstate Blood

Bank, TN in RPMI 1640 supplemented with 2g/L sodium bicarbonate, 10mg/L

hypoxanthine, 15mM HEPES, 50mg/L gentamycin sulphate, 0.5% AlbuMax II. Parasite

culture was maintained at 2.5% hematocrit at 37 °C in 90% N_2, 5% CO_2, 5% O_2.

Preparation of MßCD and MßCD-sterol complexes

Parasites were treated with 5mM MßCD diluted in appropriate medium from a stock solution

of 25mM MßCD (Catalog no. AC377110250) in PBS in appropriate medium. Sterols

[Cholesterol (Sigma, C3045-25G) , Desmosterol (Steraloids, catalog no. C3150-000),

Epicholesterol (Steraloids, catalog no. C6730-000)] were loaded onto MßCD according to

previously published protocol (33). Briefly, 10 μL of 15mg/ml sterols dissolved in ethanol

were added to 500 μL 5% w/v MßCD, heated for 10 mins at 80°C and mixed by inverting

several times until the solution was clear. The above step was repeated 4 times to add a total

of 50μL sterol stock solution. The solution was heated until the sterols were stably

incorporated in MßCD (clear solution). MßCD/sterol complexes were snap frozen in dry ice

for 2 mins. Subsequently, the MßCD/sterol complexes were lyophilized in a speed vac until

all liquid had evaporated and a fluffy powered remained at the bottom of the tube and stored

at -20°C. Immediately before use, 375μl of medium was added to the MßCD/sterol

complexes and vortexed until dissolved. The MßCD/sterol complexes were sterilized using

0.22μm syringe filter.

Treatment of parasitized erythrocytes with MBCD and reconstitution with

MBCD/sterol complexes

Parasite cultures were synchronized by treatment with 0.5M alanine for 10 minutes. Erythrocyte infected with ring stage parasites were treated with 5mM MßCD for 30 mins at 37 °C and washed 3 times with pre-warmed RPMI1640 media. Hematocrit was maintained at 2.5% throughout the culture and washing steps. The parasites were returned to normal culture conditions. In experiments involving reconstitution with MßCD/sterol complexes, parasites were first treated with MßCD as described above. Following this, the cells were incubated with 1:7 dilution of MßCD/sterol complexes for 30 mins. Parasites were washed 3 times and returned to normal culture conditions. Giemsa-stained thin blood smears were prepared 24 and 48 hours after the treatment with MßCD or after reconstitution with MßCD/sterol complexes. The culture at 48 hours was split 1:10 in normal erythrocytes. Growth of the parasites in these cultures were followed for 4 days by examining Giemsa stained thin blood smears

Hypoxanthine incorporation of ring stage erythrocyte treated with MßCD or

MßCD/sterol complexes

Ring stage erythrocyte (1.5% parasitemia) were treated with MßCD for 30 mins and washed 3 times with normal medium. The hematocrit was maintained at 2.5%. MßCD-treated ring stage erythrocytes were then treated with either MßCD/cholesterol, MßCD/desmosterol or MßCD/epicholesterol for 30 mins and washed 3 times with low hypoxanthine medium. The hematocrit was then adjusted to 1.5% by addition of low hypoxanthine medium to the cells. 200μl of cells were then plated in triplicates in a 96-well plate. Each well was pulsed with 22 μl of 0.5 μCi/ml ^{3}H-hypoxanthine (PerkinElmer NET 177) and incubated for 24 hours at 37ºC in 5% CO_2, 5% O_2, 90% N_2 chamber. Parasites were lysed by freeze/thaw and were

373   collected on filters using a cell harvester (PerkinElmer LifeSciences). This was followed by

374   addition of MicroScint O scintillation cocktail and incorporation of ^{3}H-hypoxanthine was

375   measured using a TopCount scintillation counter (PerkinElmer Life-sciences).

Parasite invasion in MßCD or MßCD/Cholesterol treated erythrocytes

We treated 500 µl of 50% hematocrit erythrocytes with 5mM MßCD for 30 mins followed by

3 washes. In addition, similarly treated erythrocytes were reconstituted with

MßCD/Cholesterol for 30 mins followed by 3 washes. Synchronized late stages trophozoites

were enriched by centrifugation over a 70% Percoll cushion (3000 RPM, 20 mins) and

washed thrice with medium. Enriched late stage trophozoites were added to MßCD-treated,

MßCD/Cholesterol-reconstituted and control erythrocytes. Giemsa-stained thin blood smears

were prepared 24 and 48 hours following the addition. The culture at 48 hours was split 1:10

in normal erythrocytes. Growth of the parasites in these cultures were followed for 4 days by

examining Giemsa-stained thin blood smears.

Live microscopy of parasites

Glass bottom culture dishes (35 mm) were coated with 0.1% poly-L-lysine overnight and

washed 3 times with PBS. Parasite culture (250µl at 2.5% hematocrit) was added to the

dishes and incubated for 30 mins to allow attachment. Culture dishes were washed 3 times

with PBS followed by addition of medium. For MßCD extrusion experiments with NF54

Rhoph2/Exp2 line, all washes, incubation and imaging was done in medium supplemented

with TMP. Parasites were attached to poly-L-lysine coated plates and incubated in RPMI

containing wheat germ agglutinin-Alexa 350 (WGA-350) (Invitrogen, Catalog no. W7024) at

5µg/ml final concentration and incubated for 12 mins to stain the erythrocyte plasma

membrane. In addition to WGA staining, PfVP1-mNG line was also stained with SYTO deep

red nuclear stain for 30 min. Parasites were treated with 5mM MßCD followed by three

washes with medium. The medium was replaced with phenol red free medium followed by

live fluorescent microscopy. Imaging was done using the Nikon Ti microscope. WGA-Alexa-

350 was visualized using DAPI filter set, mNeon green with FITC, SYTO deep red nuclear

stain with Cy5 and mRuby was visualized using TRITC filter set. Video microscopy

parameters are given in the legends of Supplementary Videos.

Inhibition of MßCD mediated extrusion by novel antimalarials.

To look at the effects of antimalarials on MßCD mediated extrusion, parasites were stained

with WGA-350 and SYTO deep red for PfVP-1mNG line or WGA-350 for NF54

Rhoph2/Exp2 line as described above and treated with $10xEC_{50}$ concentrations of

Choloroquine (150nM), KAE609 (10nM) or MMV009108(1μM) for 2.5 hours prior to 30

mins treatment with MßCD and washed thrice with medium. Medium supplemented with

drugs (at the concentration mentioned above) was added back to the dish and live fluorescent

imaging was carried out. Quantification of parasites remaining inside/outside of erythrocytes

was carried out by examining at least 100 different images of individual parasites for each

experimental condition.

Cytochalasin D (Cyt-D) treatment prior to MßCD treatment of parasites

Attached NF54 RhopH2/Exp2 trophozoites were stained with WGA-350 for 12 mins,

followed by three washes. The cells were then treated with 0.5μM Cyt-D for 45 mins. This

was followed by the addition of 5mM MßCD for 30 mins. The culture dishes were washed 3

times and live fluorescent microscopy was performed to examine parasite extrusion. In the

control condition, parasites were treated with Cyt-D without MßCD treatment.

Transmission electron microscopy of trophozoite infected MßCD treated erythrocytes

Erythrocytes infected with trophozoite stage parasites were treated with 5mM **MßCD** for 30

mins at 2.5% hematocrit followed by three washes with PBS and incubated in complete

medium for 30 mins at 37 °C. Cells were washed once in PBS by gentle centrifugation and

fixed with 2% paraformaldehyde, 2.5% glutaraldehyde in 100 mM cacodylate buffer and

were fixed for 1 hour at room temperature. Cells were gently pelleted by centrifugation,

resuspended in cacodylate buffer and immediately stored on dry ice for shipment.

Ring stage infected erythrocytes were treated with 5mM for 30 mins, washed thrice with

PBS and returned to culture at 37 °C. At 24 and 48 hours after the treatment, infected

erythrocytes were fixed and prepared for transmission electron microscopy as described

above.

Phalloidin staining of MßCD treated cultures

Infected erythrocyte cultures from the PfVP1-mNG line were treated with 5mM MßCD for

30 mins at 37 °C and washed 3 times with prewarmed RPMI media. Phalloidin staining was

carried out as described (34). Briefly, the MßCD treated cultures were stained with

Phalloidin Alexa 594 (Invitrogen; diluted 1:50 from a 200 unit ml^{-1} stock in methanol) for 5

mins at 37 °C, washed thrice with prewarmed phenol red free RPMI1640. The cells were

added to the glass slides and sealed with greased coverslips. Images were taken using the

Nikon Ti microscope. mNeongreen was visualized using the FITC channel, Phalloidin Alexa

594 was visualized using the TRITC filter set.

Acknowledgements:

We thank the members of the Vaidya lab for lively discussion and input. We thank Dr.

Hangjun Ke for providing PfVP1-mNG parasite line. This work was supported by a National

Institutes of Health grant R01 AI132508 to ABV and R00 HL133453 to J.R.B.

References:

1.   Cassera MB, Gozzo FC, D'Alexandri FL, Merino EF, del Portillo HA, Peres VJ,
Almeida IC, Eberlin MN, Wunderlich G, Wiesner J, Jomaa H, Kimura EA, Katzin
AM. 2004. The methylerythritol phosphate pathway is functionally active in all
intraerythrocytic stages of Plasmodium falciparum. J Biol Chem 279:51749-59.
2.   Gerold P, Schwarz RT. 2001. Biosynthesis of glycosphingolipids de-novo by the
human malaria parasite Plasmodium falciparum. Mol Biochem Parasitol 112:29-37.
3.   Holz GG, Jr. 1977. Lipids and the malarial parasite. Bull World Health Organ 55:237-
48.
4.   Maguire PA, Sherman IW. 1990. Phospholipid composition, cholesterol content and
cholesterol exchange in Plasmodium falciparum-infected red cells. Mol Biochem
Parasitol 38:105-12.
5.   Vial HJ, Ancelin ML, Philippot JR, Thuet MJ. 1990. Biosynthesis and dynamics of
lipids in Plasmodium-infected mature mammalian erythrocytes. Blood Cells 16:531-
55; discussion 556-61.
6.   Vial HJ, Philippot JR, Wallach DF. 1984. A reevaluation of the status of cholesterol
in erythrocytes infected by Plasmodium knowlesi and P. falciparum. Mol Biochem
Parasitol 13:53-65.
7.   Labaied M, Jayabalasingham B, Bano N, Cha SJ, Sandoval J, Guan G, Coppens I.
2011. Plasmodium salvages cholesterol internalized by LDL and synthesized de novo
in the liver. Cell Microbiol 13:569-86.
8.   Grellier P, Rigomier D, Clavey V, Fruchart JC, Schrevel J. 1991. Lipid traffic
between high density lipoproteins and Plasmodium falciparum-infected red blood
cells. J Cell Biol 112:267-77.
9.   Das S, Bhatanagar S, Morrisey JM, Daly TM, Burns JM, Jr., Coppens I, Vaidya AB.
2016. Na+ Influx Induced by New Antimalarials Causes Rapid Alterations in the
Cholesterol Content and Morphology of Plasmodium falciparum. PLoS Pathog
12:e1005647.
10.   Istvan ES, Das S, Bhatnagar S, Beck JR, Owen E, Llinas M, Ganesan SM, Niles JC,
Winzeler E, Vaidya AB, Goldberg DE. 2019. Plasmodium Niemann-Pick type C1-
related protein is a druggable target required for parasite membrane homeostasis.
Elife 8.
11.   Coppens I, Vielemeyer O. 2005. Insights into unique physiological features of neutral
lipids in Apicomplexa: from storage to potential mediation in parasite metabolic
activities. Int J Parasitol 35:597-615.
12.   Samuel BU, Mohandas N, Harrison T, McManus H, Rosse W, Reid M, Haldar K.
2001. The role of cholesterol and glycosylphosphatidylinositol-anchored proteins of
erythrocyte rafts in regulating raft protein content and malarial infection. J Biol Chem
276:29319-29.
13.   Lauer S, VanWye J, Harrison T, McManus H, Samuel BU, Hiller NL, Mohandas N,
Haldar K. 2000. Vacuolar uptake of host components, and a role for cholesterol and
sphingomyelin in malarial infection. The EMBO journal 19:3556-3564.

14.   Solebo O, Ling L, Zhou J, Fu T-M, Ke H. 2021. Plasmodium
falciparum vacuolar pyrophosphatase 1 for ring stage development and its
transition to trophozoite. bioRxiv
doi:10.1101/2021.10.25.465524:2021.10.25.465524.
15.   Ahmad M, Manzella-Lapeira J, Saggu G, Ito D, Brzostowski JA, Desai SA. 2020.
Live-Cell FRET Reveals that Malaria Nutrient Channel Proteins CLAG3 and RhopH2
Remain Associated throughout Their Tortuous Trafficking. mBio 11.
16.   Guerra FS, Sampaio LdS, Konig S, Bonamino M, Rossi MID, Costa ML, Fernandes
P, Mermelstein C. 2016. Membrane cholesterol depletion reduces breast tumor cell
migration by a mechanism that involves non-canonical Wnt signaling and IL-10
secretion. Translational Medicine Communications 1:3.
17.   Casella JF, Flanagan MD, Lin S. 1981. Cytochalasin D inhibits actin polymerization
and induces depolymerization of actin filaments formed during platelet shape change.
Nature 293:302-5.
18.   Bhatnagar S, Nicklas S, Morrisey JM, Goldberg DE, Vaidya AB. 2019. Diverse
Chemical Compounds Target Plasmodium falciparum Plasma Membrane Lipid
Homeostasis. ACS infectious diseases 5:550-558.
19.   Camus D, Hadley TJ. 1985. A Plasmodium falciparum antigen that binds to host
erythrocytes and merozoites. Science 230:553-6.
20.   Dolan SA, Miller LH, Wellems TE. 1990. Evidence for a switching mechanism in the
invasion of erythrocytes by Plasmodium falciparum. J Clin Invest 86:618-24.
21.   Cruz-Acuña M, Pacifici N, Lewis JS. 2019. Vomocytosis: Too Much Booze, Base, or
Calcium? mBio 10.
22.   Nicola AM, Robertson EJ, Albuquerque P, Derengowski Lda S, Casadevall A. 2011.
Nonlytic exocytosis of Cryptococcus neoformans from macrophages occurs in vivo
and is influenced by phagosomal pH. mBio 2.
23.   Seoane PI, May RC. 2020. Vomocytosis: What we know so far. Cell Microbiol
22:e13145.
24.   Fraser M, Jing W, Bröer S, Kurth F, Sander LE, Matuschewski K, Maier AG. 2021.
Breakdown in membrane asymmetry regulation leads to monocyte recognition of P.
falciparum-infected red blood cells. PLoS Pathog 17:e1009259.
25.   Geoghegan ND, Evelyn C, Whitehead LW, Pasternak M, McDonald P, Triglia T,
Marapana DS, Kempe D, Thompson JK, Mlodzianoski MJ, Healer J, Biro M,
Cowman AF, Rogers KL. 2021. 4D analysis of malaria parasite invasion offers
insights into erythrocyte membrane remodeling and parasitophorous vacuole
formation. Nat Commun 12:3620.
26.   Hayakawa EH, Yamaguchi K, Mori M, Nardone G. 2020. Real-time cholesterol
sorting in Plasmodium falciparum-erythrocytes as revealed by 3D label-free imaging.
Sci Rep 10:2794.
27.   Tokumasu F, Crivat G, Ackerman H, Hwang J, Wellems TE. 2014. Inward
cholesterol gradient of the membrane system in P. falciparum-infected erythrocytes
involves a dilution effect from parasite-produced lipids. Biol Open 3:529-41.
28.   Koch M, Cegla J, Jones B, Lu Y, Mallat Z, Blagborough AM, Angrisano F, Baum J.
2019. The effects of dyslipidaemia and cholesterol modulation on erythrocyte
susceptibility to malaria parasite infection. Malar J 18:381.
29.   Glushakova S, Busse BL, Garten M, Beck JR, Fairhurst RM, Goldberg DE,
Zimmerberg J. 2017. Exploitation of a newly-identified entry pathway into the
malaria parasite-infected erythrocyte to inhibit parasite egress. Sci Rep 7:12250.

30. Beck JR, Muralidharan V, Oksman A, Goldberg DE. 2014. PTEX component
HSP101 mediates export of diverse malaria effectors into host erythrocytes. Nature
511:592-5.
31. Glushakova S, Beck JR, Garten M, Busse BL, Nasamu AS, Tenkova-Heuser T,
Heuser J, Goldberg DE, Zimmerberg J. 2018. Rounding precedes rupture and
breakdown of vacuolar membranes minutes before malaria parasite egress from
erythrocytes. Cell Microbiol 20:e12868.
32. Nessel T, Beck JM, Rayatpisheh S, Jami-Alahmadi Y, Wohlschlegel JA, Goldberg
DE, Beck JR. 2020. EXP1 is required for organisation of EXP2 in the
intraerythrocytic malaria parasite vacuole. Cell Microbiol 22:e13168.
33. Luu W, Gelissen IC, Brown AJ. 2017. Manipulating Cholesterol Status Within Cells.
Methods Mol Biol 1583:41-52.
34. Collins CR, Hackett F, Atid J, Tan MSY, Blackman MJ. 2017. The Plasmodium
falciparum pseudoprotease SERA5 regulates the kinetics and efficiency of malaria
parasite egress from host erythrocytes. PLoS Pathog 13:e1006453.
556

561 **Figures**

Figure 1: Generation of the NF54 Rhoph2/EXP2 line : (A) Strategy for tagging *rhoph2* with

mNeongreen (mNG) and gel image indicating presence of clonal population expressing

rhoph2-mNG. (B) Strategy for the tagging *exp2 with mRuby tag* and agarose gel image

indicating the presence of clonal population expressing *exp2-mRuby.* Sequences of the

diagnostic primers A,B,C, and D are provided in the Materials and Methods. (C) Live

fluorescence microscopy images indicating the expected localization of the tagged proteins.

Red arrows indicate the dual localization of RhopH2-mNG on the RBC membrane as well as

the parasite in the rings and trophozoites while in the schizonts the protein is localized to

rhoptries. The EXP2-mRuby marks the PVM of the parasite. Erythrocyte membrane is stained

with WGA-Alexa-350 (Blue)

Figure 2: Trophozoites extrude out from erythrocyte upon treatment with MßCD.

(A) Still frames from Supplementary video 1 from time-lapse video microscopy performed at.

Parasites extrude out from the erythrocyte with the PVM still tethered to the erythrocyte

membrane without causing the lysis of erythrocyte (indicated by arrows). (B) Representative

images (>50) of PfVP1- mNG line after MßCD treatment. Nucleus is stained with SYTO-deep

red (Pink). (C) Representative images (>50) of NF54 RhopH2/Exp2 line after MßCD

treatment. Erythrocyte membrane is stained with WGA-Alexa-350 (Blue). Infected

erythrocytes were subjected to live fluorescent microscopy after treatment with MßCD.

Parasites extrude out from erythrocyte without causing lysis of the erythrocyte with their PPM

(B) and PVM (C) still attached to the erythrocyte as indicated by the arrows. (D) Representative

images of erythrocytes infected with ring stages of the NF54 RhopH2/Exp2 line treated with

MßCD. Ring stage parasites do not extrude out from erythrocytes. (E) Quantification of

parasites remaining inside/outside of erythrocytes after MßCD treatment was carried out by

examining at least 100 different images of individual parasites for each experimental condition

(100 for ring stages and 330 for trophozoites). (F) Upper panel: Treatment of NF54

RhopH2/Exp2 trophozoites with 0.5 µM Cyt-D for 45 mins prior to MßCD treatment does not

inhibit parasite extrusion. Lower panel: Treatment with 0.5 µM Cyt-D alone does not cause

extrusion of trophozoites but did alter PVM morphology (as indicated by arrows). (G)

(Left)Treatment with MßCD causes extrusion of *P. falciparum* stage trophozoites without

causing erythrocyte rupture unlike the control group where the trophozoite is still inside the

erythrocyte (Scale bar- 2µM, Direct magnification-5,000x). (Right) Zoomed in image of

parasite extruded out of erythrocyte after MßCD treatment shows the PVM ruptured at multiple

places indicated by black arrows. Electron microscopy of extruded *P. falciparum*. Left panels

show lower magnification views of control and MβCD-treated trophozoites. A higher

magnification view of an extruded parasite (right panel) shows that the PVM is compromised

at multiple places (arrows) (Scale bar-500nM, Direct magnification-12,000x).

Figure 3 : Treatment with PfATP4 or PfNCR1 inhibitors prior to MßCD treatment inhibited

MßCD mediated extrusion of parasites: (A) Trophozoite stage parasites from PfVP1-mNG line

were treated with KAE609 (10nM), MMV009108 (1µM) and chloroquine (150nM) prior to

treatment with MßCD. Live fluorescent imaging showed that parasite extrusion was inhibited

by prior treatment with KAE609 and MMV009108 (indicated by yellow arrows) while

treatment with chloroquine like MßCD did not inhibit this extrusion (indicated by white

arrows). Nucleus is stained with SYTO-deep red (pink). Erythrocyte membrane is stained with

WGA-Alexa-350 (Blue). Representative images from 3 independent experiments for each

condition (B) Quantification from live fluorescent microscopy experiments of parasite

extrusion upon drug treatment followed by MßCD treatment. Approximately 250 cells for each

657   experimental condition over 4 independent experiment were assessed. (C) Giemsa stained thin

658   blood smears prepared from cultures after treatment with KAE609 (10nM), MMV009108

659   (1µM) and chloroquine (150nM) prior to treatment with MßCD. (D) Quantitation of parasite

660   extrusion upon drug treatment followed by MßCD treatment from Giemsa-stained thin blodd

661   smears (~1000 cells were counted).

Figure 4: Cholesterol in the erythrocyte membrane is essential for growth and proliferation of *P. falciparum*.(A) Structure of MßCD, Cholesterol, Epicholesterol and Desmosterol. The differences between the structure of sterols are highlighted in red. (B) Giemsa-stained thin blood smears prepared from control, MßCD and erythrocytes reconstituted with cholesterol, desmosterol or epicholesterol at 24 and 48 hours after treatment. C) Parasitemia in 1:10 split

of ring stage parasites from control, MßCD (CD) or erythrocytes reconstituted with cholesterol

(CD/Cho), desmosterol(CD/Des) or epicholesterol (CD/Epi) over 4 days (~3000 cells were

counted). Based on one tailed unpaired *t-test* (* $p<0.05$). (D) Relative (^{3}H)-hypoxanthine

incorporation by ring stage parasites over a 24 hour period following treatment with MßCD or

reconstituted with cholesterol, desmosterol and epicholesterol. Based on unpaired t-test with

Welch's correction (* $p<0.05$, *** $p<0.001$). Error bars=mean±SD N=8. (E) TEM of the ring

stage erythrocyte treated with 5mM MßCD and imaged after 24 or 48 hours. (Left)After 24

hours, in erythrocyte treated with MßCD, parasites are unable to form normal trophozoites. The

nuclear membrane is indistinct, hemozoin is dispersed (indicated by arrows) and food vacuole

is small compared to control erythrocytes. (Right) After 48 hours, parasites in the control

culture were able to invade the erythrocyte and form rings while the parasites in the MßCD

treated erythrocytes undergo did not progress further and appear to disintegrate (Scale bar-

500nM, Direct magnification-15,000x).

Figure 5: Reducing the cholesterol content in the erythrocyte membrane inhibits *P. falciparum* invasion

(A) Giemsa stained thin blood smears from control and MßCD treated erythrocytes after 24 hours. While ring stages were seen in the control erythrocytes, no invasion was seen in the MßCD treated erythrocytes but merozoites appeared to remain attached to the surface as indicated by arrows. (B) Percoll enriched trophozoites were added to erythrocytes treated with MßCD or reconstituted with cholesterol and giemsa stained thin blood smears were prepared at 24 and 48 hours. Reconstituted erythrocytes were able to support parasite growth. (C) Parasitemia from 1:10 split of culture from MßCD treated and reconstituted with cholesterol in normal erythrocytes over a period of 4 days (~3000 cells were counted). Based on Mann-Whitney test (* p<0.05) (D) Percoll synchronized trophozoites from NF54 and Dd2 strains were added onto MßCD treated erythrocytes. Parasitemia from 1:10 split of these cultures by examining Giemsa-stained thin blood smears (~1000 cells were counted).

Supplementary Figure Legends

Supplementary Figure 1: Images of extruded parasites following MßCD treatment of of the NF54 RhopH2/Exp2 transgenic line.

Representative images (out of >50) of live NF54 RhopH2/Exp2 line after MßCD treatment.

Erythrocyte membrane is stained with WGA-Alexa-350 (Blue), EXP2 is red and RhoH2 is

green. Infected erythrocytes were subjected to live fluorescent microscopy after treatment with

MßCD. Scale bar is 2 μm.

Supplementary figure 2: Phase contrast images of late stage *P. falciparum* extruded after

treatment with MßCD.

Infected erythrocytes were subjected to phase contrast microscopy after treatment with

MßCD. Extruded parasites are indicated by red arrows.

Supplementary figure 3: Phalloidin staining of MßCD treated cultures from PfVP1-NG

transgenic line: (A) Representative images of uninfected and infected cells from untreated

cultures after staining with Phalloidin AlexaFluor 594. (B) Representative images of

uninfected and infected cells from MßCD treated cultures followed by staining with

Phalloidin AlexaFluor 594. (Scale bar- 5 μm). Overall, 60-70% of uninfected as well as

infected erythrocyte cytoskeletons were stained with phalloidin, indicating membrane breach

without the lysis of erythrocytes.

Supplementary figure 4: Transmission electron microscopy of the ring stage erythrocyte

treated with 5mM MßCD and imaged after 24 (Day 1) or 48 hours (Day2)

(A) After 24 hours, the trophozoite like parasites in the MßCD treated cultures (lower panel)

have indistinct membranes, compromised food vacuole and dispersed hemozoin compared to

the controls (Upper panel) (B) After 48 hours, in the control culture (Upper panel) the

parasites invaded and formed rings while the parasites in the MßCD treated cultures (Lower

panel) failed to form mature merozoites (Scale bar-500nM, Direct magnification-15,000x).

Supplementary Videos Legends

Supplementary video 1: Treatment with MßCD causes extrusion of trophozoites from

erythrocytes

Representative time lapse video microscopy of the NF54 RhopH2/Exp2 line after MßCD

treatment. Erythrocyte membrane is stained with WGA-Alexa-350 (Blue). Infected

erythrocytes were subjected to live fluorescent microscopy after treatment with MßCD.

Parasites extrude out from erythrocyte without causing lysis of the erythrocyte. Video was

acquired using the Nikon Ti microscope. The interval between each frame was 4 sec and the

video was captured for a total 20 min (93 frames, FPS:0.09, frames 59-64 shown in the

video). Images were deconvoluted by Richardson Lucy algorithm using the Nikon NIS

elements software package and converted to .avi files.

Supplementary video 2: Cyt-D treatment does not inhibit MßCD mediated parasite

extrusion

Representative time lapse video microscopy of the NF54 RhopH2/Exp2 line after MßCD

treatment. Erythrocyte membrane is stained with WGA-Alexa-350 (Blue). Infected

erythrocytes were subjected to live fluorescent microscopy after treatment with Cyt-D

followed by treatment with MßCD. Treatment with Cyt-D does not inhibit MßCD mediated

extrusion of the parasite. Video microscopy was performed as mentioned above for video 1.

The interval between each frame was 31 sec and the video was captured for a total 5 min (11

frames, FPS:0.04, all 11 frames shown in the video)

Supplementary movie 3: Treatment with MßCD does not cause extrusion of ring stage

parasites

Erythrocytes infected with ring stages of the NF54 RhopH2/Exp2 were treated with MßCD .

Erythrocyte membrane is stained with WGA-Alexa-350 (Blue). Ring stage parasites do not

extrude out after MßCD treatment. Video microscopy was performed as mentioned for video

1. The interval between each frame was 31 secs and the video was captured for a total 5 mins

(11 frames, FPS:0.04, all 11 frames shown in the video)

Supplementary Figure 1: Images of extruded parasites following MßCD treatment of of the NF54 RhopH2/Exp2 transgenic line.

Representative images (out of >50) of live NF54 RhopH2/Exp2 line after MßCD treatment. Erythrocyte membrane is stained with WGA-Alexa-350 (Blue), EXP2 is red and RhoH2 is green. Infected erythrocytes were subjected to live fluorescent microscopy after treatment with MßCD. Scale bar is 2 μm.

Supplementary figure 2: Phase contrast images of late stage *P. falciparum* extruded after treatment with MßCD.

Infected erythrocytes were subjected to phase contrast microscopy after treatment with MßCD. Extruded parasites are indicated by red arrows.

Supplementary figure 3: Phalloidin staining of MßCD treated cultures from PfVP1-NG transgenic line: (A) Representative images of uninfected and infected cells from untreated cultures after staining with Phalloidin AlexaFluor 594. (B) Representative images of uninfected and infected cells from MßCD treated cultures followed by staining with Phalloidin AlexaFluor 594. (Scale bar- 5 µm). Overall, 60-70% of uninfected as well as infected erythrocyte cytoskeletons were stained with phalloidin, indicating membrane breach without the lysis of erythrocytes.

Supplementary figure 4: Transmission electron microscopy of the ring stage erythrocyte treated with 5mM MßCD and imaged after 24 (Day 1) or 48 hours (Day2)
(A) After 24 hours, the trophozoite like parasites in the MßCD treated cultures (lower panel) have indistinct membranes, compromised food vacuole and dispersed hemozoin compared to the controls (Upper panel) (B) After 48 hours, in the control culture (Upper panel) the parasites invaded and formed rings while the parasites in the MßCD treated cultures (Lower panel) failed to form mature merozoites (Scale bar-500nM, Direct magnification-15,000x).