Arginine methylation sites on SepIVA help balance elongation and septation of

the cell wall in *Mycobacterium smegmatis*

By Angela H Freeman[1], Karen Tembiwa[1], James R Brenner[2], Michael R Chase[3], Sarah

M Fortune[3], Yasu S Morita[2], Cara C Boutte[1*]

[1]Department of Biology, University of Texas at Arlington, Arlington, Texas, USA

[2]Department of Microbiology, University of Massachusetts, Amherst, Massachusetts,

USA

[3]Department of Immunology and Infectious Disease, Harvard TH Chan School of Public

Health, Boston, Massachusetts, USA

[*]corresponding author, cara.boutte@uta.edu

<u>Abstract</u>

Growth of mycobacterial cells requires successful coordination between elongation and

septation of the cell wall. However, it is not clear which factors directly mediate this

coordination. Here, we studied the function and post-translational modification of an

essential division factor, SepIVA, in *Mycobacterium smegmatis*. We find that SepIVA is

arginine methylated, and that alteration of these methylation sites affects both septation

and polar elongation of *Msmeg.* Furthermore, we show that SepIVA regulates the

localization of MurG, and that this regulation may impact polar elongation. Finally, we

map SepIVA's two regulatory functions to different sites on the protein: the N-terminus

regulates elongation while the C-terminus regulates division. These results establish

SepIVA as a regulator of both elongation and division and characterize a physiological

role for protein arginine methylation sites for the first time in mycobacteria.

Introduction

The cell cycle of rod-shaped bacteria requires coordination of elongation and

division. Elongation involves incorporation of newly synthesized cell wall components,

including peptidoglycan (PG) into the existing cell wall. In the well-studied species *B.*

subtilis and *E. coli*, this incorporation occurs along the lateral cell walls and is directed

by the cytoskeletal factor MreB (Egan *et al.*, 2020). However, some rod-shaped

bacteria, including many Actinobacteria and some Alpha-proteobacteria, lack MreB-like

proteins and elongate by incorporating new cell wall near the cell poles (Thanky *et al.*,

2007; Aldridge *et al.*, 2012; Brown *et al.*, 2012; Cameron *et al.*, 2015; Zupan *et al.*,

2019; Zupan *et al.*, 2021). Polar elongation in mycobacteria, which includes the

pathogen *Mycobacterium tuberculosis*, requires the DivIVA protein Wag31 (Kang, 2005;

Kang *et al.*, 2008; Melzer *et al.*, 2018), through mechanisms that are currently unclear.

Many essential cell division factors are conserved from *E. coli* to mycobacteria (Hett

and Rubin, 2008; Wu *et al.*, 2018). However, because of the extra layers of the cell wall

(Jankute *et al.*, 2015), and polar growth (Thanky *et al.*, 2007) in mycobacteria, we

expect that some aspects of cell division must be different. In mycobacteria, polar

elongation and septation are spatially coincident but temporally separated; while in

lateral growers, elongation and septation are spatially distinct. Because division creates

the site of polar growth for the next generation of mycobacterial cells, we expect there to

be proteins involved in transitioning the new pole from the septal/divisive mode cell wall

metabolism (orthogonal to the long axis of the cell) to the elongative mode of cell wall

metabolism (parallel to the long axis of the cell) - though such factors have not been

described.

The intracellular membrane domain (IMD) is likely shared by the elongation and

division processes. The IMD is a membranous hub for generation of lipid-linked cell wall

precursors and it localizes to sites of cell wall synthesis: the subpolar region and septum

(Hayashi *et al.*, 2016). Cell wall precursors assembled in the IMD are thought to be

translocated into the plasma membrane associated with the cell wall (PM-CW) before

assembly into the cell wall (García-Heredia *et al.*, 2021). How precursor enzymes

associate with the IMD, and how that affects their activity is unknown.

SepIVA is essential for division and interacts with the conserved divisome protein,

FtsQ (Wu *et al.*, 2018; Jain *et al.*, 2018). *sepIVA* has also been implicated as a

determinant of beta-lactam susceptibility, indicating that it may affect peptidoglycan

metabolism (Flores *et al.*, 2005). Like Wag31, SepIVA comprises a DivIVA domain and

aligns to DivIVA from *B. subtillis* (Fig. S1). DivIVA homologs are often involved in

recruitment of other proteins to cell poles or septa (Marston and Errington, 1999;

Halbedel and Lewis, 2019; Hammond *et al.*, 2019; Hammond *et al.*, 2021). However,

none of the confirmed interactors of DivIVA-domain proteins interact with residues that

are conserved in SepIVA (Fig. S1). Depletion of Wag31 and SepIVA have opposite

effects: depletion of SepIVA leads to continued elongation and defective division (Wu *et*

al., 2018), while depletion of Wag31 leads to elongation arrest but continued division

(Kang *et al.*, 2008).

Here, we show that SepIVA is arginine methylated. Arginine methylation is known to

be involved in various cell functions in eukaryotes, such as membrane association,

protein transport, DNA binding and protein-protein interactions (Biggar and Li, 2015;

Raposo and Piller, 2018). Protein arginine methylation has been identified using mass

spectrometry in *E. coli* (Zhang *et al.*, 2018), mycobacteria (Sakatos *et al.*, 2018) and

Salmonella (Su *et al.*, 2021). Arginine methylation sites have been shown *in vitro* to

affect DNA binding of transcription factors in *Salmonella* (Su *et al.*, 2021) and

mycobacteria (Singhal *et al.*, 2020). However, the physiological roles of protein arginine

methylation have barely been studied in bacteria. Currently, there are no characterized

protein arginine methyltransferases in mycobacteria. Recent work has suggested that

the protein glutamate methyltransferase CheR in *Salmonella enterica* can somehow

also methylate arginines (Su *et al.*, 2021); there is no homolog with a predicted CheR

methyltransferase domain in mycobacteria.

In this work, we show that methyl-ablative and methyl-mimetic mutations at

certain arginine methylation sites on SepIVA regulate SepIVA's localization,

peptidoglycan metabolism, and cell elongation and division. We also show that SepIVA

is involved in regulating the localization of peptidoglycan precursor enzyme MurG, and

that arginine methylation site mutations of *sepIVA* affect MurG's localization to the

poles. This work establishes SepIVA as a regulator of both elongation and division and

indicates that it may have a role in regulating MurG. In addition, we show here that

SepIVA is arginine methylated, and that methyl-mimetic mutations at the N-terminus of

SepIVA disrupt MurG's polar localization and slow polar growth, while methyl-ablative

site mutations promote polar growth. We also find that methylation sites near the C-

terminus of SepIVA are required for normal cell division.

<u>Results</u>

Methylation site mutations on SepIVA affect growth and cell length of Msmeg

Post-translational modifications are critical for regulating the physiology of *Mtb*

and other mycobacteria (Kang *et al.*, 2008; Prisic *et al.*, 2010; Garces *et al.*, 2010; Prisic

and Husson, 2014; Boutte *et al.*, 2016; Sakatos *et al.*, 2018; Iswahyudi *et al.*, 2019;

Shamma *et al.*, 2020). Most studies thus far have focused on phosphorylation. To

identify proteins that could be regulated by other types of post-translational modification,

we re-analyzed a mass spectrometry data set of *Mtb* peptide masses that we had

previously published (Garces *et al.*, 2010) to search for post translational modifications.

We found evidence of four arginine methylations on SepIVA (Rv2927c), as well as on

many other proteins from *Mtb* (Supplemental table 1). The four methylated arginines

from *Mtb* SepIVA were R19, R105, R111, R199 and all are conserved in SepIVA from

Msmeg.

To see if SepIVA in *M. smegmatis* is also arginine methylated, we

immunoprecipitated SepIVA-strep from *Msmeg* lysates and used LCMS-IT-TOF mass

spectrometry to measure the post-translational modifications. We found evidence of

widespread arginine methylation on SepIVA's 24 arginine residues (Supplemental table

2).

To identify arginine residues in SepIVA for which methylation might have a

physiological role, we used L5 allele swapping to make several mutants of *sepIVA* in

Msmeg (Pashley and Parish, 2003). First, we integrated a wild-type *sepIVA* at the L5

site on a nourseothricin-marked vector and deleted the endogenous copy of *sepIVA.*

Then, we cloned various *sepIVA* arginine mutants into kanamycin-marked vectors, and

transformed these into the Δ*sepIVA* L5::*sepIVA* WT strain, and selected for allele

swaps. Because there were so many arginines with potential methylation sites, we

made mutants with several clustered arginines mutated at once. We grouped arginines

based on their location on the SepIVA protein, as predicted by AlphaFold (Jumper *et al.*,

2021) (Pettersen *et al.*, 2021) (Fig. S2). We made R->M methionine mutants to

represent methyl-mimetic mutations, as methionine is the closest structural mimic to a

methylated arginine (Singhal *et al.*, 2020). We made R->A alanine mutations to test the

function of the positive charge of arginines and R->K lysine mutations to serve as

methyl-ablatives that retain the positive charge on the residue. We found that several

groups of arginines were essential for *sepIVA* function, as the alanine mutant strains did

not grow (R->A mutants, Fig. 1A) in liquid culture.

Our analysis of clustered arginine mutants of *sepIVA* (Fig. 1AB) identified two

sets of mutants with interesting phenotypes. If the lysine and methionine mutants mimic

the different arginine methylation states, and if that methylation has a physiological role,

then we expect to see opposite phenotypes in the methyl-ablative (R->K) and methyl-

mimetic (R->M) strains compared to the wild-type. The *sepIVA* R3K, R19K, R31K

mutant, hereafter called the NT-K mutant, as these are the most N-terminal arginines,

exhibited slightly faster growth, and shorter cells than the wild-type, while the NT-M

mutant (*sepIVA* R3M, R19M, R31M) had slower growth (Fig. 1AB). The mutant *sepIVA*

R224K, R228K, R230K, R234K, R245K, hereafter called the CT-K mutant, had slow

growth and (Fig. 1A) very short cell length (Fig. 1B). The C-terminal methionine mutant

(*sepIVA* R224M, R228M, R230M, R234M, R245M) was not viable, likely because the

protein was unstable (Fig. 1C).

To examine whether individual arginine residues were responsible for these

growth and cell length defects, we generated *Msmeg* strains with single methylation site

mutations in the N-terminal and C-terminal arginines. We found that single methylation

site mutations near the N-terminus of *sepIVA* did not have the same effects on growth

and cell length as the triple mutant (Fig. 1DE). Since more than one methylation site

mutant near the N-terminus was needed for these phenotypes to be apparent, we used

the NT-K and NT-M triple mutants to study the potential role of arginine methylation on

SepIVA's N-terminus (Fig. 2,3,4,6). The single methylation site mutations on the C-

terminus of SepIVA had different phenotypes from the quintuple CT-K mutant (Fig.

1ABDE). Both the *sepIVA* R230K and R230M exhibited slow growth and long cells (Fig.

1DE): this indicates that these mutations likely do not represent different methylation

states and are instead both loss-of-function mutations. Another methylation site, R234,

only showed growth defects in the methionine mutant (Fig. 1DE). Since the phenotypes

of the C-terminus single site mutants were distinct from the phenotype of the CT-K

quintuple mutant, we believe the short cell length seen in the CT-K mutant was a result

of suppressor mutations.

We made all the mutants described above with C-terminal strep tags so we could

test for protein stability using western blotting. (Fig. 1CF). To test the stability of mutant

proteins that did not support growth in allele swap strains (Fig. 1A – missing bars), we

transformed the vectors carrying the mutant *sepIVA*-strep into wild-type *Msmeg* with the

wild-type *sepIVA* allele at its native locus to support growth. Using these merodiploid

strains, we found that SepIVA-strep alanine mutant proteins that did not support growth

were unstable (Fig. 1C, '*' indicates merodiploid strain). Several of the mutants had

significantly higher levels of SepIVA than the wild-type, including the NT-K, NT-M and

CT-K mutants (Fig. 1C) which exhibited phenotypes. To test whether high levels of

these mutant proteins might be responsible for the slow growth, we tested how

overexpression of the CT-K mutant affected growth rates (Fig. S3). We found that

overexpression of SepIVA CT-K did not slow growth in both WT *sepIVA* and *sepIVA*

CT-K genetic backgrounds. This result shows that the increased protein levels are not

responsible for the slow growth and short cell length of the *sepIVA* CT-K mutant. Other

variants of SepIVA-strep are stable; though protein levels between mutants vary (Fig.

1CF), these variations do not correlate with phenotypes (Fig. 1ABDE).

These data suggest that methylation sites on SepIVA appear to play a role in

how SepIVA regulates growth and division. Near the N-terminus, methyl-mimetic site

mutations cause slow growth and lead to slightly shorter cells (Fig. 1). However, near

the C-terminus of SepIVA, both R230 mutants are defective in division. At R234 the

methyl-mimetic mutation results in slow growing, long cells, while the methyl-ablative

mutation results in cells like the WT. For the remainder of this work, we focus on the N-

terminal triple mutants and the single site mutants of R234.

According to TnSeq experiments and our previous work, *sepIVA* is an essential

gene (DeJesus *et al.*, 2017; Wu *et al.*, 2018; Dragset *et al.*, 2019). In another recent

study, a mutant of *sepIVA* missing only residues 156-245, the C-terminus, was

characterized and the authors described cell division defects (Pickford *et al.*, 2020). We

generated this C-terminus truncation mutant using L5 allele swapping and were able to

generate colonies on plates. However, once in liquid culture, these mutants exhibited

highly variable or no growth (Fig. S4). We interpret this to mean that the *sepIVA* C-

terminus is largely essential, but that suppressor mutants can form easily to allow

growth in its absence.

Methylation site mutations on SepIVA affect its polar localization

We sought to determine if alterations at the arginine methylation sites could

affect SepIVA's localization. SepIVA normally localizes at the poles and septum, with

stronger localization to the old pole, and spotty localization along the lateral walls in a

pattern (Wu *et al.*, 2018) that is seen in proteins that associate with the Intracellular

Membrane Domain (IMD) (Meniche *et al.*, 2014; Hayashi *et al.*, 2016). To test this, we

made merodiploid strains of *Msmeg* with GFPmut3-SepIVA methyl-mutant alleles

expressed from the L5 site. Strains carrying GFPmut3-*sepIVA* as the only allele of

sepIVA were nonviable, so merodiploid strains were used. We grew these strains in log.

phase, and stained the cells with the fluorescent D-amino acid HADA, which

preferentially labels sites of new peptidoglycan synthesis, and allows us to identify the

faster growing, old pole as the pole that stains more brightly (Kuru *et al.*, 2012;

Baranowski *et al.*, 2018). Upon imaging the cells, we found that methylation site

mutations on SepIVA affect its distribution and total intensity (Fig. 2). Specifically, both

NT-K and NT-M mutants showed inhibited localization of SepIVA to the fast-growing

pole (Fig. 2AC), and increased localization to the cell septum. This suggests that

regulation of these methylation sites affects SepIVA distribution throughout the cell.

Methylation site mutants also exhibit differences in GFPmut3-SepIVA intensity broadly

(Fig. 2B), however, we cannot conclude whether mean GFPmut3-SepIVA signal is an

indicator of protein levels within the cell. Our findings suggest that SepIVA's localization

to the cell poles is affected by changes in the methylation sites on the N-terminus of

SepIVA. We cannot discount the possibility that GFPmut3-SepIVA localizes differently

from the untagged protein, but these results show at least that these mutations can

affect the distribution of the fusion in a merodiploid background.

Methylation site mutations on SepIVA affect peptidoglycan metabolism

SepIVA's essentiality for cell division and sequence similarity to DivIVA proteins

suggests that it regulates the cell wall. To determine if the methylation sites on SepIVA

impact metabolism of cell wall components, we stained the allele swap *sepIVA-strep*

arginine-mutant strains with NADA (Kuru *et al.*, 2015), which is incorporated into

metabolically active peptidoglycan mostly by periplasmic amino acid exchange by L,D-

transpeptidases (LDTs) (Kuru *et al.*, 2012; Baranowski *et al.*, 2018). We found that the

N-terminal methyl-mimetic site mutant, NT-M, showed slightly decreased NADA signal

at the cell poles compared to the wild-type (Fig. 3). We also found that the NT-K mutant

showed increased, and the NT-M mutant showed decreased, overall NADA

incorporation compared to the wild-type (Fig. 3B). To observe differences in distribution

of NADA, we looked at NADA incorporation across the length of the cell. Based on

normalized polar NADA intensities, there is less NADA incorporation at both poles in the

NT-M mutant (Fig. 3CD). These data suggest SepIVA affects peptidoglycan

metabolism, though they do not address whether these effects are direct or indirect.

SepIVA had previously been shown to be essential for division (Wu *et al.*, 2018), but

these differences in polar NADA incorporation indicate that SepIVA may also be a

regulator of polar peptidoglycan metabolism.

Methylation site mutations on SepIVA affect polar elongation

To determine if the changes in NADA staining in the *sepIVA* methyl-mutants

were due to alterations in LDT-mediated peptidoglycan remodeling (Baranowski *et al.*,

2018) or due to insertion of new cell wall at the poles, we measured polar elongation in

the *sepIVA* wild-type and mutant strains. We developed a method of measuring

elongation using a pulse of HADA (blue), 1.5 hours of outgrowth in media, and another

pulse of NADA (green) stain. We then measured the distance of the NADA-stained but

HADA-clear poles, which represents the polar growth during the 1.5-hour outgrowth

(Fig. 4A).

First, we used this method to determine if depletion of SepIVA affects cell

elongation. SepIVA depletion leads to a slight increase in net polar elongation (Fig. 4B).

Because cell division is inhibited in the SepIVA-depletion condition, the increase in net

polar elongation could be due to the increased age of the poles in non-dividing cells:

older poles in mycobacteria grow faster (Aldridge *et al.*, 2012), so cells that are not

creating new poles through division will have older poles overall. From these data we

conclude that SepIVA is not essential for elongation.

To determine if methylation sites on SepIVA could affect elongation, we

performed pulse-chase-pulse staining on *sepIVA-strep* wild-type and methylation site

mutant allele swap strains. We observed increased elongation in the NT-K methyl-

ablative mutants, while the NT-M methyl-mimetic mutant showed decreased polar

elongation (Fig. 4C). Since our methyl-ablative and methyl-mimetic site mutations show

opposing effects on polar elongation, this suggests that methylation of SepIVA at these

sites regulates polar elongation. We also performed this elongation assay on single

methylation site mutants (Fig. S5). We found that individual N-terminal arginines had a

similar trend - with the lysine mutants exhibiting greater elongation than the methionine

mutants – but none of the single mutants had a statistically significant effect. From

these results, we conclude that methylation at the R3, R19 and R31 residues work

together in regulating elongation. These results show that SepIVA is an elongation

regulator; since it is not required for elongation (Fig. 4B) we suggest that its function

may be to help toggle growth between the poles and septum.

SepIVA affects MurG localization

The Intracellular Membrane Domain (IMD) is a membrane domain that can be

separated by sucrose density centrifugation from the cytoplasmic fractions and from the

PM-CW (plasma membrane attached to the cell wall) fraction, which contains PBPs and

other periplasmic cell wall enzymes (Hayashi *et al.*, 2016). Microscopy from our

previous study showed that GFPmut3-SepIVA has a similar localization pattern as the

IMD-associated arabinogalactan precursor enzyme GlfT2 (Wu *et al.*, 2018). To

determine if SepIVA is associated with the IMD, we performed sucrose gradient

fractionation on an *Msmeg* strain with a C-terminal strep tag on SepIVA (Fig. 5A). We

found that SepIVA was found in the cytoplasmic fractions 1-2, as well as fraction 3.

GlfT2 was previously shown to be found restricted to the IMD in fractions 4-6 (Hayashi

et al., 2018). MurG, the final cytoplasmic enzyme in PG precursor synthesis, which also

has a typical IMD-localization pattern (Meniche *et al.*, 2014; García-Heredia *et al.*,

2021), is found partially in the IMD and partially in the cytoplasmic fractions (García-

Heredia *et al.*, 2021). Thus, SepIVA and MurG have some overlap in their density-

gradient distribution. These results show that SepIVA is not associated strongly with the

IMD; however, the localization pattern of GFPmut3-SepIVA indicates that it may

associate indirectly or weakly.

Because other DivIVA homologs are involved in recruiting proteins to their site of

activity (Marston and Errington, 1999; Halbedel and Lewis, 2019), we hypothesized that

perhaps SepIVA could be involved in recruiting proteins to its locations near the IMD

and at the septum. To test this, we looked at the effects of SepIVA depletion on

localization of GlfT2-mRFP and MurG-Dendra2. We built an *Msmeg* strain where the

only copy of *sepIVA* has a C-terminal DAS degradation tag, which targets a protein to

be proteolyzed by ClpP when the SspB adaptor is expressed by adding an Atc inducer

(Kim *et al.*, 2011; Wu *et al.*, 2018). We transformed this depletion strain with either of

the localization constructs. We then added Atc to induce the degradation of SepIVA-

DAS, or did not add Atc in the control, and imaged the cells. We found that depletion of

SepIVA had mild effects on GlfT2-mRFP localization (Fig. 5B-D). However, we found

that MurG-Dendra2 localization was highly sensitive to SepIVA levels. In the strain with

SepIVA-DAS and no *sspB*, MurG-Dendra2 had a localization pattern similar to the

normal IMD pattern that has been described before (Meniche *et al.*, 2014; Hayashi *et*

al., 2016; García-Heredia *et al.*, 2021) (Fig. 5EH). When we transformed the *sspB*

vector into this strain but did not induce *sspB* expression, we expect a moderate

decrease in SepIVA levels due to leaky SspB expression, and we see complete

delocalization of MurG-Dendra2 (Fig. 5GH). When we induce SspB to strongly deplete

SepIVA-DAS, MurG-Dendra2 localizes in puncta that are randomly distributed around

the cell (Fig. 5GH). We observed moderate changes in mean intensity of GlfT2-mRFP,

and large changes in MurG-Dendra2 intensity between the SepIVA depleted and wild-

type strains (Fig. 5CG); we do not know the significance of these changes, and here

focus instead on the changes in distribution (Fig. 5H).

Because SepIVA has different effects on GlfT2-RFP and MurG-Dendra2, we

conclude that SepIVA does not affect IMD structure globally and is not required for IMD-

association of GlfT2. Our results indicate, however, that SepIVA could have a role

modulating MurG's localization and/ or association with the membrane.

Methylation site mutants on SepIVA affect polar localization of MurG.

We determined that depletion of SepIVA affects localization of MurG more strongly

than GlfT2 (Fig. 5). To determine whether arginine methylation site mutations in *sepIVA*

affect MurG or GlfT2 localization, we imaged each fluorescent protein fusion in *sepIVA*

wild-type and methyl-mutant strains allele swap strains. We found that the distribution of

GlfT2-mRFP around the cell was unaffected by methylation site mutations (Fig.

6ABCD). However, MurG-Dendra2 localization was less polar and more septal in the

methyl-mimetic NT-M mutant (Fig. 6EFGH) compared to the methyl-ablative mutant and

the WT. This suggests that an unmethylated N-terminus of SepIVA may stimulate

MurG's association with the growing pole (Fig 6GH), possibly by recruiting it to that site

(Fig. 2) to promote polar PG synthesis and polar elongation (Fig. 3,4). SepIVA with a

methylated N-terminus may instead decrease MurG's association to the poles (Fig.

6GH), which results in decreased polar peptidoglycan metabolism (Fig. 3CD) and

elongation (Fig. 4C) resulting in slower growth (Fig. 1A). We are observing a correlation

between MurG polar association (Fig. 6GH) and polar growth (Fig. 4C) in these strains,

which could be due to SepIVA directly or indirectly regulating MurG.

To test if arginine methylation site mutants of *sepIVA* affect the FtsZ ring, we

expressed *ftsZ-mcherry2b* in methylation mutants. Our results show that arginine

methylation of SepIVA does not affect FtsZ-mcherry2B localization (Fig. S6).

Residues near the C-terminus of SepIVA function in cell division

Our preliminary work on individual arginine mutations at the C-terminus of

sepIVA indicated that these residues affect cell division (Fig. 1). We further

characterized the *sepIVA* R234 methylation site mutants. We observed the methyl-

ablative methylation site mutant, R234K, is like the WT in growth rate and cell length,

while the methyl-mimetic R234M has slower growth and longer cell length compared to

the WT (Fig. 7ABC). The *sepIVA* R234M cells also elongate slightly more than the WT

(Fig. 7D), though this is likely to be from the increased age of poles in cells with

inhibited cell division, as in Fig. 4B. Thus, the *sepIVA* R234M strain displays inhibited

cell division. The R234M strain has higher net peptidoglycan metabolism, as measured

by NADA staining, than the WT, while the R234K has lower peptidoglycan metabolism

than the WT (Fig. 7E). Because the two R234 mutants have opposing phenotypes, they

may mimic the methylated and unmethylated states of SepIVA at R234. The data from

these mutants suggests that arginine methylation at R234 inhibits division, while the

unmethylated divides like the WT, suggesting that in the conditions measured, SepIVA

is likely not methylated at this residue. The *sepIVA* R234 mutant strains both have

increased MurG-Dendra2 fluorescent signal, but comparable distributions to the wild-

type. Here, MurG signal does not correlate with difference in cell wall metabolism or

growth (Fig. 7BCDGH), suggesting that regulation of cell division through the C-

terminus of SepIVA does not involve regulating MurG localization. Our data show that

the R234 methylation site is crucial for SepIVA's role in cell division.

<u>Discussion</u>

Although SepIVA is a DivIVA protein, the preliminary data on this factor indicate that

its function is distinct from other described DivIVA proteins. All other described DivIVA

proteins localize specifically to the curved membranes at poles and septa (Ramamurthi

and Losick, 2009). SepIVA has some localization to the septum, but it mostly localizes

in a ring around the cell near the pole, but not at the pole tip (Wu *et al.*, 2018) (Fig. 2).

This shows that SepIVA is not associated with primarily with curved membranes like

other DivIVA proteins. Other described DivIVA proteins have been shown to have

functions in promoting polar elongation (Kang *et al.*, 2008; Letek *et al.*, 2008; Hempel *et*

al., 2008; Melzer *et al.*, 2018), helping to inhibit cell division (Marston and Errington,

1999; Eswara *et al.*, 2018) and/or to bind and regulate transcription factors

(Eswaramoorthy *et al.*, 2014), cell wall enzymes (Cleverley *et al.*, 2019), cell division

regulators (Eswara *et al.*, 2018), and an envelope saccharide transporter (Hammond *et*

al., 2022). None of these DivIVA proteins are essential for cell division, as SepIVA is

(Wu *et al.*, 2018). While some individual residues on DivIVA homologs have been

characterized as being involved in these functions, none of these residues are

conserved in SepIVA (Fig. S1). The low conservation between SepIVA and other

DivIVA proteins, the difference in localization, and the wide variety of functions and

binding partners across species make it impossible to use homology to predict the

molecular functionality of SepIVA.

Mycobacteria must regulate peptidoglycan precursor synthesis in coordination

with cell wall expansion and remodeling needs. Our data indicate that SepIVA likely

helps regulate the peptidoglycan layer (Fig. 3) possibly partly through control of MurG

(Fig. 5,6), which is the final enzyme in the construction of lipidII before it is flipped into

the periplasm (Gee *et al.*, 2012; Egan *et al.*, 2020). In *E. coli*, MurG interacts with both

elongation and division factors (Laddomada *et al.*, 2016; Egan *et al.*, 2020). According

to our data, mild SepIVA depletion results in dispersion of MurG, and strong SepIVA

depletion results in mislocalization of MurG (Fig. 5G-J). We speculate that SepIVA is

somehow affecting MurG's association with the Intracellular Membrane Domain (IMD)

(Hayashi *et al.*, 2016; García-Heredia *et al.*, 2021) or other regulators that help cell wall

metabolism transition between septal and elongative modes – our data do not indicate

whether this regulation is direct or indirect. A third possibility is that SepIVA depletion is

causing the type of cell wall damage that has previously been shown to cause MurG to

re-localize to the lateral walls (Melzer *et al.*, 2021). The arginine methylation sites on the

N-terminus of SepIVA appear to toggle its effect on MurG localization in a way that

correlates with polar elongation: the strain with the *sepIVA* N-terminal methyl-mimetic

mutations have less polar growth (Fig. 4C) and less MurG at the poles (Fig. 6F). If

SepIVA's role in elongation regulation is to regulate MurG's association and dispersal

from the IMD domain at the poles - directly or indirectly - then it appears that when

SepIVA is methylated at the N-terminus, it is more active in dispersing MurG from the

poles.

Arginine methylation sites near the C-terminus of SepIVA, specifically R234,

seem to play a role in regulating cell division (Fig. 7). The methyl-mimetic R234M

mutant showed inhibited division (Fig. 7) while the methyl-ablative, R234K, did not.

Methylation at this arginine residue could be inhibiting cell division. These effects on cell

division revealed through mutations at the C-terminus appear to be independent of

MurG localization (Fig. 7G).

Studies on protein arginine methylation done in eukaryotes support its

importance in cell cycle progression (Biggar and Li, 2015; Raposo and Piller, 2018). Our

data establish that protein arginine methylation sites on SepIVA affect the mycobacterial

cell cycle as well. We have found that the arginine methylation sites on SepIVA affect

both the elongation and division stages of the cell cycle. This work establishes protein

arginine methylation as a physiologically important post-translational modification in

bacteria and identifies SepIVA as a potential mediator between the elongation and

division processes.

Methods and Materials

Bacterial strains and culture conditions.

Mycobacterium smegmatis (mc^{2}155) was cultured in 7H9 medium (Becton,

Dickinson, Franklin Lakes, NJ) with 5 g/L bovine serum albumin (BSA), 2 g/L glucose,

0.85 g/L NaCl, 0.003 g/L catalase, 0.2% glycerol, and 0.05% Tween 80 or plated on LB

agar. For *M. smeg* cultures, antibiotic concentrations were: 20 µg/mL nourseothricin, 20

µg/mL zeocin, 20 µg/mL streptomycin, 25 µg/mL kanamycin, 50 µg/mL hygromycin, and

30 µg/mL apramycin. For *E. coli* cultures, antibiotic concentrations were: 40 µg/mL

nourseothricin, 25 µg/mL zeocin, 50 µg/mL streptomycin, 50 µg/mL kanamycin, 100

µg/mL hygromycin, and 50 µg/mL apramycin *E. coli* TOP10 or DH5α were used for

cloning.

Strain construction

Because it is essential in *M. smegmatis, sepIVA* mutants were constructed by

first integrating an additional copy of *sepIVA* at the L5 integration site. After

complementing *sepIVA* at the L5 site, *sepIVA* at its native locus was knocked out via

recombineering (van Kessel and Hatfull, 2008), as previously described (Boutte *et al.*,

2016). Mutant copies of *sepIVA* replaced wildtype *sepIVA* at the L5 site via allelic

exchange (Pashley and Parish, 2003; Kieser *et al.*, 2015).

sepIVA depletion strains were constructed as previously described (Wu *et al.*,

2018). Merodiploid expression constructs of GlfT2-mRFP, FtsZ-mcherry2B, and MurG-

Dendra2 were integrated at the Tweety integrase site (Pham *et al.*, n.d.). The

merodiploid construct expressing GFPmut3-SepIVA was integrated at the L5 integrase

site. All strains used are listed in Table S3 of the supplementary material, along with

plasmids listed in Table S4 and primers listed in Table S5.

Growth rate assay

Biological replicates of *sepIVA* arginine mutants were grown to log phase in 7H9

medium. Cultures were diluted to $OD_{600} = 0.1$ in 200 µL 7H9 medium in a non-treated

96-well plate. Optical density at 600 nm was measured over the next 18-24 hours using

a plate reader (BioTek Synergy neo2 multi-mode reader) at 37°C, shaking continuously.

Population doubling times were determined using the exponential growth equation and

least squares regression fitting method in GraphPad Prism (version 9.2). P values were

calculated using ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a

single pooled variance.

Western blots

Log phase cultures (10 mL) were pelleted and resuspended in 500 µL PBS + 1

mM phenylmethylsulfonyl fluoride (PMSF). Cells were lysed using glass beads

(MiniBeadBeater-16, model 607, Biospec). Supernatants from cell lysates were run on

4–12% NuPAGE Bis Tris precast gels (Life Technologies, Beverley, MA) using MES

running buffer (Life Technologies, Beverly, MA). Proteins were transferred onto

polyvinylidene difluoride (PVDF) membranes (GE Healthcare). Strep-tagged proteins

were detected using rabbit anti-Strep-tag II antibody (1:1000, Abcam, ab76949) in Tris-

buffered saline with Tween 20 (TBST) with 0.5% BSA and goat anti-rabbit IgG (H+L)

horseradish peroxide (HRP)-conjugated secondary antibody (1:10,000, Thermo Fisher

Scientific 31460) in TBST. Tagged proteins were visualized using chemiluminescent

reagents (Thermo Fisher Scientific 34579).

Microscopy

Cells were imaged using a Nikon Ti-2 widefield epifluorescence microscope with

a Photometrics Prime 95B camera and a Plan Apo 100X, 1.45 -numerical-aperture

objective. Green-fluorescent images for GFPmut3 or Dendra2 localization or NADA

staining were taken with a 470/40nm excitation filter and a 525/50nm emission filter.

Blue-fluorescent images for HADA staining were taken with a 350/50nm excitation filter

and a 460/50nm emission filter. Red-fluorescent images for mRFP and mcherry2B

localization were taken with a 560/40nm excitation filter and a 630/70 emission filter.

Images were captured using NIS Elements software. Images were analyzed using FIJI

and MicrobeJ (Ducret *et al.*, 2016). For cell detection using MicrobeJ, parameters for

width and area were set. V-snapping cells were cut at the septum so daughter cells

could be detected as individual cells. Overlapping cells were excluded from analysis.

Cell lengths, mean intensities, maximum intensities, and minimum intensities of

at least 250 cells per genotype were quantified using MicrobeJ. Mean intensity profiles

were plotted using the "XStatProfile" plotting tool in MicrobeJ. P-values were calculated

using ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a single

pooled variance using GraphPad Prism (v9.2).

Localization of cell wall proteins upon SepIVA depletion.

For depletion of SepIVA, log phase cells (10 mL) were incubated (37°C) with 500

ng/mL Anhydrotetracycline (ATc) for 7-9 hours. For induction of GlfT2-mRFP, 1 ng/mL

IVN were added, and cells were incubated (37°C) for 5 hours. Cells were fixed for

microscopy. Cells were fixed for microscopy with 1.6% paraformaldehyde for 10

minutes and then washed and resuspended in PBS tween80. Microscopy images were

analyzed in ImageJ and MicrobeJ. P-values calculated by unpaired t-test using

GraphPad Prism (v9.2).

Fluorescent staining

HADA (ThermoFisher) and NADA (ThermoFisher) fluorescent D-alanine amino

acids were used to stain the peptidoglycan cell wall layer (Kuru *et al.*, 2012). For PG

analysis, NADA stain was added to 1 mL exponentially growing cells at a final

concentration of 1 ug/mL for 3 minutes at room temperature before washing twice in

PBST. For pulse-chase experiments, HADA (1 ug/mL) was added for 15 minutes before

cells were washed twice in 7H9, incubated (37°C) for 1.5 hours, and stained again using

NADA (1 ug/mL) for 3 minutes at room temperature. Cells were washed twice in PBST

and imaged. Images were analyzed in ImageJ (NIH). The length of the poles and the

length of the entire cell were measured manually for each cell.

Sucrose Density Gradient Fractionation

Sucrose density gradient fractionation was performed as previously described

(1{Citation}5). In brief, cells were grown to log phase and lysed by nitrogen cavitation.

Lysate was then placed on top of a 20-50% sucrose gradient and centrifuged at 35,000

rpm for 6 hours at 4°C. Twelve one-mL fractions were collected and analyzed.

Immunoprecipitation and mass spectrometry

We previously performed LCQ-MS/MS to identify peptides from *Mtb* cell lysates

(Garces *et al.*, 2010). Here, we re-analyzed those data to search for modifications in

peptides, using Andromeda software.

SepIVA-strep was immunoprecipitated from CB1223. Cells were grown to late

log. phase, pelleted and resuspended in PBS with PMSF at 1 mM, lysed by bead

beating, and clarified by centrifugation. Immunoprecipitation was performed with

MagStrep type 3 XT beads (Iba Life Sciences, Gottingen, Germany) according to the

manufacturers protocol. Liquid was evaporated from the immunoprecipitants, and then

digested by Trypsin/Lys-C Mix, Mass Spec Grade protease (Promega), according to the

manufacturer's protocols for reduction, alkylation and two-step in-solution digestion of

proteolytically resistant proteins. The digested peptide solution was then cleaned by

C18-Ziptips (Sigma Aldrich) according to the manufacturers protocol.

LC-MS analysis was performed on LTQ Velos pro mass spectrometer (Thermo

Scientific, CA, USA) combined with a UHPLC (UltiMate 3000, Dionex, USA). The

digested peptides were separated by a nano viper analytical C18 column. (Acclaim

pepMap, 150 mm × 75µm, 2µm, 100 Å, Thermo Scientific, CA, USA). A 60 min gradient

method was used to separate the digested peptides (0–3 min 4.0%B, 3–50 min 4.0–

50.0% B, 50–50.1 min 50–90% B,50.1–55 min 90% B, 55–55.1 min 90–4% B, 55.1–60

min 4% B; mobile phase A: 0.1% FA in water; mobile phase B: 0.1% FA in 95%

acetonitrile, 5% water). 5µL of digested samples was injected at 300 nL/min flow. The

nano-electrospray ionization (ESI) source used with a fixed spray voltage at 2.4 kV and

a heated capillary temperature at 275 °C. A full MS spectrum obtained in normal scan

mode from 350 to 2000 m/z mass range. The data-dependent acquisition was

performed to get the MS/MS spectra for the 5 most abundant ions. The software

Proteome Discoverer (Thermo Scientific, USA) was used to search and identify the

peptides. LC-MS was performed at the Shimadzu Center for Advanced Analytical

Chemistry at UT Arlington.

Acknowledgements

This work was funded by grant R15GM131317 from the NIH. We thank Heather

Lake who contributed to optimization of some experiments, and Arash Emami Saleh

who aided in data processing.

Aldridge, B.B., Fernandez-Suarez, M., Heller, D., Ambravaneswaran, V., Irimia, D., Toner, M.,
and Fortune, S.M. (2012) Asymmetry and Aging of Mycobacterial Cells Lead to Variable
Growth and Antibiotic Susceptibility. *Science* **335**: 100–104.

Baranowski, C., Welsh, M.A., Sham, L.-T., Eskandarian, H.A., Lim, H.C., Kieser, K.J., *et al.*
(2018) Maturing Mycobacterium smegmatis peptidoglycan requires non-canonical crosslinks to
maintain shape. *eLife* **7**: e37516.

Biggar, K.K., and Li, S.S.-C. (2015) Non-histone protein methylation as a regulator of cellular
signalling and function. *Nat Rev Mol Cell Biol* **16**: 5–17.

Boutte, C.C., Baer, C.E., Papavinasasundaram, K., Liu, W., Chase, M.R., Meniche, X., *et al.*
(2016) A cytoplasmic peptidoglycan amidase homologue controls mycobacterial cell wall
synthesis. *eLife* **5** https://elifesciences.org/articles/14590. Accessed February 6, 2019.

Brown, P.J.B., Pedro, M.A. de, Kysela, D.T., Van der Henst, C., Kim, J., De Bolle, X., *et al.*
(2012) Polar growth in the Alphaproteobacterial order Rhizobiales. *Proceedings of the National*
Academy of Sciences **109**: 1697–1701.

Cameron, T.A., Zupan, J.R., and Zambryski, P.C. (2015) The essential features and modes of
bacterial polar growth. *Trends in Microbiology* **23**: 347–353.

Cleverley, R.M., Rutter, Z.J., Rismondo, J., Corona, F., Tsui, H.-C.T., Alatawi, F.A., *et al.*
(2019) The cell cycle regulator GpsB functions as cytosolic adaptor for multiple cell wall
enzymes. *Nat Commun* **10**: 261.

DeJesus, M.A., Gerrick, E.R., Xu, W., Park, S.W., Long, J.E., Boutte, C.C., *et al.* (2017)
Comprehensive Essentiality Analysis of the *Mycobacterium tuberculosis* Genome via Saturating
Transposon Mutagenesis. *mBio* **8** http://mbio.asm.org/lookup/doi/10.1128/mBio.02133-16.
Accessed February 6, 2019.

Dragset, M.S., Ioerger, T.R., Zhang, Y.J., Mærk, M., Ginbot, Z., Sacchettini, J.C., *et al.* (2019)
Genome-wide Phenotypic Profiling Identifies and Categorizes Genes Required for
Mycobacterial Low Iron Fitness. *Sci Rep* **9**: 11394.

Ducret, A., Quardokus, E.M., and Brun, Y.V. (2016) MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. *Nature Microbiology* **1**: 16077.

Egan, A.J.F., Errington, J., and Vollmer, W. (2020) Regulation of peptidoglycan synthesis and
remodelling. *Nat Rev Microbiol* http://www.nature.com/articles/s41579-020-0366-3. Accessed
May 18, 2020.

Eswara, P.J., Brzozowski, R.S., Viola, M.G., Graham, G., Spanoudis, C., Trebino, C., *et al.*
(2018) An essential Staphylococcus aureus cell division protein directly regulates FtsZ
dynamics. *eLife* **7** https://elifesciences.org/articles/38856. Accessed February 6, 2019.

Eswaramoorthy, P., Winter, P.W., Wawrzusin, P., York, A.G., Shroff, H., and Ramamurthi, K.S.
(2014) Asymmetric Division and Differential Gene Expression during a Bacterial Developmental
Program Requires DivIVA. *PLoS Genet* **10**: e1004526.

Flores, A.R., Parsons, L.M., and Pavelka, M.S. (2005) Characterization of Novel Mycobacterium
tuberculosis and Mycobacterium smegmatis Mutants Hypersusceptible to -Lactam Antibiotics.
Journal of Bacteriology **187**: 1892–1900.

Garces, A., Atmakuri, K., Chase, M.R., Woodworth, J.S., Krastins, B., Rothchild, A.C., *et al.*
(2010) EspA Acts as a Critical Mediator of ESX1-Dependent Virulence in Mycobacterium
tuberculosis by Affecting Bacterial Cell Wall Integrity. *PLOS Pathogens* **6**: e1000957.

García-Heredia, A., Kado, T., Sein, C.E., Puffal, J., Osman, S.H., Judd, J., *et al.* (2021)
Membrane-partitioned cell wall synthesis in mycobacteria. *eLife* **10**: e60263.

Gee, C.L., Papavinasasundaram, K.G., Blair, S.R., Baer, C.E., Falick, A.M., King, D.S., *et al.*
(2012) A Phosphorylated Pseudokinase Complex Controls Cell Wall Synthesis in Mycobacteria.
Sci Signal **5**: ra7–ra7.

Halbedel, S., and Lewis, R.J. (2019) Structural basis for interaction of DivIVA/GpsB proteins
with their ligands. *Molecular Microbiology* **111**: 1404–1415.

Hammond, L.R., Khan, S.J., Sacco, M.D., Spanoudis, C., Hough, A., Chen, Y., and Eswara, P.J.
(2021) GpsB coordinates cell division and cell surface decoration by wall teichoic acids in
Staphylococcus aureus. Microbiology, .
http://biorxiv.org/lookup/doi/10.1101/2021.09.29.462461. Accessed October 4, 2021.

Hammond, L.R., Sacco, M.D., Khan, S.J., Spanoudis, C., Hough-Neidig, A., Chen, Y., and
Eswara, P.J. (2022) GpsB Coordinates Cell Division and Cell Surface Decoration by Wall
Teichoic Acids in Staphylococcus aureus. *Microbiol Spectr* **10**: e01413-22.

Hammond, L.R., White, M.L., and Eswara, P.J. (2019) ¡vIVA la DivIVA! *J Bacteriol* JB.00245-
19, jb;JB.00245-19v1.

Hayashi, J.M., Luo, C.-Y., Mayfield, J.A., Hsu, T., Fukuda, T., Walfield, A.L., *et al.* (2016)
Spatially distinct and metabolically active membrane domain in mycobacteria. *PNAS* **113**: 5400–
5405.

Hayashi, J.M., Richardson, K., Melzer, E.S., Sandler, S.J., Aldridge, B.B., Siegrist, M.S., and
Morita, Y.S. (2018) Stress-Induced Reorganization of the Mycobacterial Membrane Domain.
mBio **9** http://mbio.asm.org/lookup/doi/10.1128/mBio.01823-17. Accessed February 6, 2019.

Hempel, A.M., Wang, S. -b., Letek, M., Gil, J.A., and Flardh, K. (2008) Assemblies of DivIVA
Mark Sites for Hyphal Branching and Can Establish New Zones of Cell Wall Growth in
Streptomyces coelicolor. *Journal of Bacteriology* **190**: 7579–7583.

Hett, E.C., and Rubin, E.J. (2008) Bacterial Growth and Cell Division: a Mycobacterial
Perspective. *Microbiology and Molecular Biology Reviews* **72**: 126–156.

Iswahyudi, Mukamolova, G.V., Straatman-Iwanowska, A.A., Allcock, N., Ajuh, P., Turapov, O.,
and O'Hare, H.M. (2019) Mycobacterial phosphatase PstP regulates global serine threonine
phosphorylation and cell division. *Sci Rep* **9**: 8337.

Jain, P., Malakar, B., Khan, M.Z., Lochab, S., Singh, A., and Nandicoori, V.K. (2018)
Delineating FtsQ-mediated regulation of cell division in *Mycobacterium tuberculosis*. *Journal of*
Biological Chemistry **293**: 12331–12349.

Jankute, M., Cox, J.A.G., Harrison, J., and Besra, G.S. (2015) Assembly of the Mycobacterial
Cell Wall. *Annual Review of Microbiology* **69**: 405–423.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., *et al.* (2021) Highly
accurate protein structure prediction with AlphaFold. *Nature*
http://www.nature.com/articles/s41586-021-03819-2. Accessed August 2, 2021.

Kang, C.-M. (2005) The Mycobacterium tuberculosis serine/threonine kinases PknA and PknB:
substrate identification and regulation of cell shape. *Genes & Development* **19**: 1692–1704.

Kang, C.-M., Nyayapathy, S., Lee, J.-Y., Suh, J.-W., and Husson, R.N. (2008) Wag31, a
homologue of the cell division protein DivIVA, regulates growth, morphology and polar cell
wall synthesis in mycobacteria. *Microbiology* **154**: 725–735.

Kessel, J.C. van, and Hatfull, G.F. (2008) Mycobacterial Recombineering. *Methods in Molecular*
Biology (Clifton, NJ) **435**: 203–215.

Kieser, K.J., Boutte, C.C., Kester, J.C., Baer, C.E., Barczak, A.K., Meniche, X., *et al.* (2015)
Phosphorylation of the Peptidoglycan Synthase PonA1 Governs the Rate of Polar Elongation in
Mycobacteria. *PLOS Pathogens* **11**: e1005010.

Kim, J.-H., Wei, J.-R., Wallach, J.B., Robbins, R.S., Rubin, E.J., and Schnappinger, D. (2011)
Protein inactivation in mycobacteria by controlled proteolysis and its application to deplete the
beta subunit of RNA polymerase. *Nucleic Acids Res* **39**: 2210–2220.

Kuru, E., Hughes, H.V., Brown, P.J., Hall, E., Tekkam, S., Cava, F., *et al.* (2012) In Situ Probing
of Newly Synthesized Peptidoglycan in Live Bacteria with Fluorescent D -Amino Acids.
Angewandte Chemie International Edition **51**: 12519–12523.

Kuru, E., Tekkam, S., Hall, E., Brun, Y.V., and Van Nieuwenhze, M.S. (2015) Synthesis of
fluorescent D-amino acids and their use for probing peptidoglycan synthesis and bacterial growth
in situ. *Nat Protoc* **10**: 33–52.

Laddomada, F., Miyachiro, M.M., and Dessen, A. (2016) Structural Insights into Protein-Protein
Interactions Involved in Bacterial Cell Wall Biogenesis. *Antibiotics (Basel)* **5**: 14.

Letek, M., Ordonez, E., Vaquera, J., Margolin, W., Flardh, K., Mateos, L.M., and Gil, J.A.
(2008) DivIVA Is Required for Polar Growth in the MreB-Lacking Rod-Shaped Actinomycete
Corynebacterium glutamicum. *Journal of Bacteriology* **190**: 3283–3292.

Marston, A.L., and Errington, J. (1999) Selection of the midcell division site in Bacillus subtilis through MinD-dependent polar localization and activation of MinC. *Molecular Microbiology* **33**: 84–96.

Melzer, E.S., Kado, T., García-Heredia, A., Gupta, K.R., Meniche, X., Morita, Y.S., *et al.* (2021) Cell wall damage reveals spatial flexibility in peptidoglycan synthesis and a non-redundant role for RodA in mycobacteria. Microbiology, . http://biorxiv.org/lookup/doi/10.1101/2021.10.26.465981. Accessed November 8, 2021.

Melzer, E.S., Sein, C.E., Chambers, J.J., and Sloan Siegrist, M. (2018) DivIVA concentrates mycobacterial cell envelope assembly for initiation and stabilization of polar growth. *Cytoskeleton* **75**: 498–507.

Meniche, X., Otten, R., Siegrist, M.S., Baer, C.E., Murphy, K.C., Bertozzi, C.R., and Sassetti, C.M. (2014) Subpolar addition of new cell wall is directed by DivIVA in mycobacteria. *Proceedings of the National Academy of Sciences* **111**: E3243–E3251.

Pashley, C.A., and Parish, T. (2003) Efficient switching of mycobacteriophage L5-based integrating plasmids in *Mycobacterium tuberculosis*. *FEMS Microbiology Letters* **229**: 211–215.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll, T.I., *et al.* (2021) UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci* **30**: 70–82.

Pham, T.T., Jacobs-Sera, D., Pedulla, M.L., Hendrix, R.W., and Hatfull, G.F.Y. 2007 Comparative genomic analysis of mycobacteriophage Tweety: evolutionary insights and construction of compatible site-specific integration vectors for mycobacteria. *Microbiology* **153**: 2711–2723.

Pickford, H., Alcock, E., Singh, A., Kelemen, G., and Bhatt, A. (2020) A mycobacterial DivIVA domain-containing protein involved in cell length and septation. *Microbiology* https://www.microbiologyresearch.org/content/journal/micro/10.1099/mic.0.000952. Accessed July 18, 2020.

Prisic, S., Dankwa, S., Schwartz, D., Chou, M.F., Locasale, J.W., Kang, C.-M., *et al.* (2010) Extensive phosphorylation with overlapping specificity by Mycobacterium tuberculosis serine/threonine protein kinases. *Proceedings of the National Academy of Sciences* **107**: 7521–7526.

Prisic, S., and Husson, R.N. (2014) Mycobacterium tuberculosis Serine/Threonine Protein Kinases. *Microbiology Spectrum* **2** http://www.asmscience.org/content/journal/microbiolspec/10.1128/microbiolspec.MGM2-0006-2013. Accessed February 6, 2019.

Ramamurthi, K.S., and Losick, R. (2009) Negative membrane curvature as a cue for subcellular localization of a bacterial protein. *Proceedings of the National Academy of Sciences* **106**: 13541–13545.

Raposo, A.E., and Piller, S.C. (2018) Protein arginine methylation: an emerging regulator of the
cell cycle. *Cell Div* **13**: 3.

Sakatos, A., Babunovic, G.H., Chase, M.R., Dills, A., Leszyk, J., Rosebrock, T., *et al.* (2018)
Posttranslational modification of a histone-like protein regulates phenotypic resistance to
isoniazid in mycobacteria. *Science Advances* **4**: eaao1478.

Shamma, F., Papavinasasundaram, K., Quintanilla, S.Y., Bandekar, A., Sassetti, C., and Boutte,
C.C. (2020) Phosphorylation on PstP Regulates Cell Wall Metabolism and Antibiotic Tolerance
in *Mycobacterium smegmatis*. *J Bacteriol* **203**: e00563-20, /jb/203/4/JB.00563-20.atom.

Singhal, A., Virmani, R., Naz, S., Arora, G., Gaur, M., Kundu, P., *et al.* (2020) Methylation of
two-component response regulator MtrA in mycobacteria negatively modulates its DNA binding
and transcriptional activation. *Biochemical Journal* **477**: 4473–4489.

Su, Y., Li, J., Zhang, W., Ni, J., Huang, R., Wang, Z., *et al.* (2021) Methylation of PhoP by
CheR Regulates *Salmonella* Virulence. *mBio* https://journals.asm.org/doi/10.1128/mBio.02099-
21. Accessed September 28, 2021.

Thanky, N.R., Young, D.B., and Robertson, B.D. (2007) Unusual features of the cell cycle in
mycobacteria: Polar-restricted growth and the snapping-model of cell division. *Tuberculosis* **87**:
231–236.

Wu, K.J., Zhang, J., Baranowski, C., Leung, V., Rego, E.H., Morita, Y.S., *et al.* (2018)
Characterization of Conserved and Novel Septal Factors in *Mycobacterium smegmatis*. *Journal*
of Bacteriology **200**: e00649-17, /jb/200/6/e00649-17.atom.

Zhang, M., Xu, J.-Y., Hu, H., Ye, B.-C., and Tan, M. (2018) Systematic Proteomic Analysis of
Protein Methylation in Prokaryotes and Eukaryotes Revealed Distinct Substrate Specificity.
PROTEOMICS **18**: 1700300.

Zupan, J., Guo, Z., Biddle, T., and Zambryski, P. (2021) *Agrobacterium tumefaciens* Growth
Pole Ring Protein: C Terminus and Internal Apolipoprotein Homologous Domains Are Essential
for Function and Subcellular Localization. *mBio* **12**: e00764-21, /mbio/12/3/mBio.00764-
21.atom.

Zupan, J.R., Grangeon, R., Robalino-Espinosa, J.S., Garnica, N., and Zambryski, P. (2019)
GROWTH POLE RING protein forms a 200-nm-diameter ring structure essential for polar
growth and rod shape in Agrobacterium tumefaciens. *PNAS* 201905900.

Figure 1. Methylation site mutations on SepIVA affect growth and cell length of
Msmeg. **(A)** Doubling time of allele-swap *M. smeg* cells expressing WT or arginine
methylation site mutants of SepIVA. WT SepIVA represents the L5 complemented

strain. Lighter color outlined bars represent arginine to alanine mutants. No outline,

darker color bars represent methyl-ablative methylation site mutants. Black outline bars

represent methyl-mimetic methylation site mutants. Strains without bars were non-

viable. At least 3 biological replicates of each strain were used. Error bars represent

standard deviation. ****, $P = <0.0001$. **(B)** Cell lengths of SepIVA WT or arginine

methylation site mutants. Bars represent mean. At least 300 cells over three biological

replicates of each strain were analyzed. *, $P = 0.0439$; ****, $P = <0.0001$. **(C)** α-strep

western blot of L5::*sepIVA*-strep WT or arginine methylation site mutants. mc^{2}155

without a strep tag was used as a negative control (lane 2). The non-specific band

serves as a loading control. '*' denotes merodiploid strains. Merodiploid strains were

used for SepIVA variants did not support growth as allele swaps. **(D)** Doubling time of

allele-swap *M. smeg* cells expressing WT or single arginine methylation site mutants of

SepIVA. WT SepIVA is the strain with WT *sepIVA*-strep at the L5 site and *sepIVA*

deleted from its native locus – this is the isogenic control. No outline, darker color bars

represent methyl-ablative methylation site mutants. Black outline bars represent methyl-

mimetic methylation site mutants. Strains without bars were non-viable. At least 3

biological replicates of each strain were used. Error bars represent standard deviation.

****, $P = <0.0001$. **(E)** Cell lengths of SepIVA WT or single arginine methylation site

mutants. Bars represent mean. At least 300 cells over three biological replicates of each

strain were analyzed. *, $P = 0.0351$; ****, $P = <0.0001$. **(F)** α-strep western blot of

L5::*sepIVA*-strep WT or single arginine methylation site mutants. mc^{2}155 without a strep

tag was used as a negative control (lane 2). The non-specific band serves as a loading

control. P-values were calculated using ordinary one-way ANOVA, Dunnett's multiple

comparisons test, with a single pooled variance in GraphPad Prism (v9.2). At least two

western blots were performed for each strain shown, and representative results are

displayed.

Figure 2. Methylation site mutations affect polar localization of SepIVA. (A) Images
of *Msmeg* cells expressing L5::*GFPmut3-sepIVA* WT and methylation site mutants.
Strains are merodiploid. Phase images are on the top row. Fluorescence images of
GFPmut3-SepIVA are in the middle panel. HADA images are on the bottom row. Scale
bar applies to all images. Pictures of several cells from images processed identically
were pasted together. **(B)** Mean intensity values per cell of cells expressing
L5::*GFPmut3-sepIVA* WT and methylation mutants. At least 250+ cells across three
biological replicates of each mutant were analyzed in MicrobeJ. Bar represents mean
intensity value between cells. ****, *P = <0.0001.* P-values were calculated using
ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a single pooled
variance in GraphPad Prism (v9.2). **(C)** Normalized Intensity profiles of GFP-SepIVA

signal of WT and methylation site mutants. The solid line represents the mean intensity

value across the relative position of the cell. Normalized intensity values were

generated by dividing each position's intensity value by the mean intensity value. Cells

were pole-sorted such that the brighter pole in the HADA channel was set to 0 on the X

axis.

Figure 3. Methylation site mutations of SepIVA affect peptidoglycan metabolism.

(A) Images of M. smeg cells expressing *L5::sepIVA* WT and methylation site mutants.

Scale bar applies to all images. Pictures of several cells from images processed

identically were pasted together. **(B)** Mean NADA intensity values of *M. smeg* cells

expressing *L5::sepIVA* WT and methylation site mutants. At least 250+ cells across

three biological replicates of each mutant were analyzed in MicrobeJ. Bar represents

mean intensity value. *****, P = <0.0001.* P-values were calculated using ordinary one-

way ANOVA, Dunnett's multiple comparisons test, with a single pooled variance in

GraphPad Prism (v9.2). **(C)** Normalized intensity profiles of NADA signal of WT and

methylation mutants. The solid line represents the mean intensity value across the

relative position of the cell. Intensity values were normalized by dividing each strain's

intensity value by its mean intensity value. Cells were pole-sorted such that the brighter

pole in the NADA channel was set to 0 on the X axis. **(D)** Normalized polar NADA

intensities of WT and methylation site mutants. Intensity values were normalized by

dividing each cell's maximum polar intensity value by the mean intensity of the whole

cell. *****, P = <0.0001.* P-values were calculated using ordinary one-way ANOVA,

Dunnett's multiple comparisons test, with a single pooled variance in GraphPad Prism

(v9.2).

Figure 4. Methylation site mutations on SepIVA affect polar elongation. (A)
Schematic of pulse-chase-pulse staining method. We stained growing cells with blue
peptidoglycan dye HADA, then returned them to media for another 1.5 hours of growth,
then stained them again with green dye NADA. We false-colored the green signal as
red in these images to make the contrast clearer. **(B)** Length of poles after SepIVA
depletion. +SepIVA (black dots) represent cells expressing SepIVA, -SepIVA (black-
outlined dots) represent cells with depleted SepIVA protein. At least 100 cells across
three biological replicates were analyzed. Bar represents median. Scale bar applies to
all images. *****, P = <0.0001*. P-value calculated by unpaired t-test using GraphPad
Prism (v9.2). **(C)** Length of poles elongation after 1.5 hours in cells expressing *sepIVA*
WT and methylation site allele swap mutants. At least 100 cells across three biological
replicates were analyzed. Bar represents mean polar elongation. *, P = 0.0461, ****, P =
<0.0001.* P-values were calculated using ordinary one-way ANOVA, Dunnett's multiple
comparisons test, with a single pooled variance in GraphPad Prism (v9.2).

Figure 5. SepIVA has the same membrane association as MurG, and it affects

MurG localization. (A) anti-Strep western blot of sucrose gradient fractions. Fractions

4-6 contain intracellular membrane domain (IMD) proteins. The green bracket denotes

fractionation of MurG-Dendra2, as shown in Garcia-Heredia *et al.* 2021. **(B)**

Micrographs of cells expressing or depleting SepIVA and induced for GlfT2-mRFP.

Scale bar is 5 microns and applies to all images. Pictures of several cells from images

processed identically were pasted together. **(C)** Mean GlfT2-mRFP intensity values of

Msmeg cells expressing or depleting SepIVA and induced for GlfT2-mRFP. At least

250+ cells across three biological replicates of each mutant were analyzed in MicrobeJ.

Bar represents mean intensity value. ****, P = <0.0001.* P-value calculated by unpaired

t-test using GraphPad Prism (v9.2). **(D)** Normalized intensity profiles of GlfT2-mRFP

signal of cells expressing and depleting SepIVA and induced for GlfT2-mRFP. The solid

line represents the mean intensity value across the relative position of the cell. Cells

were pole-sorted such that the brighter pole in the HADA channel was set to 0 on the X

axis. **(E)** Micrographs of cells expressing *sepIVA::FLAG-das.* This strain lacks inducible

sspB, therefore, SepIVA cannot be depleted in this strain. We refer to this strain as

++SepIVA. These images are 12-bit. Scale bar is 5 microns and applies to all images.

Pictures of several cells from images processed identically were pasted together. **(F)**

Micrographs of cells expressing or depleting SepIVA while constitutively expressing

MurG-Dendra2. These images are 16-bit. Scale bar is 5 microns and applies to all

images. Pictures of several cells from images processed identically were pasted

together. **(G)** Mean MurG-Dendra2 intensity values of *Msmeg* cells expressing or

depleting SepIVA while constitutively expressing MurG-Dendra2. At least 250+ cells

across three biological replicates of each mutant were analyzed in MicrobeJ. Bar

represents mean intensity value. ****, P = <0.0001.* P-value calculated by unpaired t-test

using GraphPad Prism (v9.2). **(H)** Normalized intensity profiles of MurG-Dendra2 signal

of cells expressing and depleting SepIVA. The solid line represents the mean intensity

value across the relative position of the cell. Cells were pole-sorted such that the

brighter pole in the HADA channel was set to 0 on the X axis.

Figure 6. Methylation site mutations on SepIVA polar localization of MurG. (A)
Images of *Msmeg* cells expressing L5::*sepIVA* WT or methylation site mutants and
induced for GlfT2-mRFP. Scale bar is 5 microns and applies to all images. Pictures of
several cells from images processed identically were pasted together. **(B)** Mean GlfT2-
mRFP intensity values of *Msmeg* cells expressing *L5::sepIVA* WT or methylation site
mutants and induced for GlfT2-mRFP. At least 250+ cells across three biological

replicates of each mutant were analyzed in MicrobeJ. Bar represents mean intensity

value. ****, *P = <0.0001.* **(C)** Normalized intensity profiles of GlfT2-mRFP signal of cells

expressing *L5::sepIVA* WT or methylation site mutants and induced for GlfT2-mRFP.

The solid line represents the mean intensity value across the relative position of the cell.

Cells were pole-sorted such that the brighter pole in the HADA channel was set to 0 on

the X axis. **(D)** Normalized polar GlfT2-mRFP intensities of WT and methylation site

mutants. Intensity values were normalized by dividing each cell's maximum polar

intensity value by the mean intensity of the whole cell. *, *P = 0.0111.* **(E)** Images of

Msmeg cells expressing L5::*sepIVA* WT or methylation site mutants and constitutively

expressing MurG-Dendra2. Scale bar is 5 microns and applies to all images. Pictures of

several cells from images processed identically were pasted together. **(F)** Mean MurG-

Dendra2 intensity values of *Msmeg* cells expressing *L5::sepIVA* WT and methylation

site mutants. At least 250+ cells across three biological replicates of each mutant were

analyzed in MicrobeJ. Bar represents mean intensity value. ****, *P = <0.0001.* **(G)**

Normalized intensity profiles of MurG-Dendra2 signal of cells expressing *L5::sepIVA* WT

and methylation site mutants. The solid line represents the mean intensity value across

the relative position of the cell. Cells were pole-sorted such that the brighter pole in the

HADA channel was set to 0 on the X axis. **(H)** Normalized polar MurG-Dendra2

intensities of WT and methylation site mutants. Intensity values were normalized by

dividing each cell's maximum polar intensity value by the mean intensity of the whole

cell. ***, *P = 0.0002.* P-values were calculated using ordinary one-way ANOVA,

Dunnett's multiple comparisons test, with a single pooled variance in GraphPad Prism

(v9.2).

Figure 7. Methylation site R234 of SepIVA affects cell division.

(A) Images of *Msmeg* cells expressing L5::*sepIVA* WT and R234 methylation site

mutants. Scale bar is 5 microns and applies to all images. Pictures of several cells from

images processed identically were pasted together. **(B)** Doubling time of *Msmeg* cells

expressing WT or R234 methylation site mutants of SepIVA. No outline, darker color

bars represent methyl-ablative methylation site mutants. Black outline bars represent

methyl-mimetic methylation site mutants. At least 3 biological replicates of each strain

were used. Error bars represent standard deviation. **, *P = 0.0012.* **(C)** Cell lengths of

SepIVA WT or R234 methylation site mutants. Bars represent mean. At least 300 cells

over three biological replicates of each strain were analyzed. ****, *P = <0.0001.* **(D)** Net

polar elongation after 1.5 hours in cells expressing *L5::sepIVA* WT or R234 methylation

site mutants. Bar represents mean polar elongation. At least 100 cells across three

biological replicates were analyzed. **, *P = 0.0066.* **(E)** Mean NADA intensity values of

Msmeg cells expressing *L5::sepIVA* WT or R234 methylation site mutants. Bar

represents mean intensity value. At least 250+ cells across three biological replicates of

each mutant were analyzed in MicrobeJ. *P = <0.0001.* **(F)** Normalized intensity profiles

of NADA signal of cells expressing *L5::sepIVA* WT or R234 methylation site mutants.

The solid line represents the mean intensity value across the relative position of the cell.

Cells were pole-sorted such that the brighter pole in the NADA channel was set to 0 on

the X axis. **(G)** Mean MurG-Dendra2 intensity values of *Msmeg* cells expressing

L5::sepIVA WT or R234 methylation site mutants. Bar represents mean intensity value.

At least 250+ cells across three biological replicates of each mutant were analyzed in

MicrobeJ. ****, *P = <0.0001.* All P-values were calculated using ordinary one-way

ANOVA, Dunnett's multiple comparisons test, with a single pooled variance in

GraphPad Prism (v9.2). **(H)** Normalized intensity profiles of MurG-Dendra2 signal of

cells expressing *L5::sepIVA* WT or R234 methylation site mutants. The solid line

represents the mean intensity value across the relative position of the cell. Cells were

pole-sorted such that the brighter pole in the HADA channel was set to 0 on the X axis.

892
893