Arginine methylation helps SepIVA balance regulation of septation and

elongation in *Mycobacterium smegmatis*

By Angela H Freeman[1], Karen Tembiwa[1], James R Brenner[2], Michael R Chase[3], Sarah

M Fortune[3], Yasu Morita[2], Cara C Boutte[1*]

[1]Department of Biology, University of Texas at Arlington, Arlington, Texas, USA

[2]Department of Microbiology, University of Massachusetts, Amherst, Massachusetts,

USA

[3]Department of Immunology and Infectious Disease, Harvard TH Chan School of Public

Health, Boston, Massachusetts, USA

[*]corresponding author, cara.boutte@uta.edu

Abstract

Growth of mycobacterial cells requires successful coordination between

elongation and division of the cell wall. However, it is not clear which factors directly

mediate this coordination. Here, we studied the function and post-translational

modification of an essential division factor, SepIVA, in *Mycobacterium smegmatis*. We

find that SepIVA is arginine methylated, and that these modifications alter both division

and polar elongation of *Msmeg.* Furthermore, SepIVA impacts the localization of MurG.

Polar localization of MurG correlates with polar elongation in arginine methylation

mutants of *sepIVA*. These results establish SepIVA as a regulator of both elongation

and division, and characterize a physiological role for protein arginine methylation for

the first time in bacteria.

Introduction

The cell cycle of rod-shaped bacteria requires coordination of elongation and

division. Elongation involves incorporation of newly synthesized cell wall components,

such as peptidoglycan (PG) into the existing cell wall. In the well-studied species *B.*

subtilis and *E. coli*, this incorporation occurs along the lateral cell wall and is directed by

the cytoskeletal factor MreB (1). However, some rod-shaped bacteria, including many

Actinobacteria and some Alpha-proteobacteria, lack MreB-like proteins and elongate by

incorporating new cell wall at the cell poles (2–7). Polar elongation in mycobacteria,

which includes the pathogen *Mycobacterium tuberculosis*, requires the DivIVA protein

Wag31 (8–10). Wag31 directs subpolar PG synthesis from its localization site at the cell

pole (11) through mechanisms that are currently unclear.

Many essential cell division factors are conserved from *E. coli* to mycobacteria (12, 13).

However, because of the distinct features of mycobacteria, such as the extra layers of

the cell wall (14), and polar growth (2), we expect that some aspects of cell division

must be different as well. In mycobacteria, polar elongation and septation are spatially

coincident but temporally separated; while in lateral growers, elongation and septation

are spatially distinct. Because division creates the site of polar growth for the next

generation of mycobacterial cells, we expect there to be proteins involved in

transitioning between division and elongation, though such factors have not been

described.

One structure that is likely shared by the elongation and division processes is the

intracellular membrane domain (IMD). The IMD is a membranous hub for generation of

lipid-linked cell wall precursors and it localizes to sites of cell wall synthesis: the

subpolar region and septum (15). Precursors assembled in the IMD are thought to be

flipped into the plasma membrane associated with the cell wall (PM-CW) before

assembly into the cell wall (16). Depletion of Wag31 resulted in delocalization of IMD

markers and accumulation of peptidoglycan precursor lipid II in the IMD (16), suggesting

that Wag31 helps define and regulate the IMD. How precursor enzymes associate with

the IMD, and how that affects their activity is unknown.

SepIVA is essential for division and interacts with conserved divisome proteins (13,

17). *sepIVA* has also been implicated as a determinant of beta-lactam susceptibility,

indicating that it may affect peptidoglycan metabolism (18). Like Wag31, SepIVA

comprises a DivIVA domain. DivIVA homologs are often involved in recruitment of other

proteins to a site of interest (19–21). Depletion of Wag31 and SepIVA have opposite

effects: depletion of SepIVA leads to continued elongation and defective division (13),

while depletion of Wag31 leads to elongation arrest (8).

Here, we show that SepIVA is arginine methylated. Arginine methylation is known to

be involved in various cell functions in eukaryotes, such as membrane association,

protein transport, DNA binding and protein-protein interactions (22, 23). Protein arginine

methylation has been identified using mass spectrometry in *E. coli* (24) mycobacteria

(25) and *Salmonella* (26). Arginine methylation sites have been shown *in vitro* to affect

DNA binding of transcription factors in *Salmonella* (26) and mycobacteria (27).

However, protein arginine methylation has never been shown to affect a phenotype of

any bacteria.

In this work, we show that arginine methylation on SepIVA regulates SepIVA's

localization, PG metabolism, and cell elongation and division. The arginine methylations

also affect the localization of PG precursor enzyme MurG. This work establishes

SepIVA as a regulator of both elongation and division and demonstrates that arginine

methylation has roles beyond modulating DNA binding in bacteria and can critically

affect phenotypes such as cell growth and division.

Results

Arginine methylation of SepIVA affects growth and cell length of Msmeg

Post-translational modifications are critical for regulating the physiology of *Mtb*

and other mycobacteria (8, 25, 28–33). Most studies thus far have focused on

phosphorylation. To identify proteins that could be regulated by other types of post-

translational modification, we re-analyzed a mass spectrometry data set of *Mtb* proteins

that we had previously published (30) to search for post translational modification. We

found four arginine methylations on SepIVA (Rv2927c), as well as in many other

proteins from *Mtb* (Supplemental table 1).

To see if SepIVA in *M. smegmatis* is also arginine methylated, we

immunoprecipitated SepIVA-strep from *Msmeg* lysates and used LCMS-IT-TOF mass

spectrometry to measure the post-translational modifications. We found evidence of at

least some methylations on nearly every arginine in SepIVA (Supplemental table 2).

To determine if the arginine methylations on SepIVA have a physiological role,

we built strains of *Msmeg* in which the only copy of *sepIVA*, at the L5 site, is replaced

with mutants where groups of the arginines are replaced with either alanines, to probe

the function of the positive charges, or lysines, to probe the function of the methylations.

We found several sets of arginines in which the positive charges were essential (R->A

mutants, Fig. 1A). One mutant with arginine to lysine mutations showed that the

methylations at the C-terminus were important for normal growth (R->K mutant R224,

R228, R230, R234, R254, hereafter called the CT-K mutant) (Fig. 1A) and for normal

cell length (Fig. 1B). We were also interested in the arginines at the N-terminus, as the

R->K mutant (R3, R19, R31, hereafter called the NT-K mutant) exhibited slightly faster

growth than the wild-type, though it was not statistically significant. To further probe the

function of the arginine methylations at the C and N termini, we made arginine to

methionine mutations at these sites (R->M), as methionine serves as a methyl-mimetic.

We found the N-terminal methionine mutant (hereafter NT-M) had slower growth than

the wild-type and shorter cell length (Fig. 1AB). The C-terminal methionine mutant was

not stably viable in liquid culture, likely because the protein was unstable (Fig. 1C).

We made all the mutants described above with C-terminal strep tags so we could

test for protein stability. We found that the non-viable mutants (Fig. 1A) had unstable

SepIVA protein (Fig. 1C). Several of the mutants had significantly higher levels of

SepIVA than the wild-type, including the NT-K, NT-M and CT-K mutants (Fig. 1C) which

exhibited phenotypes. In order to test whether increased stability of these mutant

proteins might be responsible for the phenotypes, we tested how overexpression of the

CT-K mutant – which had the most pronounced phenotypes – affected growth rates

(Fig. S1). We found that overexpression of SepIVA CT-K did not slow growth in any

genetic background. This result shows that it is the lysine mutations, not the increased

protein levels that are responsible for the slow growth and short cell length of the

sepIVA CT-K mutant.

These data suggest that the ability to methylate the C-terminus of SepIVA is

required for normal growth rate and cell length; however, the N-terminus does not need

to be methylated to sustain normal growth and cell length, and methylation there

actually slows growth and leads to slightly shorter cells. For the remainder of this work,

we focus on the N and C terminal mutants.

Methylation of SepIVA affects it polar localization

SepIVA's localization pattern (13) mimics that of other IMD-associated proteins.

We therefore reasoned that SepIVA may associate with membranes, even though it

does not have predicted transmembrane regions. Because arginines are positively

charged, they could possibly mediate interactions with negatively charged phospholipid

membranes. We sought to determine if the arginine methylations could affect SepIVA's

IMD-like localization pattern, which has bright localization especially at the growing pole.

To test this, we made merodiploid strains of *Msmeg* with GFPmut3-SepIVA methyl-

mutant alleles expressed from the L5 site. We found that methylation of SepIVA affects

its distribution and total intensity (Fig. 2). Specifically, methylation at the N-terminus, in

the NT-M mutant, inhibits SepIVA's localization to the fast-growing pole (Fig. 2ACD),

and has similar signal at both poles (Fig. 2CD). Wild-type cells and mutants that cannot

be methylated at the N-terminus (NT-K) have brighter GFPmut3-SepIVA intensity at the

fast-growing pole than the slow-growing pole, as determined by brighter HADA staining

(Fig. 2CD). The inability to be methylated at the C-terminus, in the CT-K mutant, nearly

equalizes SepIVA distribution between the poles (Fig. 2C). Methylation mutants also

exhibit differences in GFPmut3-SepIVA intensity broadly (Fig. 2B), but these do not

necessarily correlate with western blot results (Fig. 1C), so we cannot conclude whether

GFPmut3-SepIVA signal is an indicator of protein stability. Our findings suggest that

SepIVA's localization to the cell poles is regulated by methylation on the N-terminus of

SepIVA.

Arginine methylation of SepIVA affects PG metabolism

SepIVA's essentiality for cell division and sequence similarity to DivIVA proteins

indicates that it regulates the cell wall. To determine if the arginine methylations on

SepIVA impact metabolism of cell wall components, we used fluorescent dyes that stain

these components when they are metabolically active. We stained log. phase cultures

with the fluorescent D-amino acid, HADA (34), which incorporates into the

peptidoglycan, and fluorescent trehalose DMN-tre (35), which stains the mycolic acid

layer of the cell wall (Fig. 3A). We found that arginine methylation of SepIVA affects PG

metabolism more strongly than MA metabolism, indicating that it affects PG more

directly (Fig. 3BCD). Specifically, methylation at the N-terminus seems to inhibit PG

metabolism, as the NT-M mutants exhibit the dimmest HADA staining (Fig. 3BC).

Interestingly, the NT-K *sepIVA* mutant, which had brighter GFPmut3-SepIVA signal at

the fast-growing pole also had brighter HADA signal at the poles. These data suggest

that SepIVA is a regulator of polar PG metabolism, despite its role as an essential cell

division factor. Furthermore, methylation at the N-terminus of SepIVA appears to inhibit

polar PG metabolism, while a mutant that cannot be methylated at the N-terminus, NT-

K, has increased PG metabolism over wild-type, especially at the poles.

We also observed differences in septal HADA staining between the wild-type and
CT-K strain (Fig. 3E). In *sepIVA* wild-type cells, HADA-stained septa are restricted
mostly to cells in the larger half of the cell length distribution. However, cells lacking
methylation at the C-terminus of *sepIVA* (CT-K mutants) show septal HADA staining in
cells of much shorter lengths. The very short average cell lengths in CT-K cell (Fig. 1B)
also support the model that, when the C-terminus of SepIVA cannot be methylated,
septation is cued a shorter cell lengths. Thus, arginine methylation of *sepIVA* impacts
PG metabolism at both the poles and septa.
176

Arginine methylation of SepIVA affects polar elongation

To determine if the changes in HADA staining in the *sepIVA* methyl-mutants
were due to alterations in the activity of LDT-mediated peptidoglycan remodeling (36) or
due to insertion of new peptidoglycan at the cell poles, we measured polar elongation in
the *sepIVA* wild-type and mutant strains. We developed a method of measuring
elongation using a pulse of HADA (blue), 1.5 hours of outgrowth in media, and another
pulse of NADA (green) stain. We then measured the distance of the NADA-stained but
HADA-clear poles, which represents the polar growth during the 1.5-hour outgrowth
(Fig. 4A).
First, we used this method to determine if depletion of SepIVA affects cell
elongation. SepIVA depletion leads to a slight increase in net polar elongation, with the
increase restricted to the slower-growing poles (Fig. 4B). Because cell division is
inhibited in the SepIVA-depletion condition, the slower-growing poles there are older
than the slower poles in the SepIVA-induced condition. Mycobacterial cell poles typically
do not elongate at new poles for a period after division, then begin elongating at the
same rate as old poles (37). From these data we conclude that SepIVA is not essential
for elongation, and that it is not required for pole maturation.

To determine if methylation of SepIVA affects elongation, we performed pulse-

chase-pulse staining on *sepIVA* wild-type and mutant strains. We found that methylation

at the N-terminus inhibits elongation of both poles, and lack of methylation at N-

terminus stimulates elongation (Fig. 4C). We also found that lack of methylation at the

C-terminus inhibits elongation. Methylation at the two termini of SepIVA thus appear to

have opposite effects on elongation. These results show that SepIVA is an elongation

regulator; since it is not required for elongation (Fig. 4B) we conclude that its function

may be to help toggle growth between the poles and septum. We observe that HADA

staining results match elongation results in the N-terminal mutants: the brighter polar

HADA staining (Fig. 3AB) correlated with greater elongation (Fig. 4C). Thus, we

conclude that the HADA is reflecting differences in newly synthesized PG, not just

differences in LDT-mediated remodeling, and that arginine methylation at the N-

terminus of SepIVA inhibits polar growth, while lack of methylation stimulates polar

growth.

SepIVA affects MurG localization

Our previous study showed that GFPmut3-SepIVA has a similar localization

pattern as the IMD-associated arabinogalactan precursor enzyme GlfT2 (13). The IMD

is a membrane domain that can be separated by sucrose density centrifugation from the

cytoplasmic fractions and from the PM-CW (plasma membrane-cell wall) fraction, which

contains PBPs and other periplasmic cell wall enzymes. To determine if SepIVA is truly

associated with the IMD, we performed sucrose gradient fractionation on an *Msmeg*

strain with a C-terminal strep tag on SepIVA. We found that SepIVA was found in the

cytoplasmic fractions 1-3. GlfT2 was previously shown to be found restricted to the IMD

in fractions 4-6 (38). MurG, the final cytoplasmic enzyme in PG precursor synthesis,

which also has a typical IMD-localization pattern (11, 16), is found partially in the IMD

and partially in the cytoplasmic fractions (16). Thus, SepIVA and MurG have some

overlap in their density-gradient distribution.

Because other DivIVA homologs are involved in recruiting proteins to their site of

activity (19, 20), we hypothesized that perhaps SepIVA could be involved in recruiting

proteins to its locations near the IMD and at the septum. To test this, we looked at the

effects of SepIVA depletion on localization of GlfT2-mRFP and MurG-Dendra2. We

depleted SepIVA in a strain that has a C-terminal DAS tag on the only copy of *sepIVA*,

and then induced *sspB* to direct SepIVA-DAS to be proteolyzed by ClpP (13, 39). We

found that depletion of SepIVA had little to no effect on GlfT2-mRFP localization (Fig.

5BC). However, we found that MurG-Dendra2 was more evenly distributed in cells that

are not inducing *sspB and* was in a more punctate pattern in cells induced for *sspB* and

depleted for SepIVA (Fig. 5DE). We conclude that SepIVA does not affect IMD structure

globally and is not required for IMD-association of GlfT2. Our results indicate, however,

that SepIVA could have a role in dispersing MurG from certain sites, or in modulating its

stability.

The delocalization of MurG-Dendra2 in the SepIVA-DAS strain was not observed

in a SepIVA-DAS strain that did not carry the *sspB* expression plasmid (Fig. S2).

Therefore, the delocalization of MurG-Dendra2 is not due to the DAS tag, but to leaky

expression of SspB in the uninduced condition. These data indicate that MurG-Dendra2

localization is sensitive to subtle changes in SepIVA levels. When SepIVA levels are

moderately decreased, MurG-Dendra2 is delocalized and appear cytoplasmic (Fig. 5D,

top), but when SepIVA is significantly depleted, MurG-Dendra2 localizes in puncta

which could be associated with the IMD (Fig. 5D, bottom).

Arginine methylation of SepIVA affects polar localization of MurG and GlfT2.

We determined that depletion of SepIVA affects localization of MurG and but not

GlfT2 (Fig. 5). To determine whether arginine methylation of SepIVA affects MurG or

GlfT2 localization, we localized both proteins independently in *sepIVA* wild-type and
methyl-mutant strains. According to our results, lack of methylation of SepIVA at the C-
terminus inhibits GlfT2-mRFP and MurG-Dendra2's polar localization (Fig. 6ACDF),
however, the N-terminus does not affect GlfT2-mRFP localization (Fig. 6ABC).
Furthermore, methylation of SepIVA at the N-terminus (NT-M) inhibits MurG's
association with the growing pole (Fig. 6DEF). When sepIVA is unmethylated at the N-
terminus (NT-K), MurG has a stronger localization to the growing pole. This suggests
that SepIVA that is unmethylated at the N-terminus stimulates MurG's association with
the growing pole (Fig 6F), possibly by recruiting it to that site (Fig. 2) and somehow
activating it to promote polar PG synthesis (Fig. 3,4).
To ensure that division defects were not due to effects of arginine methylation of
SepIVA on the FtsZ ring, we expressed *ftsZ-mcherry2b* in methylation mutants. Our
results show that arginine methylation of SepIVA does not affect FtsZ-mcherry2B
localization (Fig. S3A).
265
<u>Discussion</u>
Mycobacteria must regulate not only peptidoglycan precursor synthesis and
assembly, but also the flow of precursors for arabinogalactan and mycolic acid
synthesis. Little is known about how the various synthesis pathways for these different
cell wall layers are coordinated. However, since the PG layer is the closest to the inner
membrane, we expect that PG precursor synthesis is the first to be regulated. Our data
indicate that SepIVA is likely directly regulating PG precursor synthesis, which likely has
an indirect effect on the other layers including the mycomembrane (Fig. 3B). Our data
indicate that SepIVA regulates MurG (Fig. 5,6), which is the final enzyme to work on PG
precursors before they are flipped into the periplasm (1, 40), and is therefore an
important site of regulation for the PG synthesis pathway. We have yet to determine
whether this regulation is direct or indirect.

The IMD is the site of synthesis of lipid-linked cell wall precursors (15, 16)

supporting its involvement in both elongation and division. However, we do not currently

understand how the IMD is reorganized during the cell cycle to allow for septation and

the creation of the new poles. In *E. coli*, MurG interacts with both elongation and

division factors (1, 41). We expect this must be the case in mycobacteria as well, and

that MurG must need to re-localize at the new IMD at the septum. Our data show that

SepIVA is helping to regulate both elongation (Fig.4) and division (Fig. 3F), and that it

regulates MurG localization. SepIVA's effects on the localization of MurG could be

critical for its function in modulation of both elongation and division.

According to our data, SepIVA depletion results in accumulation or aggregation

of MurG. We speculate that SepIVA is somehow affecting MurG's association with the

IMD. In *E. coli,* MurG associates with, and depends on the MreBCD proteins for its

localization to sites of elongation (42). It appears that MurG$_{Ec}$ is distributed

cytoplasmically in the absence of *mreBCD* (42), similarly to how MurG is apparently

distributed cytoplasmically in the *sepIVA*-DAS giles::*sspB* strain (Fig. 5D). This implies

that MurG proteins are regulated and localized by cytoplasmic factors, and that they can

potentially sometimes be dissociated from the membrane. We expect that a

cytoplasmically localized MurG protein would be less active, because it is separated

from a high concentration of its substrate, lipidI. Because we also see greater MurG-

Dendra2 signal when SepIVA is depleted, it is possible that SepIVA is regulating MurG

by destabilizing it, or directing it to proteolysis. These may not be separate phenomena.

Relocation to the cytoplasm may be sufficient to destabilize a membrane-associated

protein.

Regardless of how exactly SepIVA is affecting MurG's localization, our data

indicate that arginine methylation on SepIVA is involved in regulating MurG. MurG

localization to the pole correlates with polar growth efficiency (Fig. 4, 6). The *sepIVA*

NT-K mutant has higher MurG-Dendra2 polar signal (Fig. 6), as well has greater PG

metabolism (Fig. 3) and polar growth (Fig. 4) than the wild-type, while the NT-M and

CT-K mutants both have less polar MurG-Dendra2 signal as well as less PG

metabolism and polar growth (Fig. 3,4,6). The fact that polar localization of MurG is

correlated with higher PG metabolism and polar growth also supports the idea that

MurG is at the IMD and active when it is localized to the pole. If SepIVA's role in

elongation regulation is to regulate MurG's association and dispersal from the IMD

domain at the poles, then it appears that when SepIVA is methylated at the N-terminus,

or unmethylated at the C-terminus, it is more active in dispersing MurG from the poles.

Our current model for SepIVA's arginine methylation is that the two poles are

methylated and demethylated at different times during the cell cycle. We expect that

during polar growth, the N-terminus is unmethylated and the C-terminus is methylated.

The unmethylated N-terminus promotes MurG association with the pole (Fig. 6DF) and

polar elongation (Fig. 4), while the methylation of the C-terminus inhibits septation (Fig.

1B, 3E). When the cell is ready to begin septation, we expect that it demethylates the C-

terminus of SepIVA, and this cues a slowing of elongation (Fig. 4) and initiation of

septation (Fig. 3E), possibly partially through re-localization of MurG (Fig. 6).

Methylation of the N-terminus (Fig. 4) may occur either to slow polar growth during

division, or possibly to inhibit polar growth at the new pole.

Studies on protein arginine methylation done in eukaryotes supports its

importance in cell cycle progression (22, 23). Our data establish protein arginine

methylation as a regulator of the bacterial cell cycle as well. We have found that the

arginine methylation on SepIVA affects both the elongation and division stages of the

cell cycle. This work establishes protein arginine methylation as a physiologically

important post-translational modification in bacteria and identifies SepIVA as a mediator

between the elongation and division processes.

Methods and Materials

Bacterial strains and culture conditions.

Mycobacterium smegmatis (mc^{2}155) was cultured in 7H9 medium (Becton,

Dickinson, Franklin Lakes, NJ) with 5 g/L bovine serum albumin (BSA), 2 g/L glucose,

0.85 g/L NaCl, 0.003 g/L catalase, 0.2% glycerol, and 0.05% Tween 80 or plated on LB

agar. For *M. smeg* cultures, antibiotic concentrations were: 20 µg/mL nourseothricin, 20

µg/mL zeocin, 20 µg/mL streptomycin, 25 µg/mL kanamycin, 50 µg/mL hygromycin, and

30 µg/mL apramycin. For *E. coli* cultures, antibiotic concentrations were: 40 µg/mL

nourseothricin, 25 µg/mL zeocin, 50 µg/mL streptomycin, 50 µg/mL kanamycin, 100

µg/mL hygromycin, and 50 µg/mL apramycin *E. coli* TOP10 or DH5α were used for

cloning.

Strain construction

Because it is essential in *M. smegmatis, sepIVA* mutants were constructed by

first integrating an additional copy of *sepIVA* at the L5 integration site. After

complementing *sepIVA* at the L5 site, *sepIVA* at its native locus was knocked out via

recombineering (43), as previously described (31). Mutant copies of *sepIVA* replaced

wildtype *sepIVA* at the L5 site via allelic exchange (44, 45).

sepIVA depletion strains were constructed as previously described (13).

Merodiploid expression constructs of GlfT2-mRFP, FtsZ-mcherry2B, and MurG-

Dendra2 were integrated at the Tweety integrase site (46). The merodiploid construct

expressing GFPmut3-SepIVA was integrated at the L5 integrase site. All strains used

are listed in Table S3 of the supplementary material, along with plasmids listed in Table

S4 and primers listed in Table S5.

Growth rate assay

Biological replicates of *sepIVA* arginine mutants were grown to log phase in 7H9

medium. Cultures were diluted to OD$_{600}$ = 0.1 in 200 µL 7H9 medium in a non-treated

96-well plate. Optical density at 600 nm was measured over the next 18-24 hours using

a plate reader (BioTek Synergy neo2 multi-mode reader) at 37°C, shaking continuously.

Population doubling times were determined using the exponential growth equation and

least squares regression fitting method in GraphPad Prism (version 9.2). P values were

calculated using ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a

single pooled variance.

Western blots

Log phase cultures (10 mL) were pelleted and resuspended in 500 µL PBS + 1

mM phenylmethylsulfonyl fluoride (PMSF). Cells were lysed using glass beads

(MiniBeadBeater-16, model 607, Biospec). Supernatants from cell lysates were run on

4–12% NuPAGE Bis Tris precast gels (Life Technologies, Beverley, MA) using MES

running buffer (Life Technologies, Beverly, MA). Proteins were transferred onto

polyvinylidene difluoride (PVDF) membranes (GE Healthcare). Strep-tagged proteins

were detected using rabbit anti-Strep-tag II antibody (1:1000, Abcam, ab76949) in Tris-

buffered saline with Tween 20 (TBST) with 0.5% BSA and goat anti-rabbit IgG (H+L)

horseradish peroxidase (HRP)-conjugated secondary antibody (1:10,000, Thermo Fisher

Scientific 31460) in TBST.

Microscopy

Cells were imaged using a Nikon Ti-2 widefield epifluorescence microscope with

a Photometrics Prime 95B camera and a Plan Apo 100X, 1.45 -numerical-aperture (NA)

lens objective. Green-fluorescent images for GFPmut3 or Dendra2 localization or DMN-

tre staining were taken with a 470/40nm excitation filter and a 525/50nm emission filter.

Blue-fluorescent images for HADA staining were taken with a 350/50nm excitation filter

and a 460/50nm emission filter. Red-fluorescent images for mRFP and mcherry2B

localization were taken with a 560/40nm excitation filter and a 630/70 emission filter.

Images were captured using NIS Elements software. Images were analyzed using FIJI

and MicrobeJ (47). For cell detection using MicrobeJ, parameters for width and area

were set. V-snapping cells were cut at the septum so daughter cells could be detected
as individual cells. Overlapping cells were excluded from analysis.

Cell lengths, mean intensities, maximum intensities and minimum intensities of at
least 250 cells per genotype were quantified using MicrobeJ. Mean intensity profiles
were plotted using the "XStatProfile" plotting tool in MicrobeJ. P-values were calculated
using ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a single
pooled variance using GraphPad Prism (v9.2).

Localization of cell wall proteins upon SepIVA depletion.

For depletion of SepIVA, log phase cells (10 mL) were incubated (37°C) with 500
ng/mL Anhydrotetracycline (ATc) for 7-9 hours. For induction of GlfT2-mRFP, 1 ng/mL
IVN were added, and cells were incubated (37°C) for 5 hours. Cells were fixed for
microscopy. Cells were fixed for microscopy with 1.6% paraformaldehyde for 10
minutes and then washed and resuspended in PBS tween80. Microscopy images were
analyzed in ImageJ and MicrobeJ. P-values calculated by unpaired t-test using
GraphPad Prism (v9.2).

Fluorescent staining

HADA and NADA fluorescent D-alanine amino acids were used to stain the
peptidoglycan cell wall layer (48). For PG analysis, HADA stain was added to 1 mL
exponentially growing cells at a final concentration of 1 ug/mL for 2 minutes at room
temperature before washing twice in PBST. For pulse-chase experiments, HADA (1
ug/mL) was added for 10 minutes before cells were washed twice in 7H9, incubated
(37°C) for 2 hours, and stained again using NADA (1 ug/mL) for 2 minutes. Cells were
washed twice in PBST and imaged. Images were analyzed in ImageJ (NIH). The length
of the poles and the length of the entire cell were measured manually for each cell.

DMN-tre stain was used to visualize the trehalose mycolic acid layer of the cell
wall (35). DMN-tre stain was added to 500 µL log phase culture at final concentration 1

µg/mL. Cultures were incubated (37°C) for 30 minutes, washed once with 7H9 and once

with PBST before imaging.

Immunoprecipitation and mass spectrometry

We previously performed LCQ-MS/MS to identify peptides from *Mtb* cell lysates

(30). Here, we re-analyzed those data to search for modifications in peptides, using

Andromeda software.

SepIVA-strep was immunoprecipitated from CB1223. Cells were grown to late

log. phase, pelleted and frozen. Cell pellets were resuspended in PBS with PMSF at 1

mM, lysed by bead beating, and clarified by centrifugation. Immunoprecipitation was

performed with MagStrep type 3 XT beads (Iba Life Sciences, Gottingen, Germany)

according to the manufacturers protocol. Liquid was evaporated from the

immunoprecipitants, and then digested by Trypsin/Lys-C Mix, Mass Spec Grade

protease (Promega), according to the manufacturer's protocols for reduction, alkylation

and two-step in-solution digestion of proteolytically resistant proteins. The digested

peptide solution was then cleaned by C18-Ziptips (Sigma Aldrich) according to the

manufacturers protocol.

LC-MS analysis was performed on LTQ Velos pro mass spectrometer (Thermo

Scientific, CA, USA) combined with a UHPLC (UltiMate 3000, Dionex, USA). The

digested peptides were separated by a nano viper analytical C18 column. (Acclaim

pepMap, 150 mm × 75µm, 2µm, 100 Å, Thermo Scientific, CA, USA). A 60 min gradient

method was used to separate the digested peptides (0–3 min 4.0%B, 3–50 min 4.0–

50.0% B, 50–50.1 min 50–90% B,50.1–55 min 90% B, 55–55.1 min 90–4% B, 55.1–60

min 4% B; mobile phase A: 0.1% FA in water; mobile phase B: 0.1% FA in 95%

acetonitrile, 5% water). 5µL of digested samples was injected at 300 nL/min flow. The

nano-electrospray ionization (ESI) source used with a fixed spray voltage at 2.4 kV and

a heated capillary temperature at 275 °C. A full MS spectrum obtained in normal scan

mode from 350 to 2000 m/z mass range. The data-dependent acquisition was

performed to get the MS/MS spectra for the 5 most abundant ions. The software

Proteome Discoverer (Thermo Scientific, USA) was used to search and identify the

peptides. LC-MS was performed at the Shimadzu Center for Advanced Analytical

Chemistry at UT Arlington.

Acknowledgements

This work was funded by grant R15GM131317 from the NIH. We thank Heather

Lake who contributed to optimization of some experiments, and Arash Emami Saleh

who aided in data processing.

<u>References</u>

1.   Egan AJF, Errington J, Vollmer W. 2020. Regulation of peptidoglycan synthesis

and remodelling. Nat Rev Microbiol https://doi.org/10.1038/s41579-020-0366-3.

2.   Thanky NR, Young DB, Robertson BD. 2007. Unusual features of the cell cycle in

mycobacteria: Polar-restricted growth and the snapping-model of cell division.

Tuberculosis 87:231–236.

3.   Aldridge BB, Fernandez-Suarez M, Heller D, Ambravaneswaran V, Irimia D, Toner

M, Fortune SM. 2012. Asymmetry and Aging of Mycobacterial Cells Lead to

Variable Growth and Antibiotic Susceptibility. Science 335:100–104.

4.   Brown PJB, de Pedro MA, Kysela DT, Van der Henst C, Kim J, De Bolle X, Fuqua

C, Brun YV. 2012. Polar growth in the Alphaproteobacterial order Rhizobiales.

Proceedings of the National Academy of Sciences 109:1697–1701.

5.   Cameron TA, Zupan JR, Zambryski PC. 2015. The essential features and modes of

bacterial polar growth. Trends in Microbiology 23:347–353.

6.   Zupan JR, Grangeon R, Robalino-Espinosa JS, Garnica N, Zambryski P. 2019.
GROWTH POLE RING protein forms a 200-nm-diameter ring structure essential
for polar growth and rod shape in Agrobacterium tumefaciens. PNAS 201905900.

7.   Zupan J, Guo Z, Biddle T, Zambryski P. 2021. *Agrobacterium tumefaciens* Growth
Pole Ring Protein: C Terminus and Internal Apolipoprotein Homologous Domains
Are Essential for Function and Subcellular Localization. mBio 12:e00764-21,
/mbio/12/3/mBio.00764-21.atom.

8.   Kang C-M, Nyayapathy S, Lee J-Y, Suh J-W, Husson RN. 2008. Wag31, a
homologue of the cell division protein DivIVA, regulates growth, morphology and
polar cell wall synthesis in mycobacteria. Microbiology 154:725–735.

9.   Kang C-M. 2005. The Mycobacterium tuberculosis serine/threonine kinases PknA
and PknB: substrate identification and regulation of cell shape. Genes &
Development 19:1692–1704.

10.   Melzer ES, Sein CE, Chambers JJ, Sloan Siegrist M. 2018. DivIVA concentrates
mycobacterial cell envelope assembly for initiation and stabilization of polar growth.
Cytoskeleton 75:498–507.

11.   Meniche X, Otten R, Siegrist MS, Baer CE, Murphy KC, Bertozzi CR, Sassetti CM.
2014. Subpolar addition of new cell wall is directed by DivIVA in mycobacteria.
Proceedings of the National Academy of Sciences 111:E3243–E3251.

12.   Hett EC, Rubin EJ. 2008. Bacterial Growth and Cell Division: a Mycobacterial
Perspective. Microbiology and Molecular Biology Reviews 72:126–156.

13.   Wu KJ, Zhang J, Baranowski C, Leung V, Rego EH, Morita YS, Rubin EJ, Boutte
CC. 2018. Characterization of Conserved and Novel Septal Factors in

Mycobacterium smegmatis. Journal of Bacteriology 200:e00649-17,

/jb/200/6/e00649-17.atom.

14.   Jankute M, Cox JAG, Harrison J, Besra GS. 2015. Assembly of the Mycobacterial

Cell Wall. Annual Review of Microbiology 69:405–423.

15.   Hayashi JM, Luo C-Y, Mayfield JA, Hsu T, Fukuda T, Walfield AL, Giffen SR,

Leszyk JD, Baer CE, Bennion OT, Madduri A, Shaffer SA, Aldridge BB, Sassetti

CM, Sandler SJ, Kinoshita T, Moody DB, Morita YS. 2016. Spatially distinct and

metabolically active membrane domain in mycobacteria. PNAS 113:5400–5405.

16.   García-Heredia A, Kado T, Sein CE, Puffal J, Osman SH, Judd J, Gray TA, Morita

YS, Siegrist MS. 2021. Membrane-partitioned cell wall synthesis in mycobacteria.

eLife 10:e60263.

17.   Jain P, Malakar B, Khan MZ, Lochab S, Singh A, Nandicoori VK. 2018. Delineating

FtsQ-mediated regulation of cell division in *Mycobacterium tuberculosis*. Journal of

Biological Chemistry 293:12331–12349.

18.   Flores AR, Parsons LM, Pavelka MS. 2005. Characterization of Novel

Mycobacterium tuberculosis and Mycobacterium smegmatis Mutants

Hypersusceptible to -Lactam Antibiotics. Journal of Bacteriology 187:1892–1900.

19.   Marston AL, Errington J. 1999. Selection of the midcell division site in Bacillus

subtilis through MinD-dependent polar localization and activation of MinC.

Molecular Microbiology 33:84–96.

20.   Halbedel S, Lewis RJ. 2019. Structural basis for interaction of DivIVA/GpsB

proteins with their ligands. Molecular Microbiology 111:1404–1415.

21. Hammond LR, Khan SJ, Sacco MD, Spanoudis C, Hough A, Chen Y, Eswara PJ.
2021. GpsB coordinates cell division and cell surface decoration by wall teichoic
acids in *Staphylococcus aureus*. preprint, Microbiology.

22. Raposo AE, Piller SC. 2018. Protein arginine methylation: an emerging regulator of
the cell cycle. Cell Div 13:3.

23. Biggar KK, Li SS-C. 2015. Non-histone protein methylation as a regulator of cellular
signalling and function. Nat Rev Mol Cell Biol 16:5–17.

24. Zhang M, Xu J-Y, Hu H, Ye B-C, Tan M. 2018. Systematic Proteomic Analysis of
Protein Methylation in Prokaryotes and Eukaryotes Revealed Distinct Substrate
Specificity. PROTEOMICS 18:1700300.

25. Sakatos A, Babunovic GH, Chase MR, Dills A, Leszyk J, Rosebrock T, Bryson B,
Fortune SM. 2018. Posttranslational modification of a histone-like protein regulates
phenotypic resistance to isoniazid in mycobacteria. Science Advances 4:eaao1478.

26. Su Y, Li J, Zhang W, Ni J, Huang R, Wang Z, Cheng S, Wang Y, Tian Z, Zhou Q,
Lin D, Wu W, Tang CM, Liu X, Lu J, Yao Y-F. 2021. Methylation of PhoP by CheR
Regulates *Salmonella* Virulence. mBio https://doi.org/10.1128/mBio.02099-21.

27. Singhal A, Virmani R, Naz S, Arora G, Gaur M, Kundu P, Sajid A, Misra R, Dabla
A, Kumar S, Nellissery J, Molle V, Gerth U, Swaroop A, Sharma K, Nandicoori VK,
Singh Y. 2020. Methylation of two-component response regulator MtrA in
mycobacteria negatively modulates its DNA binding and transcriptional activation.
Biochemical Journal 477:4473–4489.

28. Prisic S, Husson RN. 2014. Mycobacterium tuberculosis Serine/Threonine Protein
Kinases. Microbiology Spectrum 2.

29. Prisic S, Dankwa S, Schwartz D, Chou MF, Locasale JW, Kang C-M, Bemis G,
Church GM, Steen H, Husson RN. 2010. Extensive phosphorylation with
overlapping specificity by Mycobacterium tuberculosis serine/ threonine protein
kinases. Proceedings of the National Academy of Sciences 107:7521–7526.

30. Garces A, Atmakuri K, Chase MR, Woodworth JS, Krastins B, Rothchild AC,
Ramsdell TL, Lopez MF, Behar SM, Sarracino DA, Fortune SM. 2010. EspA Acts
as a Critical Mediator of ESX1-Dependent Virulence in Mycobacterium tuberculosis
by Affecting Bacterial Cell Wall Integrity. PLOS Pathogens 6:e1000957.

31. Boutte CC, Baer CE, Papavinasasundaram K, Liu W, Chase MR, Meniche X,
Fortune SM, Sassetti CM, Ioerger TR, Rubin EJ. 2016. A cytoplasmic
peptidoglycan amidase homologue controls mycobacterial cell wall synthesis. eLife
5.

32. Iswahyudi, Mukamolova GV, Straatman-Iwanowska AA, Allcock N, Ajuh P, Turapov
O, O'Hare HM. 2019. Mycobacterial phosphatase PstP regulates global serine
threonine phosphorylation and cell division. Sci Rep 9:8337.

33. Shamma F, Papavinasasundaram K, Quintanilla SY, Bandekar A, Sassetti C,
Boutte CC. 2020. Phosphorylation on PstP Regulates Cell Wall Metabolism and
Antibiotic Tolerance in *Mycobacterium smegmatis*. J Bacteriol 203:e00563-20,
/jb/203/4/JB.00563-20.atom.

34. Kuru E, Tekkam S, Hall E, Brun YV, Van Nieuwenhze MS. 2015. Synthesis of
fluorescent D-amino acids and their use for probing peptidoglycan synthesis and
bacterial growth in situ. Nat Protoc 10:33–52.

35. Kamariza M, Shieh P, Ealand CS, Peters JS, Chu B, Rodriguez-Rivera FP, Sait
MRB, Treuren WV, Martinson N, Kalscheuer R, Kana BD, Bertozzi CR. 2018.

562   Rapid detection of Mycobacterium tuberculosis in sputum with a solvatochromic

563   trehalose probe. Science Translational Medicine 10:eaam6310.

564 36. Baranowski C, Welsh MA, Sham L-T, Eskandarian HA, Lim HC, Kieser KJ, Wagner

565   JC, McKinney JD, Fantner GE, Ioerger TR, Walker S, Bernhardt TG, Rubin EJ,

566   Rego EH. 2018. Maturing Mycobacterium smegmatis peptidoglycan requires non-

567   canonical crosslinks to maintain shape. eLife 7:e37516.

568 37. Hannebelle MTM, Ven JXY, Toniolo C, Eskandarian HA, Vuaridel-Thurre G,

569   McKinney JD, Fantner GE. 2020. A biphasic growth model for cell pole elongation

570   in mycobacteria. Nat Commun 11:452.

571 38. Hayashi JM, Richardson K, Melzer ES, Sandler SJ, Aldridge BB, Siegrist MS,

572   Morita YS. 2018. Stress-Induced Reorganization of the Mycobacterial Membrane

573   Domain. mBio 9.

574 39. Kim J-H, Wei J-R, Wallach JB, Robbins RS, Rubin EJ, Schnappinger D. 2011.

575   Protein inactivation in mycobacteria by controlled proteolysis and its application to

576   deplete the beta subunit of RNA polymerase. Nucleic Acids Res 39:2210–2220.

577 40. Gee CL, Papavinasasundaram KG, Blair SR, Baer CE, Falick AM, King DS, Griffin

578   JE, Venghatakrishnan H, Zukauskas A, Wei J-R, Dhiman RK, Crick DC, Rubin EJ,

579   Sassetti CM, Alber T. 2012. A Phosphorylated Pseudokinase Complex Controls

580   Cell Wall Synthesis in Mycobacteria. Sci Signal 5:ra7–ra7.

581 41. Laddomada F, Miyachiro MM, Dessen A. 2016. Structural Insights into Protein-

582   Protein Interactions Involved in Bacterial Cell Wall Biogenesis. Antibiotics (Basel)

583   5:14.

584 42. Mohammadi T, Karczmarek A, Crouvoisier M, Bouhss A, Mengin-Lecreulx D, den

585   Blaauwen T. 2007. The essential peptidoglycan glycosyltransferase MurG forms a

complex with proteins involved in lateral envelope growth as well as with proteins
involved in cell division in Escherichia coli. Mol Microbiol 65:1106–1121.

43. van Kessel JC, Hatfull GF. 2008. Mycobacterial Recombineering. Methods in
Molecular Biology (Clifton, NJ) 435:203–215.

44. Pashley CA, Parish T. 2003. Efficient switching of mycobacteriophage L5-based
integrating plasmids in *Mycobacterium tuberculosis*. FEMS Microbiology Letters
229:211–215.

45. Kieser KJ, Boutte CC, Kester JC, Baer CE, Barczak AK, Meniche X, Chao MC,
Rego EH, Sassetti CM, Fortune SM, Rubin EJ. 2015. Phosphorylation of the
Peptidoglycan Synthase PonA1 Governs the Rate of Polar Elongation in
Mycobacteria. PLOS Pathogens 11:e1005010.

46. Pham TT, Jacobs-Sera D, Pedulla ML, Hendrix RW, Hatfull GFY 2007.
Comparative genomic analysis of mycobacteriophage Tweety: evolutionary insights
and construction of compatible site-specific integration vectors for mycobacteria.
Microbiology 153:2711–2723.

47. Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. Nature Microbiology 1:16077.

48. Kuru E, Hughes HV, Brown PJ, Hall E, Tekkam S, Cava F, de Pedro MA, Brun YV,
VanNieuwenhze MS. 2012. In Situ Probing of Newly Synthesized Peptidoglycan in
Live Bacteria with Fluorescent D -Amino Acids. Angewandte Chemie International
Edition 51:12519–12523.

Figure 1. Arginine methylation of SepIVA affects growth and cell length of *M.smeg.*
(A) Doubling time of *M. smeg* cells expressing WT or arginine methylation mutants of SepIVA. Black outline, lighter color bars represent arginine to alanine mutants. No outline, darker color bars represent methyl-ablative mutants. Red outline bars represent methyl-mimetic mutants. Strains without bars were non-viable. At least 3 biological replicates of each strain were used. Error bars represent standard deviation. ****, $P = <0.0001$.
(B) Cell lengths of SepIVA WT and arginine methylation mutants. Bars represent mean. At least 300 cells over three biological replicates of each strain were analyzed. *, $P = 0.0439$; ****, $P = <0.0001$.
(C) α-strep western blot of L5::*sepIVA*-strep WT or arginine methylation mutants. mc^{2}155 without a strep tag was used as a negative control (lane 2). '*' denotes merodiploid strains. The non-specific band serves as a loading control.

Figure 2. Methylation of SepIVA affects its polar localization. (A) Images of *M. smeg* cells expressing L5::*GFPmut3-sepIVA* WT and arginine methylation mutants. Strains are merodiploid. Phase images are on the top. Fluorescent images are on the bottom. Scale bar applies to all images. **(B)** Mean intensity values of 250+ cells expressing L5::*GFPmut3-sepIVA* WT and arginine methylation mutants. Bar represents mean intensity value between cells. The adjusted p-values of each mutants compared to the WT are: NT-K = <0.0001; NT-M = <0.0001; and CT-K = 0.0002. Three biological replicate strains were imaged for each genotype. **(C)** Normalized intensity of poles of cells expressing L5::*GFPmut3-sepIVA* WT and arginine methylation mutants. Each dot represents ratio of pole intensity value over mean intensity value of the cell. Filled in dots represent the faster (brighter) growing pole, empty dots represent the slower (dimmer) growing pole. Bar represents median intensity value. At least 250+ cells across three biological replicates of each mutant were used. The adjusted p-values of the mutant fast poles compared to the WT fast pole are: NT-K = <0.0001; NT-M = <0.0001; and CT-K = <0.0001. The adjusted p-values of the mutant slow poles compared to the WT slow pole are: NT-K = 0.9510; NT-M = <0.0001; and CT-K =

<0.0001. **(D)** Intensity profiles of GFP-SepIVA signal of WT and methylation mutants.

The solid line represents the mean intensity value, and the shaded area represents

standard deviation. At least 250+ cells across three biological replicates of each mutant

were analyzed in MicrobeJ. P-values were calculated using ordinary one-way ANOVA,

Dunnett's multiple comparisons test, with a single pooled variance in GraphPad Prism

(v9.2).

Figure 3. Arginine Methylation of SepIVA affects PG metabolism. (A) Images of M.

smeg cells expressing *L5::sepIVA* WT and arginine methylation mutants. Pictures of

several cells from images processed identically were pasted together. Scale bar applies

to all images. **(B)** Intensity profiles of HADA (blue) and DMN-tre (green) signal of WT

and methylation mutants. The solid line represents the mean intensity value, and the

shaded area represents standard deviation. At least 250+ cells across three biological

replicates of each mutant were analyzed in MicrobeJ. Cells were pole-sorted such that

the brighter pole in the HADA channel was set to 0 on the X axis. **(C and D)** Mean

HADA **(C)** and DMN-tre **(D)** intensity values of M. smeg cells expressing *L5::sepIVA* WT

and arginine methylation mutants. Bar represents mean intensity value. ****, P =

<0.0001. P-values were calculated using ordinary one-way ANOVA, Dunnett's multiple

comparisons test, with a single pooled variance in GraphPad Prism (v9.2). **(E)**

Demographs of HADA intensity across the length of the cell of WT SepIVA and CT-K

mutant. Demographs for other methylation mutants can be found in supplemental figure

2. At least 250+ cells across three biological replicates of each mutant were analyzed in

MicrobeJ. Cells were pole-sorted such that the brighter pole in the HADA channel was

oriented to the top along the Y axis.

A. Cell staining method

HADA → Growth x 1.5 hrs → NADA

Pulse 1 | Pulse 2 | Both
5 um

B. Polar elongation after SepIVA depletion

● Fast pole
○ Slow pole
▲ Sum of poles

Length (um)

+SepIVA   -SepIVA

+SepIVA
5 um

-SepIVA

C. Polar elongation in 1.5 hours

Length (um)

● Fast pole
○ Slow pole

WT SepIVA   NT-K   NT-M   CT-K

WT SepIVA | NT-K | NT-M | CT-K
5 um

Figure 4. Arginine methylation of SepIVA affects polar elongation.
(A) Schematic of pulse-chase-pulse staining method. **(B)** Length of poles after SepIVA depletion. Filled in dots represent the faster (brighter) growing pole, empty dots represent the slower (dimmer) growing pole. Triangles represent the sum of poles or total elongation. At least 100 cells across three biological replicates were analyzed. Bar represents median. Scale bar applies to all images. **(C)** Length of poles elongation after 1.5 hours in cells expressing L5::*sepIVA* WT and arginine methylation mutants. At least 100 cells across three biological replicates were analyzed. Bar represents median polar elongation. The adjusted p-values of each fast pole mutant compared to the WT are: NT-K = <0.0001; NT-M = 0.0458; and CT-K = <0.0001. The adjusted p-values of each slow pole mutant compared to the WT are: NT-K = 0.0446; NT-M = 0.0118; and CT-K = <0.0001. Representative cells are shown. Scale bar applies to all images. P-values were

708   calculated using ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a

709   single pooled variance in GraphPad Prism (v9.2).

Figure 5. SepIVA has the same membrane association as MurG, and it affects

MurG localization. (A) anti-Strep (top) western blot after sucrose gradient

fractionations. Fractions 4-6 contain intracellular membrane domain (IMD) proteins. **(B)**

Micrographs of cells expressing or depleting SepIVA and induced for GlfT2-mRFP.

Scale bar is 5 microns and applies to all images. **(C)** GlfT2-mRFP distribution

represented by coefficient of variation values. Coefficient of variation was calculated

based on fluorescence intensity across the relative position of each cell. Bar represents

median coefficient. **(D)** Micrographs of cells expressing or depleting SepIVA and

induced for MurG-Dendra2. Scale bar is 5 microns and applies to all images. **(E)** MurG-

728   Dendra2 distribution represented by coefficient of variation values. Coefficient of
729   variation was calculated based on fluorescence intensity across the relative position of
730   each cell. Bar represents median coefficient. P-value calculated by unpaired t-test using
731   GraphPad Prism (v9.2).

A. Localization of GlfT2-mRFP in *M. smegmatis* cells

B. Mean GlfT2-mRFP intensity

C. Intensity profiles

D. Localization of MurG-Dendra2 in *M. smegmatis* cells

E. Mean MurG-Dendra2 intensity

F. Intensity profiles

Figure 6. Arginine methylation of SepIVA affects MurG's localization to the poles.

(A) Images of *M. smeg* cells expressing L5::*sepIVA* WT and arginine methylation

mutants and induced for GlfT2-mRFP. Pictures of several cells from images processed

identically were pasted together. Scale bar applies to all images. **(B)** Mean intensity

values of 250+ cells expressing L5::*sepIVA* WT and arginine methylation mutants and

induced for GlfT2-mRFP. Bar represents mean intensity value between cells. The

adjusted p-values of each mutants compared to the WT are: NT-K = 0.9289; NT-M =

<0.0001; and CT-K = <0.0001. At least 2 biological replicate strains were imaged for

each genotype. **(C)** Intensity profiles of GlfT2-mRFP signal of WT and methylation

mutants. The solid line represents the mean intensity value, and the shaded area

represents standard deviation. At least 250+ cells across three biological replicates of

each mutant were analyzed in MicrobeJ. Cells were pole-sorted such that the brighter

pole in the HADA channel was set to 0 on the X axis. **(D)** Images of *M. smeg* cells

expressing L5::*sepIVA* WT and arginine methylation mutants and TW::*murG-dendra2*.

Pictures of several cells from images processed identically were pasted together. Scale

bar applies to all images. **(E)** Mean intensity values of 250+ cells expressing L5::*sepIVA*

WT and arginine methylation mutants and TW::*murG-dendra2*. Bar represents mean

intensity value between cells. The adjusted p-values of each mutants compared to the

WT are: NT-K = <0.0001; NT-M = <0.0001; and CT-K = <0.0001. Three biological

replicate strains were imaged for each genotype. P-values were calculated using

ordinary one-way ANOVA, Dunnett's multiple comparisons test, with a single pooled

variance in GraphPad Prism (v9.2). **(F)** Intensity profiles of MurG-Dendra2 signal of WT

and methylation mutants. The solid line represents the mean intensity value, and the

shaded area represents standard deviation. At least 250+ cells across three biological

replicates of each mutant were analyzed in MicrobeJ. Cells were pole-sorted such that

the brighter pole in the HADA channel was set to 0 on the X axis.

761