Evolution of increased early-life fecundity but faster reproductive senescence in

***Drosophila melanogaster* females through adaptation to chronic live-yeast deprivation**

Purbasha Dasgupta[1], purbasha17@iiserbpr.ac.in

Subhasish Halder[1], subhasish@iiserbpr.ac.in

Debapriya Dari[1], debapriyadari1287@gmail.com

Nabeel P.[1], nabeelp2121@gmail.com

Sai Samhitha Vajja[2], vsais18@iiserb.ac.in

Bodhisatta Nandy*[1], nandy@iiserbpr.ac.in

*Corresponding author

Present address:

1. Indian Institute of Science Education and Research Berhampur, Transit Campus, Govt. ITI

Building, NH 59, Engineering School Junction, Ganjam, Berhampur 760 010, Odisha, India.

Contact number: +91-680-222 7755

2. Indian Institute of Science Education and Research Bhopal, Bhopal Bypass Rd, Bhauri, Madhya

Pradesh 462066.

Key words: Nutritional deprivation, fitness trade-offs, antagonistic pleiotropy, experimental

evolution.

Running title: Female adaptation to live-yeast deprivation

Word count: 5,504 excluding abstract, table and figure captions, acknowledgement and

references.

Figure count: 3

Table count: 2

Supplementary information: A separate file that includes three figures and six tables.

Abstract

Dietary restriction is a common ecological challenge that limits reproduction. Yet only a few studies have explored adaptation under chronic protein deprivation. We subjected four replicate laboratory-adapted populations (YLB) of *Drosophila melanogaster* to a complete deprivation of live-yeast to mimic diet restricted ecology. In this insect, live-yeast is a critical source of protein that strongly affect reproductive output, especially in females. Following 24 generations of experimental evolution, compared to their matched controls (BL), females from YLB populations showed increase in reproductive output early in life, both in presence and absence of live-yeast. The observed increase in reproductive output was not associated with any accommodating alteration in average egg size; and development time, pre-adult survivorship, and body mass at eclosion of the progeny. Interestingly, adult lifespan was also found to be unaffected. However, YLB females were found to have a significantly faster rate of reproductive senescence albeit without any change in a measure of lifetime reproductive output. Taken together, adaptation to LYD ecology shows that reproductive output can evolve without affecting lifespan, suggesting that widely observed reproduction-survival trade-off is not universal. Populations can optimize fitness by fine tuning the scheduling of reproduction even when lifetime reproductive output is constrained.

1. Introduction

The ability of an organism to cope with a nutritional challenge is essential to adaptation in a given ecology (Kolss et al 2009, Zhang et al. 2019). Beyond providing the essential energy to sustain life, diet determines survival by affecting resistance to stress and other ecological challenges, including pathogens (Chippindale 1993, Field et al. 2002, Andersen et al. 2010, Sisodia and Singh 2012, Miller and Cotter 2017, Weldon et al. 2019). In addition, it also affects reproductive output and is central to a variety of trade-offs between components of life history (Chippindale 1993, Tatar & Carey 1995, Chapman & Partridge 1996, Marden et al. 2003, Adler & Bonduriansky 2014). Adaptation to even the simplest of nutritional challenges is expected to be complex in nature, involving a network of life history, physiological and behavioural traits. In response to one form of nutritional challenge, often referred to as dietary restriction (DR), organisms ranging from yeast to mammals have been found to elicit a complex physiological response that culminates in the partial or complete shutdown of reproduction, and extension of lifespan (Nakagawa et al. 2012, Fontana & Partridge 2015). Though the physiology and molecular genetic basis of DR response are well understood (Fontana et al. 2010, Fontana & Partridge 2015), there is considerable debate on the evolutionary significance of such a short term response, and the long term consequence of a chronic DR and other forms of nutritional challenges (Adler & Bonduriansky 2014). Further, little is known about quantitative genetic variance and covariance for traits that determine the DR response (but see Rikke et al. 2010, Jin et al. 2020).

Dietary restriction of one or more essential nutrients without causing malnutrition, has been widely reported to trigger an extension of lifespan in *D. melanogaster,* possibly through lowered reproduction (Mair et al. 2005, Partridge et al. 2005). This plastic response of life history to DR has been investigated in *D. melanogaster* by limiting yeast supplement on food

(Partridge et al. 1987, Chippindale et al.1993) or by dilution of nutrients in food (Chapman

and Partridge, 1996). Mair et al. (2005) showed that simultaneous dilution of sugar and heat-

killed yeast in their food resulted in lifespan extension in laboratory populations. Moreover,

the lifespan extension effect was more prominent in response to yeast restriction than an

equivalent reduction in sucrose, emphasizing the importance of yeast in generating DR

response in flies (Mair et al. 2005). Apart from such DR, other forms of nutritional

challenges, including extreme starvation, have also been studied using the fruit fly system

(Hoffmann & Harshman 1999, Parkash & Munjal 2000, Partridge et al. 2005, Rion &

Kawecki 2007). Similar to DR, both evolutionary and plastic responses to starvation (with

some exceptions) involve a physiological shift to 'survival mode' marked by greater response

to stress and longer life-span at the cost of reduced fecundity (reviewed in Rion & Kawecki

2007).

Resource re-allocation hypothesis, a theory aimed at explaining the plasticity of life history in

response to DR (simply, DR response hereon), proposes that such a response is an adaptive

diversion of resources from reproduction to somatic maintenance to maximise the chance of

survival under a period of temporary resource scarcity (Shanley & Kirkwood 1999, Moatt et

al. 2020). Such resource re-allocation, however, can only lead to higher fitness when DR

condition prevails for a shorter duration (Zajitschek et al. 2016 & 2019, Moatt et al. 2020). In

face of chronic DR, continuing for several generations, natural selection should favour a

higher investment in reproduction at the cost of survival (Zajitschek et al. 2016, 2019). There

have been a few recent investigations on adaptation to chronic DR using *D.*

melanogaster (Kolss et al. 2009, Vijendravarma & Kawecki 2013, Zajitschek et al. 2016,

2019). Kolss et al. (2009) found that after 29-64 generations of selection under larval DR, the

populations had evolved better egg-adult survivorship and faster development in DR food

indicating adaptation to the novel diet. Zajitschek et al. (2016, 2019) investigated adaptation

to prolonged DR in adults, and contrary to the prediction from re-allocation theory, they

found that males from the DR selected populations had higher reproductive output on a DR

food without any effect on longevity even after 25 generations of selection (Zajitschek,

2016). The females on the other hand, after 50 generations of selection, had reduced

longevity without any effect on fecundity (Zajitschek, 2019).

Both laboratory cultures and wild populations of fruit flies, often need live-yeast as a dietary

protein source for reproduction (Begon 1982, Starmer 1982, Chippindale et al., 1993;

Anagnostou, Dorsch & Rohlfs 2010). Importantly, besides affecting female reproductive

output, such yeast supplement impacts adult lifespan and starvation resistance (Chippindale et

al. 1993), oviposition site preference (Becher et al 2012), and may even affect offspring

survival and fitness components (Rohlfs & Hoffmeister 2005, Anagnostou, Dorsch & Rohlfs

2010). Given the strong fitness consequence of feeding on live-yeast, the lack of

experimental data on adaptation to live-yeast deprivation (LYD) in the available literature is

rather surprising. Interestingly, because live-yeast acts as dietary protein supplement

primarily for females (Chippindale et al. 1993, Nandy et al. 2012), it is possible for LYD to

represent a sex-specific selective environment.

Here, we subjected four replicate populations of *D. melanogaster* to chronic LYD in order to

investigate the life-history and behavioural adaptations to such ecology. After 23 generations

of selection, we addressed the following questions to assess adaptation to such an ecology:

(a) Considering female fecundity to be the prime target of selection under LYD, do evolved

females show a clear sign of adaptation to this new maintenance regime? (b) Given that

female reproductive output in *D. melanogaster* is often influenced by their mates in addition

to their own physiology and/or behaviour, does male effect on female fecundity evolve in

response to LYD? (c) If adaptation to LYD involves changes in reproductive output, what

consequences do such adaptation have on life-history traits besides fecundity?

2. Materials and methods

<u>Study populations and selection regime</u>

Four new experimental populations (referred to as YLB_{1-4} here on) were derived from four

replicate baseline populations BL_{1-4} (Figure S1). The BL populations (Nandy et al. 2016) are

outbred, wild type populations of *D. melanogaster*. They are maintained under a 14-day

discrete generation cycle, under 24-hour light, at 25 °C (±1) ambient temperature, and on a

standard banana-sugarcane jaggery-yeast medium. Flies are grown in standard vials (25 mm

diameter × 96 mm height) for 12 days. Larval density is controlled at ~70 per vial with 8 ml

food. On day 12 of the generation cycle, adults are transferred to plexiglass cages (23cm ×

20cm × 15 cm) with a food plate smeared with an *ad-lib* quantity of live yeast (a paste made

with water). On the day 14, the flies are given fresh food for oviposition and eggs laid during

an 18-hour window (fitness window) are collected to start the next generation. BL

populations were maintained in the laboratory under this regime for almost five years prior to

the derivation of the experimental YLB populations. The maintenance of the YLB

populations is identical to that of the BL except for the absence of live-yeast supplement in

the YLB ecology. All assays reported here were conducted after 23-25 generations of

experimental evolution. To equalize any potential non-genetic parental effect, all populations

were passed through one generation of common rearing ecology (Rose 1984). During this, a

subset of the experimental populations was generated by collecting eggs from the stock BL_x

and YLB_x populations. These populations were maintained under a regime that exactly

matched that of the BL-regime. Experimental flies were generated by collecting eggs from

these subset populations. All fly handling, including the collection of virgin flies, wherever

necessary, was done under light CO_2 anaesthesia.

<u>Experiment 1: Juvenile traits and early-life fecundity in females</u>

After 23 generations of experimental evolution, we assessed whether the YLB populations

have adaptively diverged from the ancestral BL populations. Our initial observation

suggested fecundity to be the most likely target trait of selection (data not shown). Therefore,

as the first set of assays, we quantified female early-life fecundity under a range of yeasting

conditions. In addition, since fecundity is often a function of body size and related juvenile

traits, we also investigated these traits in a parallel assay.

Female early-life fecundity and a three important juvenile traits – pre-adult (egg-to-adult)

development time, juvenile (egg-to-adult) survivorship and dry weight at eclosion were

measured between 23-25 generations. Following one generation of common rearing (see

above), eggs were collected from the population cages in exact numbers. We allowed an

oviposition window of about one hour. Such a short window of oviposition ensured the

developmental synchrony of the eggs collected for the experiment. Eggs were transferred

from the oviposition plate to an egg-collection surface made of solidified agar gel (1% Agar-

agar solution) using a fine brush. Eggs were carefully counted and a small piece of Agar gel

containing 70 eggs was cut out and transferred to a fresh food vial having 6-8ml food. Ten

such vials were set up for each population. Each vial was labelled individually and placed

into the incubator. As an added precaution, the vials in the incubator were shuffled twice

every day to randomize any positional effect. The vials were monitored carefully for the signs

of eclosion. Beginning from the eclosion of the first individual, the eclosing adults were

counted and sexed every 6 hours until all the pupae eclosed. This data was used to derive the

average measure of egg-to-adult development time (i.e., pre-adult development time) for

males and females in each vial. Out of the 70 eggs initially cultured, the proportion that

finally eclosed from a vial was considered as the measure of egg-to-adult survivorship (i.e.,

juvenile survivorship). Three vials (one from the YLB_1; two from BL_2) were excluded from

the analysis due to mishandling. For each population (BL_x and YLB_x), 50 males and 50

females were collected for dry whole body weight measurement. They were collected within

6-hours of eclosion, and were immediately frozen at -20 °C. They were later dried at 60°C for

48 hr and weighed in groups of five using Shimadzu AUW220D to the nearest 0.01 mg. The

mean body weight of each group was calculated and taken as a unit of analysis.

For early-life fecundity assay, flies were raised and collected as virgins as mentioned above.

Virgin flies were held in same-sex vials in groups of 12 in each vial. On the day 12 post egg-

collection (i.e., corresponding to 2-3 day post-eclosion), one vial each of males and females

(i.e., 12 males and 12 females) belonging to the same regime, were transferred to a fresh food

vial assigned to one of the three live-yeast supplementation treatments - No yeast

(N), Limiting yeast (L: 4.5 mg yeast) and High yeast (H: 9 mg yeast). Twelve such vials were

set up per live-yeast supplementation treatment for a population. Live-yeast was dissolved in

tap water to make a suspension, which was then dispensed on the top of the food in a vial.

The suspension was allowed to air-dry before the introduction of the experimental flies. The

males and the females were allowed to interact in these vials for 48 hours. Following this, on

day 4-5 post-eclosion flies in each treatment vial were anaesthetized under light CO_2 - six

females were randomly separated and transferred to a vial for egg size measurement, while

the other six were individually transferred to six oviposition tubes (12×75 mm) for fecundity

measurement. For the former, females were allowed to oviposit for 16 hours in a vial,

following which they were discarded. The vial with eggs was frozen at -20 °C for a minimum

of one day. For egg size measurement, eggs were mounted on a glass slide on their dorsal

side and photographed using a stereozoom microscope and the area of the two-dimensional

elliptical outline of the eggs were measured in Image J software. This area was taken as a

proxy for the size of each egg. A given egg was measured three times and the average of

these three measurements was taken as the unit of analysis. Females were allowed to oviposit

in these tubes for 16 hours. Females from the oviposition tubes were then discarded and

number of eggs was counted. We removed a few vials from the analysis due to a variety of

handling errors, including escaped flies and broken tube/vial, during the assay. The final

number of vials used for the analysis ranged from 9-12 vials per live-yeast treatment for a

population.

Experiment 2: Female × male effect on female fitness

To further compare the evolution of female fitness, we quantified female early-life

reproductive output (progeny count = number of surviving adult offspring produced by a

female) during a 16-hour window on the day 4-5 post-eclosion. This window corresponded to

the 18-hour fitness window on the 14th day of the generation cycle and therefore, is a good

measure of Darwinian fitness in this system. Importantly, female fitness, especially in *D.*

melanogaster, is affected by both female fecundity and male factors such as seminal fluid

composition (Chapman et al. 1995; Arnqvist & Nilsson 2000; Wolfner 2009). Therefore, it is

important to assess male and female specific effect of selection on female fitness. To test this

effect, we combined YLB and BL males and females in a full factorial design and measured

early-life reproductive output of females in the above-mentioned protocol. These assays were

carried out after 26-29 generations of selection. Experimental flies were generated following

a protocol similar to that stated above. Males and females were collected as virgins and held

in single-sex vials in groups of six in each vial until day 12 post-egg collection. On the day

12 of the generation cycle (i.e., 2-3 days post-eclosion), sexes were combined to set up adult

interaction vials having six individuals of each sex in a vial. Four types of combinations (viz.,

female×male: BL×YLB; BL×BL; YLB×YLB; YLB×BL), each having 15 replicate vials,

were set up. A few flies escaped during the setting up of one vial in the third block with

$YLB_3 \times YLB_3$ combination and hence, it was removed from the experiment. These adult

interaction vials were left undisturbed for 48 hours. On the day 14, females from each of

these vials were sorted under light CO_2 anaesthesia, and individually transferred to

oviposition test tubes (one female per tube) where they were allowed to oviposit for 17 hours.

After 17 hours, females from the oviposition tubes were discarded and the tubes were kept in

the incubator for the development and emergence of the progeny. After the emergence of all

the progeny, the test tubes were frozen at -20 °C for the final count.

<u>Experiment 3: Longevity and age-specific fecundity</u>

To test reproduction vs. lifespan trade-off theory, longevity and fecundity of the experimental

females were measured after 26 generations of selection. To assess age related changes in

fecundity, we measured fecundity of the test females in regular intervals. Similar to the other

assays described above, all populations were passed through one generation of common

rearing. The eggs were collected from the common rearing cages and cultured under standard

density of ~70 per 6-8ml food in a vial. Adult flies were collected as virgins within six hours

of eclosion and held in single-sex vials in groups of 12 in standard food vials. These vials

were left undisturbed for two days. On the day 3 post eclosion (i.e., on the day 12 post egg

collection), experimental vials were set up by transferring the flies into fresh food vials. Two

sets of vials were set up for each population – (a) Continuously exposed (CE): 12 males and

12 females placed (without anaesthesia) in a vial and (b) Virgin females (VF): 12 virgin

females placed in a vial. For a population, 12 vials were set up for each of CE and VF. Except

for the fecundity count days (see below), flies were transferred into fresh food vials every

alternate day without anaesthesia. Dead flies were sexed and counted during every transfer.

In the course of the repeated vial-to-vial transfers, a few flies escaped at various time points.

These flies were right censored for the survivorship analysis. For analysis of mean longevity,

vials where more than four flies escaped were removed from the final analysis. Thus mean

longevity analysis included data from 8-12 vials per treatment for each population. In this

study, we have focused on the response to selection on female longevity only.

Fecundity of the CE females were measured at regular intervals. Starting from day 5 post-

eclosion (i.e. day 14 post-egg collection), every fifth day, the flies were transferred into fresh

food vials (fecundity vials), where they were allowed a window of 17-hours before being

transferred to fresh food vials again. The number of eggs in these fecundity vials was

counted. This was done at day 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, and 60 (by this time-

point all females had died). For each time point, the number of eggs in a vial was divided by

the number of surviving females to calculate per female fecundity. This was taken as the unit

of analysis. However, the final analysis of age-specific fecundity was done excluding the last

few age points (post day 40) to avoid an unbalanced design arising due to the death of most

individuals towards the later age points. Additionally, the cumulative fecundity of females

was also quantified across all the above-mentioned age points till death.

<u>Data analysis:</u>

The entire study used a total of eight populations. Populations with replicate identity (for

example, BL_1 and YLB_1) were phylogenetically closer (see Figure S1) compared to those

with different replicate identity. In addition, during an assay a given replicate set (for

example, BL_1 and YLB_1) was handled together whereas different replicate sets were handled

independently, often on different days. Thus data from replicate sets were treated as statistical

blocks and was modelled as a random effect in all the analyses. The entire study, thus

involved four statistical blocks. Within each block, vials were the unit of replication and vial

values / means were used as the unit of analysis / response variable. Except for adult

survivorship and age-specific fecundity, results of all assays were analysed using mixed-

model analysis of variance (ANOVA). In case of the analysis of juvenile traits, selection

regime and sex (wherever applicable) were treated as fixed effects. For the early-life female

fecundity assay and egg size measurement, selection regime and live-yeast concentration

treatment were considered as fixed effects. For the analysis of the Experiment 2 early-life

progeny count data, male (selection) regime and female (selection) regime were used as fixed

effects. For analysing female mean longevity data from Experiment 3, selection regime and

mating status were treated as fixed effects. Cumulative life-time fecundity data was analysed

using selection regime as fixed factor. These analyses were performed on Statistica (version

10, Statsoft, Tulsa, OK, USA).

For female survivorship analysis, we applied a mixed Cox's Proportional Hazard models,

using the `coxme` package in R version 4.0.2 (Gupta et al. 2013, Therneau et al. 2015). The

effects of regime, mating status and their interaction on survivorship was analysed and block

was fitted as a random intercept. Cox partial likelihood estimates were compared across

treatment and selection regime.

Age-specific fecundity was analysed as linear mixed-effects models with the `lme4` package

in R version 4.0.2 (Bates et al. 2014). Further, to obtain the p-values for the `lmer` models

with degrees of freedom based on the Satterthwaite approximation, we used the R package

`lmerTest` (Kuznetsova et al. 2017). Regime and time-point (age) were fitted as categorical

fixed effects. Block as well as all interactions involving it were fitted as random effects. In

addition, vial ID (replicate vial identity) was nested within block and was fitted as a random

effect. The nested random effect accounts for the fact that subsequent egg counts from the

same group of flies within a particular vial are not independent (Harrison et al. 2018). Post-

hoc analysis was done using Tukey's HSD with the `emmeans` package (Lenth 2018). To

measure reproductive senescence, rates at which fecundity declined with age, after reaching a

peak, were quantified for both regimes. Rate of decline in fecundity was quantified across

seven time-points (τ), starting from the peak, i.e. day 11 post-eclosion until day 41. For a

particular age-point, the mean of per-capita fecundity (ε) of a population was calculated using

the observed fecundity across all the vials within that population. A linear model ($\varepsilon \sim b_p\tau +$

c_p, where b_p is population specific slope and c_p is population specific intercept) was fitted to

these data using the `lm` function in R to obtain a b-coefficient (i.e., slope) for each of the

eight populations (Fig. 3b). The slope of the regression line served as a proxy for rate of

reproductive senescence. Finally, these b-coefficients, without the sign, were analysed using

mixed model ANOVA in Statistica, taking selection regime as a fixed effect and block as a

random effect.

3. Results

<u>Experiment 1: Juvenile traits and female fecundity</u>

We did not find a significant effect of selection regime on juvenile traits including

development time, pre-adult (egg-to-adult) survivorship and dry body weight at the time of

eclosion (Table-S1 and S2). Analysis of the early-life fecundity data indicated significant

effects of the selection regime and yeast concentration, however, the effect of the selection

regime × yeast concentration interaction was not significant (Table 1). Compared to the BL

females, YLB females had 8.7-22.89 % higher mean per-capita fecundity across all live-yeast

concentrations in the assay (Fig.1). Moreover, the higher early-life fecundity output of YLB

females was consistent across a gradient of live-yeast concentration (Fig.1). The effect of

selection regime on egg size was non-significant (Table S1 and S2). Hence, there was no

evidence of differential provisioning in eggs produced by the experimental females.

Figure 1: Female early-life fecundity from Experiment 1. Number of eggs laid during a 16-hour assay window by 4-5 day

old (post-eclosion) females following the 48 hour live-yeast treatment was measured. This was done under three live-yeast

conditions - no yeast (N), limiting yeast (L) and high yeast (H). Data points represent mean per-capita fecundity of selected

(YLB) and the ancestral (BL) females. Six of the twelve test females in a vial were used for fecundity measurement. Mean

per-capita fecundity in a replicate vial was calculated using these six females. Three factor mixed model ANOVA indicated

significant effects of selection regime and live-yeast treatment. However, the effect of selection regime × live-yeast

treatment interaction was not significant.

<u>Experiment 2: Female × male effect on female fitness</u>

The analysis of the early-life progeny output data from Experiment 2 indicated a significant

effect female selection regime. However, neither the male selection regime nor the interaction

between these two fixed factors were found to have significant effects on progeny count

 (Table 1, Fig. 2). YLB females produced significantly more progeny irrespective of the

regime of the males they interacted with (Fig. 2). Thus, the higher early-life reproductive

output in YLB's seems to be independent of any male influence.

Figure 2: Female × male effect on early-life reproductive output in females as measured in Experiment 2. Number of

progeny produced during a 17-hour assay window by 4-5 day old (post-eclosion) BL and YLB females, mated and held with

BL and YLB males (full factorial combination), was measured. Data points represent per capita progeny output from

selected (YLB) and ancestral (BL) females. All six females in each replicate vial were allowed to produce progeny. Vials

mean, which was calculated over these six test females, was used as the unit of analysis. Two factor mixed model ANOVA

indicated a significant effect of female regime, but effect of male regime on progeny count was not significant, nor was that

of the male regime × female regime interaction.

Traits	Effects	SS	DF	MS	DEN DF	DEN MS	F	p
Early life fecundity	Selection Regime (SR)	947.8	1	947.8	3.01	7.47	126.83	**<0.01**
	Live-yeast concentration (YC)	51192.3	2	25596.2	6.00	1043.57	24.52	**<0.01**
	Block	19814.0	3	6604.7	5.59	1008.10	6.55	**0.03**
	SR × YC	23.0	2	11.5	6.01	42.40	0.27	0.77
	SR × Block	22.4	3	7.5	6.02	42.41	0.17	0.90
	YC × Block	6266.1	6	1044.4	6.00	42.39	24.63	**<0.01**
	SR × YC × Block	254.3	6	42.4	247.00	59.14	0.71	0.64
Progeny count	Female regime	411.13	1	411.14	3.00	7.40	55.55	**<0.01**
	Male regime	0.99	1	0.99	3.00	36.84	0.03	0.88
	Block	3149.10	3	1049.70	0.64	20.48	51.26	0.26
	Female regime × Male regime	7.18	1	7.18	3.00	23.76	0.30	0.62
	Female regime × Block	22.20	3	7.40	3	23.76	0.31	0.82
	Male regime × Block	110.51	3	36.84	3	23.76	1.55	0.36
	Fem. reg. × Male reg. × Block	71.28	3	23.76	222	20.71	1.15	0.33

356 **Table 1:** Results of the analyses on early-life fecundity (from experiment 1) and progeny output (from experiment 2) in

357 females. Fecundity data was analysed using three factor mixed-model ANOVA with selection regime and live-yeast

358 concentration as fixed factors, and block as random factor. Progeny count data from experiment 2 was analysed using three

359 factor mixed-model ANOVA with female regime and male regime as fixed factors, and block as random factor. Statistically

360 significant p-values are highlighted in boldface.

 Experiment 3: Longevity and age-specific fecundity

We did not find a significant effect of selection regime on survival probability (Table S3) and

mean longevity (Table S4) of females. Mating status, however, had a significant impact.

Virgin females had a better survival rate compared to the mated females (Fig. S2).

Unsurprisingly, analysis of the mean longevity indicated that mating status had a significant

effect on mean longevity (Table S4). Virgin females were found to have 67.58 % longer

mean longevity compared to the mated females for both BL and YLB selection regimes (Fig.

S3,Table S4). Results of these analyses can be found in the supplementary information.

Analysis of the cumulative fecundity of females quantified across different age points till

death revealed that there was no significant effect of selection regime on per-capita fecundity,

however, the effect of the selection regime × time point interaction was found to be

significant (Table 3a, Fig.3a). Tukey's adjusted p-value obtained by pairwise comparisons

suggested that, as compared to the ancestors, YLBs have significantly higher fecundity at the

age of 6th and 11th day PE. Interestingly, there was a clear trend of YLBs having higher mean

per-capita fecundity in initial age points compared to BLs (through age- 6, 11, 16, 21 and 26

days PE). The trend, however, was found to be reversed at the later time points, with BLs

showing comparatively higher fecundity (through age- 31, 36 and 41 days PE). The detailed

outcome of the post-hoc analysis is mentioned in the supplementary information

(Supplementary sheet, Table- S5). Further, a comparison of the slopes (b-coefficients) of the

linear regression fit of age-specific fecundity on time for the two regimes suggested a faster

rate of decline in fecundity in YLBs compared to that of the BLs (mean ±SE: b_{BL}= -2.03

±0.41; b_{YLB} = -3.36 ±0.13, Fig. 3b). Two factor mixed model ANOVA on the b-coefficient

indicated a significant effect of selection regime (Table 3b). Considering the b-coefficient to

be a measure of reproductive senescence, YLB females showed faster reproductive

senescence compared to that of the BL females. The cumulative fecundity of females,

quantified across all the above-mentioned age points, did not differ between the two regimes

(Table 3c, mean ±SE; BL: 129.22 ±4.52; YLB: 127.91 ±4.37).

Figure 3: Age-specific fecundity of females measured in Experiment 3. Number of eggs laid during 17 hour windows on

day 5, 10, 15, 20, 25, 30, 35, 40 were measured during the longevity assay. Only CE (continuously exposed) vials were used

for this purpose. Per capita fecundity was calculated for a vial by diving the fecundity count by the number of surviving

females, which was twelve to start with. (a) Mean age-specific per capita fecundity analysis indicated a significant effect of

selection regime × age interaction. (b) Slope (b-coefficient) of the linear decline in per capita fecundity with age following

the peak fecundity was computed for each population by fitting a linear model. The two factor mixed model ANOVA on the

b-coefficients indicated a significant effect of selection regime.

Trait	Effects	SS	DF	MS	Denominator DF	F	p
(a) Age-specific fecundity	Selection Regime (SR)	8.2	1	8.18	59.03	0.39	0.53
	Time point	6531.9	7	933.13	22.27	44.44	**< 0.01**
	SR × Time point	986.2	7	140.88	22.41	6.71	**<0.01**
(b) b-coefficient	Selection Regime	4.1	1	4.1	3	13.68	**0.03**
	Block	1.4	3	0.47	3	1.54	0.37
(c) Cumulative Fecundity	Selection Regime (SR)	35	1	35	3.01	0.06	0.82
	Block	4089	3	1363	3	2.26	0.26
	SR × Block	1806	3	602	76	0.73	0.53

Table 3: Results of the analyses of age-specific fecundity measured in Experiment 2. (a) Age-specific fecundity count was analysed using linear mixed-effect model, where selection regime and time point (age) were fitted as categorical fixed effects, and block as random effect. (b) For each population, age-specific fecundity data was fitted in a linear model ($\varepsilon \sim b_p\tau + c_p$, where b_p is population specific slope and c_p is population specific intercept) to compute b-coefficient (slope). These b-coefficient values (i.e., slope values without the sign) were analysed using two factor mixed model ANOVA with selection regime and block as fixed and random factors respectively. (c) Cumulative fecundity, i.e., fecundity count summed over all age points, were analysed using a two factor mixed model ANOVA with selection regime and block as fixed and random factors respectively. Statistically significant p-values are mentioned in bold face.

4. Discussion

In *D. melanogaster,* live-yeast is a strong stimulus for female fecundity both in wild and laboratory-adapted *D. melanogaster* (Begon 1982, Stramer 1982, Chippindale et al. 1993, Simmons & Bradley 1997, Anagnostou, Dorsch & Rohlfs 2010). In absence of live-yeast supplement females usually produce far fewer eggs. Therefore, in absence of live-yeast fecundity selection is expected to work on unstimulated fecundity of females. Selection imposed by such an ecology is expected to be female-specific at least to some extent. This is because adult females forage extensively on live-yeast, whereas males do not (Simmons & Bradley 1997). After 23 generations of laboratory evolution, we found clear evidence of increased early-life female fecundity. The results thus indicate the presence of substantial standing genetic variation in traits that determine unstimulated early-life fecundity in females. Several hundreds of generations of laboratory domestication, thus, have not eroded such variation. What maintains such standing genetic variation is thus an open question.

Interestingly, higher early-life fecundity of the evolved females was observed under unstimulated assay condition (i.e., in absence of yeast supplement) as well as low and high concentrations of yeast supplement. We also showed that this increased early-life fecundity is reflected on the reproductive output (i.e., progeny number) of the females. Taken together,

these results suggest that the YLB females have adapted to the live-yeast supplement

deprived ecology. Though males in *D. melanogaster* are known to affect female fecundity

(Wolfner et al. 2009), the results of experiment-3 rule out this possibility, further suggesting

that this to be a female adaptation. There are several non-mutually exclusive ways in which

YLB females could achieve higher early-life fecundity. In the following section, we explore

these different possibilities.

In principle, females can adapt to such nutritional deprivation either by increasing acquisition

and/or utilization of dietary protein. The selection did not affect body mass at eclosion and

juvenile traits, such as development time and egg-to-adult survivorship. Thus, there was no

indication of a change in juvenile stage resource acquisition. Alternatively, a change in the

relative abundance of different macronutrients, especially total protein and/or lipid content, in

the adult female body may also explain our results. This idea, however, remains to be tested

with additional experimental data. Theoretically, higher fecundity can be achieved by trading

off with per-capita provisioning, i.e., egg size (Hein et al. 2018, also see Vijendravarma et al.

2010). We, however, did not find any evidence supporting this number vs. size trade-off as

there was no effect of selection on egg size. This is perhaps not surprising as previous

investigations on *D. melanogaster* found egg size and number to be genetically uncorrelated

(Schwarzkopf et al. 1999).

Surprisingly, we did not find a negative impact of the increased early-life fecundity on

longevity, as predicted by the trade-off theory. A trade-off between reproduction and lifespan

is pervasive in nature and has been widely reported even in laboratory systems (Rose 1984,

Chippindale et al. 1993, Zwaan et al. 1995, Maklakov et al. 2017, Flatt 2011, Hensen et al.

2013). Several attempts to test this prediction, however, have been uninformative (Dudash &

Fenster 1997, Kotiaho & Simmons 2003). More recently, laboratory adaptation to chronic

dietary yeast deprivation resulted in *D. melanogaster* females with reduced lifespan without a

corresponding increase in reproductive output (Zajitschek et al. 2019). Thus, the lack of

evidence of reproduction-longevity trade-off in our populations is an addition to a growing

list of evidence challenging the generality of this theory (Le Bourg and Minois 1996,

Partridge et al. 2005, Grandison et al. 2009, Flatt 2011, Zajitschek et al. 2016, 2019).

Importantly, it is possible that this trade-off is more sensitive to total reproductive allocation.

If YLB females have evolved age-specific reproductive allocation without any change in total

lifetime reproductive investment, the expected decline in longevity may not evolve. This is

particularly important because there is no fitness consequence of altered longevity in

YLB/BL ecology and therefore, longevity is free to evolve only as a correlated response. On

the other hand, early life fecundity selection is a prime target of the YLB selection,

irrespective of the correlated consequences expressed later in life. Therefore, it is not

unreasonable to expect age-specific changes in fecundity to evolve in our selection regime.

We will return to this theory later in this section.

In addition to adult lifespan, changes in several adult traits may also result in increased

fecundity in females. There are at least four such possibilities, which are not mutually

exclusive. First, the simplest possibility is that an increased foraging behaviour can also result

in more resources being available for egg production (McCaffery 1975, Ives 1981, Barnes et

al. 2008). Secondly, a reduction in unnecessary wastage of resources may make more

resources available for egg production. For example, *D. melanogaster* females seem to

achieve higher fecundity by showing lower spontaneous locomotion (Long & Rice 2006).

Third, better resource recycling or reduced physiological wastage can potentially explain

increased fecundity (Adler & Bonduriansky 2014). For example, certain kinds of nutritional

deprivation seem to influence autophagy and scavenging of molecules – processes that may

be viewed as the physiology of resource recycling (Scott et al. 2004, Barth et al. 2011,

Cuervo & Macian 2012, Mason et al. 2018). More data are needed to test these theories. A

fourth way by which YLB females could have achieved higher fecundity is through an

alteration in the reproductive schedule without altering the overall lifetime reproductive

output – an idea we explore next.

The evolution of early life fitness components at the cost of late-life fitness has been one of

the central issues in life-history theory (Stearns 1992). In the laboratory ecology of 14-day

discrete generation, the YLB populations are selected for early-life fitness (see Promislow

and Tatar 1998, Houle and Rowe 2003). Hence, it is not surprising to find the evolution of

early-life fecundity in YLB females. The associated observation of a faster decrease in

fecundity without a significant change in mean lifetime egg output is somewhat surprising,

but it seems to fit with the antagonistic pleiotropy (AP) theory of ageing (Williams 1957).

Antagonistic pleiotropy theory has been in the centre stage especially due to its success in

explaining the evolution of organismal ageing. Early investigations

on *Drosophila* populations substantiated the theory (Rose 1984, Rauser et al. 2006), and have

been shown to explain both somatic as well as reproductive senescence. Multiple

investigations also reported a substantial genetic variation in age-specific fecundity in *D.*

melanogaster (for example, Tatar et al. 1996, Durham et al. 2014). More recently, using *C.*

elegans, Anderson et al. (2011) found that genetically heterogeneous populations selected for

early-life fecundity evolved increased early but reduced late-life reproductive output without

suffering a total fitness or lifespan cost. Thus, it is not unreasonable to suggest that adaptation

seen in our YLB populations involves the evolution of early-life beneficial alleles with

detrimental pleiotropic effects on late-life reproductive output.

5. Conclusion

Adaptive evolution of early-life fecundity in YLB females seems to support the hitherto reported phenomenon of the release of hidden genetic variation in novel ecology or stress (Badyaev 2005). Several past investigations have shown that novel environmental challenges or stresses (e.g., heat shock) can generate new trait variances or convert otherwise phenotypically neutral alleles to non-neutral (Hall 1990, Ratner et al. 1992, Kondrashov and Houle 1996, Wessler 1996, Leips & Mackay 2000, Queitsch et al. 2002, Dworkin et al. 2003, Nedelcu & Michod 2003, Ruden et al. 2003, Sollars et al. 2003). Such novel genetic variance can then be subjected to selective pressures leading to adaptation. It is possible that under LYD ecology, hidden variation in fecundity traits might have been expressed, followed by its directional selection in the YLB regime. Further, there is a growing literature on substantial G×E interaction on fitness-related traits, including fecundity (see Leroi et al. 1994, Fry et al. 1998, Burns et al. 2012, Pallares et al. 2021). Pallares et al. (2021) reported that dietary sugar enrichment has a profound effect on the fitness landscape of the fecundity and longevity traits in *D. melanogaster*. Fitness consequences of many fitness affecting alleles were found to be much stronger in the altered diet compared to the standard diet (Pallares et al. 2021). Our results could also be affected by a similar diet dependence of fitness consequence of fecundity traits in this system. Selectively neutral alleles that affect unstimulated (by live-yeast) fecundity could have strong fitness effects under LYD ecology and hence, be subjected to directional selection, at least in the initial generations of the YLB history. Many of these theories will need further investigation with genomic and transcriptomic data.

6. Acknowledgement

We thank Daniel E. L. Promislow and Benjamin Harrison of University of Washington for

their constructive criticism and suggestions on an earlier version of the manuscript.

Integrating their suggestions have greatly improved the quality of the manuscript, including

the clarity of the presentations. We are thankful to Tanya Verma for her help during the

experiments and population maintenance. The experiments reported here was supported by

the Core Research Grant (file no. CRG/2019/002460) from Science and Engineering

Research Board, Department of Science and Technology, Government of India. PD thanks

Council for Scientific and Industrial Research, Government of India for financial support in

the form of Junior and Senior Research Fellowship. NP thanks Summer Research Fellowship

Programme of Jawaharlal Nehru Centre for Advanced Scientific Research, Bengaluru for

financial assistance during his summer internship.

7. References

Adler, M. I., & Bonduriansky, R. (2014). Why do the well-fed appear to die young? A new

evolutionary hypothesis for the effect of dietary restriction on lifespan. *Bioessays*, *36*(5), 439-

450.

Andersen, L. H., Kristensen, T. N., Loeschcke, V., Toft, S., &Mayntz, D. (2010). Protein and

carbohydrate composition of larval food affects tolerance to thermal stress and desiccation in

adult *Drosophila melanogaster*. *Journal of insect physiology*, *56*(4), 336-340.

Anderson, J. L., Reynolds, R. M., Morran, L. T., Tolman-Thompson, J., & Phillips, P. C.

(2011). Experimental evolution reveals antagonistic pleiotropy in reproductive timing but not

life span in *Caenorhabditis elegans*. *Journals of Gerontology Series A: Biomedical Sciences

and Medical Sciences*, *66*(12), 1300-1308.

Anagnostou, C., Dorsch, M., &Rohlfs, M. (2010). Influence of dietary yeasts on *Drosophila

melanogaster* life-history traits. *EntomologiaExperimentalis et Applicata*, *136*(1), 1-11.

Arnqvist, G., & Nilsson, T. (2000). The evolution of polyandry: multiple mating and female

fitness in insects. Animal behaviour, 60(2), 145-164.

Barnes, A. I., Wigby, S., Boone, J. M., Partridge, L., & Chapman, T. (2008). Feeding,

fecundity and lifespan in female *Drosophila melanogaster*. *Proceedings of the Royal Society*

B: Biological Sciences, *275*(1643), 1675-1683.

Barth, J. M. I., Szabad, J., Hafen, E., & Köhler, K. (2011). Autophagy in *Drosophila* ovaries

is induced by starvation and is required for oogenesis. *Cell Death & Differentiation*, *18*(6),

915-924.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models

using lme4. *arXiv preprint arXiv:1406.5823*.

Begon, M., Ashburner, M., Carson, H. L., & Thompson, J. J. N. (1982). The genetics and

biology of *Drosophila.Academic Press, London, UK*, 345–384.

Becher, P. G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S. &

Bengtsson, M. (2012). Yeast, not fruit volatiles mediate *Drosophila melanogaster* attraction,

oviposition and development. *Functional Ecology*, *26*(4), 822-828.

Burns, J. G., Svetec, N., Rowe, L., Mery, F., Dolan, M. J., Boyce, W. T., & Sokolowski, M.

B. (2012). Gene–environment interplay in *Drosophila melanogaster*: Chronic food

deprivation in early life affects adult exploratory and fitness traits. *Proceedings of the*

National Academy of Sciences, *109*(Supplement 2), 17239-17244.

Chapman, T., Liddle, L. F., Kalb, J. M., Wolfner, M. F., & Partridge, L. (1995). Cost of

mating in Drosophila melanogaster females is mediated by male accessory gland products.

Nature, *373*(6511), 241-244.

Chapman, T., & Partridge, L. (1996). Female fitness in *Drosophila melanogaster*: an

interaction between the effect of nutrition and of encounter rate with males. *Proceedings of*

the Royal Society of London. Series B: Biological Sciences, *263*(1371), 755-759.

Chippindale, A. K., Leroi, A. M., Kim, S. B., & Rose, M. R. (1993). Phenotypic plasticity

and selection in Drosophila life-history evolution. I. Nutrition and the cost of reproduction.

Journal of Evolutionary Biology, 6(2), 171-193.

Cuervo, A. M., &Macian, F. (2012). Autophagy, nutrition and immunology. *Molecular*

aspects of medicine, *33*(1), 2-13.

Dudash, M. R., & Fenster, C. B. (1997). Multiyear study of pollen limitation and cost of

reproduction in the iteroparous Silene virginica. *Ecology*, *78*(2), 484-493.

Durham, M. F., Magwire, M. M., Stone, E. A., &Leips, J. (2014). Genome-wide analysis in

Drosophila reveals age-specific effects of SNPs on fitness traits. *Nature*

communications, *5*(1), 1-8.

Dworkin, I., Palsson, A., Birdsall, K., & Gibson, G. (2003). Evidence that Egfr contributes to

cryptic genetic variation for photoreceptor determination in natural populations of Drosophila

melanogaster. Current Biology, 13(21), 1888-1893.

Field, C. J., Johnson, I. R., &Schley, P. D. (2002). Nutrients and their role in host resistance

to infection. *Journal of leukocyte biology*, *71*(1), 16-32.

Flatt, T. (2011). Survival costs of reproduction in *Drosophila*. *Experimental*

gerontology, *46*(5), 369-375.

Fontana, L., Partridge, L., & Longo, V. D. (2010). Dietary restriction, growth factors and

aging: from yeast to humans. *Science (New York, NY)*, *328*(5976), 321.

Fontana, L., & Partridge, L. (2015). Promoting health and longevity through diet: from model

organisms to humans. *Cell*, *161*(1), 106-118.

Fry, J. D., Nuzhdin, S. V., Pasyukova, E. G., & Mackay, T. F. (1998). QTL mapping of

genotype–environment interaction for fitness in *Drosophila melanogaster*. *Genetics*

Research, *71*(2), 133-141.

Grandison, R. C., Piper, M. D., & Partridge, L. (2009). Amino-acid imbalance explains

extension of lifespan by dietary restriction in *Drosophila. Nature*, *462*(7276), 1061-1064.

Gupta, V., Ali, Z. S., & Prasad, N. G. (2013). Sexual activity increases resistance against

Pseudomonas entomophila in male *Drosophila melanogaster*. *BMC Evolutionary*

Biology, *13*(1), 1-11.

Hall, B. G. (1990). Spontaneous point mutations that occur more often when advantageous

than when neutral. Genetics, 126(1), 5-16.

Harrison, X. A., Donaldson, L., Correa-Cano, M. E., Evans, J., Fisher, D. N., Goodwin, C. E.,

... & Inger, R. (2018). A brief introduction to mixed effects modelling and multi-model

inference in ecology. *PeerJ*, *6*, e4794.

Hansen, M., Flatt, T., & Aguilaniu, H. (2013). Reproduction, fat metabolism, and life span:

what is the connection? *Cell metabolism, 17*(1), 10-19.

Hein, N., Brendel, M. R., Feilhauer, H., Finch, O. D., & Löffler, J. (2018). Egg size versus

egg number trade-off in the alpine-tundra wolf spider, Pardosa palustris (Araneae:

Lycosidae). Polar Biology, 41(8), 1607-1617.

Hoffmann, A. A., & Harshman, L. G. (1999). Desiccation and starvation resistance in

Drosophila: patterns of variation at the species, population and intrapopulation

levels. *Heredity*, *83*(6), 637-643.

Houle, D., & Rowe, L. (2003). Natural selection in a bottle. The American Naturalist, 161(1),

50-67.

Ives, P. M. (1981). Feeding and egg production of two species of coccinellids in the

laboratory. *The Canadian Entomologist*, *113*(11), 999-1005.

Jin, K., Wilson, K. A., Beck, J. N., Nelson, C. S., Brownridge III, G. W., Harrison, B. R., ...

& Promislow, D. (2020). Genetic and metabolomic architecture of variation in diet

restriction-mediated lifespan extension in Drosophila. PLoS genetics, 16(7), e1008835.

Kassambara, A., Kosinski, M., Biecek, P., & Fabian, S. (2017). Package

'survminer'. *Drawing Survival Curves using 'ggplot2'(R package version 03 1)*.

Kondrashov, A. S., & Houle, D. (1994). Genotype—environment interactions and the

estimation of the genomic mutation rate in Drosophila melanogaster. Proceedings of the

Royal Society of London. Series B: Biological Sciences, 258(1353), 221-227.

Kolss, M., Vijendravarma, R.K., Schwaller, G.& Kawecki, T.J.(2009). Life-history

consequences of adaptation to larval nutritional stress in *Drosophila. Evolution63,*2389–

2401.

Kotiaho, J. S., & Simmons, L. W. (2003). Longevity cost of reproduction for males but no

longevity cost of mating or courtship for females in the male-dimorphic dung beetle

Onthophagusbinodis. Journal of Insect Physiology, *49*(9), 817-822.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. (2017). lmerTest package: tests in

linear mixed effects models. *Journal of statistical software*, *82*(1), 1-26.

Le Bourg, E., &Minois, N. (1996). Failure to confirm increased longevity in *Drosophila*

melanogaster submitted to a food restriction procedure. *The Journals of Gerontology Series*

A: Biological Sciences and Medical Sciences, *51*(4), B280-B283.

Leips, J., & Mackay, T. F. (2000). Quantitative trait loci for life span in Drosophila

melanogaster: interactions with genetic background and larval density. *Genetics, 155*(4),

1773-1788.

Lenth, R., Singmann, H., Love, J., Buerkner, P., &Herve, M. (2018). Emmeans: Estimated

marginal means, aka least-squares means. *R package version*, *1*(1), 3.

Leroi, A. M., Chippindale, A. K., & Rose, M. R. (2004). Long-term laboratory evolution of a

genetic life-history trade-off in *Drosophila melanogaster*. 1. The role of genotype-by-

environment interaction. In *Methuselah Flies: A Case Study in the Evolution of Aging* (pp.

26-39).

Long, T. A., & Rice, W. R. (2007). Adult locomotory activity mediates intralocus sexual

conflict in a laboratory-adapted population of *Drosophila melanogaster. Proceedings of the*

Royal Society B: Biological Sciences, *274*(1629), 3105-3112.

Mair, W., Piper, M. D. W., & Partridge, L. (2005). Calories do not explain extension of life

span by dietary restriction in *Drosophila. PLoS biology*, *3*(7), e223.

Maklakov, A. A., Carlsson, H., Denbaum, P., Lind, M. I., Mautz, B., Hinas, A., &Immler, S.

(2017). Antagonistically pleiotropic allele increases lifespan and late-life reproduction at the

cost of early-life reproduction and individual fitness. *Proceedings of the Royal Society B:*

Biological Sciences, *284*(1856), 20170376.

Marden, J. H., Rogina, B., Montooth, K. L., & Helfand, S. L. (2003). Conditional tradeoffs

between aging and organismal performance of Indy long-lived mutant flies. *Proceedings of*

the National Academy of Sciences, *100*(6), 3369-3373.

Mason, J. S., Wileman, T., & Chapman, T. (2018). Lifespan extension without fertility

reduction following dietary addition of the autophagy activator Torin1 in *Drosophila*

melanogaster. PloS one, *13*(1), e0190105.

McCaffery, A. R. (1975). Food quality and quantity in relation to egg production in

*Locustamigratoria*migratorioides. *Journal of Insect Physiology*, *21*(9), 1551-1558.

Miller, C. V., & Cotter, S. C. (2018). Resistance and tolerance: the role of nutrients on

pathogen dynamics and infection outcomes in an insect host. *Journal of Animal*

Ecology, *87*(2), 500-510.

Moatt, J. P., Savola, E., Regan, J. C., Nussey, D. H., & Walling, C. A. (2020). Lifespan

extension via dietary restriction: time to reconsider the evolutionary mechanisms?.

BioEssays, 42(8), 1900241.

Nandy, B., Dasgupta, P., Halder, S., &Verma, T. (2016). Plasticity in aggression and the

correlated changes in the cost of reproduction in male *Drosophila melanogaster. Animal*

Behaviour, 114, 3-9.

Nedelcu, A. M., & Michod, R. E. (2003). Sex as a response to oxidative stress: the effect of

antioxidants on sexual induction in a facultatively sexual lineage. *Proceedings of the Royal*

Society of London. Series B: Biological Sciences, 270(suppl_2), S136-S139.

Nakagawa, S., Lagisz, M., Hector, K. L., & Spencer, H. G. (2012). Comparative and meta-

analytic insights into life extension via dietary restriction. *Aging cell, 11*(3), 401-409.

Pallares, L. F., Lea, A. J., Han, C., Filippova, E. V., Andolfatto, P., &Ayroles, J. F. (2021).

Diet unmasks genetic variants that regulate lifespan in outbred *Drosophila. bioRxiv*, 2020-10.

Parkash, R., & Munjal, A. K. (2000). Evidence of independent climatic selection for

desiccation and starvation tolerance in Indian tropical populations of *Drosophila*

melanogaster. Evolutionary Ecology Research, 2(5), 685-699.

Partridge, L., Green, A., & Fowler, K. (1987). Effects of egg-production and of exposure to

males on female survival in *Drosophila melanogaster. Journal of Insect Physiology, 33*(10),

745-749.

Partridge, L., Piper, M. D., &Mair, W. (2005). Dietary restriction in*Drosophila. Mechanisms*

of ageing and development, 126(9), 938-950.

Promislow, D. E., & Tatar, M. (1998). Mutation and senescence: where genetics and

demography meet. *Genetica, 102*, 299-314.

Queitsch, C., Sangster, T. A., & Lindquist, S. (2002). Hsp90 as a capacitor of phenotypic

variation. *Nature, 417*(6889), 618-624.

Ratner, V. A., Zabanov, S. A., Kolesnikova, O. V., & Vasilyeva, L. A. (1992). Induction of

the mobile genetic element Dm-412 transpositions in the Drosophila genome by heat shock

treatment. Proceedings of the National Academy of Sciences, 89(12), 5650-5654.

Rauser, C. L., Tierney, J. J., Gunion, S. M., Covarrubias, G. M., Mueller, L. D., & Rose, M.

R. (2006). Evolution of late-life fecundity in *Drosophila melanogaster*. *Journal of*

Evolutionary Biology, *19*(1), 289-301.

Rikke, B. A., Liao, C. Y., McQueen, M. B., Nelson, J. F., & Johnson, T. E. (2010). Genetic

dissection of dietary restriction in mice supports the metabolic efficiency model of life

extension. Experimental gerontology, 45(9), 691-701.

Rion, S., & Kawecki, T. J. (2007). Evolutionary biology of starvation resistance: what we

have learned from *Drosophila*. *Journal of evolutionary biology*, *20*(5), 1655-1664.

Rohlfs, M., & Hoffmeister, T. S. (2005). Maternal effects increase survival probability in

Drosophila subobscura larvae. *EntomologiaExperimentalis et Applicata*, *117*(1), 51-58.

Rose, M. R. (1984). Laboratory evolution of postponed senescence in *Drosophila*

melanogaster. *Evolution*, 1004-1010.

Ruden, D. M., Garfinkel, M. D., Sollars, V. E., & Lu, X. (2003, October). Waddington's

widget: Hsp90 and the inheritance of acquired characters. In Seminars in cell &

developmental biology (Vol. 14, No. 5, pp. 301-310). Academic Press.

Schwarzkopf, L., Blows, M. W., & Caley, M. J. (1999). Life-history consequences of

divergent selection on egg size in Drosophila melanogaster. The American Naturalist, 154(3),

333-340.

Scott, R. C., Schuldiner, O., & Neufeld, T. P. (2004). Role and regulation of starvation-

induced autophagy in the *Drosophila* fat body. *Developmental cell*, *7*(2), 167-178.

Shanley, D. P., & Kirkwood, T. B. (2000). Calorie restriction and aging: A life-history

analysis. Evolution, 54(3), 740-750.

Sisodia, S., & Singh, B. N. (2012). Experimental evidence for nutrition regulated stress

resistance in *Drosophila ananassae*.*PLoS ONE 7*(10): e46131.

Simmons, F. H., & Bradley, T. J. (1997). An analysis of resource allocation in response to

dietary yeast in *Drosophila melanogaster*. *Journal of insect physiology*, *43*(8), 779-788.

Sollars, V., Lu, X., Xiao, L., Wang, X., Garfinkel, M. D., & Ruden, D. M. (2003). Evidence

for an epigenetic mechanism by which Hsp90 acts as a capacitor for morphological evolution.

Nature genetics, 33(1), 70-74.

Starmer, W. T. (1981). A comparison of *Drosophila* habitats according to the physiological

attributes of the associated yeast communities. *Evolution*, 38-52.

Stearns, S.C. (1992) The evolution of life histories. *Oxford University Press, Oxford*.

Tatar, M., & Carey, J. R. (1995). Nutrition mediates reproductive trade-offs with age-specific

mortality in the beetle *Callosobruchus maculatus*. *Ecology*, *76*(7), 2066-2073.

Tatar, M., Promislow, D. E., Khazaeli, A. A., & Curtsinger, J. W. (1996). Age-specific

patterns of genetic variance in Drosophila melanogaster. II. Fecundity and its genetic

covariance with age-specific mortality. Genetics, 143(2), 849-858.

Therneau, T. (2015). Mixed effects Cox models. *CRAN repository*.

Vijendravarma, R. K., Narasimha, S., & Kawecki, T. J. (2010). Effects of parental larval diet

on egg size and offspring traits in Drosophila. *Biology letters, 6(2)*, 238-241.

Vijendravarma, R. K., & Kawecki, T. J. (2013). Epistasis and maternal effects in

experimental adaptation to chronic nutritional stress in *Drosophila*. *Journal of evolutionary*

biology, *26*(12), 2566-2580.

Weldon, C. W., Mnguni, S., Démares, F., du Rand, E. E., Malod, K., Manrakhan, A., &

Nicolson, S. W. (2019). Adult diet does not compensate for impact of a poor larval diet on

stress resistance in a tephritid fruit fly. *Journal of Experimental Biology*, *222*(6), jeb192534.

Wessler, S. R. (1996). Turned on by stress. Plant retrotransposons. *Current biology: CB, 6*(8),

959-961.

Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of

senescence. *evolution*, 398-411.

Wolfner, M. F. (2009). Battle and ballet: molecular interactions between the sexes in

Drosophila. Journal of Heredity, 100(4), 399-410.

Zajitschek, F., Zajitschek, S. R., Canton, C., Georgolopoulos, G., Friberg, U., &Maklakov, A.

A. (2016). Evolution under dietary restriction increases male reproductive performance

without survival cost. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1825),

20152726.

Zajitschek, F., Georgolopoulos, G., Vourlou, A., Ericsson, M., Zajitschek, S. R., Friberg, U.,

&Maklakov, A. A. (2019). Evolution under dietary restriction decouples survival from

fecundity in *Drosophila melanogaster* females. *The Journals of Gerontology: Series*

A, *74*(10), 1542-1548.

Zwaan, B., Bijlsma, R., & Hoekstra, R. F. (1995). Direct selection on life span in *Drosophila*

melanogaster. *Evolution*, *49*(4), 649-659