Evolution of a novel female reproductive strategy in *Drosophila melanogaster*

populations subjected to long term protein restriction

Running title: Female adaptation to yeast deprivation

Purbasha Dasgupta[1], Subhasish Halder[1], Debapriya Dari[1], Nabeel P.[1], Sai Samhitha Vajja[1, 2]

Bodhisatta Nandy*[1]

Affiliation:

[1]Indian Institute of Science Education and Research Berhampur, Transit Campus, Govt. ITI Building,

NH 59, Engineering School Junction, Ganjam, Berhampur 760 010, Odisha, India.

*Corresponding author: nandy@iiserbpr.ac.in, contact number: +91-680-222 7755

Key words: Nutritional deprivation, fitness trade-offs, antagonistic pleiotropy, experimental

evolution.

Word count: 5,974 excluding abstract, table and figure captions, acknowledgement and

references.

Figure count: 3

Table count: 2

Supplementary information: A separate file that includes three figures and six tables.

Author contributions: BN conceived the idea, designed the experiment, analysed and

interpreted the data, wrote the manuscript. PD designed and performed the experiments,

analysed data, prepared the manuscript. SH, DD, NP, and SSV performed various assays and

collected data. BN and PD revised the manuscript and wrote the responses to reviewers'

comments.

Abstract

Reproductive output is often constrained by availability of macronutrients, especially protein.

Long term protein restriction, therefore, is expected to select for traits that maximize

reproductive output in spite of such nutritional challenge. We subjected four replicate

populations of *Drosophila melanogaster* to a complete deprivation of live-yeast supplement,

thereby mimicking a protein restricted ecology. As yeast supplement is a key source of

protein, such an ecology is expected to strongly limit reproductive output, especially in

females. Following 24 generations of experimental evolution, compared to their matched

controls, females from experimentally evolved populations showed increase in reproductive

output early in life, both in presence and absence of yeast supplement. The observed increase

in reproductive output was not associated with any accommodating alteration in egg size; and

development time, pre-adult survivorship, and body mass at eclosion of the progeny. There

was no evidence for evolution of lifespan and lifelong cumulative reproductive output in

females. However, females from experiment regime were found to have a significantly faster

rate of reproductive senescence, as indicated by a faster rate of age related decline in

reproductive output following the attainment of the reproductive peak. Therefore, adaptation

to yeast deprivation ecology in our study involved a novel reproductive strategy whereby

females attained higher reproductive output early in life followed by faster reproductive

aging. To the best of our knowledge, this set of results is one of the most clear demonstration

of optimization of fitness by fine tuning of reproductive schedule during adaptation to a

prolonged nutritional deprivation.

1. Introduction

The ability of an organism to reproduce in spite of nutritional challenges is essential to adaptation in a given ecology (Kolss et al 2009, Zhang et al. 2019). Beyond providing the essential energy to sustain life, nutrients, especially macronutrients such as proteins and carbohydrates, determine survival by affecting resistance to stress and other ecological challenges, including pathogens (Chippindale 1993, Field et al. 2002, Andersen et al. 2010, Sisodia and Singh 2012, Miller and Cotter 2017, Weldon et al. 2019). Access to macronutrients also affect reproductive output and is central to a variety of trade-offs concerning reproduction (Chippindale 1993, Tatar & Carey 1995, Chapman & Partridge 1996, Marden et al. 2003, Adler & Bonduriansky 2014). This is evident in response to a form of nutritional challenge, wherein one or more essential macronutrients is restricted without causing malnutrition, organisms have been found to elicit a complex physiological response that culminates in the partial or complete shutdown of reproduction, resulting in an extension of lifespan (Mair et al. 2005, Partridge et al. 2005, Nakagawa et al. 2012, Fontana & Partridge 2015). Temporarily shutting down reproduction could be advantageous if it allows the organism to tide over the period of nutritional shortage, following which reproduction can resume. However, such a reproductive strategy is expected to be detrimental if nutritional shortage is long lasting, spanning generations. There is no reason to not expect such long term nutritional shortage. It is not clear how organisms adaptively optimize reproduction when faced with such selective pressure (Adler & Bonduriansky 2014). Perhaps the most famous example of adaptation to such dietary shortage is the evolution of Finch populations in the Galapagos archipelago following an intense selection imposed by a prolonged period of draught that led to a severe qualitative and quantitative depletion of the food supply, resulting in strong natural selection (Boag & Grant 1981). It resulted in a complex evolutionary response in morphology, physiology and behaviour of the finch populations (for

example, see Schluter & Grant 1984). Thus, undoubtedly diet can drive adaptive evolution.

Recent studies on nutritional deprivation have shown that deprivation of specific

macronutrients, especially protein, can pose specific effects on fitness components – limiting

reproduction (Maklakov et al. 2008, Lee et al. 2008, Fanson et al. 2009, Fanson and Taylor

2012, Short et al. 2020), stress resistance (Sisodia & Singh 2012), memory and learning

(Mery & Kawecki 2005). However, surprisingly little is known about how populations

evolving under nutritional deprivation, specifically protein deprivation, optimize

reproduction. In addition, there is also a serious dearth of literature on genetic variance and

covariance for traits that could allow or impede such adaptation (but see Rikke et al. 2010, Jin

et al. 2020).

Apart from other model and non-model organisms, *Drosophila melanogaster* has been

extensively used to investigate the plasticity in life history and the underlying physiological

response to dietary restriction. Among the macronutrients, while carbohydrate is the primary

caloric source, protein is a major determinant of reproductive output. Both in laboratory

cultures and in wild populations, Drosophilid fruit flies acquire protein by foraging on yeast

(Begon 1982, Starmer 1982, Chippindale et al., 1993; Anagnostou, Dorsch & Rohlfs 2010).

Restricted access to yeast, therefore, lead to protein starvation, and is known to result in

reduced reproductive output, especially in females (Lee et al. 2008). Yeast deprivation in

laboratory systems of *Drosophila* sp. is usually achieved either by limiting yeast supplement

on food (Partridge et al. 1987, Chippindale et al.1993) or by reducing the amount of yeast in

the food (Chapman and Partridge, 1996) – both leading to reduced fecundity in females. Such

reduction in reproduction, triggered by dietary restriction, has been frequently found to be

accompanied with increased lifespan, inspiring a large body of experimental research. There

is, now, ample evidence to suggest that such lifespan extension is almost entirely due to yeast

restriction and not caloric or carbohydrate restriction (Zimmerman et al. 2003, Mair et al.

2005, Piper et al. 2005). Further, alteration in the dietary yeast to sucrose ratio, within certain

limits, without changing the caloric value of food has been found to generate a physiological

shift away from reproduction (Lee et al. 2008). Other nutritional stresses, including extreme

starvation, have also been shown to lead to a physiological shift resulting in extension of

lifespan, often at the cost of reduced fecundity in females (Hoffmann & Harshman 1999,

Parkash & Munjal 2000, Partridge et al. 2005, Rion & Kawecki 2007). Although adaptive

during a transient nutritional deprivation (Shanley & Kirkwood 1999, Adler & Bonduriansky

2014, Moatt et al. 2020), such a physiological response of diversion of resources away from

reproduction is an unlikely solution for prolonged nutritional scarcity. In such case, natural

selection may favour a higher investment in reproduction even at the cost of survival,

especially for short lived organisms (Williams 1957, Bonduriansky et al. 2008).

Experimental evolution using fruit flies has been a powerful approach to study a variety of

evolutionary genetic problems, including adaptation to nutritional challenges (Prasad and

Joshi 2003, Garland and Rose 2009). Kolss et al. (2009) showed that *D. melanogaster*

populations can adapt to nutritional deprivation by evolving better egg-adult survivorship and

faster development. More recent studies investigated adaptation to yeast-restricted adult diet

to find that males under such a regime evolve higher reproductive output on a DR food

without any significant change in lifespan (Zajitschek, 2016). However, the females in the

same study on the other hand, evolved shorter lifespan with no alteration in their reproductive

output (Zajitschek, 2019). Importantly, the outcome of experimental evolution critically

depends on the specific context of the experimental selection pressure and the definition of

fitness in the experimental population (Rose et al. 1996, Garland and Rose 2009). Hence, the

issue of reproductive adaptation under nutritional deprivation is only beginning to emerge as

a research programme with plenty of open questions.

Here we investigated adaptation of *D. melanogaster* populations to yeast deprivation

specifically targeting yeast supplement in their diet instead of heat-killed yeast in their food.

Importantly, besides affecting female reproductive output, yeast supplement is known to

impact adult lifespan and starvation resistance (Chippindale et al. 1993), oviposition site

preference (Becher et al 2012), and may even affect offspring survival and fitness

components (Rohlfs & Hoffmeister 2005, Anagnostou, Dorsch & Rohlfs 2010). Given the

strong fitness consequence of access to yeast supplement, the lack of experimental data on

adaptation to yeast deprivation in the available literature is rather surprising. Interestingly,

because yeast acts as dietary protein supplement primarily for females (Chippindale et al.

1993, Nandy et al. 2012), it is possible for yeast deprivation to represent a sex-specific

selective environment, which can lead to sex specific adaptations as found by Zajitschek et al.

(2016, 2019).

To investigate adaptation to yeast deprivation, we subjected four replicate populations of *D.*

melanogaster to a complete withdrawal of yeast supplement from their diet. After 23

generations of experimental evolution, we investigated the effect of experimental evolution

on (a) female fecundity in absence of yeast supplement as well as under varying access to the

same, (b) a measure of female fitness, (c) pre-adult traits that can affect female fecundity, and

(d) longevity and age specific reproductive output in females. Keeping the adaptation theory

in mind, we predicted female fecundity and reproductive output to improve in the selected

populations compared to the ancestral base populations, triggering correlated changes in the

life history.

2. Materials and methods

<u>Study populations and selection regime</u>

Four new experimental populations (referred to as YLB_{1-4} here on) were derived from four

replicate baseline populations BL_{1-4} (Figure S1). The BL populations (Nandy et al. 2016) are

outbred, wild type populations of *D. melanogaster*. They are maintained under a 14-day

discrete generation cycle, under 24-hour light, at 25 °C ($\pm$1) ambient temperature, on a

standard banana-sugarcane jaggery-yeast medium, with a population size of ~2,800. Food is

prepared by boiling mashed ripe banana, barley, sugarcane jaggery, and baker's yeast in

water. Agar-agar is used as solidifying agent and p-hydroxybenzoic acid dissolved in ethanol

is used as a preservative. The complete recipe with the cooking method can be found in

Supplementary information (Table S1). Flies are grown in standard vials (25 mm diameter $\times$

96 mm height) for 12 days. Larval density is controlled at ~70 per vial with 8 ml food and 40

such vials are maintained for each replicate population. On day 12 of the generation cycle,

~2,800 adults are transferred to each plexiglass cage (23cm $\times$ 20cm $\times$ 15 cm) with a food

plate smeared with an *ad-lib* quantity of live yeast (a paste made with water). On day 14 post-

egg collection, the flies are given fresh food for oviposition and eggs laid during an 18-hour

window (fitness window) are collected to start the next generation. BL populations were

maintained in the laboratory under this regime for almost five years prior to the derivation of

the experimental YLB populations. From each of the four replicate populations of BL (BL_{1-}

$_4$), a corresponding YLB population (YLB_{1-4}) was derived by collecting 40 vials of eggs with

a standard egg density (i.e., approximately 2800 eggs were used to found each YLB

population). The maintenance of the YLB populations is identical to that of the BL except for

the absence of live-yeast supplement in the YLB ecology. All assays reported here were

conducted after 23-25 generations of experimental evolution. To equalize any potential non-

genetic parental effect, all populations were passed through one generation of common

rearing ecology (Rose 1984). During this, a subset of the experimental populations was

generated by collecting eggs from the stock BL_x and YLB_x populations. These populations

were maintained under a regime that exactly matched that of the BL-regime (i.e. both BL and

YLB were provided with *ad-lib* live-yeast). Experimental flies were generated by collecting

eggs from these subset populations. During the assays, all fly handling, including the

collection of virgin flies, wherever necessary, was done under light CO_2 anaesthesia. A

schematic representation of the overall experimental plan can be found in the supplementary

information (Fig. S2).

Experiment 1: Juvenile traits and early-life fecundity in females

After 23 generations of experimental evolution, we assessed whether the YLB populations

have adaptively diverged from the ancestral BL populations. Our initial observation

suggested fecundity to be the most likely target trait of selection (data not shown). Therefore,

as the first set of assays, we quantified female early-life fecundity under a range of yeast

concentrations. In addition, since fecundity is often a function of body size and related

juvenile traits, we measured these traits.

Female early-life fecundity and three juvenile traits – pre-adult (egg-to-adult) development

time, juvenile (egg-to-adult) survivorship and dry weight at eclosion were measured between

23-25 generations. Following one generation of common rearing (see above), eggs were

collected from the population cages in exact numbers. For egg-collection, two small food

plates were introduced in a cage, and were left undisturbed for about one hour to allow

oviposition. Such a short window of oviposition ensured the developmental synchrony of the

eggs collected for the experiment. Eggs were randomly scooped out from these oviposition

plates, and transferred to an egg-collection surface made of solidified agar gel (1% Agar-agar

solution) using a fine brush. Eggs were carefully counted and a small piece of Agar gel

containing 70 eggs was cut out and transferred to a fresh food vial having 8ml food. Ten such

vials were set up for each population. Each vial was labelled individually and placed into the

incubator. As an added precaution, the vials in the incubator were shuffled twice every day to

randomize any positional effect. The vials were monitored carefully for the signs of eclosion.

Beginning from the eclosion of the first individual, the eclosing adults were counted and

sexed every 6 hours until all the pupae eclosed. This data was used to derive the average

measure of egg-to-adult development time (i.e., pre-adult development time) for males and

females in each vial. Out of the 70 eggs initially cultured, the proportion that finally eclosed

from a vial was considered as the measure of egg-to-adult survivorship (i.e., juvenile

survivorship). Three vials (one from the YLB_1; two from BL_2) were excluded from the

analysis due to mishandling. For each population (BL_x and YLB_x), 50 males and 50 females

were collected for dry whole body weight measurement. They were collected within 6-hours

of eclosion, and were immediately frozen at -20 °C. They were later dried at 60 °C for 48 hr

and weighed in groups of five using Shimadzu AUW220D to the nearest 0.01 mg. The mean

body weight of each group was calculated and taken as a unit of analysis.

For early-life fecundity assay, flies were raised and collected as virgins as mentioned above.

Virgin flies were held in same-sex vials in groups of 12 in each vial. On the day12 post-egg

collection (i.e., corresponding to 2-3 day post-eclosion), one vial each of males and females

(i.e., 12 males and 12 females) belonging to the same regime, were transferred to a fresh food

vial assigned to one of the three live-yeast supplementation treatments - No yeast

(N), Limiting yeast (L: each vial containing 4.5 mg yeast suspended in 50 μl of water)

and High yeast (H: each vial containing 9 mg yeast suspended in 50 μl of water). Twelve

such vials were set up per yeast supplementation treatment for a population. For setting up L

and H treatments, yeast was dissolved in tap water to make a suspension. It was then

dispensed at a point on the rim of the food meniscus in a vial to ensure unfettered access to

both food and the live-yeast . The suspension was allowed to air-dry before the introduction

of the experimental flies. The males and the females were allowed to interact in these vials

for 48 hours. Following this, on day 4-5 post-eclosion, flies in each treatment vial were

anaesthetized under light CO_2 -six females were randomly separated and transferred to a vial

for egg size measurement, while the other six were individually transferred to six oviposition

tubes (12×75 mm) for fecundity measurement. For the former, females were allowed to

oviposit for 16 hours in a vial, following which they were discarded. The vial with eggs was

frozen at -20 °C for a minimum of one day. For egg size measurement, eggs were mounted

on a glass slide on their dorsal side and photographed using a stereozoom microscope and the

area of the two-dimensional elliptical outline of the eggs were measured in Image J software.

This area was taken as a proxy for the size of each egg. A given egg was measured three

times and the average of these three measurements was taken as the unit of analysis. Females

were allowed to oviposit in these tubes for 16 hours. Females from the oviposition tubes were

then discarded and number of eggs was counted. We removed a few vials from the analysis

due to a variety of handling errors, including escaped flies and broken tube/vial, during the

assay. The final number of vials used for the analysis ranged from 9-12 vials per live-yeast

treatment for a population.

<u>Experiment 2: Female × male effect on female fitness</u>

To further assess the response of female reproductive output, and eventually fitness, we

quantified female early-life reproductive output (progeny count = number of surviving adult

offspring produced by a female) during a 16-hour window on the day 4-5 post-eclosion.

Given that most of the flies eclose during day9-10 (i.e., 9-10[th] day of the generation cycle

starting from egg stage), this window corresponded to the 18-hour fitness window on the 14[th]

day of the generation cycle and therefore, is a good measure of Darwinian fitness in this

system (see Rice et al. 2006 for the measure of fitness in laboratory adapted populations).

Importantly, female fitness, especially in *D. melanogaster*, is affected by both female

fecundity and male factors such as seminal fluid composition (Chapman et al. 1995; Arnqvist

& Nilsson 2000; Wolfner 2009). Therefore, it is important to assess male and female specific

effect of selection on female fitness. To test this effect, we combined YLB and BL males and

females in a full factorial design and measured early-life reproductive output (i.e., progeny

count) of females in the above-mentioned protocol. These assays were carried out after 26-29

generations of selection. Experimental flies were generated following a protocol similar to

that followed in Experiment 1. Males and females were collected as virgins (on around day 9-

10 post-egg collection) and held in single-sex vials in groups of six in each vial until day 12

post-egg collection. On the day12 of the generation cycle (i.e., 2-3 days post-eclosion), sexes

were combined to set up adult interaction vials having six individuals of each sex in a vial.

Four types of combinations (viz., female×male: BL×YLB; BL×BL; YLB×YLB; YLB×BL),

each having 15 replicate vials, were set up. A few flies escaped during the setting up of one

vial in the third block with $YLB_3 \times YLB_3$ combination and hence, it was removed from the

experiment. These adult interaction vials were left undisturbed for 48 hours. On the day 14,

females from each of these vials were sorted under light CO_2anaesthesia, and individually

transferred to oviposition test tubes (one female per tube) where they were allowed to

oviposit for 17 hours. After 17 hours, females from the oviposition tubes were discarded and

the tubes were kept in the incubator for the development and emergence of the progeny. After

the emergence of all the progeny, the test tubes were frozen at -20 °C for the final count.

Experiment 3: Longevity and age-specific fecundity

To test reproduction vs. lifespan trade-off theory, longevity and fecundity of the experimental

females were measured after 26 generations of selection. To assess age related changes in

fecundity, we measured fecundity of the test females in regular intervals. Similar to the other

assays described above, all populations were passed through one generation of common

rearing. The eggs were collected from the common rearing cages and cultured under standard

density of ~70 per 6-8ml food in a vial. Adult flies were collected as virgins within six hours

of eclosion and held in single-sex vials in groups of 12 in standard food vials. These vials

were left undisturbed for two days. On the day 3 post eclosion (i.e., on the day 12 post egg

collection), experimental vials were set up by transferring the flies into fresh food vials. Two

sets of vials were set up for each population – (a) Continuously exposed (CE): 12 males and

12 females placed (without anaesthesia) in a vial and (b) Virgin females (VF):12 virgin

females placed in a vial. For a population, 12 vials were set up for each of CE and VF. Except

for the fecundity count days (see below), flies were transferred into fresh food vials every

alternate day without anaesthesia. Dead flies were sexed and counted during every transfer.

In the course of the repeated vial-to-vial transfers, a few flies escaped at various time points.

These flies were right censored for the survivorship analysis. For analysis of mean longevity,

vials where more than four flies escaped were removed from the final analysis. Thus mean

longevity analysis included data from 8-12 vials per treatment for each population. In this

study, we have focused on the response to selection on female longevity only.

Fecundity of the CE females were measured at regular intervals. Starting from day5 post-

eclosion (i.e. day 14 post-egg collection), every fifth day, the flies were transferred into fresh

food vials (fecundity vials), where they were allowed a window of 17-hours before being

transferred to fresh food vials again. The number of eggs in these fecundity vials was

counted. This was done at day5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, and 60 (by this time-

point all females had died).For each time point, the number of eggs in a vial was divided by

the number of surviving females to calculate per female fecundity. This was taken as the unit

of analysis. However, the final analysis of age-specific fecundity was done excluding the last

few age points (post-day40) to avoid an unbalanced design arising due to the death of most

individuals towards the later age points. Additionally, the cumulative fecundity of females

was also quantified across all the above-mentioned age points till death.

<u>Data analysis:</u>

The entire study used a total of eight populations. Populations with replicate identity (for

example, BL_1 and YLB_1) were phylogenetically closer (see Figure S1) compared to those

with different replicate identity. In addition, during an assay a given replicate set (for

example, BL_1 and YLB_1) was handled together whereas different replicate sets were handled

independently, often on different days. Thus data from replicate sets were treated as statistical

blocks and was modelled as a random effect in all the analyses. The entire studythus involved

four statistical blocks. Within each block, vials were the unit of replication and vial values /

means were used as the unit of analysis / response variable. Except for adult survivorship and

age-specific fecundity, results of all assays were analysed using mixed-model analysis of

variance (ANOVA) on Statistica (version 10, Statsoft, Tulsa, OK, USA). In case of the

analysis of juvenile traits, selection regime and sex (wherever applicable) were treated as

fixed effects. For the early-life female fecundity assay and egg size measurement, selection

regime and live-yeast concentration treatment were considered as fixed effects. For the

analysis of the early-life progeny count data from Experiment2, male (selection) regime and

female (selection) regime were used as fixed effects. For analysing female mean longevity

data from Experiment3, selection regime and mating status were treated as fixed effects.

Cumulative life-time fecundity data was analysed using selection regime as fixed factor.

For female survivorship analysis, we applied a mixed Cox's Proportional Hazard models,

using the `coxme` package in R version 4.0.2 (Gupta et al. 2013, Therneau et al. 2015). The

effects of regime, mating status and their interaction on survivorship was analysed and block

was fitted as a random intercept. A few flies those escaped during the course of the assay

were censored. Cox partial likelihood estimates were compared across treatment and

selection regime.

Age-specific fecundity was analysed as linear mixed-effects models with the `lme4` package

in R version 4.0.2 (Bates et al. 2014). Further, to obtain the p-values for the `lmer` models

with degrees of freedom based on the Satterthwaite approximation, we used the R package

`lmerTest` (Kuznetsova et al. 2017). Regime and time-point (age) were fitted as categorical

fixed effects. Block as well as all interactions involving it were fitted as random effects. In

addition, vial ID (replicate vial identity) was nested within block and was fitted as a random

effect. The nested random effect accounts for the fact that subsequent egg counts from the

same group of flies within a particular vial are not independent (Harrison et al. 2018). Post-

hoc analysis was done using Tukey's HSD with the `emmeans` package (Lenth 2018). To

measure reproductive senescence, rates at which fecundity declined with age, after reaching a

peak, were quantified for both regimes. Rate of decline in fecundity was quantified across

seven time-points (τ), starting from the peak, i.e. day10 post-eclosion until day40. For a

particular age-point, the mean of per-capita fecundity (ε) of a population was calculated using

the observed fecundity across all the vials within that population. A linear model ($\varepsilon \sim b_p\tau +$

c_p, where b_p is population specific slope and c_p is population specific intercept) was fitted to

these data using the `lm` function in R to obtain a b-coefficient (i.e., slope) for each of the

eight populations (Fig. S4 of supplementary information). The slope of the regression line

served as a proxy for rate of reproductive senescence. Finally, these b-coefficients, without

the sign, were analysed using mixed model ANOVA in Statistica, taking selection regime as

a fixed effect and block as a random effect.

3. Results

Experiment 1: Juvenile traits and female fecundity

We did not find a significant effect of selection regime on juvenile traits including

development time, pre-adult (egg-to-adult) survivorship and dry body weight at the time of

eclosion (Table S2). Analysis of the early-life fecundity data indicated significant effects of

selection regime and yeast concentration (Table 1). The effect of selection regime × yeast

concentration interaction was not significant (Table 1). Compared to the BL females, YLB

females had 8.7-22.89 % higher mean per-capita fecundity across all live-yeast

concentrations in the assay(Fig.1). Moreover, the higher early-life fecundity output of YLB

females was consistent across a gradient of live-yeast concentration (Fig.1). The effect of

selection regime on egg size was non-significant (Table S2). Hence, there was no evidence of

differential provisioning in eggs produced by the experimental females.

Figure 1: Female early-life fecundity from Experiment 1. Number of eggs laid during a 16-hour assay window by 4-5 day old (post-eclosion) females following the 48 hour live-yeast treatment was measured. This was done under three live-yeast conditions - no yeast (N), limiting yeast (L) and high yeast (H). The data distributions are shown as scatters. Pink circles represent data from BL (control regime) females, and teal triangles represent that from YLB (experimental regime). The black circle in each scatter represent the mean, which is shown along with the standard error (vertical error bars). Six of the twelve test females in a vial were used for fecundity measurement. Mean per-capita fecundity in a replicate vial was calculated using these six females. Three factor mixed model ANOVA indicated significant effects of selection regime and live-yeast treatment. However, the effect of selection regime × live-yeast treatment interaction was not significant.

Experiment 2: Female × male effect on female fitness

The analysis of the early-life progeny output data from Experiment 2 indicated a significant effect female selection regime. Neither the male selection regime, nor the interaction between these two fixed factors were found to have significant effects on progeny count (Table 1, Fig. 2). YLB females produced significantly more progeny irrespective of the regime of the males

they interacted with (Fig. 2). Thus, the higher early-life reproductive output in YLB's seems

to be independent of any male influence.

Figure 2: Female × male effect on early-life reproductive output in females as measured in Experiment 2. Number of

progeny produced during a 17-hour assay window by 4-5 day old (post-eclosion) BL and YLB females, mated and held with

BL and YLB males (full factorial combination), was measured. The data distributions are shown as scatters. Pink circles

represent data from BL (control regime) females, and teal triangles represent that from YLB (experimental regime). The

black circle in each scatter represent the mean, which is shown along with the standard error (vertical error bars). All six

females in each replicate vial were allowed to produce progeny. Vials mean, which was calculated over these six test

females, was used as the unit of analysis. Two factor mixed model ANOVA indicated a significant effect of female regime,

but effect of male regime on progeny count was not significant, nor was that of the male regime × female regime interaction.

Traits	Effects	SS	DF	MS	DEN DF	DEN MS	F	p
Early life fecundity	Selection Regime (SR)	947.8	1	947.8	3.01	7.47	126.83	**<0.01**
	Live-yeast concentration (YC)	51192.3	2	25596.2	6.00	1043.57	24.52	**<0.01**
	Block	19814.0	3	6604.7	5.59	1008.10	6.55	**0.03**
	SR × YC	23.0	2	11.5	6.01	42.40	0.27	0.77
	SR × Block	22.4	3	7.5	6.02	42.41	0.17	0.90
	YC × Block	6266.1	6	1044.4	6.00	42.39	24.63	**<0.01**
	SR × YC × Block	254.3	6	42.4	247.00	59.14	0.71	0.64
Progeny count	Female regime	411.13	1	411.14	3.00	7.40	55.55	**<0.01**
	Male regime	0.99	1	0.99	3.00	36.84	0.03	0.88
	Block	3149.10	3	1049.70	0.64	20.48	51.26	0.26
	Female regime × Male regime	7.18	1	7.18	3.00	23.76	0.30	0.62
	Female regime × Block	22.20	3	7.40	3	23.76	0.31	0.82
	Male regime × Block	110.51	3	36.84	3	23.76	1.55	0.36
	Fem. reg. × Male reg. × Block	71.28	3	23.76	222	20.71	1.15	0.33

Table 1: Results of the analyses on early-life fecundity(from experiment 1) and progeny output (from experiment 2) in females. Fecundity data was analysed using three factor mixed-model ANOVA with selection regime and live-yeast concentration as fixed factors, and block as random factor. Progeny count data from experiment 2 was analysed using three factor mixed-model ANOVA with female regime and male regime as fixed factors, and block as random factor. Statistically significant p-values are highlighted in boldface.

Experiment 3: Longevity and age-specific fecundity

We did not find a significant effect of selection regime on survival probability (Table S4) and mean longevity (Table S5) of females. Mating status, however, had a significant impact. Virgin females had a better survival rate compared to the mated females (Fig. S3). Unsurprisingly, analysis of the mean longevity indicated that mating status had a significant

effect on mean longevity (Table S5). Virgin females were found to have 67.58 % longer

mean longevity compared to the mated females for both BL and YLB selection regimes (Fig.

S4, Table S4). Results of these analyses can be found in the supplementary information.

Analysis of the cumulative fecundity of females quantified across different age points till

death revealed that there was no significant effect of selection regime on per-capita fecundity,

however, the effect of the selection regime×time point interaction was found to be significant

(Table 3a, Fig.3a). Tukey's adjusted p-value obtained by pairwise comparisons suggested

that, as compared to the ancestors, YLBs have significantly higher fecundity at the age of

day5 and day11 post-eclosion. Interestingly, there was a clear trend of YLBs having higher

mean per-capita fecundity in initial age points compared to BLs (through age- 5, 10, 15, 20

and 25 days post-eclosion). The trend, however, was found to be reversed at the later time

points, with BLs showing comparatively higher fecundity (through age- 30, 35 and 40 days

post-eclosion). The detailed outcome of the post-hoc analysis is mentioned in the

supplementary information (Supplementary sheet, Table- S6). Further, a comparison of the

slopes (b-coefficients) of the linear regression fit of age-specific fecundity on time for the two

regimes suggested a faster rate of decline in fecundity in YLBs compared to that of the BLs

(mean ±SE: b_{BL}=-2.03 ±0.41; b_{YLB}=-3.36 ±0.13, Fig. S5). Two factor mixed model ANOVA

on the b-coefficient indicated a significant effect of selection regime (Table 3b). Considering

the b-coefficient to be a measure of reproductive senescence, YLB females showed faster

reproductive senescence compared to that of the BL females. The cumulative fecundity of

females, quantified across all the above-mentioned age points, did not differ between the two

regimes (Fig. 3b, Table 3c)

Figure 3: Age-specific and cumulative fecundity of females as measured in Experiment 3. Number of eggs laid during

17 hour windows on day 5, 10, 15, 20, 25, 30, 35, 40 were measured during the longevity assay. Only CE (continuously

exposed) vials were used for this purpose. Per capita fecundity was calculated for a vial by diving the fecundity count by the

number of surviving females, which was twelve to start with. (a) Mean age-specific per capita fecundity analysis indicated a

significant effect of selection regime × age interaction. Pink circles represent data from BL (control regime) females, and

teal triangles represent that from YLB (experimental regime). Data points represent mean per capita fecundity at the

corresponding age point. The vertical error bars represent standard error of means. (b) The mean cumulative per-capita

fecundity of females, quantified across all the above-mentioned age points. The data distributions are shown as scatters. Pink

circles represent data from BL (control regime) females, and teal triangles represent that from YLB (experimental regime).

The black circle in each scatter represent the mean, which is shown along with the standard error (vertical error bars).

Analysis using two factor mixed model ANOVA indicated no significant difference in mean cumulative per-capita fecundity

between the females of the two regimes.

Trait	Effects	SS	DF	MS	Denominator DF	F	p
(a) Age-specific fecundity	Selection Regime (SR)	8.2	1	8.18	59.03	0.39	0.53
	Time point	6531.9	7	933.13	22.27	44.44	**< 0.01**
	SR × Time point	986.2	7	140.88	22.41	6.71	**<0.01**
(b) b-coefficient	Selection Regime	4.1	1	4.1	3	13.68	**0.03**
	Block	1.4	3	0.47	3	1.54	0.37
(c) Cumulative Fecundity	Selection Regime (SR)	35	1	35	3.01	0.06	0.82
	Block	4089	3	1363	3	2.26	0.26
	SR × Block	1806	3	602	76	0.73	0.53

Table 3: Results of the analyses of age-specific fecundity measured in Experiment 2. (a) Age-specific fecundity count was

analysed using linear mixed-effect model, where selection regimeand time point (age) were fitted as categorical fixed

effects, and block as random effect.(b) For each population, age-specific fecundity data was fitted in a linear model ($\varepsilon \sim b_p\tau$

$+ c_p$, where b_p is population specific slope and c_p is population specific intercept) to compute b-coefficient (slope). These b-

coefficient values (i.e., slope values without the sign) were analysed using two factor mixed model ANOVA with selection

regime and block as fixed and random factors respectively. (c) Cumulative fecundity, i.e., fecundity count summed over all

age points, were analysed using a two factor mixed model ANOVA with selection regime and block as fixed and random

factors respectively. Statistically significant p-values are mentioned in bold face.

4. Discussion

Yeast being the main source of protein, especially in a laboratory ecology (Chippindale et al.

1993, Lee et al. 2008), a culture protocol devoid of yeast supplement poses a significant

deprivation of dietary protein, limiting reproductive output. Following twenty three

generations of laboratory evolution under such a condition, we found clear evidence of

evolution of fecundity traits indicating adaptive changes. Fecundity and progeny output in

females most relevant to fitness showed a significant increase. However, there was no

evidence to suggest a change in either lifetime reproductive out or lifespan. Instead,

scheduling of reproductive output showed an interesting shift, wherein early life reproductive

output was found to be significantly higher, with a faster drop as the females grew older,

indicating faster reproductive senescence. Further, higher early life fecundity of females from

the experimental regime was observed in absence of yeast supplement as well as in presence

of low and high concentrations of the same, suggesting that plasticity in fecundity in response

to variation in protein availability did not evolve. There are a number of possible ways in

which females in the experimental populations could have attained higher early life

reproductive output. We discuss these possibilities and the broader implication of our finding

in the following section.

There are at least four different ways in which early-life fecundity and reproductive output

could have increased in females. First, increased larval resource acquisition and/or storage

can significantly contribute to the total resource available for reproduction, particularly early

in life. In *D. melanogaster,* yeast-starvation at the larval stage dramatically brings down

female fecundity such that females grown under complete yeast starvation can only remain

reproductive for about 5-6 days, clearly indicating the importance of larval resource

acquisition on early life reproductive output (Robertson & Sang 1944). However, we did not

find any evidence upholding this theory. Neither juvenile traits (development time, egg-to-

adult survivorship) nor body mass at eclosion was affected by experimental evolution.

Alternatively, an increase in the relative abundance of different macronutrients, especially

total protein content in adult females without affecting total body mass, could translate into

increased fecundity. This idea of more fine scale physiological adaptation, however, remains

to be tested with additional experimental data.

Secondly, number of offspring produced can be increased if investment in each offspring is

reduced (Hein et al. 2018, also see Vijendravarma et al. 2010). We, however, did not find any

evidence supporting such number vs. size trade-off as there was no effect of selection on egg

size even though number of eggs produced by the females increased substantially. This is

perhaps not surprising as previous investigations on *D. melanogaster* found egg size and

number to be genetically uncorrelated (Schwarzkopf et al. 1999). Hence, egg production rate

can evolve without affecting egg size.

A third possibility involves resource reallocation across life history traits such that higher

amount of resources are made available to egg production. For example, resources invested in

somatic maintenance could be reallocated to reproduction, especially when there is strong

selection on higher reproductive output and little or no selection on longevity traits, such as in

short lived organisms. Such a trade-off between reproduction and lifespan is pervasive in

nature and has been widely reported even in laboratory systems (Rose 1984, Chippindale et

al. 1993, Zwaan et al. 1995, Flatt 2011, Hansen et al. 2013, Maklakov et al. 2017). This

theory is also inadequate in explaining our findings on fecundity evolution as we did not find

a significant effect of experimental evolution on female longevity. This, however, is not

surprising as several attempts to test reproduction-longevity trade-off have been either

inconclusive or have failed (Le Bourg and Minois 1996, Dudash & Fenster 1997, Kotiaho &

Simmons 2003, Partridge et al. 2005, Grandison et al. 2009, Zajitschek et al. 2016, 2019).

There is a growing consensus that longevity and fecundity traits are, at least to some extent,

genetically independent of each other and the trade-off between them can be uncoupled

(reviewed in Flatt 2011). Importantly, in the present investigation we did not find any

evidence suggesting change in total lifetime reproductive output. If reproduction-longevity

trade-off is more sensitive to the total lifetime reproductive effort, a lack of response in the

longevity traits is not unconceivable.

A fourth, and perhaps the most plausible scenario is the change in reproductive schedule

without any alteration in life time reproductive output. Importantly, due to the 14 day discrete

generation cycle, early life fecundity is a major determinant of female fitness in our

laboratory ecology (see similar arguments in Promislow and Tatar 1998, Houle and Rowe

2003, Prasad & Joshi 2003, Rice et al. 2006). Both survival and reproduction beyond day14

of the generation cycle (i.e., approximately 4-5 days of adult age) have no contribution to

fitness. Hence, early life fitness components, including fecundity, are exclusive targets of

selection under such an ecology. This view of the population demography would predict the

evolution of early life fitness regardless of its consequences to the late life performance

(Stearns 1992). This theory fits perfectly with our age specific fecundity results, wherein

females from the evolved populations were found to have significantly higher fecundity. The

associated observation of a faster decrease in fecundity without a significant change in mean

lifetime egg output seems to fit antagonistic pleiotropy theory of ageing (Williams 1957).

Early investigations on *Drosophila* populations substantiated the theory (Rose 1984, Rauser

et al. 2006), and have since been shown to explain both somatic as well as reproductive

senescence. Multiple investigations also reported a substantial genetic variation in age-

specific fecundity in *D. melanogaster* (for example, Tatar et al. 1996, Durham et al. 2014).

More recently, using *C. elegans,* Anderson et al. (2011) found that genetically heterogeneous

populations selected for early-life fecundity evolved increased early but reduced late-life

reproductive output without suffering a total fitness or lifespan cost. Thus, it is not

unreasonable to suggest that adaptation seen in our YLB populations involves the evolution

of early-life beneficial alleles with detrimental pleiotropic effects on late-life reproductive

output.

Adaptive evolution of early-life fecundity in YLB females seems to support the hitherto

reported phenomenon of the release of hidden genetic variation in novel ecology or stress

(Badyaev 2005). Several past investigations have shown that novel environmental challenges

or stresses (e.g., heat shock) can generate new trait variances or convert otherwise

phenotypically neutral alleles to non-neutral (Hall 1990, Ratner et al. 1992, Kondrashov and

Houle 1996, Wessler 1996, Leips & Mackay 2000, Queitsch et al. 2002, Dworkin et al. 2003,

Nedelcu & Michod 2003, Ruden et al. 2003, Sollars et al. 2003). Such novel genetic variance

can then be subjected to selective pressures leading to adaptation. Further, there is a growing

literature on substantial G×E interaction on fitness-related traits, including fecundity (see

Leroi et al. 1994, Fry et al. 1998, Burns et al. 2012, Pallares et al. 2021). Most notably,

Pallares et al. (2021) reported that dietary sugar enrichment to have a profound effect on the

fitness landscape of the fecundity and longevity traits in *D. melanogaster*. Fitness

consequences of many alleles were found to be much stronger in the altered diet compared to

the standard diet (Pallares et al. 2021). Our results could also be affected by a similar diet

dependence of fitness consequence of fecundity traits in this system. Alleles that affect

unstimulated (by live yeast supplement) fecundity could be selectively neutral in the ancestral

BL populations, but have strong fitness effects under yeast deprived ecology of the YLB

populations and hence, be subjected to directional selection.

The identity of physiological traits that could be subjected to selection during such an

adaptation would depend on the proximate processes affecting early life fecundity. Therefore,

the proximate mechanistic explanation of higher early-life fecundity in the evolved females is

an important issue. We set aside this problem as dealing with it will require additional data.

However, there are several plausible theories to explore. For example, increased foraging

behaviour (McCaffery 1975, Ives 1981, Barnes et al. 2008), reduction in wastage of resources

in non-essential activities such as, spontaneous locomotion (Long & Rice 2006), better

resource recycling (Scott et al. 2004, Barth et al. 2011, Cuervo & Macian 2012, Adler &

Bonduriansky 2014, Mason et al. 2018) can all lead to increased resource availability for egg

production in females.

As discussed earlier, in case of short term food shortage, the strategy to divert resources away

from reproduction so that the temporary famine can be overcome, as suggested by adaptive

resource reallocation theory, could be adaptive. Here we have shown that selection can

favour novel reproductive strategies that might involve increased reproduction, if the famine

is long lasting. For a short lived species, as our results suggests, that could lead to increased

reproductive output at the cost of late life fertility.

6. Acknowledgements:

We thank Daniel E. L. Promislow and Benjamin Harrison of University of Washington for

their constructive criticism and suggestions on an earlier version of the manuscript.

Integrating their suggestions have greatly improved the quality of the manuscript, including

the clarity of the presentations. We thank the two anonymous reviewers of a previous version

of the manuscript for their immensely helpful and constructive criticisms. Incorporation of

their suggestions have greatly improved the quality of the manuscript. We are thankful to

Tanya Verma for her help during the experiments and population maintenance. The

experiments reported here was supported by the Core Research Grant (file no.

CRG/2019/002460) from Science and Engineering Research Board, Department of Science

and Technology, Government of India. PD thanks Council for Scientific and Industrial

Research, Government of India for financial support in the form of Junior and Senior

Research Fellowship. NP thanks Summer Research Fellowship Programme of Jawaharlal

7. References

Adler, M. I., & Bonduriansky, R. (2014). Why do the well-fed appear to die young? A new

evolutionary hypothesis for the effect of dietary restriction on lifespan. *Bioessays*, *36*, 439-

450.

Anagnostou, C., Dorsch, M., & Rohlfs, M. (2010). Influence of dietary yeasts on *Drosophila*

melanogaster life-history traits. *Entomologia Experimentalis et Applicata*, *136*, 1-11.

Andersen, L. H., Kristensen, T. N., Loeschcke, V., Toft, S., &Mayntz, D. (2010). Protein and

carbohydrate composition of larval food affects tolerance to thermal stress and desiccation in

adult *Drosophila melanogaster*. *Journal of insect physiology*, *56*, 336-340.

Anderson, J. L., Reynolds, R. M., Morran, L. T., Tolman-Thompson, J., & Phillips, P. C.

(2011). Experimental evolution reveals antagonistic pleiotropy in reproductive timing but not

life span in *Caenorhabditis elegans*. *Journals of Gerontology Series A: Biomedical Sciences*

and Medical Sciences, 66, 1300-1308.Arnqvist, G., & Nilsson, T. (2000). The evolution of

polyandry: multiple mating and female fitness in insects. *Animal behaviour*, 60, 145-164.

Badyaev, A. V. (2005). Stress-induced variation in evolution: from behavioural plasticity to

genetic assimilation. *Proceedings of the Royal Society B: Biological Sciences*, 272, 877-886.

Barnes, A. I., Wigby, S., Boone, J. M., Partridge, L., & Chapman, T. (2008). Feeding,

fecundity and lifespan in female *Drosophila melanogaster*. *Proceedings of the Royal Society*

B: Biological Sciences, *275*, 1675-1683.

Barth, J. M. I., Szabad, J., Hafen, E., & Köhler, K. (2011). Autophagy in *Drosophila* ovaries

is induced by starvation and is required for oogenesis. *Cell Death & Differentiation*, 18, 915-

924.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models

using lme4. *arXiv preprint arXiv:1406.5823*.

Begon M (1982) Yeasts and Drosophila. In: Ashburner M, Carson HL, Thompson J, editors.

The genetics and biology of Drosophila. New York: Academic Press. 345–384.

Becher, P. G., Flick, G., Rozpędowska, E., Schmidt, A., Hagman, A., Lebreton, S. &

Bengtsson, M. (2012). Yeast, not fruit volatiles mediate *Drosophila melanogaster* attraction,

oviposition and development. *Functional Ecology*, *26*, 822-828.

Boag, P. T., & Grant, P. R. (1981). Intense natural selection in a population of Darwin's

finches (Geospizinae) in the Galapagos. *Science*, 214, 82-85.

Bonduriansky, R., Maklakov, A., Zajitschek, F., & Brooks, R. (2008). Sexual selection,

sexual conflict and the evolution of ageing and life span. *Functional ecology*, 22, 443-453.

Burns, J. G., Svetec, N., Rowe, L., Mery, F., Dolan, M. J., Boyce, W. T., & Sokolowski, M.

B. (2012). Gene–environment interplay in *Drosophila melanogaster*: Chronic food

deprivation in early life affects adult exploratory and fitness traits. *Proceedings of the*

National Academy of Sciences USA, 109, 17239-17244.

Chapman, T., Liddle, L. F., Kalb, J. M., Wolfner, M. F., & Partridge, L. (1995). Cost of

mating in Drosophila melanogaster females is mediated by male accessory gland products.

Nature, 373, 241-244.

Chapman, T., & Partridge, L. (1996). Female fitness in *Drosophila melanogaster*: an

interaction between the effect of nutrition and of encounter rate with males. *Proceedings of*

the Royal Society of London. Series B: Biological Sciences, 263, 755-759.

Chippindale, A. K., Leroi, A. M., Kim, S. B., & Rose, M. R. (1993). Phenotypic plasticity

and selection in Drosophila life-history evolution. I. Nutrition and the cost of reproduction.

Journal of Evolutionary Biology, 6, 171-193.

Cuervo, A. M., & Macian, F. (2012). Autophagy, nutrition and immunology. *Molecular*

aspects of medicine, 33, 2-13.

Dudash, M. R., & Fenster, C. B. (1997). Multiyear study of pollen limitation and cost of

reproduction in the iteroparous Silene virginica. *Ecology*, 78, 484-493.

Durham, M. F., Magwire, M. M., Stone, E. A., &Leips, J. (2014). Genome-wide analysis in

Drosophila reveals age-specific effects of SNPs on fitness traits. *Nature communications*, 5,

1-8.

Dworkin, I., Palsson, A., Birdsall, K., & Gibson, G. (2003). Evidence that Egfr contributes to

cryptic genetic variation for photoreceptor determination in natural populations of Drosophila

melanogaster. *Current Biology*, 13, 1888-1893.

Fanson, B. G., Weldon, C. W., Pérez☐Staples, D., Simpson, S. J., & Taylor, P. W. (2009).

Nutrients, not caloric restriction, extend lifespan in Queensland fruit flies (Bactrocera tryoni).

Aging cell, 8, 514-523.

Fanson, B. G., & Taylor, P. W. (2012). Protein: carbohydrate ratios explain life span patterns

found in Queensland fruit fly on diets varying in yeast: sugar ratios. *Age*, 34, 1361-1368.

Field, C. J., Johnson, I. R., &Schley, P. D. (2002). Nutrients and their role in host resistance

to infection. *Journal of leukocyte biology*, 71, 16-32.

Flatt, T. (2011). Survival costs of reproduction in *Drosophila*. *Experimental gerontology*, 46,

369-375.

Fontana, L., Partridge, L., & Longo, V. D. (2010). Dietary restriction, growth factors and

aging: from yeast to humans. *Science*, 328, 321.

Fontana, L., & Partridge, L. (2015). Promoting health and longevity through diet: from model

organisms to humans. *Cell*, 161, 106-118.

Fry, J. D., Nuzhdin, S. V., Pasyukova, E. G., & Mackay, T. F. (1998). QTL mapping of

genotype–environment interaction for fitness in *Drosophila melanogaster*. *Genetics*

Research, 71, 133-141.

Garland T, Rose MR. 2009 Experimental evolution: concepts, methods, and applications of

selection experiments. Berkeley, CA: University of California Press.

Grandison, R. C., Piper, M. D., & Partridge, L. (2009). Amino-acid imbalance explains

extension of lifespan by dietary restriction in *Drosophila. Nature*, 462, 1061-1064.

Gupta, V., Ali, Z. S., & Prasad, N. G. (2013). Sexual activity increases resistance against

Pseudomonas entomophila in male *Drosophila melanogaster*. *BMC Evolutionary*

Biology, 13, 1-11.

Hall, B. G. (1990). Spontaneous point mutations that occur more often when advantageous

than when neutral. *Genetics*, 126, 5-16.

Harrison, X. A., Donaldson, L., Correa-Cano, M. E., Evans, J., Fisher, D. N., Goodwin, C. E.,

... & Inger, R. (2018). A brief introduction to mixed effects modelling and multi-model

inference in ecology. *PeerJ*, 6, e4794.

Hansen, M., Flatt, T., & Aguilaniu, H. (2013). Reproduction, fat metabolism, and life span:

what is the connection? *Cell metabolism,* 17, 10-19.

Hein, N., Brendel, M. R., Feilhauer, H., Finch, O. D., & Löffler, J. (2018). Egg size versus

egg number trade-off in the alpine-tundra wolf spider, Pardosa palustris (Araneae:

Lycosidae). *Polar Biology*, 41, 1607-1617.

Hoffmann, A. A., & Harshman, L. G. (1999). Desiccation and starvation resistance in

Drosophila: patterns of variation at the species, population and intrapopulation

levels. *Heredity*, 83, 637-643.

Houle, D., & Rowe, L. (2003). Natural selection in a bottle. *The American Naturalist*, 161,

50-67.

Ives, P. M. (1981). Feeding and egg production of two species of coccinellids in the

laboratory. *The Canadian Entomologist*, 113, 999-1005.

Jin, K., Wilson, K. A., Beck, J. N., Nelson, C. S., Brownridge III, G. W., Harrison, B. R., ...

& Promislow, D. (2020). Genetic and metabolomic architecture of variation in diet

restriction-mediated lifespan extension in Drosophila. *PLoS Genetics*, 16, e1008835.

Kassambara, A., Kosinski, M., Biecek, P., & Fabian, S. (2017). Package

'survminer'. *Drawing Survival Curves using 'ggplot2'(R package version 03 1)*.

Kondrashov, A. S., & Houle, D. (1994). Genotype-environment interactions and the

estimation of the genomic mutation rate in *Drosophila melanogaster*. *Proceedings of the*

Royal Society of London. Series B: Biological Sciences, 258, 221-227.

Kolss, M., Vijendravarma, R.K., Schwaller, G.& Kawecki, T.J. (2009). Life-history

consequences of adaptation to larval nutritional stress in *Drosophila*. *Evolution* 63, 2389-

2401.

Kotiaho, J. S., & Simmons, L. W. (2003). Longevity cost of reproduction for males but no

longevity cost of mating or courtship for females in the male-dimorphic dung beetle

Onthophagus binodis. *Journal of Insect Physiology*, 49, 817-822.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. (2017). lmerTest package: tests in

linear mixed effects models. *Journal of statistical software*, 82, 1-26.

Le Bourg, E., &Minois, N. (1996). Failure to confirm increased longevity in *Drosophila*

melanogaster submitted to a food restriction procedure. *The Journals of Gerontology Series*

A: Biological Sciences and Medical Sciences, 51, B280-B283.

Lee, K. P., Simpson, S. J., Clissold, F. J., Brooks, R., Ballard, J. W. O., Taylor, P. W., ... &

Raubenheimer, D. (2008). Lifespan and reproduction in Drosophila: new insights from

nutritional geometry. *Proceedings of the National Academy of Sciences USA*, 105, 2498-

2503.

Leips, J., & Mackay, T. F. (2000). Quantitative trait loci for life span in Drosophila

melanogaster: interactions with genetic background and larval density. *Genetics,* 155, 1773-

1788.

Lenth, R., Singmann, H., Love, J., Buerkner, P., &Herve, M. (2018). Emmeans: Estimated

marginal means, aka least-squares means. *R package version*, 1, 3.

Leroi, A. M., Chippindale, A. K., & Rose, M. R. (2004). Long-term laboratory evolution of a

genetic life-history trade-off in *Drosophila melanogaster*. 1. The role of genotype-by-

environment interaction. In *Methuselah Flies: A Case Study in the Evolution of Aging* (pp.

26-39).

Long, T. A., & Rice, W. R. (2007). Adult locomotory activity mediates intralocus sexual

conflict in a laboratory-adapted population of *Drosophila melanogaster*. *Proceedings of the

Royal Society B: Biological Sciences*, 274, 3105-3112.

Mair, W., Piper, M. D. W., & Partridge, L. (2005). Calories do not explain extension of life

span by dietary restriction in *Drosophila*. *PLoS Biology*, 3, e223.

Maklakov, A. A., Carlsson, H., Denbaum, P., Lind, M. I., Mautz, B., Hinas, A., &Immler, S.

(2017). Antagonistically pleiotropic allele increases lifespan and late-life reproduction at the

cost of early-life reproduction and individual fitness. *Proceedings of the Royal Society B:

Biological Sciences*, 284, 20170376.

Maklakov, A. A., Simpson, S. J., Zajitschek, F., Hall, M. D., Dessmann, J., Clissold, F., ... &

Brooks, R. C. (2008). Sex-specific fitness effects of nutrient intake on reproduction and

lifespan. *Current Biology*, 18, 1062-1066.

Marden, J. H., Rogina, B., Montooth, K. L., & Helfand, S. L. (2003). Conditional tradeoffs

between aging and organismal performance of Indy long-lived mutant flies. *Proceedings of

the National Academy of Sciences USA*, *100*, 3369-3373.

Mason, J. S., Wileman, T., & Chapman, T. (2018). Lifespan extension without fertility

reduction following dietary addition of the autophagy activator Torin1 in *Drosophila*

melanogaster. PloS One, *13*, e0190105.

McCaffery, A. R. (1975). Food quality and quantity in relation to egg production in *Locusta*

migratoria migratorioides. *Journal of Insect Physiology*, *21*, 1551-1558.

Mery, F., & Kawecki, T. J. (2005). A cost of long-term memory in *Drosophila*. *Science*, 308,

1148-1148.

Miller, C. V., & Cotter, S. C. (2018). Resistance and tolerance: the role of nutrients on

pathogen dynamics and infection outcomes in an insect host. *Journal of Animal Ecology*, 87,

500-510.

Moatt, J. P., Savola, E., Regan, J. C., Nussey, D. H., & Walling, C. A. (2020). Lifespan

extension via dietary restriction: time to reconsider the evolutionary mechanisms?.

BioEssays, 42, 1900241.

Nandy, B., Joshi, A., Ali, Z. S., Sen, S., & Prasad, N. G. (2012). Degree of adaptive male

mate choice is positively correlated with female quality variance. *Scientific Reports*, 2, 1-8.

Nandy, B., Dasgupta, P., Halder, S., & Verma, T. (2016). Plasticity in aggression and the

correlated changes in the cost of reproduction in male *Drosophila melanogaster*. *Animal*

Behaviour, 114, 3-9.

Nedelcu, A. M., & Michod, R. E. (2003). Sex as a response to oxidative stress: the effect of

antioxidants on sexual induction in a facultatively sexual lineage. *Proceedings of the Royal*

Society of London. Series B: Biological Sciences, 270, S136-S139.

Nakagawa, S., Lagisz, M., Hector, K. L., & Spencer, H. G. (2012). Comparative and

meta‐analytic insights into life extension via dietary restriction. *Aging cell*, 11, 401-409.

Pallares, L. F., Lea, A. J., Han, C., Filippova, E. V., Andolfatto, P., &Ayroles, J. F. (2021).

Diet unmasks genetic variants that regulate lifespan in outbred *Drosophila*. *BioRxiv*, 2020-10.

Parkash, R., & Munjal, A. K. (2000). Evidence of independent climatic selection for

desiccation and starvation tolerance in Indian tropical populations of *Drosophila*

melanogaster. *Evolutionary Ecology Research*, 2, 685-699.

Partridge, L., Green, A., & Fowler, K. (1987). Effects of egg-production and of exposure to

males on female survival in *Drosophila melanogaster*. *Journal of Insect Physiology*, 33, 745-

749.

Partridge, L., Piper, M. D., & Mair, W. (2005). Dietary restriction in *Drosophila*.

Mechanisms of ageing and development, 126, 938-950.

Piper, M. D., Mair, W., & Partridge, L. (2005). Counting the calories: the role of specific

nutrients in extension of life span by food restriction. *The Journals of Gerontology Series A:*

Biological Sciences and Medical Sciences, 60, 549-555.

Prasad, N. G., & Joshi, A. (2003). What have two decades of laboratory life-history evolution

studies on Drosophila melanogaster taught us? *Journal of Genetics*, 82, 45-76.

Promislow, D. E., & Tatar, M. (1998). Mutation and senescence: where genetics and

demography meet. *Genetica,* 102, 299-314.

Queitsch, C., Sangster, T. A., & Lindquist, S. (2002). Hsp90 as a capacitor of phenotypic

variation. *Nature,* 417, 618-624.

Ratner, V. A., Zabanov, S. A., Kolesnikova, O. V., & Vasilyeva, L. A. (1992). Induction of

the mobile genetic element Dm-412 transpositions in the Drosophila genome by heat shock

treatment. *Proceedings of the National Academy of Sciences*, 89, 5650-5654.

Rauser, C. L., Tierney, J. J., Gunion, S. M., Covarrubias, G. M., Mueller, L. D., & Rose, M.

R. (2006). Evolution of late-life fecundity in *Drosophila melanogaster*. *Journal of*

Evolutionary Biology, 19, 289-301.

Rice, W. R., Stewart, A. D., Morrow, E. H., Linder, J. E., Orteiza, N., & Byrne, P. G. (2006).

Assessing sexual conflict in the Drosophila melanogaster laboratory model system.

Philosophical Transactions of the Royal Society B: Biological Sciences, 361, 287-299.

Rikke, B. A., Liao, C. Y., McQueen, M. B., Nelson, J. F., & Johnson, T. E. (2010). Genetic

dissection of dietary restriction in mice supports the metabolic efficiency model of life

extension. *Experimental gerontology*, 45, 691-701.

Rion, S., & Kawecki, T. J. (2007). Evolutionary biology of starvation resistance: what we

have learned from *Drosophila*. *Journal of Evolutionary Biology*, 20, 1655-1664.

Robertson, F. W., & Sang, J. H. (1944). The ecological determinants of population growth in

a *Drosophila* culture. I. Fecundity of adult flies. *Proceedings of the Royal Society of London.*

Series B-Biological Sciences, 132, 258-277.

Rohlfs, M., & Hoffmeister, T. S. (2005). Maternal effects increase survival probability in

Drosophila subobscura larvae. *Entomologia Experimentalis et Applicata*, 117, 51-58.

Rose, M. R. (1984). Laboratory evolution of postponed senescence in *Drosophila*

melanogaster. *Evolution*, 38, 1004-1010.

Ruden, D. M., Garfinkel, M. D., Sollars, V. E., & Lu, X. (2003). Waddington's widget:

Hsp90 and the inheritance of acquired characters. *Seminars in Cell & Developmental Biology*

14, 301-310.

Schluter, D., & Grant, P. R. (1984). Ecological correlates of morphological evolution in a

Darwin's finch, *Geospiza difficilis*. *Evolution*, 38, 856-869.

Schwarzkopf, L., Blows, M. W., & Caley, M. J. (1999). Life-history consequences of

divergent selection on egg size in Drosophila melanogaster. *The American Naturalist*, 154,

333-340.

Scott, R. C., Schuldiner, O., & Neufeld, T. P. (2004). Role and regulation of starvation-

induced autophagy in the *Drosophila* fat body. *Developmental cell*, 7, 167-178.

Shanley, D. P., & Kirkwood, T. B. (2000). Calorie restriction and aging: A life‐history

analysis. *Evolution*, 54, 740-750.

Short, C. A., Hatle, J. D., & Hahn, D. A. (2020). Protein stores regulate when reproductive

displays begin in the male Caribbean fruit fly. *Frontiers in physiology*, 11, 991.

Sisodia, S., & Singh, B. N. (2012). Experimental evidence for nutrition regulated stress

resistance in *Drosophila ananassae*. *PLoS One* 7: e46131.

Simmons, F. H., & Bradley, T. J. (1997). An analysis of resource allocation in response to

dietary yeast in *Drosophila melanogaster*. *Journal of insect physiology*, 43, 779-788.

Sollars, V., Lu, X., Xiao, L., Wang, X., Garfinkel, M. D., & Ruden, D. M. (2003). Evidence

for an epigenetic mechanism by which Hsp90 acts as a capacitor for morphological evolution.

Nature Genetics, 33, 70-74.

Starmer, W. T. (1981). A comparison of *Drosophila* habitats according to the physiological

attributes of the associated yeast communities. *Evolution*, 35, 38-52.

Stearns, S.C. (1992) The evolution of life histories. *Oxford University Press, Oxford*.

Tatar, M., & Carey, J. R. (1995). Nutrition mediates reproductive trade‐offs with age‐specific

mortality in the beetle *Callosobruchus maculatus*. *Ecology*, 76, 2066-2073.

Tatar, M., Promislow, D. E., Khazaeli, A. A., & Curtsinger, J. W. (1996). Age-specific

patterns of genetic variance in Drosophila melanogaster. II. Fecundity and its genetic

covariance with age-specific mortality. *Genetics*, 143, 849-858.

Therneau, T. (2015). Mixed effects Cox models. *CRAN repository*.

Vijendravarma, R. K., Narasimha, S., & Kawecki, T. J. (2010). Effects of parental larval diet

on egg size and offspring traits in Drosophila. *Biology Letters,* 6, 238-241.

Vijendravarma, R. K., & Kawecki, T. J. (2013). Epistasis and maternal effects in

experimental adaptation to chronic nutritional stress in *Drosophila*. *Journal of Evolutionary*

Biology, 26, 2566-2580.

Weldon, C. W., Mnguni, S., Démares, F., du Rand, E. E., Malod, K., Manrakhan, A., &

Nicolson, S. W. (2019). Adult diet does not compensate for impact of a poor larval diet on

stress resistance in a tephritid fruit fly. *Journal of Experimental Biology*, 222, jeb192534.

Wessler, S. R. (1996). Turned on by stress. Plant retrotransposons. *Current Biology,* 6, 959-

961.

Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of senescence.

Evolution, 11, 398-411.

Wolfner, M. F. (2009). Battle and ballet: molecular interactions between the sexes in

Drosophila. *Journal of Heredity*, 100, 399-410.

Zajitschek, F., Zajitschek, S. R., Canton, C., Georgolopoulos, G., Friberg, U., & Maklakov,

A. A. (2016). Evolution under dietary restriction increases male reproductive performance

without survival cost. *Proceedings of the Royal Society B: Biological Sciences*, 283,

20152726.

Zajitschek, F., Georgolopoulos, G., Vourlou, A., Ericsson, M., Zajitschek, S. R., Friberg, U.,

& Maklakov, A. A. (2019). Evolution under dietary restriction decouples survival from

fecundity in *Drosophila melanogaster* females. *The Journals of Gerontology: Series A*, 74,

1542-1548.

Zhang, D. W., Xiao, Z. J., Zeng, B. P., Li, K., & Tang, Y. L. (2019). Insect behavior and

physiological adaptation mechanisms under starvation stress. *Frontiers in physiology*, 10,

163.

Zimmerman, J. A., Malloy, V., Krajcik, R., & Orentreich, N. (2003). Nutritional control of

aging. Experimental gerontology, 38, 47-52.

Zwaan, B., Bijlsma, R., & Hoekstra, R. F. (1995). Direct selection on life span in *Drosophila*

melanogaster. *Evolution*, 49, 649-659.