Multiple regulatory mechanisms control the production of CmrRST, an atypical

signal transduction system in *Clostridioides difficile*

Elizabeth M. Garrett[1], Anchal Mehra[1], Ognjen Sekulovic[2], Rita Tamayo[1]*

[1] Department of Microbiology and Immunology, University of North Carolina Chapel Hill, 125

Mason Farm Road CB#7290, Chapel Hill, NC, 27599, USA

[2] Department of Molecular Biology and Microbiology, Tufts University School of Medicine,

136 Harrison Avenue, Boston, MA, 02111, USA

*Corresponding Author

Rita Tamayo, PhD

125 Mason Farm Rd, CB# 7290

Chapel Hill, NC, 27599-7290

(919) 843-2864

Rita_tamayo@med.unc.edu

ABSTRACT

Clostridioides difficile, an intestinal pathogen and leading cause of nosocomial infection, exhibits extensive phenotypic heterogeneity through phase variation by site-specific recombination. The signal transduction system CmrRST, which encodes two response regulators (CmrR and CmrT) and a sensor kinase (CmrS), impacts *C. difficile* cell and colony morphology, surface and swimming motility, biofilm formation, and virulence in an animal model. CmrRST is subject to phase variation through site-specific recombination and reversible inversion of the '*cmr* switch', and expression of *cmrRST* is also regulated by c-di-GMP through a riboswitch. The goal of this study was to determine how the *cmr* switch and c-di-GMP work together to regulate *cmrRST* expression. We generated "phase locked" strains by mutating key residues in the right inverted repeat flanking the *cmr* switch. Phenotypic characterization of these phase locked *cmr*-ON and -OFF strains demonstrates that they cannot switch between rough and smooth colony morphologies, respectively, or other CmrRST-associated phenotypes. Manipulation of c-di-GMP levels in these mutants showed that c-di-GMP promotes *cmrRST* expression and associated phenotypes independent of *cmr* switch orientation. We identified multiple promoters controlling *cmrRST* transcription, including one within the ON orientation of *cmr* switch and another that is positively autoregulated by CmrR. Overall, this work reveals a complex regulatory network that governs *cmrRST* expression and a unique intersection of phase variation and c-di-GMP signaling. These findings suggest that multiple environmental signals impact the production of this signaling transduction system.

IMPORTANCE

Clostridioides difficile is a leading cause of hospital-acquired intestinal infections in the

U.S. The CmrRST signal transduction system controls numerous physiological traits

and processes in *C. difficile*, including cell and colony morphology, motility, biofilm

formation, and virulence. Here we define the complex, multi-level regulation of *cmrRST*

expression, including stochastic control through phase variation, modulation by the

second messenger c-di-GMP, and positive autoregulation by CmrR. The results of this

study suggest that multiple, distinct environmental stimuli and selective pressures must

be integrated to appropriately control *cmrRST* expression.

INTRODUCTION

The ability to adapt to environmental changes is critical to bacterial survival,

including that of pathogens, which can face rapidly changing conditions and stresses

during infection (1-4). Sense-and-respond adaptation strategies often involve two-

component systems (TCS) that consist of a sensor kinase and a cognate response

regulator. In response to an activating signal, such as binding of a ligand or a change in

pH, the sensor kinase autophosphorylates and activates the response regulator through

transfer of the phosphoryl group (5,6). Response regulators have a wide range of

functions, and many control transcription through direct binding of DNA. The resulting

transcriptional changes contribute to adaptation of the bacterium to the environmental

stimulus sensed by the sensor kinase. Bacteria can also adapt to stimuli via intracellular

small molecules such as cyclic diguanylate (c-di-GMP) (7,8). The intracellular level of c-

di-GMP is modulated by the opposing activities of diguanylate cyclases and

phosphodiesterases that synthesize and degrade c-di-GMP, respectively; the

production and function of these enzymes is controlled by environmental signals (9,10).

C-di-GMP is then recognized by specific protein or RNA receptors (riboswitches) that

mediate the adaptive response (11-13).

In contrast, diversification of phenotypes in a bacterial population serves as a

bet-hedging strategy to help ensure survival of the population as a whole. The

development of phenotypically distinct variants, independent of environmental

conditions, improves the odds that a subpopulation survives a sudden stress (2,14).

Phase variation, a mechanism of generating phenotypic heterogeneity, occurs through

reversible genetic changes that typically cause an ON/OFF phenotypic "switch" (15,16).

Several mechanisms of phase variation have been described including conservative

site-specific recombination, in which a sequence-specific recombinase binds inverted

repeats and mediates inversion of the intervening DNA. The invertible DNA element

contains regulatory information, such as a promoter, that impacts the expression of

adjacent genes. In a well-characterized example in *Escherichia coli*, phase variation of

fimbriae production is mediated by the *fimS* invertible element, which contains a

promoter that drives transcription of the fimbrial genes when properly oriented (17-19).

Additional mechanisms have been recently described. Phase variation of the cell wall

protein CwpV in *Clostridioides difficile* occurs as a result of site-specific recombination

of a sequence mapping to the 5' untranslated region of *cwpV*. In one orientation, the

invertible DNA sequence results in the formation of an intrinsic terminator in the mRNA

when in one orientation, preventing expression of the downstream gene *cwpV*; the

intrinsic terminator does not form in the mRNA with the sequence in the inverse

orientation, allowing *cwpV* transcription to occur (20). Phase variation of flagellum and

toxin biosynthesis in *C. difficile* also occurs through a mRNA-mediated mechanism,

where one orientation of the invertible sequence yields an mRNA permissive for

transcriptional readthrough, while the other orientation results in Rho-mediated

transcription termination (21-24).

Clostridioides difficile is an intestinal pathogen and a leading cause of

nosocomial infections in the U.S. *C. difficile* infection (CDI) can result in mild to severe

diarrhea and potentially fatal complications such as pseudomembranous colitis, toxic

megacolon, and sepsis. Recent work has shown that *C. difficile* contains multiple

invertible DNA elements flanked by inverted repeats (including those noted above),

indicating a considerable capacity for phenotypic heterogeneity through phase variation

(25,26). Three of these invertible elements have been demonstrated to regulate

downstream genes and related phenotypes in a phase variable manner. The Cdi4

invertible element, also called the flagellar switch, modulates expression of the *flgB*

flagellar operon resulting in phase variation of flagella (21,22). This operon encodes the

sigma factor SigD, which promotes the transcription of flagellar genes as well as

transcription of *tcdR*, which encodes a direct activator of the *C. difficile* toxin genes *tcdA*

and *tcdB* (27,28). Accordingly, the production of these toxins is also phase variable (21-

23). The Cdi1 invertible element mediates phase variation of CwpV, which contributes

to phage resistance (20,29-31). Finally, the Cdi6 invertible element, here called the "*cmr*

switch*", regulates the expression of *cmrRST* in a phase variable manner (25,32). The

cmrRST operon encodes a putative non-canonical TCS with two DNA-binding response

regulators (CmrR and CmrT) and a sensor kinase (CmrS) (32). Both CmrR and CmrT

contain phosphoreceiver and DNA binding domains suggesting that they function as

transcriptional regulators. Phase variation of CmrRST allows *C. difficile* to switch

between rough and smooth colony morphologies that differ in several physiological

characteristics. Through unknown mechanisms, CmrRST positively regulates type IV

pilus-independent surface motility and cell elongation, and it negatively regulates

swimming motility and biofilm formation. Furthermore, a *cmrR* mutant strain is deficient

for colonization and shows attenuated virulence in a hamster model of infection,

indicating a role for this regulatory system in CDI.

C-di-GMP riboswitches are widespread in the *C. difficile* genome and appear to

be the primary mechanism of c-di-GMP regulation in *C. difficile* (11,12,33,34). Upstream

of the *cmr* switch are encoded a σ^A-dependent promoter followed by a class II c-di-GMP

riboswitch that positively regulates *cmrRST* transcription in response to c-di-GMP

(Figure 1A) (33,34). Increasing c-di-GMP levels results in the formation of the rough

colony morphology, consistent with increased *cmrRST* expression (32). The relative

contributions of the c-di-GMP riboswitch and the *cmr* switch to controlling expression of

cmrRST, and therefore to the phenotypes controlled by this system, are unknown.

One challenge to the study of phase variation is its stochastic nature, which adds

uncontrolled variation into an otherwise controlled experiment. Phase locked strains in

which the invertible element is prevented from inverting can be a useful tool with which

to study phase variable systems (21,35). Our previous work on CmrRST relied primarily

on characterization of wild-type (WT) *C. difficile* rough and smooth colony isolates,

which have a strong bias for the ON and OFF *cmr* switch orientations, respectively, but

remain capable of switch inversion and phenotypic switching (32). In this study, we

characterized *cmrRST* regulation through the interplay of c-di-GMP and the *cmr* switch

by generating phase-locked *cmr* OFF and *cmr* ON mutants. Through phenotypic

characterization of these phase locked strains and analysis of the effects of c-di-GMP

and *cmr* switch orientation, we found that that these regulatory features act

independently. We determined that phase variable regulation occurs through inversion

of a promoter encoded in the *cmr* switch. In addition, we demonstrate that CmrR

positively autoregulates expression of *cmrRST* via an additional promoter located

between the *cmr* switch and the *cmrR* start codon. The results of this study indicate that

cmrRST expression is subject to regulation both through sense-and-respond

mechanisms and phase variation, highlighting the potential importance of this system to

C. difficile physiology through the variety of activating signals.

RESULTS

Generation and phenotypic characterization of phase locked *cmr* ON and *cmr*

OFF strains

To characterize the dual regulation of the *cmrRST* system by phase variation and

c-di-GMP, we first generated phase- locked strains with the *cmr* switch fixed in the ON

or OFF orientation. Phase locking can be achieved in multiple ways. One strategy is to

delete the site-specific recombinase responsible for inversion. In *C. difficile* R20291, the

RecV recombinase is required for inversion of the *cmr* switch (25). However, RecV

mediates inversion of multiple sequences, thus a mutation in *recV* has pleiotropic

effects (20,21,25). Instead, we chose to phase lock the *cmr* switch by mutating an

inverted repeat sequence to prevent site-specific recombination from occurring (36).

The exact nucleotides at which recombination and inversion of the *cmr* switch occurs

were previously identified (25). We used allelic exchange to delete the nucleotide at the

site of inversion in the right inverted repeat (RIR) (position 3,736,177) in the R20291

genome, as well as one additional nucleotide on each side (Figure 1B). Prior work

determined that mutating the equivalent residues is the flagellar switch RIR eliminated

switch inversion resulting on phase-locked strains (37). We recovered independent

mutants with the *cmr* switch in either orientation and designated them *cmr*-Δ3 ON and

cmr-Δ3 OFF. These mutants were subjected to orientation specific PCR to confirm that

they are genotypically phase locked. In contrast to the WT parent that yielded both *cmr*

ON and OFF products indicating heterogeneity in *cmr* switch orientation in that

population of cells, *cmr*-Δ3 ON exclusively yielded a PCR product corresponding to the

ON orientation, and *cmr*-Δ3 OFF only yielded the OFF orientation product (Figure 1C).

Previous work showed that R20291 rough and smooth colony variants correlate

with *cmr* ON and OFF gene expression and phenotypes, respectively (32). Specifically,

the *cmr* ON, rough colony variant exhibits greater surface growth and reduced

swimming motility compared to the *cmr* OFF, smooth colony variant. Furthermore,

ectopic expression of *cmrR* and *cmrT* individually stimulated rough colony formation and

surface motility while inhibiting flagellum-mediated swimming motility, although only

cmrT was required for rough colony development and surface motility. To conclusively

determine the effect of *cmr* switch orientation on gene expression, we measured the

abundance of the *cmrR* and *cmrT* transcripts in the *cmr*-Δ3 ON and *cmr*-Δ3 OFF

mutants. Non-locked rough and smooth isolates were included as controls (32). Both

cmrR and *cmrT* transcripts were 7- to 10-fold more abundant in *C. difficile* with the *cmr*

switch in the ON orientation, whether in the naturally arising rough colony variant or in

cmr-Δ3 ON (Figure 1D). Therefore, the three nucleotides at the site of RecV-mediated

recombination are required for *cmr* switch inversion, and the ON orientation

corresponding to the inverse of the published R20291 genome promotes expression of

cmrRST.

The *cmr*-Δ3 phase-locked mutants were then evaluated for the phenotypes

previously associated with *cmrRST* expression. As a control we included WT R20291,

which is capable of CmrRST phase variation and generation of both rough and smooth

colonies, as well as a Δ*cmrR*Δ*cmrT* double mutant that forms only smooth colonies

(Figure 2A). The *cmr*-Δ3 OFF mutant formed exclusively smooth colonies, similar to

Δ*cmrR*Δ*cmrT*. *cmr*-Δ3 ON formed rough colonies that resemble those in WT, though

they appeared to have less defined topology (Figure 2A). Also consistent with increased

cmrRST expression, *cmr*-Δ3 ON displayed greater surface motility than WT, while *cmr*-

Δ3 OFF had decreased surface motility more similar to that of Δ*cmrR*Δ*cmrT* (Figure 2B,

C). Finally, *cmr*-Δ3 ON showed decreased swimming motility and biofilm formation

compared to all other strains (Figure 2D, E). Together, these results reinforce the model

that the *cmr* switch orientation modulates expression of *cmrRST* and the assignment of

the respective ON and OFF switch orientations.

C-di-GMP regulates *cmrRST* expression independently of the *cmr* switch

Upstream of the *cmr* switch is a σ^A-dependent promoter followed by a c-di-GMP

riboswitch sequence, adding another regulatory layer to the transcription of *cmrRST*

(33,34). The two regulatory features could act independently with one regulatory

element dominant over the other, or they might act coordinately such that both the

riboswitch and *cmr* switch must be in the "on" state to allow *cmrRST* transcription. To

distinguish between these possibilities, we modulated c-di-GMP levels in the phase-

locked *cmr*-Δ3 ON and -OFF strains by overexpressing the diguanylate cyclase *dccA*

(pP$_{tet}$::*dccA*) to increase intracellular c-di-GMP or the catalytic EAL domain of the

phosphodiesterase *pdcA*, which hydrolyzes c-di-GMP, to reduce c-di-GMP (pP$_{tet}$::EAL)

(10,38,39). Increasing c-di-GMP through overexpression of *dccA* resulted in rough

colony formation in *cmr*-Δ3 OFF, which normally forms smooth colonies (Figure 3A).

Additionally, increasing c-di-GMP enhanced the rough colony appearance of *cmr*-Δ3

ON. In contrast, reduction of c-di-GMP through overexpression of the EAL domain did

not visibly affect the morphology of *cmr*-Δ3 ON or -OFF, suggesting that the impact of

EAL overexpression on c-di-GMP levels was insufficient to alter expression of *cmrRST*

compared to the WT with vector under these conditions. Overexpression of *dccA* led to

a significant 2- to 3-fold increase in the abundance of the *cmrS* transcript, which served

as a marker for *cmrRST* expression, in both *cmr*-Δ3 ON and -OFF, consistent with the

colony morphology phenotypes observed (Figure 3B).

These methods assess changes in *cmrRST* expression as a population average.

To determine the effects of the c-di-GMP riboswitch and the *cmr* switch on the

heterogeneity of *cmrRST* expression among individual bacteria, we used a strain in

which *cmrR* was replaced by a codon-optimized *SNAP*-tag gene reporter,

Δ*cmrR*::SNAP (25). The pP$_{tet}$::*dccA* plasmid and vector control were introduced into this

strain. In Δ*cmrR*::SNAP with vector and in the uninduced control, approximately 1% of

the population fluoresced (Figure 3C). This result is similar to prior analyses showing

that a minority of WT cells express *cmrRST* (25). Induction of *dccA* increased

fluorescence-positive cells to > 95%. This result was not due to a shift in *cmr* switch

orientation; by qPCR, the orientation of the *cmr* switch was not significantly different

between WT overexpressing *dccA* and the vector control (Figure S1). Therefore, the

higher percentage of fluorescent cells reflects *cmrRST* expression in response to c-di-

GMP. Together, these data indicate that expression of *cmrRST* from the upstream σ^A-

dependent promoter and c-di-GMP riboswitch is not dependent on the orientation of the

cmr switch. Rather, high c-di-GMP levels increase *cmrRST* expression regardless of

cmr switch orientation.

The *cmr* switch contains a promoter in the ON orientation

The best characterized phase variation regulatory mechanisms involve a

promoter encoded within the invertible element that drives expression of adjacent genes

when in the proper orientation (19). Our data suggest that *cmrRST* transcription

originates from multiple sites in a manner dependent on internal c-di-GMP

concentrations and the orientation of the *cmr* switch. Therefore, we sought to map the

transcriptional start sites (TSS) in strains reflecting these different states. Instead of

using *cmr-Δ3* phase locked strains, which are missing three nucleotides of unknown

significance to transcription, we used strains in which the *cmr* switch is locked by a

mutation in *recV*, designated *recV cmr*-ON and *recV cmr*-OFF (Figure S2). Additionally,

dccA was overexpressed in *recV cmr*-OFF to identify any TSS associated with

increased c-di-GMP and transcriptional readthrough via the riboswitch. Using 5' RACE,

four total TSS were identified in *recV cmr*-ON, *recV cmr*-OFF, and *recV cmr*-OFF

pDccA. In *recV cmr*-OFF with and without *dccA* expression, a TSS designated TSS1

was identified 699 nt upstream of the *cmrR* start codon, located upstream of the c-di-

GMP riboswitch (Figure 4A, Figure S3). TSS1 was not detected in *recV cmr*-ON, but

this may be due to lower relative abundance of the TSS1 transcript in this strain

compared to transcripts from other initiation sites. Another TSS, designated TSS2, was

found in *recV cmr*-ON, located 336 nt upstream of *cmrR* and mapping within *cmr* switch

in the ON orientation. A third TSS, designated TSS3, was identified 342 nt upstream of

the *cmrR* start codon in both *recV cmr*-OFF strains. This position maps TSS3 to within

the *cmr* switch, appearing only in transcripts from the *cmr*-OFF strains. In all three

strains, a fourth TSS (TSS4) was identified 90 nt upstream from the *cmrR* start codon.

The 5' RACE results confirm an expected TSS upstream of the c-di-GMP

riboswitch and suggest that expression of *cmrRST* can originate from multiple additional

sites. TSS2 within the *cmr* switch in the ON orientation was identified, however we

unexpectedly identified TSS3 within the *cmr* switch in the OFF orientation. To elucidate

how phase variable *cmrRST* expression is mediated, we examined the functionality of

the newly identified TSS. We created a series of transcriptional reporter fusions of the

alkaline phosphatase (AP) gene *phoZ* to regions immediately upstream of *cmrRST*

(Figure 4B) (40). Two of the fusions contain the upstream region from the left inverted

repeat (LIR) to the start of the *cmrR* coding sequence. These fusions differ in the

orientation of the invertible element: ON (pMC123*::cmr*ON/TSS4-*phoZ*) encompasses

TSS2, and OFF (pMC123*::cmr*OFF/TSS4-*phoZ*) encompasses TSS3. Both constructs

also include TSS4. Additionally, we included a reporter with the region from the 3' end

of the RIR to the start of the *cmrR* coding sequence (pMC123*::*TSS4-*phoZ*), which

contains TSS4, as well as a promoterless control (vector). To prevent inversion of the

cmr switch present in the reporters once in *C. difficile*, we assayed AP activity in a *recV*

mutant lacking the recombinase necessary for inversion (25). *C. difficile* with

pMC123*::*TSS4-*phoZ* (Figure 4B, construct 1) or pMC123*::cmr*OFF/TSS4-*phoZ*

(construct 2) showed no significant difference in activity compared to the promoterless

control (Figure 4B). In contrast, *C. difficile* with pMC123*::cmr*ON/TSS4-*phoZ* (construct

3) exhibited 20-fold higher AP activity (Figure 4B). These results indicate the presence

of a functional promoter when the *cmr* switch is in the ON orientation but not the OFF

orientation.

CmrR positively autoregulates expression of *cmrRST*

Response regulators often have the property of autoregulating their expression

(41). To determine whether CmrR and/or CmrT have the ability to transcriptionally

regulate *cmrRST*, we overexpressed *cmrR* (pCmrR) and *cmrT* (pCmrT) in *C. difficile*

and analyzed *cmrRST* expression by qRT-PCR. Overexpression of *cmrR* resulted in a

significant 7-fold increase in transcript abundance of *cmrS* and *cmrT* as compared to

the vector control (Figure 5A). In contrast, overexpression of *cmrT* had no effect on

cmrR or *cmrS* transcript abundance. These results indicate that CmrR, but not CmrT,

transcriptionally regulates the *cmrRST* operon.

To assess whether CmrR-mediated regulation is dependent on the orientation of

the invertible element, we overexpressed *cmrR* from a plasmid in the phase-locked *recV*

cmr-ON and *cmr*-OFF mutants. We then used qRT-PCR measuring *cmrS* to assess

expression. Overexpression of *cmrR* significantly increased *cmrS* transcript abundance

compared to the respective vector control in both *recV cmr*-OFF and *cmr*-ON mutants

(Figure 5B). While overexpression of *cmrR* in *recV cmr*-OFF resulted in over a 20-fold

increase in *cmrS* transcript abundance relative to vector control, the *cmrS* transcript

increased only about 2-fold in *recV cmr*-ON pCmrR relative to vector control.

We considered the possibility that CmrR regulates *cmrRST* expression from one

of the previously identified TSS. To address this possibility, we used qRT-PCR to

measure the transcript abundance by probing sequences immediately downstream of

TSS1, the *cmr* switch containing TSS2 or TSS3, and TSS4, as well as within *cmrS*

(Figure 5C). Because transcript abundance differed substantially depending on the

region measured, we expressed the data normalized to *rpoC* transcript abundance

rather than the ΔΔCt method. Overexpression of *cmrR* did not alter transcript

abundance when probing immediately downstream of TSS1, regardless of *cmr* switch

orientation, indicating that CmrR-mediated autoregulation occurs downstream of the

TSS1 promoter (Figure 5D). In contrast, when probing immediately downstream of the

cmr switch (TSS2/TSS3), *cmrR* overexpression resulted in 11-fold higher transcript

abundance in *recV cmr*-OFF relative to the vector control, though the difference did not

reach statistical significance (Figure 5D). Transcript abundance was 23-fold higher in

recV cmr-ON compared to *recV cmr*-OFF (vector controls, $p < 0.01$), and *cmrR*

overexpression in the *cmr*-ON background led to a further 3-fold increase. Similar

changes were observed when probing downstream of TSS4: *cmrR* overexpression

increased transcript abundance 31-fold in *recV cmr*-OFF and 60-fold higher in *recV cmr*-

ON compared to the *recV cmr*-OFF vector control (Figure 5D). These results suggest

that CmrR autoregulates *cmrRST* expression from the promoter corresponding to TSS2,

TSS4, or both. Notably, transcript abundance was 29- and 36-fold higher when

measuring downstream of TSS4 compared to TSS3 (*cmr*-OFF) and TSS2 (*cmr*-ON),

respectively, indicating that the TSS2 promoter is weaker than the TSS4 promoter

under these conditions.

CmrR autoregulation of *cmrRST* expression requires the TSS4 promoter

To further define the promoter autoregulated by CmrR, we generated reporter

strains in which different portions of the *cmrRST* regulatory region were transcriptionally

fused to a *phoZ* (alkaline phosphatase, AP) reporter gene (40). We first constructed a

strain in which *cmrR,* under the control of the P$_{tet}$ promoter, was inserted at an ectopic

site on the RT1693 (*recV cmr*-OFF) chromosome. This strain allowed ATc-inducible

expression of *cmrR* (Figure S4), prevented unintended inversion of the *cmr* switch

(Figure S2), and permitted the use of plasmid-borne *phoZ* reporters. Figure 6 shows the

combinations of the TSS1/c-di-GMP riboswitch, the *cmr* switch orientation (TSS2/ON or

TSS3/OFF), and the region encompassing TSS4.

The promoter corresponding to TSS1, located upstream the c-di-GMP riboswitch,

displayed similar transcriptional activity regardless of ATc (pRT2566), indicating that

CmrR does not autoregulate expression at this site. This result is consistent with the

qRT-PCR analysis of the native *cmrRST* site (Figure 5), and it further suggests that

cmrR overexpression does not impact global intracellular c-di-GMP levels. The

promoter corresponding to TSS3 (pRT2567), present in the *cmr* OFF switch orientation,

yielded no detectable AP activity when isolated from upstream (TSS1) and downstream

(TSS4) sequences and was not affected by overexpression of *cmrR*. Adding the

downstream sequence to the *cmr* OFF switch region (TSS3 + TSS4, pRT2514) resulted

in no AP activity in the absence of ATc but led to 14-fold increased AP activity when

cmrR was overexpressed.

Interestingly, the construct containing only the *cmr* ON switch sequence (TSS2,

pRT2568) also lacked transcriptional activity with and without *cmrR* overexpression.

The addition of the downstream sequence (TSS2 + TSS4, pRT2515) yielded baseline

AP activity 52-fold higher than the TSS2-only reporter and also conferred CmrR-

mediated autoactivation. Finally, the reporter containing the region downstream of the

cmr switch alone (TSS4, pRT2497) did not show AP activity in the absence of ATc but

did exhibit 19-fold activation by *cmrR* overexpression. The baseline AP activity and

degree of activation by CmrR was comparable to that from pRT2514 (TSS3 + TSS4),

confirming that there is no active promoter in the *cmr* switch when in the OFF

orientation under these conditions. Overall, only the constructs with the TSS4 region

resulted in increased activity upon *cmrR* overexpression, indicating that the TSS4

promoter is required for CmrR-mediated autoregulation of *cmrRST*. Further, the highest

transcriptional activity in the absence of *cmrR* overexpression was seen in the *cmr* ON

construct in which TSS2 and TSS4 are both present, suggesting additive effects of the

promoters or the disruption of an important sequence in the constructs separating the

two elements.

cmrRST and its upstream regulatory elements are well conserved in *C. difficile*

C. difficile strains from multiple ribotypes form rough and smooth colonies

associated with phase variable *cmrRST* expression (32). To examine the potential

scope of CmrRST phase variation and function in *C. difficile*, we used NCBI BLAST to

determine the extent of conservation of *cmrRST* and its upstream regulatory region

using R20291 as the reference sequence. The analysis examined the 895 bases 5' of

cmrR, which contains a c-di-GMP riboswitch and the *cmr* switch (Figure 1A), and the

cmrRST coding sequences. Among 71 *C. difficile* whole genome sequences available in

NCBI (taxonomic ID 1496), *C. difficile* strains shared high sequence identity to the

reference for both the riboswitch (98.9% ± 1.2) and the *cmr* invertible element (98.1% ±

1.4) (Figure S5). This similarity is comparable to the average percent sequence identity

for the *cmrRST* operon (98.7% ± 1.1). These results indicate that the c-di-GMP

riboswitch and invertible element upstream of *cmrRST* are highly conserved across

many *C. difficile* strains from divergent ribotypes and underscores the importance of this

operon and its regulation to *C. difficile* physiology.

Discussion

In this study, we demonstrate that multiple regulatory mechanisms control the

transcription of *cmrRST*. Our results support a model in which, under basal c-di-GMP

conditions, the orientation of the *cmr* switch determines the expression level of *cmrRST*.

This phase variation mechanism involves the reversible inversion of a promoter within

the *cmr* switch sequence. Increasing intracellular c-di-GMP augments *cmrRST*

expression independent of *cmr* switch orientation. Our data also show that expression

of *cmrRST* is also subject to autoregulation by CmrR, which occurs at an additional

promoter downstream of the *cmr* switch and is enhanced by the orientation of the *cmr*

switch. Therefore, multiple environmental signals may impact *cmrRST* expression

through c-di-GMP signaling, phase variation, and the activation of the response

regulator CmrR. CmrRST has important roles in *C. difficile* cell and colony morphology,

motility, biofilm formation, and virulence, suggesting multiple contexts in which distinct

environmental stimuli and selective pressures must be integrated to appropriately

control *cmrRST* expression at the single-cell and population levels. This work

represents the first account of a unique intersection of regulatory mechanisms

controlling the expression of a signal transduction system that broadly impacts *C.*

difficile physiology and disease development.

To disentangle the roles of c-di-GMP and phase variation in *cmrRST*

transcription, we phase locked the *cmr* switch by deleting three nucleotides in the RIR,

one of which was identified as the precise site of recombination during inversion of the

cmr switch (25). Phenotypic analysis of phase locked R20291 *cmr-*Δ3 ON and *cmr-* Δ3

OFF mutants showed that they behave similarly to WT rough and smooth populations,

respectively. Expression of *cmrRST* in the locked ON mutant was equivalent to that of

WT rough isolates, while expression in the *cmr-*Δ3 OFF mutant was similar to that of

WT smooth isolates. Consistent with these results, *cmr-*Δ3 ON yielded rough colonies,

increased surface motility and decreased swimming motility and biofilm formation as

compared to WT smooth isolates, the *cmr*-OFF mutant, and the Δ*cmrR*Δ*cmrT* mutant.

These data indicate that orientation of the *cmr* switch controls expression of *cmrRST*

and further confirms the role of CmrRST in these phenotypes. Interestingly, while *cmr-*

Δ3 ON forms rough colonies, they are not identical to those formed in WT populations,

and there is still some heterogeneity of colony morphology. This observation may reflect

differences in activity of CmrR or CmrT or that other factors contribute to colony

morphology. Other work has suggested heterogeneity in c-di-GMP levels among

individual *C. difficile* cells (21,24), so c-di-GMP may also result in colony morphology

differences via modulation of *cmrRST* expression.

By artificially manipulating intracellular c-di-GMP levels in the *cmr* locked

mutants, we found that c-di-GMP promotes *cmrRST* expression regardless of the

orientation of the *cmr* switch. These findings are consistent with observed phenotypes;

the *cmr*-Δ3 OFF mutant exhibited surface motility intermediate between that of WT and

Δ*cmrR*Δ*cmrT* mutant. In the *cmr*-Δ3 OFF mutant, *cmrRST* could still be expressed from

the TSS1 promoter upstream of the invertible element if c-di-GMP levels are sufficiently

high. Intracellular c-di-GMP levels have been shown to increase in *C. difficile* with

growth on a surface (39), which is consistent with c-di-GMP serving as a signal to

enhance CmrRST-mediated surface motility regardless of *cmr* switch orientation.

Multiple new TSS, in addition to the previously identified σ^A-dependent promoter

corresponding to TSS1 upstream of the c-di-GMP riboswitch (33,34), were identified

upstream of *cmrRST* using 5' RACE: TSS2 was identified within the *cmr* switch in *cmr*-

ON *C. difficile*, TSS3 was detected within the *cmr* switch in *cmr*-OFF, and TSS4 was

found between the *cmr* switch and the translational start of *cmrR* in both strains. TSS3

may have been a spurious result, as the associated putative promoter lacked detectable

transcriptional activity, and no -10/-35 sites similar to σ^{70} consensus sequences were

identifiable upstream of TSS3. The TSS2 region, when isolated from the TSS1 and

TSS4 regions, also lacked detectable transcriptional activity using the AP assay, and

transcript abundance measured by qRT-PCR was similarly low for the region

immediately downstream of TSS2. TSS2 may nonetheless indicate a weak promoter,

supported by the presence of -10 and -35 sites upstream of TSS2 (Figure S3) and its

responsiveness to CmrR. The level of TSS2 promoter activity, even with CmrR

activation, may have been below the detectable limit using the *phoZ* reporter. A longer

region containing both TSS2 and TSS4 showed maximal reporter activity, indicating that

both sequences must be present to allow optimal transcription initiation. This effect was

not observed for the equivalent TSS3 + TSS4 region, suggesting that the ON *cmr* switch

sequence contains information that enhances expression. Further, only the reporters

containing TSS4 exhibited CmrR-mediated autoactivation. The TSS4 region alone or

with the upstream TSS3 *cmr* OFF sequence showed equivalent increases in activity

upon over-expression of *cmrR*. Together these results support a model (Figure 7) in

which the TSS1 promoter (P1) acts independently of the downstream promoters and

yields an mRNA with a c-di-GMP riboswitch that enhances transcription readthrough

when the c-di-GMP ligand is bound. The *cmr* switch in the ON state contains a weak

promoter (P2) corresponding to TSS2 that can be activated by CmrR; the inverse

orientation of the *cmr* switch lacks a promoter. An additional promoter (P4) downstream

of the *cmr* switch is also activated by CmrR and strongly contributes to autoactivation of

cmrRST, particularly when the *cmr* switch is in the ON orientation. These regulatory

elements are poised to modulate *cmrRST* expression in response to distinct

environmental stimuli. Interestingly, CmrT does not regulate the expression of *cmrRST*,

suggesting that CmrR and CmrT have different DNA specificities and, accordingly,

distinct regulons. Future work will identify the genes directly and indirectly regulated by

CmrR, CmrT, or both as well as determine their consensus DNA binding sites, which

will elucidate the functions of these co-expressed response regulators.

In other bacteria, the reversible inversion of a promoter via site-specific DNA

recombination is common mechanism of phase variable expression of adjacent genes

(15,42). The *cmr* switch is the first instance of this mechanism of regulation in *C.*

difficile, though the inversion of the TSS2 promoter may be less impactful to *cmrRST*

transcription than other features of the *cmr* ON sequence combined with the

downstream TSS4 promoter. Notably, the region upstream of the *flgB* operon has a

similar arrangement to that of *cmrRST*, with a c-di-GMP riboswitch preceding the switch

that undergoes site-specific recombination. the flagellar switch that mediates phase

variation (21,27,34). However, the riboswitch upstream of *flgB* negatively regulates

transcriptional readthrough, so binding of c-di-GMP by the riboswitch leads to

transcription termination precluding synthesis of longer transcripts containing the

flagellar switch regardless of its orientation. Thus, phase variation and c-di-GMP

regulation of flagellar gene expression are linked, in contrast with the *cmrRST* system in

which c-di-GMP and the *cmr* switch independently modulate expression.

In summary, this work demonstrates that *cmrRST* expression is subject to

multilayered regulation with multiple potential inputs from environmental signals. The

complexity of this regulatory network suggests that *cmrRST* expression, and therefore

its transcriptional targets, requires careful control. Further work that defines the signals

which promote *cmrRST* expression will provide important insights into the role of this

TCS in *C. difficile* physiology and pathogenesis.

MATERIALS AND METHODS

Growth and maintenance of bacterial strains

481   Table S1 lists strains and plasmids used in this study. *C. difficile* R20291 and

482 derivative strains were maintained in an anaerobic environment of 90% N_2, 5% CO_2,

483 and 5% H_2. *C. difficile* strains were grown statically at 37 °C in BHI, BHIS (37 g/L Bacto

484 brain heart infusion, 5 g/L yeast extract), or in Tryptone Yeast (TY; 30 g/L Bacto

485 tryptone, 20 g/L yeast extract, 1 g/L thioglycolate) medium as indicated. *E. coli* strains

486 were grown in Luria Bertani medium at 37 °C with aeration. Antibiotics were used where

487 indicated at the following concentrations: chloramphenicol (Cm), 10 μg/mL;

488 thiamphenicol (Tm), 10 μg/mL; kanamycin (Kan), 100 μg/mL; ampicillin (Amp), 100

489 μg/mL.

491 **Construction of bacterial strains and plasmids**

492   Table S2 lists primers used in this study. Detailed methods for the generation of

493 plasmids and strains used in this study are provided in Supplemental Methods. Strain

494 and plasmid information is listed in Table S1.

496 **Quantitative reverse-transcriptase PCR**

497   For analysis of *cmrR* and *cmrT* expression in *C. difficile* R20291 (WT), *cmr-*Δ3

498 ON and *cmr-*Δ3 OFF, these strains were grown overnight (16 h) in TY medium, and 5

499 μL were spotted on BHIS-agar. After 24 hours, growth was collected, suspended in 1:1

500 ethanol:acetone, and stored at -80 ºC for subsequent RNA isolation. For analysis of

501 transcript abundance in R20291 strains carrying pCmrR, pCmrT, or vector, these strains

502 were grown overnight in TY-Tm medium. Cultures were diluted 1:30 in BHIS-Tm broth.

503 After two hours growth, ATc was added to induce gene expression (WT with vector or

pCmrR, 10 ng/mL; WT with pCmrT, 2 ng/mL). Samples were collected at mid-

exponential phase and saved in 1:1 ethanol:acetone at -80 °C.

RNA was extracted as described previously (43) and treated with TURBO DNA-

free™ Kit (Life Technologies) to remove contaminating genomic DNA. cDNA was

synthesized using the High-Capacity cDNA Reverse Transcription Kit (Applied

Biosystems) using the manufacturer's protocols (22,38). Real-time PCR was performed

using SensiFAST SYBR & Fluorescein Kit (Bioline) as previously described (25,32).

Data were analyzed using *rpoC* as the reference gene. Primers are as follows: *rpoC*,

R850/R851; *cmrR*, R2298/R2299; *cmrT*, R2537/R2538; *cmrS*, R2539/R2540; TSS1,

R2745/R2746; TSS2/3, R2751/R2804; TSS4, R2803/R2716.

Orientation specific PCR

Orientation specific PCR was done as previously described (21,32). Colonies

grown on BHIS-agar were boiled to produce lysate for PCR using primers R2270/R2271

to detect the ON orientation (241 bp product) and R2271/R2272 to detect the OFF

orientation (140 bp product). PCR products were separated on a TAE-1.5% agarose

gel.

Quantification of switch orientation by quantitative PCR

R20291 *cmrR*::SNAP pP$_{tet}$::*dccA* (RT2500) and vector control (RT2501) were

grown overnight in TY-Tm broth then diluted 1:30 in BHIS-Tm containing 20 ng/μL to

induce *dccA* expression. After growth to mid-exponential phase, genomic DNA was

purified by phenol:chloroform:isopropanol extraction and ethanol precipitation. qPCR

was performed as previously described with 100 ng of DNA per 20 μL reaction and 100

nM primers (32). Data were analyzed using the ΔΔCt method, with *rpoA* as the

reference gene and the indicated control condition. Primers are as follows: ON

orientation, R2270/R2271; OFF orientation, R2271/R2272; and *rpoA,* R2273/R2274.

Microscopy

To image whole colonies, strains were grown on BHIS-Tm agar for 24 hours. For

cmr-Δ3 ON/OFF with pP$_{tet}$::*dccA*, pP$_{tet}$::EAL, or vector, ATc (20 ng/mL) was included in

the agar. Colonies were imaged using a Keyence BZ-X810 microscope or Zeiss Stereo

Discovery V8 dissecting microscope with a glass stage.

Fluorescence microscopy of SNAP-labelled strains was performed as previously

described (25). R20291 *cmrR*::SNAP pP$_{tet}$::*dccA* or vector control were grown overnight

in TY-Tm broth. Cultures were diluted 1:30 in BHIS-Tm. After two hours of growth, ATc

(20 ng/mL) was added for induction. Mid-exponential phase samples were pelleted,

washed, and incubated with SNAP-Cell TMR-Star (New England Biolabs) at 37 ºC for

30 minutes. Samples were further washed and mounted on 1% agarose pads. Imaging

was done using a Keyence BZ-X810 microscope with an Olympus 100X PlanFluor

objective. Bright field and visualization of red fluorescence with a Chroma filter

(ET545/30x, EM620/60m) was done using the same image capture settings for all

samples. Cells were counted on at least three fields from two biological replicates for

each strain and condition.

Phenotype assays

Motility and biofilm assays were done as previously described (32,38); further

details are provided in Supplemental Methods.

Alkaline phosphatase assays

554   Strains carrying *phoZ* reporters were grown in BHIS medium to OD_{600} ~ 1.0, then

555   1.5 mL of culture was collected, pelleted and frozen at -20 °C. Samples were thawed,

556   and AP activity using the substrate *p*-nitrophenyl phosphate was measured as

557   previously described (21,40).

5' rapid amplification of cDNA ends (RACE)

R20291 *recV cmr*-OFF (RT1693), *recV cmr*-OFF pDccA (RT2184), and *recV*

cmr-ON (RT2520) were grown in BHIS to mid-exponential phase. RT2184 was grown

with Tm and induced with 1 μg/mL nisin (induces *dccA* expression from the P*cpr*

promoter). Samples were collected and frozen in 1:1 ethanol:acetone. RNA was

extracted and purified using the RNeasy Mini Kit (Qiagen). cDNA was synthesized with

the 5' RACE System for Rapid Amplification of cDNA Ends kit (Thermo Fisher) using the

manufacturer's protocol. Briefly, cDNA was synthesized using primer R2792 and

SuperScript II reverse transcriptase. The samples were treated with RNase Mix then

column purified and treated with TdT. The products were then PCR amplified with the

provided anchor primer and primer R2793, then subjected to Sanger sequencing.

Acknowledgements

References

1. Trastoy, R., Manso, T., Fernandez-Garcia, L., Blasco, L., Ambroa, A., Perez Del Molino, M. L., Bou, G., Garcia-Contreras, R., Wood, T. K., and Tomas, M. (2018) Mechanisms of Bacterial Tolerance and Persistence in the Gastrointestinal and Respiratory Environments. *Clin Microbiol Rev* **31**, e00023-00018
2. Magdanova, L. A., and Golyasnaya, N. V. (2013) Heterogeneity as an Adaptive Trait of Microbial Populations. *Microbiology* **82**, 1-10
3. Casadesus, J., and Low, D. A. (2013) Programmed Heterogeneity: Epigenetic Mechanisms in Bacteria. *Journal of Biological Chemistry* **288**, 13929-13935
4. Fang, F. C., Frawley, E. R., Tapscott, T., and Vazquez-Torres, A. (2016) Discrimination and Integration of Stress Signals by Pathogenic Bacteria. *Cell Host & Microbe* **20**, 144-153
5. Bourret, R. B., and Silversmith, R. E. (2010) Two-component signal transduction. *Current Opinion in Microbiology* **13**, 113-115
6. Krell, T., Lacal, J., Busch, A., Silva-Jimenez, H., Guazzaroni, M. E., and Ramos, J. L. (2010) Bacterial Sensor Kinases: Diversity in the Recognition of Environmental Signals. *Annual Review of Microbiology, Vol 64, 2010* **64**, 539-559
7. Hengge, R., Grundling, A., Jenal, U., Ryan, R., and Yildiz, F. (2016) Bacterial Signal Transduction by Cyclic Di-GMP and Other Nucleotide Second Messengers. *Journal of Bacteriology* **198**, 15-26
8. Purcell, E. B., and Tamayo, R. (2016) Cyclic diguanylate signaling in Gram-positive bacteria. *FEMS microbiology reviews* **40**, 753-773
9. Hengge, R. (2009) Principles of c-di-GMP signalling in bacteria. *Nature reviews.Microbiology* **7**, 263-273
10. Purcell, E. B., McKee, R. W., Courson, D. S., Garrett, E. M., McBride, S. M., Cheney, R. E., and Tamayo, R. (2017) A nutrient-regulated cyclic diguanylate phosphodiesterase controls *Clostridium difficile* biofilm and toxin production during stationary phase. *Infect Immun* **85**, pii: e00347-00317
11. Sudarsan, N., Lee, E. R., Weinberg, Z., Moy, R. H., Kim, J. N., Link, K. H., and Breaker, R. R. (2008) Riboswitches in eubacteria sense the second messenger cyclic di-GMP. *Science (New York, N.Y.)* **321**, 411-413
12. Smith, K. D., Lipchock, S. V., Ames, T. D., Wang, J., Breaker, R. R., and Strobel, S. A. (2009) Structural basis of ligand binding by a c-di-GMP riboswitch. *Nature structural & molecular biology* **16**, 1218-1223
13. Chou, S. H., and Galperin, M. Y. (2016) Diversity of Cyclic Di-GMP-Binding Proteins and Mechanisms. *J Bacteriol* **198**, 32-46
14. Ryall, B., Eydallin, G., and Ferenci, T. (2012) Culture History and Population Heterogeneity as Determinants of Bacterial Adaptation: the Adaptomics of a Single Environmental Transition. *Microbiol Mol Biol R* **76**, 597-625
15. van der Woude, M. W., and Baumler, A. J. (2004) Phase and antigenic variation in bacteria. *Clin Microbiol Rev* **17**, 581-611
16. van der Woude, M. W. (2011) Phase variation: how to create and coordinate population diversity. *Current opinion in microbiology* **14**, 205-211
17. Abraham, J. M., Freitag, C. S., Clements, J. R., and Eisenstein, B. I. (1985) An invertible element of DNA controls phase variation of type 1 fimbriae of Escherichia coli. *Proceedings of the National Academy of Sciences of the United States of America* **82**, 5724-5727
18. Klemm, P. (1986) Two regulatory fim genes, fimB and fimE, control the phase variation of type 1 fimbriae in Escherichia coli. *The EMBO journal* **5**, 1389-1393

19.   Blomfield, I., and van der Woude, M. (2007) Regulation of Fimbrial Expression. *EcoSal*
Plus **2**, 10.1128/ecosal.1122.1124.1122.1122

20.   Emerson, J. E., Reynolds, C. B., Fagan, R. P., Shaw, H. A., Goulding, D., and
Fairweather, N. F. (2009) A novel genetic switch controls phase variable expression of
CwpV, a *Clostridium difficile* cell wall protein. *Mol Microbiol* **74**, 541-556

21.   Anjuwon-Foster, B. R., and Tamayo, R. (2017) A genetic switch controls the production
of flagella and toxins in *Clostridium difficile. PLoS Genet* **13**, e1006701

22.   Anjuwon-Foster, B. R., Maldonado-Vazquez, N., and Tamayo, R. (2018)
Characterization of flagellar and toxin phase variation in *Clostridiodes difficile* ribotype
012 isolates. *J Bacteriol* **200**, e00056-00018

23.   Anjuwon-Foster, B. R., and Tamayo, R. (2018) Phase variation of *Clostridium difficile*
virulence factors. *Gut Microbes* **9**, 76-83

24.   Trzilova, D., Anjuwon-Foster, B. R., Rivera, D. T., and Tamayo, R. (2020) Rho factor
mediates flagellum and toxin phase variation and impacts virulence in *Clostridioides*
difficile. Plos Pathog **16**, e1008708

25.   Sekulovic, O., Mathias Garrett, E., Bourgeois, J., Tamayo, R., Shen, A., and Camilli, A.
(2018) Genome-wide detection of conservative site-specific recombination in bacteria.
PLoS Genet **14**, e1007332

26.   Sekulovic, O., Bourgeois, J., Shen, A., and Camilli, A. (2019) Expanding the repertoire of
conservative site-specific recombination in *Clostridioides difficile. Anaerobe* **60**, 102073

27.   McKee, R. W., Mangalea, M. R., Purcell, E. B., Borchardt, E. K., and Tamayo, R. (2013)
The second messenger cyclic di-GMP regulates *Clostridium difficile* toxin production by
controlling expression of *sigD. J Bacteriol* **195**, 5174-5185

28.   El Meouche, I., Peltier, J., Monot, M., Soutourina, O., Pestel-Caron, M., Dupuy, B., and
Pons, J. L. (2013) Characterization of the SigD regulon of *C. difficile* and its positive
control of toxin production through the regulation of tcdR. *PLoS One* **8**, e83748

29.   Reynolds, C. B., Emerson, J. E., de la Riva, L., Fagan, R. P., and Fairweather, N. F.
(2011) The *Clostridium difficile* cell wall protein CwpV is antigenically variable between
strains, but exhibits conserved aggregation-promoting function. *PLoS Pathog* **7**,
e1002024

30.   Sekulovic, O., Ospina Bedoya, M., Fivian-Hughes, A. S., Fairweather, N. F., and Fortier,
L. C. (2015) The *Clostridium difficile* cell wall protein CwpV confers phase-variable
phage resistance. *Mol Microbiol* **98**, 329-342

31.   Sekulovic, O., and Fortier, L. C. (2015) Global transcriptional response of Clostridium
difficile carrying the CD38 prophage. *Applied and Environmental Microbiology* **81**, 1364-
1374

32.   Garrett, E. M., Sekulovic, O., Wetzel, D., Jones, J. B., Edwards, A. N., Vargas-Cuebas,
G., McBride, S. M., and Tamayo, R. (2019) Phase variation of a signal transduction
system controls *Clostridioides difficile* colony morphology, motility, and virulence. *Plos*
Biol **17**, e3000379

33.   Soutourina, O. A., Monot, M., Boudry, P., Saujet, L., Pichon, C., Sismeiro, O.,
Semenova, E., Severinov, K., Le Bouguenec, C., Coppee, J. Y., Dupuy, B., and Martin-
Verstraete, I. (2013) Genome-wide identification of regulatory RNAs in the human
pathogen Clostridium difficile. *PLoS genetics* **9**, e1003493

34.   McKee, R. W., Harvest, C. K., and Tamayo, R. (2018) Cyclic Diguanylate Regulates
Virulence Factor Genes via Multiple Riboswitches in *Clostridium difficile. mSphere* **3**,
e00423-00418

35.   Snyder, J. A., Lloyd, A. L., Lockatell, C. V., Johnson, D. E., and Mobley, H. L. (2006)
Role of phase variation of type 1 fimbriae in a uropathogenic *Escherichia coli* cystitis
isolate during urinary tract infection. *Infect Immun* **74**, 1387-1393

36. Grindley, N. D. F., Whiteson, K. L., and Rice, P. A. (2006) Mechanisms of site-specific
recombination. *Annu Rev Biochem* **75**, 567-605
37. Trzilova, D., Warren, M. A. H., Gadda, N. C., Williams, C. L., and Tamayo, R. (2021)
Flagellum and toxin phase variation impacts colonization and disease development in a
mouse model of *Clostridioides difficile* infection. *bioRxiv*
38. Purcell, E. B., McKee, R. W., McBride, S. M., Waters, C. M., and Tamayo, R. (2012)
Cyclic diguanylate inversely regulates motility and aggregation in *Clostridium difficile*. *J*
Bacteriol **194**, 3307-3316
39. Purcell, E. B., McKee, R. W., Bordeleau, E., Burrus, V., and Tamayo, R. (2015)
Regulation of Type IV Pili Contributes to Surface Behaviors of Historical and Epidemic
Strains of *Clostridium difficile*. *Journal of Bacteriology* **198**, 565-577
40. Edwards, A. N., Pascual, R. A., Childress, K. O., Nawrocki, K. L., Woods, E. C., and
McBride, S. M. (2015) An alkaline phosphatase reporter for use in Clostridium difficile.
Anaerobe **32**, 98-104
41. Groisman, E. A. (2016) Feedback Control of Two-Component Regulatory Systems.
Annu Rev Microbiol **70**, 103-124
42. Jiang, X., Hall, A. B., Arthur, T. D., Plichta, D. R., Covington, C. T., Poyet, M., Crothers,
J., Moses, P. L., Tolonen, A. C., Vlamakis, H., Alm, E. J., and Xavier, R. J. (2019)
Invertible promoters mediate bacterial phase variation, antibiotic resistance, and host
adaptation in the gut. *Science* **363**, 181-187
43. Bouillaut, L., McBride, S. M., and Sorg, J. A. (2011) Genetic Manipulation of *Clostridium*
difficile. *Curr Protoc Microbiol* **Unit 9A.2**, 1-17

Figure 1. Generation of genetically phase-locked strains. (A) Diagram of the

cmrRST locus. The previously identified σ^A-dependent promoter and c-di-GMP

riboswitch Cdi-2-2 are indicated. Black triangles represent the inverted repeats flanking

the invertible DNA sequence (*cmr* switch). The OFF orientation corresponds to the

sequence present in the R20291 reference genome (FN545816); the ON orientation

corresponds to the inverse of that sequence. (B) Sequence of the right inverted repeat

(RIR) of the *cmr* switch. Gray shading indicates sequence identity between the

imperfect inverted repeats of the RIR in each orientation. Three nucleotides (shown in

red) were deleted from the RIR to lock the *cmr* switch in the ON and OFF orientations.

The asterisk (*) indicates the nucleotide at the site of recombination. (C) Orientation-

specific PCR to detect each orientation of the *cmr* switch in WT, *cmr-Δ3* OFF, and *cmr-*

Δ3 ON strains. (D) qRT-PCR analysis for the *cmrR* and *cmrT* transcripts in WT rough

and smooth isolates, *cmr-Δ3* OFF and *cmr-Δ3* ON. Data from four biological replicates

were analyzed using the ΔΔCt method with *rpoC* as the reference gene and

normalization to the WT smooth samples. Shown are means and standard error. #$p <$

0.0001, two-way ANOVA with Sidak's multiple comparisons post-test comparing to the

WT/smooth condition.

Figure 2. Mutating the RIR of the *cmr* switch results in phenotypically phase-locked strains. (A) Colony morphology of WT rough and smooth isolates, *cmr*-Δ3 OFF, *cmr*-Δ3 ON, and Δ*cmrR*Δ*cmrT*. Shown are representative images after 24 hours growth on BHIS agar. (B) Surface motility of WT, *cmr*-Δ3 OFF, *cmr*-Δ3 ON, and Δ*cmrR*Δ*cmrT* quantified by measuring the diameter of growth after 72 hours. (C) Representative images of surface motility after 72 hours of growth. (D) Swimming motility of WT, *cmr*-Δ3 OFF, *cmr*-Δ3 ON, and Δ*cmrR*Δ*cmrT* mutant quantified by measuring the diameter of growth in 0.5x BHIS 0.3% agar after 48 hours. E) Biofilm formation after 24 hours in BHIS-1% glucose-50 mM sodium phosphate, quantified by crystal violet staining. (B, D, E) Shown are means and standard error, with symbols representing independent samples from at least 2 independent experiments. *p < 0.05, ****p < 0.0001, one-way ANOVA with Dunnett's multiple comparisons post-test compared to the WT.

Figure 3. Regulation of *cmrRST* by c-di-GMP occurs independent of *cmr* switch

orientation. (A) Colony morphology of *cmr*-Δ3 OFF and *cmr*-Δ3 ON containing either

vector, pP$_{tet}$::*dccA*, or pP$_{tet}$::EAL. Strains were grown on BHIS-agar with 20 ng/ml ATc to

induce DGC and EAL gene expression. (B) qRT-PCR analysis of *cmrR* transcript

abundance in *cmr*-Δ3 OFF and *cmr*-Δ3 ON strains carrying pP$_{tet}$::*dccA* or vector control

grown in BHIS broth with 20 ng/ml ATc. Data were analyzed using the ΔΔCt method

with *rpoC* as the reference gene and normalization to *cmr*-Δ3 OFF with vector. Shown

are means with standard error of 6 biological replicates from two independent

experiments. *p < 0.05, ****p < 0.0001, one-way ANOVA with Tukey's multiple

comparisons post-test. C) Representative images of Δ*cmrR*::SNAP carrying pP$_{tet}$::*dccA*

or vector control was grown in BHIS broth with 20 ng/ml ATc. Samples were imaged at

60x magnification by light and fluorescence microscopy following staining with SNAP-

Cell TMR-Star substrate. Percentages of total cells that were fluorescence-positive were

determined from at least three fields each from two biological replicates. Shown are

means and standard deviations.

Figure 4. Mapping of transcriptional start sites (TSS) upstream of *cmrRST*. (A) Diagram of TSS identified by 5' RACE using RNA derived from *cmr* ON and *cmr* OFF templates. (B) Alkaline phosphatase reporters used to detect promoter activity. Brackets indicate the regions from the native *cmr* locus that were used to make transcriptional reporters. Construct 1 contains the region between the 3' end of the right inverted repeat and the 5' end of *cmrR* coding sequence (pMC123*::TSS4-phoZ*). Construct 2 includes the region in construct 1 plus the *cmr* switch in the OFF orientation (pMC123*::cmr*OFF-*phoZ*). Construct/TSS4 3 is identical to construct 2 but with the *cmr* switch in the ON orientation (pMC123*::cmr*ON/TSS4-*phoZ*). Shown are the means and standard deviations from 12 biological replicates from four independent experiments. ****p < 0.0001, one-way ANOVA with Dunnett's multiple comparisons test.

Figure 5. CmrR positively regulates *cmrRST* expression. qRT-PCR measurement of

cmrRST transcript abundance. Strains were grown with ATc induction (10 ng/mL for

vector and pCmrR, 2 ng/mL for pCmrT) prior to RNA isolation. (A) WT with pCmrR,

pCmrT, and vector control. n.d., not determined; *cmrR* and *cmrT* were not measured in

strains overexpressing those genes. ****p < 0.0001, unpaired t-test. (B) qRT-PCR

analysis of *cmrS* transcript abundance of *recV cmr* OFF and *cmr* ON, each with pCmrR
or vector control. The *recV cmr*-OFF with vector strain serves as the reference condition
for ΔΔCt analysis. Shown are means and standard deviations of 4 biological replicates.
(C) Diagram showing regions of transcript measured in (D). (D) qRT-PCR analysis of
cmrRST transcripts with values normalized to *rpoC*. (B, D) Shown are means and
standard error from independent replicates. $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p <$
0.0001, one-way ANOVA with Tukey's multiple comparisons.

770
Figure 6. CmrR autoregulation occurs at the TSS4 promoter and is enhanced by

the *cmr* ON switch sequence. Alkaline phosphatase reporter activity for regions of the

cmr 5' UTR containing one or more putative promoter in *C. difficile* with ATc-inducible

Ptet::*cmrR* at an ectopic site. (Left) Diagrams of the plasmid-borne reporter fusions

carried by the strain. (Right) AP reporter activity after growth with or without 10 ng/mL

ATc to induce cmrR expression (Figure S4). Shown are means and standard deviations

from three independent experiments. ****$p < 0.0001$, two-way ANOVA and Sidak's post-

test.

Figure 7. Model of multilevel regulation of *cmrRST* transcription. Three promoters control the expression of *cmrRST*. Promoter P1 corresponding to TSS1 generates an mRNA with a c-di-GMP riboswitch that is permissive for transcriptional readthrough when c-di-GMP is bound. The *cmr* switch contains a weak promoter (P2) corresponding the TSS2 when in the ON orientation. An additional promoter, P4, lies between the *cmr* switch and the *cmrRST* coding sequences. Transcription from P4 is positively regulated by CmrR. P1 and P2 independently regulate *cmrRST* transcription, while P4 activity is enhanced when the *cmr* switch is in the ON orientation.

 SUPPLEMENTAL FIGURES

Figure S1. C-di-GMP does not cause inversion of the *cmr* switch. Δ*cmrR*::SNAP

carrying pP$_{tet}$::*dccA* or vector control was grown to mid-exponential phase in BHIS broth

with or without 20 ng/ml ATc. gDNA was collected for qPCR analysis of *cmr* switch

orientation. Data are expressed as the percent OFF orientation. Shown are the means

and standard deviations of six biological replicates from two independent experiments.

No significant differences, two-way ANOVA with Tukey's multiple comparisons.

Figure S2. The *cmr* switch is phase locked in *recV*-deficient strains. Orientation-

specific PCR to detect each orientation of the *cmr* switch in WT, *recV cmr*-OFF, and

recV cmr-ON.

Figure S3. Identification and mapping of multiple transcriptional start sites

upstream of *cmrRST*. Map of TSS identified by 5' RACE in *recV cmr*-ON, *recV cmr*-

OFF and *recV cmr*-OFF pDccA. Gray highlight indicates the Cdi-2-2 riboswitch

sequence. Red and blue text denote the OFF and ON orientations of the *cmr* switch,

respectively. Bold text indicates the inverted repeats. Green highlights mark TSS.

Underlined text indicates putative -10/-35 sites. Yellow highlight denotes the *cmrR* start

codon.

Figure S4. Inducible expression of *cmrR* from an ectopic chromosomal site.

R20291 *recV* CDR2492::P$_{tet}$::*cmrR* was grown to mid-exponential phase in BHIS

medium with and without 10 µg/mL ATc to induce *cmrR* expression. qRT-PCR was

used to measure *cmrR* transcript abundance using 3 different primer sets to distinguish

cmrR expression from the native site (primers R3111/R3112), from the ectopic site

(primers R3113/R3112), and total *cmrR* (primers R2298/R2299). Data are expressed as

relative transcript abundance in the ATc condition compared to that without ATc, with

normalization to *rpoC* as the reference strain. Induction with ATc resulted in a 9.0-fold

increase in *cmrR* mRNA from the ectopic site and a 2.6-fold increase from the native

cmrRST locus, with a cumulative 12.7-fold increase in *cmrR* transcript. Shown are

means and standard error. * $p < 0.05$, *** $p < 0.001$, **** $p < 0.0001$ by unpaired t-test.

Figure S5. *cmrRST* and upstream regulatory sequences are highly conserved

across *C. difficile* strains and ribotypes. Heat map showing percent sequence

identity of 71 sequenced *C. difficile* strains in the NCBI database compared to the

R20291 reference sequence (NCBI accession number FN545816.1). If the strain has

been ribotyped, the ribotype is indicated on the left. For the *cmr* switch, both

orientations were used as the reference and the highest percent sequence identity is

shown. The following nucleotides were used as the reference sequence from R20291:

riboswitch, 3,736,512-3,736,863; *cmr* switch, 3,735,969-3,736,513; *cmrRST*, 3,732,474-

3,735,968.